30692011924956

WITHDRAWN FROM
COLGATE UNIVERSITY LIBRARY

Complete Solutions Manual
for
MULTIVARIABLE CALCULUS
EIGHTH EDITION

DAN CLEGG
Palomar College

BARBARA FRANK
Cape Fear Community College

CENGAGE Learning

Australia · Brazil · Mexico · Singapore · United Kingdom · United States

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

© 2016 Cengage Learning

WCN: 01-100-101

ALL RIGHTS RESERVED. No part of this work covered by the copyright herein may be reproduced, transmitted, stored, or used in any form or by any means graphic, electronic, or mechanical, including but not limited to photocopying, recording, scanning, digitizing, taping, Web distribution, information networks, or information storage and retrieval systems, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without the prior written permission of the publisher except as may be permitted by the license terms below.

For product information and technology assistance, contact us at
Cengage Learning Customer & Sales Support,
1-800-354-9706

For permission to use material from this text or product, submit all requests online at **www.cengage.com/permissions.** Further permissions questions can be e-mailed to **permissionrequest@cengage.com.**

ISBN: 978-1-305-27611-6

Cengage Learning
20 Channel Center Street
Boston, MA 02210
USA

Cengage Learning is a leading provider of customized learning solutions with employees residing in nearly 40 different countries and sales in more than 125 countries around the world. Find your local representative at: **www.cengage.com**.

Cengage Learning products are represented in Canada by Nelson Education, Ltd.

To learn more about Cengage Learning Solutions, visit **www.cengage.com**.

Purchase any of our products at your local college store or at our preferred online store **www.cengagebrain.com**.

NOTE: UNDER NO CIRCUMSTANCES MAY THIS MATERIAL OR ANY PORTION THEREOF BE SOLD, LICENSED, AUCTIONED, OR OTHERWISE REDISTRIBUTED EXCEPT AS MAY BE PERMITTED BY THE LICENSE TERMS HEREIN.

Dear Professor or Other Supplement Recipient:

Cengage Learning has provided you with this product (the "Supplement") for your review and, to the extent that you adopt the associated textbook for use in connection with your course (the "Course"), you and your students who purchase the textbook may use the Supplement as described below. Cengage Learning has established these use limitations in response to concerns raised by authors, professors, and other users regarding the pedagogical problems stemming from unlimited distribution of Supplements.

Cengage Learning hereby grants you a nontransferable license to use the Supplement in connection with the Course, subject to the following conditions. The Supplement is for your personal, noncommercial use only and may not be reproduced, posted electronically or distributed, except that portions of the Supplement may be provided to your students IN PRINT FORM ONLY in connection with your instruction of the Course, so long as such students are advised that they may not copy or distribute any portion of the Supplement to any third party. You may not sell, license, auction, or otherwise redistribute the Supplement in any form. We ask that you take reasonable steps to protect the Supplement from unauthorized use, reproduction, or distribution. Your use of the Supplement indicates your acceptance of the conditions set forth in this Agreement. If you do not accept these conditions, you must return the Supplement unused within 30 days of receipt.

All rights (including without limitation, copyrights, patents, and trade secrets) in the Supplement are and will remain the sole and exclusive property of Cengage Learning and/or its licensors. The Supplement is furnished by Cengage Learning on an "as is" basis without any warranties, express or implied. This Agreement will be governed by and construed pursuant to the laws of the State of New York, without regard to such State's conflict of law rules.

Thank you for your assistance in helping to safeguard the integrity of the content contained in this Supplement. We trust you find the Supplement a useful teaching tool.

Printed in the United States of America
Print Number: 01 Print Year: 2015

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

PREFACE

This *Complete Solutions Manual* contains detailed solutions to all exercises in the text *Multivariable Calculus,* Eighth Edition (Chapters 10–17 of *Calculus,* Eighth Edition, and *Calculus: Early Transcendentals,* Eighth Edition) by James Stewart. A *Student Solutions Manual* is also available, which contains solutions to the odd-numbered exercises in each chapter section, review section, True-False Quiz, and Problems Plus section. (It does not, however, include solutions to any of the projects.)

Because of differences between the regular version and the *Early Transcendentals* version of the text, some references are given in a dual format. In these cases, users of the *Early Transcendentals* text should use the references denoted by "ET."

While we have extended every effort to ensure the accuracy of the solutions presented, we would appreciate correspondence regarding any errors that may exist. Other suggestions or comments are also welcome, and can be sent to dan clegg at `dclegg@palomar.edu` or in care of the publisher: Cengage Learning, 20 Channel Center Street, Boston MA 02210.

We would like to thank James Stewart for entrusting us with the writing of this manual and offering suggestions, Gina Sanders for reviewing solutions for accuracy and style, and the staff of TECH-arts for typesetting and producing this manual as well as creating the illustrations. We also thank Richard Stratton, Neha Taleja, Samantha Lugtu, Stacy Green, and Terry Boyle of Cengage Learning, for their trust, assistance, and patience.

DAN CLEGG
Palomar College

BARBARA FRANK
Cape Fear Community College

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Digitized by the Internet Archive
in 2022 with funding from
Kahle/Austin Foundation

https://archive.org/details/completesolution0000cleg

ABBREVIATIONS AND SYMBOLS

CD	concave downward
CU	concave upward
D	the domain of f
FDT	First Derivative Test
HA	horizontal asymptote(s)
I	interval of convergence
IP	inflection point(s)
R	radius of convergence
VA	vertical asymptote(s)
$\overset{\text{CAS}}{=}$	indicates the use of a computer algebra system.
$\overset{\text{PR}}{\Longrightarrow}$	indicates the use of the Product Rule.
$\overset{\text{QR}}{\Longrightarrow}$	indicates the use of the Quotient Rule.
$\overset{\text{CR}}{\Longrightarrow}$	indicates the use of the Chain Rule.
$\overset{\text{H}}{=}$	indicates the use of l'Hospital's Rule.
$\overset{j}{=}$	indicates the use of Formula j in the Table of Integrals in the back endpapers.
$\overset{\text{s}}{=}$	indicates the use of the substitution $\{u = \sin x,\ du = \cos x\ dx\}$.
$\overset{\text{c}}{=}$	indicates the use of the substitution $\{u = \cos x,\ du = -\sin x\ dx\}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

CONTENTS

10 PARAMETRIC EQUATIONS AND POLAR COORDINATES 1

10.1 Curves Defined by Parametric Equations 1

Laboratory Project ▫ Running Circles Around Circles 15

10.2 Calculus with Parametric Curves 18

Laboratory Project ▫ Bézier Curves 32

10.3 Polar Coordinates 33

Laboratory Project ▫ Families of Polar Curves 47

10.4 Areas and Lengths in Polar Coordinates 50

10.5 Conic Sections 62

10.6 Conic Sections in Polar Coordinates 73

Review 79

Problems Plus 91

11 INFINITE SEQUENCES AND SERIES 95

11.1 Sequences 95

Laboratory Project ▫ Logistic Sequences 108

11.2 Series 111

11.3 The Integral Test and Estimates of Sums 128

11.4 The Comparison Tests 137

11.5 Alternating Series 143

11.6 Absolute Convergence and the Ratio and Root Tests 148

11.7 Strategy for Testing Series 156

11.8 Power Series 160

11.9 Representations of Functions as Power Series 169

11.10 Taylor and Maclaurin Series 178

Laboratory Project ▫ An Elusive Limit 195

11.11 Applications of Taylor Polynomials 196

Applied Project ▫ Radiation from the Stars 210

Review 211

Problems Plus 223

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

12 VECTORS AND THE GEOMETRY OF SPACE 235

12.1 Three-Dimensional Coordinate Systems 235
12.2 Vectors 242
12.3 The Dot Product 251
12.4 The Cross Product 260
Discovery Project ▫ The Geometry of a Tetrahedron 271
12.5 Equations of Lines and Planes 273
Laboratory Project ▫ Putting 3D in Perspective 285
12.6 Cylinders and Quadric Surfaces 287
Review 298

Problems Plus 307

13 VECTOR FUNCTIONS 313

13.1 Vector Functions and Space Curves 313
13.2 Derivatives and Integrals of Vector Functions 325
13.3 Arc Length and Curvature 335
13.4 Motion in Space: Velocity and Acceleration 351
Applied Project ▫ Kepler's Laws 362
Review 363

Problems Plus 371

14 PARTIAL DERIVATIVES 377

14.1 Functions of Several Variables 377
14.2 Limits and Continuity 395
14.3 Partial Derivatives 403
14.4 Tangent Planes and Linear Approximations 421
Applied Project ▫ The Speedo LZR Racer 431
14.5 The Chain Rule 432
14.6 Directional Derivatives and the Gradient Vector 445
14.7 Maximum and Minimum Values 458
Applied Project ▫ Designing a Dumpster 479
Discovery Project ▫ Quadratic Approximations and Critical Points 481

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.8 Lagrange Multipliers 484
Applied Project ▫ Rocket Science 497
Applied Project ▫ Hydro-Turbine Optimization 500
Review 502

Problems Plus 517

□ 15 MULTIPLE INTEGRALS 523

15.1 Double Integrals over Rectangles 523
15.2 Double Integrals over General Regions 532
15.3 Double Integrals in Polar Coordinates 545
15.4 Applications of Double Integrals 552
15.5 Surface Area 563
15.6 Triple Integrals 567
Discovery Project ▫ Volumes of Hyperspheres 584
15.7 Triple Integrals in Cylindrical Coordinates 585
Discovery Project ▫ The Intersection of Three Cylinders 592
15.8 Triple Integrals in Spherical Coordinates 594
Applied Project ▫ Roller Derby 605
15.9 Change of Variables in Multiple Integrals 606
Review 612

Problems Plus 625

□ 16 VECTOR CALCULUS 633

16.1 Vector Fields 633
16.2 Line Integrals 638
16.3 The Fundamental Theorem for Line Integrals 648
16.4 Green's Theorem 654
16.5 Curl and Divergence 661
16.6 Parametric Surfaces and Their Areas 670
16.7 Surface Integrals 684
16.8 Stokes' Theorem 695
16.9 The Divergence Theorem 699
Review 705

Problems Plus 713

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

□ 17 SECOND-ORDER DIFFERENTIAL EQUATIONS 719

17.1 Second-Order Linear Equations 719

17.2 Nonhomogeneous Linear Equations 723

17.3 Applications of Second-Order Differential Equations 729

17.4 Series Solutions 733

Review 738

□ APPENDIX 743

Complex Numbers 743

© 20160 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10 ☐ PARAMETRIC EQUATIONS AND POLAR COORDINATES

10.1 Curves Defined by Parametric Equations

1. $x = 1 - t^2, \quad y = 2t - t^2, \quad -1 \le t \le 2$

t	-1	0	1	2
x	0	1	0	-3
y	-3	0	1	0



2. $x = t^3 + t, \quad y = t^2 + 2, \quad -2 \le t \le 2$

t	-2	-1	0	1	2
x	-10	-2	0	2	10
y	6	3	2	3	6	



3. $x = t + \sin t, \quad y = \cos t, \quad -\pi \le t \le \pi$

t	$-\pi$	$-\pi/2$	0	$\pi/2$	π
x	$-\pi$	$-\pi/2+1$	0	$\pi/2+1$	π
y	-1	0	1	0	-1



4. $x = e^{-t} + t, \quad y = e^t - t, \quad -2 \le t \le 2$

t	-2	-1	0	1	2
x	$e^2 - 2$ 5.39	$e - 1$ 1.72	1	$e^{-1} + 1$ 1.37	$e^{-2} + 2$ 2.14
y	$e^{-2} + 2$ 2.14	$e^{-1} + 1$ 1.37	1	$e - 1$ 1.72	$e^2 - 2$ 5.39



5. $x = 2t - 1, \quad y = \frac{1}{2}t + 1$

(a)

t	-4	-2	0	2	4
x	-9	-5	-1	3	7
y	-1	0	1	2	3



(b) $x = 2t - 1 \quad \Rightarrow \quad 2t = x + 1 \quad \Rightarrow \quad t = \frac{1}{2}x + \frac{1}{2}$, so

$$y = \tfrac{1}{2}t + 1 = \tfrac{1}{2}\left(\tfrac{1}{2}x + \tfrac{1}{2}\right) + 1 = \tfrac{1}{4}x + \tfrac{1}{4} + 1 \quad \Rightarrow \quad y = \tfrac{1}{4}x + \tfrac{5}{4}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $x = 3t + 2, \quad y = 2t + 3$

(a)

t	-4	-2	0	2	4
x	-10	-4	2	8	14
y	-5	-1	3	7	11



(b) $x = 3t + 2 \quad \Rightarrow \quad 3t = x - 2 \quad \Rightarrow \quad t = \frac{1}{3}x - \frac{2}{3}$, so

$y = 2t + 3 = 2\left(\frac{1}{3}x - \frac{2}{3}\right) + 3 = \frac{2}{3}x - \frac{4}{3} + 3 \quad \Rightarrow \quad y = \frac{2}{3}x + \frac{5}{3}$

7. $x = t^2 - 3, \quad y = t + 2, \quad -3 \le t \le 3$

(a)

t	-3	-1	1	3
x	6	-2	-2	6
y	-1	1	3	5



(b) $y = t + 2 \quad \Rightarrow \quad t = y - 2$, so

$x = t^2 - 3 = (y - 2)^2 - 3 = y^2 - 4y + 4 - 3 \quad \Rightarrow$

$x = y^2 - 4y + 1, \; -1 \le y \le 5$

8. $x = \sin t, \quad y = 1 - \cos t, \quad 0 \le t \le 2\pi$

(a)

t	0	$\pi/2$	π	$3\pi/2$	2π
x	0	1	0	-1	0
y	0	1	2	1	0



(b) $x = \sin t, y = 1 - \cos t \quad [\text{or } y - 1 = -\cos t] \quad \Rightarrow$

$x^2 + (y - 1)^2 = (\sin t)^2 + (-\cos t)^2 \quad \Rightarrow \quad x^2 + (y - 1)^2 = 1.$

As t varies from 0 to 2π, the circle with center $(0, 1)$ and radius 1 is traced out.

9. $x = \sqrt{t}, \; y = 1 - t$

(a)

t	0	1	2	3	4
x	0	1	1.414	1.732	2
y	1	0	-1	-2	-3



(b) $x = \sqrt{t} \quad \Rightarrow \quad t = x^2 \quad \Rightarrow \quad y = 1 - t = 1 - x^2$. Since $t \ge 0$, $x \ge 0$.

So the curve is the right half of the parabola $y = 1 - x^2$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $x = t^2, y = t^3$

(a)

t	-2	-1	0	1	2
x	4	1	0	1	4
y	-8	-1	0	1	8



(b) $y = t^3 \;\Rightarrow\; t = \sqrt[3]{y} \;\Rightarrow\; x = t^2 = \left(\sqrt[3]{y}\right)^2 = y^{2/3}$. $t \in \mathbb{R}, y \in \mathbb{R}, x \geq 0$.

11. (a) $x = \sin\frac{1}{2}\theta, y = \cos\frac{1}{2}\theta, -\pi \leq \theta \leq \pi$.

$x^2 + y^2 = \sin^2\frac{1}{2}\theta + \cos^2\frac{1}{2}\theta = 1$. For $-\pi \leq \theta \leq 0$, we have $-1 \leq x \leq 0$ and $0 \leq y \leq 1$. For $0 < \theta \leq \pi$, we have $0 < x \leq 1$ and $1 > y \geq 0$. The graph is a semicircle.

(b)



12. (a) $x = \frac{1}{2}\cos\theta, y = 2\sin\theta, 0 \leq \theta \leq \pi$.

$(2x)^2 + \left(\frac{1}{2}y\right)^2 = \cos^2\theta + \sin^2\theta = 1 \;\Rightarrow\; 4x^2 + \frac{1}{4}y^2 = 1 \;\Rightarrow$

$\dfrac{x^2}{(1/2)^2} + \dfrac{y^2}{2^2} = 1$, which is an equation of an ellipse with x-intercepts $\pm\frac{1}{2}$ and y-intercepts ± 2. For $0 \leq \theta \leq \pi/2$, we have $\frac{1}{2} \geq x \geq 0$ and $0 \leq y \leq 2$. For $\pi/2 < \theta \leq \pi$, we have $0 > x \geq -\frac{1}{2}$ and $2 > y \geq 0$. So the graph is the top half of the ellipse.

(b)



13. (a) $x = \sin t, y = \csc t, 0 < t < \frac{\pi}{2}$. $y = \csc t = \dfrac{1}{\sin t} = \dfrac{1}{x}$.

For $0 < t < \frac{\pi}{2}$, we have $0 < x < 1$ and $y > 1$. Thus, the curve is the portion of the hyperbola $y = 1/x$ with $y > 1$.

(b)



14. (a) $y = e^{-2t} = (e^t)^{-2} = x^{-2} = 1/x^2$ for $x > 0$ since $x = e^t$.

(b)



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

15. (a) $y = \ln t \quad \Rightarrow \quad t = e^y$, so $x = t^2 = (e^y)^2 = e^{2y}$.

(b)



16. (a) $x = \sqrt{t+1} \quad \Rightarrow \quad x^2 = t+1 \quad \Rightarrow \quad t = x^2 - 1$.

$y = \sqrt{t-1} = \sqrt{(x^2-1)-1} = \sqrt{x^2-2}$. The curve is the part of the hyperbola $x^2 - y^2 = 2$ with $x \geq \sqrt{2}$ and $y \geq 0$.

(b)



17. (a) $x = \sinh t, y = \cosh t \quad \Rightarrow \quad y^2 - x^2 = \cosh^2 t - \sinh^2 t = 1$.

Since $y = \cosh t \geq 1$, we have the upper branch of the hyperbola $y^2 - x^2 = 1$.

(b)



18. (a) $x = \tan^2\theta, y = \sec\theta, -\pi/2 < \theta < \pi/2$.

$1 + \tan^2\theta = \sec^2\theta \quad \Rightarrow \quad 1 + x = y^2 \quad \Rightarrow \quad x = y^2 - 1$. For $-\pi/2 < \theta \leq 0$, we have $x \geq 0$ and $y \geq 1$. For $0 < \theta < \pi/2$, we have $0 < x$ and $1 < y$. Thus, the curve is the portion of the parabola $x = y^2 - 1$ in the first quadrant. As θ increases from $-\pi/2$ to 0, the point (x, y) approaches $(0, 1)$ along the parabola. As θ increases from 0 to $\pi/2$, the point (x, y) retreats from $(0, 1)$ along the parabola.

(b)



19. $x = 5 + 2\cos\pi t, y = 3 + 2\sin\pi t \quad \Rightarrow \quad \cos\pi t = \dfrac{x-5}{2}, \sin\pi t = \dfrac{y-3}{2}$. $\cos^2(\pi t) + \sin^2(\pi t) = 1 \quad \Rightarrow$

$\left(\dfrac{x-5}{2}\right)^2 + \left(\dfrac{y-3}{2}\right)^2 = 1$. The motion of the particle takes place on a circle centered at $(5, 3)$ with a radius 2. As t goes from 1 to 2, the particle starts at the point $(3, 3)$ and moves counterclockwise along the circle $\left(\dfrac{x-5}{2}\right)^2 + \left(\dfrac{y-3}{2}\right)^2 = 1$ to $(7, 3)$ [one-half of a circle].

20. $x = 2 + \sin t, y = 1 + 3\cos t \quad \Rightarrow \quad \sin t = x - 2, \cos t = \dfrac{y-1}{3}$. $\sin^2 t + \cos^2 t = 1 \quad \Rightarrow \quad (x-2)^2 + \left(\dfrac{y-1}{3}\right)^2 = 1$.

The motion of the particle takes place on an ellipse centered at $(2, 1)$. As t goes from $\pi/2$ to 2π, the particle starts at the point $(3, 1)$ and moves counterclockwise three-fourths of the way around the ellipse to $(2, 4)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $x = 5\sin t, y = 2\cos t \Rightarrow \sin t = \frac{x}{5}, \cos t = \frac{y}{2}$. $\sin^2 t + \cos^2 t = 1 \Rightarrow \left(\frac{x}{5}\right)^2 + \left(\frac{y}{2}\right)^2 = 1$. The motion of the particle takes place on an ellipse centered at $(0, 0)$. As t goes from $-\pi$ to 5π, the particle starts at the point $(0, -2)$ and moves clockwise around the ellipse 3 times.

22. $y = \cos^2 t = 1 - \sin^2 t = 1 - x^2$. The motion of the particle takes place on the parabola $y = 1 - x^2$. As t goes from -2π to $-\pi$, the particle starts at the point $(0, 1)$, moves to $(1, 0)$, and goes back to $(0, 1)$. As t goes from $-\pi$ to 0, the particle moves to $(-1, 0)$ and goes back to $(0, 1)$. The particle repeats this motion as t goes from 0 to 2π.

23. We must have $1 \le x \le 4$ and $2 \le y \le 3$. So the graph of the curve must be contained in the rectangle $[1, 4]$ by $[2, 3]$.

24. (a) From the first graph, we have $1 \le x \le 2$. From the second graph, we have $-1 \le y \le 1$. The only choice that satisfies either of those conditions is III.

(b) From the first graph, the values of x cycle through the values from -2 to 2 four times. From the second graph, the values of y cycle through the values from -2 to 2 six times. Choice I satisfies these conditions.

(c) From the first graph, the values of x cycle through the values from -2 to 2 three times. From the second graph, we have $0 \le y \le 2$. Choice IV satisfies these conditions.

(d) From the first graph, the values of x cycle through the values from -2 to 2 two times. From the second graph, the values of y do the same thing. Choice II satisfies these conditions.

25. When $t = -1$, $(x, y) = (1, 1)$. As t increases to 0, x and y both decrease to 0. As t increases from 0 to 1, x increases from 0 to 1 and y decreases from 0 to -1. As t increases beyond 1, x continues to increase and y continues to decrease. For $t < -1$, x and y are both positive and decreasing. We could achieve greater accuracy by estimating x- and y-values for selected values of t from the given graphs and plotting the corresponding points.



26. When $t = -1$, $(x, y) = (0, 0)$. As t increases to 0, x increases from 0 to 1, while y first decreases to -1 and then increases to 0. As t increases from 0 to 1, x decreases from 1 to 0, while y first increases to 1 and then decreases to 0. We could achieve greater accuracy by estimating x- and y-values for selected values of t from the given graphs and plotting the corresponding points.



27. When $t = -1$, $(x, y) = (0, 1)$. As t increases to 0, x increases from 0 to 1 and y decreases from 1 to 0. As t increases from 0 to 1, the curve is retraced in the opposite direction with x decreasing from 1 to 0 and y increasing from 0 to 1. We could achieve greater accuracy by estimating x- and y-values for selected values of t from the given graphs and plotting the corresponding points.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

28. (a) $x = t^4 - t + 1 = (t^4 + 1) - t > 0$ [think of the graphs of $y = t^4 + 1$ and $y = t$] and $y = t^2 \geq 0$, so these equations are matched with graph V.

(b) $y = \sqrt{t} \geq 0$. $x = t^2 - 2t = t(t-2)$ is negative for $0 < t < 2$, so these equations are matched with graph I.

(c) $x = \sin 2t$ has period $2\pi/2 = \pi$. Note that $y(t + 2\pi) = \sin[t + 2\pi + \sin 2(t + 2\pi)] = \sin(t + 2\pi + \sin 2t) = \sin(t + \sin 2t) = y(t)$, so y has period 2π. These equations match graph II since x cycles through the values -1 to 1 twice as y cycles through those values once.

(d) $x = \cos 5t$ has period $2\pi/5$ and $y = \sin 2t$ has period π, so x will take on the values -1 to 1, and then 1 to -1, before y takes on the values -1 to 1. Note that when $t = 0$, $(x, y) = (1, 0)$. These equations are matched with graph VI.

(e) $x = t + \sin 4t$, $y = t^2 + \cos 3t$. As t becomes large, t and t^2 become the dominant terms in the expressions for x and y, so the graph will look like the graph of $y = x^2$, but with oscillations. These equations are matched with graph IV.

(f) $x = \dfrac{\sin 2t}{4 + t^2}$, $y = \dfrac{\cos 2t}{4 + t^2}$. As $t \to \infty$, x and y both approach 0. These equations are matched with graph III.

29. Use $y = t$ and $x = t - 2\sin \pi t$ with a t-interval of $[-\pi, \pi]$.



30. Use $x_1 = t$, $y_1 = t^3 - 4t$ and $x_2 = t^3 - 4t$, $y_2 = t$ with a t-interval of $[-3, 3]$. There are 9 points of intersection; $(0, 0)$ is fairly obvious. The point in quadrant I is approximately $(2.2, 2.2)$, and by symmetry, the point in quadrant III is approximately $(-2.2, -2.2)$. The other six points are approximately $(\mp 1.9, \pm 0.5)$, $(\mp 1.7, \pm 1.7)$, and $(\mp 0.5, \pm 1.9)$.



31. (a) $x = x_1 + (x_2 - x_1)t$, $y = y_1 + (y_2 - y_1)t$, $0 \leq t \leq 1$. Clearly the curve passes through $P_1(x_1, y_1)$ when $t = 0$ and through $P_2(x_2, y_2)$ when $t = 1$. For $0 < t < 1$, x is strictly between x_1 and x_2 and y is strictly between y_1 and y_2. For every value of t, x and y satisfy the relation $y - y_1 = \dfrac{y_2 - y_1}{x_2 - x_1}(x - x_1)$, which is the equation of the line through $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$.

Finally, any point (x, y) on that line satisfies $\dfrac{y - y_1}{y_2 - y_1} = \dfrac{x - x_1}{x_2 - x_1}$; if we call that common value t, then the given parametric equations yield the point (x, y); and any (x, y) on the line between $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$ yields a value of t in $[0, 1]$. So the given parametric equations exactly specify the line segment from $P_1(x_1, y_1)$ to $P_2(x_2, y_2)$.

(b) $x = -2 + [3 - (-2)]t = -2 + 5t$ and $y = 7 + (-1 - 7)t = 7 - 8t$ for $0 \leq t \leq 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. For the side of the triangle from A to B, use $(x_1, y_1) = (1, 1)$ and $(x_2, y_2) = (4, 2)$. Hence, the equations are



$$x = x_1 + (x_2 - x_1)\,t = 1 + (4 - 1)\,t = 1 + 3t,$$
$$y = y_1 + (y_2 - y_1)\,t = 1 + (2 - 1)\,t = 1 + t.$$

Graphing $x = 1 + 3t$ and $y = 1 + t$ with $0 \le t \le 1$ gives us the side of the triangle from A to B. Similarly, for the side BC we use $x = 4 - 3t$ and $y = 2 + 3t$, and for the side AC we use $x = 1$ and $y = 1 + 4t$.

33. The circle $x^2 + (y - 1)^2 = 4$ has center $(0, 1)$ and radius 2, so by Example 4 it can be represented by $x = 2\cos t$, $y = 1 + 2\sin t$, $0 \le t \le 2\pi$. This representation gives us the circle with a counterclockwise orientation starting at $(2, 1)$.

(a) To get a clockwise orientation, we could change the equations to $x = 2\cos t$, $y = 1 - 2\sin t$, $0 \le t \le 2\pi$.

(b) To get three times around in the counterclockwise direction, we use the original equations $x = 2\cos t$, $y = 1 + 2\sin t$ with the domain expanded to $0 \le t \le 6\pi$.

(c) To start at $(0, 3)$ using the original equations, we must have $x_1 = 0$; that is, $2\cos t = 0$. Hence, $t = \frac{\pi}{2}$. So we use $x = 2\cos t$, $y = 1 + 2\sin t$, $\frac{\pi}{2} \le t \le \frac{3\pi}{2}$.

Alternatively, if we want t to start at 0, we could change the equations of the curve. For example, we could use $x = -2\sin t$, $y = 1 + 2\cos t$, $0 \le t \le \pi$.

34. (a) Let $x^2/a^2 = \sin^2 t$ and $y^2/b^2 = \cos^2 t$ to obtain $x = a\sin t$ and $y = b\cos t$ with $0 \le t \le 2\pi$ as possible parametric equations for the ellipse $x^2/a^2 + y^2/b^2 = 1$.



(b) The equations are $x = 3\sin t$ and $y = b\cos t$ for $b \in \{1, 2, 4, 8\}$.

(c) As b increases, the ellipse stretches vertically.

35. *Big circle:* It's centered at $(2, 2)$ with a radius of 2, so by Example 4, parametric equations are

$$x = 2 + 2\cos t, \qquad y = 2 + 2\sin t, \qquad 0 \le t \le 2\pi$$

Small circles: They are centered at $(1, 3)$ and $(3, 3)$ with a radius of 0.1. By Example 4, parametric equations are

(left) $\quad x = 1 + 0.1\cos t, \quad y = 3 + 0.1\sin t, \quad 0 \le t \le 2\pi$

and *(right)* $\quad x = 3 + 0.1\cos t, \quad y = 3 + 0.1\sin t, \quad 0 \le t \le 2\pi$

Semicircle: It's the lower half of a circle centered at $(2, 2)$ with radius 1. By Example 4, parametric equations are

$$x = 2 + 1\cos t, \qquad y = 2 + 1\sin t, \qquad \pi \le t \le 2\pi$$

To get all four graphs on the same screen with a typical graphing calculator, we need to change the last t-interval to $[0, 2\pi]$ in order to match the others. We can do this by changing t to $0.5t$. This change gives us the upper half. There are several ways to get the lower half—one is to change the "+" to a "−" in the y-assignment, giving us

$$x = 2 + 1\cos(0.5t), \qquad y = 2 - 1\sin(0.5t), \qquad 0 \le t \le 2\pi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

36. If you are using a calculator or computer that can overlay graphs (using multiple t-intervals), the following is appropriate.

Left side: $x = 1$ and y goes from 1.5 to 4, so use

$$x = 1, \qquad y = t, \qquad 1.5 \le t \le 4$$

Right side: $x = 10$ and y goes from 1.5 to 4, so use

$$x = 10, \qquad y = t, \qquad 1.5 \le t \le 4$$

Bottom: x goes from 1 to 10 and $y = 1.5$, so use

$$x = t, \qquad y = 1.5, \qquad 1 \le t \le 10$$

Handle: It starts at $(10, 4)$ and ends at $(13, 7)$, so use

$$x = 10 + t, \qquad y = 4 + t, \qquad 0 \le t \le 3$$

Left wheel: It's centered at $(3, 1)$, has a radius of 1, and appears to go about $30°$ above the horizontal, so use

$$x = 3 + 1\cos t, \qquad y = 1 + 1\sin t, \qquad \tfrac{5\pi}{6} \le t \le \tfrac{13\pi}{6}$$

Right wheel: Similar to the left wheel with center $(8, 1)$, so use

$$x = 8 + 1\cos t, \qquad y = 1 + 1\sin t, \qquad \tfrac{5\pi}{6} \le t \le \tfrac{13\pi}{6}$$

If you are using a calculator or computer that cannot overlay graphs (using one t-interval), the following is appropriate. We'll start by picking the t-interval $[0, 2.5]$ since it easily matches the t-values for the two sides. We now need to find parametric equations for all graphs with $0 \le t \le 2.5$.

Left side: $x = 1$ and y goes from 1.5 to 4, so use

$$x = 1, \qquad y = 1.5 + t, \qquad 0 \le t \le 2.5$$

Right side: $x = 10$ and y goes from 1.5 to 4, so use

$$x = 10, \qquad y = 1.5 + t, \qquad 0 \le t \le 2.5$$

Bottom: x goes from 1 to 10 and $y = 1.5$, so use

$$x = 1 + 3.6t, \qquad y = 1.5, \qquad 0 \le t \le 2.5$$

To get the x-assignment, think of creating a linear function such that when $t = 0$, $x = 1$ and when $t = 2.5$, $x = 10$. We can use the point-slope form of a line with $(t_1, x_1) = (0, 1)$ and $(t_2, x_2) = (2.5, 10)$.

$x - 1 = \dfrac{10 - 1}{2.5 - 0}(t - 0) \quad \Rightarrow \quad x = 1 + 3.6t.$

Handle: It starts at $(10, 4)$ and ends at $(13, 7)$, so use

$$x = 10 + 1.2t, \qquad y = 4 + 1.2t, \qquad 0 \le t \le 2.5$$

$(t_1, x_1) = (0, 10)$ and $(t_2, x_2) = (2.5, 13)$ gives us $x - 10 = \dfrac{13 - 10}{2.5 - 0}(t - 0) \quad \Rightarrow \quad x = 10 + 1.2t.$

$(t_1, y_1) = (0, 4)$ and $(t_2, y_2) = (2.5, 7)$ gives us $y - 4 = \dfrac{7 - 4}{2.5 - 0}(t - 0) \quad \Rightarrow \quad y = 4 + 1.2t.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Left wheel: It's centered at $(3, 1)$, has a radius of 1, and appears to go about $30°$ above the horizontal, so use

$$x = 3 + 1\cos\left(\tfrac{8\pi}{15}t + \tfrac{5\pi}{6}\right), \qquad y = 1 + 1\sin\left(\tfrac{8\pi}{15}t + \tfrac{5\pi}{6}\right), \qquad 0 \le t \le 2.5$$

$(t_1, \theta_1) = \left(0, \tfrac{5\pi}{6}\right)$ and $(t_2, \theta_2) = \left(\tfrac{5}{2}, \tfrac{13\pi}{6}\right)$ gives us $\theta - \tfrac{5\pi}{6} = \dfrac{\frac{13\pi}{6} - \frac{5\pi}{6}}{\frac{5}{2} - 0}(t - 0) \quad \Rightarrow \quad \theta = \tfrac{5\pi}{6} + \tfrac{8\pi}{15}t.$

Right wheel: Similar to the left wheel with center $(8, 1)$, so use

$$x = 8 + 1\cos\left(\tfrac{8\pi}{15}t + \tfrac{5\pi}{6}\right), \qquad y = 1 + 1\sin\left(\tfrac{8\pi}{15}t + \tfrac{5\pi}{6}\right), \qquad 0 \le t \le 2.5$$

37. (a) $x = t^3 \quad \Rightarrow \quad t = x^{1/3}$, so $y = t^2 = x^{2/3}$.

We get the entire curve $y = x^{2/3}$ traversed in a left to right direction.



(b) $x = t^6 \quad \Rightarrow \quad t = x^{1/6}$, so $y = t^4 = x^{4/6} = x^{2/3}$.

Since $x = t^6 \ge 0$, we only get the right half of the curve $y = x^{2/3}$.



(c) $x = e^{-3t} = (e^{-t})^3 \quad [\text{so } e^{-t} = x^{1/3}]$,
$y = e^{-2t} = (e^{-t})^2 = (x^{1/3})^2 = x^{2/3}$.
If $t < 0$, then x and y are both larger than 1. If $t > 0$, then x and y are between 0 and 1. Since $x > 0$ and $y > 0$, the curve never quite reaches the origin.



38. (a) $x = t$, so $y = t^{-2} = x^{-2}$. We get the entire curve $y = 1/x^2$ traversed in a left-to-right direction.



(b) $x = \cos t$, $y = \sec^2 t = \dfrac{1}{\cos^2 t} = \dfrac{1}{x^2}$. Since $\sec t \ge 1$, we only get the parts of the curve $y = 1/x^2$ with $y \ge 1$. We get the first quadrant portion of the curve when $x > 0$, that is, $\cos t > 0$, and we get the second quadrant portion of the curve when $x < 0$, that is, $\cos t < 0$.



(c) $x = e^t$, $y = e^{-2t} = (e^t)^{-2} = x^{-2}$. Since e^t and e^{-2t} are both positive, we only get the first quadrant portion of the curve $y = 1/x^2$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. The case $\frac{\pi}{2} < \theta < \pi$ is illustrated. C has coordinates $(r\theta, r)$ as in Example 7, and Q has coordinates $(r\theta, r + r\cos(\pi - \theta)) = (r\theta, r(1 - \cos\theta))$ [since $\cos(\pi - \alpha) = \cos\pi\cos\alpha + \sin\pi\sin\alpha = -\cos\alpha$], so P has coordinates $(r\theta - r\sin(\pi - \theta), r(1 - \cos\theta)) = (r(\theta - \sin\theta), r(1 - \cos\theta))$ [since $\sin(\pi - \alpha) = \sin\pi\cos\alpha - \cos\pi\sin\alpha = \sin\alpha$]. Again we have the parametric equations $x = r(\theta - \sin\theta)$, $y = r(1 - \cos\theta)$.



40. The first two diagrams depict the case $\pi < \theta < \frac{3\pi}{2}$, $d < r$. As in Example 7, C has coordinates $(r\theta, r)$. Now Q (in the second diagram) has coordinates $(r\theta, r + d\cos(\theta - \pi)) = (r\theta, r - d\cos\theta)$, so a typical point P of the trochoid has coordinates $(r\theta + d\sin(\theta - \pi), r - d\cos\theta)$. That is, P has coordinates (x, y), where $x = r\theta - d\sin\theta$ and $y = r - d\cos\theta$. When $d = r$, these equations agree with those of the cycloid.







41. It is apparent that $x = |OQ|$ and $y = |QP| = |ST|$. From the diagram, $x = |OQ| = a\cos\theta$ and $y = |ST| = b\sin\theta$. Thus, the parametric equations are $x = a\cos\theta$ and $y = b\sin\theta$. To eliminate θ we rearrange: $\sin\theta = y/b \;\Rightarrow\; \sin^2\theta = (y/b)^2$ and $\cos\theta = x/a \;\Rightarrow\; \cos^2\theta = (x/a)^2$. Adding the two equations: $\sin^2\theta + \cos^2\theta = 1 = x^2/a^2 + y^2/b^2$. Thus, we have an ellipse.



42. A has coordinates $(a\cos\theta, a\sin\theta)$. Since OA is perpendicular to AB, ΔOAB is a right triangle and B has coordinates $(a\sec\theta, 0)$. It follows that P has coordinates $(a\sec\theta, b\sin\theta)$. Thus, the parametric equations are $x = a\sec\theta$, $y = b\sin\theta$.

43. $C = (2a\cot\theta, 2a)$, so the x-coordinate of P is $x = 2a\cot\theta$. Let $B = (0, 2a)$. Then $\angle OAB$ is a right angle and $\angle OBA = \theta$, so $|OA| = 2a\sin\theta$ and $A = ((2a\sin\theta)\cos\theta, (2a\sin\theta)\sin\theta)$. Thus, the y-coordinate of P is $y = 2a\sin^2\theta$.



44. (a) Let θ be the angle of inclination of segment OP. Then $|OB| = \dfrac{2a}{\cos\theta}$.

Let $C = (2a, 0)$. Then by use of right triangle OAC we see that $|OA| = 2a\cos\theta$.

Now

$$|OP| = |AB| = |OB| - |OA|$$

$$= 2a\left(\frac{1}{\cos\theta} - \cos\theta\right) = 2a\,\frac{1 - \cos^2\theta}{\cos\theta} = 2a\,\frac{\sin^2\theta}{\cos\theta} = 2a\sin\theta\,\tan\theta$$

So P has coordinates $x = 2a\sin\theta\,\tan\theta\cdot\cos\theta = 2a\sin^2\theta$ and $y = 2a\sin\theta\,\tan\theta\cdot\sin\theta = 2a\sin^2\theta\,\tan\theta$.

(b)



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. (a)



There are 2 points of intersection:

$(-3, 0)$ and approximately $(-2.1, 1.4)$.

(b) A collision point occurs when $x_1 = x_2$ and $y_1 = y_2$ for the same t. So solve the equations:

$$3\sin t = -3 + \cos t \quad \textbf{(1)}$$

$$2\cos t = 1 + \sin t \quad \textbf{(2)}$$

From **(2)**, $\sin t = 2\cos t - 1$. Substituting into **(1)**, we get $3(2\cos t - 1) = -3 + \cos t \;\Rightarrow\; 5\cos t = 0 \;\; (\star) \;\Rightarrow$ $\cos t = 0 \;\Rightarrow\; t = \frac{\pi}{2}$ or $\frac{3\pi}{2}$. We check that $t = \frac{3\pi}{2}$ satisfies **(1)** and **(2)** but $t = \frac{\pi}{2}$ does not. So the only collision point occurs when $t = \frac{3\pi}{2}$, and this gives the point $(-3, 0)$. [We could check our work by graphing x_1 and x_2 together as functions of t and, on another plot, y_1 and y_2 as functions of t. If we do so, we see that the only value of t for which *both* pairs of graphs intersect is $t = \frac{3\pi}{2}$.]

(c) The circle is centered at $(3, 1)$ instead of $(-3, 1)$. There are still 2 intersection points: $(3, 0)$ and $(2.1, 1.4)$, but there are no collision points, since $(\star)$ in part (b) becomes $5\cos t = 6 \;\Rightarrow\; \cos t = \frac{6}{5} > 1$.

46. (a) If $\alpha = 30°$ and $v_0 = 500$ m/s, then the equations become $x = (500\cos 30°)t = 250\sqrt{3}t$ and $y = (500\sin 30°)t - \frac{1}{2}(9.8)t^2 = 250t - 4.9t^2$. $y = 0$ when $t = 0$ (when the gun is fired) and again when $t = \frac{250}{4.9} \approx 51$ s. Then $x = \left(250\sqrt{3}\right)\left(\frac{250}{4.9}\right) \approx 22{,}092$ m, so the bullet hits the ground about 22 km from the gun.

The formula for y is quadratic in t. To find the maximum y-value, we will complete the square:

$$y = -4.9\left(t^2 - \tfrac{250}{4.9}t\right) = -4.9\left[t^2 - \tfrac{250}{4.9}t + \left(\tfrac{125}{4.9}\right)^2\right] + \tfrac{125^2}{4.9} = -4.9\left(t - \tfrac{125}{4.9}\right)^2 + \tfrac{125^2}{4.9} \le \tfrac{125^2}{4.9}$$

with equality when $t = \frac{125}{4.9}$ s, so the maximum height attained is $\frac{125^2}{4.9} \approx 3189$ m.

(b)



As α $(0° < \alpha < 90°)$ increases up to $45°$, the projectile attains a greater height and a greater range. As α increases past $45°$, the projectile attains a greater height, but its range decreases.

(c) $x = (v_0\cos\alpha)t \;\Rightarrow\; t = \dfrac{x}{v_0\cos\alpha}$.

$$y = (v_0\sin\alpha)t - \tfrac{1}{2}gt^2 \;\Rightarrow\; y = (v_0\sin\alpha)\frac{x}{v_0\cos\alpha} - \frac{g}{2}\left(\frac{x}{v_0\cos\alpha}\right)^2 = (\tan\alpha)x - \left(\frac{g}{2v_0^2\cos^2\alpha}\right)x^2,$$

which is the equation of a parabola (quadratic in x).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

47. $x = t^2, y = t^3 - ct$. We use a graphing device to produce the graphs for various values of c with $-\pi \le t \le \pi$. Note that all the members of the family are symmetric about the x-axis. For $c < 0$, the graph does not cross itself, but for $c = 0$ it has a cusp at $(0, 0)$ and for $c > 0$ the graph crosses itself at $x = c$, so the loop grows larger as c increases.





48. $x = 2ct - 4t^3, y = -ct^2 + 3t^4$. We use a graphing device to produce the graphs for various values of c with $-\pi \le t \le \pi$. Note that all the members of the family are symmetric about the y-axis. When $c < 0$, the graph resembles that of a polynomial of even degree, but when $c = 0$ there is a corner at the origin, and when $c > 0$, the graph crosses itself at the origin, and has two cusps below the x-axis. The size of the "swallowtail" increases as c increases.



49. $x = t + a\cos t, y = t + a\sin t, a > 0$. From the first figure, we see that curves roughly follow the line $y = x$, and they start having loops when a is between 1.4 and 1.6. The loops increase in size as a increases.



While not required, the following is a solution to determine the *exact* values for which the curve has a loop, that is, we seek the values of a for which there exist parameter values t and u such that $t < u$ and $(t + a\cos t, t + a\sin t) = (u + a\cos u, u + a\sin u)$.



In the diagram at the left, T denotes the point (t, t), U the point (u, u), and P the point $(t + a\cos t, t + a\sin t) = (u + a\cos u, u + a\sin u)$. Since $\overline{PT} = \overline{PU} = a$, the triangle PTU is isosceles. Therefore its base angles, $\alpha = \angle PTU$ and $\beta = \angle PUT$ are equal. Since $\alpha = t - \frac{\pi}{4}$ and $\beta = 2\pi - \frac{3\pi}{4} - u = \frac{5\pi}{4} - u$, the relation $\alpha = \beta$ implies that $u + t = \frac{3\pi}{2}$ **(1)**.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Since $\overline{TU} = \text{distance}((t,t),(u,u)) = \sqrt{2(u-t)^2} = \sqrt{2}\,(u-t)$, we see that $\cos\alpha = \dfrac{\frac{1}{2}\overline{TU}}{\overline{PT}} = \dfrac{(u-t)/\sqrt{2}}{a}$, so $u - t = \sqrt{2}\,a\cos\alpha$, that is, $u - t = \sqrt{2}\,a\cos\left(t - \frac{\pi}{4}\right)$ **(2)**. Now $\cos\left(t - \frac{\pi}{4}\right) = \sin\left[\frac{\pi}{2} - \left(t - \frac{\pi}{4}\right)\right] = \sin\left(\frac{3\pi}{4} - t\right)$, so we can rewrite **(2)** as $u - t = \sqrt{2}\,a\sin\left(\frac{3\pi}{4} - t\right)$ **(2′)**. Subtracting **(2′)** from **(1)** and dividing by 2, we obtain $t = \frac{3\pi}{4} - \frac{\sqrt{2}}{2}a\sin\left(\frac{3\pi}{4} - t\right)$, or $\frac{3\pi}{4} - t = \frac{a}{\sqrt{2}}\sin\left(\frac{3\pi}{4} - t\right)$ **(3)**.



Since $a > 0$ and $t < u$, it follows from **(2′)** that $\sin\left(\frac{3\pi}{4} - t\right) > 0$. Thus from **(3)** we see that $t < \frac{3\pi}{4}$. [We have implicitly assumed that $0 < t < \pi$ by the way we drew our diagram, but we lost no generality by doing so since replacing t by $t + 2\pi$ merely increases x and y by 2π. The curve's basic shape repeats every time we change t by 2π.] Solving for a in **(3)**, we get $a = \dfrac{\sqrt{2}\left(\frac{3\pi}{4} - t\right)}{\sin\left(\frac{3\pi}{4} - t\right)}$. Write $z = \frac{3\pi}{4} - t$. Then $a = \dfrac{\sqrt{2}\,z}{\sin z}$, where $z > 0$. Now $\sin z < z$ for $z > 0$, so $a > \sqrt{2}$. $\left[\text{As } z \to 0^+, \text{ that is, as } t \to \left(\frac{3\pi}{4}\right)^-,\ a \to \sqrt{2}\right]$.

50. Consider the curves $x = \sin t + \sin nt$, $y = \cos t + \cos nt$, where n is a positive integer. For $n = 1$, we get a circle of radius 2 centered at the origin. For $n > 1$, we get a curve lying on or inside that circle that traces out $n - 1$ loops as t ranges from 0 to 2π.

Note:
$$\begin{aligned} x^2 + y^2 &= (\sin t + \sin nt)^2 + (\cos t + \cos nt)^2 \\ &= \sin^2 t + 2\sin t\,\sin nt + \sin^2 nt + \cos^2 t + 2\cos t\,\cos nt + \cos^2 nt \\ &= (\sin^2 t + \cos^2 t) + (\sin^2 nt + \cos^2 nt) + 2(\cos t\,\cos nt + \sin t\,\sin nt) \\ &= 1 + 1 + 2\cos(t - nt) = 2 + 2\cos((1-n)t) \le 4 = 2^2, \end{aligned}$$

with equality for $n = 1$. This shows that each curve lies on or inside the curve for $n = 1$, which is a circle of radius 2 centered at the origin.



$n = 1$



$n = 2$



$n = 3$



$n = 5$

51. Note that all the Lissajous figures are symmetric about the x-axis. The parameters a and b simply stretch the graph in the x- and y-directions respectively. For $a = b = n = 1$ the graph is simply a circle with radius 1. For $n = 2$ the graph crosses

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

itself at the origin and there are loops above and below the x-axis. In general, the figures have $n-1$ points of intersection, all of which are on the y-axis, and a total of n closed loops.



$a = b = 1$



$n = 2$



$n = 3$

52. $x = \cos t,\ y = \sin t - \sin ct$. If $c = 1$, then $y = 0$, and the curve is simply the line segment from $(-1, 0)$ to $(1, 0)$. The graphs are shown for $c = 2, 3, 4$ and 5.









It is easy to see that all the curves lie in the rectangle $[-1, 1]$ by $[-2, 2]$. When c is an integer, $x(t + 2\pi) = x(t)$ and $y(t + 2\pi) = y(t)$, so the curve is closed. When c is a positive integer greater than 1, the curve intersects the x-axis $c + 1$ times and has c loops (one of which degenerates to a tangency at the origin when c is an odd integer of the form $4k + 1$).

As c increases, the curve's loops become thinner, but stay in the region bounded by the semicircles $y = \pm(1 + \sqrt{1 - x^2})$ and the line segments from $(-1, -1)$ to $(-1, 1)$ and from $(1, -1)$ to $(1, 1)$. This is true because $|y| = |\sin t - \sin ct| \le |\sin t| + |\sin ct| \le \sqrt{1 - x^2} + 1$. This curve appears to fill the entire region when c is very large, as shown in the figure for $c = 1000$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

When c is a fraction, we get a variety of shapes with multiple loops, but always within the same region. For some fractional values, such as $c = 2.359$, the curve again appears to fill the region.



LABORATORY PROJECT Running Circles Around Circles

1. The center Q of the smaller circle has coordinates $((a-b)\cos\theta, (a-b)\sin\theta)$. Arc PS on circle C has length $a\theta$ since it is equal in length to arc AS (the smaller circle rolls without slipping against the larger.)



Thus, $\angle PQS = \dfrac{a}{b}\theta$ and $\angle PQT = \dfrac{a}{b}\theta - \theta$, so P has coordinates

$$x = (a-b)\cos\theta + b\cos(\angle PQT) = (a-b)\cos\theta + b\cos\left(\frac{a-b}{b}\theta\right)$$

and $$y = (a-b)\sin\theta - b\sin(\angle PQT) = (a-b)\sin\theta - b\sin\left(\frac{a-b}{b}\theta\right).$$

2. With $b = 1$ and a a positive integer greater than 2, we obtain a hypocycloid of a cusps. Shown in the figure is the graph for $a = 4$. Let $a = 4$ and $b = 1$. Using the sum identities to expand $\cos 3\theta$ and $\sin 3\theta$, we obtain



$$x = 3\cos\theta + \cos 3\theta = 3\cos\theta + \left(4\cos^3\theta - 3\cos\theta\right) = 4\cos^3\theta$$

and $$y = 3\sin\theta - \sin 3\theta = 3\sin\theta - \left(3\sin\theta - 4\sin^3\theta\right) = 4\sin^3\theta.$$

3. The graphs at the right are obtained with $b = 1$ and $a = \frac{1}{2}, \frac{1}{3}, \frac{1}{4}$, and $\frac{1}{10}$ with $-2\pi \le \theta \le 2\pi$. We conclude that as the denominator d increases, the graph gets smaller, but maintains the basic shape shown.

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Letting $d = 2$ and $n = 3$, 5, and 7 with $-2\pi \le \theta \le 2\pi$ gives us the following:



So if d is held constant and n varies, we get a graph with n cusps (assuming n/d is in lowest form). When $n = d + 1$, we obtain a hypocycloid of n cusps. As n increases, we must expand the range of θ in order to get a closed curve. The following graphs have $a = \frac{3}{2}$, $\frac{5}{4}$, and $\frac{11}{10}$.



4. If $b = 1$, the equations for the hypocycloid are

$$x = (a-1)\cos\theta + \cos((a-1)\theta) \qquad y = (a-1)\sin\theta - \sin((a-1)\theta)$$

which is a hypocycloid of a cusps (from Problem 2). In general, if $a > 1$, we get a figure with cusps on the "outside ring" and if $a < 1$, the cusps are on the "inside ring". In any case, as the values of θ get larger, we get a figure that looks more and more like a washer. If we were to graph the hypocycloid for all values of θ, every point on the washer would eventually be arbitrarily close to a point on the curve.



$a = \sqrt{2}, \quad -10\pi \le \theta \le 10\pi$

$a = e - 2, \quad 0 \le \theta \le 446$

5. The center Q of the smaller circle has coordinates $((a+b)\cos\theta, (a+b)\sin\theta)$.



Arc PS has length $a\theta$ (as in Problem 1), so that $\angle PQS = \dfrac{a\theta}{b}$, $\angle PQR = \pi - \dfrac{a\theta}{b}$,

and $\angle PQT = \pi - \dfrac{a\theta}{b} - \theta = \pi - \left(\dfrac{a+b}{b}\right)\theta$ since $\angle RQT = \theta$.

Thus, the coordinates of P are

$$x = (a+b)\cos\theta + b\cos\left(\pi - \frac{a+b}{b}\theta\right) = (a+b)\cos\theta - b\cos\left(\frac{a+b}{b}\theta\right)$$

and $$y = (a+b)\sin\theta - b\sin\left(\pi - \frac{a+b}{b}\theta\right) = (a+b)\sin\theta - b\sin\left(\frac{a+b}{b}\theta\right).$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. Let $b = 1$ and the equations become

$$x = (a+1)\cos\theta - \cos((a+1)\theta) \qquad y = (a+1)\sin\theta - \sin((a+1)\theta)$$

If $a = 1$, we have a cardioid. If a is a positive integer greater than 1, we get the graph of an "a-leafed clover", with cusps that are a units from the origin. (Some of the pairs of figures are not to scale.)



$a = 3, -2\pi \le \theta \le 2\pi$



$a = 10, -2\pi \le \theta \le 2\pi$

If $a = n/d$ with $n = 1$, we obtain a figure that does not increase in size and requires $-d\pi \le \theta \le d\pi$ to be a closed curve traced exactly once.

$a = \frac{1}{4}, -4\pi \le \theta \le 4\pi$

$a = \frac{1}{7}, -7\pi \le \theta \le 7\pi$

Next, we keep d constant and let n vary. As n increases, so does the size of the figure. There is an n-pointed star in the middle.



$a = \frac{2}{5}, -5\pi \le \theta \le 5\pi$



$a = \frac{7}{5}, -5\pi \le \theta \le 5\pi$

Now if $n = d + 1$ we obtain figures similar to the previous ones, but the size of the figure does not increase.



$a = \frac{4}{3}, -3\pi \le \theta \le 3\pi$



$a = \frac{7}{6}, -6\pi \le \theta \le 6\pi$

If a is irrational, we get washers that increase in size as a increases.



$a = \sqrt{2}, 0 \le \theta \le 200$

$a = e - 2, 0 \le \theta \le 446$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

10.2 Calculus with Parametric Curves

1. $x = \dfrac{t}{1+t},\ y = \sqrt{1+t} \ \Rightarrow\ \dfrac{dy}{dt} = \dfrac{1}{2}(1+t)^{-1/2} = \dfrac{1}{2\sqrt{1+t}},\ \dfrac{dx}{dt} = \dfrac{(1+t)(1)-t(1)}{(1+t)^2} = \dfrac{1}{(1+t)^2}$, and

$\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{1/(2\sqrt{1+t})}{1/(1+t)^2} = \dfrac{(1+t)^2}{2\sqrt{1+t}} = \dfrac{1}{2}(1+t)^{3/2}$.

2. $x = te^t,\ y = t + \sin t \ \Rightarrow\ \dfrac{dy}{dt} = 1 + \cos t,\ \dfrac{dx}{dt} = te^t + e^t = e^t(t+1)$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{1+\cos t}{e^t(t+1)}$.

3. $x = t^3 + 1,\ y = t^4 + t;\ \ t = -1.\ \dfrac{dy}{dt} = 4t^3 + 1,\ \dfrac{dx}{dt} = 3t^2$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{4t^3+1}{3t^2}$. When $t = -1$, $(x, y) = (0, 0)$ and $dy/dx = -3/3 = -1$, so an equation of the tangent to the curve at the point corresponding to $t = -1$ is $y - 0 = -1(x - 0)$, or $y = -x$.

4. $x = \sqrt{t},\ y = t^2 - 2t;\ \ t = 4.\ \dfrac{dy}{dt} = 2t - 2,\ \dfrac{dx}{dt} = \dfrac{1}{2\sqrt{t}}$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = (2t-2)2\sqrt{t} = 4(t-1)\sqrt{t}$. When $t = 4$, $(x, y) = (2, 8)$ and $dy/dx = 4(3)(2) = 24$, so an equation of the tangent to the curve at the point corresponding to $t = 4$ is $y - 8 = 24(x - 2)$, or $y = 24x - 40$.

5. $x = t\cos t,\ y = t\sin t;\ t = \pi.\ \ \dfrac{dy}{dt} = t\cos t + \sin t,\ \dfrac{dx}{dt} = t(-\sin t) + \cos t$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{t\cos t + \sin t}{-t\sin t + \cos t}$.

When $t = \pi$, $(x, y) = (-\pi, 0)$ and $dy/dx = -\pi/(-1) = \pi$, so an equation of the tangent to the curve at the point corresponding to $t = \pi$ is $y - 0 = \pi[x - (-\pi)]$, or $y = \pi x + \pi^2$.

6. $x = e^t \sin \pi t,\ y = e^{2t};\ \ t = 0.\ \dfrac{dy}{dt} = 2e^{2t},\ \dfrac{dx}{dt} = e^t(\pi\cos \pi t) + (\sin \pi t)e^t = e^t(\pi\cos \pi t + \sin \pi t)$, and

$\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2e^{2t}}{e^t(\pi\cos \pi t + \sin \pi t)} = \dfrac{2e^t}{\pi\cos \pi t + \sin \pi t}$. When $t = 0$, $(x, y) = (0, 1)$ and $dy/dx = 2/\pi$, so an equation of the tangent to the curve at the point corresponding to $t = 0$ is $y - 1 = \frac{2}{\pi}(x - 0)$, or $y = \frac{2}{\pi}x + 1$.

7. (a) $x = 1 + \ln t,\ y = t^2 + 2;\ (1, 3).\ \ \dfrac{dy}{dt} = 2t,\ \dfrac{dx}{dt} = \dfrac{1}{t}$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t}{1/t} = 2t^2$. At $(1, 3)$,

$x = 1 + \ln t = 1 \ \Rightarrow\ \ln t = 0 \ \Rightarrow\ t = 1$ and $\dfrac{dy}{dx} = 2$, so an equation of the tangent is $y - 3 = 2(x - 1)$, or $y = 2x + 1$.

(b) $x = 1 + \ln t \ \Rightarrow\ \ln t = x - 1 \ \Rightarrow\ t = e^{x-1}$, so $y = t^2 + 2 = (e^{x-1})^2 + 2 = e^{2x-2} + 2$, and $y' = e^{2x-2} \cdot 2$. At $(1, 3)$, $y' = e^{2(1)-2} \cdot 2 = 2$, so an equation of the tangent is $y - 3 = 2(x - 1)$, or $y = 2x + 1$.

8. (a) $x = 1 + \sqrt{t},\ y = e^{t^2};\ (2, e).\ \ \dfrac{dy}{dt} = e^{t^2} \cdot 2t,\ \dfrac{dx}{dt} = \dfrac{1}{2\sqrt{t}}$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2te^{t^2}}{1/(2\sqrt{t})} = 4t^{3/2}e^{t^2}$. At $(2, e)$,

$x = 1 + \sqrt{t} = 2 \ \Rightarrow\ \sqrt{t} = 1 \ \Rightarrow\ t = 1$ and $\dfrac{dy}{dx} = 4e$, so an equation of the tangent is $y - e = 4e(x - 2)$, or $y = 4ex - 7e$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $x = 1 + \sqrt{t} \;\Rightarrow\; \sqrt{t} = x - 1 \;\Rightarrow\; t = (x-1)^2$, so $y = e^{t^2} = e^{(x-1)^4}$, and $y' = e^{(x-1)^4} \cdot 4(x-1)^3$.

At $(2, e)$, $y' = e \cdot 4 = 4e$, so an equation of the tangent is $y - e = 4e(x-2)$, or $y = 4ex - 7e$.

9. $x = t^2 - t,\ y = t^2 + t + 1;\ (0, 3)$. $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t+1}{2t-1}$. To find the value of t corresponding to the point $(0, 3)$, solve $x = 0 \;\Rightarrow$ $t^2 - t = 0 \;\Rightarrow\; t(t-1) = 0 \;\Rightarrow\; t = 0$ or $t = 1$. Only $t = 1$ gives $y = 3$. With $t = 1$, $dy/dx = 3$, and an equation of the tangent is $y - 3 = 3(x - 0)$, or $y = 3x + 3$.



10. $x = \sin \pi t,\ y = t^2 + t;\ (0, 2)$. $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t+1}{\pi \cos \pi t}$. To find the value of t corresponding to the point $(0, 2)$, solve $y = 2 \;\Rightarrow$ $t^2 + t - 2 = 0 \;\Rightarrow\; (t+2)(t-1) = 0 \;\Rightarrow\; t = -2$ or $t = 1$. Either value gives $dy/dx = -3/\pi$, so an equation of the tangent is $y - 2 = -\frac{3}{\pi}(x - 0)$, or $y = -\frac{3}{\pi}x + 2$.



11. $x = t^2 + 1,\ y = t^2 + t \;\Rightarrow\; \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t+1}{2t} = 1 + \dfrac{1}{2t} \;\Rightarrow\; \dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{-1/(2t^2)}{2t} = -\dfrac{1}{4t^3}$.

The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $t < 0$.

12. $x = t^3 + 1,\ y = t^2 - t \;\Rightarrow\; \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t-1}{3t^2} = \dfrac{2}{3t} - \dfrac{1}{3t^2} \;\Rightarrow$

$\dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{-\dfrac{2}{3t^2} + \dfrac{2}{3t^3}}{3t^2} = \dfrac{\dfrac{2-2t}{3t^3}}{3t^2} = \dfrac{2(1-t)}{9t^5}$. The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $0 < t < 1$.

13. $x = e^t,\ y = te^{-t} \;\Rightarrow\; \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{-te^{-t} + e^{-t}}{e^t} = \dfrac{e^{-t}(1-t)}{e^t} = e^{-2t}(1-t) \;\Rightarrow$

$\dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{e^{-2t}(-1) + (1-t)(-2e^{-2t})}{e^t} = \dfrac{e^{-2t}(-1-2+2t)}{e^t} = e^{-3t}(2t-3)$. The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $t > \frac{3}{2}$.

14. $x = t^2 + 1,\ y = e^t - 1 \;\Rightarrow\; \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{e^t}{2t} \;\Rightarrow\; \dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{\dfrac{2te^t - e^t \cdot 2}{(2t)^2}}{2t} = \dfrac{2e^t(t-1)}{(2t)^3} = \dfrac{e^t(t-1)}{4t^3}$.

The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $t < 0$ or $t > 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

15. $x = t - \ln t,\ y = t + \ln t$ [note that $t > 0$] $\Rightarrow$ $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{1+1/t}{1-1/t} = \dfrac{t+1}{t-1}$ $\Rightarrow$

$\dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{\dfrac{(t-1)(1)-(t+1)(1)}{(t-1)^2}}{(t-1)/t} = \dfrac{-2t}{(t-1)^3}$. The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $0 < t < 1$.

16. $x = \cos t,\ y = \sin 2t,\ 0 < t < \pi$ $\Rightarrow$ $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2\cos 2t}{-\sin t}$ $\Rightarrow$

$$\frac{d^2y}{dx^2} = \frac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \frac{\dfrac{(-\sin t)(-4\sin 2t) - (2\cos 2t)(-\cos t)}{(-\sin t)^2}}{-\sin t} = \frac{(\sin t)(8\sin t\,\cos t) + [2(1-2\sin^2 t)](\cos t)}{(-\sin t)\sin^2 t}$$

$$= \frac{(\cos t)(8\sin^2 t + 2 - 4\sin^2 t)}{(-\sin t)\sin^2 t} = -\frac{\cos t}{\sin t}\cdot\frac{4\sin^2 t + 2}{\sin^2 t} \quad [\,(-\cot t)\cdot \text{positive expression}]$$

The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $-\cot t > 0 \iff \cot t < 0 \iff \frac{\pi}{2} < t < \pi$.

17. $x = t^3 - 3t,\ y = t^2 - 3$. $\dfrac{dy}{dt} = 2t$, so $\dfrac{dy}{dt} = 0 \iff t = 0 \iff$ $(x, y) = (0, -3)$. $\dfrac{dx}{dt} = 3t^2 - 3 = 3(t+1)(t-1)$, so $\dfrac{dx}{dt} = 0 \iff$ $t = -1$ or $1 \iff (x, y) = (2, -2)$ or $(-2, -2)$. The curve has a horizontal tangent at $(0, -3)$ and vertical tangents at $(2, -2)$ and $(-2, -2)$.



18. $x = t^3 - 3t,\ y = t^3 - 3t^2$. $\dfrac{dy}{dt} = 3t^2 - 6t = 3t(t-2)$, so $\dfrac{dy}{dt} = 0 \iff$ $t = 0$ or $2 \iff (x, y) = (0, 0)$ or $(2, -4)$. $\dfrac{dx}{dt} = 3t^2 - 3 = 3(t+1)(t-1)$, so $\dfrac{dx}{dt} = 0 \iff t = -1$ or $1 \iff (x, y) = (2, -4)$ or $(-2, -2)$. The curve has horizontal tangents at $(0, 0)$ and $(2, -4)$, and vertical tangents at $(2, -4)$ and $(-2, -2)$.



19. $x = \cos\theta,\ y = \cos 3\theta$. The whole curve is traced out for $0 \le \theta \le \pi$.

$\dfrac{dy}{d\theta} = -3\sin 3\theta$, so $\dfrac{dy}{d\theta} = 0 \iff \sin 3\theta = 0 \iff 3\theta = 0, \pi, 2\pi$, or $3\pi \iff$

$\theta = 0, \frac{\pi}{3}, \frac{2\pi}{3}$, or $\pi \iff (x, y) = (1, 1), \left(\frac{1}{2}, -1\right), \left(-\frac{1}{2}, 1\right)$, or $(-1, -1)$.

$\dfrac{dx}{d\theta} = -\sin\theta$, so $\dfrac{dx}{d\theta} = 0 \iff \sin\theta = 0 \iff \theta = 0$ or $\pi \iff$

$(x, y) = (1, 1)$ or $(-1, -1)$. Both $\dfrac{dy}{d\theta}$ and $\dfrac{dx}{d\theta}$ equal 0 when $\theta = 0$ and π.



To find the slope when $\theta = 0$, we find $\lim_{\theta\to 0}\dfrac{dy}{dx} = \lim_{\theta\to 0}\dfrac{-3\sin 3\theta}{-\sin\theta} \overset{\text{H}}{=} \lim_{\theta\to 0}\dfrac{-9\cos 3\theta}{-\cos\theta} = 9$, which is the same slope when $\theta = \pi$.

Thus, the curve has horizontal tangents at $\left(\frac{1}{2}, -1\right)$ and $\left(-\frac{1}{2}, 1\right)$, and there are no vertical tangents.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. $x = e^{\sin\theta}$, $y = e^{\cos\theta}$. The whole curve is traced out for $0 \le \theta < 2\pi$.



$\dfrac{dy}{d\theta} = -\sin\theta\, e^{\cos\theta}$, so $\dfrac{dy}{d\theta} = \theta \;\Leftrightarrow\; \sin\theta = 0 \;\Leftrightarrow\; \theta = 0 \text{ or } \pi \;\Leftrightarrow$ $(x, y) = (1, e)$ or $(1, 1/e)$. $\dfrac{dx}{d\theta} = \cos\theta\, e^{\sin\theta}$, so $\dfrac{dx}{d\theta} = 0 \;\Leftrightarrow\; \cos\theta = 0 \;\Leftrightarrow$ $\theta = \frac{\pi}{2}$ or $\frac{3\pi}{2} \;\Leftrightarrow\; (x, y) = (e, 1)$ or $(1/e, 1)$. The curve has horizontal tangents at $(1, e)$ and $(1, 1/e)$, and vertical tangents at $(e, 1)$ and $(1/e, 1)$.

21. From the graph, it appears that the rightmost point on the curve $x = t - t^6$, $y = e^t$ is about $(0.6, 2)$. To find the exact coordinates, we find the value of t for which the graph has a vertical tangent, that is, $0 = dx/dt = 1 - 6t^5 \;\Leftrightarrow\; t = 1/\sqrt[5]{6}$. Hence, the rightmost point is



$$\left(1/\sqrt[5]{6} - 1/\left(6\sqrt[5]{6}\right), e^{1/\sqrt[5]{6}}\right) = \left(5\cdot 6^{-6/5}, e^{6^{-1/5}}\right) \approx (0.58, 2.01).$$

22. From the graph, it appears that the lowest point and the leftmost point on the curve $x = t^4 - 2t$, $y = t + t^4$ are $(1.5, -0.5)$ and $(-1.2, 1.2)$, respectively. To find the exact coordinates, we solve $dy/dt = 0$ (horizontal tangents) and $dx/dt = 0$ (vertical tangents).



$\dfrac{dy}{dt} = 0 \;\Leftrightarrow\; 1 + 4t^3 = 0 \;\Leftrightarrow\; t = -\dfrac{1}{\sqrt[3]{4}}$, so the lowest point is

$$\left(\frac{1}{\sqrt[3]{256}} + \frac{2}{\sqrt[3]{4}}, -\frac{1}{\sqrt[3]{4}} + \frac{1}{\sqrt[3]{256}}\right) = \left(\frac{9}{\sqrt[3]{256}}, -\frac{3}{\sqrt[3]{256}}\right) \approx (1.42, -0.47).$$

$\dfrac{dx}{dt} = 0 \;\Leftrightarrow\; 4t^3 - 2 = 0 \;\Leftrightarrow\; t = \dfrac{1}{\sqrt[3]{2}}$, so the leftmost point is

$$\left(\frac{1}{\sqrt[3]{16}} - \frac{2}{\sqrt[3]{2}}, \frac{1}{\sqrt[3]{2}} + \frac{1}{\sqrt[3]{16}}\right) = \left(-\frac{3}{\sqrt[3]{16}}, \frac{3}{\sqrt[3]{16}}\right) \approx (-1.19, 1.19).$$

23. We graph the curve $x = t^4 - 2t^3 - 2t^2$, $y = t^3 - t$ in the viewing rectangle $[-2, 1.1]$ by $[-0.5, 0.5]$. This rectangle corresponds approximately to $t \in [-1, 0.8]$.





We estimate that the curve has horizontal tangents at about $(-1, -0.4)$ and $(-0.17, 0.39)$ and vertical tangents at about $(0, 0)$ and $(-0.19, 0.37)$. We calculate $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{3t^2 - 1}{4t^3 - 6t^2 - 4t}$. The horizontal tangents occur when $dy/dt = 3t^2 - 1 = 0 \;\Leftrightarrow\; t = \pm\frac{1}{\sqrt{3}}$, so both horizontal tangents are shown in our graph. The vertical tangents occur when

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$dx/dt = 2t(2t^2 - 3t - 2) = 0 \quad \Leftrightarrow \quad 2t(2t+1)(t-2) = 0 \quad \Leftrightarrow \quad t = 0, -\frac{1}{2}$ or 2. It seems that we have missed one vertical tangent, and indeed if we plot the curve on the t-interval $[-1.2, 2.2]$ we see that there is another vertical tangent at $(-8, 6)$.

24. We graph the curve $x = t^4 + 4t^3 - 8t^2$, $y = 2t^2 - t$ in the viewing rectangle $[-3.7, 0.2]$ by $[-0.2, 1.4]$. It appears that there is a horizontal tangent at about $(-0.4, -0.1)$, and vertical tangents at about $(-3, 1)$ and $(0, 0)$.





We calculate $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{4t-1}{4t^3 + 12t^2 - 16t}$, so there is a horizontal tangent where $dy/dt = 4t - 1 = 0 \quad \Leftrightarrow \quad t = \frac{1}{4}$. This point (the lowest point) is shown in the first graph. There are vertical tangents where $dx/dt = 4t^3 + 12t^2 - 16t = 0 \quad \Leftrightarrow$ $4t(t^2 + 3t - 4) = 0 \quad \Leftrightarrow \quad 4t(t+4)(t-1) = 0$. We have missed one vertical tangent corresponding to $t = -4$, and if we plot the graph for $t \in [-5, 3]$, we see that the curve has another vertical tangent line at approximately $(-128, 36)$.

25. $x = \cos t, \ y = \sin t \cos t. \quad dx/dt = -\sin t,$ $dy/dt = -\sin^2 t + \cos^2 t = \cos 2t. \ (x, y) = (0, 0) \quad \Leftrightarrow \quad \cos t = 0 \quad \Leftrightarrow \quad t$ is an odd multiple of $\frac{\pi}{2}$. When $t = \frac{\pi}{2}$, $dx/dt = -1$ and $dy/dt = -1$, so $dy/dx = 1$. When $t = \frac{3\pi}{2}$, $dx/dt = 1$ and $dy/dt = -1$. So $dy/dx = -1$. Thus, $y = x$ and $y = -x$ are both tangent to the curve at $(0, 0)$.



26. $x = -2\cos t, \ y = \sin t + \sin 2t$. From the graph, it appears that the curve crosses itself at the point $(1, 0)$. If this is true, then $x = 1 \quad \Leftrightarrow$ $-2\cos t = 1 \quad \Leftrightarrow \quad \cos t = -\frac{1}{2} \quad \Leftrightarrow \quad t = \frac{2\pi}{3}$ or $\frac{4\pi}{3}$ for $0 \le t \le 2\pi$. Substituting either value of t into y gives $y = 0$, confirming that $(1, 0)$ is the point where the curve crosses itself. $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{\cos t + 2\cos 2t}{2\sin t}$.



When $t = \dfrac{2\pi}{3}$, $\dfrac{dy}{dx} = \dfrac{-1/2 + 2(-1/2)}{2(\sqrt{3}/2)} = \dfrac{-3/2}{\sqrt{3}} = -\dfrac{\sqrt{3}}{2}$, so an equation of the tangent line is $y - 0 = -\dfrac{\sqrt{3}}{2}(x - 1)$, or $y = -\dfrac{\sqrt{3}}{2}x + \dfrac{\sqrt{3}}{2}$. Similarly, when $t = \dfrac{4\pi}{3}$, an equation of the tangent line is $y = \dfrac{\sqrt{3}}{2}x - \dfrac{\sqrt{3}}{2}$.

27. $x = r\theta - d\sin\theta, \ y = r - d\cos\theta.$

(a) $\dfrac{dx}{d\theta} = r - d\cos\theta, \ \dfrac{dy}{d\theta} = d\sin\theta$, so $\dfrac{dy}{dx} = \dfrac{d\sin\theta}{r - d\cos\theta}$.

(b) If $0 < d < r$, then $|d\cos\theta| \le d < r$, so $r - d\cos\theta \ge r - d > 0$. This shows that $dx/d\theta$ never vanishes, so the trochoid can have no vertical tangent if $d < r$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $x = a\cos^3\theta,\ y = a\sin^3\theta$.

(a) $\dfrac{dx}{d\theta} = -3a\cos^2\theta\sin\theta$, $\dfrac{dy}{d\theta} = 3a\sin^2\theta\cos\theta$, so $\dfrac{dy}{dx} = -\dfrac{\sin\theta}{\cos\theta} = -\tan\theta$.

(b) The tangent is horizontal $\Leftrightarrow$ $dy/dx = 0$ $\Leftrightarrow$ $\tan\theta = 0$ $\Leftrightarrow$ $\theta = n\pi$ $\Leftrightarrow$ $(x, y) = (\pm a, 0)$.

The tangent is vertical $\Leftrightarrow$ $\cos\theta = 0$ $\Leftrightarrow$ θ is an odd multiple of $\frac{\pi}{2}$ $\Leftrightarrow$ $(x, y) = (0, \pm a)$.

(c) $dy/dx = \pm 1$ $\Leftrightarrow$ $\tan\theta = \pm 1$ $\Leftrightarrow$ θ is an odd multiple of $\frac{\pi}{4}$ $\Leftrightarrow$ $(x, y) = \left(\pm\frac{\sqrt{2}}{4}a, \pm\frac{\sqrt{2}}{4}a\right)$

[All sign choices are valid.]

29. $x = 3t^2 + 1,\ y = t^3 - 1 \ \Rightarrow\ \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{3t^2}{6t} = \dfrac{t}{2}$. The tangent line has slope $\dfrac{1}{2}$ when $\dfrac{t}{2} = \dfrac{1}{2}$ $\Leftrightarrow$ $t = 1$, so the point is $(4, 0)$.

30. $x = 3t^2 + 1$, $y = 2t^3 + 1$, $\dfrac{dx}{dt} = 6t$, $\dfrac{dy}{dt} = 6t^2$, so $\dfrac{dy}{dx} = \dfrac{6t^2}{6t} = t$ [even where $t = 0$].

So at the point corresponding to parameter value t, an equation of the tangent line is $y - (2t^3 + 1) = t[x - (3t^2 + 1)]$. If this line is to pass through $(4, 3)$, we must have $3 - (2t^3 + 1) = t[4 - (3t^2 + 1)]$ $\Leftrightarrow$ $2t^3 - 2 = 3t^3 - 3t$ $\Leftrightarrow$ $t^3 - 3t + 2 = 0$ $\Leftrightarrow$ $(t - 1)^2(t + 2) = 0$ $\Leftrightarrow$ $t = 1$ or -2. Hence, the desired equations are $y - 3 = x - 4$, or $y = x - 1$, tangent to the curve at $(4, 3)$, and $y - (-15) = -2(x - 13)$, or $y = -2x + 11$, tangent to the curve at $(13, -15)$.

31. By symmetry of the ellipse about the x- and y-axes,

$$A = 4\int_0^a y\,dx = 4\int_{\pi/2}^{0} b\sin\theta\,(-a\sin\theta)\,d\theta = 4ab\int_0^{\pi/2}\sin^2\theta\,d\theta = 4ab\int_0^{\pi/2}\tfrac{1}{2}(1 - \cos 2\theta)\,d\theta$$

$$= 2ab\left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/2} = 2ab\left(\tfrac{\pi}{2}\right) = \pi ab$$

32. The curve $x = t^2 - 2t = t(t - 2)$, $y = \sqrt{t}$ intersects the y-axis when $x = 0$, that is, when $t = 0$ and $t = 2$. The corresponding values of y are 0 and $\sqrt{2}$. The shaded area is given by



$$\int_{y=0}^{y=\sqrt{2}} (x_R - x_L)\,dy = \int_{t=0}^{t=2} [0 - x(t)]\,y'(t)\,dt = -\int_0^2 (t^2 - 2t)\left(\frac{1}{2\sqrt{t}}\,dt\right)$$

$$= -\int_0^2 \left(\tfrac{1}{2}t^{3/2} - t^{1/2}\right)dt = -\left[\tfrac{1}{5}t^{5/2} - \tfrac{2}{3}t^{3/2}\right]_0^2$$

$$= -\left(\tfrac{1}{5}\cdot 2^{5/2} - \tfrac{2}{3}\cdot 2^{3/2}\right) = -2^{1/2}\left(\tfrac{4}{5} - \tfrac{4}{3}\right)$$

$$= -\sqrt{2}\left(-\tfrac{8}{15}\right) = \tfrac{8}{15}\sqrt{2}$$

33. The curve $x = t^3 + 1$, $y = 2t - t^2 = t(2 - t)$ intersects the x-axis when $y = 0$, that is, when $t = 0$ and $t = 2$. The corresponding values of x are 1 and 9. The shaded area is given by



$$\int_{x=1}^{x=9} (y_T - y_B)\,dx = \int_{t=0}^{t=2} [y(t) - 0]\,x'(t)\,dt = \int_0^2 (2t - t^2)(3t^2)\,dt$$

$$= 3\int_0^2 (2t^3 - t^4)\,dt = 3\left[\tfrac{1}{2}t^4 - \tfrac{1}{5}t^5\right]_0^2 = 3\left(8 - \tfrac{32}{5}\right) = \tfrac{24}{5}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

34. By symmetry, $A = 4\int_0^a y\,dx = 4\int_{\pi/2}^0 a\sin^3\theta(-3a\cos^2\theta\,\sin\theta)\,d\theta = 12a^2\int_0^{\pi/2}\sin^4\theta\,\cos^2\theta\,d\theta$. Now

$$\begin{aligned}\int \sin^4\theta\,\cos^2\theta\,d\theta &= \int \sin^2\theta\left(\tfrac{1}{4}\sin^2 2\theta\right)d\theta = \tfrac{1}{8}\int(1-\cos 2\theta)\sin^2 2\theta\,d\theta \\ &= \tfrac{1}{8}\int\left[\tfrac{1}{2}(1-\cos 4\theta) - \sin^2 2\theta\,\cos 2\theta\right]d\theta = \tfrac{1}{16}\theta - \tfrac{1}{64}\sin 4\theta - \tfrac{1}{48}\sin^3 2\theta + C\end{aligned}$$

so $\int_0^{\pi/2}\sin^4\theta\,\cos^2\theta\,d\theta = \left[\frac{1}{16}\theta - \frac{1}{64}\sin 4\theta - \frac{1}{48}\sin^3 2\theta\right]_0^{\pi/2} = \frac{\pi}{32}$. Thus, $A = 12a^2\left(\frac{\pi}{32}\right) = \frac{3}{8}\pi a^2$.

35. $x = r\theta - d\sin\theta,\ y = r - d\cos\theta$.

$$\begin{aligned}A &= \int_0^{2\pi r} y\,dx = \int_0^{2\pi}(r-d\cos\theta)(r-d\cos\theta)\,d\theta = \int_0^{2\pi}(r^2 - 2dr\cos\theta + d^2\cos^2\theta)\,d\theta \\ &= \left[r^2\theta - 2dr\sin\theta + \tfrac{1}{2}d^2\left(\theta + \tfrac{1}{2}\sin 2\theta\right)\right]_0^{2\pi} = 2\pi r^2 + \pi d^2\end{aligned}$$

36. (a) By symmetry, the area of $\mathcal{R}$ is twice the area inside $\mathcal{R}$ above the x-axis. The top half of the loop is described by $x = t^2,\ y = t^3 - 3t,\ -\sqrt{3} \le t \le 0$, so, using the Substitution Rule with $y = t^3 - 3t$ and $dx = 2t\,dt$, we find that

$$\begin{aligned}\text{area} &= 2\int_0^3 y\,dx = 2\int_0^{-\sqrt{3}}(t^3-3t)2t\,dt = 2\int_0^{-\sqrt{3}}(2t^4 - 6t^2)\,dt = 2\left[\tfrac{2}{5}t^5 - 2t^3\right]_0^{-\sqrt{3}} \\ &= 2\left[\tfrac{2}{5}(-3^{1/2})^5 - 2(-3^{1/2})^3\right] = 2\left[\tfrac{2}{5}\left(-9\sqrt{3}\right) - 2\left(-3\sqrt{3}\right)\right] = \tfrac{24}{5}\sqrt{3}\end{aligned}$$

(b) Here we use the formula for disks and use the Substitution Rule as in part (a):

$$\begin{aligned}\text{volume} &= \pi\int_0^3 y^2\,dx = \pi\int_0^{-\sqrt{3}}(t^3-3t)^2 2t\,dt = 2\pi\int_0^{-\sqrt{3}}(t^6 - 6t^4 + 9t^2)t\,dt = 2\pi\left[\tfrac{1}{8}t^8 - t^6 + \tfrac{9}{4}t^4\right]_0^{-\sqrt{3}} \\ &= 2\pi\left[\tfrac{1}{8}(-3^{1/2})^8 - (-3^{1/2})^6 + \tfrac{9}{4}(-3^{1/2})^4\right] = 2\pi\left[\tfrac{81}{8} - 27 + \tfrac{81}{4}\right] = \tfrac{27}{4}\pi\end{aligned}$$

(c) By symmetry, the y-coordinate of the centroid is 0. To find the x-coordinate, we note that it is the same as the x-coordinate of the centroid of the top half of $\mathcal{R}$, the area of which is $\frac{1}{2}\cdot\frac{24}{5}\sqrt{3} = \frac{12}{5}\sqrt{3}$. So, using Formula 8.3.8 with $A = \frac{12}{5}\sqrt{3}$, we get

$$\begin{aligned}\overline{x} &= \tfrac{5}{12\sqrt{3}}\int_0^3 xy\,dx = \tfrac{5}{12\sqrt{3}}\int_0^{-\sqrt{3}} t^2(t^3-3t)2t\,dt = \tfrac{5}{6\sqrt{3}}\left[\tfrac{1}{7}t^7 - \tfrac{3}{5}t^5\right]_0^{-\sqrt{3}} \\ &= \tfrac{5}{6\sqrt{3}}\left[\tfrac{1}{7}(-3^{1/2})^7 - \tfrac{3}{5}(-3^{1/2})^5\right] = \tfrac{5}{6\sqrt{3}}\left[-\tfrac{27}{7}\sqrt{3} + \tfrac{27}{5}\sqrt{3}\right] = \tfrac{9}{7}\end{aligned}$$

So the coordinates of the centroid of $\mathcal{R}$ are $(x, y) = \left(\frac{9}{7}, 0\right)$.

37. $x = t + e^{-t},\ y = t - e^{-t},\ 0 \le t \le 2$. $dx/dt = 1 - e^{-t}$ and $dy/dt = 1 + e^{-t}$, so

$(dx/dt)^2 + (dy/dt)^2 = (1 - e^{-t})^2 + (1 + e^{-t})^2 = 1 - 2e^{-t} + e^{-2t} + 1 + 2e^{-t} + e^{-2t} = 2 + 2e^{-2t}$.

Thus, $L = \int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^2 \sqrt{2 + 2e^{-2t}}\,dt \approx 3.1416$.

38. $x = t^2 - t,\ y = t^4,\ 1 \le t \le 4$. $dx/dt = 2t - 1$ and $dy/dt = 4t^3$, so

$(dx/dt)^2 + (dy/dt)^2 = (2t-1)^2 + (4t^3)^2 = 4t^2 - 4t + 1 + 16t^6$.

Thus, $L = \int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_1^4 \sqrt{16t^6 + 4t^2 - 4t + 1}\,dt \approx 255.3756$.

39. $x = t - 2\sin t,\ y = 1 - 2\cos t,\ 0 \le t \le 4\pi$. $dx/dt = 1 - 2\cos t$ and $dy/dt = 2\sin t$, so

$(dx/dt)^2 + (dy/dt)^2 = (1 - 2\cos t)^2 + (2\sin t)^2 = 1 - 4\cos t + 4\cos^2 t + 4\sin^2 t = 5 - 4\cos t$.

Thus, $L = \int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^{4\pi}\sqrt{5 - 4\cos t}\,dt \approx 26.7298$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $x = t + \sqrt{t},\ y = t - \sqrt{t},\ 0 \le t \le 1.$ $\dfrac{dx}{dt} = 1 + \dfrac{1}{2\sqrt{t}}$ and $\dfrac{dy}{dt} = 1 - \dfrac{1}{2\sqrt{t}}$, so

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = \left(1 + \frac{1}{2\sqrt{t}}\right)^2 + \left(1 - \frac{1}{2\sqrt{t}}\right)^2 = 1 + \frac{1}{\sqrt{t}} + \frac{1}{4t} + 1 - \frac{1}{\sqrt{t}} + \frac{1}{4t} = 2 + \frac{1}{2t}.$$

Thus, $L = \displaystyle\int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^1 \sqrt{2 + \frac{1}{2t}}\,dt = \lim_{t\to 0^+} \int_t^1 \sqrt{2 + \frac{1}{2t}}\,dt \approx 2.0915.$

41. $x = 1 + 3t^2,\ y = 4 + 2t^3,\ 0 \le t \le 1.$ $dx/dt = 6t$ and $dy/dt = 6t^2$, so $(dx/dt)^2 + (dy/dt)^2 = 36t^2 + 36t^4$.

Thus, $L = \displaystyle\int_0^1 \sqrt{36t^2 + 36t^4}\,dt = \int_0^1 6t\sqrt{1 + t^2}\,dt = 6\int_1^2 \sqrt{u}\,\left(\tfrac{1}{2}\,du\right)$ $[u = 1 + t^2,\ du = 2t\,dt]$

$$= 3\left[\tfrac{2}{3}u^{3/2}\right]_1^2 = 2(2^{3/2} - 1) = 2\left(2\sqrt{2} - 1\right)$$

42. $x = e^t - t,\ y = 4e^{t/2},\ 0 \le t \le 2.$ $dx/dt = e^t - 1$ and $dy/dt = 2e^{t/2}$, so

$(dx/dt)^2 + (dy/dt)^2 = (e^t - 1)^2 + (2e^{t/2})^2 = e^{2t} - 2e^t + 1 + 4e^t = e^{2t} + 2e^t + 1 = (e^t + 1)^2$. Thus,

$$L = \int_0^2 \sqrt{(e^t + 1)^2}\,dt = \int_0^2 \left|e^t + 1\right|\,dt = \int_0^2 (e^t + 1)\,dt = \left[e^t + t\right]_0^2 = (e^2 + 2) - (1 + 0) = e^2 + 1.$$

43. $x = t\sin t,\ y = t\cos t,\ 0 \le t \le 1.$ $\dfrac{dx}{dt} = t\cos t + \sin t$ and $\dfrac{dy}{dt} = -t\sin t + \cos t$, so

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = t^2\cos^2 t + 2t\sin t\,\cos t + \sin^2 t + t^2\sin^2 t - 2t\sin t\,\cos t + \cos^2 t$$
$$= t^2(\cos^2 t + \sin^2 t) + \sin^2 t + \cos^2 t = t^2 + 1.$$

Thus, $L = \int_0^1 \sqrt{t^2 + 1}\,dt \overset{21}{=} \left[\frac{1}{2}t\sqrt{t^2 + 1} + \frac{1}{2}\ln\left(t + \sqrt{t^2 + 1}\right)\right]_0^1 = \frac{1}{2}\sqrt{2} + \frac{1}{2}\ln\left(1 + \sqrt{2}\right)$.

44. $x = 3\cos t - \cos 3t,\ y = 3\sin t - \sin 3t,\ 0 \le t \le \pi.$ $\dfrac{dx}{dt} = -3\sin t + 3\sin 3t$ and $\dfrac{dy}{dt} = 3\cos t - 3\cos 3t$, so

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 9\sin^2 t - 18\sin t\,\sin 3t + 9\sin^2(3t) + 9\cos^2 t - 18\cos t\,\cos 3t + 9\cos^2(3t)$$
$$= 9(\cos^2 t + \sin^2 t) - 18(\cos t\,\cos 3t + \sin t\,\sin 3t) + 9[\cos^2(3t) + \sin^2(3t)]$$
$$= 9(1) - 18\cos(t - 3t) + 9(1) = 18 - 18\cos(-2t) = 18(1 - \cos 2t)$$
$$= 18[1 - (1 - 2\sin^2 t)] = 36\sin^2 t.$$

Thus, $L = \int_0^\pi \sqrt{36\sin^2 t}\,dt = 6\int_0^\pi |\sin t|\,dt = 6\int_0^\pi \sin t\,dt = -6\left[\cos t\right]_0^\pi = -6\,(-1 - 1) = 12.$

45.



$x = e^t\cos t,\ y = e^t\sin t,\ 0 \le t \le \pi.$

$$\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2 = [e^t(\cos t - \sin t)]^2 + [e^t(\sin t + \cos t)]^2$$
$$= (e^t)^2(\cos^2 t - 2\cos t\,\sin t + \sin^2 t)$$
$$+ (e^t)^2(\sin^2 t + 2\sin t\,\cos t + \cos^2 t$$
$$= e^{2t}(2\cos^2 t + 2\sin^2 t) = 2e^{2t}$$

Thus, $L = \int_0^\pi \sqrt{2e^{2t}}\,dt = \int_0^\pi \sqrt{2}\,e^t\,dt = \sqrt{2}\,\left[\,e^t\,\right]_0^\pi = \sqrt{2}\,(e^\pi - 1).$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

46. $x = \cos t + \ln(\tan \frac{1}{2}t), \quad y = \sin t, \quad \pi/4 \le t \le 3\pi/4.$

$\dfrac{dx}{dt} = -\sin t + \dfrac{\frac{1}{2}\sec^2(t/2)}{\tan(t/2)} = -\sin t + \dfrac{1}{2\sin(t/2)\cos(t/2)} = -\sin t + \dfrac{1}{\sin t}$ and $\dfrac{dy}{dt} = \cos t$, so

$\left(\dfrac{dx}{dt}\right)^2 + \left(\dfrac{dy}{dt}\right)^2 = \sin^2 t - 2 + \dfrac{1}{\sin^2 t} + \cos^2 t = 1 - 2 + \csc^2 t = \cot^2 t$. Thus,

$$L = \int_{\pi/4}^{3\pi/4} |\cot t|\, dt = 2\int_{\pi/4}^{\pi/2} \cot t\, dt$$
$$= 2\Big[\ln|\sin t|\Big]_{\pi/4}^{\pi/2} = 2\left(\ln 1 - \ln\frac{1}{\sqrt{2}}\right)$$
$$= 2\big(0 + \ln\sqrt{2}\,\big) = 2\big(\tfrac{1}{2}\ln 2\big) = \ln 2.$$



47.



The figure shows the curve $x = \sin t + \sin 1.5t$, $y = \cos t$ for $0 \le t \le 4\pi$.

$dx/dt = \cos t + 1.5\cos 1.5t$ and $dy/dt = -\sin t$, so

$(dx/dt)^2 + (dy/dt)^2 = \cos^2 t + 3\cos t\,\cos 1.5t + 2.25\cos^2 1.5t + \sin^2 t$.

Thus, $L = \int_0^{4\pi} \sqrt{1 + 3\cos t\,\cos 1.5t + 2.25\cos^2 1.5t}\, dt \approx 16.7102$.

48. $x = 3t - t^3$, $y = 3t^2$. $dx/dt = 3 - 3t^2$ and $dy/dt = 6t$, so

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (3 - 3t^2)^2 + (6t)^2 = (3 + 3t^2)^2$$

and the length of the loop is given by

$$L = \int_{-\sqrt{3}}^{\sqrt{3}} (3 + 3t^2)\, dt = 2\int_0^{\sqrt{3}} (3 + 3t^2)\, dt = 2\left[3t + t^3\right]_0^{\sqrt{3}}$$
$$= 2\big(3\sqrt{3} + 3\sqrt{3}\,\big) = 12\sqrt{3}$$



49. $x = t - e^t$, $y = t + e^t$, $-6 \le t \le 6$.

$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (1 - e^t)^2 + (1 + e^t)^2 = (1 - 2e^t + e^{2t}) + (1 + 2e^t + e^{2t}) = 2 + 2e^{2t}$, so $L = \int_{-6}^{6} \sqrt{2 + 2e^{2t}}\, dt$.

Set $f(t) = \sqrt{2 + 2e^{2t}}$. Then by Simpson's Rule with $n = 6$ and $\Delta t = \frac{6-(-6)}{6} = 2$, we get

$L \approx \frac{2}{3}[f(-6) + 4f(-4) + 2f(-2) + 4f(0) + 2f(2) + 4f(4) + f(6)] \approx 612.3053$.

50. $x = 2a\cot\theta \;\Rightarrow\; dx/d\theta = -2a\csc^2\theta$ and $y = 2a\sin^2\theta \;\Rightarrow\; dy/d\theta = 4a\sin\theta\,\cos\theta = 2a\sin 2\theta$.

So $L = \int_{\pi/4}^{\pi/2} \sqrt{4a^2\csc^4\theta + 4a^2\sin^2 2\theta}\, d\theta = 2a\int_{\pi/4}^{\pi/2} \sqrt{\csc^4\theta + \sin^2 2\theta}\, d\theta$. Using Simpson's Rule with

$n = 4$, $\Delta\theta = \frac{\pi/2 - \pi/4}{4} = \frac{\pi}{16}$, and $f(\theta) = \sqrt{\csc^4\theta + \sin^2 2\theta}$, we get

$L \approx 2a \cdot S_4 = (2a)\frac{\pi}{16\cdot 3}\left[f\left(\frac{\pi}{4}\right) + 4f\left(\frac{5\pi}{16}\right) + 2f\left(\frac{3\pi}{8}\right) + 4f\left(\frac{7\pi}{16}\right) + f\left(\frac{\pi}{2}\right)\right] \approx 2.2605a$.

51. $x = \sin^2 t$, $y = \cos^2 t$, $0 \le t \le 3\pi$.

$(dx/dt)^2 + (dy/dt)^2 = (2\sin t\,\cos t)^2 + (-2\cos t\sin t)^2 = 8\sin^2 t\cos^2 t = 2\sin^2 2t \;\Rightarrow$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Distance $= \int_0^{3\pi} \sqrt{2}\,|\sin 2t|\,dt = 6\sqrt{2}\int_0^{\pi/2} \sin 2t\,dt$ [by symmetry] $= -3\sqrt{2}\left[\cos 2t\right]_0^{\pi/2} = -3\sqrt{2}\,(-1-1) = 6\sqrt{2}$.

The full curve is traversed as t goes from 0 to $\frac{\pi}{2}$, because the curve is the segment of $x+y=1$ that lies in the first quadrant (since $x, y \geq 0$), and this segment is completely traversed as t goes from 0 to $\frac{\pi}{2}$. Thus, $L = \int_0^{\pi/2} \sin 2t\,dt = \sqrt{2}$, as above.

52. $x = \cos^2 t,\ y = \cos t,\ 0 \leq t \leq 4\pi$. $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (-2\cos t \sin t)^2 + (-\sin t)^2 = \sin^2 t\,(4\cos^2 t + 1)$

$$\begin{aligned}
\text{Distance} &= \int_0^{4\pi} |\sin t|\sqrt{4\cos^2 t + 1}\,dt = 4\int_0^{\pi} \sin t\sqrt{4\cos^2 t + 1}\,dt \\
&= -4\int_1^{-1} \sqrt{4u^2+1}\,du \quad [u = \cos t,\ du = -\sin t\,dt] \quad = 4\int_{-1}^{1} \sqrt{4u^2+1}\,du \\
&= 8\int_0^{1} \sqrt{4u^2+1}\,du = 8\int_0^{\tan^{-1} 2} \sec\theta \cdot \tfrac{1}{2}\sec^2\theta\,d\theta \qquad [2u = \tan\theta,\ 2\,du = \sec^2\theta\,d\theta] \\
&= 4\int_0^{\tan^{-1} 2} \sec^3\theta\,d\theta \overset{71}{=} \left[2\sec\theta\tan\theta + 2\ln|\sec\theta + \tan\theta|\right]_0^{\tan^{-1} 2} = 4\sqrt{5} + 2\ln\left(\sqrt{5}+2\right)
\end{aligned}$$

Thus, $L = \int_0^{\pi} |\sin t|\sqrt{4\cos^2 t + 1}\,dt = \sqrt{5} + \frac{1}{2}\ln\left(\sqrt{5}+2\right)$.

53. $x = a\sin\theta,\ y = b\cos\theta,\ 0 \leq \theta \leq 2\pi$.

$$\begin{aligned}
\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 &= (a\cos\theta)^2 + (-b\sin\theta)^2 = a^2\cos^2\theta + b^2\sin^2\theta = a^2(1-\sin^2\theta) + b^2\sin^2\theta \\
&= a^2 - (a^2 - b^2)\sin^2\theta = a^2 - c^2\sin^2\theta = a^2\left(1 - \frac{c^2}{a^2}\sin^2\theta\right) = a^2(1 - e^2\sin^2\theta)
\end{aligned}$$

So $L = 4\int_0^{\pi/2} \sqrt{a^2\left(1 - e^2\sin^2\theta\right)}\,d\theta$ [by symmetry] $= 4a\int_0^{\pi/2} \sqrt{1 - e^2\sin^2\theta}\,d\theta$.

54. $x = a\cos^3\theta,\ y = a\sin^3\theta$.



$$\begin{aligned}
\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 &= (-3a\cos^2\theta\,\sin\theta)^2 + (3a\sin^2\theta\,\cos\theta)^2 \\
&= 9a^2\cos^4\theta\,\sin^2\theta + 9a^2\sin^4\theta\,\cos^2\theta \\
&= 9a^2\sin^2\theta\,\cos^2\theta(\cos^2\theta + \sin^2\theta) = 9a^2\sin^2\theta\,\cos^2\theta.
\end{aligned}$$

The graph has four-fold symmetry and the curve in the first quadrant corresponds to $0 \leq \theta \leq \pi/2$. Thus,

$$\begin{aligned}
L &= 4\int_0^{\pi/2} 3a\sin\theta\,\cos\theta\,d\theta \qquad [\text{since } a > 0 \text{ and } \sin\theta \text{ and } \cos\theta \text{ are positive for } 0 \leq \theta \leq \pi/2] \\
&= 12a\left[\tfrac{1}{2}\sin^2\theta\right]_0^{\pi/2} = 12a\left(\tfrac{1}{2} - 0\right) = 6a
\end{aligned}$$

55. (a) $x = 11\cos t - 4\cos(11t/2),\ y = 11\sin t - 4\sin(11t/2)$.



Notice that $0 \leq t \leq 2\pi$ does not give the complete curve because $x(0) \neq x(2\pi)$. In fact, we must take $t \in [0, 4\pi]$ in order to obtain the complete curve, since the first term in each of the parametric equations has period 2π and the second has period $\frac{2\pi}{11/2} = \frac{4\pi}{11}$, and the least common integer multiple of these two numbers is 4π.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) We use the CAS to find the derivatives dx/dt and dy/dt, and then use Theorem 5 to find the arc length. Recent versions of Maple express the integral $\int_0^{4\pi} \sqrt{(dx/dt)^2 + (dy/dt)^2}\, dt$ as $88E\left(2\sqrt{2}\, i\right)$, where $E(x)$ is the elliptic integral $\int_0^1 \frac{\sqrt{1-x^2t^2}}{\sqrt{1-t^2}}\, dt$ and i is the imaginary number $\sqrt{-1}$.

Some earlier versions of Maple (as well as Mathematica) cannot do the integral exactly, so we use the command `evalf(Int(sqrt(diff(x,t)^2+diff(y,t)^2),t=0..4*Pi));` to estimate the length, and find that the arc length is approximately 294.03. Derive's `Para_arc_length` function in the utility file `Int_apps` simplifies the integral to $11\int_0^{4\pi} \sqrt{-4\cos t\, \cos\left(\frac{11t}{2}\right) - 4\sin t\, \sin\left(\frac{11t}{2}\right) + 5}\, dt$.

56. (a) It appears that as $t \to \infty$, $(x, y) \to \left(\frac{1}{2}, \frac{1}{2}\right)$, and as $t \to -\infty$, $(x, y) \to \left(-\frac{1}{2}, -\frac{1}{2}\right)$.



(b) By the Fundamental Theorem of Calculus, $dx/dt = \cos\left(\frac{\pi}{2}t^2\right)$ and $dy/dt = \sin\left(\frac{\pi}{2}t^2\right)$, so by Theorem 5, the length of the curve from the origin to the point with parameter value t is

$$L = \int_0^t \sqrt{\left(\tfrac{dx}{du}\right)^2 + \left(\tfrac{dy}{du}\right)^2}\, du = \int_0^t \sqrt{\cos^2\left(\tfrac{\pi}{2}u^2\right) + \sin^2\left(\tfrac{\pi}{2}u^2\right)}\, du$$
$$= \int_0^t 1\, du = t \qquad [\text{or } -t \text{ if } t < 0]$$

We have used u as the dummy variable so as not to confuse it with the upper limit of integration.

57. $x = t\sin t,\ y = t\cos t,\ 0 \le t \le \pi/2$. $dx/dt = t\cos t + \sin t$ and $dy/dt = -t\sin t + \cos t$, so

$$(dx/dt)^2 + (dy/dt)^2 = t^2\cos^2 t + 2t\sin t\, \cos t + \sin^2 t + t^2\sin^2 t - 2t\sin t\, \cos t + \cos^2 t$$
$$= t^2(\cos^2 t + \sin^2 t) + \sin^2 t + \cos^2 t = t^2 + 1$$

$$S = \int 2\pi y\, ds = \int_0^{\pi/2} 2\pi t\cos t\sqrt{t^2+1}\, dt \approx 4.7394.$$

58. $x = \sin t,\ y = \sin 2t,\ 0 \le t \le \pi/2$. $dx/dt = \cos t$ and $dy/dt = 2\cos 2t$, so $(dx/dt)^2 + (dy/dt)^2 = \cos^2 t + 4\cos^2 2t$.

$$S = \int 2\pi y\, ds = \int_0^{\pi/2} 2\pi \sin 2t\sqrt{\cos^2 t + 4\cos^2 2t}\, dt \approx 8.0285.$$

59. $x = t + e^t,\ y = e^{-t},\ 0 \le t \le 1$.

$dx/dt = 1 + e^t$ and $dy/dt = -e^{-t}$, so $(dx/dt)^2 + (dy/dt)^2 = (1+e^t)^2 + (-e^{-t})^2 = 1 + 2e^t + e^{2t} + e^{-2t}$.

$$S = \int 2\pi y\, ds = \int_0^1 2\pi e^{-t}\sqrt{1 + 2e^t + e^{2t} + e^{-2t}}\, dt \approx 10.6705.$$

60. $x = t^2 - t^3,\ y = t + t^4,\ 0 \le t \le 1$.

$(dx/dt)^2 + (dy/dt)^2 = (2t - 3t^2)^2 + (1 + 4t^3)^2 = 4t^2 - 12t^3 + 9t^4 + 1 + 8t^3 + 16t^6$, so

$$S = \int 2\pi y\, ds = \int_0^1 2\pi(t + t^4)\sqrt{16t^6 + 9t^4 - 4t^3 + 4t^2 + 1}\, dt \approx 12.7176.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. $x = t^3,\ y = t^2,\ 0 \le t \le 1.\quad \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (3t^2)^2 + (2t)^2 = 9t^4 + 4t^2.$

$$S = \int_0^1 2\pi y\sqrt{\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2}\,dt = \int_0^1 2\pi t^2\sqrt{9t^4+4t^2}\,dt = 2\pi\int_0^1 t^2\sqrt{t^2(9t^2+4)}\,dt$$

$$= 2\pi\int_4^{13}\left(\frac{u-4}{9}\right)\sqrt{u}\,\left(\tfrac{1}{18}\,du\right) \quad \left[\begin{array}{l} u = 9t^2+4,\ t^2 = (u-4)/9, \\ du = 18t\,dt,\ \text{so } t\,dt = \frac{1}{18}\,du \end{array}\right] \quad = \frac{2\pi}{9\cdot 18}\int_4^{13}(u^{3/2} - 4u^{1/2})\,du$$

$$= \tfrac{\pi}{81}\left[\tfrac{2}{5}u^{5/2} - \tfrac{8}{3}u^{3/2}\right]_4^{13} = \tfrac{\pi}{81}\cdot\tfrac{2}{15}\left[3u^{5/2} - 20u^{3/2}\right]_4^{13}$$

$$= \tfrac{2\pi}{1215}\left[\left(3\cdot 13^2\sqrt{13} - 20\cdot 13\sqrt{13}\right) - (3\cdot 32 - 20\cdot 8)\right] = \tfrac{2\pi}{1215}\left(247\sqrt{13} + 64\right)$$

62. $x = 2t^2 + 1/t,\ y = 8\sqrt{t},\ 1 \le t \le 3.$

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = \left(4t - \frac{1}{t^2}\right)^2 + \left(\frac{4}{\sqrt{t}}\right)^2 = 16t^2 - \frac{8}{t} + \frac{1}{t^4} + \frac{16}{t} = 16t^2 + \frac{8}{t} + \frac{1}{t^4} = \left(4t + \frac{1}{t^2}\right)^2.$$

$$S = \int_1^3 2\pi y\sqrt{\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2}\,dt = \int_1^3 2\pi\left(8\sqrt{t}\right)\sqrt{\left(4t + \frac{1}{t^2}\right)^2}\,dt = 16\pi\int_1^3 t^{1/2}(4t + t^{-2})\,dt$$

$$= 16\pi\int_1^3 (4t^{3/2} + t^{-3/2})\,dt = 16\pi\left[\tfrac{8}{5}t^{5/2} - 2t^{-1/2}\right]_1^3 = 16\pi\left[\left(\tfrac{72}{5}\sqrt{3} - \tfrac{2}{3}\sqrt{3}\right) - \left(\tfrac{8}{5} - 2\right)\right]$$

$$= 16\pi\left(\tfrac{206}{15}\sqrt{3} + \tfrac{6}{15}\right) = \tfrac{32\pi}{15}\left(103\sqrt{3} + 3\right)$$

63. $x = a\cos^3\theta,\ y = a\sin^3\theta,\ 0 \le \theta \le \frac{\pi}{2}.\quad \left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2 = (-3a\cos^2\theta\,\sin\theta)^2 + (3a\sin^2\theta\,\cos\theta)^2 = 9a^2\sin^2\theta\,\cos^2\theta.$

$$S = \int_0^{\pi/2} 2\pi\cdot a\sin^3\theta\cdot 3a\sin\theta\,\cos\theta\,d\theta = 6\pi a^2\int_0^{\pi/2}\sin^4\theta\,\cos\theta\,d\theta = \tfrac{6}{5}\pi a^2\left[\sin^5\theta\right]_0^{\pi/2} = \tfrac{6}{5}\pi a^2$$

64. $x = 2\cos\theta - \cos 2\theta,\ y = 2\sin\theta - \sin 2\theta \ \Rightarrow$

$$\left(\tfrac{dx}{d\theta}\right)^2 + \left(\tfrac{dy}{d\theta}\right)^2 = (-2\sin\theta + 2\sin 2\theta)^2 + (2\cos\theta - 2\cos 2\theta)^2$$

$$= 4[(\sin^2\theta - 2\sin\theta\,\sin 2\theta + \sin^2 2\theta) + (\cos^2\theta - 2\cos\theta\,\cos 2\theta + \cos^2 2\theta)]$$

$$= 4[1 + 1 - 2(\cos 2\theta\cos\theta + \sin 2\theta\sin\theta)] = 8[1 - \cos(2\theta - \theta)] = 8(1 - \cos\theta)$$

We plot the graph with parameter interval $[0, 2\pi]$, and see that we should only integrate between 0 and π. (If the interval $[0, 2\pi]$ were taken, the surface of revolution would be generated twice.) Also note that $y = 2\sin\theta - \sin 2\theta = 2\sin\theta(1 - \cos\theta)$. So



$$S = \int_0^{\pi} 2\pi\cdot 2\sin\theta(1-\cos\theta)\,2\sqrt{2}\sqrt{1-\cos\theta}\,d\theta$$

$$= 8\sqrt{2}\pi\int_0^{\pi}(1-\cos\theta)^{3/2}\sin\theta\,d\theta = 8\sqrt{2}\pi\int_0^2\sqrt{u^3}\,du \quad \left[\begin{array}{l} u = 1-\cos\theta, \\ du = \sin\theta\,d\theta \end{array}\right]$$

$$= 8\sqrt{2}\pi\left[\left(\tfrac{2}{5}\right)u^{5/2}\right]_0^2 = \tfrac{16}{5}\sqrt{2}\pi(2^{5/2}) = \tfrac{128}{5}\pi$$

65. $x = 3t^2,\ y = 2t^3,\ 0 \le t \le 5 \ \Rightarrow\ \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (6t)^2 + (6t^2)^2 = 36t^2(1+t^2) \ \Rightarrow$

$$S = \int_0^5 2\pi x\sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^5 2\pi(3t^2)6t\sqrt{1+t^2}\,dt = 18\pi\int_0^5 t^2\sqrt{1+t^2}\,2t\,dt$$

$$= 18\pi\int_1^{26}(u-1)\sqrt{u}\,du \quad \left[\begin{array}{l} u = 1+t^2, \\ du = 2t\,dt \end{array}\right] \quad = 18\pi\int_1^{26}(u^{3/2} - u^{1/2})\,du = 18\pi\left[\tfrac{2}{5}u^{5/2} - \tfrac{2}{3}u^{3/2}\right]_1^{26}$$

$$= 18\pi\left[\left(\tfrac{2}{5}\cdot 676\sqrt{26} - \tfrac{2}{3}\cdot 26\sqrt{26}\right) - \left(\tfrac{2}{5} - \tfrac{2}{3}\right)\right] = \tfrac{24}{5}\pi\left(949\sqrt{26} + 1\right)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

66. $x = e^t - t,\ y = 4e^{t/2},\ 0 \le t \le 1.\ \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (e^t-1)^2 + (2e^{t/2})^2 = e^{2t} + 2e^t + 1 = (e^t+1)^2.$

$$S = \int_0^1 2\pi(e^t - t)\sqrt{(e^t-1)^2 + (2e^{t/2})^2}\,dt = \int_0^1 2\pi(e^t - t)(e^t+1)d$$
$$= 2\pi\left[\tfrac{1}{2}e^{2t} + e^t - (t-1)e^t - \tfrac{1}{2}t^2\right]_0^1 = \pi(e^2 + 2e - 6)$$

67. If f' is continuous and $f'(t) \neq 0$ for $a \le t \le b$, then either $f'(t) > 0$ for all t in $[a,b]$ or $f'(t) < 0$ for all t in $[a,b]$. Thus, f is monotonic (in fact, strictly increasing or strictly decreasing) on $[a,b]$. It follows that f has an inverse. Set $F = g \circ f^{-1}$, that is, define F by $F(x) = g(f^{-1}(x))$. Then $x = f(t) \ \Rightarrow\ f^{-1}(x) = t$, so $y = g(t) = g(f^{-1}(x)) = F(x)$.

68. By Formula 8.2.5 with $y = F(x)$, $S = \int_a^b 2\pi F(x)\sqrt{1 + [F'(x)]^2}\,dx$. But by Formula 10.2.1,

$1 + [F'(x)]^2 = 1 + \left(\dfrac{dy}{dx}\right)^2 = 1 + \left(\dfrac{dy/dt}{dx/dt}\right)^2 = \dfrac{(dx/dt)^2 + (dy/dt)^2}{(dx/dt)^2}$. Using the Substitution Rule with $x = x(t)$,

where $a = x(\alpha)$ and $b = x(\beta)$, we have $\left[\text{since } dx = \dfrac{dx}{dt}\,dt\right]$

$S = \displaystyle\int_\alpha^\beta 2\pi\, F(x(t))\sqrt{\frac{(dx/dt)^2 + (dy/dt)^2}{(dx/dt)^2}}\,\frac{dx}{dt}\,dt = \int_\alpha^\beta 2\pi y\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt$, which is Formula 10.2.6.

69. (a) $\phi = \tan^{-1}\left(\dfrac{dy}{dx}\right) \ \Rightarrow\ \dfrac{d\phi}{dt} = \dfrac{d}{dt}\tan^{-1}\left(\dfrac{dy}{dx}\right) = \dfrac{1}{1 + (dy/dx)^2}\left[\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)\right]$. But $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{\dot y}{\dot x} \ \Rightarrow$

$\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right) = \dfrac{d}{dt}\left(\dfrac{\dot y}{\dot x}\right) = \dfrac{\ddot y\dot x - \ddot x\dot y}{\dot x^2} \ \Rightarrow\ \dfrac{d\phi}{dt} = \dfrac{1}{1 + (\dot y/\dot x)^2}\left(\dfrac{\ddot y\dot x - \ddot x\dot y}{\dot x^2}\right) = \dfrac{\dot x\ddot y - \ddot x\dot y}{\dot x^2 + \dot y^2}$. Using the Chain Rule, and the

fact that $s = \displaystyle\int_0^t \sqrt{\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2}\,dt \ \Rightarrow\ \tfrac{ds}{dt} = \sqrt{\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2} = \left(\dot x^2 + \dot y^2\right)^{1/2}$, we have that

$\dfrac{d\phi}{ds} = \dfrac{d\phi/dt}{ds/dt} = \left(\dfrac{\dot x\ddot y - \ddot x\dot y}{\dot x^2 + \dot y^2}\right)\dfrac{1}{(\dot x^2 + \dot y^2)^{1/2}} = \dfrac{\dot x\ddot y - \ddot x\dot y}{(\dot x^2 + \dot y^2)^{3/2}}$. So $\kappa = \left|\dfrac{d\phi}{ds}\right| = \left|\dfrac{\dot x\ddot y - \ddot x\dot y}{(\dot x^2 + \dot y^2)^{3/2}}\right| = \dfrac{|\dot x\ddot y - \ddot x\dot y|}{(\dot x^2 + \dot y^2)^{3/2}}$.

(b) $x = x$ and $y = f(x) \ \Rightarrow\ \dot x = 1, \ddot x = 0$ and $\dot y = \dfrac{dy}{dx}, \ddot y = \dfrac{d^2y}{dx^2}$.

So $\kappa = \dfrac{|1\cdot(d^2y/dx^2) - 0\cdot(dy/dx)|}{[1 + (dy/dx)^2]^{3/2}} = \dfrac{|d^2y/dx^2|}{[1 + (dy/dx)^2]^{3/2}}$.

70. (a) $y = x^2 \ \Rightarrow\ \dfrac{dy}{dx} = 2x \ \Rightarrow\ \dfrac{d^2y}{dx^2} = 2$. So $\kappa = \dfrac{|d^2y/dx^2|}{[1 + (dy/dx)^2]^{3/2}} = \dfrac{2}{(1 + 4x^2)^{3/2}}$, and at $(1,1)$,

$\kappa = \dfrac{2}{5^{3/2}} = \dfrac{2}{5\sqrt{5}}$.

(b) $\kappa' = \dfrac{d\kappa}{dx} = -3(1 + 4x^2)^{-5/2}(8x) = 0 \ \Leftrightarrow\ x = 0 \ \Rightarrow\ y = 0$. This is a maximum since $\kappa' > 0$ for $x < 0$ and

$\kappa' < 0$ for $x > 0$. So the parabola $y = x^2$ has maximum curvature at the origin.

71. $x = \theta - \sin\theta \ \Rightarrow\ \dot x = 1 - \cos\theta \ \Rightarrow\ \ddot x = \sin\theta$, and $y = 1 - \cos\theta \ \Rightarrow\ \dot y = \sin\theta \ \Rightarrow\ \ddot y = \cos\theta$. Therefore,

$\kappa = \dfrac{|\cos\theta - \cos^2\theta - \sin^2\theta|}{[(1 - \cos\theta)^2 + \sin^2\theta]^{3/2}} = \dfrac{|\cos\theta - (\cos^2\theta + \sin^2\theta)|}{(1 - 2\cos\theta + \cos^2\theta + \sin^2\theta)^{3/2}} = \dfrac{|\cos\theta - 1|}{(2 - 2\cos\theta)^{3/2}}$. The top of the arch is

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

characterized by a horizontal tangent, and from Example 2(b) in Section 10.2, the tangent is horizontal when $\theta = (2n-1)\pi$, so take $n = 1$ and substitute $\theta = \pi$ into the expression for κ: $\kappa = \dfrac{|\cos\pi - 1|}{(2 - 2\cos\pi)^{3/2}} = \dfrac{|-1-1|}{[2-2(-1)]^{3/2}} = \dfrac{1}{4}$.

72. (a) Every straight line has parametrizations of the form $x = a + vt$, $y = b + wt$, where a, b are arbitrary and v, $w \neq 0$. For example, a straight line passing through distinct points (a, b) and (c, d) can be described as the parametrized curve $x = a + (c-a)t$, $y = b + (d-b)t$. Starting with $x = a + vt$, $y = b + wt$, we compute $\dot{x} = v$, $\dot{y} = w$, $\ddot{x} = \ddot{y} = 0$, and $\kappa = \dfrac{|v \cdot 0 - w \cdot 0|}{(v^2 + w^2)^{3/2}} = 0$.

(b) Parametric equations for a circle of radius r are $x = r\cos\theta$ and $y = r\sin\theta$. We can take the center to be the origin. So $\dot{x} = -r\sin\theta \;\Rightarrow\; \ddot{x} = -r\cos\theta$ and $\dot{y} = r\cos\theta \;\Rightarrow\; \ddot{y} = -r\sin\theta$. Therefore, $\kappa = \dfrac{|r^2\sin^2\theta + r^2\cos^2\theta|}{(r^2\sin^2\theta + r^2\cos^2\theta)^{3/2}} = \dfrac{r^2}{r^3} = \dfrac{1}{r}$. And so for any θ (and thus any point), $\kappa = \dfrac{1}{r}$.

73. The coordinates of T are $(r\cos\theta, r\sin\theta)$. Since TP was unwound from arc TA, TP has length $r\theta$. Also $\angle PTQ = \angle PTR - \angle QTR = \frac{1}{2}\pi - \theta$, so P has coordinates $x = r\cos\theta + r\theta\cos\left(\frac{1}{2}\pi - \theta\right) = r(\cos\theta + \theta\sin\theta)$, $y = r\sin\theta - r\theta\sin\left(\frac{1}{2}\pi - \theta\right) = r(\sin\theta - \theta\cos\theta)$.



74. If the cow walks with the rope taut, it traces out the portion of the involute in Exercise 73 corresponding to the range $0 \le \theta \le \pi$, arriving at the point $(-r, \pi r)$ when $\theta = \pi$. With the rope now fully extended, the cow walks in a semicircle of radius πr, arriving at $(-r, -\pi r)$. Finally, the cow traces out another portion of the involute, namely the reflection about the x-axis of the initial involute path. (This corresponds to the range $-\pi \le \theta \le 0$.) Referring to the figure, we see that the total grazing area is $2(A_1 + A_3)$. A_3 is one-quarter of the area of a circle of radius πr, so $A_3 = \frac{1}{4}\pi(\pi r)^2 = \frac{1}{4}\pi^3 r^2$. We will compute $A_1 + A_2$ and then subtract $A_2 = \frac{1}{2}\pi r^2$ to obtain A_1.



To find $A_1 + A_2$, first note that the rightmost point of the involute is $\left(\frac{\pi}{2}r, r\right)$. [To see this, note that $dx/d\theta = 0$ when $\theta = 0$ or $\frac{\pi}{2}$. $\theta = 0$ corresponds to the cusp at $(r, 0)$ and $\theta = \frac{\pi}{2}$ corresponds to $\left(\frac{\pi}{2}r, r\right)$.] The leftmost point of the involute is $(-r, \pi r)$. Thus, $A_1 + A_2 = \int_{\theta=\pi}^{\pi/2} y\,dx - \int_{\theta=0}^{\pi/2} y\,dx = \int_{\theta=\pi}^{0} y\,dx$.

Now $y\,dx = r(\sin\theta - \theta\cos\theta)\,r\theta\cos\theta\,d\theta = r^2(\theta\sin\theta\,\cos\theta - \theta^2\cos^2\theta)d\theta$. Integrate: $(1/r^2)\int y\,dx = -\theta\cos^2\theta - \frac{1}{2}\left(\theta^2 - 1\right)\sin\theta\,\cos\theta - \frac{1}{6}\theta^3 + \frac{1}{2}\theta + C$. This enables us to compute

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$$A_1 + A_2 = r^2\left[-\theta\cos^2\theta - \tfrac{1}{2}(\theta^2-1)\sin\theta\cos\theta - \tfrac{1}{6}\theta^3 + \tfrac{1}{2}\theta\right]_{\pi}^{0} = r^2\left[0 - \left(-\pi - \frac{\pi^3}{6} + \frac{\pi}{2}\right)\right] = r^2\left(\frac{\pi}{2} + \frac{\pi^3}{6}\right)$$

Therefore, $A_1 = (A_1 + A_2) - A_2 = \frac{1}{6}\pi^3 r^2$, so the grazing area is $2(A_1 + A_3) = 2\left(\frac{1}{6}\pi^3 r^2 + \frac{1}{4}\pi^3 r^2\right) = \frac{5}{6}\pi^3 r^2$.

LABORATORY PROJECT Bézier Curves

1. The parametric equations for a cubic Bézier curve are

$$x = x_0(1-t)^3 + 3x_1t(1-t)^2 + 3x_2t^2(1-t) + x_3t^3$$
$$y = y_0(1-t)^3 + 3y_1t(1-t)^2 + 3y_2t^2(1-t) + y_3t^3$$

where $0 \le t \le 1$. We are given the points $P_0(x_0, y_0) = (4, 1)$, $P_1(x_1, y_1) = (28, 48)$, $P_2(x_2, y_2) = (50, 42)$, and $P_3(x_3, y_3) = (40, 5)$. The curve is then given by

$$x(t) = 4(1-t)^3 + 3\cdot 28t(1-t)^2 + 3\cdot 50t^2(1-t) + 40t^3$$
$$y(t) = 1(1-t)^3 + 3\cdot 48t(1-t)^2 + 3\cdot 42t^2(1-t) + 5t^3$$

where $0 \le t \le 1$. The line segments are of the form $x = x_0 + (x_1 - x_0)t$, $y = y_0 + (y_1 - y_0)t$:

P_0P_1 $\quad x = 4 + 24t, \quad y = 1 + 47t$

P_1P_2 $\quad x = 28 + 22t, \quad y = 48 - 6t$

P_2P_3 $\quad x = 50 - 10t, \quad y = 42 - 37t$



2. It suffices to show that the slope of the tangent at P_0 is the same as that of line segment P_0P_1, namely $\dfrac{y_1 - y_0}{x_1 - x_0}$.

We calculate the slope of the tangent to the Bézier curve:

$$\frac{dy/dt}{dx/dt} = \frac{-3y_0(1-t)^2 + 3y_1\left[-2t(1-t) + (1-t)^2\right] + 3y_2\left[-t^2 + (2t)(1-t)\right] + 3y_3t^2}{-3x_0^2(1-t) + 3x_1[-2t(1-t) + (1-t)^2] + 3x_2[-t^2 + (2t)(1-t)] + 3x_3t^2}$$

At point P_0, $t = 0$, so the slope of the tangent is $\dfrac{-3y_0 + 3y_1}{-3x_0 + 3x_1} = \dfrac{y_1 - y_0}{x_1 - x_0}$. So the tangent to the curve at P_0 passes through P_1. Similarly, the slope of the tangent at point P_3 [where $t = 1$] is $\dfrac{-3y_2 + 3y_3}{-3x_2 + 3x_3} = \dfrac{y_3 - y_2}{x_3 - x_2}$, which is also the slope of line P_2P_3.

3. It seems that if P_1 were to the right of P_2, a loop would appear.

We try setting $P_1 = (110, 30)$, and the resulting curve does indeed have a loop.



4. Based on the behavior of the Bézier curve in Problems 1–3, we suspect that the four control points should be in an exaggerated C shape. We try $P_0(10, 12)$, $P_1(4, 15)$, $P_2(4, 5)$, and $P_3(10, 8)$, and these produce a decent C. If you are using a CAS, it may be necessary to instruct it to make the x- and y-scales the same so as not to distort the figure (this is called a "constrained projection" in Maple.)



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. We use the same P_0 and P_1 as in Problem 4, and use part of our C as the top of an S. To prevent the center line from slanting up too much, we move P_2 up to $(4, 6)$ and P_3 down and to the left, to $(8, 7)$. In order to have a smooth joint between the top and bottom halves of the S (and a symmetric S), we determine points P_4, P_5, and P_6 by rotating points P_2, P_1, and P_0 about the center of the letter (point P_3). The points are therefore $P_4(12, 8)$, $P_5(12, -1)$, and $P_6(6, 2)$.



10.3 Polar Coordinates

1. (a) $\left(1, \frac{\pi}{4}\right)$



By adding 2π to $\frac{\pi}{4}$, we obtain the point $\left(1, \frac{9\pi}{4}\right)$, which satisfies the $r > 0$ requirement. The direction opposite $\frac{\pi}{4}$ is $\frac{5\pi}{4}$, so $\left(-1, \frac{5\pi}{4}\right)$ is a point that satisfies the $r < 0$ requirement.

(b) $\left(-2, \frac{3\pi}{2}\right)$



$r > 0$: $\left(-(-2), \frac{3\pi}{2} - \pi\right) = \left(2, \frac{\pi}{2}\right)$

$r < 0$: $\left(-2, \frac{3\pi}{2} + 2\pi\right) = \left(-2, \frac{7\pi}{2}\right)$

(c) $\left(3, -\frac{\pi}{3}\right)$



$r > 0$: $\left(3, -\frac{\pi}{3} + 2\pi\right) = \left(3, \frac{5\pi}{3}\right)$

$r < 0$: $\left(-3, -\frac{\pi}{3} + \pi\right) = \left(-3, \frac{2\pi}{3}\right)$

2. (a) $\left(2, \frac{5\pi}{6}\right)$



$r > 0$: $\left(2, \frac{5\pi}{6} + 2\pi\right) = \left(2, \frac{17\pi}{6}\right)$

$r < 0$: $\left(-2, \frac{5\pi}{6} - \pi\right) = \left(-2, -\frac{\pi}{6}\right)$

(b) $\left(1, -\frac{2\pi}{3}\right)$



$r > 0$: $\left(1, -\frac{2\pi}{3} + 2\pi\right) = \left(1, \frac{4\pi}{3}\right)$

$r < 0$: $\left(-1, -\frac{2\pi}{3} + \pi\right) = \left(-1, \frac{\pi}{3}\right)$

(c) $\left(-1, \frac{5\pi}{4}\right)$



$r > 0$: $\left(-(-1), \frac{5\pi}{4} - \pi\right) = \left(1, \frac{\pi}{4}\right)$

$r < 0$: $\left(-1, \frac{5\pi}{4} - 2\pi\right) = \left(-1, -\frac{3\pi}{4}\right)$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

3. (a)



$x = 2\cos\frac{3\pi}{2} = 2(0) = 0$ and $y = 2\sin\frac{3\pi}{2} = 2(-1) = -2$ give us the Cartesian coordinates $(0, -2)$.

(b)



$x = \sqrt{2}\cos\dfrac{\pi}{4} = \sqrt{2}\left(\dfrac{1}{\sqrt{2}}\right) = 1$ and $y = \sqrt{2}\sin\dfrac{\pi}{4} = \sqrt{2}\left(\dfrac{1}{\sqrt{2}}\right) = 1$ give us the Cartesian coordinates $(1, 1)$.

(c)



$x = -1\cos\left(-\dfrac{\pi}{6}\right) = -1\left(\dfrac{\sqrt{3}}{2}\right) = -\dfrac{\sqrt{3}}{2}$ and

$y = -1\sin\left(-\dfrac{\pi}{6}\right) = -1\left(-\dfrac{1}{2}\right) = \dfrac{1}{2}$ give us the Cartesian coordinates $\left(-\dfrac{\sqrt{3}}{2}, \dfrac{1}{2}\right)$.

4. (a)



$x = 4\cos\dfrac{4\pi}{3} = 4\left(-\dfrac{1}{2}\right) = -2$ and

$y = 4\sin\dfrac{4\pi}{3} = 4\left(-\dfrac{\sqrt{3}}{2}\right) = -2\sqrt{3}$ give us the Cartesian coordinates $\left(-2, -2\sqrt{3}\right)$.

(b)



$x = -2\cos\dfrac{3\pi}{4} = -2\left(-\dfrac{\sqrt{2}}{2}\right) = \sqrt{2}$ and

$y = -2\sin\dfrac{3\pi}{4} = -2\left(\dfrac{\sqrt{2}}{2}\right) = -\sqrt{2}$ give us the Cartesian coordinates $\left(\sqrt{2}, -\sqrt{2}\right)$.

(c)



$x = -3\cos\left(-\dfrac{\pi}{3}\right) = -3\left(\dfrac{1}{2}\right) = -\dfrac{3}{2}$ and

$y = -3\sin\left(-\dfrac{\pi}{3}\right) = -3\left(-\dfrac{\sqrt{3}}{2}\right) = \dfrac{3\sqrt{3}}{2}$ give us the Cartesian coordinates $\left(-\dfrac{3}{2}, \dfrac{3\sqrt{3}}{2}\right)$.

5. (a) $x = -4$ and $y = 4 \;\Rightarrow\; r = \sqrt{(-4)^2 + 4^2} = 4\sqrt{2}$ and $\tan\theta = \frac{4}{-4} = -1 \quad [\theta = -\frac{\pi}{4} + n\pi]$. Since $(-4, 4)$ is in the second quadrant, the polar coordinates are (i) $\left(4\sqrt{2}, \frac{3\pi}{4}\right)$ and (ii) $\left(-4\sqrt{2}, \frac{7\pi}{4}\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $x = 3$ and $y = 3\sqrt{3} \;\Rightarrow\; r = \sqrt{3^2 + (3\sqrt{3})^2} = \sqrt{9+27} = 6$ and $\tan\theta = \dfrac{3\sqrt{3}}{3} = \sqrt{3} \quad [\theta = \frac{\pi}{3} + n\pi]$.

Since $(3, 3\sqrt{3})$ is in the first quadrant, the polar coordinates are (i) $(6, \frac{\pi}{3})$ and (ii) $(-6, \frac{4\pi}{3})$.

6. (a) $x = \sqrt{3}$ and $y = -1 \;\Rightarrow\; r = \sqrt{(\sqrt{3})^2 + (-1)^2} = 2$ and $\tan\theta = \dfrac{-1}{\sqrt{3}} \quad [\theta = -\frac{\pi}{6} + n\pi]$. Since $(\sqrt{3}, -1)$ is in the fourth quadrant, the polar coordinates are (i) $(2, \frac{11\pi}{6})$ and (ii) $(-2, \frac{5\pi}{6})$.

(b) $x = -6$ and $y = 0 \;\Rightarrow\; r = \sqrt{(-6)^2 + 0^2} = 6$ and $\tan\theta = \frac{0}{-6} = 0 \quad [\theta = n\pi]$. Since $(-6, 0)$ is on the negative x-axis, the polar coordinates are (i) $(6, \pi)$ and (ii) $(-6, 0)$.

7. $r \geq 1$. The curve $r = 1$ represents a circle with center O and radius 1. So $r \geq 1$ represents the region on or outside the circle. Note that θ can take on any value.



8. $0 \leq r < 2, \; \pi \leq \theta \leq 3\pi/2$. This is the region inside the circle $r = 2$ in the third quadrant.



9. $r \geq 0, \; \pi/4 \leq \theta \leq 3\pi/4$.
$\theta = k$ represents a line through O.



10. $1 \leq r \leq 3, \; \pi/6 < \theta < 5\pi/6$



11. $2 < r < 3, \; \frac{5\pi}{3} \leq \theta \leq \frac{7\pi}{3}$



12. $r \geq 1, \; \pi \leq \theta \leq 2\pi$



13. Converting the polar coordinates $(4, \frac{4\pi}{3})$ and $(6, \frac{5\pi}{3})$ to Cartesian coordinates gives us $(4\cos\frac{4\pi}{3}, 4\sin\frac{4\pi}{3}) = (-2, -2\sqrt{3})$ and $(6\cos\frac{5\pi}{3}, 6\sin\frac{5\pi}{3}) = (3, -3\sqrt{3})$. Now use the distance formula

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{[3-(-2)]^2 + [-3\sqrt{3} - (-2\sqrt{3})]^2}$$
$$= \sqrt{5^2 + (-\sqrt{3})^2} = \sqrt{25+3} = \sqrt{28} = 2\sqrt{7}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

14. The points (r_1, θ_1) and (r_2, θ_2) in Cartesian coordinates are $(r_1 \cos\theta_1, r_1 \sin\theta_1)$ and $(r_2 \cos\theta_2, r_2 \sin\theta_2)$, respectively. The *square* of the distance between them is

$$\begin{aligned}(r_2\cos\theta_2 - r_1\cos\theta_1)^2 &+ (r_2\sin\theta_2 - r_1\sin\theta_1)^2 \\ &= \left(r_2^2\cos^2\theta_2 - 2r_1r_2\cos\theta_1\cos\theta_2 + r_1^2\cos^2\theta_1\right) + \left(r_2^2\sin^2\theta_2 - 2r_1r_2\sin\theta_1\sin\theta_2 + r_1^2\sin^2\theta_1\right) \\ &= r_1^2\left(\sin^2\theta_1 + \cos^2\theta_1\right) + r_2^2\left(\sin^2\theta_2 + \cos^2\theta_2\right) - 2r_1r_2(\cos\theta_1\cos\theta_2 + \sin\theta_1\sin\theta_2) \\ &= r_1^2 - 2r_1r_2\cos(\theta_1 - \theta_2) + r_2^2,\end{aligned}$$

so the distance between them is $\sqrt{r_1^2 - 2r_1r_2\cos(\theta_1 - \theta_2) + r_2^2}$.

15. $r^2 = 5 \quad \Leftrightarrow \quad x^2 + y^2 = 5$, a circle of radius $\sqrt{5}$ centered at the origin.

16. $r = 4\sec\theta \quad \Leftrightarrow \quad \dfrac{r}{\sec\theta} = 4 \quad \Leftrightarrow \quad r\cos\theta = 4 \quad \Leftrightarrow \quad x = 4$, a vertical line.

17. $r = 5\cos\theta \quad \Rightarrow \quad r^2 = 5r\cos\theta \quad \Leftrightarrow \quad x^2 + y^2 = 5x \quad \Leftrightarrow \quad x^2 - 5x + \frac{25}{4} + y^2 = \frac{25}{4} \quad \Leftrightarrow \quad \left(x - \frac{5}{2}\right)^2 + y^2 = \frac{25}{4}$, a circle of radius $\frac{5}{2}$ centered at $\left(\frac{5}{2}, 0\right)$. The first two equations are actually equivalent since $r^2 = 5r\cos\theta \quad \Rightarrow$ $r(r - 5\cos\theta) = 0 \quad \Rightarrow \quad r = 0$ or $r = 5\cos\theta$. But $r = 5\cos\theta$ gives the point $r = 0$ (the pole) when $\theta = 0$. Thus, the equation $r = 5\cos\theta$ is equivalent to the compound condition ($r = 0$ or $r = 5\cos\theta$).

18. $\theta = \dfrac{\pi}{3} \quad \Rightarrow \quad \tan\theta = \tan\dfrac{\pi}{3} \quad \Rightarrow \quad \dfrac{y}{x} = \sqrt{3} \quad \Leftrightarrow \quad y = \sqrt{3}\,x$, a line through the origin.

19. $r^2\cos 2\theta = 1 \quad \Leftrightarrow \quad r^2(\cos^2\theta - \sin^2\theta) = 1 \quad \Leftrightarrow \quad (r\cos\theta)^2 - (r\sin\theta)^2 = 1 \quad \Leftrightarrow \quad x^2 - y^2 = 1$, a hyperbola centered at the origin with foci on the x-axis.

20. $r^2\sin 2\theta = 1 \quad \Leftrightarrow \quad r^2(2\sin\theta\cos\theta) = 1 \quad \Leftrightarrow \quad 2(r\cos\theta)(r\sin\theta) = 1 \quad \Leftrightarrow \quad 2xy = 1 \quad \Leftrightarrow \quad xy = \frac{1}{2}$, a hyperbola centered at the origin with foci on the line $y = x$.

21. $y = 2 \quad \Leftrightarrow \quad r\sin\theta = 2 \quad \Leftrightarrow \quad r = \dfrac{2}{\sin\theta} \quad \Leftrightarrow \quad r = 2\csc\theta$

22. $y = x \quad \Rightarrow \quad \dfrac{y}{x} = 1 \ [x \neq 0] \quad \Rightarrow \quad \tan\theta = 1 \quad \Rightarrow \quad \theta = \tan^{-1}1 \quad \Rightarrow \quad \theta = \dfrac{\pi}{4}$ or $\dfrac{5\pi}{4}$ [either includes the pole]

23. $y = 1 + 3x \quad \Leftrightarrow \quad r\sin\theta = 1 + 3r\cos\theta \quad \Leftrightarrow \quad r\sin\theta - 3r\cos\theta = 1 \quad \Leftrightarrow \quad r(\sin\theta - 3\cos\theta) = 1 \quad \Leftrightarrow$ $r = \dfrac{1}{\sin\theta - 3\cos\theta}$

24. $4y^2 = x \quad \Leftrightarrow \quad 4(r\sin\theta)^2 = r\cos\theta \quad \Leftrightarrow \quad 4r^2\sin^2\theta - r\cos\theta = 0 \quad \Leftrightarrow \quad r(4r\sin^2\theta - \cos\theta) = 0 \quad \Leftrightarrow \quad r = 0$ or $r = \dfrac{\cos\theta}{4\sin^2\theta} \quad \Leftrightarrow \quad r = 0$ or $r = \frac{1}{4}\cot\theta\csc\theta$. $r = 0$ is included in $r = \frac{1}{4}\cot\theta\csc\theta$ when $\theta = \frac{\pi}{2}$, so the curve is represented by the single equation $r = \frac{1}{4}\cot\theta\csc\theta$.

25. $x^2 + y^2 = 2cx \quad \Leftrightarrow \quad r^2 = 2cr\cos\theta \quad \Leftrightarrow \quad r^2 - 2cr\cos\theta = 0 \quad \Leftrightarrow \quad r(r - 2c\cos\theta) = 0 \quad \Leftrightarrow \quad r = 0$ or $r = 2c\cos\theta$. $r = 0$ is included in $r = 2c\cos\theta$ when $\theta = \frac{\pi}{2} + n\pi$, so the curve is represented by the single equation $r = 2c\cos\theta$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $x^2 - y^2 = 4 \iff (r\cos\theta)^2 - (r\sin\theta)^2 = 4 \iff r^2\cos^2\theta - r^2\sin^2\theta = 4 \iff r^2(\cos^2\theta - \sin^2\theta) = 4 \iff r^2\cos 2\theta = 4$

27. (a) The description leads immediately to the polar equation $\theta = \frac{\pi}{6}$, and the Cartesian equation $y = \tan\left(\frac{\pi}{6}\right)x = \frac{1}{\sqrt{3}}x$ is slightly more difficult to derive.

(b) The easier description here is the Cartesian equation $x = 3$.

28. (a) Because its center is not at the origin, it is more easily described by its Cartesian equation, $(x-2)^2 + (y-3)^2 = 5^2$.

(b) This circle is more easily given in polar coordinates: $r = 4$. The Cartesian equation is also simple: $x^2 + y^2 = 16$.

29. $r = -2\sin\theta$





30. $r = 1 - \cos\theta$





31. $r = 2(1 + \cos\theta)$





32. $r = 1 + 2\cos\theta$





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. $r = \theta, \quad \theta \geq 0$



34. $r = \theta^2, \ -2\pi \leq \theta \leq 2\pi$



35. $r = 3\cos 3\theta$



36. $r = -\sin 5\theta$



37. $r = 2\cos 4\theta$



38. $r = 2\sin 6\theta$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. $r = 1 + 3\cos\theta$





40. $r = 1 + 5\sin\theta$





41. $r^2 = 9\sin 2\theta$





42. $r^2 = \cos 4\theta$





43. $r = 2 + \sin 3\theta$





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

44. $r^2\theta = 1 \quad \Leftrightarrow \quad r = \pm 1/\sqrt{\theta}$ for $\theta > 0$



45. $r = \sin(\theta/2)$



46. $r = \cos(\theta/3)$



47. For $\theta = 0, \pi$, and 2π, r has its minimum value of about 0.5. For $\theta = \frac{\pi}{2}$ and $\frac{3\pi}{2}$, r attains its maximum value of 2. We see that the graph has a similar shape for $0 \le \theta \le \pi$ and $\pi \le \theta \le 2\pi$.



48. The given graph has a maximum of 2 for $\theta = 0$, a minimum of 1 for $\theta = \frac{\pi}{4}$, and then a maximum of 2 for $\theta = \frac{\pi}{2}$. This pattern is repeated 4 times for $0 \le \theta \le 2\pi$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. $x = r\cos\theta = (4 + 2\sec\theta)\cos\theta = 4\cos\theta + 2$. Now, $r \to \infty \;\Rightarrow$ $(4 + 2\sec\theta) \to \infty \;\Rightarrow\; \theta \to \left(\frac{\pi}{2}\right)^-$ or $\theta \to \left(\frac{3\pi}{2}\right)^+$ [since we need only consider $0 \le \theta < 2\pi$], so $\lim\limits_{r\to\infty} x = \lim\limits_{\theta\to\pi/2^-} (4\cos\theta + 2) = 2$. Also, $r \to -\infty \;\Rightarrow\; (4 + 2\sec\theta) \to -\infty \;\Rightarrow\; \theta \to \left(\frac{\pi}{2}\right)^+$ or $\theta \to \left(\frac{3\pi}{2}\right)^-$, so $\lim\limits_{r\to-\infty} x = \lim\limits_{\theta\to\pi/2^+} (4\cos\theta + 2) = 2$. Therefore, $\lim\limits_{r\to\pm\infty} x = 2 \;\Rightarrow\; x = 2$ is a vertical asymptote.



50. $y = r\sin\theta = 2\sin\theta - \csc\theta\sin\theta = 2\sin\theta - 1$.

$r \to \infty \;\Rightarrow\; (2 - \csc\theta) \to \infty \;\Rightarrow$ $\csc\theta \to -\infty \;\Rightarrow\; \theta \to \pi^+$ [since we need only consider $0 \le \theta < 2\pi$] and so $\lim\limits_{r\to\infty} y = \lim\limits_{\theta\to\pi^+} 2\sin\theta - 1 = -1$.





Also $r \to -\infty \;\Rightarrow\; (2 - \csc\theta) \to -\infty \;\Rightarrow\; \csc\theta \to \infty \;\Rightarrow\; \theta \to \pi^-$ and so $\lim\limits_{r\to-\infty} x = \lim\limits_{\theta\to\pi^-} 2\sin\theta - 1 = -1$.

Therefore $\lim\limits_{r\to\pm\infty} y = -1 \;\Rightarrow\; y = -1$ is a horizontal asymptote.

51. To show that $x = 1$ is an asymptote we must prove $\lim\limits_{r\to\pm\infty} x = 1$.

$x = (r)\cos\theta = (\sin\theta\tan\theta)\cos\theta = \sin^2\theta$. Now, $r \to \infty \;\Rightarrow\; \sin\theta\tan\theta \to \infty \;\Rightarrow$ $\theta \to \left(\frac{\pi}{2}\right)^-$, so $\lim\limits_{r\to\infty} x = \lim\limits_{\theta\to\pi/2^-} \sin^2\theta = 1$. Also, $r \to -\infty \;\Rightarrow\; \sin\theta\tan\theta \to -\infty \;\Rightarrow$ $\theta \to \left(\frac{\pi}{2}\right)^+$, so $\lim\limits_{r\to-\infty} x = \lim\limits_{\theta\to\pi/2^+} \sin^2\theta = 1$. Therefore, $\lim\limits_{r\to\pm\infty} x = 1 \;\Rightarrow\; x = 1$ is a vertical asymptote. Also notice that $x = \sin^2\theta \ge 0$ for all θ, and $x = \sin^2\theta \le 1$ for all θ. And $x \ne 1$, since the curve is not defined at odd multiples of $\frac{\pi}{2}$. Therefore, the curve lies entirely within the vertical strip $0 \le x < 1$.



52. The equation is $(x^2 + y^2)^3 = 4x^2y^2$, but using polar coordinates we know that $x^2 + y^2 = r^2$ and $x = r\cos\theta$ and $y = r\sin\theta$. Substituting into the given equation: $r^6 = 4r^2\cos^2\theta\, r^2\sin^2\theta \;\Rightarrow\; r^2 = 4\cos^2\theta\sin^2\theta \;\Rightarrow$ $r = \pm 2\cos\theta\sin\theta = \pm\sin 2\theta$. $r = \pm\sin 2\theta$ is sketched at right.



53. (a) We see that the curve $r = 1 + c\sin\theta$ crosses itself at the origin, where $r = 0$ (in fact the inner loop corresponds to negative r-values,) so we solve the equation of the limaçon for $r = 0 \;\Leftrightarrow\; c\sin\theta = -1 \;\Leftrightarrow\; \sin\theta = -1/c$. Now if $|c| < 1$, then this equation has no solution and hence there is no inner loop. But if $c < -1$, then on the interval $(0, 2\pi)$ the equation has the two solutions $\theta = \sin^{-1}(-1/c)$ and $\theta = \pi - \sin^{-1}(-1/c)$, and if $c > 1$, the solutions are $\theta = \pi + \sin^{-1}(1/c)$ and $\theta = 2\pi - \sin^{-1}(1/c)$. In each case, $r < 0$ for θ between the two solutions, indicating a loop.

(b) For $0 < c < 1$, the dimple (if it exists) is characterized by the fact that y has a local maximum at $\theta = \frac{3\pi}{2}$. So we determine for what c-values $\dfrac{d^2y}{d\theta^2}$ is negative at $\theta = \frac{3\pi}{2}$, since by the Second Derivative Test this indicates a maximum:

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$y = r\sin\theta = \sin\theta + c\sin^2\theta \;\Rightarrow\; \dfrac{dy}{d\theta} = \cos\theta + 2c\sin\theta\cos\theta = \cos\theta + c\sin 2\theta \;\Rightarrow\; \dfrac{d^2y}{d\theta^2} = -\sin\theta + 2c\cos 2\theta.$

At $\theta = \frac{3\pi}{2}$, this is equal to $-(-1) + 2c(-1) = 1 - 2c$, which is negative only for $c > \frac{1}{2}$. A similar argument shows that for $-1 < c < 0$, y only has a local minimum at $\theta = \frac{\pi}{2}$ (indicating a dimple) for $c < -\frac{1}{2}$.

54. (a) $r = \ln\theta,\ 1 \le \theta \le 6\pi$. r increases as θ increases and there are almost three full revolutions. The graph must be either III or VI. As θ increases, r grows slowly in VI and quickly in III. Since $r = \ln\theta$ grows slowly, its graph must be VI.

(b) $r = \theta^2,\ 0 \le \theta \le 8\pi$. See part (a). This is graph III.

(c) The graph of $r = \cos 3\theta$ is a three-leaved rose, which is graph II.

(d) Since $-1 \le \cos 3\theta \le 1$, $1 \le 2 + \cos 3\theta \le 3$, so $r = 2 + \cos 3\theta$ is never 0; that is, the curve never intersects the pole. The graph must be I or IV. For $0 \le \theta \le 2\pi$, the graph assumes its minimum r-value of 1 three times, at $\theta = \frac{\pi}{3}$, π, and $\frac{5\pi}{3}$, so it must be graph IV.

(e) $r = \cos(\theta/2)$. For $\theta = 0$, $r = 1$, and as θ increases to π, r decreases to 0. Only graph V satisfies those values.

(f) $r = 2 + \cos(3\theta/2)$. As in part (d), this graph never intersects the pole, so it must be graph I.

55. $r = 2\cos\theta \;\Rightarrow\; x = r\cos\theta = 2\cos^2\theta,\ y = r\sin\theta = 2\sin\theta\cos\theta = \sin 2\theta \;\Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{2\cos 2\theta}{2\cdot 2\cos\theta(-\sin\theta)} = \frac{\cos 2\theta}{-\sin 2\theta} = -\cot 2\theta$$

When $\theta = \dfrac{\pi}{3}$, $\dfrac{dy}{dx} = -\cot\left(2\cdot\dfrac{\pi}{3}\right) = \cot\dfrac{\pi}{3} = \dfrac{1}{\sqrt{3}}$. [*Another method:* Use Equation 3.]

56. $r = 2 + \sin 3\theta \;\Rightarrow\; x = r\cos\theta = (2+\sin 3\theta)\cos\theta,\ y = r\sin\theta = (2+\sin 3\theta)\sin\theta \;\Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{(2+\sin 3\theta)\cos\theta + \sin\theta(3\cos 3\theta)}{(2+\sin 3\theta)(-\sin\theta) + \cos\theta(3\cos 3\theta)}$$

When $\theta = \dfrac{\pi}{4}$, $\dfrac{dy}{dx} = \dfrac{\left(2+\sin\frac{3\pi}{4}\right)\cos\frac{\pi}{4} + \sin\frac{\pi}{4}\left(3\cos\frac{3\pi}{4}\right)}{\left(2+\sin\frac{3\pi}{4}\right)\left(-\sin\frac{\pi}{4}\right) + \cos\frac{\pi}{4}\left(3\cos\frac{3\pi}{4}\right)} = \dfrac{\left(2+\frac{\sqrt{2}}{2}\right)\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}\cdot 3\left(-\frac{\sqrt{2}}{2}\right)}{\left(2+\frac{\sqrt{2}}{2}\right)\left(-\frac{\sqrt{2}}{2}\right) + \frac{\sqrt{2}}{2}\cdot 3\left(-\frac{\sqrt{2}}{2}\right)}$

$$= \frac{\sqrt{2} + \frac{1}{2} - \frac{3}{2}}{-\sqrt{2} - \frac{1}{2} - \frac{3}{2}} = \frac{\sqrt{2}-1}{-\sqrt{2}-2}, \text{ or, equivalently, } 2 - \frac{3}{2}\sqrt{2}.$$

57. $r = 1/\theta \;\Rightarrow\; x = r\cos\theta = (\cos\theta)/\theta,\ y = r\sin\theta = (\sin\theta)/\theta \;\Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\sin\theta(-1/\theta^2) + (1/\theta)\cos\theta}{\cos\theta(-1/\theta^2) - (1/\theta)\sin\theta}\cdot\frac{\theta^2}{\theta^2} = \frac{-\sin\theta + \theta\cos\theta}{-\cos\theta - \theta\sin\theta}$$

When $\theta = \pi$, $\dfrac{dy}{dx} = \dfrac{-0 + \pi(-1)}{-(-1) - \pi(0)} = \dfrac{-\pi}{1} = -\pi$.

58. $r = \cos(\theta/3) \;\Rightarrow\; x = r\cos\theta = \cos(\theta/3)\cos\theta,\ y = r\sin\theta = \cos(\theta/3)\sin\theta \;\Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\cos(\theta/3)\cos\theta + \sin\theta\left(-\frac{1}{3}\sin(\theta/3)\right)}{\cos(\theta/3)(-\sin\theta) + \cos\theta\left(-\frac{1}{3}\sin(\theta/3)\right)}$$

When $\theta = \pi$, $\dfrac{dy}{dx} = \dfrac{\frac{1}{2}(-1) + (0)\left(-\sqrt{3}/6\right)}{\frac{1}{2}(0) + (-1)\left(-\sqrt{3}/6\right)} = \dfrac{-1/2}{\sqrt{3}/6} = -\dfrac{3}{\sqrt{3}} = -\sqrt{3}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $r = \cos 2\theta \;\Rightarrow\; x = r\cos\theta = \cos 2\theta\cos\theta,\ y = r\sin\theta = \cos 2\theta\sin\theta \;\Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\cos 2\theta\cos\theta + \sin\theta\,(-2\sin 2\theta)}{\cos 2\theta\,(-\sin\theta) + \cos\theta\,(-2\sin 2\theta)}$$

When $\theta = \dfrac{\pi}{4}$, $\dfrac{dy}{dx} = \dfrac{0\left(\sqrt{2}/2\right) + \left(\sqrt{2}/2\right)(-2)}{0\left(-\sqrt{2}/2\right) + \left(\sqrt{2}/2\right)(-2)} = \dfrac{-\sqrt{2}}{-\sqrt{2}} = 1.$

60. $r = 1 + 2\cos\theta \;\Rightarrow\; x = r\cos\theta = (1 + 2\cos\theta)\cos\theta,\ y = r\sin\theta = (1 + 2\cos\theta)\sin\theta \;\Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{(1 + 2\cos\theta)\cos\theta + \sin\theta\,(-2\sin\theta)}{(1 + 2\cos\theta)(-\sin\theta) + \cos\theta\,(-2\sin\theta)}$$

When $\theta = \dfrac{\pi}{3}$, $\dfrac{dy}{dx} = \dfrac{2\left(\frac{1}{2}\right) + \left(\sqrt{3}/2\right)\left(-\sqrt{3}\right)}{2\left(-\sqrt{3}/2\right) + \frac{1}{2}\left(-\sqrt{3}\right)} \cdot \dfrac{2}{2} = \dfrac{2 - 3}{-2\sqrt{3} - \sqrt{3}} = \dfrac{-1}{-3\sqrt{3}} = \dfrac{\sqrt{3}}{9}.$

61. $r = 3\cos\theta \;\Rightarrow\; x = r\cos\theta = 3\cos\theta\cos\theta,\ y = r\sin\theta = 3\cos\theta\sin\theta \;\Rightarrow$

$\frac{dy}{d\theta} = -3\sin^2\theta + 3\cos^2\theta = 3\cos 2\theta = 0 \;\Rightarrow\; 2\theta = \frac{\pi}{2}$ or $\frac{3\pi}{2} \;\Leftrightarrow\; \theta = \frac{\pi}{4}$ or $\frac{3\pi}{4}$.

So the tangent is horizontal at $\left(\frac{3}{\sqrt{2}}, \frac{\pi}{4}\right)$ and $\left(-\frac{3}{\sqrt{2}}, \frac{3\pi}{4}\right)$ $\left[\text{same as } \left(\frac{3}{\sqrt{2}}, -\frac{\pi}{4}\right)\right]$.

$\frac{dx}{d\theta} = -6\sin\theta\cos\theta = -3\sin 2\theta = 0 \;\Rightarrow\; 2\theta = 0$ or $\pi \;\Leftrightarrow\; \theta = 0$ or $\frac{\pi}{2}$. So the tangent is vertical at $(3, 0)$ and $\left(0, \frac{\pi}{2}\right)$.

62. $r = 1 - \sin\theta \;\Rightarrow\; x = r\cos\theta = \cos\theta\,(1 - \sin\theta),\ y = r\sin\theta = \sin\theta\,(1 - \sin\theta) \;\Rightarrow$

$\frac{dy}{d\theta} = \sin\theta\,(-\cos\theta) + (1 - \sin\theta)\cos\theta = \cos\theta\,(1 - 2\sin\theta) = 0 \;\Rightarrow\; \cos\theta = 0$ or $\sin\theta = \frac{1}{2} \;\Rightarrow$

$\theta = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}$, or $\frac{3\pi}{2} \;\Rightarrow$ horizontal tangent at $\left(\frac{1}{2}, \frac{\pi}{6}\right)$, $\left(\frac{1}{2}, \frac{5\pi}{6}\right)$, and $\left(2, \frac{3\pi}{2}\right)$.

$$\frac{dx}{d\theta} = \cos\theta\,(-\cos\theta) + (1 - \sin\theta)(-\sin\theta) = -\cos^2\theta - \sin\theta + \sin^2\theta = 2\sin^2\theta - \sin\theta - 1$$
$$= (2\sin\theta + 1)(\sin\theta - 1) = 0 \;\Rightarrow$$

$\sin\theta = -\frac{1}{2}$ or $1 \;\Rightarrow\; \theta = \frac{7\pi}{6}, \frac{11\pi}{6}$, or $\frac{\pi}{2} \;\Rightarrow$ vertical tangent at $\left(\frac{3}{2}, \frac{7\pi}{6}\right)$, $\left(\frac{3}{2}, \frac{11\pi}{6}\right)$, and $\left(0, \frac{\pi}{2}\right)$.

Note that the tangent is vertical, not horizontal, when $\theta = \frac{\pi}{2}$, since

$$\lim_{\theta\to(\pi/2)^-} \frac{dy/d\theta}{dx/d\theta} = \lim_{\theta\to(\pi/2)^-} \frac{\cos\theta\,(1 - 2\sin\theta)}{(2\sin\theta + 1)(\sin\theta - 1)} = \infty \text{ and } \lim_{\theta\to(\pi/2)^+} \frac{dy/d\theta}{dx/d\theta} = -\infty.$$

63. $r = 1 + \cos\theta \;\Rightarrow\; x = r\cos\theta = \cos\theta\,(1 + \cos\theta),\ y = r\sin\theta = \sin\theta\,(1 + \cos\theta) \;\Rightarrow$

$\frac{dy}{d\theta} = (1 + \cos\theta)\cos\theta - \sin^2\theta = 2\cos^2\theta + \cos\theta - 1 = (2\cos\theta - 1)(\cos\theta + 1) = 0 \;\Rightarrow\; \cos\theta = \frac{1}{2}$ or $-1 \;\Rightarrow$

$\theta = \frac{\pi}{3}, \pi$, or $\frac{5\pi}{3} \;\Rightarrow$ horizontal tangent at $\left(\frac{3}{2}, \frac{\pi}{3}\right)$, $(0, \pi)$, and $\left(\frac{3}{2}, \frac{5\pi}{3}\right)$.

$\frac{dx}{d\theta} = -(1 + \cos\theta)\sin\theta - \cos\theta\sin\theta = -\sin\theta\,(1 + 2\cos\theta) = 0 \;\Rightarrow\; \sin\theta = 0$ or $\cos\theta = -\frac{1}{2} \;\Rightarrow$

$\theta = 0, \pi, \frac{2\pi}{3}$, or $\frac{4\pi}{3} \;\Rightarrow$ vertical tangent at $(2, 0)$, $\left(\frac{1}{2}, \frac{2\pi}{3}\right)$, and $\left(\frac{1}{2}, \frac{4\pi}{3}\right)$.

Note that the tangent is horizontal, not vertical when $\theta = \pi$, since $\lim\limits_{\theta\to\pi} \dfrac{dy/d\theta}{dx/d\theta} = 0.$

64. $r = e^\theta \;\Rightarrow\; x = r\cos\theta = e^\theta\cos\theta,\ y = r\sin\theta = e^\theta\sin\theta \;\Rightarrow$

$\frac{dy}{d\theta} = e^\theta\sin\theta + e^\theta\cos\theta = e^\theta(\sin\theta + \cos\theta) = 0 \;\Rightarrow\; \sin\theta = -\cos\theta \;\Rightarrow\; \tan\theta = -1 \;\Rightarrow$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$\theta = -\frac{1}{4}\pi + n\pi$ [n any integer] $\Rightarrow$ horizontal tangents at $\left(e^{\pi(n-1/4)}, \pi\left(n - \frac{1}{4}\right)\right)$.

$\frac{dx}{d\theta} = e^\theta \cos\theta - e^\theta \sin\theta = e^\theta(\cos\theta - \sin\theta) = 0 \Rightarrow \sin\theta = \cos\theta \Rightarrow \tan\theta = 1 \Rightarrow$

$\theta = \frac{1}{4}\pi + n\pi$ [n any integer] $\Rightarrow$ vertical tangents at $\left(e^{\pi(n+1/4)}, \pi\left(n + \frac{1}{4}\right)\right)$.

65. $r = a\sin\theta + b\cos\theta \Rightarrow r^2 = ar\sin\theta + br\cos\theta \Rightarrow x^2 + y^2 = ay + bx \Rightarrow$

$x^2 - bx + \left(\frac{1}{2}b\right)^2 + y^2 - ay + \left(\frac{1}{2}a\right)^2 = \left(\frac{1}{2}b\right)^2 + \left(\frac{1}{2}a\right)^2 \Rightarrow \left(x - \frac{1}{2}b\right)^2 + \left(y - \frac{1}{2}a\right)^2 = \frac{1}{4}(a^2 + b^2)$, and this is a circle with center $\left(\frac{1}{2}b, \frac{1}{2}a\right)$ and radius $\frac{1}{2}\sqrt{a^2 + b^2}$.

66. These curves are circles which intersect at the origin and at $\left(\frac{1}{\sqrt{2}}a, \frac{\pi}{4}\right)$. At the origin, the first circle has a horizontal tangent and the second a vertical one, so the tangents are perpendicular here. For the first circle [$r = a\sin\theta$], $dy/d\theta = a\cos\theta\,\sin\theta + a\sin\theta\,\cos\theta = a\sin 2\theta = a$ at $\theta = \frac{\pi}{4}$ and $dx/d\theta = a\cos^2\theta - a\sin^2\theta = a\cos 2\theta = 0$ at $\theta = \frac{\pi}{4}$, so the tangent here is vertical. Similarly, for the second circle [$r = a\cos\theta$], $dy/d\theta = a\cos 2\theta = 0$ and $dx/d\theta = -a\sin 2\theta = -a$ at $\theta = \frac{\pi}{4}$, so the tangent is horizontal, and again the tangents are perpendicular.

67. $r = 1 + 2\sin(\theta/2)$. The parameter interval is $[0, 4\pi]$.



68. $r = \sqrt{1 - 0.8\sin^2\theta}$. The parameter interval is $[0, 2\pi]$.



69. $r = e^{\sin\theta} - 2\cos(4\theta)$.

The parameter interval is $[0, 2\pi]$.



70. $r = |\tan\theta|^{|\cot\theta|}$.

The parameter interval $[0, \pi]$ produces the heart-shaped valentine curve shown in the first window.

The complete curve, including the reflected heart, is produced by the parameter interval $[0, 2\pi]$, but perhaps you'll agree that the first curve is more appropriate.





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

71. $r = 1 + \cos^{999}\theta$. The parameter interval is $[0, 2\pi]$.



72. $r = 2 + \cos(9\theta/4)$. The parameter interval is $[0, 8\pi]$.



73. It appears that the graph of $r = 1 + \sin\left(\theta - \frac{\pi}{6}\right)$ is the same shape as the graph of $r = 1 + \sin\theta$, but rotated counterclockwise about the origin by $\frac{\pi}{6}$. Similarly, the graph of $r = 1 + \sin\left(\theta - \frac{\pi}{3}\right)$ is rotated by $\frac{\pi}{3}$. In general, the graph of $r = f(\theta - \alpha)$ is the same shape as that of $r = f(\theta)$, but rotated counterclockwise through α about the origin. That is, for any point (r_0, θ_0) on the curve $r = f(\theta)$, the point $(r_0, \theta_0 + \alpha)$ is on the curve $r = f(\theta - \alpha)$, since $r_0 = f(\theta_0) = f((\theta_0 + \alpha) - \alpha)$.



74.



From the graph, the highest points seem to have $y \approx 0.77$. To find the exact value, we solve $dy/d\theta = 0$. $y = r\sin\theta = \sin\theta\,\sin 2\theta \;\Rightarrow$

$$\begin{aligned} dy/d\theta &= 2\sin\theta\,\cos 2\theta + \cos\theta\,\sin 2\theta \\ &= 2\sin\theta\,(2\cos^2\theta - 1) + \cos\theta\,(2\sin\theta\,\cos\theta) \\ &= 2\sin\theta\,(3\cos^2\theta - 1) \end{aligned}$$

In the first quadrant, this is 0 when $\cos\theta = \frac{1}{\sqrt{3}} \;\Leftrightarrow\; \sin\theta = \sqrt{\frac{2}{3}} \;\Leftrightarrow$ $y = 2\sin^2\theta\cos\theta = 2 \cdot \frac{2}{3} \cdot \frac{1}{\sqrt{3}} = \frac{4}{9}\sqrt{3} \approx 0.77$.

75. Consider curves with polar equation $r = 1 + c\cos\theta$, where c is a real number. If $c = 0$, we get a circle of radius 1 centered at the pole. For $0 < c \le 0.5$, the curve gets slightly larger, moves right, and flattens out a bit on the left side. For $0.5 < c < 1$, the left side has a dimple shape. For $c = 1$, the dimple becomes a cusp. For $c > 1$, there is an internal loop. For $c \ge 0$, the rightmost point on the curve is $(1 + c, 0)$. For $c < 0$, the curves are reflections through the vertical axis of the curves with $c > 0$.



$c = 0.25$



$c = 0.75$



$c = 1$



$c = 2$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

76. Consider the polar curves $r = 1 + \cos^n \theta$, where n is a positive integer. First, let **n be an even positive integer**. The first figure shows that the curve has a peanut shape for $n = 2$, but as n increases, the ends are squeezed. As n becomes large, the curves look more and more like the unit circle, but with spikes to the points $(2, 0)$ and $(2, \pi)$.



The second figure shows r as a function of θ in Cartesian coordinates for the same values of n. We can see that for large n, the graph is similar to the graph of $y = 1$, but with spikes to $y = 2$ for $x = 0$, π, and 2π. (Note that when $0 < \cos\theta < 1$, $\cos^{1000}\theta$ is very small.)



Next, let **n be an odd positive integer**. The third figure shows that the curve is a cardioid for $n = 1$, but as n increases, the heart shape becomes more pronounced. As n becomes large, the curves again look more like the unit circle, but with an outward spike to $(2, 0)$ and an inward spike to $(0, \pi)$.



The fourth figure shows r as a function of θ in Cartesian coordinates for the same values of n. We can see that for large n, the graph is similar to the graph of $y = 1$, but spikes to $y = 2$ for $x = 0$ and π, and to $y = 0$ for $x = \pi$.



77. $\tan\psi = \tan(\phi - \theta) = \dfrac{\tan\phi - \tan\theta}{1 + \tan\phi\,\tan\theta} = \dfrac{\dfrac{dy}{dx} - \tan\theta}{1 + \dfrac{dy}{dx}\tan\theta} = \dfrac{\dfrac{dy/d\theta}{dx/d\theta} - \tan\theta}{1 + \dfrac{dy/d\theta}{dx/d\theta}\tan\theta}$

$$= \frac{\dfrac{dy}{d\theta} - \dfrac{dx}{d\theta}\tan\theta}{\dfrac{dx}{d\theta} + \dfrac{dy}{d\theta}\tan\theta} = \frac{\left(\dfrac{dr}{d\theta}\sin\theta + r\cos\theta\right) - \tan\theta\left(\dfrac{dr}{d\theta}\cos\theta - r\sin\theta\right)}{\left(\dfrac{dr}{d\theta}\cos\theta - r\sin\theta\right) + \tan\theta\left(\dfrac{dr}{d\theta}\sin\theta + r\cos\theta\right)} = \frac{r\cos\theta + r\cdot\dfrac{\sin^2\theta}{\cos\theta}}{\dfrac{dr}{d\theta}\cos\theta + \dfrac{dr}{d\theta}\cdot\dfrac{\sin^2\theta}{\cos\theta}}$$

$$= \frac{r\cos^2\theta + r\sin^2\theta}{\dfrac{dr}{d\theta}\cos^2\theta + \dfrac{dr}{d\theta}\sin^2\theta} = \frac{r}{dr/d\theta}$$

78. (a) $r = e^\theta \;\Rightarrow\; dr/d\theta = e^\theta$, so by Exercise 77, $\tan\psi = r/e^\theta = 1 \;\Rightarrow\; \psi = \arctan 1 = \frac{\pi}{4}$.



(b) The Cartesian equation of the tangent line at $(1, 0)$ is $y = x - 1$, and that of the tangent line at $(0, e^{\pi/2})$ is $y = e^{\pi/2} - x$.

(c) Let a be the tangent of the angle between the tangent and radial lines, that is, $a = \tan\psi$. Then, by Exercise 77, $a = \dfrac{r}{dr/d\theta} \;\Rightarrow\; \dfrac{dr}{d\theta} = \dfrac{1}{a}r \;\rightarrow$ $r = Ce^{\theta/a}$ (by Theorem 9.4.2).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

LABORATORY PROJECT Families of Polar Curves

1. (a) $r = \sin n\theta$.

$n = 2$ $n = 3$ $n = 4$ $n = 5$

From the graphs, it seems that when n is even, the number of loops in the curve (called a rose) is $2n$, and when n is odd, the number of loops is simply n. This is because in the case of n odd, every point on the graph is traversed twice, due to the fact that

$$r(\theta + \pi) = \sin[n(\theta + \pi)] = \sin n\theta \cos n\pi + \cos n\theta \sin n\pi = \begin{cases} \sin n\theta & \text{if } n \text{ is even} \\ -\sin n\theta & \text{if } n \text{ is odd} \end{cases}$$

(b) The graph of $r = |\sin n\theta|$ has $2n$ loops whether n is odd or even, since $r(\theta + \pi) = r(\theta)$.



$n = 2$ $n = 3$ $n = 4$ $n = 5$

2. $r = 1 + c \sin n\theta$. We vary n while keeping c constant at 2. As n changes, the curves change in the same way as those in Exercise 1: the number of loops increases. Note that if n is even, the smaller loops are outside the larger ones; if n is odd, they are inside.

$c = 2$



$n = 2$ $n = 3$ $n = 4$ $n = 5$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Now we vary c while keeping $n = 3$. As c increases toward 0, the entire graph gets smaller (the graphs below are not to scale) and the smaller loops shrink in relation to the large ones. At $c = -1$, the small loops disappear entirely, and for $-1 < c < 1$, the graph is a simple, closed curve (at $c = 0$ it is a circle). As c continues to increase, the same changes are seen, but in reverse order, since $1 + (-c)\sin n\theta = 1 + c\sin n(\theta + \pi)$, so the graph for $c = c_0$ is the same as that for $c = -c_0$, with a rotation through π. As $c \to \infty$, the smaller loops get relatively closer in size to the large ones. Note that the distance between the outermost points of corresponding inner and outer loops is always 2. Maple's `animate` command (or Mathematica's `Animate`) is very useful for seeing the changes that occur as c varies.

$n = 3$



$c = -4$ $c = -1.4$ $c = -1$ $c = -0.8$



$c = -0.2$ $c = 0$ $c = 0.5$ $c = 8$

3. $r = \dfrac{1 - a\cos\theta}{1 + a\cos\theta}$. We start with $a = 0$, since in this case the curve is simply the circle $r = 1$.

As a increases, the graph moves to the left, and its right side becomes flattened. As a increases through about 0.4, the right side seems to grow a dimple, which upon closer investigation (with narrower θ-ranges) seems to appear at $a \approx 0.42$ [the actual value is $\sqrt{2} - 1$]. As $a \to 1$, this dimple becomes more pronounced, and the curve begins to stretch out horizontally, until at $a = 1$ the denominator vanishes at $\theta = \pi$, and the dimple becomes an actual cusp. For $a > 1$ we must choose our parameter interval carefully, since $r \to \infty$ as $1 + a\cos\theta \to 0 \quad \Leftrightarrow \quad \theta \to \pm\cos^{-1}(-1/a)$. As a increases from 1, the curve splits into two parts. The left part has a loop, which grows larger as a increases, and the right part grows broader vertically, and its left tip develops a dimple when $a \approx 2.42$ [actually, $\sqrt{2} + 1$]. As a increases, the dimple grows more and more pronounced. If $a < 0$, we get the same graph as we do for the corresponding positive a-value, but with a rotation through π about the pole, as happened when c was replaced with $-c$ in Exercise 2.

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.



$a = 0$



$a = 0.3$



$a = 0.41, |\theta| \le 0.5$



$a = 0.42, |\theta| \le 0.5$



$a = 0.9, |\theta| \le 0.5$



$a = 1, |\theta| \le 0.1$



$a = 2$



$a = 2.41, |\theta - \pi| \le 0.2$



$a = 2.42, |\theta - \pi| \le 0.2$



$a = 4$

4. Most graphing devices cannot plot implicit polar equations, so we must first find an explicit expression (or expressions) for r in terms of θ, a, and c. We note that the given equation, $r^4 - 2c^2r^2\cos 2\theta + c^4 - a^4 = 0$, is a quadratic in r^2, so we use the quadratic formula and find that

$$r^2 = \frac{2c^2\cos 2\theta \pm \sqrt{4c^4\cos^2 2\theta - 4(c^4 - a^4)}}{2} = c^2\cos 2\theta \pm \sqrt{a^4 - c^4\sin^2 2\theta}$$

so $r = \pm\sqrt{c^2\cos 2\theta \pm \sqrt{a^4 - c^4\sin^2 2\theta}}$. So for each graph, we must plot four curves to be sure of plotting all the points which satisfy the given equation. Note that all four functions have period π.

We start with the case $a = c = 1$, and the resulting curve resembles the symbol for infinity. If we let a decrease, the curve splits into two symmetric parts, and as a decreases further, the parts become smaller, further apart, and rounder. If instead we let a increase from 1, the two lobes of the curve join together, and as a increases further they continue to merge, until at

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$a \approx 1.4$, the graph no longer has dimples, and has an oval shape. As $a \to \infty$, the oval becomes larger and rounder, since the c^2 and c^4 terms lose their significance. Note that the shape of the graph seems to depend only on the ratio c/a, while the size of the graph varies as c and a jointly increase.



10.4 Areas and Lengths in Polar Coordinates

1. $r = e^{-\theta/4}$, $\pi/2 \le \theta \le \pi$.

$$A = \int_{\pi/2}^{\pi} \tfrac{1}{2} r^2 \, d\theta = \int_{\pi/2}^{\pi} \tfrac{1}{2} (e^{-\theta/4})^2 \, d\theta = \int_{\pi/2}^{\pi} \tfrac{1}{2} e^{-\theta/2} \, d\theta = \tfrac{1}{2} \left[-2e^{-\theta/2} \right]_{\pi/2}^{\pi} = -1(e^{-\pi/2} - e^{-\pi/4}) = e^{-\pi/4} - e^{-\pi/2}$$

2. $r = \cos\theta$, $0 \le \theta \le \pi/6$.

$$A = \int_0^{\pi/6} \tfrac{1}{2} r^2 \, d\theta = \int_0^{\pi/6} \tfrac{1}{2} \cos^2\theta \, d\theta = \tfrac{1}{2} \int_0^{\pi/6} \tfrac{1}{2} (1 + \cos 2\theta) \, d\theta = \tfrac{1}{4} \left[\theta + \tfrac{1}{2} \sin 2\theta \right]_0^{\pi/6}$$
$$= \tfrac{1}{4} \left(\tfrac{\pi}{6} + \tfrac{1}{2} \cdot \tfrac{1}{2} \sqrt{3} \right) = \tfrac{\pi}{24} + \tfrac{1}{16} \sqrt{3}$$

3. $r = \sin\theta + \cos\theta$, $0 \le \theta \le \pi$.

$$A = \int_0^{\pi} \frac{1}{2} r^2 \, d\theta = \int_0^{\pi} \frac{1}{2} (\sin\theta + \cos\theta)^2 \, d\theta = \int_0^{\pi} \frac{1}{2} (\sin^2\theta + 2\sin\theta\cos\theta + \cos^2\theta) \, d\theta = \int_0^{\pi} \frac{1}{2} (1 + \sin 2\theta) \, d\theta$$
$$= \tfrac{1}{2} \left[\theta - \tfrac{1}{2} \cos 2\theta \right]_0^{\pi} = \tfrac{1}{2} \left[\left(\pi - \tfrac{1}{2} \right) - \left(0 - \tfrac{1}{2} \right) \right] = \tfrac{\pi}{2}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $r = 1/\theta,\ \pi/2 \le \theta \le 2\pi.$

$$A = \int_{\pi/2}^{2\pi} \frac{1}{2} r^2\, d\theta = \int_{\pi/2}^{2\pi} \frac{1}{2}\left(\frac{1}{\theta}\right)^2 d\theta = \int_{\pi/2}^{2\pi} \frac{1}{2}\theta^{-2}\, d\theta = \frac{1}{2}\left[-\frac{1}{\theta}\right]_{\pi/2}^{2\pi}$$
$$= \tfrac{1}{2}\left(-\tfrac{1}{2\pi} + \tfrac{2}{\pi}\right) = \tfrac{1}{2}\left(-\tfrac{1}{2\pi} + \tfrac{4}{2\pi}\right) = \tfrac{3}{4\pi}$$

5. $r^2 = \sin 2\theta,\ 0 \le \theta \le \pi/2.$

$$A = \int_0^{\pi/2} \tfrac{1}{2} r^2\, d\theta = \int_0^{\pi/2} \tfrac{1}{2}\sin 2\theta\, d\theta = \left[-\tfrac{1}{4}\cos 2\theta\right]_0^{\pi/2} = -\tfrac{1}{4}(\cos\pi - \cos 0) = -\tfrac{1}{4}(-1-1) = \tfrac{1}{2}$$

6. $r = 2 + \cos\theta,\ \pi/2 \le \theta \le \pi.$

$$A = \int_{\pi/2}^{\pi} \tfrac{1}{2} r^2\, d\theta = \int_{\pi/2}^{\pi} \tfrac{1}{2}(2+\cos\theta)^2\, d\theta = \int_{\pi/2}^{\pi} \tfrac{1}{2}(4 + 4\cos\theta + \cos^2\theta)\, d\theta = \int_{\pi/2}^{\pi} \tfrac{1}{2}[4 + 4\cos\theta + \tfrac{1}{2}(1+\cos 2\theta)]\, d\theta$$
$$= \int_{\pi/2}^{\pi} \left(\tfrac{9}{4} + 2\cos\theta + \tfrac{1}{4}\cos 2\theta\right) d\theta = \left[\tfrac{9}{4}\theta + 2\sin\theta + \tfrac{1}{8}\sin 2\theta\right]_{\pi/2}^{\pi} = \left(\tfrac{9\pi}{4} + 0 + 0\right) - \left(\tfrac{9\pi}{8} + 2 + 0\right) = \tfrac{9\pi}{8} - 2$$

7. $r = 4 + 3\sin\theta,\ -\tfrac{\pi}{2} \le \theta \le \tfrac{\pi}{2}.$

$$A = \int_{-\pi/2}^{\pi/2} \tfrac{1}{2}((4+3\sin\theta)^2\, d\theta = \tfrac{1}{2}\int_{-\pi/2}^{\pi/2} (16 + 24\sin\theta + 9\sin^2\theta)\, d\theta$$
$$= \tfrac{1}{2}\int_{-\pi/2}^{\pi/2} (16 + 9\sin^2\theta)\, d\theta \qquad \text{[by Theorem 4.5.6(b) [ET 5.5.7(b)]]}$$
$$= \tfrac{1}{2}\cdot 2\int_0^{\pi/2} \left[16 + 9\cdot\tfrac{1}{2}(1-\cos 2\theta)\right] d\theta \qquad \text{[by Theorem 4.5.6(a) [ET 5.5.7(a)]]}$$
$$= \int_0^{\pi/2} \left(\tfrac{41}{2} - \tfrac{9}{2}\cos 2\theta\right) d\theta = \left[\tfrac{41}{2}\theta - \tfrac{9}{4}\sin 2\theta\right]_0^{\pi/2} = \left(\tfrac{41\pi}{4} - 0\right) - (0-0) = \tfrac{41\pi}{4}$$

8. $r = \sqrt{\ln\theta},\ 1 \le \theta \le 2\pi.$

$$A = \int_1^{2\pi} \tfrac{1}{2}\left(\sqrt{\ln\theta}\right)^2 d\theta = \int_1^{2\pi} \tfrac{1}{2}\ln\theta\, d\theta = \left[\tfrac{1}{2}\theta\ln\theta\right]_1^{2\pi} - \int_1^{2\pi} \tfrac{1}{2}\, d\theta \qquad \left[\begin{array}{ll} u = \ln\theta, & dv = \frac{1}{2}\, d\theta \\ du = (1/\theta)\, d\theta, & v = \frac{1}{2}\theta \end{array}\right]$$
$$= [\pi\ln(2\pi) - 0] - \left[\tfrac{1}{2}\theta\right]_1^{2\pi} = \pi\ln(2\pi) - \pi + \tfrac{1}{2}$$

9. The area is bounded by $r = 2\sin\theta$ for $\theta = 0$ to $\theta = \pi$.

$$A = \int_0^{\pi} \tfrac{1}{2} r^2\, d\theta = \tfrac{1}{2}\int_0^{\pi} (2\sin\theta)^2\, d\theta = \tfrac{1}{2}\int_0^{\pi} 4\sin^2\theta\, d\theta$$
$$= 2\int_0^{\pi} \tfrac{1}{2}(1-\cos 2\theta) d\theta = \left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\pi} = \pi$$

Also, note that this is a circle with radius 1, so its area is $\pi(1)^2 = \pi$.



10. $A = \int_0^{2\pi} \tfrac{1}{2} r^2\, d\theta = \int_0^{2\pi} \tfrac{1}{2}(1-\sin\theta)^2\, d\theta$

$$= \tfrac{1}{2}\int_0^{2\pi} (1 - 2\sin\theta + \sin^2\theta)\, d\theta = \tfrac{1}{2}\int_0^{2\pi} \left[1 - 2\sin\theta + \tfrac{1}{2}(1-\cos 2\theta)\right] d\theta$$
$$= \tfrac{1}{2}\int_0^{2\pi} \left(\tfrac{3}{2} - 2\sin\theta - \tfrac{1}{2}\cos 2\theta\right) d\theta = \tfrac{1}{2}\left[\tfrac{3}{2}\theta + 2\cos\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}$$
$$= \tfrac{1}{2}[(3\pi + 2) - (2)] = \tfrac{3\pi}{2}$$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $A = \int_0^{2\pi} \frac{1}{2}r^2\,d\theta = \int_0^{2\pi} \frac{1}{2}(3+2\cos\theta)^2\,d\theta = \frac{1}{2}\int_0^{2\pi} (9+12\cos\theta+4\cos^2\theta)\,d\theta$

$= \frac{1}{2}\int_0^{2\pi} \left[9+12\cos\theta+4\cdot\frac{1}{2}(1+\cos 2\theta)\right] d\theta$

$= \frac{1}{2}\int_0^{2\pi} (11+12\cos\theta+2\cos 2\theta)\,d\theta = \frac{1}{2}\left[11\theta+12\sin\theta+\sin 2\theta\right]_0^{2\pi}$

$= \frac{1}{2}(22\pi) = 11\pi$



12. $A = \int_0^{2\pi} \frac{1}{2}r^2\,d\theta = \int_0^{2\pi} \frac{1}{2}(2-\cos\theta)^2\,d\theta = \int_0^{2\pi} \frac{1}{2}(4-4\cos\theta+\cos^2\theta)\,d\theta$

$= \int_0^{2\pi} \frac{1}{2}\left[4-4\cos\theta+\frac{1}{2}(1+\cos 2\theta)\right] d\theta = \int_0^{2\pi} \left(\frac{9}{4}-2\cos\theta+\frac{1}{4}\cos 2\theta\right) d\theta$

$= \left[\frac{9}{4}\theta-2\sin\theta+\frac{1}{8}\sin 2\theta\right]_0^{2\pi} = \left(\frac{9\pi}{2}-0+0\right)-(0-0+0) = \frac{9\pi}{2}$



13. $A = \int_0^{2\pi} \frac{1}{2}r^2\,d\theta = \int_0^{2\pi} \frac{1}{2}(2+\sin 4\theta)^2\,d\theta = \frac{1}{2}\int_0^{2\pi} (4+4\sin 4\theta+\sin^2 4\theta)\,d\theta$

$= \frac{1}{2}\int_0^{2\pi} \left[4+4\sin 4\theta+\frac{1}{2}(1-\cos 8\theta)\right] d\theta$

$= \frac{1}{2}\int_0^{2\pi} \left(\frac{9}{2}+4\sin 4\theta-\frac{1}{2}\cos 8\theta\right) d\theta = \frac{1}{2}\left[\frac{9}{2}\theta-\cos 4\theta-\frac{1}{16}\sin 8\theta\right]_0^{2\pi}$

$= \frac{1}{2}[(9\pi-1)-(-1)] = \frac{9}{2}\pi$



14. $A = \int_0^{2\pi} \frac{1}{2}r^2\,d\theta = \int_0^{2\pi} \frac{1}{2}(3-2\cos 4\theta)^2\,d\theta = \frac{1}{2}\int_0^{2\pi} (9-12\cos 4\theta+4\cos^2 4\theta)\,d\theta$

$= \frac{1}{2}\int_0^{2\pi} \left[9-12\cos 4\theta+4\cdot\frac{1}{2}(1+\cos 8\theta)\right] d\theta$

$= \frac{1}{2}\int_0^{2\pi} (11-12\cos 4\theta+2\cos 8\theta)\,d\theta = \frac{1}{2}\left[11\theta-3\sin 4\theta+\frac{1}{4}\sin 8\theta\right]_0^{2\pi}$

$= \frac{1}{2}(22\pi) = 11\pi$



15. $A = \int_0^{2\pi} \frac{1}{2}r^2\,d\theta = \int_0^{2\pi} \frac{1}{2}\left(\sqrt{1+\cos^2 5\theta}\right)^2 d\theta$

$= \frac{1}{2}\int_0^{2\pi} (1+\cos^2 5\theta)\,d\theta = \frac{1}{2}\int_0^{2\pi} \left[1+\frac{1}{2}(1+\cos 10\theta)\right] d\theta$

$= \frac{1}{2}\left[\frac{3}{2}\theta+\frac{1}{20}\sin 10\theta\right]_0^{2\pi} = \frac{1}{2}(3\pi) = \frac{3}{2}\pi$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. $A = \int_0^{2\pi} \frac{1}{2}r^2\,d\theta = \int_0^{2\pi} \frac{1}{2}(1+5\sin 6\theta)^2\,d\theta$

$= \frac{1}{2}\int_0^{2\pi} (1 + 10\sin 6\theta + 25\sin^2 6\theta)\,d\theta$

$= \frac{1}{2}\int_0^{2\pi} \left[1 + 10\sin 6\theta + 25\cdot\tfrac{1}{2}(1-\cos 12\theta)\right] d\theta$

$= \frac{1}{2}\int_0^{2\pi} \left[\tfrac{27}{2} + 10\sin 6\theta - \tfrac{25}{2}\cos 12\theta\right] d\theta = \frac{1}{2}\left[\tfrac{27}{2}\theta - \tfrac{5}{3}\cos 6\theta - \tfrac{25}{24}\sin 12\theta\right]_0^{2\pi}$

$= \frac{1}{2}\left[\left(27\pi - \tfrac{5}{3}\right) - \left(-\tfrac{5}{3}\right)\right] = \tfrac{27}{2}\pi$



17. The curve passes through the pole when $r = 0 \;\Rightarrow\; 4\cos 3\theta = 0 \;\Rightarrow\; \cos 3\theta = 0 \;\Rightarrow\; 3\theta = \frac{\pi}{2} + \pi n \;\Rightarrow$ $\theta = \frac{\pi}{6} + \frac{\pi}{3}n$. The part of the shaded loop above the polar axis is traced out for $\theta = 0$ to $\theta = \pi/6$, so we'll use $-\pi/6$ and $\pi/6$ as our limits of integration.

$A = \int_{-\pi/6}^{\pi/6} \frac{1}{2}(4\cos 3\theta)^2\,d\theta = 2\int_0^{\pi/6} \frac{1}{2}(16\cos^2 3\theta)\,d\theta$

$= 16\int_0^{\pi/6} \frac{1}{2}(1+\cos 6\theta)\,d\theta = 8\left[\theta + \frac{1}{6}\sin 6\theta\right]_0^{\pi/6} = 8\left(\frac{\pi}{6}\right) = \frac{4}{3}\pi$



18. The curve given by $r^2 = 4\cos 2\theta$ passes through the pole when $r = 0 \;\Rightarrow\; 4\cos 2\theta = 0 \;\Rightarrow\; \cos 2\theta = 0 \;\Rightarrow$ $2\theta = \frac{\pi}{2} + \pi n \;\Rightarrow\; \theta = \frac{\pi}{4} + \frac{\pi}{2}n$. The part of the shaded loop above the polar axis is traced out for $\theta = 0$ to $\theta = \pi/4$, so we'll use $-\pi/4$ to $\pi/4$ as our limits of integration.

$A = \int_{-\pi/4}^{\pi/4} \frac{1}{2}(4\cos 2\theta)\,d\theta = 2\int_0^{\pi/4} 2\cos 2\theta\,d\theta = 2\Big[\sin 2\theta\Big]_0^{\pi/4}$

$= 2\sin\frac{\pi}{2} = 2(1) = 2$



19. $r = 0 \;\Rightarrow\; \sin 4\theta = 0 \;\Rightarrow\; 4\theta = \pi n \;\Rightarrow\; \theta = \frac{\pi}{4}n.$

$A = \int_0^{\pi/4} \frac{1}{2}(\sin 4\theta)^2\,d\theta = \frac{1}{2}\int_0^{\pi/4} \sin^2 4\theta\,d\theta = \frac{1}{2}\int_0^{\pi/4} \frac{1}{2}(1-\cos 8\theta)\,d\theta$

$= \frac{1}{4}\left[\theta - \frac{1}{8}\sin 8\theta\right]_0^{\pi/4} = \frac{1}{4}\left(\frac{\pi}{4}\right) = \frac{1}{16}\pi$



20. $r = 0 \;\Rightarrow\; 2\sin 5\theta = 0 \;\Rightarrow\; \sin 5\theta = 0 \;\Rightarrow\; 5\theta = \pi n \;\Rightarrow\; \theta = \frac{\pi}{5}n.$

$A = \int_0^{\pi/5} \frac{1}{2}(2\sin 5\theta)^2\,d\theta = \frac{1}{2}\int_0^{\pi/5} 4\sin^2 5\theta\,d\theta$

$= 2\int_0^{\pi/5} \frac{1}{2}(1-\cos 10\theta)\,d\theta = \left[\theta - \frac{1}{10}\sin 10\theta\right]_0^{\pi/5} = \frac{\pi}{5}$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

21.





This is a limaçon, with inner loop traced out between $\theta = \frac{7\pi}{6}$ and $\frac{11\pi}{6}$ [found by solving $r = 0$].

$$A = 2\int_{7\pi/6}^{3\pi/2} \tfrac{1}{2}(1+2\sin\theta)^2\,d\theta = \int_{7\pi/6}^{3\pi/2} \left(1+4\sin\theta+4\sin^2\theta\right)d\theta = \int_{7\pi/6}^{3\pi/2} \left[1+4\sin\theta+4\cdot\tfrac{1}{2}(1-\cos 2\theta)\right]d\theta$$

$$= \left[\theta - 4\cos\theta + 2\theta - \sin 2\theta\right]_{7\pi/6}^{3\pi/2} = \left(\tfrac{9\pi}{2}\right) - \left(\tfrac{7\pi}{2} + 2\sqrt{3} - \tfrac{\sqrt{3}}{2}\right) = \pi - \tfrac{3\sqrt{3}}{2}$$

22. To determine when the strophoid $r = 2\cos\theta - \sec\theta$ passes through the pole, we solve

$$r = 0 \;\Rightarrow\; 2\cos\theta - \frac{1}{\cos\theta} = 0 \;\Rightarrow\; 2\cos^2\theta - 1 = 0 \;\Rightarrow\; \cos^2\theta = \frac{1}{2} \;\Rightarrow$$

$$\cos\theta = \pm\frac{1}{\sqrt{2}} \;\Rightarrow\; \theta = \tfrac{\pi}{4} \text{ or } \theta = \tfrac{3\pi}{4} \text{ for } 0 \le \theta \le \pi \text{ with } \theta \ne \tfrac{\pi}{2}.$$



$$A = 2\int_0^{\pi/4} \tfrac{1}{2}(2\cos\theta - \sec\theta)^2\,d\theta = \int_0^{\pi/4} (4\cos^2\theta - 4 + \sec^2\theta)\,d\theta$$

$$= \int_0^{\pi/4} \left[4\cdot\tfrac{1}{2}(1+\cos 2\theta) - 4 + \sec^2\theta\right]d\theta = \int_0^{\pi/4} (-2 + 2\cos 2\theta + \sec^2\theta)\,d\theta$$

$$= \left[-2\theta + \sin 2\theta + \tan\theta\right]_0^{\pi/4} = \left(-\tfrac{\pi}{2} + 1 + 1\right) - 0 = 2 - \tfrac{\pi}{2}$$

23. $4\sin\theta = 2 \;\Rightarrow\; \sin\theta = \frac{1}{2} \;\Rightarrow\; \theta = \frac{\pi}{6} \text{ or } \frac{5\pi}{6} \;\Rightarrow$

$$A = \int_{\pi/6}^{5\pi/6} \tfrac{1}{2}\left[(4\sin\theta)^2 - 2^2\right]d\theta = 2\int_{\pi/6}^{\pi/2} \tfrac{1}{2}(16\sin^2\theta - 4)\,d\theta$$

$$= \int_{\pi/6}^{\pi/2} \left[16\cdot\tfrac{1}{2}(1-\cos 2\theta) - 4\right]d\theta = \int_{\pi/6}^{\pi/2} (4 - 8\cos 2\theta)\,d\theta$$

$$= \left[4\theta - 4\sin 2\theta\right]_{\pi/6}^{\pi/2} = (2\pi - 0) - \left(\tfrac{2\pi}{3} - 2\sqrt{3}\right) = \tfrac{4\pi}{3} + 2\sqrt{3}$$



24. $1 - \sin\theta = 1 \;\Rightarrow\; \sin\theta = 0 \;\Rightarrow\; \theta = 0 \text{ or } \pi \;\Rightarrow$

$$A = \int_{\pi}^{2\pi} \tfrac{1}{2}\left[(1-\sin\theta)^2 - 1\right]d\theta = \tfrac{1}{2}\int_{\pi}^{2\pi} (\sin^2\theta - 2\sin\theta)\,d\theta$$

$$-\tfrac{1}{4}\int_{\pi}^{2\pi} (1 - \cos 2\theta - 4\sin\theta)\,d\theta = \tfrac{1}{4}\left[\theta - \tfrac{1}{2}\sin 2\theta + 4\cos\theta\right]_{\pi}^{2\pi}$$

$$-\tfrac{1}{4}\pi + 2$$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. To find the area inside the leminiscate $r^2 = 8\cos 2\theta$ and outside the circle $r = 2$, we first note that the two curves intersect when $r^2 = 8\cos 2\theta$ and $r = 2$, that is, when $\cos 2\theta = \frac{1}{2}$. For $-\pi < \theta \le \pi$, $\cos 2\theta = \frac{1}{2} \Leftrightarrow 2\theta = \pm\pi/3$ or $\pm 5\pi/3 \Leftrightarrow \theta = \pm\pi/6$ or $\pm 5\pi/6$. The figure shows that the desired area is 4 times the area between the curves from 0 to $\pi/6$. Thus,



$$A = 4\int_0^{\pi/6} \left[\tfrac{1}{2}(8\cos 2\theta) - \tfrac{1}{2}(2)^2\right] d\theta = 8\int_0^{\pi/6} (2\cos 2\theta - 1)\, d\theta$$

$$= 8\left[\sin 2\theta - \theta\right]_0^{\pi/6} = 8\left(\sqrt{3}/2 - \pi/6\right) = 4\sqrt{3} - 4\pi/3$$

26. $3\sin\theta = 1 + \sin\theta \Rightarrow \sin\theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{6}$ or $\frac{5\pi}{6} \Rightarrow$



$$A = \int_{\pi/6}^{5\pi/6} \tfrac{1}{2}[(3\sin\theta)^2 - (1+\sin\theta)^2]\, d\theta$$

$$= 2\int_{\pi/6}^{\pi/2} \tfrac{1}{2}(9\sin^2\theta - 1 - 2\sin\theta - \sin^2\theta)\, d\theta$$

$$= \int_{\pi/6}^{\pi/2} (8\sin^2\theta - 1 - 2\sin\theta)\, d\theta$$

$$= \int_{\pi/6}^{\pi/2} \left[8\cdot\tfrac{1}{2}(1-\cos 2\theta) - 1 - 2\sin\theta\right] d\theta = \int_{\pi/6}^{\pi/2} (3 - 4\cos 2\theta - 2\sin\theta)\, d\theta$$

$$= \left[3\theta - 2\sin 2\theta + 2\cos\theta\right]_{\pi/6}^{\pi/2} = \left(\tfrac{3\pi}{2} - 0 + 0\right) - \left(\tfrac{\pi}{2} - \sqrt{3} + \sqrt{3}\right) = \pi$$

27. $3\cos\theta = 1 + \cos\theta \Leftrightarrow \cos\theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$ or $-\frac{\pi}{3}$.



$$A = 2\int_0^{\pi/3} \tfrac{1}{2}[(3\cos\theta)^2 - (1+\cos\theta)^2]\, d\theta$$

$$= \int_0^{\pi/3} (8\cos^2\theta - 2\cos\theta - 1)\, d\theta = \int_0^{\pi/3} [4(1+\cos 2\theta) - 2\cos\theta - 1]\, d\theta$$

$$= \int_0^{\pi/3} (3 + 4\cos 2\theta - 2\cos\theta)\, d\theta = \left[3\theta + 2\sin 2\theta - 2\sin\theta\right]_0^{\pi/3}$$

$$= \pi + \sqrt{3} - \sqrt{3} = \pi$$

28. $3\sin\theta = 2 - \sin\theta \Rightarrow 4\sin\theta = 2 \Rightarrow \sin\theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{6}$ or $\frac{5\pi}{6}$.



$$A = 2\int_{\pi/6}^{\pi/2} \tfrac{1}{2}[(3\sin\theta)^2 - (2-\sin\theta)^2]\, d\theta$$

$$= \int_{\pi/6}^{\pi/2} (9\sin^2\theta - 4 + 4\sin\theta - \sin^2\theta]\, d\theta$$

$$= \int_{\pi/6}^{\pi/2} (8\sin^2\theta + 4\sin\theta - 4)\, d\theta$$

$$= 4\int_{\pi/6}^{\pi/2} \left[2\cdot\tfrac{1}{2}(1-\cos 2\theta) + \sin\theta - 1\right] d\theta$$

$$= 4\int_{\pi/6}^{\pi/2} (\sin\theta - \cos 2\theta)\, d\theta = 4\left[-\cos\theta - \tfrac{1}{2}\sin 2\theta\right]_{\pi/6}^{\pi/2}$$

$$= 4\left[(0-0) - \left(-\tfrac{\sqrt{3}}{2} - \tfrac{\sqrt{3}}{4}\right)\right] = 4\left(\tfrac{3\sqrt{3}}{4}\right) = 3\sqrt{3}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

29. $3\sin\theta = 3\cos\theta \;\Rightarrow\; \dfrac{3\sin\theta}{3\cos\theta} = 1 \;\Rightarrow\; \tan\theta = 1 \;\Rightarrow\; \theta = \frac{\pi}{4} \;\Rightarrow$

$$A = 2\int_0^{\pi/4} \tfrac{1}{2}(3\sin\theta)^2\,d\theta = \int_0^{\pi/4} 9\sin^2\theta\,d\theta = \int_0^{\pi/4} 9\cdot\tfrac{1}{2}(1-\cos 2\theta)\,d\theta$$

$$= \int_0^{\pi/4} \left(\tfrac{9}{2} - \tfrac{9}{2}\cos 2\theta\right) d\theta = \left[\tfrac{9}{2}\theta - \tfrac{9}{4}\sin 2\theta\right]_0^{\pi/4} = \left(\tfrac{9\pi}{8} - \tfrac{9}{4}\right) - (0-0)$$

$$= \tfrac{9\pi}{8} - \tfrac{9}{4}$$



30. $A = 4\int_0^{\pi/2} \tfrac{1}{2}(1-\cos\theta)^2\,d\theta = 2\int_0^{\pi/2}(1 - 2\cos\theta + \cos^2\theta)\,d\theta$

$$= 2\int_0^{\pi/2}\left[1 - 2\cos\theta + \tfrac{1}{2}(1+\cos 2\theta)\right] d\theta$$

$$= 2\int_0^{\pi/2}\left(\tfrac{3}{2} - 2\cos\theta + \tfrac{1}{2}\cos 2\theta\right) d\theta = \int_0^{\pi/2}(3 - 4\cos\theta + \cos 2\theta)\,d\theta$$

$$= \left[3\theta - 4\sin\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/2} = \tfrac{3\pi}{2} - 4$$



31. $\sin 2\theta = \cos 2\theta \;\Rightarrow\; \dfrac{\sin 2\theta}{\cos 2\theta} = 1 \;\Rightarrow\; \tan 2\theta = 1 \;\Rightarrow\; 2\theta = \frac{\pi}{4} \;\Rightarrow$

$\theta = \frac{\pi}{8} \;\Rightarrow$

$$A = 8\cdot 2\int_0^{\pi/8} \tfrac{1}{2}\sin^2 2\theta\,d\theta = 8\int_0^{\pi/8} \tfrac{1}{2}(1-\cos 4\theta)\,d\theta$$

$$= 4\left[\theta - \tfrac{1}{4}\sin 4\theta\right]_0^{\pi/8} = 4\left(\tfrac{\pi}{8} - \tfrac{1}{4}\cdot 1\right) = \tfrac{\pi}{2} - 1$$



32. $3 + 2\cos\theta = 3 + 2\sin\theta \;\Rightarrow\; \cos\theta = \sin\theta \;\Rightarrow\; \theta = \frac{\pi}{4}$ or $\frac{5\pi}{4}$.

$$A = 2\int_{\pi/4}^{5\pi/4} \tfrac{1}{2}(3+2\cos\theta)^2\,d\theta = \int_{\pi/4}^{5\pi/4}(9 + 12\cos\theta + 4\cos^2\theta)\,d\theta$$

$$= \int_{\pi/4}^{5\pi/4}\left[9 + 12\cos\theta + 4\cdot\tfrac{1}{2}(1+\cos 2\theta)\right] d\theta$$

$$= \int_{\pi/4}^{5\pi/4}(11 + 12\cos\theta + 2\cos 2\theta)\,d\theta = \left[11\theta + 12\sin\theta + \sin 2\theta\right]_{\pi/4}^{5\pi/4}$$

$$= \left(\tfrac{55\pi}{4} - 6\sqrt{2} + 1\right) - \left(\tfrac{11\pi}{4} + 6\sqrt{2} + 1\right) = 11\pi - 12\sqrt{2}$$



33. From the figure, we see that the shaded region is 4 times the shaded region from $\theta = 0$ to $\theta = \pi/4$. $r^2 = 2\sin 2\theta$ and $r = 1 \;\Rightarrow$

$2\sin 2\theta = 1^2 \;\Rightarrow\; \sin 2\theta = \frac{1}{2} \;\Rightarrow\; 2\theta = \frac{\pi}{6} \;\Rightarrow\; \theta = \frac{\pi}{12}$.

$$A = 4\int_0^{\pi/12} \tfrac{1}{2}(2\sin 2\theta)\,d\theta + 4\int_{\pi/12}^{\pi/4} \tfrac{1}{2}(1)^2\,d\theta$$

$$= \int_0^{\pi/12} 4\sin 2\theta\,d\theta + \int_{\pi/12}^{\pi/4} 2\,d\theta = \left[-2\cos 2\theta\right]_0^{\pi/12} + \left[2\theta\right]_{\pi/12}^{\pi/4}$$

$$= \left(-\sqrt{3} + 2\right) + \left(\tfrac{\pi}{2} - \tfrac{\pi}{6}\right) = -\sqrt{3} + 2 + \tfrac{\pi}{3}$$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. Let $\alpha = \tan^{-1}(b/a)$. Then



$$\begin{aligned} A &= \int_0^{\alpha} \tfrac{1}{2}(a\sin\theta)^2\,d\theta + \int_{\alpha}^{\pi/2} \tfrac{1}{2}(b\cos\theta)^2\,d\theta \\ &= \tfrac{1}{4}a^2\left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\alpha} + \tfrac{1}{4}b^2\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_{\alpha}^{\pi/2} \\ &= \tfrac{1}{4}\alpha(a^2 - b^2) + \tfrac{1}{8}\pi b^2 - \tfrac{1}{4}(a^2 + b^2)(\sin\alpha\cos\alpha) \\ &= \tfrac{1}{4}(a^2 - b^2)\tan^{-1}(b/a) + \tfrac{1}{8}\pi b^2 - \tfrac{1}{4}ab \end{aligned}$$

35. The darker shaded region (from $\theta = 0$ to $\theta = 2\pi/3$) represents $\frac{1}{2}$ of the desired area plus $\frac{1}{2}$ of the area of the inner loop. From this area, we'll subtract $\frac{1}{2}$ of the area of the inner loop (the lighter shaded region from $\theta = 2\pi/3$ to $\theta = \pi$), and then double that difference to obtain the desired area.



$$\begin{aligned} A &= 2\left[\int_0^{2\pi/3} \tfrac{1}{2}\left(\tfrac{1}{2} + \cos\theta\right)^2\,d\theta - \int_{2\pi/3}^{\pi} \tfrac{1}{2}\left(\tfrac{1}{2} + \cos\theta\right)^2\,d\theta\right] \\ &= \int_0^{2\pi/3} \left(\tfrac{1}{4} + \cos\theta + \cos^2\theta\right)d\theta - \int_{2\pi/3}^{\pi} \left(\tfrac{1}{4} + \cos\theta + \cos^2\theta\right)d\theta \\ &= \int_0^{2\pi/3} \left[\tfrac{1}{4} + \cos\theta + \tfrac{1}{2}(1 + \cos 2\theta)\right]d\theta \\ &\qquad - \int_{2\pi/3}^{\pi} \left[\tfrac{1}{4} + \cos\theta + \tfrac{1}{2}(1 + \cos 2\theta)\right]d\theta \\ &= \left[\frac{\theta}{4} + \sin\theta + \frac{\theta}{2} + \frac{\sin 2\theta}{4}\right]_0^{2\pi/3} - \left[\frac{\theta}{4} + \sin\theta + \frac{\theta}{2} + \frac{\sin 2\theta}{4}\right]_{2\pi/3}^{\pi} \\ &= \left(\tfrac{\pi}{6} + \tfrac{\sqrt{3}}{2} + \tfrac{\pi}{3} - \tfrac{\sqrt{3}}{8}\right) - \left(\tfrac{\pi}{4} + \tfrac{\pi}{2}\right) + \left(\tfrac{\pi}{6} + \tfrac{\sqrt{3}}{2} + \tfrac{\pi}{3} - \tfrac{\sqrt{3}}{8}\right) \\ &= \tfrac{\pi}{4} + \tfrac{3}{4}\sqrt{3} = \tfrac{1}{4}\left(\pi + 3\sqrt{3}\right) \end{aligned}$$

36. $r = 0 \;\Rightarrow\; 1 + 2\cos 3\theta = 0 \;\Rightarrow\; \cos 3\theta = -\frac{1}{2} \;\Rightarrow\; 3\theta = \frac{2\pi}{3}, \frac{4\pi}{3}$ [for $0 \le 3\theta \le 2\pi$] $\Rightarrow\; \theta = \frac{2\pi}{9}, \frac{4\pi}{9}$. The darker shaded region (from $\theta = 0$ to $\theta = 2\pi/9$) represents $\frac{1}{2}$ of the desired area plus $\frac{1}{2}$ of the area of the inner loop. From this area, we'll subtract $\frac{1}{2}$ of the area of the inner loop (the lighter shaded region from $\theta = 2\pi/9$ to $\theta = \pi/3$), and then double that difference to obtain the desired area.



$$A = 2\left[\int_0^{2\pi/9} \tfrac{1}{2}(1 + 2\cos 3\theta)^2\,d\theta - \int_{2\pi/9}^{\pi/3} \tfrac{1}{2}(1 + 2\cos 3\theta)^2\,d\theta\right]$$

Now
$$\begin{aligned} r^2 &= (1 + 2\cos 3\theta)^2 = 1 + 4\cos 3\theta + 4\cos^2 3\theta = 1 + 4\cos 3\theta + 4\cdot\tfrac{1}{2}(1 + \cos 6\theta) \\ &= 1 + 4\cos 3\theta + 2 + 2\cos 6\theta = 3 + 4\cos 3\theta + 2\cos 6\theta \end{aligned}$$

and $\int r^2 d\theta = 3\theta + \frac{4}{3}\sin 3\theta + \frac{1}{3}\sin 6\theta + C$, so

$$\begin{aligned} A &= \left[3\theta + \tfrac{4}{3}\sin 3\theta + \tfrac{1}{3}\sin 6\theta\right]_0^{2\pi/9} - \left[3\theta + \tfrac{4}{3}\sin 3\theta + \tfrac{1}{3}\sin 6\theta\right]_{2\pi/9}^{\pi/3} \\ &= \left[\left(\tfrac{2\pi}{3} + \tfrac{4}{3}\cdot\tfrac{\sqrt{3}}{2} + \tfrac{1}{3}\cdot\tfrac{-\sqrt{3}}{2}\right) - 0\right] - \left[(\pi + 0 + 0) - \left(\tfrac{2\pi}{3} + \tfrac{4}{3}\cdot\tfrac{\sqrt{3}}{2} + \tfrac{1}{3}\cdot\tfrac{-\sqrt{3}}{2}\right)\right] \\ &= \tfrac{4\pi}{3} + \tfrac{4}{3}\sqrt{3} - \tfrac{1}{3}\sqrt{3} - \pi = \tfrac{\pi}{3} + \sqrt{3} \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

37. The pole is a point of intersection. $\sin\theta = 1 - \sin\theta \;\Rightarrow\; 2\sin\theta = 1 \;\Rightarrow\; \sin\theta = \frac{1}{2} \;\Rightarrow\; \theta = \frac{\pi}{6}$ or $\frac{5\pi}{6}$. So the other points of intersection are $\left(\frac{1}{2}, \frac{\pi}{6}\right)$ and $\left(\frac{1}{2}, \frac{5\pi}{6}\right)$.



38. The pole is a point of intersection. $1 + \cos\theta = 1 - \sin\theta \;\Rightarrow\; \cos\theta = -\sin\theta \;\Rightarrow\; \dfrac{\cos\theta}{\sin\theta} = -1 \;\Rightarrow\; \cot\theta = -1 \;\Rightarrow\; \theta = \frac{3\pi}{4}$ or $\frac{7\pi}{4}$. So the other points of intersection are $\left(1 - \frac{1}{2}\sqrt{2}, \frac{3\pi}{4}\right)$ and $\left(1 + \frac{1}{2}\sqrt{2}, \frac{7\pi}{4}\right)$.



39. $2\sin 2\theta = 1 \;\Rightarrow\; \sin 2\theta = \frac{1}{2} \;\Rightarrow\; 2\theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{13\pi}{6}$, or $\frac{17\pi}{6}$.

By symmetry, the eight points of intersection are given by $(1, \theta)$, where $\theta = \frac{\pi}{12}, \frac{5\pi}{12}, \frac{13\pi}{12}$, and $\frac{17\pi}{12}$, and $(-1, \theta)$, where $\theta = \frac{7\pi}{12}, \frac{11\pi}{12}, \frac{19\pi}{12}$, and $\frac{23\pi}{12}$.

[There are many ways to describe these points.]



40. Clearly the pole lies on both curves. $\sin 3\theta = \cos 3\theta \;\Rightarrow\; \tan 3\theta = 1 \;\Rightarrow\; 3\theta = \frac{\pi}{4} + n\pi$ [n any integer] $\;\Rightarrow\; \theta = \frac{\pi}{12} + \frac{\pi}{3}n \;\Rightarrow\; \theta = \frac{\pi}{12}, \frac{5\pi}{12}$, or $\frac{3\pi}{4}$, so the three remaining intersection points are $\left(\frac{1}{\sqrt{2}}, \frac{\pi}{12}\right)$, $\left(-\frac{1}{\sqrt{2}}, \frac{5\pi}{12}\right)$, and $\left(\frac{1}{\sqrt{2}}, \frac{3\pi}{4}\right)$.



41. The pole is a point of intersection. $\sin\theta = \sin 2\theta = 2\sin\theta\cos\theta \;\Leftrightarrow\; \sin\theta(1 - 2\cos\theta) = 0 \;\Leftrightarrow\; \sin\theta = 0$ or $\cos\theta = \frac{1}{2} \;\Rightarrow\; \theta = 0, \pi, \frac{\pi}{3}$, or $-\frac{\pi}{3} \;\Rightarrow$ the other intersection points are $\left(\frac{\sqrt{3}}{2}, \frac{\pi}{3}\right)$ and $\left(\frac{\sqrt{3}}{2}, \frac{2\pi}{3}\right)$ [by symmetry].



42. Clearly the pole is a point of intersection. $\sin 2\theta = \cos 2\theta \;\Rightarrow\; \tan 2\theta = 1 \;\Rightarrow\; 2\theta = \frac{\pi}{4} + 2n\pi$ [since $\sin 2\theta$ and $\cos 2\theta$ must be positive in the equations] $\;\Rightarrow\; \theta = \frac{\pi}{8} + n\pi \;\Rightarrow\; \theta = \frac{\pi}{8}$ or $\frac{9\pi}{8}$. So the curves also intersect at $\left(\frac{1}{\sqrt[4]{2}}, \frac{\pi}{8}\right)$ and $\left(\frac{1}{\sqrt[4]{2}}, \frac{9\pi}{8}\right)$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43.





From the first graph, we see that the pole is one point of intersection. By zooming in or using the cursor, we find the θ-values of the intersection points to be $\alpha \approx 0.88786 \approx 0.89$ and $\pi - \alpha \approx 2.25$. (The first of these values may be more easily estimated by plotting $y = 1 + \sin x$ and $y = 2x$ in rectangular coordinates; see the second graph.) By symmetry, the total area contained is twice the area contained in the first quadrant, that is,

$$A = 2\int_0^{\alpha} \tfrac{1}{2}(2\theta)^2\,d\theta + 2\int_{\alpha}^{\pi/2} \tfrac{1}{2}(1+\sin\theta)^2\,d\theta = \int_0^{\alpha} 4\theta^2\,d\theta + \int_{\alpha}^{\pi/2} \left[1 + 2\sin\theta + \tfrac{1}{2}(1-\cos 2\theta)\right] d\theta$$

$$= \left[\tfrac{4}{3}\theta^3\right]_0^{\alpha} + \left[\theta - 2\cos\theta + \left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right)\right]_{\alpha}^{\pi/2} = \tfrac{4}{3}\alpha^3 + \left[\left(\tfrac{\pi}{2} + \tfrac{\pi}{4}\right) - \left(\alpha - 2\cos\alpha + \tfrac{1}{2}\alpha - \tfrac{1}{4}\sin 2\alpha\right)\right] \approx 3.4645$$

44.



We need to find the shaded area A in the figure. The horizontal line representing the front of the stage has equation $y = 4 \Leftrightarrow r\sin\theta = 4 \Rightarrow r = 4/\sin\theta$. This line intersects the curve $r = 8 + 8\sin\theta$ when $8 + 8\sin\theta = \dfrac{4}{\sin\theta} \Rightarrow$

$8\sin\theta + 8\sin^2\theta = 4 \Rightarrow 2\sin^2\theta + 2\sin\theta - 1 = 0 \Rightarrow$

$$\sin\theta = \frac{-2 \pm \sqrt{4+8}}{4} = \frac{-2 \pm 2\sqrt{3}}{4} = \frac{-1+\sqrt{3}}{2} \quad \text{[the other value is less than } -1\text{]} \quad \Rightarrow \quad \theta = \sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right).$$

This angle is about $21.5°$ and is denoted by α in the figure.

$$A = 2\int_{\alpha}^{\pi/2} \tfrac{1}{2}(8+8\sin\theta)^2\,d\theta - 2\int_{\alpha}^{\pi/2} \tfrac{1}{2}(4\csc\theta)^2\,d\theta = 64\int_{\alpha}^{\pi/2}(1 + 2\sin\theta + \sin^2\theta)\,d\theta - 16\int_{\alpha}^{\pi/2}\csc^2\theta\,d\theta$$

$$= 64\int_{\alpha}^{\pi/2}\left(1 + 2\sin\theta + \tfrac{1}{2} - \tfrac{1}{2}\cos 2\theta\right)\,d\theta + 16\int_{\alpha}^{\pi/2}(-\csc^2\theta)\,d\theta = 64\left[\tfrac{3}{2}\theta - 2\cos\theta - \tfrac{1}{4}\sin 2\theta\right]_{\alpha}^{\pi/2} + 16\left[\cot\theta\right]_{\alpha}^{\pi/2}$$

$$= 16\left[6\theta - 8\cos\theta - \sin 2\theta + \cot\theta\right]_{\alpha}^{\pi/a} = 16[(3\pi - 0 - 0 + 0) - (6\alpha - 8\cos\alpha - \sin 2\alpha + \cot\alpha)]$$

$$= 48\pi - 96\alpha + 128\cos\alpha + 16\sin 2\alpha - 16\cot\alpha$$

From the figure, $x^2 + \left(\sqrt{3}-1\right)^2 = 2^2 \Rightarrow x^2 = 4 - \left(3 - 2\sqrt{3} + 1\right) \Rightarrow$ $x^2 = 2\sqrt{3} = \sqrt{12}$, so $x = \sqrt{2\sqrt{3}} = \sqrt[4]{12}$. Using the trigonometric relationships for a right triangle and the identity $\sin 2\alpha = 2\sin\alpha\cos\alpha$, we continue:



$$A = 48\pi - 96\alpha + 128 \cdot \frac{\sqrt[4]{12}}{2} + 16 \cdot 2 \cdot \frac{\sqrt{3}-1}{2} \cdot \frac{\sqrt[4]{12}}{2} - 16 \cdot \frac{\sqrt[4]{12}}{\sqrt{3}-1} \cdot \frac{\sqrt{3}+1}{\sqrt{3}+1}$$

$$= 48\pi - 96\alpha + 64\sqrt[4]{12} + 8\sqrt[4]{12}\left(\sqrt{3}-1\right) - 8\sqrt[4]{12}\left(\sqrt{3}+1\right) = 48\pi + 48\sqrt[4]{12} - 96\sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right)$$

$$\approx 204.16 \text{ m}^2$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

45. $L = \int_a^b \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_0^\pi \sqrt{(2\cos\theta)^2 + (-2\sin\theta)^2}\, d\theta$

$= \int_0^\pi \sqrt{4(\cos^2\theta + \sin^2\theta)}\, d\theta = \int_0^\pi \sqrt{4}\, d\theta = [2\theta]_0^\pi = 2\pi$

As a check, note that the curve is a circle of radius 1, so its circumference is $2\pi(1) = 2\pi$.

46. $L = \int_a^b \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{(5^\theta)^2 + (5^\theta \ln 5)^2}\, d\theta = \int_0^{2\pi} \sqrt{5^{2\theta}[1 + (\ln 5)^2]}\, d\theta$

$= \sqrt{1 + (\ln 5)^2} \int_0^{2\pi} \sqrt{5^{2\theta}}\, d\theta = \sqrt{1 + (\ln 5)^2} \int_0^{2\pi} 5^\theta\, d\theta = \sqrt{1 + (\ln 5)^2} \left[\frac{5^\theta}{\ln 5}\right]_0^{2\pi}$

$= \sqrt{1 + (\ln 5)^2} \left(\frac{5^{2\pi}}{\ln 5} - \frac{1}{\ln 5}\right) = \frac{\sqrt{1 + (\ln 5)^2}}{\ln 5}(5^{2\pi} - 1)$

47. $L = \int_a^b \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{(\theta^2)^2 + (2\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{\theta^4 + 4\theta^2}\, d\theta$

$= \int_0^{2\pi} \sqrt{\theta^2(\theta^2 + 4)}\, d\theta = \int_0^{2\pi} \theta\sqrt{\theta^2 + 4}\, d\theta$

Now let $u = \theta^2 + 4$, so that $du = 2\theta\, d\theta$ $\left[\theta\, d\theta = \frac{1}{2}\, du\right]$ and

$$\int_0^{2\pi} \theta\sqrt{\theta^2 + 4}\, d\theta = \int_4^{4\pi^2+4} \tfrac{1}{2}\sqrt{u}\, du = \tfrac{1}{2} \cdot \tfrac{2}{3}\left[u^{3/2}\right]_4^{4(\pi^2+1)} = \tfrac{1}{3}[4^{3/2}(\pi^2+1)^{3/2} - 4^{3/2}] = \tfrac{8}{3}[(\pi^2+1)^{3/2} - 1]$$

48. $L = \int_a^b \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{[2(1+\cos\theta)]^2 + (-2\sin\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{4 + 8\cos\theta + 4\cos^2\theta + 4\sin^2\theta}\, d\theta$

$= \int_0^{2\pi} \sqrt{8 + 8\cos\theta}\, d\theta = \sqrt{8} \int_0^{2\pi} \sqrt{1 + \cos\theta}\, d\theta = \sqrt{8} \int_0^{2\pi} \sqrt{2 \cdot \tfrac{1}{2}(1 + \cos\theta)}\, d\theta$

$= \sqrt{8} \int_0^{2\pi} \sqrt{2\cos^2\frac{\theta}{2}}\, d\theta = \sqrt{8}\sqrt{2} \int_0^{2\pi} \left|\cos\frac{\theta}{2}\right| d\theta = 4 \cdot 2 \int_0^{\pi} \cos\frac{\theta}{2}\, d\theta$ [by symmetry]

$= 8\left[2\sin\frac{\theta}{2}\right]_0^{\pi} = 8(2) = 16$

49. The curve $r = \cos^4(\theta/4)$ is completely traced with $0 \le \theta \le 4\pi$.



$r^2 + (dr/d\theta)^2 = [\cos^4(\theta/4)]^2 + \left[4\cos^3(\theta/4) \cdot (-\sin(\theta/4)) \cdot \tfrac{1}{4}\right]^2$

$= \cos^8(\theta/4) + \cos^6(\theta/4)\sin^2(\theta/4)$

$= \cos^6(\theta/4)[\cos^2(\theta/4) + \sin^2(\theta/4)] = \cos^6(\theta/4)$

$L = \int_0^{4\pi} \sqrt{\cos^6(\theta/4)}\, d\theta = \int_0^{4\pi} \left|\cos^3(\theta/4)\right| d\theta$

$= 2\int_0^{2\pi} \cos^3(\theta/4)\, d\theta$ [since $\cos^3(\theta/4) \ge 0$ for $0 \le \theta \le 2\pi$] $= 8\int_0^{\pi/2} \cos^3 u\, du$ $\left[u = \tfrac{1}{4}\theta\right]$

$= 8\int_0^{\pi/2} (1 - \sin^2 u)\cos u\, du = 8\int_0^1 (1 - x^2)\, dx$ $\left[\begin{matrix} x = \sin u, \\ dx = \cos u\, du \end{matrix}\right]$

$= 8\left[x - \tfrac{1}{3}x^3\right]_0^1 = 8\left(1 - \tfrac{1}{3}\right) = \tfrac{16}{3}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. The curve $r = \cos^2(\theta/2)$ is completely traced with $0 \le \theta \le 2\pi$.



$$\begin{aligned} r^2 + (dr/d\theta)^2 &= [\cos^2(\theta/2)]^2 + \left[2\cos(\theta/2)\cdot(-\sin(\theta/2))\cdot\tfrac{1}{2}\right]^2 \\ &= \cos^4(\theta/2) + \cos^2(\theta/2)\sin^2(\theta/2) \\ &= \cos^2(\theta/2)[\cos^2(\theta/2) + \sin^2(\theta/2)] \\ &= \cos^2(\theta/2) \end{aligned}$$

$$L = \int_0^{2\pi} \sqrt{\cos^2(\theta/2)}\, d\theta = \int_0^{2\pi} |\cos(\theta/2)|\, d\theta = 2\int_0^{\pi} \cos(\theta/2)\, d\theta \qquad [\text{since } \cos(\theta/2) \ge 0 \text{ for } 0 \le \theta \le \pi]$$

$$= 4\int_0^{\pi/2} \cos u\, du \quad [u = \tfrac{1}{2}\theta] \quad = 4\big[\sin u\big]_0^{\pi/2} = 4(1-0) = 4$$

51. One loop of the curve $r = \cos 2\theta$ is traced with $-\pi/4 \le \theta \le \pi/4$.

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \cos^2 2\theta + (-2\sin 2\theta)^2 = \cos^2 2\theta + 4\sin^2 2\theta = 1 + 3\sin^2 2\theta \;\Rightarrow$$

$$L = \int_{-\pi/4}^{\pi/4} \sqrt{1 + 3\sin^2 2\theta}\, d\theta \approx 2.4221.$$

52. $r^2 + \left(\dfrac{dr}{d\theta}\right)^2 = \tan^2\theta + (\sec^2\theta)^2 \;\Rightarrow\; L = \displaystyle\int_{\pi/6}^{\pi/3} \sqrt{\tan^2\theta + \sec^4\theta}\, d\theta \approx 1.2789$

53. The curve $r = \sin(6\sin\theta)$ is completely traced with $0 \le \theta \le \pi$. $r = \sin(6\sin\theta) \;\Rightarrow$

$\dfrac{dr}{d\theta} = \cos(6\sin\theta)\cdot 6\cos\theta$, so $r^2 + \left(\dfrac{dr}{d\theta}\right)^2 = \sin^2(6\sin\theta) + 36\cos^2\theta\cos^2(6\sin\theta) \;\Rightarrow$

$$L = \int_0^{\pi} \sqrt{\sin^2(6\sin\theta) + 36\cos^2\theta\,\cos^2(6\sin\theta)}\, d\theta \approx 8.0091.$$

54. The curve $r = \sin(\theta/4)$ is completely traced with $0 \le \theta \le 8\pi$. $r = \sin(\theta/4) \;\Rightarrow\; \dfrac{dr}{d\theta} = \frac{1}{4}\cos(\theta/4)$, so

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \sin^2(\theta/4) + \tfrac{1}{16}\cos^2(\theta/4) \;\Rightarrow\; L = \int_0^{8\pi} \sqrt{\sin^2(\theta/4) + \tfrac{1}{16}\cos^2(\theta/4)}\, d\theta \approx 17.1568.$$

55. (a) From (10.2.6),

$$\begin{aligned} S &= \int_a^b 2\pi y\sqrt{(dx/d\theta)^2 + (dy/d\theta)^2}\, d\theta \\ &= \int_a^b 2\pi y\sqrt{r^2 + (dr/d\theta)^2}\, d\theta \qquad [\text{from the derivation of Equation 10.4.5}] \\ &= \int_a^b 2\pi r\sin\theta\sqrt{r^2 + (dr/d\theta)^2}\, d\theta \end{aligned}$$

(b) The curve $r^2 = \cos 2\theta$ goes through the pole when $\cos 2\theta = 0 \;\Rightarrow$ $2\theta = \frac{\pi}{2} \;\Rightarrow\; \theta = \frac{\pi}{4}$. We'll rotate the curve from $\theta = 0$ to $\theta = \frac{\pi}{4}$ and double this value to obtain the total surface area generated.



$$r^2 = \cos 2\theta \;\Rightarrow\; 2r\frac{dr}{d\theta} = -2\sin 2\theta \;\Rightarrow\; \left(\frac{dr}{d\theta}\right)^2 = \frac{\sin^2 2\theta}{r^2} = \frac{\sin^2 2\theta}{\cos 2\theta}.$$

$$S = 2\int_0^{\pi/4} 2\pi\sqrt{\cos 2\theta}\sin\theta\sqrt{\cos 2\theta + (\sin^2 2\theta)/\cos 2\theta}\, d\theta = 4\pi\int_0^{\pi/4} \sqrt{\cos 2\theta}\sin\theta\sqrt{\frac{\cos^2 2\theta + \sin^2 2\theta}{\cos 2\theta}}\, d\theta$$

$$= 4\pi\int_0^{\pi/4} \sqrt{\cos 2\theta}\sin\theta\,\frac{1}{\sqrt{\cos 2\theta}}\, d\theta = 4\pi\int_0^{\pi/4} \sin\theta\, d\theta = 4\pi\big[-\cos\theta\big]_0^{\pi/4} = -4\pi\left(\tfrac{\sqrt{2}}{2} - 1\right) = 2\pi\left(2 - \sqrt{2}\right)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

56. (a) Rotation around $\theta = \frac{\pi}{2}$ is the same as rotation around the y-axis, that is, $S = \int_a^b 2\pi x \, ds$ where $ds = \sqrt{(dx/dt)^2 + (dy/dt)^2}\, dt$ for a parametric equation, and for the special case of a polar equation, $x = r\cos\theta$ and $ds = \sqrt{(dx/d\theta)^2 + (dy/d\theta)^2}\, d\theta = \sqrt{r^2 + (dr/d\theta)^2}\, d\theta$ [see the derivation of Equation 10.4.5]. Therefore, for a polar equation rotated around $\theta = \frac{\pi}{2}$, $S = \int_a^b 2\pi r\cos\theta\sqrt{r^2 + (dr/d\theta)^2}\, d\theta$.

(b) As in the solution for Exercise 55(b), we can double the surface area generated by rotating the curve from $\theta = 0$ to $\theta = \frac{\pi}{4}$ to obtain the total surface area.

$$S = 2\int_0^{\pi/4} 2\pi\sqrt{\cos 2\theta}\cos\theta\sqrt{\cos 2\theta + (\sin^2 2\theta)/\cos 2\theta}\, d\theta = 4\pi\int_0^{\pi/4}\sqrt{\cos 2\theta}\cos\theta\sqrt{\frac{\cos^2 2\theta + \sin^2 2\theta}{\cos 2\theta}}\, d\theta$$
$$= 4\pi\int_0^{\pi/4}\sqrt{\cos 2\theta}\cos\theta\,\frac{1}{\sqrt{\cos 2\theta}}\, d\theta = 4\pi\int_0^{\pi/4}\cos\theta\, d\theta = 4\pi\big[\sin\theta\big]_0^{\pi/4} = 4\pi\left(\frac{\sqrt{2}}{2} - 0\right) = 2\sqrt{2}\,\pi$$

10.5 Conic Sections

1. $x^2 = 6y$ and $x^2 = 4py \;\Rightarrow\; 4p = 6 \;\Rightarrow\; p = \frac{3}{2}$.

The vertex is $(0, 0)$, the focus is $\left(0, \frac{3}{2}\right)$, and the directrix is $y = -\frac{3}{2}$.



2. $2y^2 = 5x \;\Rightarrow\; y^2 = \frac{5}{2}x$. $4p = \frac{5}{2} \;\Rightarrow\; p = \frac{5}{8}$.

The vertex is $(0, 0)$, the focus is $\left(\frac{5}{8}, 0\right)$, and the directrix is $x = -\frac{5}{8}$.



3. $2x = -y^2 \;\Rightarrow\; y^2 = -2x$. $4p = -2 \;\Rightarrow\; p = -\frac{1}{2}$.

The vertex is $(0, 0)$, the focus is $\left(-\frac{1}{2}, 0\right)$, and the directrix is $x = \frac{1}{2}$.



4. $3x^2 + 8y = 0 \;\Rightarrow\; 3x^2 = -8y \;\Rightarrow\; x^2 = -\frac{8}{3}y$. $4p = -\frac{8}{3} \;\Rightarrow\; p = -\frac{2}{3}$. The vertex is $(0, 0)$, the focus is $\left(0, -\frac{2}{3}\right)$, and the directrix is $y = \frac{2}{3}$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $(x+2)^2 = 8(y-3)$. $4p = 8$, so $p = 2$. The vertex is $(-2, 3)$, the focus is $(-2, 5)$, and the directrix is $y = 1$.



6. $(y-2)^2 = 2x + 1 = 2\left(x + \frac{1}{2}\right)$. $4p = 2$, so $p = \frac{1}{2}$. The vertex is $\left(-\frac{1}{2}, 2\right)$, the focus is $(0, 2)$, and the directrix is $x = -1$.



7. $y^2 + 6y + 2x + 1 = 0 \Leftrightarrow y^2 + 6y = -2x - 1 \Leftrightarrow y^2 + 6y + 9 = -2x + 8 \Leftrightarrow (y+3)^2 = -2(x-4)$. $4p = -2$, so $p = -\frac{1}{2}$. The vertex is $(4, -3)$, the focus is $\left(\frac{7}{2}, -3\right)$, and the directrix is $x = \frac{9}{2}$.



8. $2x^2 - 16x - 3y + 38 = 0 \Leftrightarrow 2x^2 - 16x = 3y - 38 \Leftrightarrow 2(x^2 - 8x + 16) = 3y - 38 + 32 \Leftrightarrow 2(x-4)^2 = 3y - 6 \Leftrightarrow (x-4)^2 = \frac{3}{2}(y-2)$. $4p = \frac{3}{2}$, so $p = \frac{3}{8}$. The vertex is $(4, 2)$, the focus is $\left(4, \frac{19}{8}\right)$, and the directrix is $y = \frac{13}{8}$.



9. The equation has the form $y^2 = 4px$, where $p < 0$. Since the parabola passes through $(-1, 1)$, we have $1^2 = 4p(-1)$, so $4p = -1$ and an equation is $y^2 = -x$ or $x = -y^2$. $4p = -1$, so $p = -\frac{1}{4}$ and the focus is $\left(-\frac{1}{4}, 0\right)$ while the directrix is $x = \frac{1}{4}$.

10. The vertex is $(2, -2)$, so the equation is of the form $(x-2)^2 = 4p(y+2)$, where $p > 0$. The point $(0, 0)$ is on the parabola, so $4 = 4p(2)$ and $4p = 2$. Thus, an equation is $(x-2)^2 = 2(y+2)$. $4p = 2$, so $p = \frac{1}{2}$ and the focus is $\left(2, -\frac{3}{2}\right)$ while the directrix is $y = -\frac{5}{2}$.

11. $\frac{x^2}{2} + \frac{y^2}{4} = 1 \Rightarrow a = \sqrt{4} = 2, b = \sqrt{2}$, $c = \sqrt{a^2 - b^2} = \sqrt{4-2} = \sqrt{2}$. The ellipse is centered at $(0, 0)$, with vertices at $(0, \pm 2)$. The foci are $\left(0, \pm\sqrt{2}\right)$.



12. $\frac{x^2}{36} + \frac{y^2}{8} = 1 \Rightarrow a = \sqrt{36} = 6, b = \sqrt{8}$, $c = \sqrt{a^2 - b^2} = \sqrt{36 - 8} = \sqrt{28} = 2\sqrt{7}$. The ellipse is centered at $(0, 0)$, with vertices at $(\pm 6, 0)$. The foci are $(\pm 2\sqrt{7}, 0)$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

13. $x^2+9y^2=9 \quad \Leftrightarrow \quad \dfrac{x^2}{9}+\dfrac{y^2}{1}=1 \quad \Rightarrow \quad a=\sqrt{9}=3,$ $b=\sqrt{1}=1, c=\sqrt{a^2-b^2}=\sqrt{9-1}=\sqrt{8}=2\sqrt{2}.$ The ellipse is centered at $(0,0)$, with vertices $(\pm 3,0)$. The foci are $(\pm 2\sqrt{2},0)$.



14. $100x^2+36y^2=225 \quad \Leftrightarrow \quad \dfrac{x^2}{\frac{225}{100}}+\dfrac{y^2}{\frac{225}{36}}=1 \quad \Leftrightarrow$ $\dfrac{x^2}{\frac{9}{4}}+\dfrac{y^2}{\frac{25}{4}}=1 \quad \Rightarrow \quad a=\sqrt{\frac{25}{4}}=\frac{5}{2}, b=\sqrt{\frac{9}{4}}=\frac{3}{2},$ $c=\sqrt{a^2-b^2}=\sqrt{\frac{25}{4}-\frac{9}{4}}=2$. The ellipse is centered at $(0,0)$, with vertices $\left(0,\pm\frac{5}{2}\right)$. The foci are $(0,\pm 2)$.



15. $9x^2-18x+4y^2=27 \quad \Leftrightarrow$ $9(x^2-2x+1)+4y^2=27+9 \quad \Leftrightarrow$ $9(x-1)^2+4y^2=36 \quad \Leftrightarrow \quad \dfrac{(x-1)^2}{4}+\dfrac{y^2}{9}=1 \quad \Rightarrow$ $a=3, b=2, c=\sqrt{5} \quad \Rightarrow$ center $(1,0)$, vertices $(1,\pm 3)$, foci $\left(1,\pm\sqrt{5}\right)$



16. $x^2+3y^2+2x-12y+10=0 \quad \Leftrightarrow$ $x^2+2x+1+3(y^2-4y+4)=-10+1+12 \quad \Leftrightarrow$ $(x+1)^2+3(y-2)^2=3 \quad \Leftrightarrow$ $\dfrac{(x+1)^2}{3}+\dfrac{(y-2)^2}{1}=1 \quad \Rightarrow \quad a=\sqrt{3}, b=1,$ $c=\sqrt{2} \quad \Rightarrow$ center $(-1,2)$, vertices $\left(-1\pm\sqrt{3},2\right)$, foci $\left(-1\pm\sqrt{2},2\right)$



17. The center is $(0,0)$, $a=3$, and $b=2$, so an equation is $\dfrac{x^2}{4}+\dfrac{y^2}{9}=1$. $c=\sqrt{a^2-b^2}=\sqrt{5}$, so the foci are $\left(0,\pm\sqrt{5}\right)$.

18. The ellipse is centered at $(2,1)$, with $a=3$ and $b=2$. An equation is $\dfrac{(x-2)^2}{9}+\dfrac{(y-1)^2}{4}=1$. $c=\sqrt{a^2-b^2}=\sqrt{5}$, so the foci are $\left(2\pm\sqrt{5},1\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $\dfrac{y^2}{25} - \dfrac{x^2}{9} = 1 \;\Rightarrow\; a = 5, b = 3, c = \sqrt{25+9} = \sqrt{34} \;\Rightarrow$

center $(0, 0)$, vertices $(0, \pm 5)$, foci $\left(0, \pm\sqrt{34}\right)$, asymptotes $y = \pm\frac{5}{3}x$.

Note: It is helpful to draw a $2a$-by-$2b$ rectangle whose center is the center of the hyperbola. The asymptotes are the extended diagonals of the rectangle.



20. $\dfrac{x^2}{36} - \dfrac{y^2}{64} = 1 \;\Rightarrow\; a = 6, b = 8, c = \sqrt{36+64} = 10 \;\Rightarrow$

center $(0, 0)$, vertices $(\pm 6, 0)$, foci $(\pm 10, 0)$, asymptotes $y = \pm\frac{8}{6}x = \pm\frac{4}{3}x$



21. $x^2 - y^2 = 100 \;\Leftrightarrow\; \dfrac{x^2}{100} - \dfrac{y^2}{100} = 1 \;\Rightarrow\; a = b = 10,$

$c = \sqrt{100+100} = 10\sqrt{2} \;\Rightarrow$ center $(0, 0)$, vertices $(\pm 10, 0)$,

foci $\left(\pm 10\sqrt{2}, 0\right)$, asymptotes $y = \pm\frac{10}{10}x = \pm x$



22. $y^2 - 16x^2 = 16 \;\Leftrightarrow\; \dfrac{y^2}{16} - \dfrac{x^2}{1} = 1 \;\Rightarrow\; a = 4, b = 1,$

$c = \sqrt{16+1} = \sqrt{17} \;\Rightarrow$ center $(0, 0)$, vertices $(0, \pm 4)$,

foci $\left(0, \pm\sqrt{17}\right)$, asymptotes $y = \pm\frac{4}{1}x = \pm 4x$



23. $x^2 - y^2 + 2y = 2 \;\Leftrightarrow\; x^2 - (y^2 - 2y + 1) = 2 - 1 \;\Leftrightarrow$

$\dfrac{x^2}{1} - \dfrac{(y-1)^2}{1} = 1 \;\Rightarrow\; a = b = 1, c = \sqrt{1+1} = \sqrt{2} \;\Rightarrow$

center $(0, 1)$, vertices $(\pm 1, 1)$, foci $\left(\pm\sqrt{2}, 1\right)$,

asymptotes $y - 1 = \pm\frac{1}{1}x = \pm x$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

24. $9y^2 - 4x^2 - 36y - 8x = 4 \quad \Leftrightarrow$

$9(y^2 - 4y + 4) - 4(x^2 + 2x + 1) = 4 + 36 - 4 \quad \Leftrightarrow$

$9(y-2)^2 - 4(x+1)^2 = 36 \quad \Leftrightarrow \quad \dfrac{(y-2)^2}{4} - \dfrac{(x+1)^2}{9} = 1 \quad \Rightarrow$

$a = 2, b = 3, c = \sqrt{4+9} = \sqrt{13} \quad \Rightarrow$ center $(-1, 2)$, vertices $(-1, 2 \pm 2)$, foci $\left(-1, 2 \pm \sqrt{13}\right)$, asymptotes $y - 2 = \pm\frac{2}{3}(x+1)$.



25. $4x^2 = y^2 + 4 \quad \Leftrightarrow \quad 4x^2 - y^2 = 4 \quad \Leftrightarrow \quad \dfrac{x^2}{1} - \dfrac{y^2}{4} = 1$. This is an equation of a *hyperbola* with vertices $(\pm 1, 0)$. The foci are at $\left(\pm\sqrt{1+4}, 0\right) = \left(\pm\sqrt{5}, 0\right)$.

26. $4x^2 = y + 4 \quad \Leftrightarrow \quad x^2 = \frac{1}{4}(y+4)$. This is an equation of a *parabola* with $4p = \frac{1}{4}$, so $p = \frac{1}{16}$. The vertex is $(0, -4)$ and the focus is $\left(0, -4 + \frac{1}{16}\right) = \left(0, -\frac{63}{16}\right)$.

27. $x^2 = 4y - 2y^2 \quad \Leftrightarrow \quad x^2 + 2y^2 - 4y = 0 \quad \Leftrightarrow \quad x^2 + 2(y^2 - 2y + 1) = 2 \quad \Leftrightarrow \quad x^2 + 2(y-1)^2 = 2 \quad \Leftrightarrow$

$\dfrac{x^2}{2} + \dfrac{(y-1)^2}{1} = 1$. This is an equation of an *ellipse* with vertices at $\left(\pm\sqrt{2}, 1\right)$. The foci are at $\left(\pm\sqrt{2-1}, 1\right) = (\pm 1, 1)$.

28. $y^2 - 2 = x^2 - 2x \quad \Leftrightarrow \quad y^2 - x^2 + 2x = 2 \quad \Leftrightarrow \quad y^2 - (x^2 - 2x + 1) = 2 - 1 \quad \Leftrightarrow \quad \dfrac{y^2}{1} - \dfrac{(x-1)^2}{1} = 1$. This is an equation of a *hyperbola* with vertices $(1, \pm 1)$. The foci are at $\left(1, \pm\sqrt{1+1}\right) = \left(1, \pm\sqrt{2}\right)$.

29. $3x^2 - 6x - 2y = 1 \quad \Leftrightarrow \quad 3x^2 - 6x = 2y + 1 \quad \Leftrightarrow \quad 3(x^2 - 2x + 1) = 2y + 1 + 3 \quad \Leftrightarrow \quad 3(x-1)^2 = 2y + 4 \quad \Leftrightarrow$

$(x-1)^2 = \frac{2}{3}(y+2)$. This is an equation of a *parabola* with $4p = \frac{2}{3}$, so $p = \frac{1}{6}$. The vertex is $(1, -2)$ and the focus is $\left(1, -2 + \frac{1}{6}\right) = \left(1, -\frac{11}{6}\right)$.

30. $x^2 - 2x + 2y^2 - 8y + 7 = 0 \quad \Leftrightarrow \quad (x^2 - 2x + 1) + 2(y^2 - 4y + 4) = -7 + 1 + 8 \quad \Leftrightarrow \quad (x-1)^2 + 2(y-2)^2 = 2 \quad \Leftrightarrow$

$\dfrac{(x-1)^2}{2} + \dfrac{(y-2)^2}{1} = 1$. This is an equation of an *ellipse* with vertices at $\left(1 \pm \sqrt{2}, 2\right)$. The foci are at $\left(1 \pm \sqrt{2-1}, 2\right) = (1 \pm 1, 2)$.

31. The parabola with vertex $(0, 0)$ and focus $(1, 0)$ opens to the right and has $p = 1$, so its equation is $y^2 = 4px$, or $y^2 = 4x$.

32. The parabola with focus $(0, 0)$ and directrix $y = 6$ has vertex $(0, 3)$ and opens downward, so $p = -3$ and its equation is $(x-0)^2 = 4p(y-3)$, or $x^2 = -12(y-3)$.

33. The distance from the focus $(-4, 0)$ to the directrix $x = 2$ is $2 - (-4) = 6$, so the distance from the focus to the vertex is $\frac{1}{2}(6) = 3$ and the vertex is $(-1, 0)$. Since the focus is to the left of the vertex, $p = -3$. An equation is $y^2 = 4p(x+1) \quad \Rightarrow \quad y^2 = -12(x+1)$.

34. The parabola with vertex $(2, 3)$ and focus $(2, -1)$ opens downward and has $p = -1 - 3 = -4$, so its equation is $(x-2)^2 = 4p(y-3)$, or $(x-2)^2 = -16(y-3)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. The parabola with vertex $(3, -1)$ having a horizontal axis has equation $[y-(-1)]^2 = 4p(x-3)$. Since it passes through $(-15, 2)$, $(2+1)^2 = 4p(-15-3) \Rightarrow 9 = 4p(-18) \Rightarrow 4p = -\frac{1}{2}$. An equation is $(y+1)^2 = -\frac{1}{2}(x-3)$.

36. The parabola with vertical axis and passing through $(0, 4)$ has equation $y = ax^2 + bx + 4$. It also passes through $(1, 3)$ and $(-2, -6)$, so

$$\begin{cases} 3 = a+b+4 \\ -6 = 4a-2b+4 \end{cases} \Rightarrow \begin{cases} -1 = a+b \\ -10 = 4a-2b \end{cases} \Rightarrow \begin{cases} -1 = a+b \\ -5 = 2a-b \end{cases}$$

Adding the last two equations gives us $3a = -6$, or $a = -2$. Since $a+b = -1$, we have $b = 1$, and an equation is $y = -2x^2 + x + 4$.

37. The ellipse with foci $(\pm 2, 0)$ and vertices $(\pm 5, 0)$ has center $(0, 0)$ and a horizontal major axis, with $a = 5$ and $c = 2$, so $b^2 = a^2 - c^2 = 25 - 4 = 21$. An equation is $\dfrac{x^2}{25} + \dfrac{y^2}{21} = 1$.

38. The ellipse with foci $\left(0, \pm\sqrt{2}\right)$ and vertices $(0, \pm 2)$ has center $(0, 0)$ and a vertical major axis, with $a = 2$ and $c = \sqrt{2}$, so $b^2 = a^2 - c^2 = 4 - 2 = 2$. An equation is $\dfrac{x^2}{2} + \dfrac{y^2}{4} = 1$.

39. Since the vertices are $(0, 0)$ and $(0, 8)$, the ellipse has center $(0, 4)$ with a vertical axis and $a = 4$. The foci at $(0, 2)$ and $(0, 6)$ are 2 units from the center, so $c = 2$ and $b = \sqrt{a^2 - c^2} = \sqrt{4^2 - 2^2} = \sqrt{12}$. An equation is $\dfrac{(x-0)^2}{b^2} + \dfrac{(y-4)^2}{a^2} = 1 \Rightarrow \dfrac{x^2}{12} + \dfrac{(y-4)^2}{16} = 1$.

40. Since the foci are $(0, -1)$ and $(8, -1)$, the ellipse has center $(4, -1)$ with a horizontal axis and $c = 4$. The vertex $(9, -1)$ is 5 units from the center, so $a = 5$ and $b = \sqrt{a^2 - c^2} = \sqrt{5^2 - 4^2} = \sqrt{9}$. An equation is $\dfrac{(x-4)^2}{a^2} + \dfrac{(y+1)^2}{b^2} = 1 \Rightarrow \dfrac{(x-4)^2}{25} + \dfrac{(y+1)^2}{9} = 1$.

41. An equation of an ellipse with center $(-1, 4)$ and vertex $(-1, 0)$ is $\dfrac{(x+1)^2}{b^2} + \dfrac{(y-4)^2}{4^2} = 1$. The focus $(-1, 6)$ is 2 units from the center, so $c = 2$. Thus, $b^2 + 2^2 = 4^2 \Rightarrow b^2 = 12$, and the equation is $\dfrac{(x+1)^2}{12} + \dfrac{(y-4)^2}{16} = 1$.

42. Foci $F_1(-4, 0)$ and $F_2(4, 0) \Rightarrow c = 4$ and an equation is $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$. The ellipse passes through $P(-4, 1.8)$, so

$2a = |PF_1| + |PF_2| \Rightarrow 2a = 1.8 + \sqrt{8^2 + (1.8)^2} \Rightarrow 2a = 1.8 + 8.2 \Rightarrow a = 5$.

$b^2 = a^2 - c^2 = 25 - 16 = 9$ and the equation is $\dfrac{x^2}{25} + \dfrac{y^2}{9} = 1$.

43. An equation of a hyperbola with vertices $(\pm 3, 0)$ is $\dfrac{x^2}{3^2} - \dfrac{y^2}{b^2} = 1$. Foci $(\pm 5, 0) \Rightarrow c = 5$ and $3^2 + b^2 = 5^2 \Rightarrow$ $b^2 = 25 - 9 = 16$, so the equation is $\dfrac{x^2}{9} - \dfrac{y^2}{16} = 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

44. An equation of a hyperbola with vertices $(0, \pm 2)$ is $\dfrac{y^2}{2^2} - \dfrac{x^2}{b^2} = 1$. Foci $(0, \pm 5) \;\Rightarrow\; c = 5$ and $2^2 + b^2 = 5^2 \;\Rightarrow$ $b^2 = 25 - 4 = 21$, so the equation is $\dfrac{y^2}{4} - \dfrac{x^2}{21} = 1$.

45. The center of a hyperbola with vertices $(-3, -4)$ and $(-3, 6)$ is $(-3, 1)$, so $a = 5$ and an equation is $\dfrac{(y-1)^2}{5^2} - \dfrac{(x+3)^2}{b^2} = 1$. Foci $(-3, -7)$ and $(-3, 9) \;\Rightarrow\; c = 8$, so $5^2 + b^2 = 8^2 \;\Rightarrow\; b^2 = 64 - 25 = 39$ and the equation is $\dfrac{(y-1)^2}{25} - \dfrac{(x+3)^2}{39} = 1$.

46. The center of a hyperbola with vertices $(-1, 2)$ and $(7, 2)$ is $(3, 2)$, so $a = 4$ and an equation is $\dfrac{(x-3)^2}{4^2} - \dfrac{(y-2)^2}{b^2} = 1$. Foci $(-2, 2)$ and $(8, 2) \;\Rightarrow\; c = 5$, so $4^2 + b^2 = 5^2 \;\Rightarrow\; b^2 = 25 - 16 = 9$ and the equation is $\dfrac{(x-3)^2}{16} - \dfrac{(y-2)^2}{9} = 1$.

47. The center of a hyperbola with vertices $(\pm 3, 0)$ is $(0, 0)$, so $a = 3$ and an equation is $\dfrac{x^2}{3^2} - \dfrac{y^2}{b^2} = 1$. Asymptotes $y = \pm 2x \;\Rightarrow\; \dfrac{b}{a} = 2 \;\Rightarrow\; b = 2(3) = 6$ and the equation is $\dfrac{x^2}{9} - \dfrac{y^2}{36} = 1$.

48. The center of a hyperbola with foci $(2, 0)$ and $(2, 8)$ is $(2, 4)$, so $c = 4$ and an equation is $\dfrac{(y-4)^2}{a^2} - \dfrac{(x-2)^2}{b^2} = 1$. The asymptote $y = 3 + \frac{1}{2}x$ has slope $\frac{1}{2}$, so $\dfrac{a}{b} = \dfrac{1}{2} \;\Rightarrow\; b = 2a$ and $a^2 + b^2 = c^2 \;\Rightarrow\; a^2 + (2a)^2 = 4^2 \;\Rightarrow$ $5a^2 = 16 \;\Rightarrow\; a^2 = \frac{16}{5}$ and so $b^2 = 16 - \frac{16}{5} = \frac{64}{5}$. Thus, an equation is $\dfrac{(y-4)^2}{16/5} - \dfrac{(x-2)^2}{64/5} = 1$.

49. In Figure 8, we see that the point on the ellipse closest to a focus is the closer vertex (which is a distance $a - c$ from it) while the farthest point is the other vertex (at a distance of $a + c$). So for this lunar orbit, $(a - c) + (a + c) = 2a = (1728 + 110) + (1728 + 314)$, or $a = 1940$; and $(a + c) - (a - c) = 2c = 314 - 110$, or $c = 102$. Thus, $b^2 = a^2 - c^2 = 3{,}753{,}196$, and the equation is $\dfrac{x^2}{3{,}763{,}600} + \dfrac{y^2}{3{,}753{,}196} = 1$.

50. (a) Choose V to be the origin, with x-axis through V and F. Then F is $(p, 0)$, A is $(p, 5)$, so substituting A into the equation $y^2 = 4px$ gives $25 = 4p^2$ so $p = \frac{5}{2}$ and $y^2 = 10x$.

(b) $x = 11 \;\Rightarrow\; y = \sqrt{110} \;\Rightarrow\; |CD| = 2\sqrt{110}$

51. (a) Set up the coordinate system so that A is $(-200, 0)$ and B is $(200, 0)$.

$|PA| - |PB| = (1200)(980) = 1{,}176{,}000$ ft $= \frac{2450}{11}$ mi $= 2a \;\Rightarrow\; a = \frac{1225}{11}$, and $c = 200$ so

$b^2 = c^2 - a^2 = \dfrac{3{,}339{,}375}{121} \;\Rightarrow\; \dfrac{121x^2}{1{,}500{,}625} - \dfrac{121y^2}{3{,}339{,}375} = 1.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Due north of B $\Rightarrow$ $x = 200$ $\Rightarrow$ $\dfrac{(121)(200)^2}{1{,}500{,}625} - \dfrac{121y^2}{3{,}339{,}375} = 1$ $\Rightarrow$ $y = \dfrac{133{,}575}{539} \approx 248$ mi

52. $|PF_1| - |PF_2| = \pm 2a \Leftrightarrow \sqrt{(x+c)^2+y^2} - \sqrt{(x-c)^2+y^2} = \pm 2a \Leftrightarrow$

$\sqrt{(x+c)^2+y^2} = \sqrt{(x-c)^2+y^2} \pm 2a \Leftrightarrow (x+c)^2 + y^2 = (x-c)^2 + y^2 + 4a^2 \pm 4a\sqrt{(x-c)^2+y^2} \Leftrightarrow$

$4cx - 4a^2 = \pm 4a\sqrt{(x-c)^2+y^2} \Leftrightarrow c^2x^2 - 2a^2cx + a^4 = a^2(x^2 - 2cx + c^2 + y^2) \Leftrightarrow$

$(c^2-a^2)x^2 - a^2y^2 = a^2(c^2-a^2) \Leftrightarrow b^2x^2 - a^2y^2 = a^2b^2$ [where $b^2 = c^2 - a^2$] $\Leftrightarrow \dfrac{x^2}{a^2} - \dfrac{y^2}{b^2} = 1$

53. The function whose graph is the upper branch of this hyperbola is concave upward. The function is

$y = f(x) = a\sqrt{1 + \dfrac{x^2}{b^2}} = \dfrac{a}{b}\sqrt{b^2+x^2}$, so $y' = \dfrac{a}{b}x(b^2+x^2)^{-1/2}$ and

$y'' = \dfrac{a}{b}\left[(b^2+x^2)^{-1/2} - x^2(b^2+x^2)^{-3/2}\right] = ab(b^2+x^2)^{-3/2} > 0$ for all x, and so f is concave upward.

54. We can follow exactly the same sequence of steps as in the derivation of Formula 4, except we use the points $(1,1)$ and $(-1,-1)$ in the distance formula (first equation of that derivation) so $\sqrt{(x-1)^2+(y-1)^2} + \sqrt{(x+1)^2+(y+1)^2} = 4$ will lead (after moving the second term to the right, squaring, and simplifying) to $2\sqrt{(x+1)^2+(y+1)^2} = x + y + 4$, which, after squaring and simplifying again, leads to $3x^2 - 2xy + 3y^2 = 8$.

55. (a) If $k > 16$, then $k - 16 > 0$, and $\dfrac{x^2}{k} + \dfrac{y^2}{k-16} = 1$ is an *ellipse* since it is the sum of two squares on the left side.

(b) If $0 < k < 16$, then $k - 16 < 0$, and $\dfrac{x^2}{k} + \dfrac{y^2}{k-16} = 1$ is a *hyperbola* since it is the difference of two squares on the left side.

(c) If $k < 0$, then $k - 16 < 0$, and there is *no curve* since the left side is the sum of two negative terms, which cannot equal 1.

(d) In case (a), $a^2 = k$, $b^2 = k - 16$, and $c^2 = a^2 - b^2 = 16$, so the foci are at $(\pm 4, 0)$. In case (b), $k - 16 < 0$, so $a^2 = k$, $b^2 = 16 - k$, and $c^2 = a^2 + b^2 = 16$, and so again the foci are at $(\pm 4, 0)$.

56. (a) $y^2 = 4px \Rightarrow 2yy' = 4p \Rightarrow y' = \dfrac{2p}{y}$, so the tangent line is

$y - y_0 = \dfrac{2p}{y_0}(x - x_0) \Rightarrow yy_0 - y_0^2 = 2p(x - x_0) \Leftrightarrow$

$yy_0 - 4px_0 = 2px - 2px_0 \Rightarrow yy_0 = 2p(x + x_0)$.



(b) The x-intercept is $-x_0$.

57. $x^2 = 4py \Rightarrow 2x = 4py' \Rightarrow y' = \dfrac{x}{2p}$, so the tangent line at (x_0, y_0) is $y - \dfrac{x_0^2}{4p} = \dfrac{x_0}{2p}(x - x_0)$. This line passes through the point $(a, -p)$ on the directrix, so $-p - \dfrac{x_0^2}{4p} = \dfrac{x_0}{2p}(a - x_0) \Rightarrow -4p^2 - x_0^2 = 2ax_0 - 2x_0^2 \Leftrightarrow$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$x_0^2 - 2ax_0 - 4p^2 = 0 \;\Leftrightarrow\; x_0^2 - 2ax_0 + a^2 = a^2 + 4p^2 \;\Leftrightarrow$

$(x_0 - a)^2 = a^2 + 4p^2 \;\Leftrightarrow\; x_0 = a \pm \sqrt{a^2 + 4p^2}$. The slopes of the tangent lines at $x = a \pm \sqrt{a^2 + 4p^2}$ are $\dfrac{a \pm \sqrt{a^2 + 4p^2}}{2p}$, so the product of the two slopes is



$$\frac{a + \sqrt{a^2 + 4p^2}}{2p} \cdot \frac{a - \sqrt{a^2 + 4p^2}}{2p} = \frac{a^2 - (a^2 + 4p^2)}{4p^2} = \frac{-4p^2}{4p^2} = -1,$$

showing that the tangent lines are perpendicular.

58. Without a loss of generality, let the ellipse, hyperbola, and foci be as shown in the figure.



The curves intersect (eliminate y^2) $\Rightarrow$

$$B^2\left(\frac{x^2}{A^2} - \frac{y^2}{B^2}\right) + b^2\left(\frac{x^2}{a^2} + \frac{y^2}{b^2}\right) = B^2 + b^2 \;\Rightarrow$$

$$\frac{B^2x^2}{A^2} + \frac{b^2x^2}{a^2} = B^2 + b^2 \;\Rightarrow\; x^2\left(\frac{B^2}{A^2} + \frac{b^2}{a^2}\right) = B^2 + b^2 \;\Rightarrow$$

$$x^2 = \frac{B^2 + b^2}{\dfrac{a^2B^2 + b^2A^2}{A^2a^2}} = \frac{A^2a^2(B^2 + b^2)}{a^2B^2 + b^2A^2}.$$

Similarly, $y^2 = \dfrac{B^2b^2(a^2 - A^2)}{b^2A^2 + a^2B^2}$.

Next we find the slopes of the tangent lines of the curves: $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \;\Rightarrow\; \dfrac{2x}{a^2} + \dfrac{2yy'}{b^2} = 0 \;\Rightarrow\; \dfrac{yy'}{b^2} = -\dfrac{x}{a^2} \;\Rightarrow$

$y'_E = -\dfrac{b^2}{a^2}\dfrac{x}{y}$ and $\dfrac{x^2}{A^2} - \dfrac{y^2}{B^2} = 1 \;\Rightarrow\; \dfrac{2x}{A^2} - \dfrac{2yy'}{B^2} = 0 \;\Rightarrow\; \dfrac{yy'}{B^2} = \dfrac{x}{A^2} \;\Rightarrow\; y'_H = \dfrac{B^2}{A^2}\dfrac{x}{y}$. The product of the slopes at (x_0, y_0) is $y'_E\,y'_H = -\dfrac{b^2B^2x_0^2}{a^2A^2y_0^2} = -\dfrac{b^2B^2\left[\dfrac{A^2a^2(B^2 + b^2)}{a^2B^2 + b^2A^2}\right]}{a^2A^2\left[\dfrac{B^2b^2(a^2 - A^2)}{b^2A^2 + a^2B^2}\right]} = -\dfrac{B^2 + b^2}{a^2 - A^2}$. Since $a^2 - b^2 = c^2$ and $A^2 + B^2 = c^2$,

we have $a^2 - b^2 = A^2 + B^2 \;\Rightarrow\; a^2 - A^2 = b^2 + B^2$, so the product of the slopes is -1, and hence, the tangent lines at each point of intersection are perpendicular.

59. $9x^2 + 4y^2 = 36 \;\Leftrightarrow\; \dfrac{x^2}{4} + \dfrac{y^2}{9} = 1$. We use the parametrization $x = 2\cos t$, $y = 3\sin t$, $0 \le t \le 2\pi$. The circumference is given by

$$L = \int_0^{2\pi} \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^{2\pi} \sqrt{(-2\sin t)^2 + (3\cos t)^2}\,dt$$
$$= \int_0^{2\pi} \sqrt{4\sin^2 t + 9\cos^2 t}\,dt = \int_0^{2\pi} \sqrt{4 + 5\cos^2 t}\,dt$$

Now use Simpson's Rule with $n = 8$, $\Delta t = \dfrac{2\pi - 0}{8} = \dfrac{\pi}{4}$, and $f(t) = \sqrt{4 + 5\cos^2 t}$ to get

$$L \approx S_8 = \tfrac{\pi/4}{3}\left[f(0) + 4f\left(\tfrac{\pi}{4}\right) + 2f\left(\tfrac{\pi}{2}\right) + 4f\left(\tfrac{3\pi}{4}\right) + 2f(\pi) + 4f\left(\tfrac{5\pi}{4}\right) + 2f\left(\tfrac{3\pi}{2}\right) + 4f\left(\tfrac{7\pi}{4}\right) + f(2\pi)\right] \approx 15.9.$$

60. The length of the major axis is $2a$, so $a = \frac{1}{2}(1.18 \times 10^{10}) = 5.9 \times 10^9$. The length of the minor axis is $2b$, so $b = \frac{1}{2}(1.14 \times 10^{10}) = 5.7 \times 10^9$. An equation of the ellipse is $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$, or converting into parametric equations,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x = a\cos\theta$ and $y = b\sin\theta$. So

$$L = 4\int_0^{\pi/2} \sqrt{(dx/d\theta)^2 + (dy/d\theta)^2}\, d\theta = 4\int_0^{\pi/2} \sqrt{a^2\sin^2\theta + b^2\cos^2\theta}\, d\theta$$

Using Simpson's Rule with $n = 10$, $\Delta\theta = \frac{\pi/2 - 0}{10} = \frac{\pi}{20}$, and $f(\theta) = \sqrt{a^2\sin^2\theta + b^2\cos^2\theta}$, we get

$$L \approx 4 \cdot S_{10} = 4 \cdot \tfrac{\pi}{20\cdot 3}\left[f(0) + 4f\left(\tfrac{\pi}{20}\right) + 2f\left(\tfrac{2\pi}{20}\right) + \cdots + 2f\left(\tfrac{8\pi}{20}\right) + 4f\left(\tfrac{9\pi}{20}\right) + f\left(\tfrac{\pi}{2}\right)\right] \approx 3.64 \times 10^{10} \text{ km}$$

61. $\dfrac{x^2}{a^2} - \dfrac{y^2}{b^2} = 1 \;\Rightarrow\; \dfrac{y^2}{b^2} = \dfrac{x^2 - a^2}{a^2} \;\Rightarrow\; y = \pm\dfrac{b}{a}\sqrt{x^2 - a^2}$.



$$A = 2\int_a^c \frac{b}{a}\sqrt{x^2 - a^2}\, dx \overset{39}{=} \frac{2b}{a}\left[\frac{x}{2}\sqrt{x^2 - a^2} - \frac{a^2}{2}\ln\left|x + \sqrt{x^2 - a^2}\right|\right]_a^c$$

$$= \frac{b}{a}\left[c\sqrt{c^2 - a^2} - a^2\ln\left|c + \sqrt{c^2 - a^2}\right| + a^2\ln|a|\right]$$

Since $a^2 + b^2 = c^2$, $c^2 - a^2 = b^2$, and $\sqrt{c^2 - a^2} = b$.

$$= \frac{b}{a}\left[cb - a^2\ln(c + b) + a^2\ln a\right] = \frac{b}{a}\left[cb + a^2(\ln a - \ln(b + c))\right]$$

$$= b^2c/a + ab\ln[a/(b + c)], \text{ where } c^2 = a^2 + b^2.$$

62. (a) $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \;\Rightarrow\; \dfrac{y^2}{b^2} = \dfrac{a^2 - x^2}{a^2} \;\Rightarrow\; y = \pm\dfrac{b}{a}\sqrt{a^2 - x^2}$.



$$V = \int_{-a}^{a} \pi\left(\frac{b}{a}\sqrt{a^2 - x^2}\right)^2 dx = 2\pi\frac{b^2}{a^2}\int_0^a (a^2 - x^2)\, dx$$

$$= \frac{2\pi b^2}{a^2}\left[a^2x - \tfrac{1}{3}x^3\right]_0^a = \frac{2\pi b^2}{a^2}\left(\frac{2a^3}{3}\right) = \frac{4}{3}\pi b^2 a$$

(b) $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \;\Rightarrow\; \dfrac{x^2}{a^2} = \dfrac{b^2 - y^2}{b^2} \;\Rightarrow\; x = \pm\dfrac{a}{b}\sqrt{b^2 - y^2}$.



$$V = \int_{-b}^{b} \pi\left(\frac{a}{b}\sqrt{b^2 - y^2}\right)^2 dy = 2\pi\frac{a^2}{b^2}\int_0^b (b^2 - y^2)\, dy$$

$$= \frac{2\pi a^2}{b^2}\left[b^2y - \tfrac{1}{3}y^3\right]_0^b = \frac{2\pi a^2}{b^2}\left(\frac{2b^3}{3}\right) = \frac{4}{3}\pi a^2 b$$

63. $9x^2 + 4y^2 = 36 \;\Leftrightarrow\; \dfrac{x^2}{4} + \dfrac{y^2}{9} = 1 \;\Rightarrow\; a = 3, b = 2$. By symmetry, $\overline{x} = 0$. By Example 2 in Section 7.3, the area of the top half of the ellipse is $\frac{1}{2}(\pi ab) = 3\pi$. Solve $9x^2 + 4y^2 = 36$ for y to get an equation for the top half of the ellipse:

$9x^2 + 4y^2 = 36 \;\Leftrightarrow\; 4y^2 = 36 - 9x^2 \;\Leftrightarrow\; y^2 = \frac{9}{4}(4 - x^2) \;\Rightarrow\; y = \frac{3}{2}\sqrt{4 - x^2}$. Now

$$\overline{y} = \frac{1}{A}\int_a^b \frac{1}{2}[f(x)]^2\, dx = \frac{1}{3\pi}\int_{-2}^{2} \frac{1}{2}\left(\frac{3}{2}\sqrt{4 - x^2}\right)^2 dx = \frac{3}{8\pi}\int_{-2}^{2}(4 - x^2)\, dx$$



$$= \frac{3}{8\pi}\cdot 2\int_0^2 (4 - x^2)\, dx = \frac{3}{4\pi}\left[4x - \frac{1}{3}x^3\right]_0^2 = \frac{3}{4\pi}\left(\frac{16}{3}\right) = \frac{4}{\pi}$$

so the centroid is $(0, 4/\pi)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

64. (a) Consider the ellipse $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1$ with $a>b$, so that the major axis is the x-axis. Let the ellipse be parametrized by $x=a\cos t,\ y=b\sin t,\ 0\le t\le 2\pi$. Then

$$\left(\frac{dx}{dt}\right)^2+\left(\frac{dy}{dt}\right)^2=a^2\sin^2 t+b^2\cos^2 t=a^2(1-\cos^2 t)+b^2\cos^2 t=a^2+(b^2-a^2)\cos^2 t=a^2-c^2\cos^2 t,$$

where $c^2=a^2-b^2$. Using symmetry and rotating the ellipse about the major axis gives us surface area

$$S=\int 2\pi y\,ds=2\int_0^{\pi/2}2\pi(b\sin t)\sqrt{a^2-c^2\cos^2 t}\,dt=4\pi b\int_c^0\sqrt{a^2-u^2}\left(-\frac{1}{c}\,du\right)\qquad\left[\begin{matrix}u=c\cos t\\ du=-c\sin t\,dt\end{matrix}\right]$$

$$=\frac{4\pi b}{c}\int_0^c\sqrt{a^2-u^2}\,du\overset{30}{=}\frac{4\pi b}{c}\left[\frac{u}{2}\sqrt{a^2-u^2}+\frac{a^2}{2}\sin^{-1}\left(\frac{u}{a}\right)\right]_0^c=\frac{2\pi b}{c}\left[c\sqrt{a^2-c^2}+a^2\sin^{-1}\left(\frac{c}{a}\right)\right]$$

$$=\frac{2\pi b}{c}\left[bc+a^2\sin^{-1}\left(\frac{c}{a}\right)\right]$$

(b) As in part (a),

$$\left(\frac{dx}{dt}\right)^2+\left(\frac{dy}{dt}\right)^2=a^2\sin^2 t+b^2\cos^2 t=a^2\sin^2 t+b^2(1-\sin^2 t)=b^2+(a^2-b^2)\sin^2 t=b^2+c^2\sin^2 t.$$

Rotating about the minor axis gives us

$$S=\int 2\pi x\,ds=2\int_0^{\pi/2}2\pi(a\cos t)\sqrt{b^2+c^2\sin^2 t}\,dt=4\pi a\int_0^c\sqrt{b^2+u^2}\left(\frac{1}{c}\,du\right)\qquad\left[\begin{matrix}u=c\sin t\\ du=c\cos t\,dt\end{matrix}\right]$$

$$\overset{21}{=}\frac{4\pi a}{c}\left[\frac{u}{2}\sqrt{b^2+u^2}+\frac{b^2}{2}\ln\left(u+\sqrt{b^2+u^2}\right)\right]_0^c=\frac{2\pi a}{c}\left[c\sqrt{b^2+c^2}+b^2\ln\left(c+\sqrt{b^2+c^2}\right)-b^2\ln b\right]$$

$$=\frac{2\pi a}{c}\left[ac+b^2\ln\left(\frac{a+c}{b}\right)\right]$$

65. Differentiating implicitly, $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1\ \Rightarrow\ \dfrac{2x}{a^2}+\dfrac{2yy'}{b^2}=0\ \Rightarrow\ y'=-\dfrac{b^2x}{a^2y}$ $[y\ne 0]$. Thus, the slope of the tangent line at P is $-\dfrac{b^2x_1}{a^2y_1}$. The slope of F_1P is $\dfrac{y_1}{x_1+c}$ and of F_2P is $\dfrac{y_1}{x_1-c}$. By the formula from Problems Plus, we have

$$\tan\alpha=\frac{\dfrac{y_1}{x_1+c}+\dfrac{b^2x_1}{a^2y_1}}{1-\dfrac{b^2x_1y_1}{a^2y_1(x_1+c)}}=\frac{a^2y_1^2+b^2x_1(x_1+c)}{a^2y_1(x_1+c)-b^2x_1y_1}=\frac{a^2b^2+b^2cx_1}{c^2x_1y_1+a^2cy_1}\qquad\left[\begin{matrix}\text{using } b^2x_1^2+a^2y_1^2=a^2b^2,\\ \text{and } a^2-b^2=c^2\end{matrix}\right]$$

$$=\frac{b^2(cx_1+a^2)}{cy_1(cx_1+a^2)}=\frac{b^2}{cy_1}$$

and $$\tan\beta=\frac{-\dfrac{b^2x_1}{a^2y_1}-\dfrac{y_1}{x_1-c}}{1-\dfrac{b^2x_1y_1}{a^2y_1(x_1-c)}}=\frac{-a^2y_1^2-b^2x_1(x_1-c)}{a^2y_1(x_1-c)-b^2x_1y_1}=\frac{-a^2b^2+b^2cx_1}{c^2x_1y_1-a^2cy_1}=\frac{b^2(cx_1-a^2)}{cy_1(cx_1-a^2)}=\frac{b^2}{cy_1}$$

Thus, $\alpha=\beta$.

66. The slopes of the line segments F_1P and F_2P are $\dfrac{y_1}{x_1+c}$ and $\dfrac{y_1}{x_1-c}$, where P is (x_1,y_1). Differentiating implicitly,

$\dfrac{2x}{a^2}-\dfrac{2yy'}{b^2}=0\ \Rightarrow\ y'=\dfrac{b^2x}{a^2y}\ \Rightarrow$ the slope of the tangent at P is $\dfrac{b^2x_1}{a^2y_1}$, so by the formula in Problem 17 on text

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

page 201 [ET Problem 21 on text page 273] ,

$$\tan\alpha = \frac{\dfrac{b^2x_1}{a^2y_1} - \dfrac{y_1}{x_1+c}}{1+\dfrac{b^2x_1y_1}{a^2y_1(x_1+c)}} = \frac{b^2x_1(x_1+c)-a^2y_1^2}{a^2y_1(x_1+c)+b^2x_1y_1} = \frac{b^2(cx_1+a^2)}{cy_1(cx_1+a^2)} \quad \left[\begin{array}{c}\text{using } x_1^2/a^2 - y_1^2/b^2 = 1,\\ \text{and } a^2+b^2=c^2\end{array}\right] = \frac{b^2}{cy_1}$$

and
$$\tan\beta = \frac{-\dfrac{b^2x_1}{a^2y_1} + \dfrac{y_1}{x_1-c}}{1+\dfrac{b^2x_1y_1}{a^2y_1(x_1-c)}} = \frac{-b^2x_1(x_1-c)+a^2y_1^2}{a^2y_1(x_1-c)+b^2x_1y_1} = \frac{b^2(cx_1-a^2)}{cy_1(cx_1-a^2)} = \frac{b^2}{cy_1}$$

So $\alpha = \beta$.

10.6 Conic Sections in Polar Coordinates

1. The directrix $x = 4$ is to the right of the focus at the origin, so we use the form with "$+\ e\cos\theta$" in the denominator. (See Theorem 6 and Figure 2.) An equation of the ellipse is $r = \dfrac{ed}{1+e\cos\theta} = \dfrac{\frac{1}{2}\cdot 4}{1+\frac{1}{2}\cos\theta} = \dfrac{4}{2+\cos\theta}$.

2. The directrix $x = -3$ is to the left of the focus at the origin, so we use the form with "$-\ e\cos\theta$" in the denominator. $e = 1$ for a parabola, so an equation is $r = \dfrac{ed}{1-e\cos\theta} = \dfrac{1\cdot 3}{1-1\cos\theta} = \dfrac{3}{1-\cos\theta}$.

3. The directrix $y = 2$ is above the focus at the origin, so we use the form with "$+\ e\sin\theta$" in the denominator. An equation of the hyperbola is $r = \dfrac{ed}{1+e\sin\theta} = \dfrac{1.5(2)}{1+1.5\sin\theta} = \dfrac{6}{2+3\sin\theta}$.

4. The directrix $x = 3$ is to the right of the focus at the origin, so we use the form with "$+\ e\cos\theta$" in the denominator. An equation of the hyperbola is $r = \dfrac{ed}{1+e\cos\theta} = \dfrac{3\cdot 3}{1+3\cos\theta} = \dfrac{9}{1+3\cos\theta}$.

5. The vertex $(2,\pi)$ is to the left of the focus at the origin, so we use the form with "$-e\cos\theta$" in the denominator. An equation of the ellipse is $r = \dfrac{ed}{1-e\cos\theta}$. Using eccentricity $e = \dfrac{2}{3}$ with $\theta = \pi$ and $r = 2$, we get $2 = \dfrac{\frac{2}{3}d}{1-\frac{2}{3}(-1)} \Rightarrow$ $2 = \dfrac{2d}{5} \Rightarrow d = 5$, so we have $r = \dfrac{\frac{2}{3}(5)}{1-\frac{2}{3}\cos\theta} = \dfrac{10}{3-2\cos\theta}$.

6. The directrix $r = 4\csc\theta$ (equivalent to $r\sin\theta = 4$ or $y = 4$) is above the focus at the origin, so we will use the form with "$+e\sin\theta$" in the denominator. The distance from the focus to the directrix is $d = 4$, so an equation of the ellipse is $r = \dfrac{ed}{1+e\sin\theta} = \dfrac{(0.6)(4)}{1+0.6\sin\theta}\cdot\dfrac{5}{5} = \dfrac{12}{5+3\sin\theta}$.

7. The vertex $\left(3,\frac{\pi}{2}\right)$ is 3 units above the focus at the origin, so the directrix is 6 units above the focus ($d = 6$), and we use the form "$+e\sin\theta$" in the denominator. $e = 1$ for a parabola, so an equation is $r = \dfrac{ed}{1+e\sin\theta} = \dfrac{1(6)}{1+1\sin\theta} = \dfrac{6}{1+\sin\theta}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. The directrix $r = -2\sec\theta$ (equivalent to $r\cos\theta = -2$ or $x = -2$) is left of the focus at the origin, so we will use the form with "$-e\cos\theta$" in the denominator. The distance from the focus to the directrix is $d = 2$, so an equation of the hyperbola is $r = \dfrac{ed}{1 - e\cos\theta} = \dfrac{2(2)}{1 - 2\cos\theta} = \dfrac{4}{1 - 2\cos\theta}$.

9. $r = \dfrac{4}{5 - 4\sin\theta} \cdot \dfrac{1/5}{1/5} = \dfrac{4/5}{1 - \frac{4}{5}\sin\theta}$, where $e = \frac{4}{5}$ and $ed = \frac{4}{5} \quad \Rightarrow \quad d = 1$.

(a) Eccentricity $= e = \frac{4}{5}$

(b) Since $e = \frac{4}{5} < 1$, the conic is an ellipse.

(c) Since "$-e\sin\theta$" appears in the denominator, the directrix is below the focus at the origin, $d = |Fl| = 1$, so an equation of the directrix is $y = -1$.

(d) The vertices are $\left(4, \frac{\pi}{2}\right)$ and $\left(\frac{4}{9}, \frac{3\pi}{2}\right)$.



10. $r = \dfrac{1}{2 + \sin\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{1/2}{1 + \frac{1}{2}\sin\theta}$, where $e = \dfrac{1}{2}$ and $ed = \dfrac{1}{2} \quad \Rightarrow \quad d = 1$.

(a) Eccentricity $= e = \dfrac{1}{2}$

(b) Since $e = \dfrac{1}{2} < 1$, the conic is an ellipse.

(c) Since "$+e\sin\theta$" appears in the denominator, the directrix is above the focus at the origin, $d = |Fl| = 1$, so an equation of the directrix is $y = 1$.

(d) The vertices are $\left(\frac{1}{3}, \frac{\pi}{2}\right)$ and $\left(1, \frac{3\pi}{2}\right)$.



11. $r = \dfrac{2}{3 + 3\sin\theta} \cdot \dfrac{1/3}{1/3} = \dfrac{2/3}{1 + 1\sin\theta}$, where $e = 1$ and $ed = \frac{2}{3} \quad \Rightarrow \quad d = \frac{2}{3}$.

(a) Eccentricity $= e = 1$

(b) Since $e = 1$, the conic is a parabola.

(c) Since "$+e\sin\theta$" appears in the denominator, the directrix is above the focus at the origin. $d = |Fl| = \frac{2}{3}$, so an equation of the directrix is $y = \frac{2}{3}$.

(d) The vertex is at $\left(\frac{1}{3}, \frac{\pi}{2}\right)$, midway between the focus and directrix.



12. $r = \dfrac{5}{2 - 4\cos\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{5/2}{1 - 2\cos\theta}$, where $e = 2$ and $ed = \dfrac{5}{2} \quad \Rightarrow \quad d = \dfrac{5}{4}$.

(a) Eccentricity $= e = 2$

(b) Since $e = 2 > 1$, the conic is a hyperbola.

(c) Since "$-e\cos\theta$" appears in the denominator, the directrix is to the left the focus at the origin. $d = |Fl| = \frac{5}{4}$, so an equation of the directrix is $x = -\frac{5}{4}$.

(d) The vertices are $\left(-\frac{5}{2}, 0\right)$ and $\left(\frac{5}{6}, \pi\right)$, so the center is midway between them, that is, $\left(\frac{5}{3}, \pi\right)$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $r = \dfrac{9}{6+2\cos\theta} \cdot \dfrac{1/6}{1/6} = \dfrac{3/2}{1+\frac{1}{3}\cos\theta}$, where $e = \frac{1}{3}$ and $ed = \frac{3}{2} \Rightarrow d = \frac{9}{2}$.

(a) Eccentricity $= e = \frac{1}{3}$

(b) Since $e = \frac{1}{3} < 1$, the conic is an ellipse.

(c) Since "$+e\cos\theta$" appears in the denominator, the directrix is to the right of the focus at the origin. $d = |Fl| = \frac{9}{2}$, so an equation of the directrix is $x = \frac{9}{2}$.

(d) The vertices are $\left(\frac{9}{8}, 0\right)$ and $\left(\frac{9}{4}, \pi\right)$, so the center is midway between them, that is, $\left(\frac{9}{16}, \pi\right)$.



14. $r = \dfrac{1}{3-3\sin\theta} \cdot \dfrac{1/3}{1/3} = \dfrac{1/3}{1-1\sin\theta}$, where $e = 1$ and $ed = \dfrac{1}{3} \Rightarrow d = \dfrac{1}{3}$

(a) Eccentricity $= e = 1$

(b) Since $e = 1$, the conic is a parabola.

(c) Since "$-e\sin\theta$" appears in the denominator, the directrix is below the focus at the origin, $d = |Fl| = \frac{1}{3}$, so an equation of the directrix is $y = -\frac{1}{3}$.

(d) The vertex is at $\left(\frac{1}{6}, \frac{3\pi}{2}\right)$, midway between the focus and the directrix.



15. $r = \dfrac{3}{4-8\cos\theta} \cdot \dfrac{1/4}{1/4} = \dfrac{3/4}{1-2\cos\theta}$, where $e = 2$ and $ed = \frac{3}{4} \Rightarrow d = \frac{3}{8}$.

(a) Eccentricity $= e = 2$

(b) Since $e = 2 > 1$, the conic is a hyperbola.

(c) Since "$-e\cos\theta$" appears in the denominator, the directrix is to the left of the focus at the origin. $d = |Fl| = \frac{3}{8}$, so an equation of the directrix is $x = -\frac{3}{8}$.

(d) The vertices are $\left(-\frac{3}{4}, 0\right)$ and $\left(\frac{1}{4}, \pi\right)$, so the center is midway between them, that is, $\left(\frac{1}{2}, \pi\right)$.



16. $r = \dfrac{4}{2+3\cos\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{2}{1+\frac{3}{2}\cos\theta}$, where $e = \dfrac{3}{2}$ and $ed = 2 \Rightarrow d = \dfrac{4}{3}$.

(a) Eccentricity $= e = \dfrac{3}{2}$

(b) Since $e = \dfrac{3}{2} > 1$, the conic is a hyperbola.

(c) Since "$+e\cos\theta$" appears in the denominator, the directrix is to the right of the focus at the origin. $d = |Fl| = \frac{4}{3}$, so an equation of the directrix is $x = \frac{4}{3}$.

(d) The vertices are $\left(\frac{4}{5}, 0\right)$ and $(-4, \pi)$, so the center is midway between them, that is, $\left(\frac{8}{5}, 0\right)$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

17. (a) $r = \dfrac{1}{1-2\sin\theta}$, where $e = 2$ and $ed = 1 \;\Rightarrow\; d = \frac{1}{2}$. The eccentricity $e = 2 > 1$, so the conic is a hyperbola. Since "$-e\sin\theta$" appears in the denominator, the directrix is below the focus at the origin. $d = |Fl| = \frac{1}{2}$, so an equation of the directrix is $y = -\frac{1}{2}$. The vertices are $\left(-1, \frac{\pi}{2}\right)$ and $\left(\frac{1}{3}, \frac{3\pi}{2}\right)$, so the center is midway between them, that is, $\left(\frac{2}{3}, \frac{3\pi}{2}\right)$.



(b) By the discussion that precedes Example 4, the equation is $r = \dfrac{1}{1-2\sin\left(\theta - \frac{3\pi}{4}\right)}$.



18. $r = \dfrac{4}{5+6\cos\theta} = \dfrac{4/5}{1+\frac{6}{5}\cos\theta}$, so $e = \frac{6}{5}$ and $ed = \frac{4}{5} \;\Rightarrow\; d = \frac{2}{3}$.

An equation of the directrix is $x = \frac{2}{3} \;\Rightarrow\; r\cos\theta = \frac{2}{3} \;\Rightarrow\; r = \dfrac{2}{3\cos\theta}$.

If the hyperbola is rotated about its focus (the origin) through an angle $\pi/3$, its equation is the same as that of the original, with θ replaced by $\theta - \frac{\pi}{3}$ (see Example 4), so $r = \dfrac{4}{5+6\cos\left(\theta - \frac{\pi}{3}\right)}$.



19. For $e < 1$ the curve is an ellipse. It is nearly circular when e is close to 0. As e increases, the graph is stretched out to the right, and grows larger (that is, its right-hand focus moves to the right while its left-hand focus remains at the origin.) At $e = 1$, the curve becomes a parabola with focus at the origin.



20. (a) The value of d does not seem to affect the shape of the conic (a parabola) at all, just its size, position, and orientation (for $d < 0$ it opens upward, for $d > 0$ it opens downward).



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) We consider only positive values of e. When $0 < e < 1$, the conic is an ellipse. As $e \to 0^+$, the graph approaches perfect roundness and zero size. As e increases, the ellipse becomes more elongated, until at $e = 1$ it turns into a parabola. For $e > 1$, the conic is a hyperbola, which moves downward and gets broader as e continues to increase.



$e = 0.1$



$e = 0.5$



$e = 0.9$



$e = 1$



$e = 1.1$



$e = 1.5$



$e = 10$

21. $|PF| = e\,|Pl| \;\Rightarrow\; r = e[d - r\cos(\pi - \theta)] = e(d + r\cos\theta) \;\Rightarrow$

$r(1 - e\cos\theta) = ed \;\Rightarrow\; r = \dfrac{ed}{1 - e\cos\theta}$



22. $|PF| = e\,|Pl| \;\Rightarrow\; r = e[d - r\sin\theta] \;\Rightarrow\; r(1 + e\sin\theta) = ed \;\Rightarrow$

$r = \dfrac{ed}{1 + e\sin\theta}$



23. $|PF| = e\,|Pl| \;\Rightarrow\; r = e[d - r\sin(\theta - \pi)] = e(d + r\sin\theta) \;\Rightarrow$

$r(1 - e\sin\theta) = ed \;\Rightarrow\; r = \dfrac{ed}{1 - e\sin\theta}$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. The parabolas intersect at the two points where $\dfrac{c}{1+\cos\theta} = \dfrac{d}{1-\cos\theta} \quad \Rightarrow \quad \cos\theta = \dfrac{c-d}{c+d} \quad \Rightarrow \quad r = \dfrac{c+d}{2}$.

For the first parabola, $\dfrac{dr}{d\theta} = \dfrac{c\sin\theta}{(1+\cos\theta)^2}$, so

$$\frac{dy}{dx} = \frac{(dr/d\theta)\sin\theta + r\cos\theta}{(dr/d\theta)\cos\theta - r\sin\theta} = \frac{c\sin^2\theta + c\cos\theta(1+\cos\theta)}{c\sin\theta\cos\theta - c\sin\theta(1+\cos\theta)} = \frac{1+\cos\theta}{-\sin\theta}$$

and similarly for the second, $\dfrac{dy}{dx} = \dfrac{1-\cos\theta}{\sin\theta} = \dfrac{\sin\theta}{1+\cos\theta}$. Since the product of these slopes is -1, the parabolas intersect at right angles.

25. We are given $e = 0.093$ and $a = 2.28 \times 10^8$. By (7), we have

$$r = \frac{a(1-e^2)}{1+e\cos\theta} = \frac{2.28\times10^8[1-(0.093)^2]}{1+0.093\cos\theta} \approx \frac{2.26\times10^8}{1+0.093\cos\theta}$$

26. We are given $e = 0.048$ and $2a = 1.56 \times 10^9 \quad \Rightarrow \quad a = 7.8 \times 10^8$. By (7), we have

$$r = \frac{a(1-e^2)}{1+e\cos\theta} = \frac{7.8\times10^8[1-(0.048)^2]}{1+0.048\cos\theta} \approx \frac{7.78\times10^8}{1+0.048\cos\theta}$$

27. Here $2a =$ length of major axis $= 36.18$ AU $\quad \Rightarrow \quad a = 18.09$ AU and $e = 0.97$. By (7), the equation of the orbit is $r = \dfrac{18.09[1-(0.97)^2]}{1+0.97\cos\theta} \approx \dfrac{1.07}{1+0.97\cos\theta}$. By (8), the maximum distance from the comet to the sun is $18.09(1+0.97) \approx 35.64$ AU or about 3.314 billion miles.

28. Here $2a =$ length of major axis $= 356.5$ AU $\quad \Rightarrow \quad a = 178.25$ AU and $e = 0.9951$. By (7), the equation of the orbit is $r = \dfrac{178.25[1-(0.9951)^2]}{1+0.9951\cos\theta} \approx \dfrac{1.7426}{1+0.9951\cos\theta}$. By (8), the minimum distance from the comet to the sun is $178.25(1-0.9951) \approx 0.8734$ AU or about 81 million miles.

29. The minimum distance is at perihelion, where $4.6\times10^7 = r = a(1-e) = a(1-0.206) = a(0.794) \quad \Rightarrow$ $a = 4.6\times10^7/0.794$. So the maximum distance, which is at aphelion, is $r = a(1+e) = (4.6\times10^7/0.794)(1.206) \approx 7.0\times10^7$ km.

30. At perihelion, $r = a(1-e) = 4.43\times10^9$, and at aphelion, $r = a(1+e) = 7.37\times10^9$. Adding, we get $2a = 11.80\times10^9$, so $a = 5.90\times10^9$ km. Therefore $1+e = a(1+e)/a = \frac{7.37}{5.90} \approx 1.249$ and $e \approx 0.249$.

31. From Exercise 29, we have $e = 0.206$ and $a(1-e) = 4.6\times10^7$ km. Thus, $a = 4.6\times10^7/0.794$. From (7), we can write the equation of Mercury's orbit as $r = a\dfrac{1-e^2}{1+e\cos\theta}$. So since

$$\frac{dr}{d\theta} = \frac{a(1-e^2)e\sin\theta}{(1+e\cos\theta)^2}$$

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \frac{a^2(1-e^2)^2}{(1+e\cos\theta)^2} + \frac{a^2(1-e^2)^2e^2\sin^2\theta}{(1+e\cos\theta)^4} - \frac{a^2(1-e^2)^2}{(1+e\cos\theta)^4}(1+2e\cos\theta+e^2)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

the length of the orbit is

$$L = \int_0^{2\pi} \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = a(1-e^2) \int_0^{2\pi} \frac{\sqrt{1+e^2+2e\cos\theta}}{(1+e\cos\theta)^2}\, d\theta \approx 3.6 \times 10^8 \text{ km}$$

This seems reasonable, since Mercury's orbit is nearly circular, and the circumference of a circle of radius a is $2\pi a \approx 3.6 \times 10^8$ km.

10 Review

TRUE-FALSE QUIZ

1. False. Consider the curve defined by $x = f(t) = (t-1)^3$ and $y = g(t) = (t-1)^2$. Then $g'(t) = 2(t-1)$, so $g'(1) = 0$, but its graph has a *vertical* tangent when $t = 1$. *Note:* The statement is true if $f'(1) \neq 0$ when $g'(1) = 0$.

2. False. If $x = f(t)$ and $y = g(t)$ are twice differentiable, then $\dfrac{d^2y}{dx^2} = \dfrac{d}{dx}\left(\dfrac{dy}{dx}\right) = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{\dfrac{dx}{dt}}$.

3. False. For example, if $f(t) = \cos t$ and $g(t) = \sin t$ for $0 \le t \le 4\pi$, then the curve is a circle of radius 1, hence its length is 2π, but $\int_0^{4\pi} \sqrt{[f'(t)]^2 + [g'(t)]^2}\, dt = \int_0^{4\pi} \sqrt{(-\sin t)^2 + (\cos t)^2}\, dt = \int_0^{4\pi} 1\, dt = 4\pi$, since as t increases from 0 to 4π, the circle is traversed twice.

4. False. If $(r, \theta) = (1, \pi)$, then $(x, y) = (-1, 0)$, so $\tan^{-1}(y/x) = \tan^{-1} 0 = 0 \neq \theta$. The statement is true for points in quadrants I and IV.

5. True. The curve $r = 1 - \sin 2\theta$ is unchanged if we rotate it through $180°$ about O because $1 - \sin 2(\theta + \pi) = 1 - \sin(2\theta + 2\pi) = 1 - \sin 2\theta$. So it's unchanged if we replace r by $-r$. (See the discussion after Example 8 in Section 10.3.) In other words, it's the same curve as $r = -(1 - \sin 2\theta) = \sin 2\theta - 1$.

6. True. The polar equation $r = 2$, the Cartesian equation $x^2 + y^2 = 4$, and the parametric equations $x = 2\sin 3t$, $y = 2\cos 3t$ $[0 \le t \le 2\pi]$ all describe the circle of radius 2 centered at the origin.

7. False. The first pair of equations gives the portion of the parabola $y = x^2$ with $x \ge 0$, whereas the second pair of equations traces out the whole parabola $y = x^2$.

8. True. $y^2 = 2y + 3x \;\Leftrightarrow\; (y-1)^2 = 3x + 1 = 3\left(x + \frac{1}{3}\right) = 4\left(\frac{3}{4}\right)\left(x + \frac{1}{3}\right)$, which is the equation of a parabola with vertex $\left(-\frac{1}{3}, 1\right)$ and focus $\left(-\frac{1}{3} + \frac{3}{4}, 1\right)$, opening to the right.

9. True. By rotating and translating the parabola, we can assume it has an equation of the form $y = cx^2$, where $c > 0$. The tangent at the point (a, ca^2) is the line $y - ca^2 = 2ca(x - a)$; i.e., $y = 2cax - ca^2$. This tangent meets the parabola at the points (x, cx^2) where $cx^2 = 2cax - ca^2$. This equation is equivalent to $x^2 = 2ax - a^2$ [since $c > 0$]. But $x^2 = 2ax - a^2 \;\Leftrightarrow\; x^2 - 2ax + a^2 = 0 \;\Leftrightarrow\; (x-a)^2 = 0 \;\Leftrightarrow\; x = a \;\Leftrightarrow\; (x, cx^2) = (a, ca^2)$. This shows that each tangent meets the parabola at exactly one point.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. True. Consider a hyperbola with focus at the origin, oriented so that its polar equation is $r = \dfrac{ed}{1 + e\cos\theta}$, where $e > 1$.

The directrix is $x = d$, but along the hyperbola we have $x = r\cos\theta = \dfrac{ed\cos\theta}{1 + e\cos\theta} = d\left(\dfrac{e\cos\theta}{1 + e\cos\theta}\right) \neq d$.

EXERCISES

1. $x = t^2 + 4t$, $y = 2 - t$, $-4 \le t \le 1$. $t = 2 - y$, so

$x = (2 - y)^2 + 4(2 - y) = 4 - 4y + y^2 + 8 - 4y = y^2 - 8y + 12 \;\Leftrightarrow$

$x + 4 = y^2 - 8y + 16 = (y - 4)^2$. This is part of a parabola with vertex $(-4, 4)$, opening to the right.



2. $x = 1 + e^{2t}$, $y = e^t$.

$x = 1 + e^{2t} = 1 + (e^t)^2 = 1 + y^2$, $y > 0$.



3. $y = \sec\theta = \dfrac{1}{\cos\theta} = \dfrac{1}{x}$. Since $0 \le \theta \le \pi/2$, $0 < x \le 1$ and $y \ge 1$.

This is part of the hyperbola $y = 1/x$.



4. $x = 2\cos\theta$, $y = 1 + \sin\theta$, $\cos^2\theta + \sin^2\theta = 1 \;\Rightarrow$

$\left(\dfrac{x}{2}\right)^2 + (y - 1)^2 = 1 \;\Rightarrow\; \dfrac{x^2}{4} + (y - 1)^2 = 1$. This is an ellipse, centered at $(0, 1)$, with semimajor axis of length 2 and semiminor axis of length 1.



5. Three different sets of parametric equations for the curve $y = \sqrt{x}$ are

(i) $x = t$, $y = \sqrt{t}$

(ii) $x = t^4$, $y = t^2$

(iii) $x = \tan^2 t$, $y = \tan t$, $0 \le t < \pi/2$

There are many other sets of equations that also give this curve.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. For $t < -1$, $x > 0$ and $y < 0$ with x decreasing and y increasing. When $t = -1$, $(x, y) = (0, 0)$. When $-1 < t < 0$, we have $-1 < x < 0$ and $0 < y < 1/2$. When $t = 0$, $(x, y) = (-1, 0)$. When $0 < t < 1$, $-1 < x < 0$ and $-\frac{1}{2} < y < 0$. When $t = 1$, $(x, y) = (0, 0)$ again. When $t > 1$, both x and y are positive and increasing.



7. (a)



The Cartesian coordinates are $x = 4\cos\frac{2\pi}{3} = 4\left(-\frac{1}{2}\right) = -2$ and $y = 4\sin\frac{2\pi}{3} = 4\left(\frac{\sqrt{3}}{2}\right) = 2\sqrt{3}$, that is, the point $\left(-2, 2\sqrt{3}\right)$.

(b) Given $x = -3$ and $y = 3$, we have $r = \sqrt{(-3)^2 + 3^2} = \sqrt{18} = 3\sqrt{2}$. Also, $\tan\theta = \dfrac{y}{x} \Rightarrow \tan\theta = \dfrac{3}{-3}$, and since $(-3, 3)$ is in the second quadrant, $\theta = \frac{3\pi}{4}$. Thus, one set of polar coordinates for $(-3, 3)$ is $\left(3\sqrt{2}, \frac{3\pi}{4}\right)$, and two others are $\left(3\sqrt{2}, \frac{11\pi}{4}\right)$ and $\left(-3\sqrt{2}, \frac{7\pi}{4}\right)$.

8. $1 \le r < 2$, $\frac{\pi}{6} \le \theta \le \frac{5\pi}{6}$



9. $r = 1 + \sin\theta$. This cardioid is symmetric about the $\theta = \pi/2$ axis.





10. $r = \sin 4\theta$. This is an eight-leaved rose.





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

11. $r = \cos 3\theta$. This is a three-leaved rose. The curve is traced twice.





12. $r = 3 + \cos 3\theta$. The curve is symmetric about the horizontal axis.





13. $r = 1 + \cos 2\theta$. The curve is symmetric about the pole and both the horizontal and vertical axes.





14. $r = 2\cos(\theta/2)$. The curve is symmetric about the pole and both the horizontal and vertical axes.





15. $r = \dfrac{3}{1 + 2\sin\theta} \quad \Rightarrow \quad e = 2 > 1$, so the conic is a hyperbola. $de = 3 \quad \Rightarrow$ $d = \frac{3}{2}$ and the form "$+2\sin\theta$" imply that the directrix is above the focus at the origin and has equation $y = \frac{3}{2}$. The vertices are $\left(1, \frac{\pi}{2}\right)$ and $\left(-3, \frac{3\pi}{2}\right)$.



16. $r = \dfrac{3}{2 - 2\cos\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{3/2}{1 - 1\cos\theta} \quad \Rightarrow \quad e = 1$, so the conic is a parabola. $de = \frac{3}{2} \quad \Rightarrow \quad d = \frac{3}{2}$ and the form "$-2\cos\theta$" imply that the directrix is to the left of the focus at the origin and has equation $x = -\frac{3}{2}$. The vertex is $\left(\frac{3}{4}, \pi\right)$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $x+y=2 \;\Leftrightarrow\; r\cos\theta + r\sin\theta = 2 \;\Leftrightarrow\; r(\cos\theta+\sin\theta)=2 \;\Leftrightarrow\; r=\dfrac{2}{\cos\theta+\sin\theta}$

18. $x^2+y^2=2 \;\Rightarrow\; r^2=2 \;\Rightarrow\; r=\sqrt{2}$. [$r=-\sqrt{2}$ gives the same curve.]

19. $r=(\sin\theta)/\theta$. As $\theta\to\pm\infty$, $r\to 0$. As $\theta\to 0$, $r\to 1$. In the first figure, there are an infinite number of x-intercepts at $x=\pi n$, n a nonzero integer. These correspond to pole points in the second figure.





20. $r=\dfrac{2}{4-3\cos\theta}=\dfrac{1/2}{1-\frac{3}{4}\cos\theta} \;\Rightarrow\; e=\frac{3}{4}$ and $d=\frac{2}{3}$. The equation of the directrix is $x=-\frac{2}{3} \;\Rightarrow\; r=-2/(3\cos\theta)$. To obtain the equation of the rotated ellipse, we replace θ in the original equation with $\theta-\frac{2\pi}{3}$, and get $r=\dfrac{2}{4-3\cos\left(\theta-\frac{2\pi}{3}\right)}$.



21. $x=\ln t$, $y=1+t^2$; $t=1$. $\dfrac{dy}{dt}=2t$ and $\dfrac{dx}{dt}=\dfrac{1}{t}$, so $\dfrac{dy}{dx}=\dfrac{dy/dt}{dx/dt}=\dfrac{2t}{1/t}=2t^2$.

When $t=1$, $(x,y)=(0,2)$ and $dy/dx=2$.

22. $x=t^3+6t+1$, $y=2t-t^2$; $\quad t=-1$. $\dfrac{dy}{dx}=\dfrac{dy/dt}{dx/dt}=\dfrac{2-2t}{3t^2+6}$. When $t=-1$, $(x,y)=(-6,-3)$ and $\dfrac{dy}{dx}=\dfrac{4}{9}$.

23. $r=e^{-\theta} \;\Rightarrow\; y=r\sin\theta=e^{-\theta}\sin\theta$ and $x=r\cos\theta=e^{-\theta}\cos\theta \;\Rightarrow$

$$\frac{dy}{dx}=\frac{dy/d\theta}{dx/d\theta}=\frac{\frac{dr}{d\theta}\sin\theta+r\cos\theta}{\frac{dr}{d\theta}\cos\theta-r\sin\theta}=\frac{-e^{-\theta}\sin\theta+e^{-\theta}\cos\theta}{-e^{-\theta}\cos\theta-e^{-\theta}\sin\theta}\cdot\frac{-e^{\theta}}{-e^{\theta}}=\frac{\sin\theta-\cos\theta}{\cos\theta+\sin\theta}.$$

When $\theta=\pi$, $\dfrac{dy}{dx}=\dfrac{0-(-1)}{-1+0}=\dfrac{1}{-1}=-1$.

24. $r=3+\cos 3\theta \;\Rightarrow\; \dfrac{dy}{dx}=\dfrac{dy/d\theta}{dx/d\theta}=\dfrac{\frac{dr}{d\theta}\sin\theta+r\cos\theta}{\frac{dr}{d\theta}\cos\theta \quad r\sin\theta}=\dfrac{-3\sin 3\theta\sin\theta+(3+\cos 3\theta)\cos\theta}{-3\sin 3\theta\cos\theta-(3+\cos 3\theta)\sin\theta}$.

When $\theta=\pi/2$, $\dfrac{dy}{dx}=\dfrac{(-3)(-1)(1)+(3+0)\cdot 0}{(-3)(-1)(0)-(3+0)\cdot 1}=\dfrac{3}{-3}=-1$.

25. $x=t+\sin t$, $y=t-\cos t \;\Rightarrow\; \dfrac{dy}{dx}=\dfrac{dy/dt}{dx/dt}=\dfrac{1+\sin t}{1+\cos t} \;\Rightarrow$

$$\frac{d^2y}{dx^2}=\frac{\frac{d}{dt}\left(\frac{dy}{dx}\right)}{dx/dt}=\frac{(1+\cos t)\cos t-(1+\sin t)(-\sin t)}{(1+\cos t)^2}\Big/(1+\cos t)=\frac{\cos t+\cos^2 t+\sin t+\sin^2 t}{(1+\cos t)^3}=\frac{1+\cos t+\sin t}{(1+\cos t)^3}$$

26. $x=1+t^2$, $y=t-t^3$. $\dfrac{dy}{dt}=1-3t^2$ and $\dfrac{dx}{dt}=2t$, so $\dfrac{dy}{dx}=\dfrac{dy/dt}{dx/dt}=\dfrac{1-3t^2}{2t}=\frac{1}{2}t^{-1}-\frac{3}{2}t$.

$$\frac{d^2y}{dx^2}=\frac{d(dy/dx)/dt}{dx/dt}=\frac{-\frac{1}{2}t^{-2}-\frac{3}{2}}{2t}=-\frac{1}{4}t^{-3}-\frac{3}{4}t^{-1}=-\frac{1}{4t^3}\left(1+3t^2\right)=-\frac{3t^2+1}{4t^3}.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

27. We graph the curve $x = t^3 - 3t$, $y = t^2 + t + 1$ for $-2.2 \le t \le 1.2$.



By zooming in or using a cursor, we find that the lowest point is about $(1.4, 0.75)$. To find the exact values, we find the t-value at which

$$dy/dt = 2t + 1 = 0 \quad \Leftrightarrow \quad t = -\tfrac{1}{2} \quad \Leftrightarrow \quad (x, y) = \left(\tfrac{11}{8}, \tfrac{3}{4}\right).$$

28. We estimate the coordinates of the point of intersection to be $(-2, 3)$. In fact this is exact, since both $t = -2$ and $t = 1$ give the point $(-2, 3)$. So the area enclosed by the loop is

$$\begin{aligned}\int_{t=-2}^{t=1} y\,dx &= \int_{-2}^{1} (t^2 + t + 1)(3t^2 - 3)\,dt = \int_{-2}^{1} (3t^4 + 3t^3 - 3t - 3)\,dt \\ &= \left[\tfrac{3}{5}t^5 + \tfrac{3}{4}t^4 - \tfrac{3}{2}t^2 - 3t\right]_{-2}^{1} = \left(\tfrac{3}{5} + \tfrac{3}{4} - \tfrac{3}{2} - 3\right) - \left[-\tfrac{96}{5} + 12 - 6 - (-6)\right] = \tfrac{81}{20}\end{aligned}$$

29. $x = 2a\cos t - a\cos 2t \quad \Rightarrow \quad \dfrac{dx}{dt} = -2a\sin t + 2a\sin 2t = 2a\sin t(2\cos t - 1) = 0 \quad \Leftrightarrow$

$\sin t = 0$ or $\cos t = \frac{1}{2} \quad \Rightarrow \quad t = 0, \frac{\pi}{3}, \pi$, or $\frac{5\pi}{3}$.

$y = 2a\sin t - a\sin 2t \quad \Rightarrow \quad \dfrac{dy}{dt} = 2a\cos t - 2a\cos 2t = 2a\big(1 + \cos t - 2\cos^2 t\big) = 2a(1 - \cos t)(1 + 2\cos t) = 0 \quad \Rightarrow$

$t = 0, \frac{2\pi}{3}$, or $\frac{4\pi}{3}$.

Thus the graph has vertical tangents where $t = \frac{\pi}{3}, \pi$ and $\frac{5\pi}{3}$, and horizontal tangents where $t = \frac{2\pi}{3}$ and $\frac{4\pi}{3}$. To determine what the slope is where $t = 0$, we use l'Hospital's Rule to evaluate $\lim\limits_{t\to 0} \dfrac{dy/dt}{dx/dt} = 0$, so there is a horizontal tangent there.

t	x	y
0	a	0
$\frac{\pi}{3}$	$\frac{3}{2}a$	$\frac{\sqrt{3}}{2}a$
$\frac{2\pi}{3}$	$-\frac{1}{2}a$	$\frac{3\sqrt{3}}{2}a$
π	$-3a$	0
$\frac{4\pi}{3}$	$-\frac{1}{2}a$	$-\frac{3\sqrt{3}}{2}a$
$\frac{5\pi}{3}$	$\frac{3}{2}a$	$-\frac{\sqrt{3}}{2}a$



30. From Exercise 29, $x = 2a\cos t - a\cos 2t$, $y = 2a\sin t - a\sin 2t \quad \Rightarrow$

$$\begin{aligned}A &= 2\int_{\pi}^{0} (2a\sin t - a\sin 2t)(-2a\sin t + 2a\sin 2t)\,dt = 4a^2\int_0^{\pi} (2\sin^2 t + \sin^2 2t - 3\sin t\sin 2t)\,dt \\ &= 4a^2\int_0^{\pi} \left[(1 - \cos 2t) + \tfrac{1}{2}(1 - \cos 4t) - 6\sin^2 t\cos t\right] dt = 4a^2\left[t - \tfrac{1}{2}\sin 2t + \tfrac{1}{2}t - \tfrac{1}{8}\sin 4t - 2\sin^3 t\right]_0^{\pi} \\ &= 4a^2\left(\tfrac{3}{2}\right)\pi = 6\pi a^2\end{aligned}$$

31. The curve $r^2 = 9\cos 5\theta$ has 10 "petals." For instance, for $-\frac{\pi}{10} \le \theta \le \frac{\pi}{10}$, there are two petals, one with $r > 0$ and one with $r < 0$.

$$A = 10\int_{-\pi/10}^{\pi/10} \tfrac{1}{2}r^2\,d\theta = 5\int_{-\pi/10}^{\pi/10} 9\cos 5\theta\,d\theta = 5\cdot 9\cdot 2\int_0^{\pi/10} \cos 5\theta\,d\theta = 18\big[\sin 5\theta\big]_0^{\pi/10} = 18$$

32. $r = 1 - 3\sin\theta$. The inner loop is traced out as θ goes from $\alpha = \sin^{-1}\left(\frac{1}{3}\right)$ to $\pi - \alpha$, so

$$\begin{aligned}A &= \int_{\alpha}^{\pi-\alpha} \tfrac{1}{2}r^2\,d\theta = \int_{\alpha}^{\pi/2} (1 - 3\sin\theta)^2\,d\theta = \int_{\alpha}^{\pi/2} \left[1 - 6\sin\theta + \tfrac{9}{2}(1 - \cos 2\theta)\right] d\theta \\ &= \left[\tfrac{11}{2}\theta + 6\cos\theta - \tfrac{9}{4}\sin 2\theta\right]_{\alpha}^{\pi/2} = \tfrac{11}{4}\pi - \tfrac{11}{2}\sin^{-1}\left(\tfrac{1}{3}\right) - 3\sqrt{2}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. The curves intersect when $4\cos\theta = 2 \;\Rightarrow\; \cos\theta = \frac{1}{2} \;\Rightarrow\; \theta = \pm\frac{\pi}{3}$ for $-\pi \le \theta \le \pi$. The points of intersection are $\left(2, \frac{\pi}{3}\right)$ and $\left(2, -\frac{\pi}{3}\right)$.



34. The two curves clearly both contain the pole. For other points of intersection, $\cot\theta = 2\cos(\theta + 2n\pi)$ or $-2\cos(\theta + \pi + 2n\pi)$, both of which reduce to $\cot\theta = 2\cos\theta \;\Leftrightarrow\; \cos\theta = 2\sin\theta\cos\theta \;\Leftrightarrow\; \cos\theta(1 - 2\sin\theta) = 0 \;\Rightarrow$ $\cos\theta = 0$ or $\sin\theta = \frac{1}{2} \;\Rightarrow\; \theta = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}$ or $\frac{3\pi}{2} \;\Rightarrow$ intersection points are $\left(0, \frac{\pi}{2}\right)$, $\left(\sqrt{3}, \frac{\pi}{6}\right)$, and $\left(\sqrt{3}, \frac{11\pi}{6}\right)$.

35. The curves intersect where $2\sin\theta = \sin\theta + \cos\theta \;\Rightarrow$ $\sin\theta = \cos\theta \;\Rightarrow \theta = \frac{\pi}{4}$, and also at the origin (at which $\theta = \frac{3\pi}{4}$ on the second curve).



$$\begin{aligned} A &= \int_0^{\pi/4} \tfrac{1}{2}(2\sin\theta)^2\,d\theta + \int_{\pi/4}^{3\pi/4} \tfrac{1}{2}(\sin\theta + \cos\theta)^2\,d\theta \\ &= \int_0^{\pi/4} (1 - \cos 2\theta)\,d\theta + \tfrac{1}{2}\int_{\pi/4}^{3\pi/4} (1 + \sin 2\theta)\,d\theta \\ &= \left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/4} + \left[\tfrac{1}{2}\theta - \tfrac{1}{4}\cos 2\theta\right]_{\pi/4}^{3\pi/4} = \tfrac{1}{2}(\pi - 1) \end{aligned}$$

36.



$$\begin{aligned} A &= 2\int_{-\pi/2}^{\pi/6} \tfrac{1}{2}\left[(2 + \cos 2\theta)^2 - (2 + \sin\theta)^2\right] d\theta \\ &= \int_{-\pi/2}^{\pi/6} \left[4\cos 2\theta + \cos^2 2\theta - 4\sin\theta - \sin^2\theta\right] d\theta \\ &= \left[2\sin 2\theta + \tfrac{1}{2}\theta + \tfrac{1}{8}\sin 4\theta + 4\cos\theta - \tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta\right]_{-\pi/2}^{\pi/6} \\ &= \tfrac{51}{16}\sqrt{3} \end{aligned}$$

37. $x = 3t^2,\ y = 2t^3$.

$$\begin{aligned} L &= \int_0^2 \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^2 \sqrt{(6t)^2 + (6t^2)^2}\,dt = \int_0^2 \sqrt{36t^2 + 36t^4}\,dt = \int_0^2 \sqrt{36t^2}\,\sqrt{1 + t^2}\,dt \\ &= \int_0^2 6\,|t|\,\sqrt{1 + t^2}\,dt = 6\int_0^2 t\,\sqrt{1 + t^2}\,dt = 6\int_1^5 u^{1/2}\left(\tfrac{1}{2}du\right) \qquad \left[u = 1 + t^2,\ du = 2t\,dt\right] \\ &= 6 \cdot \tfrac{1}{2} \cdot \tfrac{2}{3}\left[u^{3/2}\right]_1^5 = 2(5^{3/2} - 1) = 2\left(5\sqrt{5} - 1\right) \end{aligned}$$

38. $x = 2 + 3t,\quad y = \cosh 3t \;\Rightarrow\; (dx/dt)^2 + (dy/dt)^2 = 3^2 + (3\sinh 3t)^2 = 9(1 + \sinh^2 3t) = 9\cosh^2 3t$, so

$$L = \int_0^1 \sqrt{9\cosh^2 3t}\,dt = \int_0^1 |3\cosh 3t|\,dt = \int_0^1 3\cosh 3t\,dt = \left[\sinh 3t\right]_0^1 = \sinh 3 - \sinh 0 = \sinh 3.$$

39.

$$\begin{aligned} L &= \int_\pi^{2\pi} \sqrt{r^2 + (dr/d\theta)^2}\,d\theta = \int_\pi^{2\pi} \sqrt{(1/\theta)^2 + (-1/\theta^2)^2}\,d\theta = \int_\pi^{2\pi} \frac{\sqrt{\theta^2 + 1}}{\theta^2}\,d\theta \\ &\overset{24}{=} \left[-\frac{\sqrt{\theta^2 + 1}}{\theta} + \ln\left(\theta + \sqrt{\theta^2 + 1}\right)\right]_\pi^{2\pi} = \frac{\sqrt{\pi^2 + 1}}{\pi} - \frac{\sqrt{4\pi^2 + 1}}{2\pi} + \ln\left(\frac{2\pi + \sqrt{4\pi^2 + 1}}{\pi + \sqrt{\pi^2 + 1}}\right) \\ &= \frac{2\sqrt{\pi^2 + 1} - \sqrt{4\pi^2 + 1}}{2\pi} + \ln\left(\frac{2\pi + \sqrt{4\pi^2 + 1}}{\pi + \sqrt{\pi^2 + 1}}\right) \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $L = \int_0^{\pi} \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_0^{\pi} \sqrt{\sin^6\left(\frac{1}{3}\theta\right) + \sin^4\left(\frac{1}{3}\theta\right)\cos^2\left(\frac{1}{3}\theta\right)}\, d\theta$

$= \int_0^{\pi} \sin^2\left(\frac{1}{3}\theta\right) d\theta = \left[\frac{1}{2}\left(\theta - \frac{3}{2}\sin\left(\frac{2}{3}\theta\right)\right)\right]_0^{\pi} = \frac{1}{2}\pi - \frac{3}{8}\sqrt{3}$

41. $x = 4\sqrt{t},\ y = \dfrac{t^3}{3} + \dfrac{1}{2t^2},\ 1 \le t \le 4 \quad \Rightarrow$

$S = \int_1^4 2\pi y \sqrt{(dx/dt)^2 + (dy/dt)^2}\, dt = \int_1^4 2\pi\left(\frac{1}{3}t^3 + \frac{1}{2}t^{-2}\right)\sqrt{\left(2/\sqrt{t}\right)^2 + (t^2 - t^{-3})^2}\, dt$

$= 2\pi \int_1^4 \left(\frac{1}{3}t^3 + \frac{1}{2}t^{-2}\right)\sqrt{(t^2 + t^{-3})^2}\, dt = 2\pi \int_1^4 \left(\frac{1}{3}t^5 + \frac{5}{6} + \frac{1}{2}t^{-5}\right) dt = 2\pi\left[\frac{1}{18}t^6 + \frac{5}{6}t - \frac{1}{8}t^{-4}\right]_1^4 = \frac{471{,}295}{1024}\pi$

42. $x = 2 + 3t, \quad y = \cosh 3t \quad \Rightarrow \quad (dx/dt)^2 + (dy/dt)^2 = 3^2 + (3\sinh 3t)^2 = 9(1 + \sinh^2 3t) = 9\cosh^2 3t$, so

$S = \int_0^1 2\pi y\, ds = \int_0^1 2\pi \cosh 3t \sqrt{9\cosh^2 3t}\, dt = \int_0^1 2\pi \cosh 3t\, |3\cosh 3t|\, dt = \int_0^1 2\pi \cosh 3t \cdot 3\cosh 3t\, dt$

$= 6\pi \int_0^1 \cosh^2 3t\, dt = 6\pi \int_0^1 \frac{1}{2}(1 + \cosh 6t)\, dt = 3\pi\left[t + \frac{1}{6}\sinh 6t\right]_0^1 = 3\pi\left(1 + \frac{1}{6}\sinh 6\right) = 3\pi + \frac{\pi}{2}\sinh 6$

43. For all c except -1, the curve is asymptotic to the line $x = 1$. For $c < -1$, the curve bulges to the right near $y = 0$. As c increases, the bulge becomes smaller, until at $c = -1$ the curve is the straight line $x = 1$. As c continues to increase, the curve bulges to the left, until at $c = 0$ there is a cusp at the origin. For $c > 0$, there is a loop to the left of the origin, whose size and roundness increase as c increases. Note that the x-intercept of the curve is always $-c$.



44. For a close to 0, the graph of $r^n = |\sin 2\theta|$ consists of four thin petals. As a increases, the petals get wider, until as $a \to \infty$, each petal occupies almost its entire quarter-circle.


$a = 0.01$


$a = 0.1$


$a = 1$


$a = 5$


$a = 10$


$a = 25$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. $\dfrac{x^2}{9}+\dfrac{y^2}{8}=1$ is an ellipse with center $(0,0)$.

$a=3,\ b=2\sqrt{2},\ c=1 \ \Rightarrow$

foci $(\pm1,0)$, vertices $(\pm3,0)$.



46. $4x^2-y^2=16 \ \Leftrightarrow \ \dfrac{x^2}{4}-\dfrac{y^2}{16}=1$ is a hyperbola with center $(0,0)$, vertices $(\pm2,0)$, $a=2$, $b=4$, $c=\sqrt{16+4}=2\sqrt{5}$, foci $\left(\pm2\sqrt{5},0\right)$ and asymptotes $y=\pm2x$.



47. $6y^2+x-36y+55=0 \ \Leftrightarrow$

$6(y^2-6y+9)=-(x+1) \ \Leftrightarrow$

$(y-3)^2=-\frac{1}{6}(x+1)$, a parabola with vertex $(-1,3)$, opening to the left, $p=-\frac{1}{24} \ \Rightarrow$ focus $\left(-\frac{25}{24},3\right)$ and directrix $x=-\frac{23}{24}$.



48. $25x^2+4y^2+50x-16y=59 \ \Leftrightarrow$

$25(x+1)^2+4(y-2)^2=100 \ \Leftrightarrow$

$\frac{1}{4}(x+1)^2+\frac{1}{25}(y-2)^2=1$ is an ellipse centered at $(-1,2)$ with foci on the line $x=-1$, vertices $(-1,7)$ and $(-1,-3)$; $a=5,\ b=2 \ \Rightarrow \ c=\sqrt{21} \ \Rightarrow$ foci $\left(-1,2\pm\sqrt{21}\right)$.



49. The ellipse with foci $(\pm4,0)$ and vertices $(\pm5,0)$ has center $(0,0)$ and a horizontal major axis, with $a=5$ and $c=4$, so $b^2=a^2-c^2=5^2-4^2=9$. An equation is $\dfrac{x^2}{25}+\dfrac{y^2}{9}=1$.

50. The distance from the focus $(2,1)$ to the directrix $x=-4$ is $2-(-4)=6$, so the distance from the focus to the vertex is $\frac{1}{2}(6)=3$ and the vertex is $(-1,1)$. Since the focus is to the right of the vertex, $p=3$. An equation is $(y-1)^2=4\cdot3[x-(-1)]$, or $(y-1)^2=12(x+1)$.

51. The center of a hyperbola with foci $(0,\pm4)$ is $(0,0)$, so $c=4$ and an equation is $\dfrac{y^2}{a^2}-\dfrac{x^2}{b^2}=1$.

The asymptote $y=3x$ has slope 3, so $\dfrac{a}{b}=\dfrac{3}{1} \ \Rightarrow \ a=3b$ and $a^2+b^2=c^2 \ \Rightarrow \ (3b)^2+b^2=4^2 \ \Rightarrow$

$10b^2=16 \ \Rightarrow \ b^2=\frac{8}{5}$ and so $a^2=16-\frac{8}{5}=\frac{72}{5}$. Thus, an equation is $\dfrac{y^2}{72/5}-\dfrac{x^2}{8/5}=1$, or $\dfrac{5y^2}{72}-\dfrac{5x^2}{8}=1$.

52. Center is $(3,0)$, and $a=\frac{8}{2}=4,\ c=2 \ \Leftrightarrow \ b=\sqrt{4^2-2^2}=\sqrt{12} \ \Rightarrow$

an equation of the ellipse is $\dfrac{(x-3)^2}{12}+\dfrac{y^2}{16}=1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

53. $x^2 + y = 100 \;\Leftrightarrow\; x^2 = -(y - 100)$ has its vertex at $(0, 100)$, so one of the vertices of the ellipse is $(0, 100)$. Another form of the equation of a parabola is $x^2 = 4p(y - 100)$ so $4p(y - 100) = -(y - 100) \;\Rightarrow\; 4p = -1 \;\Rightarrow\; p = -\frac{1}{4}$. Therefore the shared focus is found at $\left(0, \frac{399}{4}\right)$ so $2c = \frac{399}{4} - 0 \;\Rightarrow\; c = \frac{399}{8}$ and the center of the ellipse is $\left(0, \frac{399}{8}\right)$. So $a = 100 - \frac{399}{8} = \frac{401}{8}$ and $b^2 = a^2 - c^2 = \dfrac{401^2 - 399^2}{8^2} = 25$. So the equation of the ellipse is $\dfrac{x^2}{b^2} + \dfrac{\left(y - \frac{399}{8}\right)^2}{a^2} = 1 \;\Rightarrow$ $\dfrac{x^2}{25} + \dfrac{\left(y - \frac{399}{8}\right)^2}{\left(\frac{401}{8}\right)^2} = 1$, or $\dfrac{x^2}{25} + \dfrac{(8y - 399)^2}{160{,}801} = 1$.

54. $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \;\Rightarrow\; \dfrac{2x}{a^2} + \dfrac{2y}{b^2}\dfrac{dy}{dx} = 0 \;\Rightarrow\; \dfrac{dy}{dx} = -\dfrac{b^2}{a^2}\dfrac{x}{y}$. Therefore $\dfrac{dy}{dx} = m \;\Leftrightarrow\; y = -\dfrac{b^2}{a^2}\dfrac{x}{m}$. Combining this condition with $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$, we find that $x = \pm\dfrac{a^2 m}{\sqrt{a^2m^2 + b^2}}$. In other words, the two points on the ellipse where the tangent has slope m are $\left(\pm\dfrac{a^2 m}{\sqrt{a^2m^2 + b^2}}, \mp\dfrac{b^2}{\sqrt{a^2m^2 + b^2}}\right)$. The tangent lines at these points have the equations $y \pm \dfrac{b^2}{\sqrt{a^2m^2 + b^2}} = m\left(x \mp \dfrac{a^2 m}{\sqrt{a^2m^2 + b^2}}\right)$ or $y = mx \mp \dfrac{a^2m^2}{\sqrt{a^2m^2 + b^2}} \mp \dfrac{b^2}{\sqrt{a^2m^2 + b^2}} = mx \mp \sqrt{a^2m^2 + b^2}$.

55. Directrix $x = 4 \;\Rightarrow\; d = 4$, so $e = \frac{1}{3} \;\Rightarrow\; r = \dfrac{ed}{1 + e\cos\theta} = \dfrac{4}{3 + \cos\theta}$.

56. See the end of the proof of Theorem 10.6.1. If $e > 1$, then $1 - e^2 < 0$ and Equations 10.6.4 become $a^2 = \dfrac{e^2d^2}{(e^2 - 1)^2}$ and $b^2 = \dfrac{e^2d^2}{e^2 - 1}$, so $\dfrac{b^2}{a^2} = e^2 - 1$. The asymptotes $y = \pm\dfrac{b}{a}x$ have slopes $\pm\dfrac{b}{a} = \pm\sqrt{e^2 - 1}$, so the angles they make with the polar axis are $\pm\tan^{-1}\left[\sqrt{e^2 - 1}\right] = \cos^{-1}(\pm 1/e)$.

57. In polar coordinates, an equation for the circle is $r = 2a\sin\theta$. Thus, the coordinates of Q are $x = r\cos\theta = 2a\sin\theta\,\cos\theta$ and $y = r\sin\theta = 2a\sin^2\theta$. The coordinates of R are $x = 2a\cot\theta$ and $y = 2a$. Since P is the midpoint of QR, we use the midpoint formula to get $x = a(\sin\theta\,\cos\theta + \cot\theta)$ and $y = a(1 + \sin^2\theta)$.

58. (a) If (a, b) lies on the curve, then there is some parameter value t_1 such that $\dfrac{3t_1}{1 + t_1^3} = a$ and $\dfrac{3t_1^2}{1 + t_1^3} = b$. If $t_1 = 0$, the point is $(0, 0)$, which lies on the line $y = x$. If $t_1 \neq 0$, then the point corresponding to $t = \dfrac{1}{t_1}$ is given by $x = \dfrac{3(1/t_1)}{1 + (1/t_1)^3} = \dfrac{3t_1^2}{t_1^3 + 1} = b$, $y = \dfrac{3(1/t_1)^2}{1 + (1/t_1)^3} = \dfrac{3t_1}{t_1^3 + 1} = a$. So (b, a) also lies on the curve. [Another way to see this is to do part (e) first; the result is immediate.] The curve intersects the line $y = x$ when $\dfrac{3t}{1 + t^3} = \dfrac{3t^2}{1 + t^3} \;\Rightarrow$ $t = t^2 \;\Rightarrow\; t = 0$ or 1, so the points are $(0, 0)$ and $\left(\frac{3}{2}, \frac{3}{2}\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\dfrac{dy}{dt} = \dfrac{(1+t^3)(6t) - 3t^2(3t^2)}{(1+t^3)^2} = \dfrac{6t - 3t^4}{(1+t^3)^2} = 0$ when $6t - 3t^4 = 3t(2 - t^3) = 0 \quad \Rightarrow \quad t = 0$ or $t = \sqrt[3]{2}$, so there are horizontal tangents at $(0, 0)$ and $\left(\sqrt[3]{2}, \sqrt[3]{4}\right)$. Using the symmetry from part (a), we see that there are vertical tangents at $(0, 0)$ and $\left(\sqrt[3]{4}, \sqrt[3]{2}\right)$.

(c) Notice that as $t \to -1^+$, we have $x \to -\infty$ and $y \to \infty$. As $t \to -1^-$, we have $x \to \infty$ and $y \to -\infty$. Also

$y - (-x - 1) = y + x + 1 = \dfrac{3t + 3t^2 + (1+t^3)}{1+t^3} = \dfrac{(t+1)^3}{1+t^3} = \dfrac{(t+1)^2}{t^2 - t + 1} \to 0$ as $t \to -1$. So $y = -x - 1$ is a slant asymptote.

(d) $\dfrac{dx}{dt} = \dfrac{(1+t^3)(3) - 3t(3t^2)}{(1+t^3)^2} = \dfrac{3 - 6t^3}{(1+t^3)^2}$ and from part (b) we have $\dfrac{dy}{dt} = \dfrac{6t - 3t^4}{(1+t^3)^2}$. So $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{t(2 - t^3)}{1 - 2t^3}$.

Also $\dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{2(1+t^3)^4}{3(1-2t^3)^3} > 0 \quad \Leftrightarrow \quad t < \dfrac{1}{\sqrt[3]{2}}$.

So the curve is concave upward there and has a minimum point at $(0, 0)$ and a maximum point at $\left(\sqrt[3]{2}, \sqrt[3]{4}\right)$. Using this together with the information from parts (a), (b), and (c), we sketch the curve.



(e) $x^3 + y^3 = \left(\dfrac{3t}{1+t^3}\right)^3 + \left(\dfrac{3t^2}{1+t^3}\right)^3 = \dfrac{27t^3 + 27t^6}{(1+t^3)^3} = \dfrac{27t^3(1+t^3)}{(1+t^3)^3} = \dfrac{27t^3}{(1+t^3)^2}$

and $3xy = 3\left(\dfrac{3t}{1+t^3}\right)\left(\dfrac{3t^2}{1+t^3}\right) = \dfrac{27t^3}{(1+t^3)^2}$, so $x^3 + y^3 = 3xy$.

(f) We start with the equation from part (e) and substitute $x = r\cos\theta$, $y = r\sin\theta$. Then $x^3 + y^3 = 3xy \quad \Rightarrow$ $r^3\cos^3\theta + r^3\sin^3\theta = 3r^2\cos\theta\,\sin\theta$. For $r \neq 0$, this gives $r = \dfrac{3\cos\theta\,\sin\theta}{\cos^3\theta + \sin^3\theta}$. Dividing numerator and denominator by $\cos^3\theta$, we obtain $r = \dfrac{3\left(\dfrac{1}{\cos\theta}\right)\dfrac{\sin\theta}{\cos\theta}}{1 + \dfrac{\sin^3\theta}{\cos^3\theta}} = \dfrac{3\sec\theta\,\tan\theta}{1 + \tan^3\theta}$.

(g) The loop corresponds to $\theta \in \left(0, \frac{\pi}{2}\right)$, so its area is

$$A = \int_0^{\pi/2} \frac{r^2}{2}\,d\theta = \frac{1}{2}\int_0^{\pi/2} \left(\frac{3\sec\theta\,\tan\theta}{1+\tan^3\theta}\right)^2 d\theta = \frac{9}{2}\int_0^{\pi/2} \frac{\sec^2\theta\,\tan^2\theta}{(1+\tan^3\theta)^2}\,d\theta = \frac{9}{2}\int_0^{\infty} \frac{u^2\,du}{(1+u^3)^2} \quad [\text{let } u = \tan\theta]$$

$$= \lim_{b\to\infty} \tfrac{9}{2}\left[-\tfrac{1}{3}(1+u^3)^{-1}\right]_0^b = \tfrac{3}{2}$$

(h) By symmetry, the area between the folium and the line $y = -x - 1$ is equal to the enclosed area in the third quadrant, plus twice the enclosed area in the fourth quadrant. The area in the third quadrant is $\frac{1}{2}$, and since $y = -x - 1 \quad \Rightarrow$ $r\sin\theta = -r\cos\theta - 1 \quad \Rightarrow \quad r = -\dfrac{1}{\sin\theta + \cos\theta}$, the area in the fourth quadrant is

$$\frac{1}{2}\int_{-\pi/2}^{-\pi/4} \left[\left(-\frac{1}{\sin\theta + \cos\theta}\right)^2 - \left(\frac{3\sec\theta\,\tan\theta}{1+\tan^3\theta}\right)^2\right] d\theta \overset{\text{CAS}}{=} \frac{1}{2}.$$

Therefore, the total area is $\frac{1}{2} + 2\left(\frac{1}{2}\right) = \frac{3}{2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

PROBLEMS PLUS

1. See the figure. The circle with center $(-1, 0)$ and radius $\sqrt{2}$ has equation $(x+1)^2 + y^2 = 2$ and describes the circular arc from $(0, -1)$ to $(0, 1)$. Converting the equation to polar coordinates gives us



$(r\cos\theta + 1)^2 + (r\sin\theta)^2 = 2 \quad \Rightarrow$

$r^2\cos^2\theta + 2r\cos\theta + 1 + r^2\sin^2\theta = 2 \quad \Rightarrow$

$r^2(\cos^2\theta + \sin^2\theta) + 2r\cos\theta = 1 \quad \Rightarrow \quad r^2 + 2r\cos\theta = 1$. Using the quadratic formula to solve for r gives us

$$r = \frac{-2\cos\theta \pm \sqrt{4\cos^2\theta + 4}}{2} = -\cos\theta + \sqrt{\cos^2\theta + 1} \text{ for } r > 0.$$

The darkest shaded region is $\frac{1}{8}$ of the entire shaded region A, so $\frac{1}{8}A = \int_0^{\pi/4} \frac{1}{2}r^2\,d\theta = \frac{1}{2}\int_0^{\pi/4} (1 - 2r\cos\theta)\,d\theta \quad \Rightarrow$

$$\begin{aligned}
\tfrac{1}{4}A &= \int_0^{\pi/4} \left[1 - 2\cos\theta\left(-\cos\theta + \sqrt{\cos^2\theta + 1}\right)\right] d\theta = \int_0^{\pi/4} \left(1 + 2\cos^2\theta - 2\cos\theta\sqrt{\cos^2\theta + 1}\right) d\theta \\
&= \int_0^{\pi/4} \left[1 + 2\cdot\tfrac{1}{2}(1 + \cos 2\theta) - 2\cos\theta\sqrt{(1 - \sin^2\theta) + 1}\right] d\theta \\
&= \int_0^{\pi/4} (2 + \cos 2\theta)\,d\theta - 2\int_0^{\pi/4} \cos\theta\sqrt{2 - \sin^2\theta}\,d\theta \\
&= \left[2\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/4} - 2\int_0^{1/\sqrt{2}} \sqrt{2 - u^2}\,du \qquad \left[\begin{matrix} u = \sin\theta, \\ du = \cos\theta\,d\theta \end{matrix}\right] \\
&= \left(\frac{\pi}{2} + \frac{1}{2}\right) - (0 + 0) - 2\left[\frac{u}{2}\sqrt{2 - u^2} + \sin^{-1}\frac{u}{\sqrt{2}}\right]_0^{1/\sqrt{2}} \qquad \left[\begin{matrix} \text{Formula 30,} \\ a = \sqrt{2} \end{matrix}\right] \\
&= \frac{\pi}{2} + \frac{1}{2} - 2\left(\frac{1}{2\sqrt{2}}\cdot\frac{\sqrt{3}}{\sqrt{2}} + \frac{\pi}{6}\right) = \frac{\pi}{2} + \frac{1}{2} - \frac{1}{2}\sqrt{3} - \frac{\pi}{3} = \frac{\pi}{6} + \frac{1}{2} - \frac{1}{2}\sqrt{3}.
\end{aligned}$$

Thus, $A = 4\left(\frac{\pi}{6} + \frac{1}{2} - \frac{1}{2}\sqrt{3}\right) = \frac{2\pi}{3} + 2 - 2\sqrt{3}$.

2. (a) The curve $x^4 + y^4 = x^2 + y^2$ is symmetric about both axes and about the line $y = x$ (since interchanging x and y does not change the equation) so we need only consider $y \geq x \geq 0$ to begin with. Implicit differentiation gives

$4x^3 + 4y^3y' = 2x + 2yy' \quad \Rightarrow \quad y' = \dfrac{x(1 - 2x^2)}{y(2y^2 - 1)} \quad \Rightarrow \quad y' = 0$ when $x = 0$ and when $x = \pm\frac{1}{\sqrt{2}}$. If $x = 0$, then

$y^4 = y^2 \quad \Rightarrow \quad y^2(y^2 - 1) = 0 \quad \Rightarrow \quad y = 0$ or ± 1. The point $(0, 0)$ can't be a highest or lowest point because it is isolated. [If $-1 < x < 1$ and $-1 < y < 1$, then $x^4 < x^2$ and $y^4 < y^2 \quad \Rightarrow \quad x^4 + y^4 < x^2 + y^2$, except for $(0, 0)$.]

If $x = \frac{1}{\sqrt{2}}$, then $x^2 = \frac{1}{2}$, $x^4 = \frac{1}{4}$, so $\frac{1}{4} + y^4 = \frac{1}{2} + y^2 \quad \Rightarrow \quad 4y^4 - 4y^2 - 1 = 0 \quad \Rightarrow \quad y^2 = \frac{4 \pm \sqrt{16+16}}{8} = \frac{1 \pm \sqrt{2}}{2}$.

But $y^2 > 0$, so $y^2 = \frac{1+\sqrt{2}}{2} \quad \Rightarrow \quad y = \pm\sqrt{\frac{1}{2}(1 + \sqrt{2})}$. Near the point $(0, 1)$, the denominator of y' is positive and the numerator changes from negative to positive as x increases through 0, so $(0, 1)$ is a local minimum point. At

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$\left(\frac{1}{\sqrt{2}}, \sqrt{\frac{1+\sqrt{2}}{2}}\right)$, y' changes from positive to negative, so that point gives a maximum. By symmetry, the highest points on the curve are $\left(\pm\frac{1}{\sqrt{2}}, \sqrt{\frac{1+\sqrt{2}}{2}}\right)$ and the lowest points are $\left(\pm\frac{1}{\sqrt{2}}, -\sqrt{\frac{1+\sqrt{2}}{2}}\right)$.

(b) We use the information from part (a), together with symmetry with respect to the axes and the lines $y = \pm x$, to sketch the curve.



(c) In polar coordinates, $x^4 + y^4 = x^2 + y^2$ becomes $r^4\cos^4\theta + r^4\sin^4\theta = r^2$ or $r^2 = \dfrac{1}{\cos^4\theta + \sin^4\theta}$. By the symmetry shown in part (b), the area enclosed by the curve is $A = 8\displaystyle\int_0^{\pi/4} \frac{1}{2}r^2\,d\theta = 4\int_0^{\pi/4} \frac{d\theta}{\cos^4\theta + \sin^4\theta} \overset{\text{CAS}}{=} \sqrt{2}\,\pi$.

3. In terms of x and y, we have $x = r\cos\theta = (1 + c\sin\theta)\cos\theta = \cos\theta + c\sin\theta\cos\theta = \cos\theta + \frac{1}{2}c\sin 2\theta$ and $y = r\sin\theta = (1 + c\sin\theta)\sin\theta = \sin\theta + c\sin^2\theta$. Now $-1 \le \sin\theta \le 1 \;\Rightarrow\; -1 \le \sin\theta + c\sin^2\theta \le 1 + c \le 2$, so $-1 \le y \le 2$. Furthermore, $y = 2$ when $c = 1$ and $\theta = \frac{\pi}{2}$, while $y = -1$ for $c = 0$ and $\theta = \frac{3\pi}{2}$. Therefore, we need a viewing rectangle with $-1 \le y \le 2$.

To find the x-values, look at the equation $x = \cos\theta + \frac{1}{2}c\sin 2\theta$ and use the fact that $\sin 2\theta \ge 0$ for $0 \le \theta \le \frac{\pi}{2}$ and $\sin 2\theta \le 0$ for $-\frac{\pi}{2} \le \theta \le 0$. [Because $r = 1 + c\sin\theta$ is symmetric about the y-axis, we only need to consider $-\frac{\pi}{2} \le \theta \le \frac{\pi}{2}$.] So for $-\frac{\pi}{2} \le \theta \le 0$, x has a maximum value when $c = 0$ and then $x = \cos\theta$ has a maximum value of 1 at $\theta = 0$. Thus, the maximum value of x must occur on $\left[0, \frac{\pi}{2}\right]$ with $c = 1$. Then $x = \cos\theta + \frac{1}{2}\sin 2\theta \;\Rightarrow$ $\frac{dx}{d\theta} = -\sin\theta + \cos 2\theta = -\sin\theta + 1 - 2\sin^2\theta \;\Rightarrow\; \frac{dx}{d\theta} = -(2\sin\theta - 1)(\sin\theta + 1) = 0$ when $\sin\theta = -1$ or $\frac{1}{2}$ [but $\sin\theta \ne -1$ for $0 \le \theta \le \frac{\pi}{2}$]. If $\sin\theta = \frac{1}{2}$, then $\theta = \frac{\pi}{6}$ and $x = \cos\frac{\pi}{6} + \frac{1}{2}\sin\frac{\pi}{3} = \frac{3}{4}\sqrt{3}$. Thus, the maximum value of x is $\frac{3}{4}\sqrt{3}$, and, by symmetry, the minimum value is $-\frac{3}{4}\sqrt{3}$. Therefore, the smallest viewing rectangle that contains every member of the family of polar curves $r = 1 + c\sin\theta$, where $0 \le c \le 1$, is $\left[-\frac{3}{4}\sqrt{3}, \frac{3}{4}\sqrt{3}\right] \times [-1, 2]$.



4. (a) Let us find the polar equation of the path of the bug that starts in the upper right corner of the square. If the polar coordinates of this bug, at a particular moment, are (r, θ), then the polar coordinates of the bug that it is crawling toward must be $\left(r, \theta + \frac{\pi}{2}\right)$. (The next bug must be the same distance from the origin and the angle between the lines joining the bugs to the pole must be $\frac{\pi}{2}$.) The Cartesian coordinates of the first bug are $(r\cos\theta, r\sin\theta)$ and for the second bug we have $x = r\cos\left(\theta + \frac{\pi}{2}\right) = -r\sin\theta$, $y = r\sin\left(\theta + \frac{\pi}{2}\right) = r\cos\theta$. So the slope of the line joining the bugs is



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\dfrac{r\cos\theta - r\sin\theta}{-r\sin\theta - r\cos\theta} = \dfrac{\sin\theta - \cos\theta}{\sin\theta + \cos\theta}$. This must be equal to the slope of the tangent line at (r, θ), so by Equation 10.3.3 we have $\dfrac{(dr/d\theta)\sin\theta + r\cos\theta}{(dr/d\theta)\cos\theta - r\sin\theta} = \dfrac{\sin\theta - \cos\theta}{\sin\theta + \cos\theta}$. Solving for $\dfrac{dr}{d\theta}$, we get

$\dfrac{dr}{d\theta}\sin^2\theta + \dfrac{dr}{d\theta}\sin\theta\cos\theta + r\sin\theta\cos\theta + r\cos^2\theta = \dfrac{dr}{d\theta}\sin\theta\cos\theta - \dfrac{dr}{d\theta}\cos^2\theta - r\sin^2\theta + r\sin\theta\cos\theta \Rightarrow$

$\dfrac{dr}{d\theta}\left(\sin^2\theta + \cos^2\theta\right) + r\left(\cos^2\theta + \sin^2\theta\right) = 0 \;\Rightarrow\; \dfrac{dr}{d\theta} = -r$. Solving this differential equation as a separable equation (as in Section 9.3), or using Theorem 9.4.2 with $k = -1$, we get $r = Ce^{-\theta}$. To determine C we use the fact that, at its starting position, $\theta = \frac{\pi}{4}$ and $r = \frac{1}{\sqrt{2}}a$, so $\frac{1}{\sqrt{2}}a = Ce^{-\pi/4} \;\Rightarrow\; C = \frac{1}{\sqrt{2}}ae^{\pi/4}$. Therefore, a polar equation of the bug's path is $r = \frac{1}{\sqrt{2}}ae^{\pi/4}e^{-\theta}$ or $r = \frac{1}{\sqrt{2}}ae^{(\pi/4)-\theta}$.

(b) The distance traveled by this bug is $L = \int_{\pi/4}^{\infty}\sqrt{r^2 + (dr/d\theta)^2}\,d\theta$, where $\dfrac{dr}{d\theta} = \dfrac{a}{\sqrt{2}}\,e^{\pi/4}(-e^{-\theta})$ and so $r^2 + (dr/d\theta)^2 = \frac{1}{2}a^2e^{\pi/2}e^{-2\theta} + \frac{1}{2}a^2e^{\pi/2}e^{-2\theta} = a^2e^{\pi/2}e^{-2\theta}$. Thus

$$L = \int_{\pi/4}^{\infty} ae^{\pi/4}e^{-\theta}\,d\theta = ae^{\pi/4}\lim_{t\to\infty}\int_{\pi/4}^{t} e^{-\theta}\,d\theta = ae^{\pi/4}\lim_{t\to\infty}\left[-e^{-\theta}\right]_{\pi/4}^{t}$$
$$= ae^{\pi/4}\lim_{t\to\infty}\left[e^{-\pi/4} - e^{-t}\right] = ae^{\pi/4}e^{-\pi/4} = a$$

5. Without loss of generality, assume the hyperbola has equation $\dfrac{x^2}{a^2} - \dfrac{y^2}{b^2} = 1$. Use implicit differentiation to get $\dfrac{2x}{a^2} - \dfrac{2y\,y'}{b^2} = 0$, so $y' = \dfrac{b^2x}{a^2y}$. The tangent line at the point (c, d) on the hyperbola has equation $y - d = \dfrac{b^2c}{a^2d}(x - c)$. The tangent line intersects the asymptote $y = \dfrac{b}{a}x$ when $\dfrac{b}{a}x - d = \dfrac{b^2c}{a^2d}(x - c) \;\Rightarrow\; abdx - a^2d^2 = b^2cx - b^2c^2 \;\Rightarrow$ $abdx - b^2cx = a^2d^2 - b^2c^2 \;\Rightarrow\; x = \dfrac{a^2d^2 - b^2c^2}{b(ad - bc)} = \dfrac{ad + bc}{b}$ and the y-value is $\dfrac{b}{a}\dfrac{ad + bc}{b} = \dfrac{ad + bc}{a}$.

Similarly, the tangent line intersects $y = -\dfrac{b}{a}x$ at $\left(\dfrac{bc - ad}{b}, \dfrac{ad - bc}{a}\right)$. The midpoint of these intersection points is

$\left(\dfrac{1}{2}\left(\dfrac{ad + bc}{b} + \dfrac{bc - ad}{b}\right), \dfrac{1}{2}\left(\dfrac{ad + bc}{a} + \dfrac{ad - bc}{a}\right)\right) = \left(\dfrac{1}{2}\dfrac{2bc}{b}, \dfrac{1}{2}\dfrac{2ad}{a}\right) = (c, d)$, the point of tangency.

Note: If $y = 0$, then at $(\pm a, 0)$, the tangent line is $x = \pm a$, and the points of intersection are clearly equidistant from the point of tangency.

6. (a) Since the smaller circle rolls without slipping around C, the amount of arc traversed on C ($2r\theta$ in the figure) must equal the amount of arc of the smaller circle that has been in contact with C. Since the smaller circle has radius r, it must have turned through an angle of $2r\theta/r = 2\theta$. In addition to turning through an angle 2θ, the little circle has rolled through an angle θ against C. Thus, P has turned through an angle of 3θ as shown in the figure. (If the little circle had turned through an angle of 2θ with its center pinned to the x-axis,



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

then P would have turned only 2θ instead of 3θ. The movement of the little circle around C adds θ to the angle.) From the figure, we see that the center of the small circle has coordinates $(3r\cos\theta, 3r\sin\theta)$. Thus, P has coordinates (x, y), where $x = b\cos 3\theta + 3r\cos\theta$ and $y = b\sin 3\theta + 3r\sin\theta$.

(b)



$b = \frac{1}{5}r$ $\quad$ $b = \frac{2}{5}r$ $\quad$ $b = \frac{3}{5}r$ $\quad$ $b = \frac{4}{5}r$

(c) The diagram gives an alternate description of point P on the epitrochoid. Q moves around a circle of radius b, and P rotates one-third as fast with respect to Q at a distance of $3r$. Place an equilateral triangle with sides of length $3\sqrt{3}r$ so that its centroid is at Q and one vertex is at P. (The distance from the centroid to a vertex is $\frac{1}{\sqrt{3}}$ times the length of a side of the equilateral triangle.)



As θ increases by $\frac{2\pi}{3}$, the point Q travels once around the circle of radius b, returning to its original position. At the same time, P (and the rest of the triangle) rotate through an angle of $\frac{2\pi}{3}$ about Q, so P's position is occupied by another vertex. In this way, we see that the epitrochoid traced out by P is simultaneously traced out by the other two vertices as well. The whole equilateral triangle sits inside the epitrochoid (touching it only with its vertices) and each vertex traces out the curve once while the centroid moves around the circle three times.

(d) We view the epitrochoid as being traced out in the same way as in part (c), by a rotor for which the distance from its center to each vertex is $3r$, so it has radius $6r$. To show that the rotor fits inside the epitrochoid, it suffices to show that for any position of the tracing point P, there are no points on the opposite side of the rotor which are outside the epitrochoid. But the most likely case of intersection is when P is on the y-axis, so as long as the diameter of the rotor (which is $3\sqrt{3}r$) is less than the distance between the y-intercepts, the rotor will fit. The y-intercepts occur when $\theta = \frac{\pi}{2}$ or $\theta = \frac{3\pi}{2}$ $\Rightarrow$ $y = -b + 3r$ or $y = b - 3r$, so the distance between the intercepts is $(-b + 3r) - (b - 3r) = 6r - 2b$, and the rotor will fit if $3\sqrt{3}r \le 6r - 2b \quad \Leftrightarrow \quad 2b \le 6r - 3\sqrt{3}r \quad \Leftrightarrow \quad b \le \frac{3}{2}(2 - \sqrt{3})r$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11 ☐ INFINITE SEQUENCES AND SERIES

11.1 Sequences

1. (a) A sequence is an ordered list of numbers. It can also be defined as a function whose domain is the set of positive integers.

(b) The terms a_n approach 8 as n becomes large. In fact, we can make a_n as close to 8 as we like by taking n sufficiently large.

(c) The terms a_n become large as n becomes large. In fact, we can make a_n as large as we like by taking n sufficiently large.

2. (a) From Definition 1, a convergent sequence is a sequence for which $\lim_{n\to\infty} a_n$ exists. Examples: $\{1/n\}$, $\{1/2^n\}$

(b) A divergent sequence is a sequence for which $\lim_{n\to\infty} a_n$ *does not* exist. Examples: $\{n\}$, $\{\sin n\}$

3. $a_n = \dfrac{2^n}{2n+1}$, so the sequence is $\left\{\dfrac{2^1}{2(1)+1}, \dfrac{2^2}{2(2)+1}, \dfrac{2^3}{2(3)+1}, \dfrac{2^4}{2(4)+1}, \dfrac{2^5}{2(5)+1}, \ldots\right\} = \left\{\dfrac{2}{3}, \dfrac{4}{5}, \dfrac{8}{7}, \dfrac{16}{9}, \dfrac{32}{11}, \ldots\right\}$.

4. $a_n = \dfrac{n^2-1}{n^2+1}$, so the sequence is $\left\{\dfrac{1-1}{1+1}, \dfrac{4-1}{4+1}, \dfrac{9-1}{9+1}, \dfrac{16-1}{16+1}, \dfrac{25-1}{25+1}, \ldots\right\} = \left\{0, \dfrac{3}{5}, \dfrac{8}{10}, \dfrac{15}{17}, \dfrac{24}{26}, \ldots\right\}$.

5. $a_n = \dfrac{(-1)^{n-1}}{5^n}$, so the sequence is $\left\{\dfrac{1}{5^1}, \dfrac{-1}{5^2}, \dfrac{1}{5^3}, \dfrac{-1}{5^4}, \dfrac{1}{5^5}, \ldots\right\} = \left\{\dfrac{1}{5}, -\dfrac{1}{25}, \dfrac{1}{125}, -\dfrac{1}{625}, \dfrac{1}{3125}, \ldots\right\}$.

6. $a_n = \cos\dfrac{n\pi}{2}$, so the sequence is $\left\{\cos\dfrac{\pi}{2}, \cos\pi, \cos\dfrac{3\pi}{2}, \cos 2\pi, \cos\dfrac{5\pi}{2}, \ldots\right\} = \{0, -1, 0, 1, 0, \ldots\}$.

7. $a_n = \dfrac{1}{(n+1)!}$, so the sequence is $\left\{\dfrac{1}{2!}, \dfrac{1}{3!}, \dfrac{1}{4!}, \dfrac{1}{5!}, \dfrac{1}{6!}, \ldots\right\} = \left\{\dfrac{1}{2}, \dfrac{1}{6}, \dfrac{1}{24}, \dfrac{1}{120}, \dfrac{1}{720}, \ldots\right\}$.

8. $a_n = \dfrac{(-1)^n n}{n!+1}$, so $a_1 = \dfrac{(-1)^1 1}{1!+1} = \dfrac{-1}{2}$, and the sequence is

$\left\{\dfrac{-1}{2}, \dfrac{2}{2+1}, \dfrac{-3}{6+1}, \dfrac{4}{24+1}, \dfrac{-5}{120+1}, \ldots\right\} = \left\{-\dfrac{1}{2}, \dfrac{2}{3}, -\dfrac{3}{7}, \dfrac{4}{25}, -\dfrac{5}{121}, \ldots\right\}$.

9. $a_1 = 1$, $a_{n+1} = 5a_n - 3$. Each term is defined in terms of the preceding term. $a_2 = 5a_1 - 3 = 5(1) - 3 = 2$.

$a_3 = 5a_2 - 3 = 5(2) - 3 = 7$. $a_4 = 5a_3 - 3 = 5(7) - 3 = 32$. $a_5 = 5a_4 - 3 = 5(32) - 3 = 157$.

The sequence is $\{1, 2, 7, 32, 157, \ldots\}$.

10. $a_1 = 6$, $a_{n+1} = \dfrac{a_n}{n}$. $a_2 = \dfrac{a_1}{1} = \dfrac{6}{1} = 6$. $a_3 = \dfrac{a_2}{2} = \dfrac{6}{2} = 3$. $a_4 = \dfrac{a_3}{3} = \dfrac{3}{3} = 1$. $a_5 = \dfrac{a_4}{4} = \dfrac{1}{4}$.

The sequence is $\left\{6, 6, 3, 1, \frac{1}{4}, \ldots\right\}$.

11. $a_1 = 2$, $a_{n+1} = \dfrac{a_n}{1+a_n}$. $a_2 = \dfrac{a_1}{1+a_1} = \dfrac{2}{1+2} = \dfrac{2}{3}$. $a_3 = \dfrac{a_2}{1+a_2} = \dfrac{2/3}{1+2/3} = \dfrac{2}{5}$. $a_4 = \dfrac{a_3}{1+a_3} = \dfrac{2/5}{1+2/5} = \dfrac{2}{7}$.

$a_5 = \dfrac{a_4}{1+a_4} = \dfrac{2/7}{1+2/7} = \dfrac{2}{9}$. The sequence is $\left\{2, \frac{2}{3}, \frac{2}{5}, \frac{2}{7}, \frac{2}{9}, \ldots\right\}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

12. $a_1 = 2, a_2 = 1, a_{n+1} = a_n - a_{n-1}$. Each term is defined in term of the two preceding terms.

$a_3 = a_2 - a_1 = 1 - 2 = -1$. $a_4 = a_3 - a_2 = -1 - 1 = -2$. $a_5 = a_4 - a_3 = -2 - (-1) = -1$.

$a_6 = a_5 - a_4 = -1 - (-2) = 1$. The sequence is $\{2, 1, -1, -2, -1, 1, \ldots\}$.

13. $\left\{\frac{1}{2}, \frac{1}{4}, \frac{1}{6}, \frac{1}{8}, \frac{1}{10}, \ldots\right\}$. The denominator is two times the number of the term, n, so $a_n = \dfrac{1}{2n}$.

14. $\left\{4, -1, \frac{1}{4}, -\frac{1}{16}, \frac{1}{64}, \ldots\right\}$. The first term is 4 and each term is $-\frac{1}{4}$ times the preceding one, so $a_n = 4\left(-\frac{1}{4}\right)^{n-1}$.

15. $\left\{-3, 2, -\frac{4}{3}, \frac{8}{9}, -\frac{16}{27}, \ldots\right\}$. The first term is -3 and each term is $-\frac{2}{3}$ times the preceding one, so $a_n = -3\left(-\frac{2}{3}\right)^{n-1}$.

16. $\{5, 8, 11, 14, 17, \ldots\}$. Each term is larger than the preceding term by 3, so $a_n = a_1 + d(n-1) = 5 + 3(n-1) = 3n + 2$.

17. $\left\{\frac{1}{2}, -\frac{4}{3}, \frac{9}{4}, -\frac{16}{5}, \frac{25}{6}, \ldots\right\}$. The numerator of the nth term is n^2 and its denominator is $n + 1$. Including the alternating signs, we get $a_n = (-1)^{n+1}\dfrac{n^2}{n+1}$.

18. $\{1, 0, -1, 0, 1, 0, -1, 0, \ldots\}$. Two possibilities are $a_n = \sin\dfrac{n\pi}{2}$ and $a_n = \cos\dfrac{(n-1)\pi}{2}$.

19.

n	$a_n = \dfrac{3n}{1+6n}$
1	0.4286
2	0.4615
3	0.4737
4	0.4800
5	0.4839
6	0.4865
7	0.4884
8	0.4898
9	0.4909
10	0.4918



It appears that $\lim\limits_{n\to\infty} a_n = 0.5$.

$$\lim_{n\to\infty}\frac{3n}{1+6n} = \lim_{n\to\infty}\frac{(3n)/n}{(1+6n)/n} = \lim_{n\to\infty}\frac{3}{1/n+6} = \frac{3}{6} = \frac{1}{2}$$

20.

n	$a_n = 2 + \dfrac{(-1)^n}{n}$
1	1.0000
2	2.5000
3	1.6667
4	2.2500
5	1.8000
6	2.1667
7	1.8571
8	2.1250
9	1.8889
10	2.1000



It appears that $\lim\limits_{n\to\infty} a_n = 2$.

$$\lim_{n\to\infty}\left(2 + \frac{(-1)^n}{n}\right) = \lim_{n\to\infty} 2 + \lim_{n\to\infty}\frac{(-1)^n}{n} = 2 + 0 = 2 \text{ since } \lim_{n\to\infty}\frac{1}{n} = 0$$

and by Theorem 6, $\lim\limits_{n\to\infty}\dfrac{(-1)^n}{n} = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21.

n	$a_n = 1 + \left(-\frac{1}{2}\right)^n$
1	0.5000
2	1.2500
3	0.8750
4	1.0625
5	0.9688
6	1.0156
7	0.9922
8	1.0039
9	0.9980
10	1.0010



It appears that $\lim_{n\to\infty} a_n = 1$.

$\lim_{n\to\infty} \left(1 + \left(-\frac{1}{2}\right)^n\right) = \lim_{n\to\infty} 1 + \lim_{n\to\infty} \left(-\frac{1}{2}\right)^n = 1 + 0 = 1$ since $\lim_{n\to\infty} \left(-\frac{1}{2}\right)^n = 0$ by (9).

22.

n	$a_n = 1 + \dfrac{10^n}{9^n}$
1	2.1111
2	2.2346
3	2.3717
4	2.5242
5	2.6935
6	2.8817
7	3.0908
8	3.3231
9	3.5812
10	3.8680



It appears that the sequence does not have a limit.

$\lim_{n\to\infty} \dfrac{10^n}{9^n} = \lim_{n\to\infty} \left(\dfrac{10}{9}\right)^n$, which diverges by (9) since $\frac{10}{9} > 1$.

23. $a_n = \dfrac{3+5n^2}{n+n^2} = \dfrac{(3+5n^2)/n^2}{(n+n^2)/n^2} = \dfrac{5+3/n^2}{1+1/n}$, so $a_n \to \dfrac{5+0}{1+0} = 5$ as $n \to \infty$. Converges

24. $a_n = \dfrac{3+5n^2}{1+n} = \dfrac{(3+5n^2)/n}{(1+n)/n} = \dfrac{3/n+5n}{1/n+1}$, so $a_n \to \infty$ as $n \to \infty$ since $\lim_{n\to\infty} \left(\dfrac{3}{n} + 5n\right) = \infty$ and

$\lim_{n\to\infty} \left(\dfrac{1}{n} + 1\right) = 0 + 1 = 1$. Diverges

25. $a_n = \dfrac{n^4}{n^3-2n} = \dfrac{n^4/n^3}{(n^3-2n)/n^3} = \dfrac{n}{1-2/n^2}$, so $a_n \to \infty$ as $n \to \infty$ since $\lim_{n\to\infty} n = \infty$ and

$\lim_{n\to\infty} \left(1 - \dfrac{2}{n^2}\right) = 1 - 0 = 1$. Diverges

26. $a_n = 2 + (0.86)^n \to 2 + 0 = 2$ as $n \to \infty$ since $\lim_{n\to\infty} (0.86)^n = 0$ by (9) with $r = 0.86$. Converges

27. $a_n = 3^n 7^{-n} = \dfrac{3^n}{7^n} = \left(\dfrac{3}{7}\right)^n$, so $\lim_{n\to\infty} a_n = 0$ by (9) with $r = \dfrac{3}{7}$. Converges

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

28. $a_n = \dfrac{3\sqrt{n}}{\sqrt{n}+2} = \dfrac{3\sqrt{n}/\sqrt{n}}{(\sqrt{n}+2)/\sqrt{n}} = \dfrac{3}{1+2/\sqrt{n}} \to \dfrac{3}{1+0} = 3$ as $n \to \infty$. Converges

29. Because the natural exponential function is continuous at 0, Theorem 7 enables us to write

$\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} e^{-1/\sqrt{n}} = e^{\lim_{n\to\infty}(-1/\sqrt{n})} = e^0 = 1$. Converges

30. $a_n = \dfrac{4^n}{1+9^n} = \dfrac{4^n/9^n}{(1+9^n)/9^n} = \dfrac{(4/9)^n}{(1/9)^n+1} \to \dfrac{0}{0+1} = 0$ as $n \to \infty$ since $\lim\limits_{n\to\infty}\left(\dfrac{4}{9}\right)^n = 0$ and

$\lim\limits_{n\to\infty}\left(\dfrac{1}{9}\right)^n = 0$ by (9). Converges

31. $a_n = \sqrt{\dfrac{1+4n^2}{1+n^2}} = \sqrt{\dfrac{(1+4n^2)/n^2}{(1+n^2)/n^2}} = \sqrt{\dfrac{(1/n^2)+4}{(1/n^2)+1}} \to \sqrt{4} = 2$ as $n \to \infty$ since $\lim\limits_{n\to\infty}(1/n^2) = 0$. Converges

32. $a_n = \cos\left(\dfrac{n\pi}{n+1}\right) = \cos\left(\dfrac{n\pi/n}{(n+1)/n}\right) = \cos\left(\dfrac{\pi}{1+1/n}\right)$, so $a_n \to \cos\pi = -1$ as $n \to \infty$ since $\lim\limits_{n\to\infty} 1/n = 0$.

Converges

33. $a_n = \dfrac{n^2}{\sqrt{n^3+4n}} = \dfrac{n^2/\sqrt{n^3}}{\sqrt{n^3+4n}/\sqrt{n^3}} = \dfrac{\sqrt{n}}{\sqrt{1+4/n^2}}$, so $a_n \to \infty$ as $n \to \infty$ since $\lim\limits_{n\to\infty}\sqrt{n} = \infty$ and

$\lim\limits_{n\to\infty}\sqrt{1+4/n^2} = 1$. Diverges

34. If $b_n = \dfrac{2n}{n+2}$, then $\lim\limits_{n\to\infty} b_n = \lim\limits_{n\to\infty}\dfrac{(2n)/n}{(n+2)/n} = \lim\limits_{n\to\infty}\dfrac{2}{1+2/n} = \dfrac{2}{1} = 2$. Since the natural exponential function is

continuous at 2, by Theorem 7, $\lim\limits_{n\to\infty} e^{2n/(n+2)} = e^{\lim_{n\to\infty} b_n} = e^2$. Converges

35. $\lim\limits_{n\to\infty}|a_n| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)^n}{2\sqrt{n}}\right| = \dfrac{1}{2}\lim\limits_{n\to\infty}\dfrac{1}{n^{1/2}} = \dfrac{1}{2}(0) = 0$, so $\lim\limits_{n\to\infty} a_n = 0$ by (6). Converges

36. $\lim\limits_{n\to\infty}\dfrac{n}{n+\sqrt{n}} = \lim\limits_{n\to\infty}\dfrac{n/n}{(n+\sqrt{n})/n} = \lim\limits_{n\to\infty}\dfrac{1}{1+1/\sqrt{n}} = \dfrac{1}{1+0} = 1$. Thus, $a_n = \dfrac{(-1)^{n+1}n}{n+\sqrt{n}}$ has odd-numbered terms

that approach 1 and even-numbered terms that approach -1 as $n \to \infty$, and hence, the sequence $\{a_n\}$ is divergent.

37. $a_n = \dfrac{(2n-1)!}{(2n+1)!} = \dfrac{(2n-1)!}{(2n+1)(2n)(2n-1)!} = \dfrac{1}{(2n+1)(2n)} \to 0$ as $n \to \infty$. Converges

38. $a_n = \dfrac{\ln n}{\ln 2n} = \dfrac{\ln n}{\ln 2 + \ln n} = \dfrac{1}{\frac{\ln 2}{\ln n}+1} \to \dfrac{1}{0+1} = 1$ as $n \to \infty$. Converges

39. $a_n = \sin n$. This sequence diverges since the terms don't approach any particular real number as $n \to \infty$. The terms take on values between -1 and 1. Diverges

40. $a_n = \dfrac{\tan^{-1} n}{n}$. $\lim\limits_{n\to\infty}\tan^{-1} n = \lim\limits_{x\to\infty}\tan^{-1} x = \dfrac{\pi}{2}$ by (3), so $\lim\limits_{n\to\infty} a_n = 0$. Converges

41. $a_n = n^2e^{-n} = \dfrac{n^2}{e^n}$. Since $\lim\limits_{x\to\infty}\dfrac{x^2}{e^x} \overset{\text{H}}{=} \lim\limits_{x\to\infty}\dfrac{2x}{e^x} \overset{\text{H}}{=} \lim\limits_{x\to\infty}\dfrac{2}{e^x} = 0$, it follows from Theorem 3 that $\lim\limits_{n\to\infty} a_n = 0$. Converges

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $a_n = \ln(n+1) - \ln n = \ln\left(\dfrac{n+1}{n}\right) = \ln\left(1 + \dfrac{1}{n}\right) \to \ln(1) = 0$ as $n \to \infty$ because ln is continuous. Converges

43. $0 \le \dfrac{\cos^2 n}{2^n} \le \dfrac{1}{2^n}$ [since $0 \le \cos^2 n \le 1$], so since $\lim\limits_{n\to\infty} \dfrac{1}{2^n} = 0$, $\left\{\dfrac{\cos^2 n}{2^n}\right\}$ converges to 0 by the Squeeze Theorem.

44. $a_n = \sqrt[n]{2^{1+3n}} = (2^{1+3n})^{1/n} = (2^1 2^{3n})^{1/n} = 2^{1/n}2^3 = 8 \cdot 2^{1/n}$, so

$\lim\limits_{n\to\infty} a_n = 8 \lim\limits_{n\to\infty} 2^{1/n} = 8 \cdot 2^{\lim_{n\to\infty}(1/n)} = 8 \cdot 2^0 = 8$ by Theorem 7, since the function $f(x) = 2^x$ is continuous at 0.

Converges

45. $a_n = n\sin(1/n) = \dfrac{\sin(1/n)}{1/n}$. Since $\lim\limits_{x\to\infty} \dfrac{\sin(1/x)}{1/x} = \lim\limits_{t\to 0^+} \dfrac{\sin t}{t}$ [where $t = 1/x$] $= 1$, it follows from Theorem 3

that $\{a_n\}$ converges to 1.

46. $a_n = 2^{-n}\cos n\pi$. $0 \le \left|\dfrac{\cos n\pi}{2^n}\right| \le \dfrac{1}{2^n} = \left(\dfrac{1}{2}\right)^n$, so $\lim\limits_{n\to\infty} |a_n| = 0$ by (9), and $\lim\limits_{n\to\infty} a_n = 0$ by (6). Converges

47. $y = \left(1 + \dfrac{2}{x}\right)^x \quad \Rightarrow \quad \ln y = x\ln\left(1 + \dfrac{2}{x}\right)$, so

$$\lim_{x\to\infty} \ln y = \lim_{x\to\infty} \frac{\ln(1+2/x)}{1/x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{\left(\dfrac{1}{1+2/x}\right)\left(-\dfrac{2}{x^2}\right)}{-1/x^2} = \lim_{x\to\infty} \frac{2}{1+2/x} = 2 \quad \Rightarrow$$

$\lim\limits_{x\to\infty} \left(1 + \dfrac{2}{x}\right)^x = \lim\limits_{x\to\infty} e^{\ln y} = e^2$, so by Theorem 3, $\lim\limits_{n\to\infty} \left(1 + \dfrac{2}{n}\right)^n = e^2$. Converges

48. $y = x^{1/x} \quad \Rightarrow \quad \ln y = \dfrac{1}{x}\ln x$, so $\lim\limits_{x\to\infty} \ln y = \lim\limits_{x\to\infty} \dfrac{\ln x}{x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1/x}{1} = \lim\limits_{x\to\infty} \dfrac{1}{x} = 0 \quad \Rightarrow$

$\lim\limits_{x\to\infty} x^{1/x} = \lim\limits_{x\to\infty} e^{\ln y} = e^0 = 1$, so by Theorem 3, $\lim\limits_{n\to\infty} \sqrt[n]{n} = 1$. Converges

49. $a_n = \ln(2n^2+1) - \ln(n^2+1) = \ln\left(\dfrac{2n^2+1}{n^2+1}\right) = \ln\left(\dfrac{2+1/n^2}{1+1/n^2}\right) \to \ln 2$ as $n \to \infty$. Converges

50. $\lim\limits_{x\to\infty} \dfrac{(\ln x)^2}{x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{2(\ln x)(1/x)}{1} = 2\lim\limits_{x\to\infty} \dfrac{\ln x}{x} \overset{\text{H}}{=} 2\lim\limits_{x\to\infty} \dfrac{1/x}{1} = 0$, so by Theorem 3, $\lim\limits_{n\to\infty} \dfrac{(\ln n)^2}{n} = 0$. Converges

51. $a_n = \arctan(\ln n)$. Let $f(x) = \arctan(\ln x)$. Then $\lim\limits_{x\to\infty} f(x) = \frac{\pi}{2}$ since $\ln x \to \infty$ as $x \to \infty$ and arctan is continuous.

Thus, $\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} f(n) = \frac{\pi}{2}$. Converges

52. $a_n = n - \sqrt{n+1}\sqrt{n+3} = n - \sqrt{n^2+4n+3} = \dfrac{n - \sqrt{n^2+4n+3}}{1} \cdot \dfrac{n + \sqrt{n^2+4n+3}}{n + \sqrt{n^2+4n+3}}$

$$= \frac{n^2 - (n^2+4n+3)}{n + \sqrt{n^2+4n+3}} = \frac{-4n-3}{n + \sqrt{n^2+4n+3}} = \frac{(-4n-3)/n}{\left(n + \sqrt{n^2+4n+3}\right)/n} = \frac{-4 - 3/n}{1 + \sqrt{1+4/n+3/n^2}},$$

so $\lim\limits_{n\to\infty} a_n = \dfrac{-4-0}{1+\sqrt{1+0+0}} = \dfrac{-4}{2} = -2$. Converges

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

53. $\{0, 1, 0, 0, 1, 0, 0, 0, 1, \ldots\}$ diverges since the sequence takes on only two values, 0 and 1, and never stays arbitrarily close to either one (or any other value) for n sufficiently large.

54. $\left\{\frac{1}{1}, \frac{1}{3}, \frac{1}{2}, \frac{1}{4}, \frac{1}{3}, \frac{1}{5}, \frac{1}{4}, \frac{1}{6}, \ldots\right\}$. $a_{2n-1} = \dfrac{1}{n}$ and $a_{2n} = \dfrac{1}{n+2}$ for all positive integers n. $\lim\limits_{n\to\infty} a_n = 0$ since $\lim\limits_{n\to\infty} a_{2n-1} = \lim\limits_{n\to\infty} \dfrac{1}{n} = 0$ and $\lim\limits_{n\to\infty} a_{2n} = \lim\limits_{n\to\infty} \dfrac{1}{n+2} = 0$. For n sufficiently large, a_n can be made as close to 0 as we like. Converges

55. $a_n = \dfrac{n!}{2^n} = \dfrac{1}{2}\cdot\dfrac{2}{2}\cdot\dfrac{3}{2}\cdot\cdots\cdot\dfrac{(n-1)}{2}\cdot\dfrac{n}{2} \geq \dfrac{1}{2}\cdot\dfrac{n}{2}$ [for $n > 1$] $= \dfrac{n}{4} \to \infty$ as $n \to \infty$, so $\{a_n\}$ diverges.

56. $0 < |a_n| = \dfrac{3^n}{n!} = \dfrac{3}{1}\cdot\dfrac{3}{2}\cdot\dfrac{3}{3}\cdot\cdots\cdot\dfrac{3}{(n-1)}\cdot\dfrac{3}{n} \leq \dfrac{3}{1}\cdot\dfrac{3}{2}\cdot\dfrac{3}{n}$ [for $n > 2$] $= \dfrac{27}{2n} \to 0$ as $n \to \infty$, so by the Squeeze Theorem and Theorem 6, $\{(-3)^n/n!\}$ converges to 0.

57.



From the graph, it appears that the sequence $\{a_n\} = \left\{(-1)^n \dfrac{n}{n+1}\right\}$ is divergent, since it oscillates between 1 and -1 (approximately). To prove this, suppose that $\{a_n\}$ converges to L. If $b_n = \dfrac{n}{n+1}$, then $\{b_n\}$ converges to 1, and $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \dfrac{L}{1} = L$. But $\dfrac{a_n}{b_n} = (-1)^n$, so $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n}$ does not exist. This contradiction shows that $\{a_n\}$ diverges.

58.



From the graph, it appears that the sequence converges to 0.

$|a_n| = \left|\dfrac{\sin n}{n}\right| = \dfrac{|\sin n|}{|n|} \leq \dfrac{1}{n}$, so $\lim\limits_{n\to\infty} |a_n| = 0$. By (6), it follows that $\lim\limits_{n\to\infty} a_n = 0$.

59.



From the graph, it appears that the sequence converges to a number between 0.7 and 0.8.

$a_n = \arctan\left(\dfrac{n^2}{n^2+4}\right) = \arctan\left(\dfrac{n^2/n^2}{(n^2+4)/n^2}\right) = \arctan\left(\dfrac{1}{1+4/n^2}\right) \to \arctan 1 = \dfrac{\pi}{4}$ $[\approx 0.785]$ as $n \to \infty$.

60.



From the graph, it appears that the sequence converges to 5.

$$5 = \sqrt[n]{5^n} \leq \sqrt[n]{3^n + 5^n} \leq \sqrt[n]{5^n + 5^n} = \sqrt[n]{2}\,\sqrt[n]{5^n}$$

$$= \sqrt[n]{2}\cdot 5 \to 5 \text{ as } n \to \infty \quad \left[\lim_{n\to\infty} 2^{1/n} = 2^0 = 1\right]$$

Hence, $a_n \to 5$ by the Squeeze Theorem.

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Alternate solution: Let $y = (3^x + 5^x)^{1/x}$. Then

$$\lim_{x\to\infty} \ln y = \lim_{x\to\infty} \frac{\ln(3^x + 5^x)}{x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{3^x \ln 3 + 5^x \ln 5}{3^x + 5^x} = \lim_{x\to\infty} \frac{\left(\frac{3}{5}\right)^x \ln 3 + \ln 5}{\left(\frac{3}{5}\right)^x + 1} = \ln 5,$$

so $\lim_{x\to\infty} y = e^{\ln 5} = 5$, and so $\left\{\sqrt[n]{3^n + 5^n}\right\}$ converges to 5.

61.



From the graph, it appears that the sequence $\{a_n\} = \left\{\dfrac{n^2 \cos n}{1 + n^2}\right\}$ is divergent, since it oscillates between 1 and -1 (approximately). To prove this, suppose that $\{a_n\}$ converges to L. If $b_n = \dfrac{n^2}{1 + n^2}$, then $\{b_n\}$ converges to 1, and $\lim_{n\to\infty} \dfrac{a_n}{b_n} = \dfrac{L}{1} = L$. But $\dfrac{a_n}{b_n} = \cos n$, so $\lim_{n\to\infty} \dfrac{a_n}{b_n}$ does not exist. This contradiction shows that $\{a_n\}$ diverges.

62.





From the graphs, it seems that the sequence diverges. $a_n = \dfrac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{n!}$. We first prove by induction that $a_n \geq \left(\dfrac{3}{2}\right)^{n-1}$ for all n. This is clearly true for $n = 1$, so let $P(n)$ be the statement that the above is true for n. We must show it is then true for $n + 1$. $a_{n+1} = a_n \cdot \dfrac{2n+1}{n+1} \geq \left(\dfrac{3}{2}\right)^{n-1} \cdot \dfrac{2n+1}{n+1}$ (induction hypothesis). But $\dfrac{2n+1}{n+1} \geq \dfrac{3}{2}$ [since $2(2n+1) \geq 3(n+1) \iff 4n+2 \geq 3n+3 \iff n \geq 1$], and so we get that $a_{n+1} \geq \left(\frac{3}{2}\right)^{n-1} \cdot \frac{3}{2} = \left(\frac{3}{2}\right)^n$ which is $P(n+1)$. Thus, we have proved our first assertion, so since $\left\{\left(\frac{3}{2}\right)^{n-1}\right\}$ diverges [by (9)], so does the given sequence $\{a_n\}$.

63.



From the graph, it appears that the sequence approaches 0.

$$0 < a_n = \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{(2n)^n} = \frac{1}{2n} \cdot \frac{3}{2n} \cdot \frac{5}{2n} \cdot \cdots \cdot \frac{2n-1}{2n}$$

$$\leq \frac{1}{2n} \cdot (1) \cdot (1) \cdot \cdots \cdot (1) = \frac{1}{2n} \to 0 \text{ as } n \to \infty$$

So by the Squeeze Theorem, $\left\{\dfrac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{(2n)^n}\right\}$ converges to 0.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

64. (a) $a_1 = 1$, $a_{n+1} = 4 - a_n$ for $n \geq 1$. $a_1 = 1$, $a_2 = 4 - a_1 = 4 - 1 = 3$, $a_3 = 4 - a_2 = 4 - 3 = 1$, $a_4 = 4 - a_3 = 4 - 1 = 3$, $a_5 = 4 - a_4 = 4 - 3 = 1$. Since the terms of the sequence alternate between 1 and 3, the sequence is divergent.

(b) $a_1 = 2$, $a_2 = 4 - a_1 = 4 - 2 = 2$, $a_3 = 4 - a_2 = 4 - 2 = 2$. Since all of the terms are 2, $\lim_{n\to\infty} a_n = 2$ and hence, the sequence is convergent.

65. (a) $a_n = 1000(1.06)^n \;\Rightarrow\; a_1 = 1060$, $a_2 = 1123.60$, $a_3 = 1191.02$, $a_4 = 1262.48$, and $a_5 = 1338.23$.

(b) $\lim_{n\to\infty} a_n = 1000 \lim_{n\to\infty} (1.06)^n$, so the sequence diverges by (9) with $r = 1.06 > 1$.

66. (a) Substitute 1 to 6 for n in $I_n = 100\left(\dfrac{1.0025^n - 1}{0.0025} - n\right)$ to get $I_1 = \$0$, $I_2 = \$0.25$, $I_3 = \$0.75$, $I_4 = \$1.50$, $I_5 = \$2.51$, and $I_6 = \$3.76$.

(b) For two years, use $2 \cdot 12 = 24$ for n to get \$70.28.

67. (a) We are given that the initial population is 5000, so $P_0 = 5000$. The number of catfish increases by 8% per month and is decreased by 300 per month, so $P_1 = P_0 + 8\%P_0 - 300 = 1.08P_0 - 300$, $P_2 = 1.08P_1 - 300$, and so on. Thus, $P_n = 1.08P_{n-1} - 300$.

(b) Using the recursive formula with $P_0 = 5000$, we get $P_1 = 5100$, $P_2 = 5208$, $P_3 = 5325$ (rounding any portion of a catfish), $P_4 = 5451$, $P_5 = 5587$, and $P_6 = 5734$, which is the number of catfish in the pond after six months.

68. $a_{n+1} = \begin{cases} \frac{1}{2}a_n & \text{if } a_n \text{ is an even number} \\ 3a_n + 1 & \text{if } a_n \text{ is an odd number} \end{cases}$ When $a_1 = 11$, the first 40 terms are 11, 34, 17, 52, 26, 13, 40, 20, 10, 5, 16, 8, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4. When $a_1 = 25$, the first 40 terms are 25, 76, 38, 19, 58, 29, 88, 44, 22, 11, 34, 17, 52, 26, 13, 40, 20, 10, 5, 16, 8, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4. The famous Collatz conjecture is that this sequence always reaches 1, regardless of the starting point a_1.

69. If $|r| \geq 1$, then $\{r^n\}$ diverges by (9), so $\{nr^n\}$ diverges also, since $|nr^n| = n\,|r^n| \geq |r^n|$. If $|r| < 1$ then $\lim_{x\to\infty} xr^x = \lim_{x\to\infty} \dfrac{x}{r^{-x}} \overset{\text{H}}{=} \lim_{x\to\infty} \dfrac{1}{(-\ln r)\, r^{-x}} = \lim_{x\to\infty} \dfrac{r^x}{-\ln r} = 0$, so $\lim_{n\to\infty} nr^n = 0$, and hence $\{nr^n\}$ converges whenever $|r| < 1$.

70. (a) Let $\lim_{n\to\infty} a_n = L$. By Definition 2, this means that for every $\varepsilon > 0$ there is an integer N such that $|a_n - L| < \varepsilon$ whenever $n > N$. Thus, $|a_{n+1} - L| < \varepsilon$ whenever $n + 1 > N \;\Leftrightarrow\; n > N - 1$. It follows that $\lim_{n\to\infty} a_{n+1} = L$ and so $\lim_{n\to\infty} a_n = \lim_{n\to\infty} a_{n+1}$.

(b) If $L = \lim_{n\to\infty} a_n$ then $\lim_{n\to\infty} a_{n+1} = L$ also, so L must satisfy $L = 1/(1 + L) \;\Rightarrow\; L^2 + L - 1 = 0 \;\Rightarrow\; L = \frac{-1+\sqrt{5}}{2}$ (since L has to be nonnegative if it exists).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

71. Since $\{a_n\}$ is a decreasing sequence, $a_n > a_{n+1}$ for all $n > 1$. Because all of its terms lie between 5 and 8, $\{a_n\}$ is a bounded sequence. By the Monotonic Sequence Theorem, $\{a_n\}$ is convergent; that is, $\{a_n\}$ has a limit L. L must be less than 8 since $\{a_n\}$ is decreasing, so $5 \le L < 8$.

72. Since $\{a_n\} = \{\cos n\} \approx \{0.54, -0.42, -0.99, -0.65, 0.28, \ldots\}$, the sequence is not monotonic. The sequence is bounded since $-1 \le \cos n \le 1$ for all n.

73. $a_n = \dfrac{1}{2n+3}$ is decreasing since $a_{n+1} = \dfrac{1}{2(n+1)+3} = \dfrac{1}{2n+5} < \dfrac{1}{2n+3} = a_n$ for each $n \ge 1$. The sequence is bounded since $0 < a_n \le \frac{1}{5}$ for all $n \ge 1$. Note that $a_1 = \frac{1}{5}$.

74. $a_n > a_{n+1} \quad \Leftrightarrow \quad \dfrac{1-n}{2+n} > \dfrac{1-(n+1)}{2+(n+1)} \quad \Leftrightarrow \quad \dfrac{1-n}{2+n} > \dfrac{-n}{n+3} \quad \Leftrightarrow \quad -n^2-2n+3 > -n^2-2n \quad \Leftrightarrow \quad 3 > 0$, which is true for all $n \ge 1$, so $\{a_n\}$ is decreasing. Since $a_1 = 0$ and $\lim\limits_{n\to\infty} \dfrac{1-n}{2+n} = \lim\limits_{n\to\infty} \dfrac{1/n-1}{2/n+1} = -1$, the sequence is bounded $-1 < a_n \le 0$.

75. The terms of $a_n = n(-1)^n$ alternate in sign, so the sequence is not monotonic. The first five terms are -1, 2, -3, 4, and -5. Since $\lim\limits_{n\to\infty} |a_n| = \lim\limits_{n\to\infty} n = \infty$, the sequence is not bounded.

76. Since $\{a_n\} = \left\{2 + \dfrac{(-1)^n}{n}\right\} = \{1, 2\frac{1}{2}, 1\frac{2}{3}, \ldots\}$, the sequence is not monotonic. The sequence is bounded since $1 \le a_n \le \frac{5}{2}$ for all n.

77. $a_n = 3 - 2ne^{-n}$. Let $f(x) = 3 - 2xe^{-x}$. Then $f'(x) = 0 - 2[x(-e^{-x}) + e^{-x}] = 2e^{-x}(x-1)$, which is positive for $x > 1$, so f is increasing on $(1, \infty)$. It follows that the sequence $\{a_n\} = \{f(n)\}$ is increasing. The sequence is bounded below by $a_1 = 3 - 2e^{-1} \approx 2.26$ and above by 3, so the sequence is bounded.

78. $a_n = n^3 - 3n + 3$. Let $f(x) = x^3 - 3x + 3$. Then $f'(x) = 3x^2 - 3 = 3(x^2-1)$, which is positive for $x > 1$, so f is increasing on $[1, \infty)$. It follows that the sequence $\{a_n\} = \{f(n)\}$ is increasing. The sequence is bounded below by $a_1 = 1$, but is not bounded above, so it is not bounded.

79. For $\left\{\sqrt{2}, \sqrt{2\sqrt{2}}, \sqrt{2\sqrt{2\sqrt{2}}}, \ldots\right\}$, $a_1 = 2^{1/2}$, $a_2 = 2^{3/4}$, $a_3 = 2^{7/8}$, …, so $a_n = 2^{(2^n-1)/2^n} = 2^{1-(1/2^n)}$.

$\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} 2^{1-(1/2^n)} = 2^1 = 2$.

Alternate solution: Let $L = \lim\limits_{n\to\infty} a_n$. (We could show the limit exists by showing that $\{a_n\}$ is bounded and increasing.) Then L must satisfy $L = \sqrt{2 \cdot L} \quad \Rightarrow \quad L^2 = 2L \quad \Rightarrow \quad L(L-2) = 0$. $L \ne 0$ since the sequence increases, so $L = 2$.

80. (a) Let P_n be the statement that $a_{n+1} \ge a_n$ and $a_n \le 3$. P_1 is obviously true. We will assume that P_n is true and then show that as a consequence P_{n+1} must also be true. $a_{n+2} \ge a_{n+1} \quad \Leftrightarrow \quad \sqrt{2+a_{n+1}} \ge \sqrt{2+a_n} \quad \Leftrightarrow$ $2 + a_{n+1} \ge 2 + a_n \quad \Leftrightarrow \quad a_{n+1} \ge a_n$, which is the induction hypothesis. $a_{n+1} \le 3 \quad \Leftrightarrow \quad \sqrt{2+a_n} \le 3 \quad \Leftrightarrow$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$2 + a_n \le 9 \;\Leftrightarrow\; a_n \le 7$, which is certainly true because we are assuming that $a_n \le 3$. So P_n is true for all n, and so $a_1 \le a_n \le 3$ (showing that the sequence is bounded), and hence by the Monotonic Sequence Theorem, $\lim_{n\to\infty} a_n$ exists.

(b) If $L = \lim_{n\to\infty} a_n$, then $\lim_{n\to\infty} a_{n+1} = L$ also, so $L = \sqrt{2+L} \;\Rightarrow\; L^2 = 2 + L \;\Leftrightarrow\; L^2 - L - 2 = 0 \;\Leftrightarrow$ $(L+1)(L-2) = 0 \;\Leftrightarrow\; L = 2$ [since L can't be negative].

81. $a_1 = 1, a_{n+1} = 3 - \dfrac{1}{a_n}$. We show by induction that $\{a_n\}$ is increasing and bounded above by 3. Let P_n be the proposition that $a_{n+1} > a_n$ and $0 < a_n < 3$. Clearly P_1 is true. Assume that P_n is true. Then $a_{n+1} > a_n \;\Rightarrow\; \dfrac{1}{a_{n+1}} < \dfrac{1}{a_n} \;\Rightarrow$ $-\dfrac{1}{a_{n+1}} > -\dfrac{1}{a_n}$. Now $a_{n+2} = 3 - \dfrac{1}{a_{n+1}} > 3 - \dfrac{1}{a_n} = a_{n+1} \;\Leftrightarrow\; P_{n+1}$. This proves that $\{a_n\}$ is increasing and bounded above by 3, so $1 = a_1 < a_n < 3$, that is, $\{a_n\}$ is bounded, and hence convergent by the Monotonic Sequence Theorem.

If $L = \lim_{n\to\infty} a_n$, then $\lim_{n\to\infty} a_{n+1} = L$ also, so L must satisfy $L = 3 - 1/L \;\Rightarrow\; L^2 - 3L + 1 = 0 \;\Rightarrow\; L = \frac{3 \pm \sqrt{5}}{2}$. But $L > 1$, so $L = \frac{3+\sqrt{5}}{2}$.

82. $a_1 = 2, a_{n+1} = \dfrac{1}{3 - a_n}$. We use induction. Let P_n be the statement that $0 < a_{n+1} \le a_n \le 2$. Clearly P_1 is true, since $a_2 = 1/(3-2) = 1$. Now assume that P_n is true. Then $a_{n+1} \le a_n \;\Rightarrow\; -a_{n+1} \ge -a_n \;\Rightarrow\; 3 - a_{n+1} \ge 3 - a_n \;\Rightarrow$ $a_{n+2} = \dfrac{1}{3 - a_{n+1}} \le \dfrac{1}{3 - a_n} = a_{n+1}$. Also $a_{n+2} > 0$ [since $3 - a_{n+1}$ is positive] and $a_{n+1} \le 2$ by the induction hypothesis, so P_{n+1} is true. To find the limit, we use the fact that $\lim_{n\to\infty} a_n = \lim_{n\to\infty} a_{n+1} \;\Rightarrow\; L = \frac{1}{3-L} \;\Rightarrow$ $L^2 - 3L + 1 = 0 \;\Rightarrow\; L = \frac{3 \pm \sqrt{5}}{2}$. But $L \le 2$, so we must have $L = \frac{3-\sqrt{5}}{2}$.

83. (a) Let a_n be the number of rabbit pairs in the nth month. Clearly $a_1 = 1 = a_2$. In the nth month, each pair that is 2 or more months old (that is, a_{n-2} pairs) will produce a new pair to add to the a_{n-1} pairs already present. Thus, $a_n = a_{n-1} + a_{n-2}$, so that $\{a_n\} = \{f_n\}$, the Fibonacci sequence.

(b) $a_n = \dfrac{f_{n+1}}{f_n} \;\Rightarrow\; a_{n-1} = \dfrac{f_n}{f_{n-1}} = \dfrac{f_{n-1} + f_{n-2}}{f_{n-1}} = 1 + \dfrac{f_{n-2}}{f_{n-1}} = 1 + \dfrac{1}{f_{n-1}/f_{n-2}} = 1 + \dfrac{1}{a_{n-2}}$. If $L = \lim_{n\to\infty} a_n$, then $L = \lim_{n\to\infty} a_{n-1}$ and $L = \lim_{n\to\infty} a_{n-2}$, so L must satisfy $L = 1 + \dfrac{1}{L} \;\Rightarrow\; L^2 - L - 1 = 0 \;\Rightarrow\; L = \frac{1+\sqrt{5}}{2}$ [since L must be positive].

84. (a) If f is continuous, then $f(L) = f\left(\lim_{n\to\infty} a_n\right) = \lim_{n\to\infty} f(a_n) = \lim_{n\to\infty} a_{n+1} = \lim_{n\to\infty} a_n = L$ by Exercise 70(a).

(b) By repeatedly pressing the cosine key on the calculator (that is, taking cosine of the previous answer) until the displayed value stabilizes, we see that $L \approx 0.73909$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

85. (a)



From the graph, it appears that the sequence $\left\{\dfrac{n^5}{n!}\right\}$ converges to 0, that is, $\lim\limits_{n\to\infty} \dfrac{n^5}{n!} = 0$.

(b)





From the first graph, it seems that the smallest possible value of N corresponding to $\varepsilon = 0.1$ is 9, since $n^5/n! < 0.1$ whenever $n \geq 10$, but $9^5/9! > 0.1$. From the second graph, it seems that for $\varepsilon = 0.001$, the smallest possible value for N is 11 since $n^5/n! < 0.001$ whenever $n \geq 12$.

86. Let $\varepsilon > 0$ and let N be any positive integer larger than $\ln(\varepsilon)/\ln|r|$. If $n > N$, then $n > \ln(\varepsilon)/\ln|r| \quad \Rightarrow \quad n\ln|r| < \ln\varepsilon$ [since $|r| < 1 \;\Rightarrow\; \ln|r| < 0$] $\quad\Rightarrow\quad \ln(|r|^n) < \ln\varepsilon \quad\Rightarrow\quad |r|^n < \varepsilon \quad\Rightarrow\quad |r^n - 0| < \varepsilon$, and so by Definition 2, $\lim\limits_{n\to\infty} r^n = 0$.

87. Theorem 6: If $\lim\limits_{n\to\infty} |a_n| = 0$ then $\lim\limits_{n\to\infty} -|a_n| = 0$, and since $-|a_n| \leq a_n \leq |a_n|$, we have that $\lim\limits_{n\to\infty} a_n = 0$ by the Squeeze Theorem.

88. Theorem 7: If $\lim\limits_{n\to\infty} a_n = L$ and the function f is continuous at L, then $\lim\limits_{n\to\infty} f(a_n) = f(L)$.

Proof: We must show that, given a number $\varepsilon > 0$, there is an integer N such that $|f(a_n) - f(L)| < \varepsilon$ whenever $n > N$. Suppose $\varepsilon > 0$. Since f is continuous at L, there is a number $\delta > 0$ such that $|f(x) - f(L)| < \varepsilon$ if $|x - L| < \delta$. Since $\lim\limits_{n\to\infty} a_n = L$, there is an integer N such that $|a_n - L| < \delta$ if $n > N$. Suppose $n > N$. Then $0 < |a_n - L| < \delta$, so $|f(a_n) - f(L)| < \varepsilon$.

89. To Prove: If $\lim\limits_{n\to\infty} a_n = 0$ and $\{b_n\}$ is bounded, then $\lim\limits_{n\to\infty} (a_n b_n) = 0$.

Proof: Since $\{b_n\}$ is bounded, there is a positive number M such that $|b_n| \leq M$ and hence, $|a_n|\,|b_n| \leq |a_n|\,M$ for all $n \geq 1$. Let $\varepsilon > 0$ be given. Since $\lim\limits_{n\to\infty} a_n = 0$, there is an integer N such that $|a_n - 0| < \dfrac{\varepsilon}{M}$ if $n > N$. Then $|a_n b_n - 0| = |a_n b_n| = |a_n|\,|b_n| \leq |a_n|\,M = |a_n - 0|\,M < \dfrac{\varepsilon}{M} \cdot M = \varepsilon$ for all $n > N$. Since ε was arbitrary, $\lim\limits_{n\to\infty} (a_n b_n) = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

90. (a) $\dfrac{b^{n+1}-a^{n+1}}{b-a} = b^n + b^{n-1}a + b^{n-2}a^2 + b^{n-3}a^3 + \cdots + ba^{n-1} + a^n$

$< b^n + b^{n-1}b + b^{n-2}b^2 + b^{n-3}b^3 + \cdots + bb^{n-1} + b^n = (n+1)b^n$

(b) Since $b-a>0$, we have $b^{n+1}-a^{n+1} < (n+1)b^n(b-a) \;\Rightarrow\; b^{n+1}-(n+1)b^n(b-a) < a^{n+1} \;\Rightarrow$ $b^n[(n+1)a-nb] < a^{n+1}$.

(c) With this substitution, $(n+1)a-nb=1$, and so $b^n = \left(1+\dfrac{1}{n}\right)^n < a^{n+1} = \left(1+\dfrac{1}{n+1}\right)^{n+1}$.

(d) With this substitution, we get $\left(1+\dfrac{1}{2n}\right)^n\left(\dfrac{1}{2}\right) < 1 \;\Rightarrow\; \left(1+\dfrac{1}{2n}\right)^n < 2 \;\Rightarrow\; \left(1+\dfrac{1}{2n}\right)^{2n} < 4.$

(e) $a_n < a_{2n}$ since $\{a_n\}$ is increasing, so $a_n < a_{2n} < 4$.

(f) Since $\{a_n\}$ is increasing and bounded above by 4, $a_1 \le a_n \le 4$, and so $\{a_n\}$ is bounded and monotonic, and hence has a limit by the Monotonic Sequence Theorem.

91. (a) First we show that $a > a_1 > b_1 > b$.

$a_1 - b_1 = \frac{a+b}{2} - \sqrt{ab} = \frac{1}{2}\left(a - 2\sqrt{ab} + b\right) = \frac{1}{2}\left(\sqrt{a}-\sqrt{b}\right)^2 > 0$ [since $a>b$] $\;\Rightarrow\; a_1 > b_1$. Also

$a - a_1 = a - \frac{1}{2}(a+b) = \frac{1}{2}(a-b) > 0$ and $b - b_1 = b - \sqrt{ab} = \sqrt{b}\left(\sqrt{b}-\sqrt{a}\right) < 0$, so $a > a_1 > b_1 > b$. In the same way we can show that $a_1 > a_2 > b_2 > b_1$ and so the given assertion is true for $n=1$. Suppose it is true for $n=k$, that is, $a_k > a_{k+1} > b_{k+1} > b_k$. Then

$a_{k+2} - b_{k+2} = \frac{1}{2}(a_{k+1}+b_{k+1}) - \sqrt{a_{k+1}b_{k+1}} = \frac{1}{2}\left(a_{k+1} - 2\sqrt{a_{k+1}b_{k+1}} + b_{k+1}\right) = \frac{1}{2}\left(\sqrt{a_{k+1}} - \sqrt{b_{k+1}}\right)^2 > 0.$

$a_{k+1} - a_{k+2} = a_{k+1} - \frac{1}{2}(a_{k+1}+b_{k+1}) = \frac{1}{2}(a_{k+1}-b_{k+1}) > 0$, and

$b_{k+1} - b_{k+2} = b_{k+1} - \sqrt{a_{k+1}b_{k+1}} = \sqrt{b_{k+1}}\left(\sqrt{b_{k+1}} - \sqrt{a_{k+1}}\right) < 0 \;\Rightarrow\; a_{k+1} > a_{k+2} > b_{k+2} > b_{k+1}$,

so the assertion is true for $n=k+1$. Thus, it is true for all n by mathematical induction.

(b) From part (a) we have $a > a_n > a_{n+1} > b_{n+1} > b_n > b$, which shows that both sequences, $\{a_n\}$ and $\{b_n\}$, are monotonic and bounded. So they are both convergent by the Monotonic Sequence Theorem.

(c) Let $\lim\limits_{n\to\infty} a_n = \alpha$ and $\lim\limits_{n\to\infty} b_n = \beta$. Then $\lim\limits_{n\to\infty} a_{n+1} = \lim\limits_{n\to\infty} \dfrac{a_n+b_n}{2} \;\Rightarrow\; \alpha = \dfrac{\alpha+\beta}{2} \;\Rightarrow$ $2\alpha = \alpha + \beta \;\Rightarrow\; \alpha = \beta$.

92. (a) Let $\varepsilon > 0$. Since $\lim\limits_{n\to\infty} a_{2n} = L$, there exists N_1 such that $|a_{2n} - L| < \varepsilon$ for $n > N_1$. Since $\lim\limits_{n\to\infty} a_{2n+1} = L$, there exists N_2 such that $|a_{2n+1} - L| < \varepsilon$ for $n > N_2$. Let $N = \max\{2N_1, 2N_2+1\}$ and let $n > N$. If n is even, then $n = 2m$ where $m > N_1$, so $|a_n - L| = |a_{2m} - L| < \varepsilon$. If n is odd, then $n = 2m+1$, where $m > N_2$, so $|a_n - L| = |a_{2m+1} - L| < \varepsilon$. Therefore $\lim\limits_{n\to\infty} a_n = L$.

(b) $a_1 = 1$, $a_2 = 1 + \frac{1}{1+1} = \frac{3}{2} = 1.5$, $a_3 = 1 + \frac{1}{5/2} = \frac{7}{5} = 1.4$, $a_4 = 1 + \frac{1}{12/5} = \frac{17}{12} = 1.41\overline{6}$,

$a_5 = 1 + \frac{1}{29/12} = \frac{41}{29} \approx 1.413793$, $a_6 = 1 + \frac{1}{70/29} = \frac{99}{70} \approx 1.414286$, $a_7 = 1 + \frac{1}{169/70} = \frac{239}{169} \approx 1.414201$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$a_8 = 1 + \frac{1}{408/169} = \frac{577}{408} \approx 1.414216$. Notice that $a_1 < a_3 < a_5 < a_7$ and $a_2 > a_4 > a_6 > a_8$. It appears that the odd terms are increasing and the even terms are decreasing. Let's prove that $a_{2n-2} > a_{2n}$ and $a_{2n-1} < a_{2n+1}$ by mathematical induction. Suppose that $a_{2k-2} > a_{2k}$. Then $1 + a_{2k-2} > 1 + a_{2k} \Rightarrow \frac{1}{1+a_{2k-2}} < \frac{1}{1+a_{2k}} \Rightarrow$

$1 + \frac{1}{1+a_{2k-2}} < 1 + \frac{1}{1+a_{2k}} \Rightarrow a_{2k-1} < a_{2k+1} \Rightarrow 1 + a_{2k-1} < 1 + a_{2k+1} \Rightarrow$

$\frac{1}{1+a_{2k-1}} > \frac{1}{1+a_{2k+1}} \Rightarrow 1 + \frac{1}{1+a_{2k-1}} > 1 + \frac{1}{1+a_{2k+1}} \Rightarrow a_{2k} > a_{2k+2}$. We have thus shown, by induction, that the odd terms are increasing and the even terms are decreasing. Also all terms lie between 1 and 2, so both $\{a_n\}$ and $\{b_n\}$ are bounded monotonic sequences and are therefore convergent by the Monotonic Sequence Theorem. Let $\lim_{n\to\infty} a_{2n} = L$. Then $\lim_{n\to\infty} a_{2n+2} = L$ also. We have

$$a_{n+2} = 1 + \frac{1}{1 + 1 + 1/(1+a_n)} = 1 + \frac{1}{(3+2a_n)/(1+a_n)} = \frac{4+3a_n}{3+2a_n}$$

so $a_{2n+2} = \frac{4+3a_{2n}}{3+2a_{2n}}$. Taking limits of both sides, we get $L = \frac{4+3L}{3+2L} \Rightarrow 3L + 2L^2 = 4 + 3L \Rightarrow L^2 = 2 \Rightarrow$ $L = \sqrt{2}$ [since $L > 0$]. Thus, $\lim_{n\to\infty} a_{2n} = \sqrt{2}$. Similarly we find that $\lim_{n\to\infty} a_{2n+1} = \sqrt{2}$. So, by part (a), $\lim_{n\to\infty} a_n = \sqrt{2}$.

93. (a) Suppose $\{p_n\}$ converges to p. Then $p_{n+1} = \frac{bp_n}{a+p_n} \Rightarrow \lim_{n\to\infty} p_{n+1} = \frac{b \lim_{n\to\infty} p_n}{a + \lim_{n\to\infty} p_n} \Rightarrow p = \frac{bp}{a+p} \Rightarrow$

$p^2 + ap = bp \Rightarrow p(p + a - b) = 0 \Rightarrow p = 0$ or $p = b - a$.

(b) $p_{n+1} = \frac{bp_n}{a+p_n} = \frac{\left(\frac{b}{a}\right)p_n}{1 + \frac{p_n}{a}} < \left(\frac{b}{a}\right)p_n$ since $1 + \frac{p_n}{a} > 1$.

(c) By part (b), $p_1 < \left(\frac{b}{a}\right)p_0$, $p_2 < \left(\frac{b}{a}\right)p_1 < \left(\frac{b}{a}\right)^2 p_0$, $p_3 < \left(\frac{b}{a}\right)p_2 < \left(\frac{b}{a}\right)^3 p_0$, etc. In general, $p_n < \left(\frac{b}{a}\right)^n p_0$, so $\lim_{n\to\infty} p_n \le \lim_{n\to\infty} \left(\frac{b}{a}\right)^n \cdot p_0 = 0$ since $b < a$. $\left[\text{By (9), } \lim_{n\to\infty} r^n = 0 \text{ if } -1 < r < 1. \text{ Here } r = \frac{b}{a} \in (0,1).\right]$

(d) Let $a < b$. We first show, by induction, that if $p_0 < b - a$, then $p_n < b - a$ and $p_{n+1} > p_n$.

For $n = 0$, we have $p_1 - p_0 = \frac{bp_0}{a+p_0} - p_0 = \frac{p_0(b - a - p_0)}{a+p_0} > 0$ since $p_0 < b - a$. So $p_1 > p_0$.

Now we suppose the assertion is true for $n = k$, that is, $p_k < b - a$ and $p_{k+1} > p_k$. Then

$b - a - p_{k+1} = b - a - \frac{bp_k}{a+p_k} = \frac{a(b-a) + bp_k - ap_k - bp_k}{a+p_k} = \frac{a(b-a-p_k)}{a+p_k} > 0$ because $p_k < b - a$. So

$p_{k+1} < b - a$. And $p_{k+2} - p_{k+1} = \frac{bp_{k+1}}{a+p_{k+1}} - p_{k+1} = \frac{p_{k+1}(b - a - p_{k+1})}{a+p_{k+1}} > 0$ since $p_{k+1} < b - a$. Therefore, $p_{k+2} > p_{k+1}$. Thus, the assertion is true for $n = k + 1$. It is therefore true for all n by mathematical induction.

A similar proof by induction shows that if $p_0 > b - a$, then $p_n > b - a$ and $\{p_n\}$ is decreasing.

In either case the sequence $\{p_n\}$ is bounded and monotonic, so it is convergent by the Monotonic Sequence Theorem. It then follows from part (a) that $\lim_{n\to\infty} p_n = b - a$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

LABORATORY PROJECT Logistic Sequences

1. To write such a program in Maple it is best to calculate all the points first and then graph them. One possible sequence of commands [taking $p_0 = \frac{1}{2}$ and $k = 1.5$ for the difference equation] is

```
t:='t';p(0):=1/2;k:=1.5;
for j from 1 to 20  do p(j):=k*p(j-1)*(1-p(j-1)) od;
plot([seq([t,p(t)] t=0..20)],t=0..20,p=0..0.5,style=point);
```

In Mathematica, we can use the following program:

```
p[0]=1/2
k=1.5
p[j_]:=k*p[j-1]*(1-p[j-1])
P=Table[p[t],{t,20}]
ListPlot[P]
```

With $p_0 = \frac{1}{2}$ and $k = 1.5$:

n	p_n	n	p_n	n	p_n
0	0.5	7	0.3338465076	14	0.3333373303
1	0.375	8	0.3335895255	15	0.3333353318
2	0.3515625	9	0.3334613309	16	0.3333343326
3	0.3419494629	10	0.3333973076	17	0.3333338329
4	0.3375300416	11	0.3333653143	18	0.3333335831
5	0.3354052689	12	0.3333493223	19	0.3333334582
6	0.3343628617	13	0.3333413274	20	0.3333333958



With $p_0 = \frac{1}{2}$ and $k = 2.5$:

n	p_n	n	p_n	n	p_n
0	0.5	7	0.6004164790	14	0.5999967417
1	0.625	8	0.5997913269	15	0.6000016291
2	0.5859375	9	0.6001042277	16	0.5999991854
3	0.6065368651	10	0.5999478590	17	0.6000004073
4	0.5966247409	11	0.6000260637	18	0.5999997964
5	0.6016591486	12	0.5999869664	19	0.6000001018
6	0.5991635437	13	0.6000065164	20	0.5999999491



Both of these sequences seem to converge (the first to about $\frac{1}{3}$, the second to about 0.60).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

With $p_0 = \frac{7}{8}$ and $k = 1.5$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.3239166554	14	0.3332554829
1	0.1640625	8	0.3284919837	15	0.3332943990
2	0.2057189941	9	0.3308775005	16	0.3333138639
3	0.2450980344	10	0.3320963702	17	0.3333235980
4	0.2775374819	11	0.3327125567	18	0.3333284655
5	0.3007656421	12	0.3330223670	19	0.3333308994
6	0.3154585059	13	0.3331777051	20	0.3333321164



With $p_0 = \frac{7}{8}$ and $k = 2.5$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.6016572368	14	0.5999869815
1	0.2734375	8	0.5991645155	15	0.6000065088
2	0.4966735840	9	0.6004159972	16	0.5999967455
3	0.6249723374	10	0.5997915688	17	0.6000016272
4	0.5859547872	11	0.6001041070	18	0.5999991864
5	0.6065294364	12	0.5999479194	19	0.6000004068
6	0.5966286980	13	0.6000260335	20	0.5999997966



The limit of the sequence seems to depend on k, but not on p_0.

2. With $p_0 = \frac{7}{8}$ and $k = 3.2$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.5830728495	14	0.7990633827
1	0.35	8	0.7779164854	15	0.5137954979
2	0.728	9	0.5528397669	16	0.7993909896
3	0.6336512	10	0.7910654689	17	0.5131681132
4	0.7428395416	11	0.5288988570	18	0.7994451225
5	0.6112926626	12	0.7973275394	19	0.5130643795
6	0.7603646184	13	0.5171082698	20	0.7994538304



It seems that eventually the terms fluctuate between two values (about 0.5 and 0.8 in this case).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

3. With $p_0 = \frac{7}{8}$ and $k = 3.42$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.4523028596	14	0.8442074951
1	0.3740625	8	0.8472194412	15	0.4498025048
2	0.8007579316	9	0.4426802161	16	0.8463823232
3	0.5456427596	10	0.8437633929	17	0.4446659586
4	0.8478752457	11	0.4508474156	18	0.8445284520
5	0.4411212220	12	0.8467373602	19	0.4490464985
6	0.8431438501	13	0.4438243545	20	0.8461207931



With $p_0 = \frac{7}{8}$ and $k = 3.45$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.4670259170	14	0.8403376122
1	0.37734375	8	0.8587488490	15	0.4628875685
2	0.8105962830	9	0.4184824586	16	0.8577482026
3	0.5296783241	10	0.8395743720	17	0.4209559716
4	0.8594612299	11	0.4646778983	18	0.8409445432
5	0.4167173034	12	0.8581956045	19	0.4614610237
6	0.8385707740	13	0.4198508858	20	0.8573758782



From the graphs above, it seems that for k between 3.4 and 3.5, the terms eventually fluctuate between four values. In the graph below, the pattern followed by the terms is $0.395, 0.832, 0.487, 0.869, 0.395, \ldots$. Note that even for $k = 3.42$ (as in the first graph), there are four distinct "branches"; even after 1000 terms, the first and third terms in the pattern differ by about 2×10^{-9}, while the first and fifth terms differ by only 2×10^{-10}. With $p_0 = \frac{7}{8}$ and $k = 3.48$:



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4.



$p_0 = 0.5,\ k = 3.7$



$p_0 = 0.501,\ k = 3.7$



$p_0 = 0.75,\ k = 3.9$



$p_0 = 0.749,\ k = 3.9$



$p_0 = 0.5,\ k = 3.999$

From the graphs, it seems that if p_0 is changed by 0.001, the whole graph changes completely. (Note, however, that this might be partially due to accumulated round-off error in the CAS. These graphs were generated by Maple with 100-digit accuracy, and different degrees of accuracy give different graphs.) There seem to be some some fleeting patterns in these graphs, but on the whole they are certainly very chaotic. As k increases, the graph spreads out vertically, with more extreme values close to 0 or 1.

11.2 Series

1. (a) A sequence is an ordered list of numbers whereas a series is the *sum* of a list of numbers.

(b) A series is convergent if the sequence of partial sums is a convergent sequence. A series is divergent if it is not convergent.

2. $\sum_{n=1}^{\infty} a_n = 5$ means that by adding sufficiently many terms of the series we can get as close as we like to the number 5.

In other words, it means that $\lim_{n\to\infty} s_n = 5$, where s_n is the nth partial sum, that is, $\sum_{i=1}^{n} a_i$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

3. $\sum_{n=1}^{\infty} a_n = \lim_{n\to\infty} s_n = \lim_{n\to\infty} [2 - 3(0.8)^n] = \lim_{n\to\infty} 2 - 3 \lim_{n\to\infty} (0.8)^n = 2 - 3(0) = 2$

4. $\sum_{n=1}^{\infty} a_n = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \dfrac{n^2-1}{4n^2+1} = \lim_{n\to\infty} \dfrac{(n^2-1)/n^2}{(4n^2+1)/n^2} = \lim_{n\to\infty} \dfrac{1-1/n^2}{4+1/n^2} = \dfrac{1-0}{4+0} = \dfrac{1}{4}$

5. For $\sum_{n=1}^{\infty} \dfrac{1}{n^4+n^2}$, $a_n = \dfrac{1}{n^4+n^2}$. $s_1 = a_1 = \dfrac{1}{1^4+1^2} = \dfrac{1}{2} = 0.5$, $s_2 = s_1 + a_2 = \dfrac{1}{2} + \dfrac{1}{16+4} = 0.55$,

$s_3 = s_2 + a_3 \approx 0.5611$, $s_4 = s_3 + a_4 \approx 0.5648$, $s_5 = s_4 + a_5 \approx 0.5663$, $s_6 = s_5 + a_6 \approx 0.5671$,

$s_7 = s_6 + a_7 \approx 0.5675$, and $s_8 = s_7 + a_8 \approx 0.5677$. It appears that the series is convergent.

6. For $\sum_{n=1}^{\infty} \dfrac{1}{\sqrt[3]{n}}$, $a_n = \dfrac{1}{\sqrt[3]{n}}$. $s_1 = a_1 = \dfrac{1}{\sqrt[3]{1}} = 1$, $s_2 = s_1 + a_2 = 1 + \dfrac{1}{\sqrt[3]{2}} \approx 1.7937$,

$s_3 = s_2 + a_3 \approx 2.4871$, $s_4 = s_3 + a_4 \approx 3.1170$, $s_5 = s_4 + a_5 \approx 3.7018$, $s_6 = s_5 + a_6 \approx 4.2521$,

$s_7 = s_6 + a_7 \approx 4.7749$, and $s_8 = s_7 + a_8 \approx 5.2749$. It appears that the series is divergent.

7. For $\sum_{n=1}^{\infty} \sin n$, $a_n = \sin n$. $s_1 = a_1 = \sin 1 \approx 0.8415$, $s_2 = s_1 + a_2 \approx 1.7508$,

$s_3 = s_2 + a_3 \approx 1.8919$, $s_4 = s_3 + a_4 \approx 1.1351$, $s_5 = s_4 + a_5 \approx 0.1762$, $s_6 = s_5 + a_6 \approx -0.1033$,

$s_7 = s_6 + a_7 \approx 0.5537$, and $s_8 = s_7 + a_8 \approx 1.5431$. It appears that the series is divergent.

8. For $\sum_{n=1}^{\infty} \dfrac{(-1)^{n-1}}{n!}$, $a_n = (-1)^{n-1}\dfrac{1}{n!}$. $s_1 = a_1 = \dfrac{1}{1!} = 1$, $s_2 = s_1 + a_2 = 1 - \dfrac{1}{2!} = 0.5$,

$s_3 = s_2 + a_3 = 0.5 + \dfrac{1}{3!} \approx 0.6667$, $s_4 = s_3 + a_4 = 0.625$, $s_5 = s_4 + a_5 \approx 0.6333$, $s_6 = s_5 + a_6 \approx 0.6319$,

$s_7 = s_6 + a_7 \approx 0.6321$, and $s_8 = s_7 + a_8 \approx 0.6321$. It appears that the series is convergent.

9.

n	s_n
1	−2.40000
2	−1.92000
3	−2.01600
4	−1.99680
5	−2.00064
6	−1.99987
7	−2.00003
8	−1.99999
9	−2.00000
10	−2.00000



From the graph and the table, it seems that the series converges to -2. In fact, it is a geometric series with $a = -2.4$ and $r = -\frac{1}{5}$, so its sum is $\sum_{n=1}^{\infty} \dfrac{12}{(-5)^n} = \dfrac{-2.4}{1-\left(-\frac{1}{5}\right)} = \dfrac{-2.4}{1.2} = -2$.

Note that the dot corresponding to $n = 1$ is part of both $\{a_n\}$ and $\{s_n\}$.

TI-86 Note: To graph $\{a_n\}$ and $\{s_n\}$, set your calculator to Param mode and DrawDot mode. (DrawDot is under GRAPH, MORE, FORMT (F3).) Now under `E(t)=` make the assignments: `xt1=t, yt1=12/(-5)^t, xt2=t, yt2=sum seq(yt1,t,1,t,1)`. (sum and seq are under LIST, OPS (F5), MORE.) Under WIND use `1,10,1,0,10,1,-3,1,1` to obtain a graph similar to the one above. Then use TRACE (F4) to see the values.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10.

n	s_n
1	0.54030
2	0.12416
3	−0.86584
4	−1.51948
5	−1.23582
6	−0.27565
7	0.47825
8	0.33275
9	−0.57838
10	−1.41745



The series $\sum_{n=1}^{\infty} \cos n$ diverges, since its terms do not approach 0.

11.

n	s_n
1	0.44721
2	1.15432
3	1.98637
4	2.88080
5	3.80927
6	4.75796
7	5.71948
8	6.68962
9	7.66581
10	8.64639



The series $\sum_{n=1}^{\infty} \dfrac{n}{\sqrt{n^2+4}}$ diverges, since its terms do not approach 0.

12.

n	s_n
1	4.90000
2	8.33000
3	10.73100
4	12.41170
5	13.58819
6	14.41173
7	14.98821
8	15.39175
9	15.67422
10	15.87196



From the graph and the table, we see that the terms are getting smaller and may approach 0, and that the series approaches a value near 16. The series is geometric with $a_1 = 4.9$ and $r = 0.7$, so its sum is $\sum_{n=1}^{\infty} \dfrac{7^{n+1}}{10^n} = \dfrac{4.9}{1-0.7} = \dfrac{4.9}{0.3} = 16.\overline{3}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

13.

n	s_n
2	1.00000
3	1.33333
4	1.50000
5	1.60000
6	1.66667
7	1.71429
8	1.75000
9	1.77778
10	1.80000
11	1.81818



From the graph and the table, we see that the terms are getting smaller and may approach 0, and that the series may approach a number near 2. Using partial fractions, we have

$$\begin{aligned}\sum_{n=2}^{k} \frac{2}{n^2-n} &= \sum_{n=2}^{k} \left(\frac{2}{n-1} - \frac{2}{n}\right) \\ &= \left(\frac{2}{1} - \frac{2}{2}\right) + \left(\frac{2}{2} - \frac{2}{3}\right) + \left(\frac{2}{3} - \frac{2}{4}\right) \\ &\qquad + \cdots + \left(\frac{2}{k-2} - \frac{2}{k-1}\right) + \left(\frac{2}{k-1} - \frac{2}{k}\right) \\ &= 2 - \frac{2}{k}\end{aligned}$$

As $k \to \infty$, $2 - \frac{2}{k} \to 2$, so $\sum_{n=2}^{\infty} \frac{2}{n^2-n} = 2$.

14.

n	s_n
1	0.36205
2	0.51428
3	0.59407
4	0.64280
5	0.67557
6	0.69910
7	0.71680
8	0.73059
9	0.74164
10	0.75069



From the graph and the table, we see that the terms are getting smaller and may approach 0, and that the series may approach a number near 1.

$$\begin{aligned}\sum_{n=1}^{k} \left(\sin\frac{1}{n} - \sin\frac{1}{n+1}\right) &= \left(\sin 1 - \sin\frac{1}{2}\right) + \left(\sin\frac{1}{2} - \sin\frac{1}{3}\right) \\ &\qquad + \cdots + \left(\sin\frac{1}{k-1} + \sin\frac{1}{k}\right) \\ &\qquad + \left(\sin\frac{1}{k} - \sin\frac{1}{k+1}\right) \\ &= \sin 1 - \sin\frac{1}{k+1}\end{aligned}$$

As $k \to \infty$, $\sin 1 - \sin\frac{1}{k+1} \to \sin 1 - \sin 0 = \sin 1$, so

$$\sum_{n=1}^{\infty} \left(\sin\frac{1}{n} - \sin\frac{1}{n+1}\right) = \sin 1 \approx 0.84147.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. (a) $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \dfrac{2n}{3n+1} = \dfrac{2}{3}$, so the *sequence* $\{a_n\}$ is convergent by (11.1.1).

(b) Since $\lim_{n\to\infty} a_n = \frac{2}{3} \neq 0$, the *series* $\sum_{n=1}^{\infty} a_n$ is divergent by the Test for Divergence.

16. (a) Both $\sum_{i=1}^{n} a_i$ and $\sum_{j=1}^{n} a_j$ represent the sum of the first n terms of the sequence $\{a_n\}$, that is, the nth partial sum.

(b) $\sum_{i=1}^{n} a_j = \underbrace{a_j + a_j + \cdots + a_j}_{n \text{ terms}} = na_j$, which, in general, is not the same as $\sum_{i=1}^{n} a_i = a_1 + a_2 + \cdots + a_n$.

17. $3 - 4 + \frac{16}{3} - \frac{64}{9} + \cdots$ is a geometric series with ratio $r = -\frac{4}{3}$. Since $|r| = \frac{4}{3} > 1$, the series diverges.

18. $4 + 3 + \frac{9}{4} + \frac{27}{16} + \cdots$ is a geometric series with ratio $\frac{3}{4}$. Since $|r| = \frac{3}{4} < 1$, the series converges to $\dfrac{a}{1-r} = \dfrac{4}{1-3/4} = 16$.

19. $10 - 2 + 0.4 - 0.08 + \cdots$ is a geometric series with ratio $-\frac{2}{10} = -\frac{1}{5}$. Since $|r| = \frac{1}{5} < 1$, the series converges to

$$\frac{a}{1-r} = \frac{10}{1-(-1/5)} = \frac{10}{6/5} = \frac{50}{6} = \frac{25}{3}.$$

20. $2 + 0.5 + 0.125 + 0.03125 + \cdots$ is a geometric series with ratio $r = \frac{0.5}{2} = \frac{1}{4}$. Since $|r| = \frac{1}{4} < 1$, the series converges to $\dfrac{a}{1-r} = \dfrac{2}{1-1/4} = \dfrac{2}{3/4} = \dfrac{8}{3}$.

21. $\sum_{n=1}^{\infty} 12\,(0.73)^{n-1}$ is a geometric series with first term $a = 12$ and ratio $r = 0.73$. Since $|r| = 0.73 < 1$, the series converges to $\dfrac{a}{1-r} = \dfrac{12}{1-0.73} = \dfrac{12}{0.27} = \dfrac{12(100)}{27} = \dfrac{400}{9}$.

22. $\sum_{n=1}^{\infty} \dfrac{5}{\pi^n} = 5\sum_{n=1}^{\infty} \left(\dfrac{1}{\pi}\right)^n$. The latter series is geometric with $a = \dfrac{1}{\pi}$ and ratio $r = \dfrac{1}{\pi}$. Since $|r| = \dfrac{1}{\pi} < 1$, it converges to $\dfrac{1/\pi}{1-1/\pi} = \dfrac{1}{\pi - 1}$. Thus, the given series converges to $5\left(\dfrac{1}{\pi-1}\right) = \dfrac{5}{\pi-1}$.

23. $\sum_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{4^n} = \dfrac{1}{4}\sum_{n=1}^{\infty} \left(-\dfrac{3}{4}\right)^{n-1}$. The latter series is geometric with $a = 1$ and ratio $r = -\frac{3}{4}$. Since $|r| = \frac{3}{4} < 1$, it converges to $\dfrac{1}{1-(-3/4)} = \frac{4}{7}$. Thus, the given series converges to $\left(\frac{1}{4}\right)\left(\frac{4}{7}\right) = \frac{1}{7}$.

24. $\sum_{n=0}^{\infty} \dfrac{3^{n+1}}{(-2)^n} = 3\sum_{n=0}^{\infty} \left(-\frac{3}{2}\right)^n$ is a geometric series with ratio $r = -\frac{3}{2}$. Since $|r| = \frac{3}{2} > 1$, the series diverges.

25. $\sum_{n=1}^{\infty} \dfrac{e^{2n}}{6^{n-1}} = \sum_{n=1}^{\infty} \dfrac{(e^2)^n}{6^n 6^{-1}} = 6\sum_{n=1}^{\infty} \left(\dfrac{e^2}{6}\right)^n$ is a geometric series with ratio $r = \dfrac{e^2}{6}$. Since $|r| = \dfrac{e^2}{6} [\approx 1.23] > 1$, the series diverges.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

26. $\displaystyle\sum_{n=1}^{\infty}\frac{6\cdot 2^{2n-1}}{3^n}-\sum_{n=1}^{\infty}\frac{6(2^2)^n\cdot 2^{-1}}{3^n}=3\sum_{n=1}^{\infty}\left(\frac{4}{3}\right)^n$ is a geometric series with ratio $r=\frac{4}{3}$. Since $|r|=\frac{4}{3}>1$, the series diverges.

27. $\frac{1}{3}+\frac{1}{6}+\frac{1}{9}+\frac{1}{12}+\frac{1}{15}+\cdots=\sum_{n=1}^{\infty}\frac{1}{3n}=\frac{1}{3}\sum_{n=1}^{\infty}\frac{1}{n}$. This is a constant multiple of the divergent harmonic series, so it diverges.

28. $\frac{1}{3}+\frac{2}{9}+\frac{1}{27}+\frac{2}{81}+\frac{1}{243}+\frac{2}{729}+\cdots=\left(\frac{1}{3}+\frac{1}{27}+\frac{1}{243}+\cdots\right)+\left(\frac{2}{9}+\frac{2}{81}+\frac{2}{729}+\cdots\right)$, which are both convergent geometric series with sums $\dfrac{1/3}{1-1/9}=\dfrac{3}{8}$ and $\dfrac{2/9}{1-1/9}=\dfrac{1}{4}$, so the original series converges and its sum is $\frac{3}{8}+\frac{1}{4}=\frac{5}{8}$.

29. $\displaystyle\sum_{n=1}^{\infty}\frac{2+n}{1-2n}$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty}a_n=\lim_{n\to\infty}\frac{2+n}{1-2n}=\lim_{n\to\infty}\frac{2/n+1}{1/n-2}=-\frac{1}{2}\neq 0$.

30. $\displaystyle\sum_{k=1}^{\infty}\frac{k^2}{k^2-2k+5}$ diverges by the Test for Divergence since $\displaystyle\lim_{k\to\infty}\frac{k^2}{k^2-2k+5}=\lim_{k\to\infty}\frac{1}{1-2/k+5/k^2}=1\neq 0$.

31. $\displaystyle\sum_{n=1}^{\infty}3^{n+1}4^{-n}=\sum_{n=1}^{\infty}\frac{3^n\cdot 3^1}{4^n}=3\sum_{n=1}^{\infty}\left(\frac{3}{4}\right)^n$. The latter series is geometric with $a=\frac{3}{4}$ and ratio $r=\frac{3}{4}$. Since $|r|=\frac{3}{4}<1$, it converges to $\dfrac{3/4}{1-3/4}=3$. Thus, the given series converges to $3(3)=9$.

32.
$$\sum_{n=1}^{\infty}[(-0.2)^n+(0.6)^{n-1}]=\sum_{n=1}^{\infty}(-0.2)^n+\sum_{n=1}^{\infty}(0.6)^{n-1}\qquad\text{[sum of two geometric series]}$$
$$=\frac{-0.2}{1-(-0.2)}+\frac{1}{1-0.6}=-\frac{1}{6}+\frac{5}{2}=\frac{7}{3}$$

33. $\displaystyle\sum_{n=1}^{\infty}\frac{1}{4+e^{-n}}$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty}\frac{1}{4+e^{-n}}=\frac{1}{4+0}=\frac{1}{4}\neq 0$.

34. $\displaystyle\sum_{n=1}^{\infty}\frac{2^n+4^n}{e^n}$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty}\frac{2^n+4^n}{e^n}=\lim_{n\to\infty}\left(\frac{2^n}{e^n}+\frac{4^n}{e^n}\right)\geq\lim_{n\to\infty}\left(\frac{4}{e}\right)^n=\infty$ since $\dfrac{4}{e}>1$.

35. $\displaystyle\sum_{k=1}^{\infty}(\sin 100)^k$ is a geometric series with first term $a=\sin 100\ [\approx -0.506]$ and ratio $r=\sin 100$. Since $|r|<1$, the series converges to $\dfrac{\sin 100}{1-\sin 100}\approx -0.336$.

36. $\displaystyle\sum_{n=1}^{\infty}\frac{1}{1+\left(\frac{2}{3}\right)^n}$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty}\frac{1}{1+\left(\frac{2}{3}\right)^n}=\frac{1}{1+0}=1\neq 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $\sum_{n=1}^{\infty} \ln\left(\frac{n^2+1}{2n^2+1}\right)$ diverges by the Test for Divergence since

$$\lim_{n\to\infty} a_n = \lim_{n\to\infty} \ln\left(\frac{n^2+1}{2n^2+1}\right) = \ln\left(\lim_{n\to\infty} \frac{n^2+1}{2n^2+1}\right) = \ln \tfrac{1}{2} \neq 0.$$

38. $\sum_{k=0}^{\infty} (\sqrt{2})^{-k} = \sum_{k=0}^{\infty} \left(\frac{1}{\sqrt{2}}\right)^k$ is a geometric series with first term $a = \left(\frac{1}{\sqrt{2}}\right)^0 = 1$ and ratio $r = \frac{1}{\sqrt{2}}$. Since $|r| < 1$, the series converges to $\dfrac{1}{1-1/\sqrt{2}} = \dfrac{\sqrt{2}}{\sqrt{2}-1} \approx 3.414$.

39. $\sum_{n=1}^{\infty} \arctan n$ diverges by the Test for Divergence since $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \arctan n = \frac{\pi}{2} \neq 0$.

40. $\sum_{n=1}^{\infty} \left(\frac{3}{5^n} + \frac{2}{n}\right)$ diverges because $\sum_{n=1}^{\infty} \frac{2}{n} = 2\sum_{n=1}^{\infty} \frac{1}{n}$ diverges. (If it converged, then $\frac{1}{2} \cdot 2\sum_{n=1}^{\infty} \frac{1}{n}$ would also converge by Theorem 8(i), but we know from Example 9 that the harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.) If the given series converges, then the difference $\sum_{n=1}^{\infty} \left(\frac{3}{5^n} + \frac{2}{n}\right) - \sum_{n=1}^{\infty} \frac{3}{5^n}$ must converge (since $\sum_{n=1}^{\infty} \frac{3}{5^n}$ is a convergent geometric series) and equal $\sum_{n=1}^{\infty} \frac{2}{n}$, but we have just seen that $\sum_{n=1}^{\infty} \frac{2}{n}$ diverges, so the given series must also diverge.

41. $\sum_{n=1}^{\infty} \frac{1}{e^n} = \sum_{n=1}^{\infty} \left(\frac{1}{e}\right)^n$ is a geometric series with first term $a = \frac{1}{e}$ and ratio $r = \frac{1}{e}$. Since $|r| = \frac{1}{e} < 1$, the series converges to $\dfrac{1/e}{1-1/e} = \dfrac{1/e}{1-1/e} \cdot \dfrac{e}{e} = \dfrac{1}{e-1}$. By Example 8, $\sum_{n=1}^{\infty} \dfrac{1}{n(n+1)} = 1$. Thus, by Theorem 8(ii),

$$\sum_{n=1}^{\infty} \left(\frac{1}{e^n} + \frac{1}{n(n+1)}\right) = \sum_{n=1}^{\infty} \frac{1}{e^n} + \sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \frac{1}{e-1} + 1 = \frac{1}{e-1} + \frac{e-1}{e-1} = \frac{e}{e-1}.$$

42. $\sum_{n=1}^{\infty} \frac{e^n}{n^2}$ diverges by the Test for Divergence since $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{e^n}{n^2} = \lim_{x\to\infty} \frac{e^x}{x^2} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{e^x}{2x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{e^x}{2} = \infty \neq 0$.

43. Using partial fractions, the partial sums of the series $\sum_{n=2}^{\infty} \dfrac{2}{n^2-1}$ are

$$\begin{aligned} s_n &= \sum_{i=2}^{n} \frac{2}{(i-1)(i+1)} = \sum_{i=2}^{n} \left(\frac{1}{i-1} - \frac{1}{i+1}\right) \\ &= \left(1 - \frac{1}{3}\right) + \left(\frac{1}{2} - \frac{1}{4}\right) + \left(\frac{1}{3} - \frac{1}{5}\right) + \cdots + \left(\frac{1}{n-3} - \frac{1}{n-1}\right) + \left(\frac{1}{n-2} - \frac{1}{n}\right) \end{aligned}$$

This sum is a telescoping series and $s_n = 1 + \frac{1}{2} - \dfrac{1}{n-1} - \dfrac{1}{n}$.

Thus, $\sum_{n=2}^{\infty} \dfrac{2}{n^2-1} = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \left(1 + \frac{1}{2} - \dfrac{1}{n-1} - \dfrac{1}{n}\right) = \dfrac{3}{2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

44. For the series $\sum_{n=1}^{\infty} \ln \frac{n}{n+1}$,

$$s_n = (\ln 1 - \ln 2) + (\ln 2 - \ln 3) + (\ln 3 - \ln 4) + \cdots + [\ln n - \ln(n+1)] = \ln 1 - \ln(n+1) = -\ln(n+1)$$

[telescoping series]

Thus, $\lim_{n\to\infty} s_n = -\infty$, so the series is divergent.

45. For the series $\sum_{n=1}^{\infty} \frac{3}{n(n+3)}$, $s_n = \sum_{i=1}^{n} \frac{3}{i(i+3)} = \sum_{i=1}^{n} \left(\frac{1}{i} - \frac{1}{i+3}\right)$ [using partial fractions]. The latter sum is

$$\left(1 - \tfrac{1}{4}\right) + \left(\tfrac{1}{2} - \tfrac{1}{5}\right) + \left(\tfrac{1}{3} - \tfrac{1}{6}\right) + \left(\tfrac{1}{4} - \tfrac{1}{7}\right) + \cdots + \left(\tfrac{1}{n-3} - \tfrac{1}{n}\right) + \left(\tfrac{1}{n-2} - \tfrac{1}{n+1}\right) + \left(\tfrac{1}{n-1} - \tfrac{1}{n+2}\right) + \left(\tfrac{1}{n} - \tfrac{1}{n+3}\right)$$

$$= 1 + \tfrac{1}{2} + \tfrac{1}{3} - \tfrac{1}{n+1} - \tfrac{1}{n+2} - \tfrac{1}{n+3} \quad \text{[telescoping series]}$$

Thus, $\sum_{n=1}^{\infty} \frac{3}{n(n+3)} = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \left(1 + \tfrac{1}{2} + \tfrac{1}{3} - \tfrac{1}{n+1} - \tfrac{1}{n+2} - \tfrac{1}{n+3}\right) = 1 + \tfrac{1}{2} + \tfrac{1}{3} = \tfrac{11}{6}$. Converges

46. For the series $\sum_{n=4}^{\infty} \left(\frac{1}{\sqrt{n}} - \frac{1}{\sqrt{n+1}}\right)$

$$s_n = \sum_{i=4}^{n} \left(\frac{1}{\sqrt{i}} - \frac{1}{\sqrt{i+1}}\right) = \left(\frac{1}{\sqrt{4}} - \frac{1}{\sqrt{5}}\right) + \left(\frac{1}{\sqrt{5}} - \frac{1}{\sqrt{6}}\right) + \cdots + \left(\frac{1}{\sqrt{n}} - \frac{1}{\sqrt{n+1}}\right) = \frac{1}{\sqrt{4}} - \frac{1}{\sqrt{n+1}}$$

[telescoping series]

Thus, $\sum_{n=4}^{\infty} \left(\frac{1}{\sqrt{n}} - \frac{1}{\sqrt{n+1}}\right) = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \left(\frac{1}{\sqrt{4}} - \frac{1}{\sqrt{n+1}}\right) = \frac{1}{\sqrt{4}} - 0 = \frac{1}{2}$. Converges

47. For the series $\sum_{n=1}^{\infty} \left(e^{1/n} - e^{1/(n+1)}\right)$,

$$s_n = \sum_{i=1}^{n} \left(e^{1/i} - e^{1/(i+1)}\right) = (e^1 - e^{1/2}) + (e^{1/2} - e^{1/3}) + \cdots + \left(e^{1/n} - e^{1/(n+1)}\right) = e - e^{1/(n+1)}$$

[telescoping series]

Thus, $\sum_{n=1}^{\infty} \left(e^{1/n} - e^{1/(n+1)}\right) = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \left(e - e^{1/(n+1)}\right) = e - e^0 = e - 1$. Converges

48. Using partial fractions, the partial sums of the series $\sum_{n=2}^{\infty} \frac{1}{n^3 - n}$ are

$$s_n = \sum_{i=2}^{n} \frac{1}{i(i-1)(i+1)} = \sum_{i=2}^{n} \left(-\frac{1}{i} + \frac{1/2}{i-1} + \frac{1/2}{i+1}\right) = \frac{1}{2} \sum_{i=2}^{n} \left(\frac{1}{i-1} - \frac{2}{i} + \frac{1}{i+1}\right)$$

$$= \frac{1}{2}\left[\left(\frac{1}{1} - \frac{2}{2} + \frac{1}{3}\right) + \left(\frac{1}{2} - \frac{2}{3} + \frac{1}{4}\right) + \left(\frac{1}{3} - \frac{2}{4} + \frac{1}{5}\right) + \left(\frac{1}{4} - \frac{2}{5} + \frac{1}{6}\right) + \cdots \right.$$

$$\left. + \left(\frac{1}{n-3} - \frac{2}{n-2} + \frac{1}{n-1}\right) + \left(\frac{1}{n-2} - \frac{2}{n-1} + \frac{1}{n}\right) + \left(\frac{1}{n-1} - \frac{2}{n} + \frac{1}{n+1}\right)\right]$$

Note: In three consecutive expressions in parentheses, the 3rd term in the first expression plus the 2nd term in the second expression plus the 1st term in the third expression sum to 0.

$$= \frac{1}{2}\left(\frac{1}{1} - \frac{2}{2} + \frac{1}{2} + \frac{1}{n} - \frac{2}{n} + \frac{1}{n+1}\right) = \frac{1}{4} - \frac{1}{2n} + \frac{1}{2n+2}$$

Thus, $\sum_{n=2}^{\infty} \frac{1}{n^3 - n} = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \left(\frac{1}{4} - \frac{1}{2n} + \frac{1}{2n+2}\right) = \frac{1}{4}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. (a) Many people would guess that $x < 1$, but note that x consists of an infinite number of 9s.

(b) $x = 0.99999\ldots = \dfrac{9}{10} + \dfrac{9}{100} + \dfrac{9}{1000} + \dfrac{9}{10{,}000} + \cdots = \displaystyle\sum_{n=1}^{\infty} \frac{9}{10^n}$, which is a geometric series with $a_1 = 0.9$ and $r = 0.1$. Its sum is $\dfrac{0.9}{1 - 0.1} = \dfrac{0.9}{0.9} = 1$, that is, $x = 1$.

(c) The number 1 has two decimal representations, $1.00000\ldots$ and $0.99999\ldots$.

(d) Except for 0, all rational numbers that have a terminating decimal representation can be written in more than one way. For example, 0.5 can be written as $0.49999\ldots$ as well as $0.50000\ldots$.

50. $a_1 = 1, a_n = (5 - n)a_{n-1} \;\Rightarrow\; a_2 = (5-2)a_1 = 3(1) = 3,\ a_3 = (5-3)a_2 = 2(3) = 6,\ a_4 = (5-4)a_3 = 1(6) = 6,$ $a_5 = (5-5)a_4 = 0$, and all succeeding terms equal 0. Thus, $\displaystyle\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{4} a_n = 1 + 3 + 6 + 6 = 16.$

51. $0.\overline{8} = \dfrac{8}{10} + \dfrac{8}{10^2} + \cdots$ is a geometric series with $a = \dfrac{8}{10}$ and $r = \dfrac{1}{10}$. It converges to $\dfrac{a}{1 - r} = \dfrac{8/10}{1 - 1/10} = \dfrac{8}{9}$.

52. $0.\overline{46} = \dfrac{46}{100} + \dfrac{46}{100^2} + \cdots$ is a geometric series with $a = \dfrac{46}{100}$ and $r = \dfrac{1}{100}$. It converges to $\dfrac{a}{1 - r} = \dfrac{46/100}{1 - 1/100} = \dfrac{46}{99}$.

53. $2.\overline{516} = 2 + \dfrac{516}{10^3} + \dfrac{516}{10^6} + \cdots$. Now $\dfrac{516}{10^3} + \dfrac{516}{10^6} + \cdots$ is a geometric series with $a = \dfrac{516}{10^3}$ and $r = \dfrac{1}{10^3}$. It converges to $\dfrac{a}{1 - r} = \dfrac{516/10^3}{1 - 1/10^3} = \dfrac{516/10^3}{999/10^3} = \dfrac{516}{999}$. Thus, $2.\overline{516} = 2 + \dfrac{516}{999} = \dfrac{2514}{999} = \dfrac{838}{333}$.

54. $10.1\overline{35} = 10.1 + \dfrac{35}{10^3} + \dfrac{35}{10^5} + \cdots$. Now $\dfrac{35}{10^3} + \dfrac{35}{10^5} + \cdots$ is a geometric series with $a = \dfrac{35}{10^3}$ and $r = \dfrac{1}{10^2}$. It converges to $\dfrac{a}{1 - r} = \dfrac{35/10^3}{1 - 1/10^2} = \dfrac{35/10^3}{99/10^2} = \dfrac{35}{990}$. Thus, $10.1\overline{35} = 10.1 + \dfrac{35}{990} = \dfrac{9999 + 35}{990} = \dfrac{10{,}034}{990} = \dfrac{5017}{495}$.

55. $1.234\overline{567} = 1.234 + \dfrac{567}{10^6} + \dfrac{567}{10^9} + \cdots$. Now $\dfrac{567}{10^6} + \dfrac{567}{10^9} + \cdots$ is a geometric series with $a = \dfrac{567}{10^6}$ and $r = \dfrac{1}{10^3}$. It converges to $\dfrac{a}{1 - r} = \dfrac{567/10^6}{1 - 1/10^3} = \dfrac{567/10^6}{999/10^3} = \dfrac{567}{999{,}000} = \dfrac{21}{37{,}000}$. Thus,

$$1.234\overline{567} = 1.234 + \frac{21}{37{,}000} = \frac{1234}{1000} + \frac{21}{37{,}000} = \frac{45{,}658}{37{,}000} + \frac{21}{37{,}000} = \frac{45{,}679}{37{,}000}.$$

56. $5.\overline{71358} = 5 + \dfrac{71{,}358}{10^5} + \dfrac{71{,}358}{10^{10}} + \cdots$. Now $\dfrac{71{,}358}{10^5} + \dfrac{71{,}358}{10^{10}} + \cdots$ is a geometric series with $a = \dfrac{71{,}358}{10^5}$ and $r = \dfrac{1}{10^5}$. It converges to $\dfrac{a}{1 - r} = \dfrac{71{,}358/10^5}{1 - 1/10^5} = \dfrac{71{,}358/10^5}{99{,}999/10^5} = \dfrac{71{,}358}{99{,}999} = \dfrac{23{,}786}{33{,}333}$. Thus,

$$5.\overline{71358} = 5 + \frac{23{,}786}{33{,}333} = \frac{166{,}665}{33{,}333} + \frac{23{,}786}{33{,}333} = \frac{190{,}451}{33{,}333}.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

57. $\sum_{n=1}^{\infty}(-5)^n x^n = \sum_{n=1}^{\infty}(-5x)^n$ is a geometric series with $r=-5x$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $|-5x|<1$ $\Leftrightarrow$ $|x|<\frac{1}{5}$, that is, $-\frac{1}{5}<x<\frac{1}{5}$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{-5x}{1-(-5x)}=\dfrac{-5x}{1+5x}$.

58. $\sum_{n=1}^{\infty}(x+2)^n$ is a geometric series with $r=x+2$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $|x+2|<1$ $\Leftrightarrow$ $-1<x+2<1$ $\Leftrightarrow$ $-3<x<-1$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{x+2}{1-(x+2)}=\dfrac{x+2}{-x-1}$.

59. $\sum_{n=0}^{\infty}\dfrac{(x-2)^n}{3^n}=\sum_{n=0}^{\infty}\left(\dfrac{x-2}{3}\right)^n$ is a geometric series with $r=\dfrac{x-2}{3}$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $\left|\dfrac{x-2}{3}\right|<1$ $\Leftrightarrow$ $-1<\dfrac{x-2}{3}<1$ $\Leftrightarrow$ $-3<x-2<3$ $\Leftrightarrow$ $-1<x<5$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-\frac{x-2}{3}}=\dfrac{1}{\frac{3-(x-2)}{3}}=\dfrac{3}{5-x}$.

60. $\sum_{n-0}^{\infty}(-4)^n(x-5)^n=\sum_{n=0}^{\infty}[-4(x-5)]^n$ is a geometric series with $r=-4(x-5)$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $|-4(x-5)|<1$ $\Leftrightarrow$ $|x-5|<\frac{1}{4}$ $\Leftrightarrow$ $-\frac{1}{4}<x-5<\frac{1}{4}$ $\Leftrightarrow$ $\frac{19}{4}<x<\frac{21}{4}$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-[-4(x-5)]}=\dfrac{1}{4x-19}$.

61. $\sum_{n=0}^{\infty}\dfrac{2^n}{x^n}=\sum_{n=0}^{\infty}\left(\dfrac{2}{x}\right)^n$ is a geometric series with $r=\dfrac{2}{x}$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $\left|\dfrac{2}{x}\right|<1$ $\Leftrightarrow$ $2<|x|$ $\Leftrightarrow$ $x>2$ or $x<-2$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-2/x}=\dfrac{x}{x-2}$.

62. $\sum_{n=0}^{\infty}\dfrac{\sin^n x}{3^n}=\sum_{n=0}^{\infty}\left(\dfrac{\sin x}{3}\right)^n$ is a geometric series with $r=\dfrac{\sin x}{3}$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $\left|\dfrac{\sin x}{3}\right|<1$ $\Leftrightarrow$ $|\sin x|<3$, which is true for all x. Thus, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-(\sin x)/3}=\dfrac{3}{3-\sin x}$.

63. $\sum_{n=0}^{\infty}e^{nx}=\sum_{n=0}^{\infty}(e^x)^n$ is a geometric series with $r=e^x$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $|e^x|<1$ $\Leftrightarrow$ $-1<e^x<1$ $\Leftrightarrow$ $0<e^x<1$ $\Leftrightarrow$ $x<0$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-e^x}$.

64. Because $\dfrac{1}{n}\to 0$ and ln is continuous, we have $\lim_{n\to\infty}\ln\left(1+\dfrac{1}{n}\right)=\ln 1=0$.

We now show that the series $\sum_{n=1}^{\infty}\ln\left(1+\dfrac{1}{n}\right)=\sum_{n=1}^{\infty}\ln\left(\dfrac{n+1}{n}\right)=\sum_{n=1}^{\infty}[\ln(n+1)-\ln n]$ diverges.

$s_n=(\ln 2-\ln 1)+(\ln 3-\ln 2)+\cdots+(\ln(n+1)-\ln n)=\ln(n+1)-\ln 1=\ln(n+1)$.

As $n\to\infty$, $s_n=\ln(n+1)\to\infty$, so the series diverges.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

65. After defining f, We use `convert(f,parfrac);` in Maple, `Apart` in Mathematica, or `Expand Rational` and `Simplify` in Derive to find that the general term is $\dfrac{3n^2+3n+1}{(n^2+n)^3} = \dfrac{1}{n^3} - \dfrac{1}{(n+1)^3}$. So the nth partial sum is

$$s_n = \sum_{k=1}^{n}\left(\frac{1}{k^3} - \frac{1}{(k+1)^3}\right) = \left(1 - \frac{1}{2^3}\right) + \left(\frac{1}{2^3} - \frac{1}{3^3}\right) + \cdots + \left(\frac{1}{n^3} - \frac{1}{(n+1)^3}\right) = 1 - \frac{1}{(n+1)^3}$$

The series converges to $\lim_{n\to\infty} s_n = 1$. This can be confirmed by directly computing the sum using `sum(f,n=1..infinity);` (in Maple), `Sum[f,{n,1,Infinity}]` (in Mathematica), or `Calculus Sum` (from 1 to ∞) and `Simplify` (in Derive).

66. See Exercise 65 for specific CAS commands.

$\dfrac{1}{n^5-5n^3+4n} = \dfrac{1}{24(n-2)} + \dfrac{1}{24(n+2)} - \dfrac{1}{6(n-1)} - \dfrac{1}{6(n+1)} + \dfrac{1}{4n}$. So the nth partial sum is

$$\begin{aligned} s_n &= \frac{1}{24}\sum_{k=3}^{n}\left(\frac{1}{k-2} - \frac{4}{k-1} + \frac{6}{k} - \frac{4}{k+1} + \frac{1}{k+2}\right) \\ &= \frac{1}{24}\left[\left(\frac{1}{1} - \frac{4}{2} + \frac{6}{3} - \frac{4}{4} + \frac{1}{5}\right) + \cdots + \left(\frac{1}{n-2} - \frac{4}{n-1} + \frac{6}{n} - \frac{4}{n+1} + \frac{1}{n+2}\right)\right] \end{aligned}$$

The terms with denominator 5 or greater cancel, except for a few terms with n in the denominator. So as $n \to \infty$,

$$s_n \to \frac{1}{24}\left(\frac{1}{1} - \frac{3}{2} + \frac{3}{3} - \frac{1}{4}\right) = \frac{1}{24}\left(\frac{1}{4}\right) = \frac{1}{96}.$$

67. For $n = 1$, $a_1 = 0$ since $s_1 = 0$. For $n > 1$,

$$a_n = s_n - s_{n-1} = \frac{n-1}{n+1} - \frac{(n-1)-1}{(n-1)+1} = \frac{(n-1)n - (n+1)(n-2)}{(n+1)n} = \frac{2}{n(n+1)}$$

Also, $\sum_{n=1}^{\infty} a_n = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \dfrac{1-1/n}{1+1/n} = 1$.

68. $a_1 = s_1 = 3 - \frac{1}{2} = \frac{5}{2}$. For $n \neq 1$,

$$a_n = s_n - s_{n-1} = \left(3 - n2^{-n}\right) - \left[3 - (n-1)2^{-(n-1)}\right] = -\frac{n}{2^n} + \frac{n-1}{2^{n-1}}\cdot\frac{2}{2} = \frac{2(n-1)}{2^n} - \frac{n}{2^n} = \frac{n-2}{2^n}$$

Also, $\sum_{n=1}^{\infty} a_n = \lim_{n\to\infty} s_n = \lim_{n\to\infty}\left(3 - \dfrac{n}{2^n}\right) = 3$ because $\lim_{x\to\infty} \dfrac{x}{2^x} \overset{\text{H}}{=} \lim_{x\to\infty} \dfrac{1}{2^x \ln 2} = 0$.

69. (a) The quantity of the drug in the body after the first tablet is 100 mg. After the second tablet, there is 100 mg plus 20% of the first 100-mg tablet; that is, $100 + 0.20(100) = 120$ mg. After the third tablet, the quantity is $100 + 0.20(120)$ or, equivalently, $100 + 100(0.20) + 100(0.20)^2$. Either expression gives us 124 mg.

(b) From part (a), we see that $Q_{n+1} = 100 + 0.20\,Q_n$.

(c) $Q_n = 100 + 100(0.20)^1 + 100(0.20)^2 + \cdots + 100(0.20)^{n-1}$

$= \sum_{i=1}^{n} 100(0.20)^{i-1}$ [geometric with $a = 100$ and $r = 0.20$].

The quantity of the antibiotic that remains in the body in the long run is $\lim_{n\to\infty} Q_n = \dfrac{100}{1-0.20} = \dfrac{100}{4/5} = 125$ mg.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

70. (a) The concentration of the drug after the first injection is 1.5 mg/L. "Reduced by 90%" is the same as 10% remains, so the concentration after the second injection is $1.5 + 0.10(1.5) = 1.65$ mg/L. The concentration after the third injection is $1.5 + 0.10(1.65)$, or, equivalently, $1.5 + 1.5(0.10) + 1.5(0.10)^2$. Either expression gives us 1.665 mg/L.

(b) $C_n = 1.5 + 1.5(0.10)^1 + 1.5(0.10)^2 + \cdots + 1.5(0.10)^{n-1}$

$= \sum_{i=1}^{n} 1.5(0.10)^{i-1}$ [geometric with $a = 1.5$ and $r = 0.10$].

By (3), $C_n = \dfrac{1.5[1-(0.10)^n]}{1-0.10} = \dfrac{1.5}{0.9}[1-(0.10)^n] = \dfrac{5}{3}[1-(0.10)^n]$ mg/L.

(c) The limiting value of the concentration is $\lim_{n\to\infty} C_n = \lim_{n\to\infty} \frac{5}{3}[1-(0.10)^n] = \frac{5}{3}(1-0) = \frac{5}{3}$ mg/L.

71. (a) The quantity of the drug in the body after the first tablet is 150 mg. After the second tablet, there is 150 mg plus 5% of the first 150- mg tablet, that is, $[150 + 150(0.05)]$ mg. After the third tablet, the quantity is $[150 + 150(0.05) + 150(0.05)^2] = 157.875$ mg. After n tablets, the quantity (in mg) is $150 + 150(0.05) + \cdots + 150(0.05)^{n-1}$. We can use Formula 3 to write this as $\dfrac{150(1-0.05^n)}{1-0.05} = \dfrac{3000}{19}(1-0.05^n)$.

(b) The number of milligrams remaining in the body in the long run is $\lim_{n\to\infty} \left[\frac{3000}{19}(1-0.05^n)\right] = \frac{3000}{19}(1-0) \approx 157.895$, only 0.02 mg more than the amount after 3 tablets.

72. (a) The residual concentration just before the second injection is De^{-aT}; before the third, $De^{-aT} + De^{-a2T}$; before the $(n+1)$st, $De^{-aT} + De^{-a2T} + \cdots + De^{-anT}$. This sum is equal to $\dfrac{De^{-aT}\left(1-e^{-anT}\right)}{1-e^{-aT}}$ [Formula 3].

(b) The limiting pre-injection concentration is $\lim_{n\to\infty} \dfrac{De^{-aT}\left(1-e^{-anT}\right)}{1-e^{-aT}} = \dfrac{De^{-aT}(1-0)}{1-e^{-aT}} \cdot \dfrac{e^{aT}}{e^{aT}} = \dfrac{D}{e^{aT}-1}$.

(c) $\dfrac{D}{e^{aT}-1} \geq C \quad \Rightarrow \quad D \geq C\left(e^{aT}-1\right)$, so the minimal dosage is $D = C\left(e^{aT}-1\right)$.

73. (a) The first step in the chain occurs when the local government spends D dollars. The people who receive it spend a fraction c of those D dollars, that is, Dc dollars. Those who receive the Dc dollars spend a fraction c of it, that is, Dc^2 dollars. Continuing in this way, we see that the total spending after n transactions is

$S_n = D + Dc + Dc^2 + \cdots + Dc^{n-1} = \dfrac{D(1-c^n)}{1-c}$ by (3).

(b) $\lim_{n\to\infty} S_n = \lim_{n\to\infty} \dfrac{D(1-c^n)}{1-c} = \dfrac{D}{1-c} \lim_{n\to\infty} (1-c^n) = \dfrac{D}{1-c} \quad \left[\text{since } 0 < c < 1 \quad \Rightarrow \quad \lim_{n\to\infty} c^n = 0\right]$

$= \dfrac{D}{s}$ [since $c + s = 1$] $= kD$ [since $k = 1/s$]

If $c = 0.8$, then $s = 1 - c = 0.2$ and the multiplier is $k = 1/s = 5$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

74. (a) Initially, the ball falls a distance H, then rebounds a distance rH, falls rH, rebounds r^2H, falls r^2H, etc. The total distance it travels is

$$\begin{aligned} H + 2rH + 2r^2H + 2r^3H + \cdots &= H\left(1 + 2r + 2r^2 + 2r^3 + \cdots\right) = H\left[1 + 2r\left(1 + r + r^2 + \cdots\right)\right] \\ &= H\left[1 + 2r\left(\frac{1}{1-r}\right)\right] = H\left(\frac{1+r}{1-r}\right) \text{ meters} \end{aligned}$$

(b) We know that the ball falls $\frac{1}{2}gt^2$ meters in t seconds, where $g = 9.8\ \text{m/s}^2$ is the gravitational acceleration. Thus, the ball falls h meters in $t = \sqrt{2h/g}$ seconds. The total travel time in seconds is

$$\begin{aligned} \sqrt{\frac{2H}{g}} + 2\sqrt{\frac{2H}{g}r} + 2\sqrt{\frac{2H}{g}r^2} + 2\sqrt{\frac{2H}{g}r^3} + \cdots &= \sqrt{\frac{2H}{g}}\left[1 + 2\sqrt{r} + 2\sqrt{r}^{\,2} + 2\sqrt{r}^{\,3} + \cdots\right] \\ &= \sqrt{\frac{2H}{g}}\left(1 + 2\sqrt{r}\left[1 + \sqrt{r} + \sqrt{r}^{\,2} + \cdots\right]\right) \\ &= \sqrt{\frac{2H}{g}}\left[1 + 2\sqrt{r}\left(\frac{1}{1-\sqrt{r}}\right)\right] = \sqrt{\frac{2H}{g}}\,\frac{1+\sqrt{r}}{1-\sqrt{r}} \end{aligned}$$

(c) It will help to make a chart of the time for each descent and each rebound of the ball, together with the velocity just before and just after each bounce. Recall that the time in seconds needed to fall h meters is $\sqrt{2h/g}$. The ball hits the ground with velocity $-g\sqrt{2h/g} = -\sqrt{2hg}$ (taking the upward direction to be positive) and rebounds with velocity $kg\sqrt{2h/g} = k\sqrt{2hg}$, taking time $k\sqrt{2h/g}$ to reach the top of its bounce, where its velocity is 0. At that point, its height is k^2h. All these results follow from the formulas for vertical motion with gravitational acceleration $-g$:

$$\frac{d^2y}{dt^2} = -g \quad \Rightarrow \quad v = \frac{dy}{dt} = v_0 - gt \quad \Rightarrow \quad y = y_0 + v_0t - \tfrac{1}{2}gt^2.$$

number of descent	time of descent	speed before bounce	speed after bounce	time of ascent	peak height
1	$\sqrt{2H/g}$	$\sqrt{2Hg}$	$k\sqrt{2Hg}$	$k\sqrt{2H/g}$	k^2H
2	$\sqrt{2k^2H/g}$	$\sqrt{2k^2Hg}$	$k\sqrt{2k^2Hg}$	$k\sqrt{2k^2H/g}$	k^4H
3	$\sqrt{2k^4H/g}$	$\sqrt{2k^4Hg}$	$k\sqrt{2k^4Hg}$	$k\sqrt{2k^4H/g}$	k^6H
...	...	...	...	...	...

The total travel time in seconds is

$$\begin{aligned} \sqrt{\frac{2H}{g}} + k\sqrt{\frac{2H}{g}} + k\sqrt{\frac{2H}{g}} + k^2\sqrt{\frac{2H}{g}} + k^2\sqrt{\frac{2H}{g}} + \cdots &= \sqrt{\frac{2H}{g}}\left(1 + 2k + 2k^2 + 2k^3 + \cdots\right) \\ &= \sqrt{\frac{2H}{g}}\left[1 + 2k(1 + k + k^2 + \cdots)\right] \\ &= \sqrt{\frac{2H}{g}}\left[1 + 2k\left(\frac{1}{1-k}\right)\right] = \sqrt{\frac{2H}{g}}\,\frac{1+k}{1-k} \end{aligned}$$

Another method: We could use part (b). At the top of the bounce, the height is $k^2h = rh$, so $\sqrt{r} = k$ and the result follows from part (b).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

75. $\sum_{n=2}^{\infty}(1+c)^{-n}$ is a geometric series with $a=(1+c)^{-2}$ and $r=(1+c)^{-1}$, so the series converges when

$\left|(1+c)^{-1}\right|<1 \;\Leftrightarrow\; |1+c|>1 \;\Leftrightarrow\; 1+c>1$ or $1+c<-1 \;\Leftrightarrow\; c>0$ or $c<-2$. We calculate the sum of the series and set it equal to 2: $\dfrac{(1+c)^{-2}}{1-(1+c)^{-1}}=2 \;\Leftrightarrow\; \left(\dfrac{1}{1+c}\right)^2=2-2\left(\dfrac{1}{1+c}\right) \;\Leftrightarrow\; 1=2(1+c)^2-2(1+c) \;\Leftrightarrow$

$2c^2+2c-1=0 \;\Leftrightarrow\; c=\frac{-2\pm\sqrt{12}}{4}=\frac{\pm\sqrt{3}-1}{2}$. However, the negative root is inadmissible because $-2<\frac{-\sqrt{3}-1}{2}<0$.

So $c=\frac{\sqrt{3}-1}{2}$.

76. $\sum_{n=0}^{\infty}e^{nc}=\sum_{n=0}^{\infty}(e^c)^n$ is a geometric series with $a=(e^c)^0=1$ and $r=e^c$. If $e^c<1$, it has sum $\dfrac{1}{1-e^c}$, so $\dfrac{1}{1-e^c}=10 \;\Rightarrow$

$\frac{1}{10}=1-e^c \;\Rightarrow\; e^c=\frac{9}{10} \;\Rightarrow\; c=\ln\frac{9}{10}$.

77. $e^{s_n}=e^{1+\frac{1}{2}+\frac{1}{3}+\cdots+\frac{1}{n}}=e^1e^{1/2}e^{1/3}\cdots e^{1/n}>(1+1)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)\cdots\left(1+\frac{1}{n}\right) \qquad [e^x>1+x]$

$=\dfrac{2}{1}\dfrac{3}{2}\dfrac{4}{3}\cdots\dfrac{n+1}{n}=n+1$

Thus, $e^{s_n}>n+1$ and $\lim_{n\to\infty}e^{s_n}=\infty$. Since $\{s_n\}$ is increasing, $\lim_{n\to\infty}s_n=\infty$, implying that the harmonic series is divergent.

78. The area between $y=x^{n-1}$ and $y=x^n$ for $0\le x\le 1$ is

$$\int_0^1\left(x^{n-1}-x^n\right)dx=\left[\frac{x^n}{n}-\frac{x^{n+1}}{n+1}\right]_0^1=\frac{1}{n}-\frac{1}{n+1}$$
$$=\frac{(n+1)-n}{n(n+1)}=\frac{1}{n(n+1)}$$

We can see from the diagram that as $n\to\infty$, the sum of the areas between the successive curves approaches the area of the unit square, that is, 1. So $\sum_{n=1}^{\infty}\dfrac{1}{n(n+1)}=1$.



79. Let d_n be the diameter of C_n. We draw lines from the centers of the C_i to the center of D (or C), and using the Pythagorean Theorem, we can write

$1^2+\left(1-\frac{1}{2}d_1\right)^2=\left(1+\frac{1}{2}d_1\right)^2 \;\Leftrightarrow$

$1=\left(1+\frac{1}{2}d_1\right)^2-\left(1-\frac{1}{2}d_1\right)^2=2d_1$ [difference of squares] $\;\Rightarrow\; d_1=\frac{1}{2}$.

Similarly,

$1=\left(1+\frac{1}{2}d_2\right)^2-\left(1-d_1-\frac{1}{2}d_2\right)^2=2d_2+2d_1-d_1^2-d_1d_2$

$=(2-d_1)(d_1+d_2) \;\Leftrightarrow$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$d_2 = \dfrac{1}{2-d_1} - d_1 = \dfrac{(1-d_1)^2}{2-d_1}$, $1 = \left(1+\frac{1}{2}d_3\right)^2 - \left(1-d_1-d_2-\frac{1}{2}d_3\right)^2 \quad \Leftrightarrow \quad d_3 = \dfrac{[1-(d_1+d_2)]^2}{2-(d_1+d_2)}$, and in general, $d_{n+1} = \dfrac{\left(1-\sum_{i=1}^{n} d_i\right)^2}{2-\sum_{i=1}^{n} d_i}$. If we actually calculate d_2 and d_3 from the formulas above, we find that they are $\dfrac{1}{6} = \dfrac{1}{2\cdot 3}$ and $\dfrac{1}{12} = \dfrac{1}{3\cdot 4}$ respectively, so we suspect that in general, $d_n = \dfrac{1}{n(n+1)}$. To prove this, we use induction: Assume that for all $k \le n$, $d_k = \dfrac{1}{k(k+1)} = \dfrac{1}{k} - \dfrac{1}{k+1}$. Then $\sum_{i=1}^{n} d_i = 1 - \dfrac{1}{n+1} = \dfrac{n}{n+1}$ [telescoping sum]. Substituting this into our formula for d_{n+1}, we get $d_{n+1} = \dfrac{\left[1-\dfrac{n}{n+1}\right]^2}{2-\left(\dfrac{n}{n+1}\right)} = \dfrac{\dfrac{1}{(n+1)^2}}{\dfrac{n+2}{n+1}} = \dfrac{1}{(n+1)(n+2)}$, and the induction is complete.

Now, we observe that the partial sums $\sum_{i=1}^{n} d_i$ of the diameters of the circles approach 1 as $n \to \infty$; that is, $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \dfrac{1}{n(n+1)} = 1$, which is what we wanted to prove.

80. $|CD| = b\sin\theta$, $|DE| = |CD|\sin\theta = b\sin^2\theta$, $|EF| = |DE|\sin\theta = b\sin^3\theta$, ... Therefore, $|CD| + |DE| + |EF| + |FG| + \cdots = b\sum_{n=1}^{\infty} \sin^n\theta = b\left(\dfrac{\sin\theta}{1-\sin\theta}\right)$ since this is a geometric series with $r = \sin\theta$ and $|\sin\theta| < 1 \quad \left[\text{because } 0 < \theta < \frac{\pi}{2}\right]$.

81. The series $1-1+1-1+1-1+\cdots$ diverges (geometric series with $r = -1$) so we cannot say that $0 = 1-1+1-1+1-1+\cdots$.

82. If $\sum_{n=1}^{\infty} a_n$ is convergent, then $\lim_{n\to\infty} a_n = 0$ by Theorem 6, so $\lim_{n\to\infty} \dfrac{1}{a_n} \neq 0$, and so $\sum_{n=1}^{\infty} \dfrac{1}{a_n}$ is divergent by the Test for Divergence.

83. $\sum_{n=1}^{\infty} ca_n = \lim_{n\to\infty} \sum_{i=1}^{n} ca_i = \lim_{n\to\infty} c\sum_{i=1}^{n} a_i = c\lim_{n\to\infty} \sum_{i=1}^{n} a_i = c\sum_{n=1}^{\infty} a_n$, which exists by hypothesis.

84. If $\sum ca_n$ were convergent, then $\sum (1/c)(ca_n) = \sum a_n$ would be also, by Theorem 8(i). But this is not the case, so $\sum ca_n$ must diverge.

85. Suppose on the contrary that $\sum(a_n + b_n)$ converges. Then $\sum(a_n + b_n)$ and $\sum a_n$ are convergent series. So by Theorem 8(iii), $\sum[(a_n + b_n) - a_n]$ would also be convergent. But $\sum[(a_n + b_n) - a_n] = \sum b_n$, a contradiction, since $\sum b_n$ is given to be divergent.

86. No. For example, take $\sum a_n = \sum n$ and $\sum b_n = \sum(-n)$, which both diverge, yet $\sum(a_n + b_n) = \sum 0$, which converges with sum 0.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

87. The partial sums $\{s_n\}$ form an increasing sequence, since $s_n - s_{n-1} = a_n > 0$ for all n. Also, the sequence $\{s_n\}$ is bounded since $s_n \le 1000$ for all n. So by the Monotonic Sequence Theorem, the sequence of partial sums converges, that is, the series $\sum a_n$ is convergent.

88. (a) $\text{RHS} = \dfrac{1}{f_{n-1}f_n} - \dfrac{1}{f_n f_{n+1}} = \dfrac{f_n f_{n+1} - f_n f_{n-1}}{f_n^2 f_{n-1} f_{n+1}} = \dfrac{f_{n+1} - f_{n-1}}{f_n f_{n-1} f_{n+1}} = \dfrac{(f_{n-1} + f_n) - f_{n-1}}{f_n f_{n-1} f_{n+1}} = \dfrac{1}{f_{n-1} f_{n+1}} = \text{LHS}$

(b) $$\begin{aligned}\sum_{n=2}^{\infty} \frac{1}{f_{n-1}f_{n+1}} &= \sum_{n=2}^{\infty}\left(\frac{1}{f_{n-1}f_n} - \frac{1}{f_n f_{n+1}}\right) \quad \text{[from part (a)]} \\ &= \lim_{n\to\infty}\left[\left(\frac{1}{f_1 f_2} - \frac{1}{f_2 f_3}\right) + \left(\frac{1}{f_2 f_3} - \frac{1}{f_3 f_4}\right) + \left(\frac{1}{f_3 f_4} - \frac{1}{f_4 f_5}\right) + \cdots + \left(\frac{1}{f_{n-1}f_n} - \frac{1}{f_n f_{n+1}}\right)\right] \\ &= \lim_{n\to\infty}\left(\frac{1}{f_1 f_2} - \frac{1}{f_n f_{n+1}}\right) = \frac{1}{f_1 f_2} - 0 = \frac{1}{1\cdot 1} = 1 \quad \text{because } f_n \to \infty \text{ as } n \to \infty.\end{aligned}$$

(c) $$\begin{aligned}\sum_{n=2}^{\infty} \frac{f_n}{f_{n-1}f_{n+1}} &= \sum_{n=2}^{\infty}\left(\frac{f_n}{f_{n-1}f_n} - \frac{f_n}{f_n f_{n+1}}\right) \quad \text{[as above]} \\ &= \sum_{n=2}^{\infty}\left(\frac{1}{f_{n-1}} - \frac{1}{f_{n+1}}\right) \\ &= \lim_{n\to\infty}\left[\left(\frac{1}{f_1} - \frac{1}{f_3}\right) + \left(\frac{1}{f_2} - \frac{1}{f_4}\right) + \left(\frac{1}{f_3} - \frac{1}{f_5}\right) + \left(\frac{1}{f_4} - \frac{1}{f_6}\right) + \cdots + \left(\frac{1}{f_{n-1}} - \frac{1}{f_{n+1}}\right)\right] \\ &= \lim_{n\to\infty}\left(\frac{1}{f_1} + \frac{1}{f_2} - \frac{1}{f_n} - \frac{1}{f_{n+1}}\right) = 1 + 1 - 0 - 0 = 2 \quad \text{because } f_n \to \infty \text{ as } n \to \infty.\end{aligned}$$

89. (a) At the first step, only the interval $\left(\frac{1}{3}, \frac{2}{3}\right)$ (length $\frac{1}{3}$) is removed. At the second step, we remove the intervals $\left(\frac{1}{9}, \frac{2}{9}\right)$ and $\left(\frac{7}{9}, \frac{8}{9}\right)$, which have a total length of $2 \cdot \left(\frac{1}{3}\right)^2$. At the third step, we remove 2^2 intervals, each of length $\left(\frac{1}{3}\right)^3$. In general, at the nth step we remove 2^{n-1} intervals, each of length $\left(\frac{1}{3}\right)^n$, for a length of $2^{n-1} \cdot \left(\frac{1}{3}\right)^n = \frac{1}{3}\left(\frac{2}{3}\right)^{n-1}$. Thus, the total length of all removed intervals is $\sum_{n=1}^{\infty} \frac{1}{3}\left(\frac{2}{3}\right)^{n-1} = \frac{1/3}{1-2/3} = 1$ [geometric series with $a = \frac{1}{3}$ and $r = \frac{2}{3}$]. Notice that at the nth step, the leftmost interval that is removed is $\left(\left(\frac{1}{3}\right)^n, \left(\frac{2}{3}\right)^n\right)$, so we never remove 0, and 0 is in the Cantor set. Also, the rightmost interval removed is $\left(1 - \left(\frac{2}{3}\right)^n, 1 - \left(\frac{1}{3}\right)^n\right)$, so 1 is never removed. Some other numbers in the Cantor set are $\frac{1}{3}$, $\frac{2}{3}$, $\frac{1}{9}$, $\frac{2}{9}$, $\frac{7}{9}$, and $\frac{8}{9}$.

(b) The area removed at the first step is $\frac{1}{9}$; at the second step, $8 \cdot \left(\frac{1}{9}\right)^2$; at the third step, $(8)^2 \cdot \left(\frac{1}{9}\right)^3$. In general, the area removed at the nth step is $(8)^{n-1}\left(\frac{1}{9}\right)^n = \frac{1}{9}\left(\frac{8}{9}\right)^{n-1}$, so the total area of all removed squares is

$$\sum_{n=1}^{\infty} \frac{1}{9}\left(\frac{8}{9}\right)^{n-1} = \frac{1/9}{1 - 8/9} = 1.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

90. (a)

a_1	1	2	4	1	1	1000
a_2	2	3	1	4	1000	1
a_3	1.5	2.5	2.5	2.5	500.5	500.5
a_4	1.75	2.75	1.75	3.25	750.25	250.75
a_5	1.625	2.625	2.125	2.875	625.375	375.625
a_6	1.6875	2.6875	1.9375	3.0625	687.813	313.188
a_7	1.65625	2.65625	2.03125	2.96875	656.594	344.406
a_8	1.67188	2.67188	1.98438	3.01563	672.203	328.797
a_9	1.66406	2.66406	2.00781	2.99219	664.398	336.602
a_{10}	1.66797	2.66797	1.99609	3.00391	668.301	332.699
a_{11}	1.66602	2.66602	2.00195	2.99805	666.350	334.650
a_{12}	1.66699	2.66699	1.99902	3.00098	667.325	333.675

The limits seem to be $\frac{5}{3}, \frac{8}{3}$, 2, 3, 667, and 334. Note that the limits appear to be "weighted" more toward a_2. In general, we guess that the limit is $\dfrac{a_1 + 2a_2}{3}$.

(b) $a_{n+1} - a_n = \frac{1}{2}(a_n + a_{n-1}) - a_n = -\frac{1}{2}(a_n - a_{n-1}) = -\frac{1}{2}\left[\frac{1}{2}(a_{n-1} + a_{n-2}) - a_{n-1}\right]$

$= -\frac{1}{2}\left[-\frac{1}{2}(a_{n-1} - a_{n-2})\right] = \cdots = \left(-\frac{1}{2}\right)^{n-1}(a_2 - a_1)$

Note that we have used the formula $a_k = \frac{1}{2}(a_{k-1} + a_{k-2})$ a total of $n - 1$ times in this calculation, once for each k between 3 and $n + 1$. Now we can write

$$a_n = a_1 + (a_2 - a_1) + (a_3 - a_2) + \cdots + (a_{n-1} - a_{n-2}) + (a_n - a_{n-1})$$

$$= a_1 + \sum_{k=1}^{n-1}(a_{k+1} - a_k) = a_1 + \sum_{k=1}^{n-1}\left(-\tfrac{1}{2}\right)^{k-1}(a_2 - a_1)$$

and so

$$\lim_{n\to\infty} a_n = a_1 + (a_2 - a_1)\sum_{k=1}^{\infty}\left(-\tfrac{1}{2}\right)^{k-1} = a_1 + (a_2 - a_1)\left[\frac{1}{1 - (-1/2)}\right] = a_1 + \tfrac{2}{3}(a_2 - a_1) = \frac{a_1 + 2a_2}{3}.$$

91. (a) For $\displaystyle\sum_{n=1}^{\infty}\frac{n}{(n+1)!}$, $s_1 = \dfrac{1}{1\cdot 2} = \dfrac{1}{2}$, $s_2 = \dfrac{1}{2} + \dfrac{2}{1\cdot 2\cdot 3} = \dfrac{5}{6}$, $s_3 = \dfrac{5}{6} + \dfrac{3}{1\cdot 2\cdot 3\cdot 4} = \dfrac{23}{24}$,

$s_4 = \dfrac{23}{24} + \dfrac{4}{1\cdot 2\cdot 3\cdot 4\cdot 5} = \dfrac{119}{120}$. The denominators are $(n+1)!$, so a guess would be $s_n = \dfrac{(n+1)! - 1}{(n+1)!}$.

(b) For $n = 1$, $s_1 = \dfrac{1}{2} = \dfrac{2! - 1}{2!}$, so the formula holds for $n = 1$. Assume $s_k = \dfrac{(k+1)! - 1}{(k+1)!}$. Then

$$s_{k+1} = \frac{(k+1)! - 1}{(k+1)!} + \frac{k+1}{(k+2)!} = \frac{(k+1)! - 1}{(k+1)!} + \frac{k+1}{(k+1)!(k+2)} = \frac{(k+2)! - (k+2) + k + 1}{(k+2)!}$$

$$= \frac{(k+2)! - 1}{(k+2)!}$$

Thus, the formula is true for $n = k + 1$. So by induction, the guess is correct.

(c) $\displaystyle\lim_{n\to\infty} s_n = \lim_{n\to\infty}\frac{(n+1)! - 1}{(n+1)!} = \lim_{n\to\infty}\left[1 - \frac{1}{(n+1)!}\right] = 1$ and so $\displaystyle\sum_{n=1}^{\infty}\frac{n}{(n+1)!} = 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

92.



Let $r_1 =$ radius of the large circle, $r_2 =$ radius of next circle, and so on. From the figure we have $\angle BAC = 60°$ and $\cos 60° = r_1/|AB|$, so $|AB| = 2r_1$ and $|DB| = 2r_2$. Therefore, $2r_1 = r_1 + r_2 + 2r_2 \;\Rightarrow$ $r_1 = 3r_2$. In general, we have $r_{n+1} = \frac{1}{3}r_n$, so the total area is

$$A = \pi r_1^2 + 3\pi r_2^2 + 3\pi r_3^2 + \cdots = \pi r_1^2 + 3\pi r_2^2\left(1 + \frac{1}{3^2} + \frac{1}{3^4} + \frac{1}{3^6} + \cdots\right)$$

$$= \pi r_1^2 + 3\pi r_2^2 \cdot \frac{1}{1 - 1/9} = \pi r_1^2 + \tfrac{27}{8}\pi r_2^2$$

Since the sides of the triangle have length 1, $|BC| = \frac{1}{2}$ and $\tan 30° = \dfrac{r_1}{1/2}$. Thus, $r_1 = \dfrac{\tan 30°}{2} = \dfrac{1}{2\sqrt{3}} \;\Rightarrow\; r_2 = \dfrac{1}{6\sqrt{3}}$, so $A = \pi\left(\dfrac{1}{2\sqrt{3}}\right)^2 + \dfrac{27\pi}{8}\left(\dfrac{1}{6\sqrt{3}}\right)^2 = \dfrac{\pi}{12} + \dfrac{\pi}{32} = \dfrac{11\pi}{96}$. The area of the triangle is $\dfrac{\sqrt{3}}{4}$, so the circles occupy about 83.1% of the area of the triangle.

11.3 The Integral Test and Estimates of Sums

1. The picture shows that $a_2 = \dfrac{1}{2^{1.3}} < \displaystyle\int_1^2 \frac{1}{x^{1.3}}\,dx$, $a_3 = \dfrac{1}{3^{1.3}} < \displaystyle\int_2^3 \frac{1}{x^{1.3}}\,dx$, and so on, so $\displaystyle\sum_{n=2}^{\infty} \frac{1}{n^{1.3}} < \int_1^{\infty} \frac{1}{x^{1.3}}\,dx$. The integral converges by (7.8.2) with $p = 1.3 > 1$, so the series converges.



2. From the first figure, we see that $\int_1^6 f(x)\,dx < \displaystyle\sum_{i=1}^{5} a_i$. From the second figure, we see that $\displaystyle\sum_{i=2}^{6} a_i < \int_1^6 f(x)\,dx$. Thus, we have $\displaystyle\sum_{i=2}^{6} a_i < \int_1^6 f(x)\,dx < \sum_{i=1}^{5} a_i$.





3. The function $f(x) = x^{-3}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^{\infty} x^{-3}\,dx = \lim_{t\to\infty}\int_1^t x^{-3}\,dx = \lim_{t\to\infty}\left[\frac{x^{-2}}{-2}\right]_1^t = \lim_{t\to\infty}\left(-\frac{1}{2t^2} + \frac{1}{2}\right) = \frac{1}{2}.$$

Since this improper integral is convergent, the series $\displaystyle\sum_{n=1}^{\infty} n^{-3}$ is also convergent by the Integral Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. The function $f(x) = x^{-0.3}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty x^{-0.3}\,dx = \lim_{t\to\infty}\int_1^t x^{-0.3}\,dx = \lim_{t\to\infty}\left[\frac{x^{0.7}}{0.7}\right]_1^t = \lim_{t\to\infty}\left(\frac{t^{0.7}}{0.7} - \frac{1}{0.7}\right) = \infty.$$

Since this improper integral is divergent, the series $\sum_{n=1}^{\infty} n^{-0.3}$ is also divergent by the Integral Test.

5. The function $f(x) = \dfrac{2}{5x-1}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty \frac{2}{5x-1}\,dx = \lim_{t\to\infty}\int_1^t \frac{2}{5x-1}\,dx = \lim_{t\to\infty}\left[\frac{2}{5}\ln(5x-1)\right]_1^t = \lim_{t\to\infty}\left[\frac{2}{5}\ln(5t-1) - \frac{2}{5}\ln 4\right] = \infty.$$

Since this improper integral is divergent, the series $\sum_{n=1}^{\infty} \dfrac{2}{5n-1}$ is also divergent by the Integral Test.

6. The function $f(x) = \dfrac{1}{(3x-1)^4}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty \frac{1}{(3x-1)^4}\,dx = \lim_{t\to\infty}\int_1^t (3x-1)^{-4}\,dx = \lim_{t\to\infty}\left[\frac{1}{(-3)3}(3x-1)^{-3}\right]_1^t = \lim_{t\to\infty}\left[-\frac{1}{9(3t-1)^3} + \frac{1}{9\cdot 2^3}\right] = \frac{1}{72}.$$

Since this improper integral is convergent, the series $\sum_{n=1}^{\infty} \dfrac{1}{(3n-1)^4}$ is also convergent by the Integral Test.

7. The function $f(x) = \dfrac{x}{x^2+1}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$\int_1^\infty \dfrac{x}{x^2+1}\,dx = \lim_{t\to\infty}\int_1^t \dfrac{x}{x^2+1}\,dx = \lim_{t\to\infty}\left[\dfrac{1}{2}\ln(x^2+1)\right]_1^t = \dfrac{1}{2}\lim_{t\to\infty}[\ln(t^2+1) - \ln 2] = \infty$. Since this improper integral is divergent, the series $\sum_{n=1}^{\infty} \dfrac{n}{n^2+1}$ is also divergent by the Integral Test.

8. The function $f(x) = x^2 e^{-x^3}$ is continuous, positive, and decreasing ($\star$) on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty x^2 e^{-x^3}\,dx = \lim_{t\to\infty}\int_1^t x^2 e^{-x^3}\,dx = \lim_{t\to\infty}\left[-\frac{1}{3}e^{-x^3}\right]_1^t = -\frac{1}{3}\lim_{t\to\infty}\left(e^{-t^3} - e^{-1}\right) = -\frac{1}{3}\left(0 - \frac{1}{e}\right) = \frac{1}{3e}.$$

Since this improper integral is convergent, the series $\sum_{n=1}^{\infty} n^2 e^{-n^3}$ is also convergent by the Integral Test.

($\star$): $f'(x) = x^2 e^{-x^3}(-3x^2) + e^{-x^3}(2x) = xe^{-x^3}(-3x^3+2) = \dfrac{x(2-3x^3)}{e^{x^3}} < 0$ for $x > 1$

9. $\sum_{n=1}^{\infty} \dfrac{1}{n^{\sqrt{2}}}$ is a p-series with $p = \sqrt{2} > 1$, so it converges by (1).

10. $\sum_{n=3}^{\infty} n^{-0.9999} = \sum_{n=3}^{\infty} \dfrac{1}{n^{0.9999}}$ is a p-series with $p = 0.9999 < 1$, so it diverges by (1). The fact that the series begins with $n = 3$ is irrelevant when determining convergence.

11. $1 + \dfrac{1}{8} + \dfrac{1}{27} + \dfrac{1}{64} + \dfrac{1}{125} + \cdots = \sum_{n=1}^{\infty} \dfrac{1}{n^3}$. This is a p-series with $p = 3 > 1$, so it converges by (1).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $\frac{1}{5}+\frac{1}{7}+\frac{1}{9}+\frac{1}{11}+\frac{1}{13}+\cdots=\sum_{n=1}^{\infty}\frac{1}{2n+3}$. The function $f(x)=\frac{1}{2x+3}$ is continuous, positive, and decreasing on $[1,\infty)$, so the Integral Test applies.

$\int_1^{\infty}\frac{1}{2x+3}\,dx=\lim_{t\to\infty}\int_1^{t}\frac{1}{2x+3}\,dx=\lim_{t\to\infty}\left[\frac{1}{2}\ln(2x+3)\right]_1^{t}=\lim_{t\to\infty}\left[\frac{1}{2}\ln(2t+3)-\frac{1}{2}\ln 5\right]=\infty$, so the series $\sum_{n=1}^{\infty}\frac{1}{2n+3}$ diverges.

13. $\frac{1}{3}+\frac{1}{7}+\frac{1}{11}+\frac{1}{15}+\frac{1}{19}+\cdots=\sum_{n=1}^{\infty}\frac{1}{4n-1}$. The function $f(x)=\frac{1}{4x-1}$ is continuous, positive, and decreasing on $[1,\infty)$, so the Integral Test applies.

$\int_1^{\infty}\frac{1}{4x-1}\,dx=\lim_{t\to\infty}\int_1^{t}\frac{1}{4x-1}\,dx=\lim_{t\to\infty}\left[\frac{1}{4}\ln(4x-1)\right]_1^{t}=\lim_{t\to\infty}\left[\frac{1}{4}\ln(4t-1)-\frac{1}{4}\ln 3\right]=\infty$, so the series $\sum_{n=1}^{\infty}\frac{1}{4n-1}$ diverges.

14. $1+\frac{1}{2\sqrt{2}}+\frac{1}{3\sqrt{3}}+\frac{1}{4\sqrt{4}}+\frac{1}{5\sqrt{5}}+\cdots=\sum_{n=1}^{\infty}\frac{1}{n\sqrt{n}}=\sum_{n=1}^{\infty}\frac{1}{n^{3/2}}$. This is a p-series with $p=\frac{3}{2}>1$, so it converges by (1).

15. $\sum_{n=1}^{\infty}\frac{\sqrt{n}+4}{n^2}=\sum_{n=1}^{\infty}\left(\frac{\sqrt{n}}{n^2}+\frac{4}{n^2}\right)=\sum_{n=1}^{\infty}\frac{1}{n^{3/2}}+\sum_{n=1}^{\infty}\frac{4}{n^2}$. $\sum_{n=1}^{\infty}\frac{1}{n^{3/2}}$ is a convergent p-series with $p=\frac{3}{2}>1$. $\sum_{n=1}^{\infty}\frac{4}{n^2}=4\sum_{n=1}^{\infty}\frac{1}{n^2}$ is a constant multiple of a convergent p-series with $p=2>1$, so it converges. The sum of two convergent series is convergent, so the original series is convergent.

16. The function $f(x)=\frac{\sqrt{x}}{1+x^{3/2}}$ is continuous and positive on $[1,\infty)$.

$f'(x)=\dfrac{(1+x^{3/2})\left(\frac{1}{2}x^{-1/2}\right)-x^{1/2}\left(\frac{3}{2}x^{1/2}\right)}{(1+x^{3/2})^2}=\dfrac{\frac{1}{2}x^{-1/2}+\frac{1}{2}x-\frac{3}{2}x}{(1+x^{3/2})^2}=\dfrac{1-2x^{3/2}}{2\sqrt{x}(1+x^{3/2})^2}<0$ for $x\geq 1$, so f is decreasing on $[1,\infty)$, and the Integral Test applies.

$$\int_1^{\infty}\frac{\sqrt{x}}{1+x^{3/2}}\,dx=\lim_{t\to\infty}\int_1^{t}\frac{\sqrt{x}}{1+x^{3/2}}\,dx=\lim_{t\to\infty}\left[\frac{2}{3}\ln(1+x^{3/2})\right]_1^{t}\qquad\left[\begin{array}{c}\text{substitution}\\ \text{with } u=1+x^{3/2}\end{array}\right]$$
$$=\lim_{t\to\infty}\left[\frac{2}{3}\ln(1+t^{3/2})-\frac{2}{3}\ln 2\right]=\infty,$$

so the series $\sum_{n=1}^{\infty}\frac{\sqrt{n}}{1+n^{3/2}}$ diverges.

17. The function $f(x)=\frac{1}{x^2+4}$ is continuous, positive, and decreasing on $[1,\infty)$, so we can apply the Integral Test.

$$\int_1^{\infty}\frac{1}{x^2+4}\,dx=\lim_{t\to\infty}\int_1^{t}\frac{1}{x^2+4}\,dx=\lim_{t\to\infty}\left[\frac{1}{2}\tan^{-1}\frac{x}{2}\right]_1^{t}=\frac{1}{2}\lim_{t\to\infty}\left[\tan^{-1}\left(\frac{t}{2}\right)-\tan^{-1}\left(\frac{1}{2}\right)\right]$$
$$=\frac{1}{2}\left[\frac{\pi}{2}-\tan^{-1}\left(\frac{1}{2}\right)\right]$$

Therefore, the series $\sum_{n=1}^{\infty}\frac{1}{n^2+4}$ converges.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. The function $f(x) = \dfrac{1}{x^2+2x+2}$ is continuous, positive, and decreasing on $[1,\infty)$, so the Integral Test applies.

$$\int_1^\infty \frac{1}{x^2+2x+2}\,dx = \lim_{t\to\infty}\int_1^t \frac{1}{(x+1)^2+1}\,dx = \lim_{t\to\infty}\Big[\arctan(x+1)\Big]_1^t$$
$$= \lim_{t\to\infty}[\arctan(t+1) - \arctan 2] = \frac{\pi}{2} - \arctan 2,$$

so the series $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n^2+2n+2}$ converges.

19. The function $f(x) = \dfrac{x^3}{x^4+4}$ is continuous and positive on $[2,\infty)$, and is also decreasing since

$f'(x) = \dfrac{(x^4+4)(3x^2) - x^3(4x^3)}{(x^4+4)^2} = \dfrac{12x^2 - x^6}{(x^4+4)^2} = \dfrac{x^2(12-x^4)}{(x^4+4)^2} < 0$ for $x > \sqrt[4]{12} \approx 1.86$, so we can use the Integral Test on $[2,\infty)$.

$\displaystyle\int_2^\infty \frac{x^3}{x^4+4}\,dx = \lim_{t\to\infty}\int_2^t \frac{x^3}{x^4+4}\,dx = \lim_{t\to\infty}\Big[\tfrac{1}{4}\ln(x^4+4)\Big]_2^t = \lim_{t\to\infty}\Big[\tfrac{1}{4}\ln(t^4+4) - \tfrac{1}{4}\ln 20\Big] = \infty$, so the series $\displaystyle\sum_{n=2}^{\infty} \frac{n^3}{n^4+4}$ diverges, and it follows that $\displaystyle\sum_{n=1}^{\infty} \frac{n^3}{n^4+4}$ diverges as well.

20. The function $f(x) = \dfrac{3x-4}{x^2-2x} = \dfrac{2}{x} + \dfrac{1}{x-2}$ [by partial fractions] is continuous, positive, and decreasing on $[3,\infty)$ since it is the sum of two such functions, so we can apply the Integral Test.

$$\int_3^\infty \frac{3x-4}{x^2-x}\,dx = \lim_{t\to\infty}\int_3^t \left[\frac{2}{x} + \frac{1}{x-2}\right]dx = \lim_{t\to\infty}\Big[2\ln x + \ln(x-2)\Big]_3^t = \lim_{t\to\infty}[2\ln t + \ln(t-2) - 2\ln 3] = \infty.$$

The integral is divergent, so the series $\displaystyle\sum_{n=3}^{\infty} \frac{3n-4}{n^2-n}$ is divergent.

21. $f(x) = \dfrac{1}{x\ln x}$ is continuous and positive on $[2,\infty)$, and also decreasing since $f'(x) = -\dfrac{1+\ln x}{x^2(\ln x)^2} < 0$ for $x > 2$, so we can use the Integral Test. $\displaystyle\int_2^\infty \frac{1}{x\ln x}\,dx = \lim_{t\to\infty}[\ln(\ln x)]_2^t = \lim_{t\to\infty}[\ln(\ln t) - \ln(\ln 2)] = \infty$, so the series $\displaystyle\sum_{n=2}^{\infty} \frac{1}{n\ln n}$ diverges.

22. The function $f(x) = \dfrac{\ln x}{x^2}$ is continuous and positive on $[2,\infty)$, and also decreasing since

$f'(x) = \dfrac{x^2(1/x) - (\ln x)(2x)}{(x^2)^2} = \dfrac{x - 2x\ln x}{x^4} = \dfrac{1-2\ln x}{x^3} < 0$ for $x > e^{1/2} \approx 1.65$, so we can use the Integral Test on $[2,\infty)$.

$$\int_2^\infty \frac{\ln x}{x^2}\,dx = \lim_{t\to\infty}\int_2^t \frac{\ln x}{x^2}\,dx = \lim_{t\to\infty}\left(\left[-\frac{\ln x}{x}\right]_2^t + \int_2^t \frac{1}{x^2}\,dx\right) \qquad \left[\begin{array}{c}\text{by parts with}\\ u=\ln x,\ dv=(1/x^2)\,dx\end{array}\right]$$
$$= \lim_{t\to\infty}\left(-\frac{\ln t}{t} + \frac{\ln 2}{2} + \left[-\frac{1}{x}\right]_2^t\right) \overset{\text{H}}{=} \lim_{t\to\infty}\left(-\frac{1/t}{1} + \frac{\ln 2}{2} - \frac{1}{t} + \frac{1}{2}\right) = \frac{\ln 2 + 1}{2},$$

so the series $\displaystyle\sum_{n=2}^{\infty} \frac{\ln n}{n^2}$ converges.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

23. The function $f(x) = xe^{-x} = \dfrac{x}{e^x}$ is continuous and positive on $[1, \infty)$, and also decreasing since

$f'(x) = \dfrac{e^x \cdot 1 - xe^x}{(e^x)^2} = \dfrac{e^x(1-x)}{(e^x)^2} = \dfrac{1-x}{e^x} < 0$ for $x > 1$ [and $f(1) > f(2)$], so we can use the Integral Test on $[1, \infty)$.

$$\int_1^\infty xe^{-x}\,dx = \lim_{t\to\infty}\int_1^t xe^{-x}\,dx = \lim_{t\to\infty}\left(\left[-xe^{-x}\right]_1^t + \int_1^t e^{-x}\,dx\right) \qquad \left[\begin{array}{c}\text{by parts with}\\ u = x,\ dv = e^{-x}\,dx\end{array}\right]$$

$$= \lim_{t\to\infty}\left(-te^{-t} + e^{-1} + \left[-e^{-x}\right]_1^t\right) = \lim_{t\to\infty}\left(-\frac{t}{e^t} + \frac{1}{e} - \frac{1}{e^t} + \frac{1}{e}\right)$$

$$\overset{\text{H}}{=} \lim_{t\to\infty}\left(-\frac{1}{e^t} + \frac{1}{e} - 0 + \frac{1}{e}\right) = \frac{2}{e},$$

so the series $\displaystyle\sum_{k=1}^{\infty} ke^{-k}$ converges.

24. The function $f(x) = xe^{-x^2} = \dfrac{x}{e^{x^2}}$ is continuous and positive on $[1, \infty)$, and also decreasing since

$f'(x) = \dfrac{e^{x^2}\cdot 1 - xe^{x^2}\cdot 2x}{(e^{x^2})^2} = \dfrac{1-2x^2}{e^{x^2}} < 0$ for $x > \sqrt{\frac{1}{2}} \approx 0.7$, so we can use the Integral Test on $[1, \infty)$.

$\displaystyle\int_1^\infty xe^{-x^2}\,dx = \lim_{t\to\infty}\int_1^t xe^{-x^2}\,dx = \lim_{t\to\infty}\left[-\tfrac{1}{2}e^{-x^2}\right]_1^t = \lim_{t\to\infty}\left(-\tfrac{1}{2}e^{-t^2} + \tfrac{1}{2}e^{-1}\right) = \frac{1}{2e}$, so the series $\displaystyle\sum_{k=1}^{\infty} ke^{-k^2}$ converges.

25. The function $f(x) = \dfrac{1}{x^2+x^3} = \dfrac{1}{x^2} - \dfrac{1}{x} + \dfrac{1}{x+1}$ [by partial fractions] is continuous, positive and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty f(x)\,dx = \lim_{t\to\infty}\int_1^t \left(\frac{1}{x^2} - \frac{1}{x} + \frac{1}{x+1}\right)dx = \lim_{t\to\infty}\left[-\frac{1}{x} - \ln x + \ln(x+1)\right]_1^t$$

$$= \lim_{t\to\infty}\left[-\frac{1}{t} + \ln\frac{t+1}{t} + 1 - \ln 2\right] = 0 + 0 + 1 - \ln 2$$

The integral converges, so the series $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n^2+n^3}$ converges.

26. The function $f(x) = \dfrac{x}{x^4+1}$ is positive, continuous, and decreasing on $[1, \infty)$. [Note that

$f'(x) = \dfrac{x^4+1-4x^4}{(x^4+1)^2} = \dfrac{1-3x^4}{(x^4+1)^2} < 0$ on $[1, \infty)$.] Thus, we can apply the Integral Test.

$\displaystyle\int_1^\infty \frac{x}{x^4+1}\,dx = \lim_{t\to\infty}\int_1^t \frac{\frac{1}{2}(2x)}{1+(x^2)^2}\,dx = \lim_{t\to\infty}\left[\frac{1}{2}\tan^{-1}(x^2)\right]_1^t = \frac{1}{2}\lim_{t\to\infty}\left[\tan^{-1}(t^2) - \tan^{-1}1\right] = \frac{1}{2}\left(\frac{\pi}{2} - \frac{\pi}{4}\right) = \frac{\pi}{8}$

so the series $\displaystyle\sum_{n=1}^{\infty} \frac{n}{n^4+1}$ converges.

27. The function $f(x) = \dfrac{\cos \pi x}{\sqrt{x}}$ is neither positive nor decreasing on $[1, \infty)$, so the hypotheses of the Integral Test are not satisfied for the series $\displaystyle\sum_{n=1}^{\infty} \frac{\cos \pi n}{\sqrt{n}}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. The function $f(x) = \dfrac{\cos^2 x}{1+x^2}$ is not decreasing on $[1,\infty)$, so the hypotheses of the Integral Test are not satisfied for the series $\displaystyle\sum_{n=1}^{\infty} \frac{\cos^2 n}{1+n^2}$.

29. We have already shown (in Exercise 21) that when $p = 1$ the series $\displaystyle\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^p}$ diverges, so assume that $p \neq 1$.

$f(x) = \dfrac{1}{x(\ln x)^p}$ is continuous and positive on $[2,\infty)$, and $f'(x) = -\dfrac{p + \ln x}{x^2(\ln x)^{p+1}} < 0$ if $x > e^{-p}$, so that f is eventually decreasing and we can use the Integral Test.

$$\int_2^{\infty} \frac{1}{x(\ln x)^p}\,dx = \lim_{t\to\infty}\left[\frac{(\ln x)^{1-p}}{1-p}\right]_2^t \quad [\text{for } p \neq 1] = \lim_{t\to\infty}\left[\frac{(\ln t)^{1-p}}{1-p} - \frac{(\ln 2)^{1-p}}{1-p}\right]$$

This limit exists whenever $1 - p < 0 \;\Leftrightarrow\; p > 1$, so the series converges for $p > 1$.

30. $f(x) = \dfrac{1}{x \ln x\,[\ln(\ln x)]^p}$ is positive and continuous on $[3,\infty)$. For $p \geq 0$, f clearly decreases on $[3,\infty)$; and for $p < 0$, it can be verified that f is ultimately decreasing. Thus, we can apply the Integral Test.

$$I = \int_3^{\infty} \frac{dx}{x \ln x\,[\ln(\ln x)]^p} = \lim_{t\to\infty}\int_3^t \frac{[\ln(\ln x)]^{-p}}{x \ln x}\,dx = \lim_{t\to\infty}\left[\frac{[\ln(\ln x)]^{-p+1}}{-p+1}\right]_3^t \qquad [\text{for } p \neq 1]$$

$$= \lim_{t\to\infty}\left[\frac{[\ln(\ln t)]^{-p+1}}{-p+1} - \frac{[\ln(\ln 3)]^{-p+1}}{-p+1}\right],$$

which exists whenever $-p + 1 < 0 \;\Leftrightarrow\; p > 1$. If $p = 1$, then $I = \lim_{t\to\infty}\left[\ln(\ln(\ln x))\right]_3^t = \infty$. Therefore,

$\displaystyle\sum_{n=3}^{\infty} \frac{1}{n \ln n\,[\ln(\ln n)]^p}$ converges for $p > 1$.

31. Clearly the series cannot converge if $p \geq -\frac{1}{2}$, because then $\lim_{n\to\infty} n(1+n^2)^p \neq 0$. So assume $p < -\frac{1}{2}$. Then

$f(x) = x(1+x^2)^p$ is continuous, positive, and eventually decreasing on $[1,\infty)$, and we can use the Integral Test.

$$\int_1^{\infty} x(1+x^2)^p dx = \lim_{t\to\infty}\left[\frac{1}{2}\cdot\frac{(1+x^2)^{p+1}}{p+1}\right]_1^t = \frac{1}{2(p+1)}\lim_{t\to\infty}\left[(1+t^2)^{p+1} - 2^{p+1}\right].$$

This limit exists and is finite $\;\Leftrightarrow\; p + 1 < 0 \;\Leftrightarrow\; p < -1$, so the series $\displaystyle\sum_{n=1}^{\infty} n(1+n^2)^p$ converges whenever $p < -1$.

32. If $p \leq 0$, $\lim_{n\to\infty} \dfrac{\ln n}{n^p} = \infty$ and the series diverges, so assume $p > 0$. $f(x) = \dfrac{\ln x}{x^p}$ is positive and continuous and $f'(x) < 0$ for $x > e^{1/p}$, so f is eventually decreasing and we can use the Integral Test. Integration by parts gives

$$\int_1^{\infty} \frac{\ln x}{x^p}dx = \lim_{t\to\infty}\left[\frac{x^{1-p}\left[(1-p)\ln x - 1\right]}{(1-p)^2}\right]_1^t \quad (\text{for } p \neq 1) = \frac{1}{(1-p)^2}\left[\lim_{t\to\infty} t^{1-p}\left[(1-p)\ln t - 1\right] + 1\right], \text{ which exists}$$

whenever $1 - p < 0 \;\Leftrightarrow\; p > 1$. Thus, $\displaystyle\sum_{n=1}^{\infty} \frac{\ln n}{n^p}$ converges $\;\Leftrightarrow\; p > 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. Since this is a p-series with $p = x$, $\zeta(x)$ is defined when $x > 1$. Unless specified otherwise, the domain of a function f is the set of real numbers x such that the expression for $f(x)$ makes sense and defines a real number. So, in the case of a series, it's the set of real numbers x such that the series is convergent.

34. (a) $\sum_{n=2}^{\infty} \frac{1}{n^2} = \sum_{n=1}^{\infty} \frac{1}{n^2} - \frac{1}{1^2}$ [subtract a_1] $= \frac{\pi^2}{6} - 1$

(b) $\sum_{n=3}^{\infty} \frac{1}{(n+1)^2} = \sum_{n=4}^{\infty} \frac{1}{n^2} = \sum_{n=1}^{\infty} \frac{1}{n^2} - \left(\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2}\right) = \frac{\pi^2}{6} - \frac{49}{36}$

(c) $\sum_{n=1}^{\infty} \frac{1}{(2n)^2} = \sum_{n=1}^{\infty} \frac{1}{4n^2} = \frac{1}{4}\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{4}\left(\frac{\pi^2}{6}\right) = \frac{\pi^2}{24}$

35. (a) $\sum_{n=1}^{\infty} \left(\frac{3}{n}\right)^4 = \sum_{n=1}^{\infty} \frac{81}{n^4} = 81\sum_{n=1}^{\infty} \frac{1}{n^4} = 81\left(\frac{\pi^4}{90}\right) = \frac{9\pi^4}{10}$

(b) $\sum_{k=5}^{\infty} \frac{1}{(k-2)^4} = \frac{1}{3^4} + \frac{1}{4^4} + \frac{1}{5^4} + \cdots = \sum_{k=3}^{\infty} \frac{1}{k^4} = \frac{\pi^4}{90} - \left(\frac{1}{1^4} + \frac{1}{2^4}\right)$ [subtract a_1 and a_2] $= \frac{\pi^4}{90} - \frac{17}{16}$

36. (a) $f(x) = 1/x^4$ is positive and continuous and $f'(x) = -4/x^5$ is negative for $x > 0$, and so the Integral Test applies.

$\sum_{n=1}^{\infty} \frac{1}{n^4} \approx s_{10} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \cdots + \frac{1}{10^4} \approx 1.082037.$

$R_{10} \le \int_{10}^{\infty} \frac{1}{x^4}\,dx = \lim_{t\to\infty}\left[\frac{1}{-3x^3}\right]_{10}^{t} = \lim_{t\to\infty}\left(-\frac{1}{3t^3} + \frac{1}{3\,(10)^3}\right) = \frac{1}{3000}$, so the error is at most $0.000\overline{3}$.

(b) $s_{10} + \int_{11}^{\infty} \frac{1}{x^4}\,dx \le s \le s_{10} + \int_{10}^{\infty} \frac{1}{x^4}\,dx \quad \Rightarrow \quad s_{10} + \frac{1}{3(11)^3} \le s \le s_{10} + \frac{1}{3(10)^3}$ $\Rightarrow$

$1.082037 + 0.000250 = 1.082287 \le s \le 1.082037 + 0.000333 = 1.082370$, so we get $s \approx 1.08233$ with error ≤ 0.00005.

(c) The estimate in part (b) is $s \approx 1.08233$ with error ≤ 0.00005. The exact value given in Exercise 35 is $\pi^4/90 \approx 1.082323$. The difference is less than 0.00001.

(d) $R_n \le \int_{n}^{\infty} \frac{1}{x^4}\,dx = \frac{1}{3n^3}$. So $R_n < 0.00001 \quad \Rightarrow \quad \frac{1}{3n^3} < \frac{1}{10^5} \quad \Rightarrow \quad 3n^3 > 10^5 \quad \Rightarrow \quad n > \sqrt[3]{(10)^5/3} \approx 32.2$,

that is, for $n > 32$.

37. (a) $f(x) = \frac{1}{x^2}$ is positive and continuous and $f'(x) = -\frac{2}{x^3}$ is negative for $x > 0$, and so the Integral Test applies.

$\sum_{n=1}^{\infty} \frac{1}{n^2} \approx s_{10} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \cdots + \frac{1}{10^2} \approx 1.549768.$

$R_{10} \le \int_{10}^{\infty} \frac{1}{x^2}\,dx = \lim_{t\to\infty}\left[\frac{-1}{x}\right]_{10}^{t} = \lim_{t\to\infty}\left(-\frac{1}{t} + \frac{1}{10}\right) = \frac{1}{10}$, so the error is at most 0.1.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $s_{10} + \int_{11}^{\infty} \frac{1}{x^2}\,dx \le s \le s_{10} + \int_{10}^{\infty} \frac{1}{x^2}\,dx \;\Rightarrow\; s_{10} + \frac{1}{11} \le s \le s_{10} + \frac{1}{10} \;\Rightarrow$

$1.549768 + 0.090909 = 1.640677 \le s \le 1.549768 + 0.1 = 1.649768$, so we get $s \approx 1.64522$ (the average of 1.640677 and 1.649768) with error ≤ 0.005 (the maximum of $1.649768 - 1.64522$ and $1.64522 - 1.640677$, rounded up).

(c) The estimate in part (b) is $s \approx 1.64522$ with error ≤ 0.005. The exact value given in Exercise 34 is $\pi^2/6 \approx 1.644934$. The difference is less than 0.0003.

(d) $R_n \le \int_n^{\infty} \frac{1}{x^2}\,dx = \frac{1}{n}$. So $R_n < 0.001$ if $\frac{1}{n} < \frac{1}{1000} \;\Leftrightarrow\; n > 1000$.

38. $f(x) = xe^{-2x}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies. Using (2),

$$R_n \le \int_n^{\infty} xe^{-2x}\,dx = \lim_{t\to\infty}\left(\left[-\tfrac{1}{2}xe^{-2x}\right]_n^t + \int_n^t \tfrac{1}{2}e^{-2x}\,dx\right) \qquad \left[\begin{array}{c}\text{using parts with}\\ u = x,\ dv = e^{-2x}\,dx\end{array}\right]$$

$$= \lim_{t\to\infty}\left(\frac{-t}{2e^{2t}} + \frac{n}{2e^{2n}} - \frac{1}{4e^{2t}} + \frac{1}{4e^{2n}}\right) \overset{\text{H}}{=} 0 + \frac{n}{2e^{2n}} - 0 + \frac{1}{4e^{2n}} = \frac{2n+1}{4e^{2n}}$$

To be correct to four decimal places, we want $\frac{2n+1}{4e^{2n}} \le \frac{5}{10^5}$. This inequality is true for $n = 6$.

$$s_6 = \sum_{n=1}^{6} \frac{n}{e^{2n}} = \frac{1}{e^2} + \frac{2}{e^4} + \frac{3}{e^6} + \frac{4}{e^8} + \frac{5}{e^{10}} + \frac{6}{e^{12}} \approx 0.1810.$$

39. $f(x) = 1/(2x+1)^6$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies. Using (2),

$R_n \le \int_n^{\infty} (2x+1)^{-6}\,dx = \lim_{t\to\infty}\left[\frac{-1}{10(2x+1)^5}\right]_n^t = \frac{1}{10(2n+1)^5}$. To be correct to five decimal places, we want

$\frac{1}{10(2n+1)^5} \le \frac{5}{10^6} \;\Leftrightarrow\; (2n+1)^5 \ge 20{,}000 \;\Leftrightarrow\; n \ge \frac{1}{2}\left(\sqrt[5]{20{,}000} - 1\right) \approx 3.12$, so use $n = 4$.

$$s_4 = \sum_{n=1}^{4} \frac{1}{(2n+1)^6} = \frac{1}{3^6} + \frac{1}{5^6} + \frac{1}{7^6} + \frac{1}{9^6} \approx 0.001\,446 \approx 0.00145.$$

40. $f(x) = \frac{1}{x(\ln x)^2}$ is positive and continuous and $f'(x) = -\frac{\ln x + 2}{x^2(\ln x)^3}$ is negative for $x > 1$, so the Integral Test applies.

Using (2), we need $0.01 > \int_n^{\infty} \frac{dx}{x(\ln x)^2} = \lim_{t\to\infty}\left[\frac{-1}{\ln x}\right]_n^t = \frac{1}{\ln n}$. This is true for $n > e^{100}$, so we would have to add this many terms to find the sum of the series $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}$ to within 0.01, which would be problematic because $e^{100} \approx 2.7 \times 10^{43}$.

41. $\sum_{n=1}^{\infty} n^{-1.001} = \sum_{n=1}^{\infty} \frac{1}{n^{1.001}}$ is a convergent p-series with $p = 1.001 > 1$. Using (2), we get

$R_n \le \int_n^{\infty} x^{-1.001}\,dx = \lim_{t\to\infty}\left[\frac{x^{-0.001}}{-0.001}\right]_n^t = -1000 \lim_{t\to\infty}\left[\frac{1}{x^{0.001}}\right]_n^t = 1000\left(-\frac{1}{n^{0.001}}\right) = \frac{1000}{n^{0.001}}$. We want

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$R_n < 0.000\,000\,005 \quad \Leftrightarrow \quad \frac{1000}{n^{0.001}} < 5 \times 10^{-9} \quad \Leftrightarrow \quad n^{0.001} > \frac{1000}{5 \times 10^{-9}} \quad \Leftrightarrow$

$n > (2 \times 10^{11})^{1000} = 2^{1000} \times 10^{11,000} \approx 1.07 \times 10^{301} \times 10^{11,000} = 1.07 \times 10^{11,301}$.

42. (a) $f(x) = \left(\frac{\ln x}{x}\right)^2$ is continuous and positive for $x > 1$, and since $f'(x) = \frac{2(\ln x)(1 - \ln x)}{x^3} < 0$ for $x > e$, we can apply the Integral Test. Using a CAS, we get $\int_1^\infty \left(\frac{\ln x}{x}\right)^2 dx = 2$, so the series $\sum_{n=1}^{\infty} \left(\frac{\ln n}{n}\right)^2$ also converges.

(b) Since the Integral Test applies, the error in $s \approx s_n$ is $R_n \le \int_n^\infty \left(\frac{\ln x}{x}\right)^2 dx = \frac{(\ln n)^2 + 2\ln n + 2}{n}$.

(c) By graphing the functions $y_1 = \frac{(\ln x)^2 + 2\ln x + 2}{x}$ and $y_2 = 0.05$, we see that $y_1 < y_2$ for $n \ge 1373$.

(d) Using the CAS to sum the first 1373 terms, we get $s_{1373} \approx 1.94$.

43. (a) From the figure, $a_2 + a_3 + \cdots + a_n \le \int_1^n f(x)\,dx$, so with

$f(x) = \frac{1}{x}$,

$\frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n} \le \int_1^n \frac{1}{x}\,dx = \ln n$.

Thus, $s_n = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n} \le 1 + \ln n$.



(b) By part (a), $s_{10^6} \le 1 + \ln 10^6 \approx 14.82 < 15$ and

$s_{10^9} \le 1 + \ln 10^9 \approx 21.72 < 22$.

44. (a) The sum of the areas of the n rectangles in the graph to the right is

$1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}$. Now $\int_1^{n+1} \frac{dx}{x}$ is less than this sum because

the rectangles extend above the curve $y = 1/x$, so

$\int_1^{n+1} \frac{1}{x}\,dx = \ln(n+1) < 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}$, and since

$\ln n < \ln(n+1)$, $0 < 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \ln n = t_n$.



(b) The area under $f(x) = 1/x$ between $x = n$ and $x = n+1$ is

$\int_n^{n+1} \frac{dx}{x} = \ln(n+1) - \ln n$, and this is clearly greater than the area of

the inscribed rectangle in the figure to the right which is $\frac{1}{n+1}$, so

$t_n - t_{n+1} = \left[\ln(n+1) - \ln n\right] - \frac{1}{n+1} > 0$, and so $t_n > t_{n+1}$, so $\{t_n\}$ is a decreasing sequence.

(c) We have shown that $\{t_n\}$ is decreasing and that $t_n > 0$ for all n. Thus, $0 < t_n \le t_1 = 1$, so $\{t_n\}$ is a bounded monotonic sequence, and hence converges by the Monotonic Sequence Theorem.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. $b^{\ln n} = \left(e^{\ln b}\right)^{\ln n} = \left(e^{\ln n}\right)^{\ln b} = n^{\ln b} = \dfrac{1}{n^{-\ln b}}$. This is a p-series, which converges for all b such that $-\ln b > 1 \;\Leftrightarrow$ $\ln b < -1 \;\Leftrightarrow\; b < e^{-1} \;\Leftrightarrow\; b < 1/e$ [with $b > 0$].

46. For the series $\displaystyle\sum_{n=1}^{\infty}\left(\frac{c}{n} - \frac{1}{n+1}\right)$,

$$s_n = \sum_{i=1}^{n}\left(\frac{c}{i} - \frac{1}{i+1}\right) = \left(\frac{c}{1} - \frac{1}{2}\right) + \left(\frac{c}{2} - \frac{1}{3}\right) + \left(\frac{c}{3} - \frac{1}{4}\right) + \cdots + \left(\frac{c}{n} - \frac{1}{n+1}\right)$$

$$= \frac{c}{1} + \frac{c-1}{2} + \frac{c-1}{3} + \frac{c-1}{4} + \cdots + \frac{c-1}{n} - \frac{1}{n+1} = c + (c-1)\left(\frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n}\right) - \frac{1}{n+1}$$

Thus, $\displaystyle\sum_{n=1}^{\infty}\left(\frac{c}{n} - \frac{1}{n+1}\right) = \lim_{n\to\infty} s_n = \lim_{n\to\infty}\left[c + (c-1)\sum_{i=2}^{n}\frac{1}{i} - \frac{1}{n+1}\right]$. Since a constant multiple of a divergent series is divergent, the last limit exists only if $c - 1 = 0$, so the original series converges only if $c = 1$.

11.4 The Comparison Tests

1. (a) We cannot say anything about $\sum a_n$. If $a_n > b_n$ for all n and $\sum b_n$ is convergent, then $\sum a_n$ could be convergent or divergent. (See the note after Example 2.)

(b) If $a_n < b_n$ for all n, then $\sum a_n$ is convergent. [This is part (i) of the Comparison Test.]

2. (a) If $a_n > b_n$ for all n, then $\sum a_n$ is divergent. [This is part (ii) of the Comparison Test.]

(b) We cannot say anything about $\sum a_n$. If $a_n < b_n$ for all n and $\sum b_n$ is divergent, then $\sum a_n$ could be convergent or divergent.

3. $\dfrac{1}{n^3+8} < \dfrac{1}{n^3}$ for all $n \ge 1$, so $\displaystyle\sum_{n=1}^{\infty}\frac{1}{n^3+8}$ converges by comparison with $\displaystyle\sum_{n=1}^{\infty}\frac{1}{n^3}$, which converges because it is a p-series with $p = 3 > 1$.

4. $\dfrac{1}{\sqrt{n}-1} > \dfrac{1}{\sqrt{n}}$ for all $n \ge 2$, so $\displaystyle\sum_{n=2}^{\infty}\frac{1}{\sqrt{n}-1}$ diverges by comparison with $\displaystyle\sum_{n=2}^{\infty}\frac{1}{\sqrt{n}}$, which diverges because it is a p-series with $p = \frac{1}{2} \le 1$.

5. $\dfrac{n+1}{n\sqrt{n}} > \dfrac{n}{n\sqrt{n}} = \dfrac{1}{\sqrt{n}}$ for all $n \ge 1$, so $\displaystyle\sum_{n=1}^{\infty}\frac{n+1}{n\sqrt{n}}$ diverges by comparison with $\displaystyle\sum_{n=1}^{\infty}\frac{1}{\sqrt{n}}$, which diverges because it is a p-series with $p = \frac{1}{2} \le 1$.

6. $\dfrac{n-1}{n^3+1} < \dfrac{n}{n^3+1} < \dfrac{n}{n^3} = \dfrac{1}{n^2}$ for all $n \ge 1$, so $\displaystyle\sum_{n=1}^{\infty}\frac{n-1}{n^3+1}$ converges by comparison with $\displaystyle\sum_{n=1}^{\infty}\frac{1}{n^2}$, which converges because it is a p-series with $p = 2 > 1$.

7. $\dfrac{9^n}{3+10^n} < \dfrac{9^n}{10^n} = \left(\dfrac{9}{10}\right)^n$ for all $n \ge 1$. $\displaystyle\sum_{n=1}^{\infty}\left(\tfrac{9}{10}\right)^n$ is a convergent geometric series $\left(|r| = \frac{9}{10} < 1\right)$, so $\displaystyle\sum_{n=1}^{\infty}\frac{9^n}{3+10^n}$ converges by the Comparison Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $\dfrac{6^n}{5^n-1} > \dfrac{6^n}{5^n} = \left(\dfrac{6}{5}\right)^n$ for all $n \geq 1$. $\sum_{n=1}^{\infty} \left(\frac{6}{5}\right)^n$ is a divergent geometric series $\left(|r| = \frac{6}{5} > 1\right)$, so $\sum_{n=1}^{\infty} \dfrac{6^n}{5^n-1}$ diverges by the Comparison Test.

9. $\dfrac{\ln k}{k} > \dfrac{1}{k}$ for all k ≥ 3 [since $\ln k > 1$ for $k \geq 3$], so $\sum_{k=3}^{\infty} \dfrac{\ln k}{k}$ diverges by comparison with $\sum_{k=3}^{\infty} \dfrac{1}{k}$, which diverges because it is a p-series with $p = 1 \leq 1$ (the harmonic series). Thus, $\sum_{k=1}^{\infty} \dfrac{\ln k}{k}$ diverges since a finite number of terms doesn't affect the convergence or divergence of a series.

10. $\dfrac{k \sin^2 k}{1+k^3} \leq \dfrac{k}{1+k^3} < \dfrac{k}{k^3} = \dfrac{1}{k^2}$ for all $k \geq 1$, so $\sum_{k=1}^{\infty} \dfrac{k \sin^2 k}{1+k^3}$ converges by comparison with $\sum_{k=1}^{\infty} \dfrac{1}{k^2}$, which converges because it is a p-series with $p = 2 > 1$.

11. $\dfrac{\sqrt[3]{k}}{\sqrt{k^3+4k+3}} < \dfrac{\sqrt[3]{k}}{\sqrt{k^3}} = \dfrac{k^{1/3}}{k^{3/2}} = \dfrac{1}{k^{7/6}}$ for all $k \geq 1$, so $\sum_{k=1}^{\infty} \dfrac{\sqrt[3]{k}}{\sqrt{k^3+4k+3}}$ converges by comparison with $\sum_{k=1}^{\infty} \dfrac{1}{k^{7/6}}$, which converges because it is a p-series with $p = \frac{7}{6} > 1$.

12. $\dfrac{(2k-1)(k^2-1)}{(k+1)(k^2+4)^2} < \dfrac{2k(k^2)}{k(k^2)^2} = \dfrac{2k^3}{k^5} = \dfrac{2}{k^2}$ for all $k \geq 1$, so $\sum_{k=1}^{\infty} \dfrac{(2k-1)(k^2-1)}{(k+1)(k^2+4)^2}$ converges by comparison with $2\sum_{k=1}^{\infty} \dfrac{1}{k^2}$, which converges because it is a constant multiple of a p-series with $p = 2 > 1$.

13. $\dfrac{1+\cos n}{e^n} < \dfrac{2}{e^n}$ for all $n \geq 1$. $\sum_{n=1}^{\infty} \dfrac{2}{e^n}$ is a convergent geometric series $\left(|r| = \frac{1}{e} < 1\right)$, so $\sum_{n=1}^{\infty} \dfrac{1+\cos n}{e^n}$ converges by the Comparison Test.

14. $\dfrac{1}{\sqrt[3]{3n^4+1}} < \dfrac{1}{\sqrt[3]{3n^4}} < \dfrac{1}{\sqrt[3]{n^4}} = \dfrac{1}{n^{4/3}}$ for all $n \geq 1$, so $\sum_{n=1}^{\infty} \dfrac{1}{\sqrt[3]{3n^4+1}}$ converges by comparison with $\sum_{n=1}^{\infty} \dfrac{1}{n^{4/3}}$, which converges because it is a p-series with $p = \frac{4}{3} > 1$.

15. $\dfrac{4^{n+1}}{3^n-2} > \dfrac{4 \cdot 4^n}{3^n} = 4\left(\dfrac{4}{3}\right)^n$ for all $n > 1$. $\sum_{n=1}^{\infty} 4\left(\dfrac{4}{3}\right)^n = 4\sum_{n=1}^{\infty} \left(\dfrac{4}{3}\right)^n$ is a divergent geometric series $\left(|r| = \frac{4}{3} > 1\right)$, so $\sum_{n=1}^{\infty} \dfrac{4^{n+1}}{3^n-2}$ diverges by the Comparison Test.

16. $\dfrac{1}{n^n} \leq \dfrac{1}{n^2}$ for all $n \geq 1$, so $\sum_{n=1}^{\infty} \dfrac{1}{n^n}$ converges by comparison with $\sum_{n=1}^{\infty} \dfrac{1}{n^2}$, which converges because it is a p-series with $p = 2 > 1$.

17. Use the Limit Comparison Test with $a_n = \dfrac{1}{\sqrt{n^2+1}}$ and $b_n = \dfrac{1}{n}$:

$\lim_{n\to\infty} \dfrac{a_n}{b_n} = \lim_{n\to\infty} \dfrac{n}{\sqrt{n^2+1}} = \lim_{n\to\infty} \dfrac{1}{\sqrt{1+(1/n^2)}} = 1 > 0$. Since the harmonic series $\sum_{n=1}^{\infty} \dfrac{1}{n}$ diverges, so does $\sum_{n=1}^{\infty} \dfrac{1}{\sqrt{n^2+1}}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. Use the Limit Comparison Test with $a_n = \dfrac{2}{\sqrt{n}+2}$ and $b_n = \dfrac{1}{\sqrt{n}}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{2\sqrt{n}}{\sqrt{n}+2} = \lim\limits_{n\to\infty} \dfrac{2}{1+2/\sqrt{n}} = 2 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{\sqrt{n}}$ is a divergent p-series [$p = \frac{1}{2} < 1$], the series $\sum\limits_{n=1}^{\infty} \dfrac{2}{\sqrt{n}+2}$ is also divergent.

19. Use the Limit Comparison Test with $a_n = \dfrac{n+1}{n^3+n}$ and $b_n = \dfrac{1}{n^2}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{(n+1)n^2}{n(n^2+1)} = \lim\limits_{n\to\infty} \dfrac{n^2+n}{n^2+1} = \lim\limits_{n\to\infty} \dfrac{1+1/n}{1+1/n^2} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$ is a convergent p-series [$p = 2 > 1$], the series $\sum\limits_{n=1}^{\infty} \dfrac{n+1}{n^3+n}$ also converges.

20. Use the Limit Comparison Test with $a_n = \dfrac{n^2+n+1}{n^4+n^2}$ and $b_n = \dfrac{1}{n^2}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{(n^2+n+1)n^2}{n^2(n^2+1)} = \lim\limits_{n\to\infty} \dfrac{n^2+n+1}{n^2+1} = \lim\limits_{n\to\infty} \dfrac{1+1/n+1/n^2}{1+1/n^2} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$ is a convergent p-series [$p = 2 > 1$], the series $\sum\limits_{n=1}^{\infty} \dfrac{n^2+n+1}{n^4+n^2}$ also converges.

21. Use the Limit Comparison Test with $a_n = \dfrac{\sqrt{1+n}}{2+n}$ and $b_n = \dfrac{1}{\sqrt{n}}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{\sqrt{1+n}\sqrt{n}}{2+n} = \lim\limits_{n\to\infty} \dfrac{\sqrt{n^2+n}/\sqrt{n^2}}{(2+n)/n} = \lim\limits_{n\to\infty} \dfrac{\sqrt{1+1/n}}{2/n+1} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{\sqrt{n}}$ is a divergent p-series [$p = \frac{1}{2} \le 1$], the series $\sum\limits_{n=1}^{\infty} \dfrac{\sqrt{1+n}}{2+n}$ also diverges.

22. Use the Limit Comparison Test with $a_n = \dfrac{n+2}{(n+1)^3}$ and $b_n = \dfrac{1}{n^2}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n^2(n+2)}{(n+1)^3} = \lim\limits_{n\to\infty} \dfrac{1+\frac{2}{n}}{\left(1+\frac{1}{n}\right)^3} = 1 > 0$. Since $\sum\limits_{n=3}^{\infty} \dfrac{1}{n^2}$ is a convergent (partial) p-series [$p = 2 > 1$], the series $\sum\limits_{n=3}^{\infty} \dfrac{n+2}{(n+1)^3}$ also converges.

23. Use the Limit Comparison Test with $a_n = \dfrac{5+2n}{(1+n^2)^2}$ and $b_n = \dfrac{1}{n^3}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n^3(5+2n)}{(1+n^2)^2} = \lim\limits_{n\to\infty} \dfrac{5n^3+2n^4}{(1+n^2)^2} \cdot \dfrac{1/n^4}{1/(n^2)^2} = \lim\limits_{n\to\infty} \dfrac{\frac{5}{n}+2}{\left(\frac{1}{n^2}+1\right)^2} = 2 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^3}$ is a convergent p-series [$p = 3 > 1$], the series $\sum\limits_{n=1}^{\infty} \dfrac{5+2n}{(1+n^2)^2}$ also converges.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

24. $\dfrac{n+3^n}{n+2^n} > \dfrac{3^n}{n+2^n} > \dfrac{3^n}{2^n+2^n} = \dfrac{3^n}{2\cdot 2^n} = \dfrac{1}{2}\left(\dfrac{3}{2}\right)^n$, so the series $\displaystyle\sum_{n=1}^{\infty} \frac{n+3^n}{n+2^n}$ diverges by comparison with $\dfrac{1}{2}\displaystyle\sum_{n=1}^{\infty}\left(\frac{3}{2}\right)^n$, which is a constant multiple of a divergent geometric series $\left[|r| = \frac{3}{2} > 1\right]$. *Or:* Use the Limit Comparison Test with $a_n = \dfrac{n+3^n}{n+2^n}$ and $b_n = \left(\dfrac{3}{2}\right)^n$.

25. $\dfrac{e^n+1}{ne^n+1} \geq \dfrac{e^n+1}{ne^n+n} = \dfrac{e^n+1}{n(e^n+1)} = \dfrac{1}{n}$ for $n \geq 1$, so the series $\displaystyle\sum_{n=1}^{\infty} \frac{e^n+1}{ne^n+1}$ diverges by comparison with the divergent harmonic series $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n}$. *Or:* Use the Limit Comparison Test with $a_n = \dfrac{e^n+1}{ne^n+1}$ and $b_n = \dfrac{1}{n}$.

26. If $a_n = \dfrac{1}{n\sqrt{n^2-1}}$ and $b_n = \dfrac{1}{n^2}$, then

$\displaystyle\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{n^2}{n\sqrt{n^2-1}} = \lim_{n\to\infty} \frac{n/n}{\sqrt{n^2-1}/n} = \lim_{n\to\infty} \frac{1}{\sqrt{1-1/n^2}} = \frac{1}{1} = 1 > 0$, so $\displaystyle\sum_{n=2}^{\infty} \frac{1}{n\sqrt{n^2-1}}$ converges by the Limit Comparison Test with the convergent series $\displaystyle\sum_{n=2}^{\infty} \frac{1}{n^2}$.

27. Use the Limit Comparison Test with $a_n = \left(1+\dfrac{1}{n}\right)^2 e^{-n}$ and $b_n = e^{-n}$: $\displaystyle\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty}\left(1+\frac{1}{n}\right)^2 = 1 > 0$. Since $\displaystyle\sum_{n=1}^{\infty} e^{-n} = \sum_{n=1}^{\infty} \frac{1}{e^n}$ is a convergent geometric series $\left[|r| = \frac{1}{e} < 1\right]$, the series $\displaystyle\sum_{n=1}^{\infty}\left(1+\frac{1}{n}\right)^2 e^{-n}$ also converges.

28. $\dfrac{e^{1/n}}{n} > \dfrac{1}{n}$ for all $n \geq 1$, so $\displaystyle\sum_{n=1}^{\infty} \frac{e^{1/n}}{n}$ diverges by comparison with the harmonic series $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n}$.

29. Clearly $n! = n(n-1)(n-2)\cdots(3)(2) \geq 2\cdot 2\cdot 2\cdot\cdots\cdot 2\cdot 2 = 2^{n-1}$, so $\dfrac{1}{n!} \leq \dfrac{1}{2^{n-1}}$. $\displaystyle\sum_{n=1}^{\infty} \frac{1}{2^{n-1}}$ is a convergent geometric series $\left[|r| = \frac{1}{2} < 1\right]$, so $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n!}$ converges by the Comparison Test.

30. $\dfrac{n!}{n^n} = \dfrac{1\cdot 2\cdot 3\cdot\cdots\cdot(n-1)n}{n\cdot n\cdot n\cdot\cdots\cdot n\cdot n} \leq \dfrac{1}{n}\cdot\dfrac{2}{n}\cdot 1\cdot 1\cdot\cdots\cdot 1$ for $n > 2$, so since $\displaystyle\sum_{n=1}^{\infty} \frac{2}{n^2}$ converges $[p = 2 > 1]$, $\displaystyle\sum_{n=1}^{\infty} \frac{n!}{n^n}$ converges also by the Comparison Test.

31. Use the Limit Comparison Test with $a_n = \sin\left(\dfrac{1}{n}\right)$ and $b_n = \dfrac{1}{n}$. Then $\sum a_n$ and $\sum b_n$ are series with positive terms and

$\displaystyle\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{\sin(1/n)}{1/n} = \lim_{\theta\to 0} \frac{\sin\theta}{\theta} = 1 > 0$. Since $\displaystyle\sum_{n=1}^{\infty} b_n$ is the divergent harmonic series, $\displaystyle\sum_{n=1}^{\infty} \sin(1/n)$ also diverges. [Note that we could also use l'Hospital's Rule to evaluate the limit:

$\displaystyle\lim_{x\to\infty} \frac{\sin(1/x)}{1/x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{\cos(1/x)\cdot(-1/x^2)}{-1/x^2} = \lim_{x\to\infty} \cos\frac{1}{x} = \cos 0 = 1$.]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. Use the Limit Comparison Test with $a_n = \dfrac{1}{n^{1+1/n}}$ and $b_n = \dfrac{1}{n}$. $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n}{n^{1+1/n}} = \lim\limits_{n\to\infty} \dfrac{1}{n^{1/n}} = 1$

$\left[\text{since } \lim\limits_{x\to\infty} x^{1/x} = 1 \text{ by l'Hospital's Rule}\right]$, so $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ diverges [harmonic series] $\Rightarrow$ $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^{1+1/n}}$ diverges.

33. $\sum\limits_{n=1}^{10} \dfrac{1}{5+n^5} = \dfrac{1}{5+1^5} + \dfrac{1}{5+2^5} + \dfrac{1}{5+3^5} + \cdots + \dfrac{1}{5+10^5} \approx 0.19926$. Now $\dfrac{1}{5+n^5} < \dfrac{1}{n^5}$, so the error is

$$R_{10} \le T_{10} \le \int_{10}^{\infty} \frac{1}{x^5}\,dx = \lim_{t\to\infty} \int_{10}^{t} x^{-5}\,dx = \lim_{t\to\infty} \left[\frac{-1}{4x^4}\right]_{10}^{t} = \lim_{t\to\infty} \left(\frac{-1}{4t^4} + \frac{1}{40{,}000}\right) = \frac{1}{40{,}000} = 0.000\,025.$$

34. $\sum\limits_{n=1}^{10} \dfrac{e^{1/n}}{n^4} = \dfrac{e^{1/1}}{1^4} + \dfrac{e^{1/2}}{2^4} + \dfrac{e^{1/3}}{3^4} + \cdots + \dfrac{e^{1/10}}{10^4} \approx 2.84748$. Now $\dfrac{e^{1/n}}{n^4} \le \dfrac{e}{n^4}$ for $n \ge 1$, so the error is

$$R_{10} \le T_{10} \le \int_{10}^{\infty} \frac{e}{x^4}\,dx = \lim_{t\to\infty} \int_{10}^{t} ex^{-4}\,dx = \lim_{t\to\infty} \left[\frac{-e}{3x^3}\right]_{10}^{t} = \lim_{t\to\infty} \left(\frac{-e}{3t^3} + \frac{e}{3000}\right) = \frac{e}{3000} \approx 0.000\,906.$$

35. $\sum\limits_{n=1}^{10} 5^{-n}\cos^2 n = \dfrac{\cos^2 1}{5} + \dfrac{\cos^2 2}{5^2} + \dfrac{\cos^2 3}{5^3} + \cdots + \dfrac{\cos^2 10}{5^{10}} \approx 0.07393$. Now $\dfrac{\cos^2 n}{5^n} \le \dfrac{1}{5^n}$, so the error is

$$R_{10} \le T_{10} \le \int_{10}^{\infty} \frac{1}{5^x}\,dx = \lim_{t\to\infty} \int_{10}^{t} 5^{-x}\,dx = \lim_{t\to\infty} \left[-\frac{5^{-x}}{\ln 5}\right]_{10}^{t} = \lim_{t\to\infty} \left(-\frac{5^{-t}}{\ln 5} + \frac{5^{-10}}{\ln 5}\right) = \frac{1}{5^{10}\ln 5} < 6.4\times 10^{-8}.$$

36. $\sum\limits_{n=1}^{10} \dfrac{1}{3^n+4^n} = \dfrac{1}{3^1+4^1} + \dfrac{1}{3^2+4^2} + \dfrac{1}{3^3+4^3} + \cdots + \dfrac{1}{3^{10}+4^{10}} \approx 0.19788$. Now $\dfrac{1}{3^n+4^n} < \dfrac{1}{3^n+3^n} = \dfrac{1}{2\cdot 3^n}$, so the error is

$$R_{10} \le T_{10} \le \int_{10}^{\infty} \frac{1}{2\cdot 3^x}\,dx = \lim_{t\to\infty} \int_{10}^{t} \frac{1}{2}\cdot 3^{-x}\,dx = \lim_{t\to\infty} \left[-\frac{1}{2}\frac{3^{-x}}{\ln 3}\right]_{10}^{t} = \lim_{t\to\infty} \left(-\frac{1}{2}\frac{3^{-t}}{\ln 3} + \frac{1}{2}\frac{3^{-10}}{\ln 3}\right)$$
$$= \frac{1}{2\cdot 3^{10}\ln 3} < 7.7\times 10^{-6}.$$

37. Since $\dfrac{d_n}{10^n} \le \dfrac{9}{10^n}$ for each n, and since $\sum\limits_{n=1}^{\infty} \dfrac{9}{10^n}$ is a convergent geometric series $\left(|r| = \frac{1}{10} < 1\right)$, $0.d_1d_2d_3\ldots = \sum\limits_{n=1}^{\infty} \dfrac{d_n}{10^n}$ will always converge by the Comparison Test.

38. Clearly, if $p < 0$ then the series diverges, since $\lim\limits_{n\to\infty} \dfrac{1}{n^p \ln n} = \infty$. If $0 \le p \le 1$, then $n^p \ln n \le n \ln n$ $\Rightarrow$ $\dfrac{1}{n^p \ln n} \ge \dfrac{1}{n \ln n}$ and $\sum\limits_{n=2}^{\infty} \dfrac{1}{n \ln n}$ diverges (Exercise 11.3.21), so $\sum\limits_{n=2}^{\infty} \dfrac{1}{n^p \ln n}$ diverges. If $p > 1$, use the Limit Comparison Test with $a_n = \dfrac{1}{n^p \ln n}$ and $b_n = \dfrac{1}{n^p}$. $\sum\limits_{n=2}^{\infty} b_n$ converges, and $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{1}{\ln n} = 0$, so $\sum\limits_{n=2}^{\infty} \dfrac{1}{n^p \ln n}$ also converges. (Or use the Comparison Test, since $n^p \ln n > n^p$ for $n > e$.) In summary, the series converges if and only if $p > 1$.

39. Since $\sum a_n$ converges, $\lim\limits_{n\to\infty} a_n = 0$, so there exists N such that $|a_n - 0| < 1$ for all $n > N$ $\Rightarrow$ $0 \le a_n < 1$ for all $n > N$ $\Rightarrow$ $0 \le a_n^2 \le a_n$. Since $\sum a_n$ converges, so does $\sum a_n^2$ by the Comparison Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

40. (a) Since $\lim_{n\to\infty}(a_n/b_n) = 0$, there is a number $N > 0$ such that $|a_n/b_n - 0| < 1$ for all $n > N$, and so $a_n < b_n$ since a_n and b_n are positive. Thus, since $\sum b_n$ converges, so does $\sum a_n$ by the Comparison Test.

(b) (i) If $a_n = \dfrac{\ln n}{n^3}$ and $b_n = \dfrac{1}{n^2}$, then $\lim_{n\to\infty}\dfrac{a_n}{b_n} = \lim_{n\to\infty}\dfrac{\ln n}{n} = \lim_{x\to\infty}\dfrac{\ln x}{x} \overset{\text{H}}{=} \lim_{x\to\infty}\dfrac{1/x}{1} = 0$, so $\sum_{n=1}^{\infty}\dfrac{\ln n}{n^3}$ converges by part (a).

(ii) If $a_n = \dfrac{\ln n}{\sqrt{n}e^n}$ and $b_n = \dfrac{1}{e^n}$, then $\lim_{n\to\infty}\dfrac{a_n}{b_n} = \lim_{n\to\infty}\dfrac{\ln n}{\sqrt{n}} = \lim_{x\to\infty}\dfrac{\ln x}{\sqrt{x}} \overset{\text{H}}{=} \lim_{x\to\infty}\dfrac{1/x}{1/(2\sqrt{x})} = \lim_{x\to\infty}\dfrac{2}{\sqrt{x}} = 0$. Now $\sum b_n$ is a convergent geometric series with ratio $r = 1/e$ $[|r| < 1]$, so $\sum a_n$ converges by part (a).

41. (a) Since $\lim_{n\to\infty}\dfrac{a_n}{b_n} = \infty$, there is an integer N such that $\dfrac{a_n}{b_n} > 1$ whenever $n > N$. (Take $M = 1$ in Definition 11.1.5.) Then $a_n > b_n$ whenever $n > N$ and since $\sum b_n$ is divergent, $\sum a_n$ is also divergent by the Comparison Test.

(b) (i) If $a_n = \dfrac{1}{\ln n}$ and $b_n = \dfrac{1}{n}$ for $n \geq 2$, then $\lim_{n\to\infty}\dfrac{a_n}{b_n} = \lim_{n\to\infty}\dfrac{n}{\ln n} = \lim_{x\to\infty}\dfrac{x}{\ln x} \overset{\text{H}}{=} \lim_{x\to\infty}\dfrac{1}{1/x} = \lim_{x\to\infty} x = \infty$, so by part (a), $\sum_{n=2}^{\infty}\dfrac{1}{\ln n}$ is divergent.

(ii) If $a_n = \dfrac{\ln n}{n}$ and $b_n = \dfrac{1}{n}$, then $\sum_{n=1}^{\infty} b_n$ is the divergent harmonic series and $\lim_{n\to\infty}\dfrac{a_n}{b_n} = \lim_{n\to\infty}\ln n = \lim_{x\to\infty}\ln x = \infty$, so $\sum_{n=1}^{\infty} a_n$ diverges by part (a).

42. Let $a_n = \dfrac{1}{n^2}$ and $b_n = \dfrac{1}{n}$. Then $\lim_{n\to\infty}\dfrac{a_n}{b_n} = \lim_{n\to\infty}\dfrac{1}{n} = 0$, but $\sum b_n$ diverges while $\sum a_n$ converges.

43. $\lim_{n\to\infty} na_n = \lim_{n\to\infty}\dfrac{a_n}{1/n}$, so we apply the Limit Comparison Test with $b_n = \dfrac{1}{n}$. Since $\lim_{n\to\infty} na_n > 0$ we know that either both series converge or both series diverge, and we also know that $\sum_{n=1}^{\infty}\dfrac{1}{n}$ diverges [p-series with $p = 1$]. Therefore, $\sum a_n$ must be divergent.

44. First we observe that, by l'Hospital's Rule, $\lim_{x\to 0}\dfrac{\ln(1+x)}{x} = \lim_{x\to 0}\dfrac{1}{1+x} = 1$. Also, if $\sum a_n$ converges, then $\lim_{n\to\infty} a_n = 0$ by Theorem 11.2.6. Therefore, $\lim_{n\to\infty}\dfrac{\ln(1+a_n)}{a_n} = \lim_{x\to 0}\dfrac{\ln(1+x)}{x} = 1 > 0$. We are given that $\sum a_n$ is convergent and $a_n > 0$. Thus, $\sum \ln(1+a_n)$ is convergent by the Limit Comparison Test.

45. Yes. Since $\sum a_n$ is a convergent series with positive terms, $\lim_{n\to\infty} a_n = 0$ by Theorem 11.2.6, and $\sum b_n = \sum \sin(a_n)$ is a series with positive terms (for large enough n). We have $\lim_{n\to\infty}\dfrac{b_n}{a_n} = \lim_{n\to\infty}\dfrac{\sin(a_n)}{a_n} = 1 > 0$ by Theorem 2.4.2 [ET Theorem 3.3.2]. Thus, $\sum b_n$ is also convergent by the Limit Comparison Test.

46. Yes. Since $\sum a_n$ converges, its terms approach 0 as $n \to \infty$, so for some integer N, $a_n \leq 1$ for all $n \geq N$. But then $\sum_{n=1}^{\infty} a_n b_n = \sum_{n=1}^{N-1} a_n b_n + \sum_{n=N}^{\infty} a_n b_n \leq \sum_{n=1}^{N-1} a_n b_n + \sum_{n=N}^{\infty} b_n$. The first term is a finite sum, and the second term converges since $\sum_{n=1}^{\infty} b_n$ converges. So $\sum a_n b_n$ converges by the Comparison Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11.5 Alternating Series

1. (a) An alternating series is a series whose terms are alternately positive and negative.

   (b) An alternating series $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (-1)^{n-1} b_n$, where $b_n = |a_n|$, converges if $0 < b_{n+1} \le b_n$ for all n and $\lim_{n\to\infty} b_n = 0$. (This is the Alternating Series Test.)

   (c) The error involved in using the partial sum s_n as an approximation to the total sum s is the remainder $R_n = s - s_n$ and the size of the error is smaller than b_{n+1}; that is, $|R_n| \le b_{n+1}$. (This is the Alternating Series Estimation Theorem.)

2. $\frac{2}{3} - \frac{2}{5} + \frac{2}{7} - \frac{2}{9} + \frac{2}{11} - \cdots = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{2}{2n+1}$. Now $b_n = \frac{2}{2n+1} > 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

3. $-\frac{2}{5} + \frac{4}{6} - \frac{6}{7} + \frac{8}{8} - \frac{10}{9} + \cdots = \sum_{n=1}^{\infty} (-1)^n \frac{2n}{n+4}$. Now $\lim_{n\to\infty} b_n = \lim_{n\to\infty} \frac{2n}{n+4} = \lim_{n\to\infty} \frac{2}{1+4/n} = \frac{2}{1} \ne 0$. Since $\lim_{n\to\infty} a_n \ne 0$ (in fact the limit does not exist), the series diverges by the Test for Divergence.

4. $\frac{1}{\ln 3} - \frac{1}{\ln 4} + \frac{1}{\ln 5} - \frac{1}{\ln 6} + \frac{1}{\ln 7} - \cdots = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\ln(n+2)}$. Now $b_n = \frac{1}{\ln(n+2)} > 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

5. $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{3+5n} = \sum_{n=1}^{\infty} (-1)^{n-1} b_n$. Now $b_n = \frac{1}{3+5n} > 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

6. $\sum_{n=0}^{\infty} a_n = \sum_{n=0}^{\infty} \frac{(-1)^{n+1}}{\sqrt{n+1}} = \sum_{n=0}^{\infty} (-1)^{n+1} b_n$. Now $b_n = \frac{1}{\sqrt{n+1}} > 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

7. $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (-1)^n \frac{3n-1}{2n+1} = \sum_{n=1}^{\infty} (-1)^n b_n$. Now $\lim_{n\to\infty} b_n = \lim_{n\to\infty} \frac{3-1/n}{2+1/n} = \frac{3}{2} \ne 0$. Since $\lim_{n\to\infty} a_n \ne 0$ (in fact the limit does not exist), the series diverges by the Test for Divergence.

8. $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (-1)^n \frac{n^2}{n^2+n+1} = \sum_{n=1}^{\infty} (-1)^n b_n$. Now $\lim_{n\to\infty} b_n = \lim_{n\to\infty} \frac{n^2}{n^2+n+1} = \lim_{n\to\infty} \frac{1}{1+1/n+1/n^2} = 1 \ne 0$. Since $\lim_{n\to\infty} a_n \ne 0$, the series diverges by the Test for Divergence.

9. $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (-1)^n e^{-n} = \sum_{n=1}^{\infty} (-1)^n b_n$. Now $b_n = \frac{1}{e^n} > 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

10. $b_n = \dfrac{\sqrt{n}}{2n+3} > 0$ for $n \geq 1$. $\{b_n\}$ is decreasing for $n \geq 2$ since

$$\left(\frac{\sqrt{x}}{2x+3}\right)' = \frac{(2x+3)\left(\frac{1}{2}x^{-1/2}\right) - x^{1/2}(2)}{(2x+3)^2} = \frac{\frac{1}{2}x^{-1/2}[(2x+3)-4x]}{(2x+3)^2} = \frac{3-2x}{2\sqrt{x}\,(2x+3)^2} < 0 \text{ for } x > \tfrac{3}{2}.$$

Also, $\lim_{n\to\infty} b_n = \lim_{n\to\infty} \dfrac{\sqrt{n}/\sqrt{n}}{(2n+3)/\sqrt{n}} = \lim_{n\to\infty} \dfrac{1}{2\sqrt{n}+3/\sqrt{n}} = 0$. Thus, the series $\sum_{n=1}^{\infty} (-1)^n \dfrac{\sqrt{n}}{2n+3}$ converges by the Alternating Series Test.

11. $b_n = \dfrac{n^2}{n^3+4} > 0$ for $n \geq 1$. $\{b_n\}$ is decreasing for $n \geq 2$ since

$$\left(\frac{x^2}{x^3+4}\right)' = \frac{(x^3+4)(2x) - x^2(3x^2)}{(x^3+4)^2} = \frac{x(2x^3+8-3x^3)}{(x^3+4)^2} = \frac{x(8-x^3)}{(x^3+4)^2} < 0 \text{ for } x > 2. \text{ Also,}$$

$\lim_{n\to\infty} b_n = \lim_{n\to\infty} \dfrac{1/n}{1+4/n^3} = 0$. Thus, the series $\sum_{n=1}^{\infty} (-1)^{n+1} \dfrac{n^2}{n^3+4}$ converges by the Alternating Series Test.

12. $b_n = ne^{-n} = \dfrac{n}{e^n} > 0$ for $n \geq 1$. $\{b_n\}$ is decreasing for $n \geq 1$ since $(xe^{-x})' = x(-e^{-x}) + e^{-x} = e^{-x}(1-x) < 0$ for $x > 1$. Also, $\lim_{n\to\infty} b_n = 0$ since $\lim_{x\to\infty} \dfrac{x}{e^x} \overset{\text{H}}{=} \lim_{x\to\infty} \dfrac{1}{e^x} = 0$. Thus, the series $\sum_{n=1}^{\infty} (-1)^{n+1} ne^{-n}$ converges by the Alternating Series Test.

13. $\lim_{n\to\infty} b_n = \lim_{n\to\infty} e^{2/n} = e^0 = 1$, so $\lim_{n\to\infty} (-1)^{n-1} e^{2/n}$ does not exist. Thus, the series $\sum_{n=1}^{\infty} (-1)^{n-1} e^{2/n}$ diverges by the Test for Divergence.

14. $\lim_{n\to\infty} b_n = \lim_{n\to\infty} \arctan n = \frac{\pi}{2}$, so $\lim_{n\to\infty} (-1)^{n-1} \arctan n$ does not exist. Thus, the series $\sum_{n=1}^{\infty} (-1)^{n-1} \arctan n$ diverges by the Test for Divergence.

15. $a_n = \dfrac{\sin\left(n+\frac{1}{2}\right)\pi}{1+\sqrt{n}} = \dfrac{(-1)^n}{1+\sqrt{n}}$. Now $b_n = \dfrac{1}{1+\sqrt{n}} > 0$ for $n \geq 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series $\sum_{n=0}^{\infty} \dfrac{\sin\left(n+\frac{1}{2}\right)\pi}{1+\sqrt{n}}$ converges by the Alternating Series Test.

16. $a_n = \dfrac{n\cos n\pi}{2^n} = (-1)^n \dfrac{n}{2^n} = (-1)^n b_n$. $\{b_n\}$ is decreasing for $n \geq 2$ since

$(x2^{-x})' = x(-2^{-x}\ln 2) + 2^{-x} = 2^{-x}(1 - x\ln 2) < 0$ for $x > \dfrac{1}{\ln 2}$ $[\approx 1.4]$. Also, $\lim_{n\to\infty} b_n = 0$ since

$\lim_{x\to\infty} \dfrac{x}{2^x} \overset{\text{H}}{=} \lim_{x\to\infty} \dfrac{1}{2^x \ln 2} = 0$. Thus, the series $\sum_{n=1}^{\infty} \dfrac{n\cos n\pi}{2^n}$ converges by the Alternating Series Test.

17. $\sum_{n=1}^{\infty} (-1)^n \sin\left(\dfrac{\pi}{n}\right)$. $b_n = \sin\left(\dfrac{\pi}{n}\right) > 0$ for $n \geq 2$ and $\sin\left(\dfrac{\pi}{n}\right) > \sin\left(\dfrac{\pi}{n+1}\right)$, and $\lim_{n\to\infty} \sin\left(\dfrac{\pi}{n}\right) = \sin 0 = 0$, so the series converges by the Alternating Series Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $\sum_{n=1}^{\infty}(-1)^n\cos\left(\frac{\pi}{n}\right)$. $\lim_{n\to\infty}\cos\left(\frac{\pi}{n}\right)=\cos(0)=1$, so $\lim_{n\to\infty}(-1)^n\cos\left(\frac{\pi}{n}\right)$ does not exist and the series diverges by the Test for Divergence.

19. $\frac{n^n}{n!}=\frac{n\cdot n\cdot\cdots\cdot n}{1\cdot 2\cdot\cdots\cdot n}\geq n \;\Rightarrow\; \lim_{n\to\infty}\frac{n^n}{n!}=\infty \;\Rightarrow\; \lim_{n\to\infty}\frac{(-1)^n n^n}{n!}$ does not exist. So the series $\sum_{n=1}^{\infty}(-1)^n\frac{n^n}{n!}$ diverges by the Test for Divergence.

20. $b_n=\frac{\sqrt{n+1}-\sqrt{n}}{1}\cdot\frac{\sqrt{n+1}+\sqrt{n}}{\sqrt{n+1}+\sqrt{n}}=\frac{(n+1)-n}{\sqrt{n+1}+\sqrt{n}}=\frac{1}{\sqrt{n+1}+\sqrt{n}}>0$ for $n\geq 1$. $\{b_n\}$ is decreasing and $\lim_{n\to\infty}b_n=0$, so the series $\sum_{n-1}^{\infty}(-1)^n\left(\sqrt{n+1}-\sqrt{n}\right)$ converges by the Alternating Series Test.

21.



The graph gives us an estimate for the sum of the series $\sum_{n=1}^{\infty}\frac{(-0.8)^n}{n!}$ of -0.55.

$b_8=\frac{(0.8)^n}{8!}\approx 0.000\,004$, so

$$\sum_{n=1}^{\infty}\frac{(-0.8)^n}{n!}\approx s_7=\sum_{n=1}^{7}\frac{(-0.8)^n}{n!}$$

$$\approx -0.8+0.32-0.085\overline{3}+0.0170\overline{6}-0.002\,731+0.000\,364-0.000\,042\approx -0.5507$$

Adding b_8 to s_7 does not change the fourth decimal place of s_7, so the sum of the series, correct to four decimal places, is -0.5507.

22.



The graph gives us an estimate for the sum of the series $\sum_{n=1}^{\infty}(-1)^{n-1}\frac{n}{8^n}$ of 0.1.

$b_6=\frac{6}{8^6}\approx 0.000\,023$, so

$$\sum_{n=1}^{\infty}(-1)^{n-1}\frac{n}{8^n}\approx s_5=\sum_{n=1}^{5}(-1)^{n-1}\frac{n}{8^n}$$

$$\approx 0.125-0.03125+0.005\,859-0.000\,977+0.000\,153\approx 0.0988$$

Adding b_6 to s_5 does not change the fourth decimal place of s_5, so the sum of the series, correct to four decimal places, is 0.0988.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

23. The series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^6}$ satisfies (i) of the Alternating Series Test because $\frac{1}{(n+1)^6} < \frac{1}{n^6}$ and (ii) $\lim_{n\to\infty} \frac{1}{n^6} = 0$, so the series is convergent. Now $b_5 = \frac{1}{5^6} = 0.000064 > 0.00005$ and $b_6 = \frac{1}{6^6} \approx 0.00002 < 0.00005$, so by the Alternating Series Estimation Theorem, $n = 5$. (That is, since the 6th term is less than the desired error, we need to add the first 5 terms to get the sum to the desired accuracy.)

24. The series $\sum_{n=1}^{\infty} \frac{(-\frac{1}{3})^n}{n} = \sum_{n=1}^{\infty} (-1)^n \frac{1}{n3^n}$ satisfies (i) of the Alternating Series Test because $\frac{1}{(n+1)3^{n+1}} < \frac{1}{n3^n}$ and (ii) $\lim_{n\to\infty} \frac{1}{n3^n} = 0$, so the series is convergent. Now $b_5 = \frac{1}{5 \cdot 3^5} \approx 0.0008 > 0.0005$ and $b_6 = \frac{1}{6 \cdot 3^6} \approx 0.0002 < 0.0005$, so by the Alternating Series Estimation Theorem, $n = 5$. (That is, since the 6th term is less than the desired error, we need to add the first 5 terms to get the sum to the desired accuracy.)

25. The series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2 2^n}$ satisfies (i) of the Alternating Series Test because $\frac{1}{(n+1)^2 2^{n+1}} < \frac{1}{n^2 2^n}$ and (ii) $\lim_{n\to\infty} \frac{1}{n^2 2^n} = 0$, so the series is convergent. Now $b_5 = \frac{1}{5^2 2^5} = 0.00125 > 0.0005$ and $b_6 = \frac{1}{6^2 2^6} \approx 0.0004 < 0.0005$, so by the Alternating Series Estimation Theorem, $n = 5$. (That is, since the 6th term is less than the desired error, we need to add the first 5 terms to get the sum to the desired accuracy.)

26. The series $\sum_{n=1}^{\infty} \left(-\frac{1}{n}\right)^n = \sum_{n=1}^{\infty} (-1)^n \frac{1}{n^n}$ satisfies (i) of the Alternating Series Test because $\frac{1}{(n+1)^{n+1}} < \frac{1}{n^n}$ and (ii) $\lim_{n\to\infty} \frac{1}{n^n} = 0$, so the series is convergent. Now $b_5 = \frac{1}{5^5} = 0.00032 > 0.00005$ and $b_6 = \frac{1}{6^6} \approx 0.00002 < 0.00005$, so by the Alternating Series Estimation Theorem, $n = 5$. (That is, since the 6th term is less than the desired error, we need to add the first 5 terms to get the sum to the desired accuracy.)

27. $b_4 = \frac{1}{8!} = \frac{1}{40{,}320} \approx 0.000\,025$, so

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!} \approx s_3 = \sum_{n=1}^{3} \frac{(-1)^n}{(2n)!} = -\frac{1}{2} + \frac{1}{24} - \frac{1}{720} \approx -0.459\,722$$

Adding b_4 to s_3 does not change the fourth decimal place of s_3, so by the Alternating Series Estimation Theorem, the sum of the series, correct to four decimal places, is -0.4597.

28. $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^6} \approx s_9 = \frac{1}{1^6} - \frac{1}{2^6} + \frac{1}{3^6} - \frac{1}{4^6} + \frac{1}{5^6} - \frac{1}{6^6} + \frac{1}{7^6} - \frac{1}{8^6} + \frac{1}{9^6} \approx 0.985\,552$. Subtracting $b_{10} = 1/10^6$ from s_9 does not change the fourth decimal place of s_9, so by the Alternating Series Estimation Theorem, the sum of the series, correct to four decimal places, is 0.9856.

29. $\sum_{n=1}^{\infty} (-1)^n n e^{-2n} \approx s_5 = -\frac{1}{e^2} + \frac{2}{e^4} - \frac{3}{e^6} + \frac{4}{e^8} - \frac{5}{e^{10}} \approx -0.105\,025$. Adding $b_6 = 6/e^{12} \approx 0.000\,037$ to s_5 does not change the fourth decimal place of s_5, so by the Alternating Series Estimation Theorem, the sum of the series, correct to four decimal places, is -0.1050.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n4^n} \approx s_6 = \frac{1}{4} - \frac{1}{2\cdot 4^2} + \frac{1}{3\cdot 4^3} - \frac{1}{4\cdot 4^4} + \frac{1}{5\cdot 4^5} - \frac{1}{6\cdot 4^6} \approx 0.223136$. Adding $b_7 = \frac{1}{7\cdot 4^7} \approx 0.000\,0087$ to s_6 does not change the fourth decimal place of s_6, so by the Alternating Series Estimation Theorem, the sum of the series, correct to four decimal places, is 0.2231.

31. $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots + \frac{1}{49} - \frac{1}{50} + \frac{1}{51} - \frac{1}{52} + \cdots$. The 50th partial sum of this series is an underestimate, since $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = s_{50} + \left(\frac{1}{51} - \frac{1}{52}\right) + \left(\frac{1}{53} - \frac{1}{54}\right) + \cdots$, and the terms in parentheses are all positive. The result can be seen geometrically in Figure 1.

32. If $p > 0$, $\frac{1}{(n+1)^p} \le \frac{1}{n^p}$ ($\{1/n^p\}$ is decreasing) and $\lim_{n\to\infty} \frac{1}{n^p} = 0$, so the series converges by the Alternating Series Test. If $p \le 0$, $\lim_{n\to\infty} \frac{(-1)^{n-1}}{n^p}$ does not exist, so the series diverges by the Test for Divergence. Thus, $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^p}$ converges $\Leftrightarrow$ $p > 0$.

33. Clearly $b_n = \frac{1}{n+p}$ is decreasing and eventually positive and $\lim_{n\to\infty} b_n = 0$ for any p. So the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n+p}$ converges (by the Alternating Series Test) for any p for which every b_n is defined, that is, $n + p \neq 0$ for $n \ge 1$, or p is not a negative integer.

34. Let $f(x) = \frac{(\ln x)^p}{x}$. Then $f'(x) = \frac{(\ln x)^{p-1}(p - \ln x)}{x^2} < 0$ if $x > e^p$ so f is eventually decreasing for every p. Clearly $\lim_{n\to\infty} \frac{(\ln n)^p}{n} = 0$ if $p \le 0$, and if $p > 0$ we can apply l'Hospital's Rule $[\![p+1]\!]$ times to get a limit of 0 as well. So the series $\sum_{n=2}^{\infty} (-1)^{n-1} \frac{(\ln n)^p}{n}$ converges for all p (by the Alternating Series Test).

35. $\sum b_{2n} = \sum 1/(2n)^2$ clearly converges (by comparison with the p-series for $p = 2$). So suppose that $\sum (-1)^{n-1} b_n$ converges. Then by Theorem 11.2.8(ii), so does $\sum \left[(-1)^{n-1} b_n + b_n\right] = 2\left(1 + \frac{1}{3} + \frac{1}{5} + \cdots\right) = 2\sum \frac{1}{2n-1}$. But this diverges by comparison with the harmonic series, a contradiction. Therefore, $\sum (-1)^{n-1} b_n$ must diverge. The Alternating Series Test does not apply since $\{b_n\}$ is not decreasing.

36. (a) We will prove this by induction. Let $P(n)$ be the proposition that $s_{2n} = h_{2n} - h_n$. $P(1)$ is the statement $s_2 = h_2 - h_1$, which is true since $1 - \frac{1}{2} = \left(1 + \frac{1}{2}\right) - 1$. So suppose that $P(n)$ is true. We will show that $P(n+1)$ must be true as a consequence.

$$\begin{aligned} h_{2n+2} - h_{n+1} &= \left(h_{2n} + \frac{1}{2n+1} + \frac{1}{2n+2}\right) - \left(h_n + \frac{1}{n+1}\right) = (h_{2n} - h_n) + \frac{1}{2n+1} - \frac{1}{2n+2} \\ &= s_{2n} + \frac{1}{2n+1} - \frac{1}{2n+2} = s_{2n+2} \end{aligned}$$

which is $P(n+1)$, and proves that $s_{2n} = h_{2n} - h_n$ for all n.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) We know that $h_{2n} - \ln(2n) \to \gamma$ and $h_n - \ln n \to \gamma$ as $n \to \infty$. So

$s_{2n} = h_{2n} - h_n = [h_{2n} - \ln(2n)] - (h_n - \ln n) + [\ln(2n) - \ln n]$, and

$\lim_{n\to\infty} s_{2n} = \gamma - \gamma + \lim_{n\to\infty} [\ln(2n) - \ln n] = \lim_{n\to\infty} (\ln 2 + \ln n - \ln n) = \ln 2.$

11.6 Absolute Convergence and the Ratio and Root Tests

1. (a) Since $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = 8 > 1$, part (b) of the Ratio Test tells us that the series $\sum a_n$ is divergent.

(b) Since $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = 0.8 < 1$, part (a) of the Ratio Test tells us that the series $\sum a_n$ is absolutely convergent (and therefore convergent).

(c) Since $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = 1$, the Ratio Test fails and the series $\sum a_n$ might converge or it might diverge.

2. $b_n = \frac{1}{\sqrt{n}} > 0$ for $n \geq 1$, $\{b_n\}$ is decreasing for $n \geq 1$, and $\lim_{n\to\infty} b_n = 0$, so $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n}}$ converges by the Alternating Series Test. To determine absolute convergence, note that $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ diverges because it is a p-series with $p = \frac{1}{2} \leq 1$. Thus, the series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n}}$ is conditionally convergent.

3. $b_n = \frac{1}{5n+1} > 0$ for $n \geq 0$, $\{b_n\}$ is decreasing for $n \geq 0$, and $\lim_{n\to\infty} b_n = 0$, so $\sum_{n=0}^{\infty} \frac{(-1)^n}{5n+1}$ converges by the Alternating Series Test. To determine absolute convergence, choose $a_n = \frac{1}{n}$ to get

$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{1/n}{1/(5n+1)} = \lim_{n\to\infty} \frac{5n+1}{n} = 5 > 0$, so $\sum_{n=1}^{\infty} \frac{1}{5n+1}$ diverges by the Limit Comparison Test with the harmonic series. Thus, the series $\sum_{n=0}^{\infty} \frac{(-1)^n}{5n+1}$ is conditionally convergent.

4. $0 < \frac{1}{n^3+1} < \frac{1}{n^3}$ for $n \geq 1$ and $\sum_{n=1}^{\infty} \frac{1}{n^3}$ is a convergent p-series ($p = 3 > 1$), so $\sum_{n=1}^{\infty} \frac{1}{n^3+1}$ converges by comparison and the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^3+1}$ is absolutely convergent.

5. $0 < \left|\frac{\sin n}{2^n}\right| < \frac{1}{2^n}$ for $n \geq 1$ and $\sum_{n=1}^{\infty} \frac{1}{2^n}$ is a convergent geometric series ($r = \frac{1}{2} < 1$), so $\sum_{n=1}^{\infty} \left|\frac{\sin n}{2^n}\right|$ converges by comparison and the series $\sum_{n=1}^{\infty} \frac{\sin n}{2^n}$ is absolutely convergent.

6. $b_n = \frac{n}{n^2+4} > 0$ for $n \geq 1$, $\{b_n\}$ is decreasing for $n \geq 2$, and $\lim_{n\to\infty} b_n = 0$, so $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n}{n^2+4}$ converges by the Alternating Series Test. To determine absolute convergence, choose $a_n = \frac{1}{n}$ to get

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\lim_{n\to\infty}\dfrac{a_n}{b_n}=\lim_{n\to\infty}\dfrac{1/n}{n/(n^2+4)}=\lim_{n\to\infty}\dfrac{n^2+4}{n^2}=\lim_{n\to\infty}\dfrac{1+4/n^2}{1}=1>0$, so $\sum_{n=1}^{\infty}\dfrac{n}{n^2+4}$ diverges by the Limit Comparison Test with the harmonic series. Thus, the series $\sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{n}{n^2+4}$ is conditionally convergent.

7. $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\dfrac{n+1}{5^{n+1}}\cdot\dfrac{5^n}{n}\right|=\lim_{n\to\infty}\left|\dfrac{1}{5}\cdot\dfrac{n+1}{n}\right|=\dfrac{1}{5}\lim_{n\to\infty}\dfrac{1+1/n}{1}=\dfrac{1}{5}(1)=\dfrac{1}{5}<1$, so the series $\sum_{n=1}^{\infty}\dfrac{n}{5^n}$ is absolutely convergent by the Ratio Test.

8. $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\dfrac{(-2)^{n+1}}{(n+1)^2}\cdot\dfrac{n^2}{(-2)^n}\right|=\lim_{n\to\infty}\left|(-2)\dfrac{n^2}{(n+1)^2}\right|=2\lim_{n\to\infty}\dfrac{1}{(1+1/n)^2}=2(1)=2>1$, so the series $\sum_{n=1}^{\infty}\dfrac{(-2)^n}{n^2}$ is divergent by the Ratio Test.

9. $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\dfrac{(-1)^n3^{n+1}}{2^{n+1}(n+1)^3}\cdot\dfrac{2^nn^3}{(-1)^{n-1}3^n}\right|=\lim_{n\to\infty}\left|\left(-\dfrac{3}{2}\right)\dfrac{n^3}{(n+1)^3}\right|=\dfrac{3}{2}\lim_{n\to\infty}\dfrac{1}{(1+1/n)^3}=\dfrac{3}{2}(1)=\dfrac{3}{2}>1$, so the series $\sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{3^n}{2^nn^3}$ is divergent by the Ratio Test.

10. $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\dfrac{(-3)^{n+1}}{[2(n+1)+1]!}\cdot\dfrac{(2n+1)!}{(-3)^n}\right|=\lim_{n\to\infty}\left|(-3)\dfrac{1}{(2n+3)(2n+2)}\right|=3\lim_{n\to\infty}\dfrac{1}{(2n+3)(2n+2)}$

$=3(0)=0<1$

so the series $\sum_{n=0}^{\infty}\dfrac{(-3)^n}{(2n+1)!}$ is absolutely convergent by the Ratio Test.

11. $\lim_{k\to\infty}\left|\dfrac{a_{k+1}}{a_k}\right|=\lim_{k\to\infty}\left|\dfrac{1}{(k+1)!}\cdot\dfrac{k!}{1}\right|=\lim_{k\to\infty}\dfrac{1}{k+1}=0<1$, so the series $\sum_{k=1}^{\infty}\dfrac{1}{k!}$ is absolutely convergent by the Ratio Test. Since the terms of this series are positive, absolute convergence is the same as convergence.

12. $\lim_{k\to\infty}\left|\dfrac{a_{k+1}}{a_k}\right|=\lim_{k\to\infty}\left|\dfrac{(k+1)e^{-(k+1)}}{ke^{-k}}\right|=\lim_{k\to\infty}\left(\dfrac{k+1}{k}\cdot e^{-1}\right)=\dfrac{1}{e}\lim_{k\to\infty}\dfrac{1+1/k}{1}=\dfrac{1}{e}(1)=\dfrac{1}{e}<1$, so the series $\sum_{k=1}^{\infty}ke^{-k}$ is absolutely convergent by the Ratio Test. Since the terms of this series are positive, absolute convergence is the same as convergence.

13. $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left[\dfrac{10^{n+1}}{(n+2)\,4^{2n+3}}\cdot\dfrac{(n+1)\,4^{2n+1}}{10^n}\right]=\lim_{n\to\infty}\left(\dfrac{10}{4^2}\cdot\dfrac{n+1}{n+2}\right)=\dfrac{5}{8}<1$, so the series $\sum_{n=1}^{\infty}\dfrac{10^n}{(n+1)\,4^{2n+1}}$ is absolutely convergent by the Ratio Test. Since the terms of this series are positive, absolute convergence is the same as convergence.

14. $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left[\dfrac{(n+1)!}{100^{n+1}}\cdot\dfrac{100^n}{n!}\right]=\lim_{n\to\infty}\dfrac{n+1}{100}=\infty$, so the series $\sum_{n=1}^{\infty}\dfrac{n!}{100^n}$ diverges by the Ratio Test.

15. $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\dfrac{(n+1)\pi^{n+1}}{(-3)^n}\cdot\dfrac{(-3)^{n-1}}{n\pi^n}\right|=\lim_{n\to\infty}\left|\dfrac{\pi}{-3}\cdot\dfrac{n+1}{n}\right|=\dfrac{\pi}{3}\lim_{n\to\infty}\dfrac{1+1/n}{1}=\dfrac{\pi}{3}(1)=\dfrac{\pi}{3}>1$, so the series $\sum_{n=1}^{\infty}\dfrac{n\pi^n}{(-3)^{n-1}}$ diverges by the Ratio Test. *Or:* Since $\lim_{n\to\infty}|a_n|=\infty$, the series diverges by the Test for Divergence.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

16. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(n+1)^{10}}{(-10)^{n+2}}\cdot\frac{(-10)^{n+1}}{n^{10}}\right| = \lim_{n\to\infty}\left|\frac{1}{-10}\left(\frac{n+1}{n}\right)^{10}\right| = \frac{1}{10}\lim_{n\to\infty}\left(1+\frac{1}{n}\right)^{10} = \frac{1}{10}(1) = \frac{1}{10} < 1,$

so the series $\sum_{n=1}^{\infty}\frac{n^{10}}{(-10)^{n+1}}$ is absolutely convergent by the Ratio Test.

17. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{\cos[(n+1)\pi/3]}{(n+1)!}\cdot\frac{n!}{\cos(n\pi/3)}\right| = \lim_{n\to\infty}\left|\frac{\cos[(n+1)\pi/3]}{(n+1)\cos(n\pi/3)}\right| = \lim_{n\to\infty}\frac{c}{n+1} = 0 < 1$ (where $0 < c \le 2$ for all positive integers n), so the series $\sum_{n=1}^{\infty}\frac{\cos(n\pi/3)}{n!}$ is absolutely convergent by the Ratio Test.

18. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(n+1)!}{(n+1)^{n+1}}\cdot\frac{n^n}{n!}\right| = \lim_{n\to\infty}\frac{(n+1)n^n}{(n+1)^{n+1}} = \lim_{n\to\infty}\frac{n^n}{(n+1)^n} = \lim_{n\to\infty}\frac{1}{(1+1/n)^n} = \frac{1}{e} < 1$, so the series $\sum_{n=1}^{\infty}\frac{n!}{n^n}$ is absolutely convergent by the Ratio Test.

19. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(n+1)^{100}100^{n+1}}{(n+1)!}\cdot\frac{n!}{n^{100}100^n}\right| = \lim_{n\to\infty}\frac{100}{n+1}\left(\frac{n+1}{n}\right)^{100} = \lim_{n\to\infty}\frac{100}{n+1}\left(1+\frac{1}{n}\right)^{100}$

$= 0\cdot 1 = 0 < 1$

so the series $\sum_{n=1}^{\infty}\frac{n^{100}100^n}{n!}$ is absolutely convergent by the Ratio Test.

20. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{[2(n+1)]!}{[(n+1)!]^2}\cdot\frac{(n!)^2}{(2n)!}\right| = \lim_{n\to\infty}\frac{(2n+2)(2n+1)}{(n+1)(n+1)} = \lim_{n\to\infty}\frac{(2+2/n)(2+1/n)}{(1+1/n)(1+1/n)} = \frac{2\cdot 2}{1\cdot 1} = 4 > 1,$

so the series $\sum_{n=1}^{\infty}\frac{(2n)!}{(n!)^2}$ diverges by the Ratio Test.

21. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^n(n+1)!}{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)(2n+1)}\cdot\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{(-1)^{n-1}n!}\right| = \lim_{n\to\infty}\frac{n+1}{2n+1}$

$= \lim_{n\to\infty}\frac{1+1/n}{2+1/n} = \frac{1}{2} < 1,$

so the series $1 - \frac{2!}{1\cdot 3} + \frac{3!}{1\cdot 3\cdot 5} - \frac{4!}{1\cdot 3\cdot 5\cdot 7} + \cdots + (-1)^{n-1}\frac{n!}{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)} + \cdots$ is absolutely convergent by the Ratio Test.

22. $\frac{2}{3} + \frac{2\cdot 5}{3\cdot 5} + \frac{2\cdot 5\cdot 8}{3\cdot 5\cdot 7} + \frac{2\cdot 5\cdot 8\cdot 11}{3\cdot 5\cdot 7\cdot 9} + \cdots = \sum_{n=1}^{\infty}\frac{2\cdot 5\cdot 8\cdot 11\cdot\cdots\cdot(3n-1)}{3\cdot 5\cdot 7\cdot 9\cdot\cdots\cdot(2n+1)}$.

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{2\cdot 5\cdot 8\cdot\cdots\cdot(3n-1)(3n+2)}{3\cdot 5\cdot 7\cdot\cdots\cdot(2n+1)(2n+3)}\cdot\frac{3\cdot 5\cdot 7\cdot\cdots\cdot(2n+1)}{2\cdot 5\cdot 8\cdot\cdots\cdot(3n-1)}\right|$$

$$= \lim_{n\to\infty}\frac{3n+2}{2n+3} = \lim_{n\to\infty}\frac{3+2/n}{2+3/n} = \frac{3}{2} > 1,$$

so the given series diverges by the Ratio Test.

23. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)(2n+2)}{(n+1)!}\cdot\frac{n!}{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}\right| = \lim_{n\to\infty}\frac{2n+2}{n+1} = \lim_{n\to\infty}\frac{2(n+1)}{n+1} = 2 > 1$, so the series $\sum_{n=1}^{\infty}\frac{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}{n!}$ diverges by the Ratio Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{2^{n+1}\,(n+1)!}{5\cdot 8\cdot 11\cdot\cdots\cdot(3n+2)\,(3n+5)}\cdot\frac{5\cdot 8\cdot 11\cdot\cdots\cdot(3n+2)}{2^n n!}\right| = \lim_{n\to\infty}\frac{2(n+1)}{3n+5} = \frac{2}{3} < 1$, so the series $\sum_{n=1}^{\infty}(-1)^n\frac{2^n n!}{5\cdot 8\cdot 11\cdot\cdots\cdot(3n+2)}$ is absolutely convergent by the Ratio Test.

25. $\lim_{n\to\infty}\sqrt[n]{|a_n|} = \lim_{n\to\infty}\frac{n^2+1}{2n^2+1} = \lim_{n\to\infty}\frac{1+1/n^2}{2+1/n^2} = \frac{1}{2} < 1$, so the series $\sum_{n=1}^{\infty}\left(\frac{n^2+1}{2n^2+1}\right)^n$ is absolutely convergent by the Root Test.

26. $\lim_{n\to\infty}\sqrt[n]{|a_n|} = \lim_{n\to\infty}\sqrt[n]{\left|\frac{(-2)^n}{n^n}\right|} = \lim_{n\to\infty}\frac{2}{n} = 0 < 1$, so the series $\sum_{n=1}^{\infty}\frac{(-2)^n}{n^n}$ is absolutely convergent by the Root Test.

27. $\lim_{n\to\infty}\sqrt[n]{|a_n|} = \lim_{n\to\infty}\sqrt[n]{\left|\frac{(-1)^{n-1}}{(\ln n)^n}\right|} = \lim_{n\to\infty}\frac{1}{\ln n} = 0 < 1$, so the series $\sum_{n=2}^{\infty}\frac{(-1)^{n-1}}{(\ln n)^n}$ is absolutely convergent by the Root Test.

28. $\lim_{n\to\infty}\sqrt[n]{|a_n|} = \lim_{n\to\infty}\sqrt[n]{\left|\left(\frac{-2n}{n+1}\right)^{5n}\right|} = \lim_{n\to\infty}\frac{2^5 n^5}{(n+1)^5} = 32\lim_{n\to\infty}\frac{1}{\left(\frac{n+1}{n}\right)^5} = 32\lim_{n\to\infty}\frac{1}{(1+1/n)^5}$

$= 32(1) = 32 > 1,$

so the series $\sum_{n=1}^{\infty}\left(\frac{-2n}{n+1}\right)^{5n}$ diverges by the Root Test.

29. $\lim_{n\to\infty}\sqrt[n]{|a_n|} = \lim_{n\to\infty}\sqrt[n]{\left(1+\frac{1}{n}\right)^{n^2}} = \lim_{n\to\infty}\left(1+\frac{1}{n}\right)^n = e > 1$ [by Equation 6.4.9 (or 6.4*.9) [ET 3.6.6]], so the series $\sum_{n=1}^{\infty}\left(1+\frac{1}{n}\right)^{n^2}$ diverges by the Root Test.

30. $\lim_{n\to\infty}\sqrt[n]{|a_n|} = \lim_{n\to\infty}\sqrt[n]{|(\arctan n)^n|} = \lim_{n\to\infty}\arctan n = \frac{\pi}{2} > 1$, so the series $\sum_{n=0}^{\infty}(\arctan n)^n$ diverges by the Root Test.

31. $\sum_{n=2}^{\infty}\frac{(-1)^n}{\ln n}$ converges by the Alternating Series Test since $\lim_{n\to\infty}\frac{1}{\ln n} = 0$ and $\left\{\frac{1}{\ln n}\right\}$ is decreasing. Now $\ln n < n$, so $\frac{1}{\ln n} > \frac{1}{n}$, and since $\sum_{n=2}^{\infty}\frac{1}{n}$ is the divergent (partial) harmonic series, $\sum_{n=2}^{\infty}\frac{1}{\ln n}$ diverges by the Comparison Test. Thus, $\sum_{n=2}^{\infty}\frac{(-1)^n}{\ln n}$ is conditionally convergent.

32. $\lim_{n\to\infty}\sqrt[n]{|a_n|} = \lim_{n\to\infty}\sqrt[n]{\left|\left(\frac{1-n}{2+3n}\right)^n\right|} = \lim_{n\to\infty}\frac{n-1}{3n+2} = \lim_{n\to\infty}\frac{1-1/n}{3+2/n} = \frac{1}{3} < 1$, so the series $\sum_{n=1}^{\infty}\left(\frac{1-n}{2+3n}\right)^n$ is absolutely convergent by the Root Test.

33. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-9)^{n+1}}{(n+1)10^{n+2}}\cdot\frac{n10^{n+1}}{(-9)^n}\right| = \lim_{n\to\infty}\left|\frac{(-9)n}{10(n+1)}\right| = \frac{9}{10}\lim_{n\to\infty}\frac{1}{1+1/n} = \frac{9}{10}(1) = \frac{9}{10} < 1$, so the series $\sum_{n=1}^{\infty}\frac{(-9)^n}{n10^{n+1}}$ is absolutely convergent by the Ratio Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

34. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(n+1)5^{2n+2}}{10^{n+2}}\cdot\frac{10^{n+1}}{n5^{2n}}\right| = \lim_{n\to\infty}\frac{5^2(n+1)}{10n} = \frac{5}{2}\lim_{n\to\infty}\left(1+\frac{1}{n}\right) = \frac{5}{2}(1) = \frac{5}{2} > 1$, so the series $\sum_{n=1}^{\infty}\frac{n5^{2n}}{10^{n+1}}$ diverges by the Ratio Test. *Or:* Since $\lim_{n\to\infty} a_n = \infty$, the series diverges by the Test for Divergence.

35. $\lim_{n\to\infty}\sqrt[n]{|a_n|} = \lim_{n\to\infty}\sqrt[n]{\left|\left(\frac{n}{\ln n}\right)^n\right|} = \lim_{n\to\infty}\frac{n}{\ln n} = \lim_{x\to\infty}\frac{x}{\ln x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{1}{1/x} = \lim_{x\to\infty} x = \infty$, so the series $\sum_{n=2}^{\infty}\left(\frac{n}{\ln n}\right)^n$ diverges by the Root Test.

36. $\left|\frac{\sin(n\pi/6)}{1+n\sqrt{n}}\right| \le \frac{1}{1+n\sqrt{n}} < \frac{1}{n^{3/2}}$, so the series $\sum_{n=1}^{\infty}\frac{\sin(n\pi/6)}{1+n\sqrt{n}}$ converges by comparison with the convergent p-series $\sum_{n=1}^{\infty}\frac{1}{n^{3/2}}$ $\left(p=\frac{3}{2}>1\right)$. It follows that the given series is absolutely convergent.

37. $\left|\frac{(-1)^n\arctan n}{n^2}\right| < \frac{\pi/2}{n^2}$, so since $\sum_{n=1}^{\infty}\frac{\pi/2}{n^2} = \frac{\pi}{2}\sum_{n=1}^{\infty}\frac{1}{n^2}$ converges $(p=2>1)$, the given series $\sum_{n=1}^{\infty}\frac{(-1)^n\arctan n}{n^2}$ converges absolutely by the Comparison Test.

38. The function $f(x) = \frac{1}{x\ln x}$ is continuous, positive, and decreasing on $[2,\infty)$.

$\int_2^{\infty}\frac{1}{x\ln x}\,dx = \lim_{t\to\infty}\int_2^t\frac{1}{x\ln x}\,dx = \lim_{t\to\infty}[\ln(\ln x)]_2^t = \lim_{t\to\infty}[(\ln(\ln t) - \ln(\ln 2)] = \infty$, so the series $\sum_{n=2}^{\infty}\frac{(-1)^n}{n\ln n}$ diverges by the Integral Test. Now $\{b_n\} = \left\{\frac{1}{n\ln n}\right\}$ with $n\ge 2$ is a decreasing sequence of positive terms and $\lim_{n\to\infty} b_n = 0$. Thus, $\sum_{n=2}^{\infty}\frac{(-1)^n}{n\ln n}$ converges by the Alternating Series Test. It follows that $\sum_{n=2}^{\infty}\frac{(-1)^n}{n\ln n}$ is conditionally convergent.

39. By the recursive definition, $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{5n+1}{4n+3}\right| = \frac{5}{4} > 1$, so the series diverges by the Ratio Test.

40. By the recursive definition, $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{2+\cos n}{\sqrt{n}}\right| = 0 < 1$, so the series converges absolutely by the Ratio Test.

41. The series $\sum_{n=1}^{\infty}\frac{b_n^n\cos n\pi}{n} = \sum_{n=1}^{\infty}(-1)^n\frac{b_n^n}{n}$, where $b_n > 0$ for $n\ge 1$ and $\lim_{n\to\infty} b_n = \frac{1}{2}$.

$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^{n+1}b_n^{n+1}}{n+1}\cdot\frac{n}{(-1)^n b_n^n}\right| = \lim_{n\to\infty} b_n\frac{n}{n+1} = \frac{1}{2}(1) = \frac{1}{2} < 1$, so the series $\sum_{n=1}^{\infty}\frac{b_n^n\cos n\pi}{n}$ is absolutely convergent by the Ratio Test.

42. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^{n+1}(n+1)!}{(n+1)^{n+1}b_1b_2\cdots b_nb_{n+1}}\cdot\frac{n^nb_1b_2\cdots b_n}{(-1)^n\,n!}\right| = \lim_{n\to\infty}\left|\frac{(-1)(n+1)n^n}{b_{n+1}(n+1)^{n+1}}\right| = \lim_{n\to\infty}\frac{n^n}{b_{n+1}(n+1)^n}$

$= \lim_{n\to\infty}\frac{1}{b_{n+1}}\left(\frac{n}{n+1}\right)^n = \lim_{n\to\infty}\frac{1}{b_{n+1}}\left(\frac{1}{1+1/n}\right)^n = \lim_{n\to\infty}\frac{1}{b_{n+1}(1+1/n)^n} = \frac{1}{\frac{1}{2}e} = \frac{2}{e} < 1$

so the series $\sum_{n=1}^{\infty}\frac{(-1)^n\,n!}{n^nb_1b_2b_3\cdots b_n}$ is absolutely convergent by the Ratio Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. (a) $\lim_{n\to\infty}\left|\dfrac{1/(n+1)^3}{1/n^3}\right| = \lim_{n\to\infty}\dfrac{n^3}{(n+1)^3} = \lim_{n\to\infty}\dfrac{1}{(1+1/n)^3} = 1$. Inconclusive

(b) $\lim_{n\to\infty}\left|\dfrac{(n+1)}{2^{n+1}}\cdot\dfrac{2^n}{n}\right| = \lim_{n\to\infty}\dfrac{n+1}{2n} = \lim_{n\to\infty}\left(\dfrac{1}{2}+\dfrac{1}{2n}\right) = \dfrac{1}{2}$. Conclusive (convergent)

(c) $\lim_{n\to\infty}\left|\dfrac{(-3)^n}{\sqrt{n+1}}\cdot\dfrac{\sqrt{n}}{(-3)^{n-1}}\right| = 3\lim_{n\to\infty}\sqrt{\dfrac{n}{n+1}} = 3\lim_{n\to\infty}\sqrt{\dfrac{1}{1+1/n}} = 3$. Conclusive (divergent)

(d) $\lim_{n\to\infty}\left|\dfrac{\sqrt{n+1}}{1+(n+1)^2}\cdot\dfrac{1+n^2}{\sqrt{n}}\right| = \lim_{n\to\infty}\left[\sqrt{1+\dfrac{1}{n}}\cdot\dfrac{1/n^2+1}{1/n^2+(1+1/n)^2}\right] = 1$. Inconclusive

44. We use the Ratio Test:

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{[(n+1)!]^2/[k(n+1)]!}{(n!)^2/(kn)!}\right| = \lim_{n\to\infty}\left|\frac{(n+1)^2}{[k(n+1)]\,[k(n+1)-1]\cdots[kn+1]}\right|$$

Now if $k=1$, then this is equal to $\lim_{n\to\infty}\left|\dfrac{(n+1)^2}{(n+1)}\right| = \infty$, so the series diverges; if $k=2$, the limit is

$\lim_{n\to\infty}\left|\dfrac{(n+1)^2}{(2n+2)(2n+1)}\right| = \dfrac{1}{4} < 1$, so the series converges, and if $k>2$, then the highest power of n in the denominator is larger than 2, and so the limit is 0, indicating convergence. So the series converges for $k \geq 2$.

45. (a) $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\dfrac{x^{n+1}}{(n+1)!}\cdot\dfrac{n!}{x^n}\right| = \lim_{n\to\infty}\left|\dfrac{x}{n+1}\right| = |x|\lim_{n\to\infty}\dfrac{1}{n+1} = |x|\cdot 0 = 0 < 1$, so by the Ratio Test the series $\sum_{n=0}^{\infty}\dfrac{x^n}{n!}$ converges for all x.

(b) Since the series of part (a) always converges, we must have $\lim_{n\to\infty}\dfrac{x^n}{n!} = 0$ by Theorem 11.2.6.

46. (a) $R_n = a_{n+1}+a_{n+2}+a_{n+3}+a_{n+4}+\cdots = a_{n+1}\left(1+\dfrac{a_{n+2}}{a_{n+1}}+\dfrac{a_{n+3}}{a_{n+1}}+\dfrac{a_{n+4}}{a_{n+1}}+\cdots\right)$

$$= a_{n+1}\left(1+\frac{a_{n+2}}{a_{n+1}}+\frac{a_{n+3}}{a_{n+2}}\frac{a_{n+2}}{a_{n+1}}+\frac{a_{n+4}}{a_{n+3}}\frac{a_{n+3}}{a_{n+2}}\frac{a_{n+2}}{a_{n+1}}+\cdots\right)$$

$$= a_{n+1}(1+r_{n+1}+r_{n+2}r_{n+1}+r_{n+3}r_{n+2}r_{n+1}+\cdots) \quad (\star)$$

$$\leq a_{n+1}(1+r_{n+1}+r_{n+1}^2+r_{n+1}^3+\cdots) \quad [\text{since } \{r_n\} \text{ is decreasing}] \quad = \frac{a_{n+1}}{1-r_{n+1}}$$

(b) Note that since $\{r_n\}$ is increasing and $r_n \to L$ as $n\to\infty$, we have $r_n < L$ for all n. So, starting with equation $(\star)$,

$$R_n = a_{n+1}(1+r_{n+1}+r_{n+2}r_{n+1}+r_{n+3}r_{n+2}r_{n+1}+\cdots) \leq a_{n+1}(1+L+L^2+L^3+\cdots) = \frac{a_{n+1}}{1-L}.$$

47. (a) $s_5 = \sum_{n=1}^{5}\dfrac{1}{n2^n} = \dfrac{1}{2}+\dfrac{1}{8}+\dfrac{1}{24}+\dfrac{1}{64}+\dfrac{1}{160} = \dfrac{661}{960} \approx 0.68854$. Now the ratios

$r_n = \dfrac{a_{n+1}}{a_n} = \dfrac{n2^n}{(n+1)2^{n+1}} = \dfrac{n}{2(n+1)}$ form an increasing sequence, since

$r_{n+1}-r_n = \dfrac{n+1}{2(n+2)} - \dfrac{n}{2(n+1)} = \dfrac{(n+1)^2-n(n+2)}{2(n+1)(n+2)} = \dfrac{1}{2(n+1)(n+2)} > 0$. So by Exercise 46(b), the error in using s_5 is $R_5 \leq \dfrac{a_6}{1-\lim_{n\to\infty} r_n} = \dfrac{1/(6\cdot 2^6)}{1-1/2} = \dfrac{1}{192} \approx 0.00521$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) The error in using s_n as an approximation to the sum is $R_n = \dfrac{a_{n+1}}{1-\frac{1}{2}} = \dfrac{2}{(n+1)2^{n+1}}$. We want $R_n < 0.00005 \quad \Leftrightarrow$

$\dfrac{1}{(n+1)2^n} < 0.00005 \quad \Leftrightarrow \quad (n+1)2^n > 20{,}000$. To find such an n we can use trial and error or a graph. We calculate

$(11+1)2^{11} = 24{,}576$, so $s_{11} = \sum_{n=1}^{11} \dfrac{1}{n2^n} \approx 0.693109$ is within 0.00005 of the actual sum.

48. $s_{10} = \sum_{n=1}^{10} \dfrac{n}{2^n} = \dfrac{1}{2} + \dfrac{2}{4} + \dfrac{3}{8} + \cdots + \dfrac{10}{1024} \approx 1.988$. The ratios $r_n = \dfrac{a_{n+1}}{a_n} = \dfrac{n+1}{2^{n+1}} \cdot \dfrac{2^n}{n} = \dfrac{n+1}{2n} = \dfrac{1}{2}\left(1 + \dfrac{1}{n}\right)$ form a decreasing sequence, and $r_{11} = \dfrac{11+1}{2(11)} = \dfrac{12}{22} = \dfrac{6}{11} < 1$, so by Exercise 46(a), the error in using s_{10} to approximate the sum of the series $\sum_{n=1}^{\infty} \dfrac{n}{2^n}$ is $R_{10} \le \dfrac{a_{11}}{1-r_{11}} = \dfrac{\frac{11}{2048}}{1-\frac{6}{11}} = \dfrac{121}{10{,}240} \approx 0.0118$.

49. (i) Following the hint, we get that $|a_n| < r^n$ for $n > N$, and so since the geometric series $\sum_{n=1}^{\infty} r^n$ converges $[0 < r < 1]$, the series $\sum_{n=N}^{\infty} |a_n|$ converges as well by the Comparison Test, and hence so does $\sum_{n=1}^{\infty} |a_n|$, so $\sum_{n=1}^{\infty} a_n$ is absolutely convergent.

(ii) If $\lim_{n\to\infty} \sqrt[n]{|a_n|} = L > 1$, then there is an integer N such that $\sqrt[n]{|a_n|} > 1$ for all $n \ge N$, so $|a_n| > 1$ for $n \ge N$. Thus, $\lim_{n\to\infty} a_n \ne 0$, so $\sum_{n=1}^{\infty} a_n$ diverges by the Test for Divergence.

(iii) Consider $\sum_{n=1}^{\infty} \dfrac{1}{n}$ [diverges] and $\sum_{n=1}^{\infty} \dfrac{1}{n^2}$ [converges]. For each sum, $\lim_{n\to\infty} \sqrt[n]{|a_n|} = 1$, so the Root Test is inconclusive.

50. (a) $\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\dfrac{[4(n+1)]!\,[1103+26{,}390(n+1)]}{[(n+1)!]^4\,396^{4(n+1)}} \cdot \dfrac{(n!)^4\,396^{4n}}{(4n)!\,(1103+26{,}390n)}\right|$

$$= \lim_{n\to\infty} \frac{(4n+4)(4n+3)(4n+2)(4n+1)(26{,}390n+27{,}493)}{(n+1)^4\,396^4\,(26{,}390n+1103)} = \frac{4^4}{396^4} = \frac{1}{99^4} < 1,$$

so by the Ratio Test, the series $\sum_{n=0}^{\infty} \dfrac{(4n)!\,(1103+26{,}390n)}{(n!)^4\,396^{4n}}$ converges.

(b) $\dfrac{1}{\pi} = \dfrac{2\sqrt{2}}{9801} \sum_{n=0}^{\infty} \dfrac{(4n)!\,(1103+26{,}390n)}{(n!)^4\,396^{4n}}$

With the first term $(n=0)$, $\dfrac{1}{\pi} \approx \dfrac{2\sqrt{2}}{9801} \cdot \dfrac{1103}{1} \quad \Rightarrow \quad \pi \approx 3.141\,592\,73$, so we get 6 correct decimal places of π, which is 3.141 592 653 589 793 238 to 18 decimal places.

With the second term $(n=1)$, $\dfrac{1}{\pi} \approx \dfrac{2\sqrt{2}}{9801}\left(\dfrac{1103}{1} + \dfrac{4!\,(1103+26{,}390)}{396^4}\right) \quad \Rightarrow \quad \pi \approx 3.141\,592\,653\,589\,793\,878$, so we get 15 correct decimal places of π.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. (a) Since $\sum a_n$ is absolutely convergent, and since $|a_n^+| \le |a_n|$ and $|a_n^-| \le |a_n|$ (because a_n^+ and a_n^- each equal either a_n or 0), we conclude by the Comparison Test that both $\sum a_n^+$ and $\sum a_n^-$ must be absolutely convergent.

Or: Use Theorem 11.2.8.

(b) We will show by contradiction that both $\sum a_n^+$ and $\sum a_n^-$ must diverge. For suppose that $\sum a_n^+$ converged. Then so would $\sum\left(a_n^+ - \frac{1}{2}a_n\right)$ by Theorem 11.2.8. But $\sum\left(a_n^+ - \frac{1}{2}a_n\right) = \sum\left[\frac{1}{2}\left(a_n + |a_n|\right) - \frac{1}{2}a_n\right] = \frac{1}{2}\sum|a_n|$, which diverges because $\sum a_n$ is only conditionally convergent. Hence, $\sum a_n^+$ can't converge. Similarly, neither can $\sum a_n^-$.

52. Let $\sum b_n$ be the rearranged series constructed in the hint. [This series can be constructed by virtue of the result of Exercise 51(b).] This series will have partial sums s_n that oscillate in value back and forth across r. Since $\lim_{n\to\infty} a_n = 0$ (by Theorem 11.2.6), and since the size of the oscillations $|s_n - r|$ is always less than $|a_n|$ because of the way $\sum b_n$ was constructed, we have that $\sum b_n = \lim_{n\to\infty} s_n = r$.

53. Suppose that $\sum a_n$ is conditionally convergent.

(a) $\sum n^2 a_n$ is divergent: Suppose $\sum n^2 a_n$ converges. Then $\lim_{n\to\infty} n^2 a_n = 0$ by Theorem 6 in Section 11.2, so there is an integer $N > 0$ such that $n > N \;\Rightarrow\; n^2\,|a_n| < 1$. For $n > N$, we have $|a_n| < \dfrac{1}{n^2}$, so $\sum_{n>N} |a_n|$ converges by comparison with the convergent p-series $\sum_{n>N} \dfrac{1}{n^2}$. In other words, $\sum a_n$ converges absolutely, contradicting the assumption that $\sum a_n$ is conditionally convergent. This contradiction shows that $\sum n^2 a_n$ diverges.

Remark: The same argument shows that $\sum n^p a_n$ diverges for any $p > 1$.

(b) $\displaystyle\sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln n}$ is conditionally convergent. It converges by the Alternating Series Test, but does not converge absolutely

[by the Integral Test, since the function $f(x) = \dfrac{1}{x \ln x}$ is continuous, positive, and decreasing on $[2, \infty)$ and $\displaystyle\int_2^{\infty} \frac{dx}{x \ln x} = \lim_{t\to\infty} \int_2^t \frac{dx}{x \ln x} = \lim_{t\to\infty} \Big[\ln(\ln x)\Big]_2^t = \infty$]. Setting $a_n = \dfrac{(-1)^n}{n \ln n}$ for $n \ge 2$, we find that $\displaystyle\sum_{n=2}^{\infty} n a_n = \sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$ converges by the Alternating Series Test.

It is easy to find conditionally convergent series $\sum a_n$ such that $\sum n a_n$ diverges. Two examples are $\displaystyle\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$ and $\displaystyle\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n}}$, both of which converge by the Alternating Series Test and fail to converge absolutely because $\sum |a_n|$ is a p-series with $p \le 1$. In both cases, $\sum n a_n$ diverges by the Test for Divergence.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

11.7 Strategy for Testing Series

1. Use the Limit Comparison Test with $a_n = \dfrac{n^2-1}{n^3+1}$ and $b_n = \dfrac{1}{n}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{(n^2-1)n}{n^3+1} = \lim\limits_{n\to\infty} \dfrac{n^3-n}{n^3+1} = \lim\limits_{n\to\infty} \dfrac{1-1/n^2}{1+1/n^3} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ is the divergent harmonic series, the series $\sum\limits_{n=1}^{\infty} \dfrac{n^2-1}{n^3+1}$ also diverges.

2. $\dfrac{n-1}{n^3+1} < \dfrac{n}{n^3+1} < \dfrac{n}{n^3} = \dfrac{1}{n^2}$ for $n \geq 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{n-1}{n^3+1}$ converges by comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$, which converges because it is a p-series with $p = 2 > 1$.

3. $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2-1}{n^3+1} = \sum\limits_{n=1}^{\infty} (-1)^n b_n$. Now $b_n = \dfrac{n^2-1}{n^3+1} > 0$ for $n \geq 2$, $\{b_n\}$ is decreasing for $n \geq 2$, and $\lim\limits_{n\to\infty} b_n = 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2-1}{n^3+1}$ converges by the Alternating Series Test. By Exercise 1, $\sum\limits_{n=1}^{\infty} \dfrac{n^2-1}{n^3+1}$ diverges, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2-1}{n^3+1}$ is conditionally convergent.

4. $\lim\limits_{n\to\infty} |a_n| = \lim\limits_{n\to\infty} \left|(-1)^n \dfrac{n^2-1}{n^2+1}\right| = \lim\limits_{n\to\infty} \dfrac{1-1/n^2}{1+1/n^2} = 1 \neq 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2-1}{n^2+1}$ diverges by the Test for Divergence. [Note that $\lim\limits_{n\to\infty} (-1)^n \dfrac{n^2-1}{n^2+1}$ does not exist.]

5. $\lim\limits_{x\to\infty} \dfrac{e^x}{x^2} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{e^x}{2x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{e^x}{2} = \infty$, so $\lim\limits_{n\to\infty} \dfrac{e^n}{n^2} = \infty$. Thus, the series $\sum\limits_{n=1}^{\infty} \dfrac{e^n}{n^2}$ diverges by the Test for Divergence.

6. $\lim\limits_{n\to\infty} \sqrt[n]{|a_n|} = \lim\limits_{n\to\infty} \sqrt[n]{\dfrac{n^{2n}}{(1+n)^{3n}}} = \lim\limits_{n\to\infty} \dfrac{n^2}{(1+n)^3} = \lim\limits_{n\to\infty} \dfrac{1/n}{(1/n+1)^3} = \dfrac{0}{1} = 0 < 1$, so the series $\sum\limits_{n=1}^{\infty} \dfrac{n^{2n}}{(1+n)^{3n}}$ converges by the Root Test.

7. Let $f(x) = \dfrac{1}{x\sqrt{\ln x}}$. Then f is positive, continuous, and decreasing on $[2, \infty)$, so we can apply the Integral Test.

Since $\displaystyle\int \frac{1}{x\sqrt{\ln x}}\,dx \left[\begin{matrix} u = \ln x, \\ du = dx/x \end{matrix}\right] = \int u^{-1/2}\,du = 2u^{1/2} + C = 2\sqrt{\ln x} + C$, we find

$\displaystyle\int_2^{\infty} \frac{dx}{x\sqrt{\ln x}} = \lim_{t\to\infty} \int_2^t \frac{dx}{x\sqrt{\ln x}} = \lim_{t\to\infty} \left[2\sqrt{\ln x}\right]_2^t = \lim_{t\to\infty} \left(2\sqrt{\ln t} - 2\sqrt{\ln 2}\right) = \infty$. Since the integral diverges, the given series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n\sqrt{\ln n}}$ diverges.

8. $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{(n+1)^4}{4^{n+1}} \cdot \dfrac{4^n}{n^4}\right| = \lim\limits_{n\to\infty} \dfrac{(n+1)^4}{4n^4} = \dfrac{1}{4} \lim\limits_{n\to\infty} \left(1 + \dfrac{1}{n}\right)^4 = \dfrac{1}{4}(1) = \dfrac{1}{4} < 1$, so the series $\sum\limits_{n=1}^{\infty} (-1)^{n-1} \dfrac{n^4}{4^n}$ is absolutely convergent (and therefore convergent) by the Ratio Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{\pi^{2n+2}}{(2n+2)!}\cdot\frac{(2n)!}{\pi^{2n}}\right| = \lim_{n\to\infty}\frac{\pi^2}{(2n+2)(2n+1)} = 0 < 1$, so the series $\sum_{n=0}^{\infty}(-1)^n\frac{\pi^{2n}}{(2n)!}$ is absolutely convergent (and therefore convergent) by the Ratio Test.

10. Let $f(x) = x^2e^{-x^3}$. Then f is continuous and positive on $[1,\infty)$, and $f'(x) = \frac{x(2-3x^3)}{e^{x^3}} < 0$ for $x \geq 1$, so f is decreasing on $[1,\infty)$ as well, and we can apply the Integral Test. $\int_1^{\infty} x^2e^{-x^3}\,dx = \lim_{t\to\infty}\left[-\frac{1}{3}e^{-x^3}\right]_1^t = \frac{1}{3e}$, so the integral converges, and hence, the series converges.

11. $\sum_{n=1}^{\infty}\left(\frac{1}{n^3}+\frac{1}{3^n}\right) = \sum_{n=1}^{\infty}\frac{1}{n^3} + \sum_{n=1}^{\infty}\left(\frac{1}{3}\right)^n$. The first series converges since it is a p-series with $p = 3 > 1$ and the second series converges since it is geometric with $|r| = \frac{1}{3} < 1$. The sum of two convergent series is convergent.

12. $\frac{1}{k\sqrt{k^2+1}} < \frac{1}{k\sqrt{k^2}} = \frac{1}{k^2}$, so $\sum_{k=1}^{\infty}\frac{1}{k\sqrt{k^2+1}}$ converges by comparison with the convergent p-series $\sum_{k=1}^{\infty}\frac{1}{k^2}$ $(p = 2 > 1)$.

13. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{3^{n+1}(n+1)^2}{(n+1)!}\cdot\frac{n!}{3^n n^2}\right| = \lim_{n\to\infty}\frac{3(n+1)^2}{(n+1)n^2} = 3\lim_{n\to\infty}\frac{n+1}{n^2} = 0 < 1$, so the series $\sum_{n=1}^{\infty}\frac{3^n n^2}{n!}$ converges by the Ratio Test.

14. $\left|\frac{\sin 2n}{1+2^n}\right| \leq \frac{1}{1+2^n} < \frac{1}{2^n} = \left(\frac{1}{2}\right)^n$, so the series $\sum_{n=1}^{\infty}\left|\frac{\sin 2n}{1+2^n}\right|$ converges by comparison with the geometric series $\sum_{n=1}^{\infty}\left(\frac{1}{2}\right)^n$ with $|r| = \frac{1}{2} < 1$. Thus, the series $\sum_{n=1}^{\infty}\frac{\sin 2n}{1+2^n}$ converges absolutely, implying convergence.

15. $a_k = \frac{2^{k-1}3^{k+1}}{k^k} = \frac{2^k2^{-1}3^k3^1}{k^k} = \frac{3}{2}\left(\frac{2\cdot 3}{k}\right)^k$. By the Root Test, $\lim_{k\to\infty}\sqrt[k]{\left(\frac{6}{k}\right)^k} = \lim_{k\to\infty}\frac{6}{k} = 0 < 1$, so the series $\sum_{k=1}^{\infty}\left(\frac{6}{k}\right)^k$ converges. It follows from Theorem 8(i) in Section 11.2 that the given series, $\sum_{k=1}^{\infty}\frac{2^{k-1}3^{k+1}}{k^k} = \sum_{k=1}^{\infty}\frac{3}{2}\left(\frac{6}{k}\right)^k$, also converges.

16. Use the Limit Comparison Test with $a_n = \frac{\sqrt{n^4+1}}{n^3+n}$ and $b_n = \frac{1}{n}$:

$\lim_{n\to\infty}\frac{a_n}{b_n} = \lim_{n\to\infty}\frac{n\sqrt{n^4+1}}{n(n^2+1)} = \lim_{n\to\infty}\frac{\sqrt{n^4+1}/n^2}{(n^2+1)/n^2} = \lim_{n\to\infty}\frac{\sqrt{1+1/n^4}}{1+1/n^2} = 1 > 0$. Since $\sum_{n=1}^{\infty}\frac{1}{n}$ is the divergent harmonic series, the series $\sum_{n=1}^{\infty}\frac{\sqrt{n^4+1}}{n^3+n}$ also diverges.

17. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{1\cdot3\cdot5\cdot\cdots\cdot(2n-1)(2n+1)}{2\cdot5\cdot8\cdot\cdots\cdot(3n-1)(3n+2)}\cdot\frac{2\cdot5\cdot8\cdot\cdots\cdot(3n-1)}{1\cdot3\cdot5\cdot\cdots\cdot(2n-1)}\right| = \lim_{n\to\infty}\frac{2n+1}{3n+2}$

$= \lim_{n\to\infty}\frac{2+1/n}{3+2/n} = \frac{2}{3} < 1,$

so the series $\sum_{n=1}^{\infty}\frac{1\cdot3\cdot5\cdot\cdots\cdot(2n-1)}{2\cdot5\cdot8\cdot\cdots\cdot(3n-1)}$ converges by the Ratio Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

18. $b_n = \dfrac{1}{\sqrt{n}-1}$ for $n \geq 2$. $\{b_n\}$ is a decreasing sequence of positive numbers and $\lim\limits_{n\to\infty} b_n = 0$, so $\sum\limits_{n=2}^{\infty} \dfrac{(-1)^{n-1}}{\sqrt{n}-1}$ converges by the Alternating Series Test.

19. Let $f(x) = \dfrac{\ln x}{\sqrt{x}}$. Then $f'(x) = \dfrac{2-\ln x}{2x^{3/2}} < 0$ when $\ln x > 2$ or $x > e^2$, so $\dfrac{\ln n}{\sqrt{n}}$ is decreasing for $n > e^2$.

By l'Hospital's Rule, $\lim\limits_{n\to\infty} \dfrac{\ln n}{\sqrt{n}} = \lim\limits_{n\to\infty} \dfrac{1/n}{1/\left(2\sqrt{n}\right)} = \lim\limits_{n\to\infty} \dfrac{2}{\sqrt{n}} = 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{\ln n}{\sqrt{n}}$ converges by the Alternating Series Test.

20. $a_k = \dfrac{\sqrt[3]{k}-1}{k(\sqrt{k}+1)} < \dfrac{\sqrt[3]{k}}{k(\sqrt{k}+1)} < \dfrac{\sqrt[3]{k}}{k\sqrt{k}} = \dfrac{k^{1/3}}{k^{3/2}} = \dfrac{1}{k^{7/6}}$, so the series $\sum\limits_{k=1}^{\infty} \dfrac{\sqrt[3]{k}-1}{k(\sqrt{k}+1)}$ converges by comparison with the convergent p-series $\sum\limits_{k=1}^{\infty} \dfrac{1}{k^{7/6}}$ $\left(p = \frac{7}{6} > 1\right)$.

21. $\lim\limits_{n\to\infty} |a_n| = \lim\limits_{n\to\infty} \left|(-1)^n \cos(1/n^2)\right| = \lim\limits_{n\to\infty} \left|\cos(1/n^2)\right| = \cos 0 = 1$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \cos(1/n^2)$ diverges by the Test for Divergence.

22. $\lim\limits_{k\to\infty} |a_k| = \lim\limits_{k\to\infty} \left|\dfrac{1}{2+\sin k}\right| = \lim\limits_{k\to\infty} \dfrac{1}{2+\sin k}$, which does not exist (the terms vary between $\frac{1}{3}$ and 1). Thus, the series $\sum\limits_{k=1}^{\infty} \dfrac{1}{2+\sin k}$ diverges by the Test for Divergence.

23. Using the Limit Comparison Test with $a_n = \tan\left(\dfrac{1}{n}\right)$ and $b_n = \dfrac{1}{n}$, we have

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{\tan(1/n)}{1/n} = \lim\limits_{x\to\infty} \dfrac{\tan(1/x)}{1/x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{\sec^2(1/x)\cdot(-1/x^2)}{-1/x^2} = \lim\limits_{x\to\infty} \sec^2(1/x) = 1^2 = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} b_n$ is the divergent harmonic series, $\sum\limits_{n=1}^{\infty} a_n$ is also divergent.

24. $\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} \left(n \sin\dfrac{1}{n}\right) = \lim\limits_{n\to\infty} \dfrac{\sin(1/n)}{1/n} = \lim\limits_{x\to 0^+} \dfrac{\sin x}{x} = 1 \neq 0$, so the series $\sum\limits_{n=1}^{\infty} n\sin(1/n)$ diverges by the Test for Divergence.

25. Use the Ratio Test. $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{(n+1)!}{e^{(n+1)^2}} \cdot \dfrac{e^{n^2}}{n!}\right| = \lim\limits_{n\to\infty} \dfrac{(n+1)n!\cdot e^{n^2}}{e^{n^2+2n+1} n!} = \lim\limits_{n\to\infty} \dfrac{n+1}{e^{2n+1}} = 0 < 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{n!}{e^{n^2}}$ converges.

26. $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \dfrac{a_{n+1}}{a_n} = \lim\limits_{n\to\infty} \left(\dfrac{n^2+2n+2}{5^{n+1}} \cdot \dfrac{5^n}{n^2+1}\right) = \lim\limits_{n\to\infty} \left(\dfrac{1+2/n+2/n^2}{1+1/n^2} \cdot \dfrac{1}{5}\right) = \dfrac{1}{5} < 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{n^2+1}{5^n}$ converges by the Ratio Test.

27. $\displaystyle\int_2^{\infty} \frac{\ln x}{x^2}\,dx = \lim_{t\to\infty} \left[-\frac{\ln x}{x} - \frac{1}{x}\right]_1^t$ [using integration by parts] $\overset{\text{H}}{=} 1$. So $\sum\limits_{n=1}^{\infty} \dfrac{\ln n}{n^2}$ converges by the Integral Test, and since $\dfrac{k\ln k}{(k+1)^3} < \dfrac{k\ln k}{k^3} = \dfrac{\ln k}{k^2}$, the given series $\sum\limits_{k=1}^{\infty} \dfrac{k\ln k}{(k+1)^3}$ converges by the Comparison Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. Since $\left\{\frac{1}{n}\right\}$ is a decreasing sequence, $e^{1/n} \le e^{1/1} = e$ for all $n \ge 1$, and $\sum_{n=1}^{\infty} \frac{e}{n^2}$ converges ($p = 2 > 1$), so $\sum_{n=1}^{\infty} \frac{e^{1/n}}{n^2}$ converges by the Comparison Test. (Or use the Integral Test.)

29. $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (-1)^n \frac{1}{\cosh n} = \sum_{n=1}^{\infty} (-1)^n b_n$. Now $b_n = \frac{1}{\cosh n} > 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

Or: Write $\frac{1}{\cosh n} = \frac{2}{e^n + e^{-n}} < \frac{2}{e^n}$ and $\sum_{n=1}^{\infty} \frac{1}{e^n}$ is a convergent geometric series, so $\sum_{n=1}^{\infty} \frac{1}{\cosh n}$ is convergent by the Comparison Test. So $\sum_{n=1}^{\infty} (-1)^n \frac{1}{\cosh n}$ is absolutely convergent and therefore convergent.

30. Let $f(x) = \frac{\sqrt{x}}{x+5}$. Then $f(x)$ is continuous and positive on $[1, \infty)$, and since $f'(x) = \frac{5-x}{2\sqrt{x}\,(x+5)^2} < 0$ for $x > 5$, $f(x)$ is eventually decreasing, so we can use the Alternating Series Test. $\lim_{n\to\infty} \frac{\sqrt{n}}{n+5} = \lim_{n\to\infty} \frac{1}{n^{1/2} + 5n^{1/2}} = 0$, so the series $\sum_{j=1}^{\infty} (-1)^j \frac{\sqrt{j}}{j+5}$ converges.

31. $\lim_{k\to\infty} a_k = \lim_{k\to\infty} \frac{5^k}{3^k + 4^k} = [\text{divide by } 4^k]$ $\lim_{k\to\infty} \frac{(5/4)^k}{(3/4)^k + 1} = \infty$ since $\lim_{k\to\infty} \left(\frac{3}{4}\right)^k = 0$ and $\lim_{k\to\infty} \left(\frac{5}{4}\right)^k = \infty$.

Thus, $\sum_{k=1}^{\infty} \frac{5^k}{3^k + 4^k}$ diverges by the Test for Divergence.

32. $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} \sqrt[n]{\left|\frac{(n!)^n}{n^{4n}}\right|} = \lim_{n\to\infty} \frac{n!}{n^4} = \lim_{n\to\infty} \left[\frac{n}{n} \cdot \frac{n-1}{n} \cdot \frac{n-2}{n} \cdot \frac{n-3}{n} \cdot (n-4)!\right]$

$= \lim_{n\to\infty} \left[\left(1 - \frac{1}{n}\right)\left(1 - \frac{2}{n}\right)\left(1 - \frac{3}{n}\right)(n-4)!\right] = \infty,$

so the series $\sum_{n=1}^{\infty} \frac{(n!)^n}{n^{4n}}$ diverges by the Root Test.

33. $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} \left(\frac{n}{n+1}\right)^{n^2/n} = \lim_{n\to\infty} \frac{1}{[(n+1)/n]^n} = \frac{1}{\lim_{n\to\infty} (1 + 1/n)^n} = \frac{1}{e} < 1$, so the series $\sum_{n=1}^{\infty} \left(\frac{n}{n+1}\right)^{n^2}$ converges by the Root Test.

34. $0 \le n\cos^2 n \le n$, so $\frac{1}{n + n\cos^2 n} \ge \frac{1}{n+n} = \frac{1}{2n}$. Thus, $\sum_{n=1}^{\infty} \frac{1}{n + n\cos^2 n}$ diverges by comparison with $\sum_{n=1}^{\infty} \frac{1}{2n}$, which is a constant multiple of the (divergent) harmonic series.

35. $a_n = \frac{1}{n^{1+1/n}} = \frac{1}{n \cdot n^{1/n}}$, so let $b_n = \frac{1}{n}$ and use the Limit Comparison Test. $\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{1}{n^{1/n}} = 1 > 0$ (see Exercise 6.8.63 [ET 4.4.63]), so the series $\sum_{n=1}^{\infty} \frac{1}{n^{1+1/n}}$ diverges by comparison with the divergent harmonic series.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

36. Note that $(\ln n)^{\ln n} = \left(e^{\ln \ln n}\right)^{\ln n} = \left(e^{\ln n}\right)^{\ln \ln n} = n^{\ln \ln n}$ and $\ln \ln n \to \infty$ as $n \to \infty$, so $\ln \ln n > 2$ for sufficiently large n. For these n we have $(\ln n)^{\ln n} > n^2$, so $\dfrac{1}{(\ln n)^{\ln n}} < \dfrac{1}{n^2}$. Since $\sum_{n=2}^{\infty} \dfrac{1}{n^2}$ converges $[p = 2 > 1]$, so does $\sum_{n=2}^{\infty} \dfrac{1}{(\ln n)^{\ln n}}$ by the Comparison Test.

37. $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} (2^{1/n} - 1) = 1 - 1 = 0 < 1$, so the series $\sum_{n=1}^{\infty} \left(\sqrt[n]{2} - 1\right)^n$ converges by the Root Test.

38. Use the Limit Comparison Test with $a_n = \sqrt[n]{2} - 1$ and $b_n = 1/n$. Then

$$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{2^{1/n} - 1}{1/n} = \lim_{x\to\infty} \frac{2^{1/x} - 1}{1/x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{2^{1/x} \cdot \ln 2 \cdot (-1/x^2)}{-1/x^2} = \lim_{x\to\infty} (2^{1/x} \cdot \ln 2) = 1 \cdot \ln 2 = \ln 2 > 0.$$

So since $\sum_{n=1}^{\infty} b_n$ diverges (harmonic series), so does $\sum_{n=1}^{\infty} \left(\sqrt[n]{2} - 1\right)$.

Alternate solution: $\sqrt[n]{2} - 1 = \dfrac{1}{2^{(n-1)/n} + 2^{(n-2)/n} + 2^{(n-3)/n} + \cdots + 2^{1/n} + 1}$ [rationalize the numerator] $\geq \dfrac{1}{2n}$,

and since $\sum_{n=1}^{\infty} \dfrac{1}{2n} = \dfrac{1}{2} \sum_{n=1}^{\infty} \dfrac{1}{n}$ diverges (harmonic series), so does $\sum_{n=1}^{\infty} \left(\sqrt[n]{2} - 1\right)$ by the Comparison Test.

11.8 Power Series

1. A power series is a series of the form $\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots$, where x is a variable and the c_n's are constants called the coefficients of the series.

More generally, a series of the form $\sum_{n=0}^{\infty} c_n (x - a)^n = c_0 + c_1 (x - a) + c_2 (x - a)^2 + \cdots$ is called a power series in $(x - a)$ or a power series centered at a or a power series about a, where a is a constant.

2. (a) Given the power series $\sum_{n=0}^{\infty} c_n (x - a)^n$, the radius of convergence is:

(i) 0 if the series converges only when $x = a$

(ii) ∞ if the series converges for all x, or

(iii) a positive number R such that the series converges if $|x - a| < R$ and diverges if $|x - a| > R$.

In most cases, R can be found by using the Ratio Test.

(b) The interval of convergence of a power series is the interval that consists of all values of x for which the series converges. Corresponding to the cases in part (a), the interval of convergence is: (i) the single point $\{a\}$, (ii) all real numbers; that is, the real number line $(-\infty, \infty)$, or (iii) an interval with endpoints $a - R$ and $a + R$ which can contain neither, either, or both of the endpoints. In this case, we must test the series for convergence at each endpoint to determine the interval of convergence.

3. If $a_n = (-1)^n n x^n$, then

$\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\dfrac{(-1)^{n+1}(n+1)x^{n+1}}{(-1)^n n x^n}\right| = \lim_{n\to\infty} \left|(-1)\dfrac{n+1}{n} x\right| = \lim_{n\to\infty} \left[\left(1 + \dfrac{1}{n}\right)|x|\right] = |x|$. By the Ratio Test, the series $\sum_{n=1}^{\infty} (-1)^n n x^n$ converges when $|x| < 1$, so the radius of convergence $R = 1$. Now we'll check the endpoints, that is,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x = \pm 1$. Both series $\sum_{n=1}^{\infty} (-1)^n n(\pm 1)^n = \sum_{n=1}^{\infty} (\mp 1)^n n$ diverge by the Test for Divergence since $\lim_{n\to\infty} |(\mp 1)^n n| = \infty$. Thus, the interval of convergence is $I = (-1, 1)$.

4. If $a_n = \dfrac{(-1)^n x^n}{\sqrt[3]{n}}$, then

$\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\dfrac{(-1)^{n+1}x^{n+1}}{\sqrt[3]{n+1}} \cdot \dfrac{\sqrt[3]{n}}{(-1)^n x^n}\right| = \lim_{n\to\infty} \left|\dfrac{(-1)x\sqrt[3]{n}}{\sqrt[3]{n+1}}\right| = \lim_{n\to\infty} \sqrt[3]{\dfrac{1}{1+1/n}}\,|x| = |x|$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \dfrac{(-1)^n x^n}{\sqrt[3]{n}}$ converges when $|x| < 1$, so $R = 1$. When $x = 1$, the series $\sum_{n=1}^{\infty} \dfrac{(-1)^n}{\sqrt[3]{n}}$ converges by the Alternating Series Test. When $x = -1$, the series $\sum_{n=1}^{\infty} \dfrac{1}{\sqrt[3]{n}}$ diverges since it is a p-series $\left(p = \frac{1}{3} \le 1\right)$. Thus, the interval of convergence is $(-1, 1]$.

5. If $a_n = \dfrac{x^n}{2n-1}$, then $\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\dfrac{x^{n+1}}{2n+1} \cdot \dfrac{2n-1}{x^n}\right| = \lim_{n\to\infty} \left(\dfrac{2n-1}{2n+1}\,|x|\right) = \lim_{n\to\infty} \left(\dfrac{2-1/n}{2+1/n}\,|x|\right) = |x|$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \dfrac{x^n}{2n-1}$ converges when $|x| < 1$, so $R = 1$. When $x = 1$, the series $\sum_{n=1}^{\infty} \dfrac{1}{2n-1}$ diverges by comparison with $\sum_{n=1}^{\infty} \dfrac{1}{2n}$ since $\dfrac{1}{2n-1} > \dfrac{1}{2n}$ and $\dfrac{1}{2}\sum_{n=1}^{\infty} \dfrac{1}{n}$ diverges since it is a constant multiple of the harmonic series. When $x = -1$, the series $\sum_{n=1}^{\infty} \dfrac{(-1)^n}{2n-1}$ converges by the Alternating Series Test. Thus, the interval of convergence is $[-1, 1)$.

6. If $a_n = \dfrac{(-1)^n x^n}{n^2}$, then

$\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\dfrac{(-1)^{n+1}x^{n+1}}{(n+1)^2} \cdot \dfrac{n^2}{(-1)^n x^n}\right| = \lim_{n\to\infty} \left|\dfrac{(-1)xn^2}{(n+1)^2}\right| = \lim_{n\to\infty} \left[\left(\dfrac{n}{n+1}\right)^2 |x|\right] = 1^2 \cdot |x| = |x|$.

By the Ratio Test, the series $\sum_{n=1}^{\infty} \dfrac{(-1)^n x^n}{n^2}$ converges when $|x| < 1$, so $R = 1$. When $x = 1$, the series $\sum_{n=1}^{\infty} \dfrac{(-1)^n}{n^2}$ converges by the Alternating Series Test. When $x = -1$, the series $\sum_{n=1}^{\infty} \dfrac{1}{n^2}$ converges since it is a p-series with $p = 2 > 1$. Thus, the interval of convergence is $[-1, 1]$.

7. If $a_n = \dfrac{x^n}{n!}$, then $\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\dfrac{x^{n+1}}{(n+1)!} \cdot \dfrac{n!}{x^n}\right| = \lim_{n\to\infty} \left|\dfrac{x}{n+1}\right| = |x| \lim_{n\to\infty} \dfrac{1}{n+1} = |x| \cdot 0 = 0 < 1$ for *all* real x. So, by the Ratio Test, $R = \infty$ and $I = (-\infty, \infty)$.

8. Here the Root Test is easier. If $a_n = n^n x^n$, then $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} n\,|x| = \infty$ if $x \neq 0$, so $R = 0$ and $I = \{0\}$.

9. If $a_n = \dfrac{x^n}{n^4\, 4^n}$, then

$\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\dfrac{x^{n+1}}{(n+1)^4\, 4^{n+1}} \cdot \dfrac{n^4\, 4^n}{x^n}\right| = \lim_{n\to\infty} \left|\dfrac{n^4}{(n+1)^4} \cdot \dfrac{x}{4}\right| = \lim_{n\to\infty} \left(\dfrac{n}{n+1}\right)^4 \dfrac{|x|}{4} = 1^4 \cdot \dfrac{|x|}{4} = \dfrac{|x|}{4}$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \dfrac{x^n}{n^4\, 4^n}$ converges when $\dfrac{|x|}{4} < 1 \quad \Leftrightarrow \quad |x| < 4$, so $R = 4$. When $x = 4$, the series $\sum_{n=1}^{\infty} \dfrac{1}{n^4}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

converges since it is a p-series ($p = 4 > 1$). When $x = -4$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^4}$ converges by the Alternating Series Test.

Thus, the interval of convergence is $[-4, 4]$.

10. If $a_n = 2^n n^2 x^n$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{2^{n+1}(n+1)^2 x^{n+1}}{2^n n^2 x^n}\right| = \lim_{n\to\infty} 2\left(\frac{n+1}{n}\right)^2 |x| = 2|x|$. By the Ratio Test,

the series $\sum_{n=1}^{\infty} 2^n n^2 x^n$ converges when $2|x| < 1 \quad \Leftrightarrow \quad |x| < \frac{1}{2}$, so $R = \frac{1}{2}$. When $x = \pm\frac{1}{2}$, both series

$\sum_{n=1}^{\infty} 2^n n^2 \left(\pm\frac{1}{2}\right)^n = \sum_{n=1}^{\infty} (\pm 1)^n n^2$ diverge by the Test for Divergence since $\lim_{n\to\infty} \left|(\pm 1)^n n^2\right| = \infty$. Thus, the interval of

convergence is $\left(-\frac{1}{2}, \frac{1}{2}\right)$.

11. If $a_n = \frac{(-1)^n 4^n}{\sqrt{n}} x^n$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^{n+1} 4^{n+1} x^{n+1}}{\sqrt{n+1}} \cdot \frac{\sqrt{n}}{(-1)^n 4^n x^n}\right| = \lim_{n\to\infty} \sqrt{\frac{n}{n+1}} \cdot 4|x| = 4|x|$.

By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{(-1)^n 4^n}{\sqrt{n}} x^n$ converges when $4|x| < 1 \quad \Leftrightarrow \quad |x| < \frac{1}{4}$, so $R = \frac{1}{4}$. When $x = \frac{1}{4}$, the series

$\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$ converges by the Alternating Series Test. When $x = -\frac{1}{4}$, the series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ diverges since it is a p-series

$\left(p = \frac{1}{2} \le 1\right)$. Thus, the interval of convergence is $\left(-\frac{1}{4}, \frac{1}{4}\right]$.

12. If $a_n = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n5^n} x^n$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^n x^{n+1}}{(n+1)5^{n+1}} \cdot \frac{n5^n}{(-1)^{n-1} x^n}\right| = \lim_{n\to\infty} \left(\frac{n}{n+1}\right)\frac{|x|}{5} = 1 \cdot \frac{|x|}{5} = \frac{|x|}{5}$.

By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n5^n} x^n$ converges when $\frac{|x|}{5} < 1 \quad \Leftrightarrow \quad |x| < 5$, so $R = 5$. When $x = 5$, the series

$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$ converges by the Alternating Series Test. When $x = -5$, the series $\sum_{n=1}^{\infty} \frac{-1}{n}$ diverges since it is a constant

multiple of the harmonic series. Thus, the interval of convergence is $(-5, 5]$.

13. If $a_n = \frac{n}{2^n(n^2+1)} x^n$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(n+1)x^{n+1}}{2^{n+1}(n^2+2n+2)} \cdot \frac{2^n(n^2+1)}{n x^n}\right| = \lim_{n\to\infty} \frac{n^3+n^2+n+1}{n^3+2n^2+2n} \cdot \frac{|x|}{2}$$

$$= \lim_{n\to\infty} \frac{1+1/n+1/n^2+1/n^3}{1+2/n+2/n^2} \cdot \frac{|x|}{2} = \frac{|x|}{2}$$

By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{n}{2^n(n^2+1)} x^n$ converges when $\frac{|x|}{2} < 1 \quad \Leftrightarrow \quad |x| < 2$, so $R = 2$. When $x = 2$, the series

$\sum_{n=1}^{\infty} \frac{n}{n^2+1}$ diverges by the Limit Comparison Test with $b_n = \frac{1}{n}$. When $x = -2$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2+1}$ converges by the

Alternating Series Test. Thus, the interval of convergence is $[-2, 2)$.

14. If $a_n = \frac{x^{2n}}{n!}$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{2n+2}}{(n+1)!} \cdot \frac{n!}{x^{2n}}\right| = \lim_{n\to\infty} \frac{|x^2|}{n+1} = 0 < 1$ for *all* real x. So, by the Ratio Test,

$R = \infty$ and $I = (-\infty, \infty)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. If $a_n = \dfrac{(x-2)^n}{n^2+1}$, then $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(x-2)^{n+1}}{(n+1)^2+1}\cdot\dfrac{n^2+1}{(x-2)^n}\right| = |x-2|\lim\limits_{n\to\infty}\dfrac{n^2+1}{(n+1)^2+1} = |x-2|$. By the Ratio Test, the series $\sum\limits_{n=0}^{\infty}\dfrac{(x-2)^n}{n^2+1}$ converges when $|x-2|<1 \quad [R=1] \quad\Leftrightarrow\quad -1<x-2<1 \quad\Leftrightarrow\quad 1<x<3$. When $x=1$, the series $\sum\limits_{n=0}^{\infty}(-1)^n\dfrac{1}{n^2+1}$ converges by the Alternating Series Test; when $x=3$, the series $\sum\limits_{n=0}^{\infty}\dfrac{1}{n^2+1}$ converges by comparison with the p-series $\sum\limits_{n=1}^{\infty}\dfrac{1}{n^2}$ $[p=2>1]$. Thus, the interval of convergence is $I=[1,3]$.

16. If $a_n = \dfrac{(-1)^n}{(2n-1)2^n}(x-1)^n$, then

$\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)^{n+1}(x-1)^{n+1}}{(2n+1)\,2^{n+1}}\cdot\dfrac{(2n-1)\,2^n}{(-1)^n(x-1)^n}\right| = \lim\limits_{n\to\infty}\dfrac{2n-1}{2n+1}\cdot\dfrac{|x-1|}{2} = \dfrac{|x-1|}{2}$. By the Ratio Test, the series $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{(2n-1)\,2^n}(x-1)^n$ converges when $\dfrac{|x-1|}{2}<1 \quad\Leftrightarrow\quad |x-1|<2 \quad [R=2] \quad\Leftrightarrow\quad -2<x-1<2 \quad\Leftrightarrow$

$-1<x<3$. When $x=3$, the series $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{2n-1}$ converges by the Alternating Series Test. When $x=-1$, the series $\sum\limits_{n=1}^{\infty}\dfrac{1}{2n-1}$ diverges by the Limit Comparison Test with $b_n=\dfrac{1}{n}$. Thus, the interval of convergence is $(-1,3]$.

17. If $a_n = \dfrac{(x+2)^n}{2^n\,\ln n}$, then $\lim\limits_{n\to\infty}\left|\dfrac{(x+2)^{n+1}}{2^{n+1}\ln(n+1)}\cdot\dfrac{2^n\ln n}{(x+2)^n}\right| = \lim\limits_{n\to\infty}\dfrac{\ln n}{\ln(n+1)}\cdot\dfrac{|x+2|}{2} = \dfrac{|x+2|}{2}$ since

$\lim\limits_{n\to\infty}\dfrac{\ln n}{\ln(n+1)} = \lim\limits_{x\to\infty}\dfrac{\ln x}{\ln(x+1)} \overset{\text{H}}{=} \lim\limits_{x\to\infty}\dfrac{1/x}{1/(x+1)} = \lim\limits_{x\to\infty}\dfrac{x+1}{x} = \lim\limits_{x\to\infty}\left(1+\dfrac{1}{x}\right) = 1$. By the Ratio Test, the series $\sum\limits_{n=2}^{\infty}\dfrac{(x+2)^n}{2^n\,\ln n}$ converges when $\dfrac{|x+2|}{2}<1 \quad\Leftrightarrow\quad |x+2|<2 \quad [R=2] \quad\Leftrightarrow\quad -2<x+2<2 \quad\Leftrightarrow\quad -4<x<0$.

When $x=-4$, the series $\sum\limits_{n=2}^{\infty}\dfrac{(-1)^n}{\ln n}$ converges by the Alternating Series Test. When $x=0$, the series $\sum\limits_{n=2}^{\infty}\dfrac{1}{\ln n}$ diverges by the Limit Comparison Test with $b_n=\dfrac{1}{n}$ (or by comparison with the harmonic series). Thus, the interval of convergence is $[-4,0)$.

18. If $a_n = \dfrac{\sqrt{n}}{8^n}(x+6)^n$, then

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{\sqrt{n+1}\,(x+6)^{n+1}}{8^{n+1}}\cdot\frac{8^n}{\sqrt{n}(x+6)^n}\right| = \lim_{n\to\infty}\sqrt{\frac{n+1}{n}}\cdot\frac{|x+6|}{8}$$

$$= \lim_{n\to\infty}\sqrt{1+\frac{1}{n}}\cdot\frac{|x+6|}{8} = \frac{|x+6|}{8}$$

By the Ratio Test, the series $\sum\limits_{n=1}^{\infty}\dfrac{\sqrt{n}}{8^n}(x+6)^n$ converges when $\dfrac{|x+6|}{8}<1 \quad\Leftrightarrow\quad |x+6|<8 \quad [R=8] \quad\Leftrightarrow$

$-8<x+6<8 \quad\Leftrightarrow\quad -14<x<2$. When $x=2$, the series $\sum\limits_{n=1}^{\infty}\sqrt{n}$ diverges by the Test for Divergence since $\lim\limits_{n\to\infty}|a_n| = \lim\limits_{n\to\infty}\sqrt{n} = \infty > 0$. Similarly, when $x=-14$, the series $\sum\limits_{n=1}^{\infty}(-1)^n\sqrt{n}$ diverges. Thus, the interval of convergence is $(-14,2)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

19. If $a_n = \dfrac{(x-2)^n}{n^n}$, then $\lim\limits_{n\to\infty} \sqrt[n]{|a_n|} = \lim\limits_{n\to\infty} \dfrac{|x-2|}{n} = 0$, so the series converges for all x (by the Root Test).

$R = \infty$ and $I = (-\infty, \infty)$.

20. If $a_n = \dfrac{(2x-1)^n}{5^n\sqrt{n}}$, then

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(2x-1)^{n+1}}{5^{n+1}\sqrt{n+1}}\cdot\frac{5^n\sqrt{n}}{(2x-1)^n}\right| = \lim_{n\to\infty}\frac{|2x-1|}{5}\sqrt{\frac{n}{n+1}} = \lim_{n\to\infty}\frac{|2x-1|}{5}\sqrt{\frac{1}{1+1/n}} = \frac{|2x-1|}{5}.$$

By the Ratio Test, the series $\sum\limits_{n=1}^{\infty}\dfrac{(2x-1)^n}{5^n\sqrt{n}}$ converges when $\dfrac{|2x-1|}{5} < 1 \;\Leftrightarrow\; |2x-1| < 5 \;\Leftrightarrow\; \left|x-\frac{1}{2}\right| < \frac{5}{2} \;\Leftrightarrow$

$-\frac{5}{2} < x - \frac{1}{2} < \frac{5}{2} \;\Leftrightarrow\; -2 < x < 3$, so $R = \frac{5}{2}$. When $x = 3$, the series $\sum\limits_{n=1}^{\infty}\dfrac{1}{\sqrt{n}}$ is a divergent p-series $\left(p = \frac{1}{2} \le 1\right)$.

When $x = -2$, the series $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{\sqrt{n}}$ converges by the Alternating Series Test. Thus, the interval of convergence is $I = [-2, 3)$.

21. $a_n = \dfrac{n}{b^n}(x-a)^n$, where $b > 0$.

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\frac{(n+1)\,|x-a|^{n+1}}{b^{n+1}}\cdot\frac{b^n}{n\,|x-a|^n} = \lim_{n\to\infty}\left(1+\frac{1}{n}\right)\frac{|x-a|}{b} = \frac{|x-a|}{b}.$$

By the Ratio Test, the series converges when $\dfrac{|x-a|}{b} < 1 \;\Leftrightarrow\; |x-a| < b$ [so $R = b$] $\Leftrightarrow\; -b < x-a < b \;\Leftrightarrow$

$a - b < x < a + b$. When $|x-a| = b$, $\lim\limits_{n\to\infty}|a_n| = \lim\limits_{n\to\infty} n = \infty$, so the series diverges. Thus, $I = (a-b, a+b)$.

22. $a_n = \dfrac{b^n}{\ln n}(x-a)^n$, where $b > 0$.

$\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{b^{n+1}(x-a)^{n+1}}{\ln(n+1)}\cdot\dfrac{\ln n}{b^n(x-a)^n}\right| = \lim\limits_{n\to\infty}\dfrac{\ln n}{\ln(n+1)}\cdot b\,|x-a| = b\,|x-a|$ since

$\lim\limits_{n\to\infty}\dfrac{\ln n}{\ln(n+1)} = \lim\limits_{n\to\infty}\dfrac{\ln x}{\ln(x+1)} \overset{\text{H}}{=} \lim\limits_{x\to\infty}\dfrac{1/x}{1/(x+1)} = \lim\limits_{x\to\infty}\dfrac{x+1}{x} \overset{\text{H}}{=} \lim\limits_{x\to\infty}\dfrac{1}{1} = 1$. By the Ratio Test, the series

$\sum\limits_{n=2}^{\infty}\dfrac{b^n}{\ln n}(x-a)^n$ converges when $b\,|x-a| < 1 \;\Leftrightarrow\; |x-a| < \dfrac{1}{b} \;\Leftrightarrow\; -\dfrac{1}{b} < x-a < \dfrac{1}{b} \;\Leftrightarrow\; a - \dfrac{1}{b} < x < a + \dfrac{1}{b}$,

so $R = \dfrac{1}{b}$. When $x = a + \dfrac{1}{b}$, the series $\sum\limits_{n=2}^{\infty}\dfrac{1}{\ln n}$ diverges by comparison with the divergent p-series $\sum\limits_{n=2}^{\infty}\dfrac{1}{n}$ since $\dfrac{1}{\ln n} > \dfrac{1}{n}$

for $n \ge 2$. When $x = a - \dfrac{1}{b}$, the series $\sum\limits_{n=2}^{\infty}\dfrac{(-1)^n}{\ln n}$ converges by the Alternating Series Test. Thus, the interval of convergence is $I = \left[a - \dfrac{1}{b}, a + \dfrac{1}{b}\right)$.

23. If $a_n = n!\,(2x-1)^n$, then $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(n+1)!\,(2x-1)^{n+1}}{n!(2x-1)^n}\right| = \lim\limits_{n\to\infty}(n+1)\,|2x-1| \to \infty$ as $n \to \infty$

for all $x \ne \frac{1}{2}$. Since the series diverges for all $x \ne \frac{1}{2}$, $R = 0$ and $I = \left\{\frac{1}{2}\right\}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $a_n = \dfrac{n^2 x^n}{2 \cdot 4 \cdot 6 \cdot \cdots \cdot (2n)} = \dfrac{n^2 x^n}{2^n n!} = \dfrac{n x^n}{2^n (n-1)!}$, so

$\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \dfrac{(n+1)\,|x|^{n+1}}{2^{n+1} n!} \cdot \dfrac{2^n (n-1)!}{n\,|x|^n} = \lim\limits_{n\to\infty} \dfrac{n+1}{n^2} \dfrac{|x|}{2} = 0$. Thus, by the Ratio Test, the series converges for *all* real x and we have $R = \infty$ and $I = (-\infty, \infty)$.

25. If $a_n = \dfrac{(5x-4)^n}{n^3}$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(5x-4)^{n+1}}{(n+1)^3} \cdot \frac{n^3}{(5x-4)^n}\right| = \lim_{n\to\infty} |5x-4| \left(\frac{n}{n+1}\right)^3 = \lim_{n\to\infty} |5x-4| \left(\frac{1}{1+1/n}\right)^3$$
$$= |5x-4| \cdot 1 = |5x-4|$$

By the Ratio Test, $\sum\limits_{n=1}^{\infty} \dfrac{(5x-4)^n}{n^3}$ converges when $|5x-4| < 1 \;\Leftrightarrow\; \left|x - \frac{4}{5}\right| < \frac{1}{5} \;\Leftrightarrow\; -\frac{1}{5} < x - \frac{4}{5} < \frac{1}{5} \;\Leftrightarrow$

$\frac{3}{5} < x < 1$, so $R = \frac{1}{5}$. When $x = 1$, the series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^3}$ is a convergent p-series ($p = 3 > 1$). When $x = \frac{3}{5}$, the series $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^n}{n^3}$ converges by the Alternating Series Test. Thus, the interval of convergence is $I = \left[\frac{3}{5}, 1\right]$.

26. If $a_n = \dfrac{x^{2n}}{n\,(\ln n)^2}$, then $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{x^{2n+2}}{(n+1)[\ln(n+1)]^2} \cdot \dfrac{n\,(\ln n)^2}{x^{2n}}\right| = |x^2| \lim\limits_{n\to\infty} \dfrac{n\,(\ln n)^2}{(n+1)[\ln(n+1)]^2} = x^2$.

By the Ratio Test, the series $\sum\limits_{n=2}^{\infty} \dfrac{x^{2n}}{n\,(\ln n)^2}$ converges when $x^2 < 1 \;\Leftrightarrow\; |x| < 1$, so $R = 1$. When $x = \pm 1$, $x^{2n} = 1$, the series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n\,(\ln n)^2}$ converges by the Integral Test (see Exercise 11.3.29). Thus, the interval of convergence is $I = [-1, 1]$.

27. If $a_n = \dfrac{x^n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}$, then

$\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{x^{n+1}}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)(2n+1)} \cdot \dfrac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{x^n}\right| = \lim\limits_{n\to\infty} \dfrac{|x|}{2n+1} = 0 < 1$. Thus, by the Ratio Test, the series $\sum\limits_{n=1}^{\infty} \dfrac{x^n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}$ converges for *all* real x and we have $R = \infty$ and $I = (-\infty, \infty)$.

28. If $a_n = \dfrac{n!\, x^n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}$, then

$\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{(n+1)!\, x^{n+1}}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)(2n+1)} \cdot \dfrac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{n!\, x^n}\right| = \lim\limits_{n\to\infty} \dfrac{(n+1)\,|x|}{2n+1} = \frac{1}{2}\,|x|$.

By the Ratio Test, the series $\sum\limits_{n=1}^{\infty} a_n$ converges when $\frac{1}{2}\,|x| < 1 \;\Rightarrow\; |x| < 2$, so $R = 2$. When $x = \pm 2$,

$|a_n| = \dfrac{n!\, 2^n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)} = \dfrac{[1 \cdot 2 \cdot 3 \cdot \cdots \cdot n]\, 2^n}{[1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)]} = \dfrac{2 \cdot 4 \cdot 6 \cdot \cdots \cdot 2n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)} > 1$, so both endpoint series diverge by the Test for Divergence. Thus, the interval of convergence is $I = (-2, 2)$.

29. (a) We are given that the power series $\sum_{n=0}^{\infty} c_n x^n$ is convergent for $x = 4$. So by Theorem 4, it must converge for at least $-4 < x \le 4$. In particular, it converges when $x = -2$; that is, $\sum_{n=0}^{\infty} c_n (-2)^n$ is convergent.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) It does not follow that $\sum_{n=0}^{\infty} c_n(-4)^n$ is necessarily convergent. [See the comments after Theorem 4 about convergence at the endpoint of an interval. An example is $c_n = (-1)^n/(n4^n)$.]

30. We are given that the power series $\sum_{n=0}^{\infty} c_n x^n$ is convergent for $x = -4$ and divergent when $x = 6$. So by Theorem 4 it converges for at least $-4 \le x < 4$ and diverges for at least $x \ge 6$ and $x < -6$. Therefore:

(a) It converges when $x = 1$; that is, $\sum c_n$ is convergent.

(b) It diverges when $x = 8$; that is, $\sum c_n 8^n$ is divergent.

(c) It converges when $x = -3$; that is, $\sum c_n(-3^n)$ is convergent.

(d) It diverges when $x = -9$; that is, $\sum c_n(-9)^n = \sum(-1)^n c_n 9^n$ is divergent.

31. If $a_n = \dfrac{(n!)^k}{(kn)!}x^n$, then

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\frac{[(n+1)!]^k\,(kn)!}{(n!)^k\,[k(n+1)]!}|x| = \lim_{n\to\infty}\frac{(n+1)^k}{(kn+k)(kn+k-1)\cdots(kn+2)(kn+1)}|x|$$

$$= \lim_{n\to\infty}\left[\frac{(n+1)}{(kn+1)}\frac{(n+1)}{(kn+2)}\cdots\frac{(n+1)}{(kn+k)}\right]|x|$$

$$= \lim_{n\to\infty}\left[\frac{n+1}{kn+1}\right]\lim_{n\to\infty}\left[\frac{n+1}{kn+2}\right]\cdots\lim_{n\to\infty}\left[\frac{n+1}{kn+k}\right]|x|$$

$$= \left(\frac{1}{k}\right)^k|x| < 1 \quad\Leftrightarrow\quad |x| < k^k \text{ for convergence, and the radius of convergence is } R = k^k.$$

32. (a) Note that the four intervals in parts (a)–(d) have midpoint $m = \frac{1}{2}(p+q)$ and radius of convergence $r = \frac{1}{2}(q-p)$. We also know that the power series $\sum_{n=0}^{\infty} x^n$ has interval of convergence $(-1, 1)$. To change the radius of convergence to r, we can change x^n to $\left(\dfrac{x}{r}\right)^n$. To shift the midpoint of the interval of convergence, we can replace x with $x - m$. Thus, a power series whose interval of convergence is (p, q) is $\sum_{n=0}^{\infty}\left(\dfrac{x-m}{r}\right)^n$, where $m = \frac{1}{2}(p+q)$ and $r = \frac{1}{2}(q-p)$.

(b) Similar to Example 2, we know that $\sum_{n=1}^{\infty}\dfrac{x^n}{n}$ has interval of convergence $[-1, 1)$. By introducing the factor $(-1)^n$ in a_n, the interval of convergence changes to $(-1, 1]$. Now change the midpoint and radius as in part (a) to get $\sum_{n=1}^{\infty}(-1)^n\dfrac{1}{n}\left(\dfrac{x-m}{r}\right)^n$ as a power series whose interval of convergence is $(p, q]$.

(c) As in part (b), $\sum_{n=1}^{\infty}\dfrac{1}{n}\left(\dfrac{x-m}{r}\right)^n$ is a power series whose interval of convergence is $[p, q)$.

(d) If we increase the exponent on n (to say, $n = 2$), in the power series in part (c), then when $x = q$, the power series $\sum_{n=1}^{\infty}\dfrac{1}{n^2}\left(\dfrac{x-m}{r}\right)^n$ will converge by comparison to the p-series with $p = 2 > 1$, and the interval of convergence will be $[p, q]$.

33. No. If a power series is centered at a, its interval of convergence is symmetric about a. If a power series has an infinite radius of convergence, then its interval of convergence must be $(-\infty, \infty)$, not $[0, \infty)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. The partial sums of the series $\sum_{n=0}^{\infty} x^n$ definitely do not converge to $f(x) = 1/(1-x)$ for $x \geq 1$, since f is undefined at $x = 1$ and negative on $(1, \infty)$, while all the partial sums are positive on this interval. The partial sums also fail to converge to f for $x \leq -1$, since $0 < f(x) < 1$ on this interval, while the partial sums are either larger than 1 or less than 0. The partial sums seem to converge to f on $(-1, 1)$. This graphical evidence is consistent with what we know about geometric series: convergence for $|x| < 1$, divergence for $|x| \geq 1$ (see Examples 2 and 7 in Section 11.2).



35. (a) If $a_n = \dfrac{(-1)^n\, x^{2n+1}}{n!(n+1)!\, 2^{2n+1}}$, then

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{x^{2n+3}}{(n+1)!(n+2)!\, 2^{2n+3}} \cdot \frac{n!(n+1)!\, 2^{2n+1}}{x^{2n+1}}\right| = \left(\frac{x}{2}\right)^2 \lim_{n\to\infty}\frac{1}{(n+1)(n+2)} = 0 \text{ for all } x.$$

So $J_1(x)$ converges for all x and its domain is $(-\infty, \infty)$.

(b), (c) The initial terms of $J_1(x)$ up to $n = 5$ are $a_0 = \dfrac{x}{2}$, $a_1 = -\dfrac{x^3}{16}$, $a_2 = \dfrac{x^5}{384}$, $a_3 = -\dfrac{x^7}{18{,}432}$, $a_4 = \dfrac{x^9}{1{,}474{,}560}$, and $a_5 = -\dfrac{x^{11}}{176{,}947{,}200}$. The partial sums seem to approximate $J_1(x)$ well near the origin, but as $|x|$ increases, we need to take a large number of terms to get a good approximation.



36. (a) $A(x) = 1 + \sum_{n=1}^{\infty} a_n$, where $a_n = \dfrac{x^{3n}}{2 \cdot 3 \cdot 5 \cdot 6 \cdot \cdots \cdot (3n-1)(3n)}$, so $\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = |x|^3 \lim_{n\to\infty}\dfrac{1}{(3n+2)(3n+3)} = 0$ for all x, so the domain is $\mathbb{R}$.

(b), (c)



$s_0 = 1$ has been omitted from the graph. The partial sums seem to approximate $A(x)$ well near the origin, but as $|x|$ increases, we need to take a large number of terms to get a good approximation.

To plot A, we must first define $A(x)$ for the CAS. Note that for $n \geq 1$, the denominator of a_n is

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$2 \cdot 3 \cdot 5 \cdot 6 \cdot \cdots \cdot (3n-1) \cdot 3n = \dfrac{(3n)!}{1 \cdot 4 \cdot 7 \cdot \cdots \cdot (3n-2)} = \dfrac{(3n)!}{\prod_{k=1}^{n}(3k-2)}$, so $a_n = \dfrac{\prod_{k=1}^{n}(3k-2)}{(3n)!}x^{3n}$ and thus $A(x) \quad 1 + \sum_{n=1}^{\infty} \dfrac{\prod_{k=1}^{n}(3k-2)}{(3n)!}x^{3n}$. Both Maple and Mathematica are able to plot A if we define it this way, and Derive is able to produce a similar graph using a suitable partial sum of $A(x)$.

Derive, Maple and Mathematica all have two initially known Airy functions, called `AI·SERIES(z,m)` and `BI·SERIES(z,m)` from `BESSEL.MTH` in Derive and `AiryAi` and `AiryBi` in Maple and Mathematica (just `Ai` and `Bi` in older versions of Maple). However, it is very difficult to solve for A in terms of the CAS's Airy functions, although in fact $A(x) = \dfrac{\sqrt{3}\,\mathtt{AiryAi}(x) + \mathtt{AiryBi}(x)}{\sqrt{3}\,\mathtt{AiryAi}(0) + \mathtt{AiryBi}(0)}$.

37. $s_{2n-1} = 1 + 2x + x^2 + 2x^3 + x^4 + 2x^5 + \cdots + x^{2n-2} + 2x^{2n-1}$

$= 1(1+2x) + x^2(1+2x) + x^4(1+2x) + \cdots + x^{2n-2}(1+2x) = (1+2x)(1 + x^2 + x^4 + \cdots + x^{2n-2})$

$= (1+2x)\dfrac{1-x^{2n}}{1-x^2}$ [by (11.2.3) with $r = x^2$] $\quad \to \dfrac{1+2x}{1-x^2}$ as $n \to \infty$ by (11.2.4), when $|x| < 1$.

Also $s_{2n} = s_{2n-1} + x^{2n} \to \dfrac{1+2x}{1-x^2}$ since $x^{2n} \to 0$ for $|x| < 1$. Therefore, $s_n \to \dfrac{1+2x}{1-x^2}$ since s_{2n} and s_{2n-1} both approach $\dfrac{1+2x}{1-x^2}$ as $n \to \infty$. Thus, the interval of convergence is $(-1, 1)$ and $f(x) = \dfrac{1+2x}{1-x^2}$.

38. $s_{4n-1} = c_0 + c_1x + c_2x^2 + c_3x^3 + c_0x^4 + c_1x^5 + c_2x^6 + c_3x^7 + \cdots + c_3x^{4n-1}$

$= \left(c_0 + c_1x + c_2x^2 + c_3x^3\right)\left(1 + x^4 + x^8 + \cdots + x^{4n-4}\right) \to \dfrac{c_0 + c_1x + c_2x^2 + c_3x^3}{1-x^4}$ as $n \to \infty$

[by (11.2.4) with $r = x^4$] for $|x^4| < 1 \quad \Leftrightarrow \quad |x| < 1$. Also s_{4n}, s_{4n+1}, s_{4n+2} have the same limits (for example, $s_{4n} = s_{4n-1} + c_0x^{4n}$ and $x^{4n} \to 0$ for $|x| < 1$). So if at least one of c_0, c_1, c_2, and c_3 is nonzero, then the interval of convergence is $(-1, 1)$ and $f(x) = \dfrac{c_0 + c_1x + c_2x^2 + c_3x^3}{1-x^4}$.

39. We use the Root Test on the series $\sum c_nx^n$. We need $\lim_{n\to\infty} \sqrt[n]{|c_nx^n|} = |x| \lim_{n\to\infty} \sqrt[n]{|c_n|} = c\,|x| < 1$ for convergence, or $|x| < 1/c$, so $R = 1/c$.

40. Suppose $c_n \neq 0$. Applying the Ratio Test to the series $\sum c_n(x-a)^n$, we find that

$L = \lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\dfrac{c_{n+1}(x-a)^{n+1}}{c_n(x-a)^n}\right| = \lim_{n\to\infty}\dfrac{|x-a|}{|c_n/c_{n+1}|}(\star) = \dfrac{|x-a|}{\lim_{n\to\infty}|c_n/c_{n+1}|}$ (if $\lim_{n\to\infty}|c_n/c_{n+1}| \neq 0$), so the series converges when $\dfrac{|x-a|}{\lim_{n\to\infty}|c_n/c_{n+1}|} < 1 \quad \Leftrightarrow \quad |x-a| < \lim_{n\to\infty}\left|\dfrac{c_n}{c_{n+1}}\right|$. Thus, $R = \lim_{n\to\infty}\left|\dfrac{c_n}{c_{n+1}}\right|$. If $\lim_{n\to\infty}\left|\dfrac{c_n}{c_{n+1}}\right| = 0$ and $|x-a| \neq 0$, then $(\star)$ shows that $L = \infty$ and so the series diverges, and hence, $R = 0$. Thus, in all cases, $R = \lim_{n\to\infty}\left|\dfrac{c_n}{c_{n+1}}\right|$.

41. For $2 < x < 3$, $\sum c_nx^n$ diverges and $\sum d_nx^n$ converges. By Exercise 11.2.85, $\sum(c_n + d_n)\,x^n$ diverges. Since both series converge for $|x| < 2$, the radius of convergence of $\sum(c_n + d_n)\,x^n$ is 2.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. Since $\sum c_n x^n$ converges whenever $|x| < R$, $\sum c_n x^{2n} = \sum c_n (x^2)^n$ converges whenever $|x^2| < R \quad \Leftrightarrow \quad |x| < \sqrt{R}$, so the second series has radius of convergence $\sqrt{R}$.

11.9 Representations of Functions as Power Series

1. If $f(x) = \sum_{n=0}^{\infty} c_n x^n$ has radius of convergence 10, then $f'(x) = \sum_{n=1}^{\infty} n c_n x^{n-1}$ also has radius of convergence 10 by Theorem 2.

2. If $f(x) = \sum_{n=0}^{\infty} b_n x^n$ converges on $(-2, 2)$, then $\int f(x)\,dx = C + \sum_{n=0}^{\infty} \frac{b_n}{n+1} x^{n+1}$ has the same radius of convergence (by Theorem 2), but may not have the same interval of convergence—it may happen that the integrated series converges at an endpoint (or both endpoints).

3. Our goal is to write the function in the form $\frac{1}{1-r}$, and then use Equation 1 to represent the function as a sum of a power series. $f(x) = \frac{1}{1+x} = \frac{1}{1-(-x)} = \sum_{n=0}^{\infty} (-x)^n = \sum_{n=0}^{\infty} (-1)^n x^n$ with $|-x| < 1 \quad \Leftrightarrow \quad |x| < 1$, so $R = 1$ and $I = (-1, 1)$.

4. $f(x) = \frac{5}{1-4x^2} = 5\left(\frac{1}{1-4x^2}\right) = 5\sum_{n=0}^{\infty} (4x^2)^n = 5\sum_{n=0}^{\infty} 4^n x^{2n}$. The series converges when $|4x^2| < 1 \quad \Leftrightarrow$ $|x|^2 < \frac{1}{4} \quad \Leftrightarrow \quad |x| < \frac{1}{2}$, so $R = \frac{1}{2}$ and $I = \left(-\frac{1}{2}, \frac{1}{2}\right)$.

5. $f(x) = \frac{2}{3-x} = \frac{2}{3}\left(\frac{1}{1-x/3}\right) = \frac{2}{3}\sum_{n=0}^{\infty} \left(\frac{x}{3}\right)^n$ or, equivalently, $2\sum_{n=0}^{\infty} \frac{1}{3^{n+1}} x^n$. The series converges when $\left|\frac{x}{3}\right| < 1$, that is, when $|x| < 3$, so $R = 3$ and $I = (-3, 3)$.

6. $f(x) = \frac{4}{2x+3} = \frac{4}{3}\left(\frac{1}{1+2x/3}\right) = \frac{4}{3}\left(\frac{1}{1-(-2x/3)}\right) = \frac{4}{3}\sum_{n=0}^{\infty} \left(-\frac{2x}{3}\right)^n$ or, equivalently, $\sum_{n=0}^{\infty} (-1)^n \frac{2^{n+2}}{3^{n+1}} x^n$.

The series converges when $\left|-\frac{2x}{3}\right| < 1$, that is, when $|x| < \frac{3}{2}$, so $R = \frac{3}{2}$ and $I = \left(-\frac{3}{2}, \frac{3}{2}\right)$.

7. $f(x) = \frac{x^2}{x^4+16} = \frac{x^2}{16}\left(\frac{1}{1+x^4/16}\right) = \frac{x^2}{16}\left(\frac{1}{1-[-(x/2)]^4}\right) = \frac{x^2}{16}\sum_{n=0}^{\infty} \left[-\left(\frac{x}{2}\right)^4\right]^n$ or, equivalently, $\sum_{n=0}^{\infty} \frac{(-1)^n x^{4n+2}}{2^{4n+4}}$.

The series converges when $\left|-\left(\frac{x}{2}\right)^4\right| < 1 \quad \Rightarrow \quad \left|\frac{x}{2}\right| < 1 \quad \Rightarrow \quad |x| < 2$, so $R = 2$ and $I = (-2, 2)$.

8. $f(x) = \frac{x}{2x^2+1} = x\left(\frac{1}{1-(-2x^2)}\right) = x\sum_{n=0}^{\infty} (-2x^2)^n$ or, equivalently, $\sum_{n=0}^{\infty} (-1)^n 2^n x^{2n+1}$. The series converges when $|-2x^2| < 1 \quad \Rightarrow \quad |x^2| < \frac{1}{2} \quad \Rightarrow \quad |x| < \frac{1}{\sqrt{2}}$, so $R = \frac{1}{\sqrt{2}}$ and $I = \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

9. $f(x)=\dfrac{x-1}{x+2}=\dfrac{x+2-3}{x+2}=1-\dfrac{3}{x+2}=1-\dfrac{3/2}{x/2+1}=1-\dfrac{3}{2}\cdot\dfrac{1}{1-(-x/2)}$

$=1-\dfrac{3}{2}\sum_{n=0}^{\infty}\left(-\dfrac{x}{2}\right)^n=1-\dfrac{3}{2}-\dfrac{3}{2}\sum_{n=1}^{\infty}\left(-\dfrac{x}{2}\right)^n=-\dfrac{1}{2}-\sum_{n=1}^{\infty}\dfrac{(-1)^n\,3x^n}{2^{n+1}}.$

The geometric series $\sum_{n=0}^{\infty}\left(-\dfrac{x}{2}\right)^n$ converges when $\left|-\dfrac{x}{2}\right|<1 \quad\Leftrightarrow\quad |x|<2$, so $R=2$ and $I=(-2,2)$.

Alternatively, you could write $f(x)=1-3\left(\dfrac{1}{x+2}\right)$ and use the series for $\dfrac{1}{x+2}$ found in Example 2.

10. $f(x)=\dfrac{a}{x^2+a^2}\quad[a>0]\quad=\dfrac{a}{a^2}\left[\dfrac{1}{1-(-x^2/a^2)}\right]=\dfrac{1}{a}\sum_{n=0}^{\infty}\left(-\dfrac{x^2}{a^2}\right)^n=\sum_{n=0}^{\infty}\dfrac{(-1)^n x^{2n}}{a^{2n+1}}$. The geometric series $\sum_{n=0}^{\infty}\left(-\dfrac{x^2}{a^2}\right)^n$ converges when $\left|-\dfrac{x^2}{a^2}\right|<1 \quad\Leftrightarrow\quad |x|<a$, so $R=a$ and $I=(-a,a)$.

11. $f(x)=\dfrac{2x-4}{x^2-4x+3}=\dfrac{2x-4}{(x-1)(x-3)}=\dfrac{A}{x-1}+\dfrac{B}{x-3}\quad\Rightarrow\quad 2x-4=A(x-3)+B(x-1)$. Let $x=1$ to get $-2=-2A \quLeftrightarrow\quad A=1$ and $x=3$ to get $2=2B \quad\Leftrightarrow\quad B=1$. Thus,

$$\frac{2x-4}{x^2-4x+3}=\frac{1}{x-1}+\frac{1}{x-3}=\frac{-1}{1-x}+\frac{1}{-3}\left[\frac{1}{1-(x/3)}\right]=-\sum_{n=0}^{\infty}x^n-\frac{1}{3}\sum_{n=0}^{\infty}\left(\frac{x}{3}\right)^n=\sum_{n=0}^{\infty}\left(-1-\frac{1}{3^{n+1}}\right)x^n.$$

We represented f as the sum of two geometric series; the first converges for $x\in(-1,1)$ and the second converges for $x\in(-3,3)$. Thus, the sum converges for $x\in(-1,1)=I$.

12. $f(x)=\dfrac{2x+3}{x^2+3x+2}=\dfrac{2x+3}{(x+1)(x+2)}=\dfrac{A}{x+1}+\dfrac{B}{x+2}\quad\Rightarrow\quad 2x+3=A(x+2)+B(x+1)$. Let $x=-1$ to get $1=A$ and $x=-2$ to get $-1=-B \quad\Leftrightarrow\quad B=1$. Thus,

$$\frac{2x+3}{x^2+3x+2}=\frac{1}{x+1}+\frac{1}{x+2}=\frac{1}{1-(-x)}+\frac{1}{2}\left[\frac{1}{1-(-x/2)}\right]$$

$$=\sum_{n=0}^{\infty}(-x)^n+\frac{1}{2}\sum_{n=0}^{\infty}\left(-\frac{x}{2}\right)^n=\sum_{n=0}^{\infty}\left[(-1)^n\left(1+\frac{1}{2^{n+1}}\right)\right]x^n$$

We represented f as the sum of two geometric series; the first converges for $x\in(-1,1)$ and the second converges for $x\in(-2,2)$. Thus, the sum converges for $x\in(-1,1)=I$.

13. (a) $f(x)=\dfrac{1}{(1+x)^2}=\dfrac{d}{dx}\left(\dfrac{-1}{1+x}\right)=-\dfrac{d}{dx}\left[\sum_{n=0}^{\infty}(-1)^n x^n\right]$ [from Exercise 3]

$=\sum_{n=1}^{\infty}(-1)^{n+1}nx^{n-1}$ [from Theorem 2(i)] $=\sum_{n=0}^{\infty}(-1)^n(n+1)x^n$ with $R=1$.

In the last step, note that we *decreased* the initial value of the summation variable n by 1, and then *increased* each occurrence of n in the term by 1 [also note that $(-1)^{n+2}=(-1)^n$].

(b) $f(x)=\dfrac{1}{(1+x)^3}=-\dfrac{1}{2}\dfrac{d}{dx}\left[\dfrac{1}{(1+x)^2}\right]=-\dfrac{1}{2}\dfrac{d}{dx}\left[\sum_{n=0}^{\infty}(-1)^n(n+1)x^n\right]$ [from part (a)]

$=-\frac{1}{2}\sum_{n=1}^{\infty}(-1)^n(n+1)nx^{n-1}=\frac{1}{2}\sum_{n=0}^{\infty}(-1)^n(n+2)(n+1)x^n$ with $R=1$.

(c) $f(x)=\dfrac{x^2}{(1+x)^3}=x^2\cdot\dfrac{1}{(1+x)^3}=x^2\cdot\dfrac{1}{2}\sum_{n=0}^{\infty}(-1)^n(n+2)(n+1)x^n$ [from part (b)]

$=\dfrac{1}{2}\sum_{n=0}^{\infty}(-1)^n(n+2)(n+1)x^{n+2}$ [continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

To write the power series with x^n rather than x^{n+2}, we will *decrease* each occurrence of n in the term by 2 and *increase* the initial value of the summation variable by 2. This gives us $\frac{1}{2}\sum_{n=2}^{\infty}(-1)^n(n)(n-1)x^n$ with $R=1$.

14. (a) $\displaystyle\int \frac{1}{1-x}\,dx = -\ln(1-x)+C$ and

$$\int \frac{1}{1-x}\,dx = \int (1+x+x^2+\cdots)\,dx = \left(x+\frac{x^2}{2}+\frac{x^3}{3}+\cdots\right)+C = \sum_{n=1}^{\infty}\frac{x^n}{n}+C \text{ for } |x|<1.$$

So $-\ln(1-x) = \sum_{n=1}^{\infty}\frac{x^n}{n}+C$ and letting $x=0$ gives $0=C$. Thus, $f(x)=\ln(1-x)=-\sum_{n=1}^{\infty}\frac{x^n}{n}$ with $R=1$.

(b) $f(x)=x\ln(1-x)=-x\sum_{n=1}^{\infty}\frac{x^n}{n}=-\sum_{n=1}^{\infty}\frac{x^{n+1}}{n}$.

(c) Letting $x=\frac{1}{2}$ gives $\ln\frac{1}{2}=-\sum_{n=1}^{\infty}\frac{(1/2)^n}{n} \;\Rightarrow\; \ln 1-\ln 2=-\sum_{n=1}^{\infty}\frac{1^n}{n2^n} \;\Rightarrow\; \ln 2=\sum_{n=1}^{\infty}\frac{1}{n2^n}$.

15. $f(x)=\ln(5-x)=-\int\frac{dx}{5-x}=-\frac{1}{5}\int\frac{dx}{1-x/5}=-\frac{1}{5}\int\left[\sum_{n=0}^{\infty}\left(\frac{x}{5}\right)^n\right]dx=C-\frac{1}{5}\sum_{n=0}^{\infty}\frac{x^{n+1}}{5^n(n+1)}=C-\sum_{n=1}^{\infty}\frac{x^n}{n5^n}$

Putting $x=0$, we get $C=\ln 5$. The series converges for $|x/5|<1 \;\Leftrightarrow\; |x|<5$, so $R=5$.

16. $f(x)=x^2\tan^{-1}(x^3)=x^2\sum_{n=0}^{\infty}(-1)^n\frac{(x^3)^{2n+1}}{2n+1}$ [by Example 7] $=\sum_{n=0}^{\infty}(-1)^n\frac{x^{6n+3+2}}{2n+1}=\sum_{n=0}^{\infty}(-1)^n\frac{x^{6n+5}}{2n+1}$ for

$|x^3|<1 \;\Leftrightarrow\; |x|<1$, so $R=1$.

17. We know that $\frac{1}{1+4x}=\frac{1}{1-(-4x)}=\sum_{n=0}^{\infty}(-4x)^n$. Differentiating, we get

$\frac{-4}{(1+4x)^2}=\sum_{n=1}^{\infty}(-4)^n nx^{n-1}=\sum_{n=0}^{\infty}(-4)^{n+1}(n+1)x^n$, so

$$f(x)=\frac{x}{(1+4x)^2}=\frac{-x}{4}\cdot\frac{-4}{(1+4x)^2}=\frac{-x}{4}\sum_{n=0}^{\infty}(-4)^{n+1}(n+1)x^n=\sum_{n=0}^{\infty}(-1)^n4^n(n+1)x^{n+1}$$

for $|-4x|<1 \;\Leftrightarrow\; |x|<\frac{1}{4}$, so $R=\frac{1}{4}$.

18. $\frac{1}{2-x}=\frac{1}{2(1-x/2)}=\frac{1}{2}\sum_{n=0}^{\infty}\left(\frac{x}{2}\right)^n=\sum_{n=0}^{\infty}\frac{1}{2^{n+1}}x^n$. Now $\frac{d}{dx}\left(\frac{1}{2-x}\right)=\frac{d}{dx}\left(\sum_{n=0}^{\infty}\frac{1}{2^{n+1}}x^n\right) \;\Rightarrow$

$\frac{1}{(2-x)^2}=\sum_{n=1}^{\infty}\frac{1}{2^{n+1}}nx^{n-1}$ and $\frac{d}{dx}\left(\frac{1}{(2-x)^2}\right)=\frac{d}{dx}\left(\sum_{n=1}^{\infty}\frac{1}{2^{n+1}}nx^{n-1}\right) \;\Rightarrow$

$$\frac{2}{(2-x)^3}=\sum_{n=2}^{\infty}\frac{1}{2^{n+1}}n(n-1)x^{n-2}=\sum_{n=0}^{\infty}\frac{(n+2)(n+1)}{2^{n+3}}x^n.$$

Thus, $f(x)=\left(\frac{x}{2-x}\right)^3=\frac{x^3}{(2-x)^3}=\frac{x^3}{2}\cdot\frac{2}{(2-x)^3}=\frac{x^3}{2}\sum_{n=0}^{\infty}\frac{(n+2)(n+1)}{2^{n+3}}x^n=\sum_{n=0}^{\infty}\frac{(n+2)(n+1)}{2^{n+4}}x^{n+3}$

for $\left|\frac{x}{2}\right|<1 \;\Leftrightarrow\; |x|<2$, so $R=2$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

19. By Example 5, $\dfrac{1}{(1-x)^2} = \sum_{n=0}^{\infty}(n+1)x^n$. Thus,

$$f(x) = \frac{1+x}{(1-x)^2} = \frac{1}{(1-x)^2} + \frac{x}{(1-x)^2} = \sum_{n=0}^{\infty}(n+1)x^n + \sum_{n=0}^{\infty}(n+1)x^{n+1}$$

$$= \sum_{n=0}^{\infty}(n+1)x^n + \sum_{n=1}^{\infty} nx^n \qquad \text{[make the starting values equal]}$$

$$= 1 + \sum_{n=1}^{\infty}[(n+1)+n]x^n = 1 + \sum_{n=1}^{\infty}(2n+1)x^n = \sum_{n=0}^{\infty}(2n+1)x^n \text{ with } R = 1.$$

20. By Example 5, $\dfrac{1}{(1-x)^2} = \sum_{n=0}^{\infty}(n+1)x^n$, so

$\dfrac{d}{dx}\left(\dfrac{1}{(1-x)^2}\right) = \dfrac{d}{dx}\left(\sum_{n=0}^{\infty}(n+1)x^n\right) \quad \Rightarrow \quad \dfrac{2}{(1-x)^3} = \sum_{n=1}^{\infty}(n+1)nx^{n-1}$. Thus,

$$f(x) = \frac{x^2+x}{(1-x)^3} = \frac{x^2}{(1-x)^3} + \frac{x}{(1-x)^3} = \frac{x^2}{2}\cdot\frac{2}{(1-x)^3} + \frac{x}{2}\cdot\frac{2}{(1-x)^3}$$

$$= \frac{x^2}{2}\sum_{n=1}^{\infty}(n+1)nx^{n-1} + \frac{x}{2}\sum_{n=1}^{\infty}(n+1)nx^{n-1} = \sum_{n=1}^{\infty}\frac{(n+1)n}{2}x^{n+1} + \sum_{n=1}^{\infty}\frac{(n+1)n}{2}x^n$$

$$= \sum_{n=2}^{\infty}\frac{n(n-1)}{2}x^n + \sum_{n=1}^{\infty}\frac{(n+1)n}{2}x^n \qquad \text{[make the exponents on } x \text{ equal by changing an index]}$$

$$= \sum_{n=2}^{\infty}\frac{n^2-n}{2}x^n + x + \sum_{n=2}^{\infty}\frac{n^2+n}{2}x^n \qquad \text{[make the starting values equal]}$$

$$= x + \sum_{n=2}^{\infty} n^2x^n = \sum_{n=1}^{\infty} n^2x^n \text{ with } R = 1.$$

21. $f(x) = \dfrac{x^2}{x^2+1} = x^2\left(\dfrac{1}{1-(-x^2)}\right) = x^2\sum_{n=0}^{\infty}(-x^2)^n = \sum_{n=0}^{\infty}(-1)^n x^{2n+2}$. This series converges when $|-x^2| < 1 \quad \Leftrightarrow$ $x^2 < 1 \quad \Leftrightarrow \quad |x| < 1$, so $R = 1$. The partial sums are $s_1 = x^2$, $s_2 = s_1 - x^4$, $s_3 = s_2 + x^6$, $s_4 = s_3 - x^8$, $s_5 = s_4 + x^{10}, \ldots$. Note that s_1 corresponds to the first term of the infinite sum, regardless of the value of the summation variable and the value of the exponent. As n increases, $s_n(x)$ approximates f better on the interval of convergence, which is $(-1, 1)$.



22. From Example 6, we have $\ln(1+x) = \sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{x^n}{n}$ with $|x| < 1$, so $f(x) = \ln(1+x^4) = \sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{x^{4n}}{n}$ with $|x^4| < 1 \quad \Leftrightarrow \quad |x| < 1 \quad [R = 1]$. The partial sums are $s_1 = x^4$, $s_2 = s_1 - \frac{1}{2}x^8$, $s_3 = s_2 + \frac{1}{3}x^{12}$, $s_4 = s_3 - \frac{1}{4}x^{16}$, $s_5 = s_4 + \frac{1}{5}x^{20}, \ldots$. Note that s_1 corresponds to the first term of the infinite sum, regardless of the value of the summation variable and the value of the exponent. As n increases, $s_n(x)$ approximates f better on the interval of convergence, which is $[-1, 1]$. (When $x = \pm 1$, the series is the convergent alternating harmonic series.)



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $f(x) = \ln\left(\dfrac{1+x}{1-x}\right) = \ln(1+x) - \ln(1-x) = \displaystyle\int \frac{dx}{1+x} + \int \frac{dx}{1-x} = \int \frac{dx}{1-(-x)} + \int \frac{dx}{1-x}$

$$= \int \left[\sum_{n=0}^{\infty} (-1)^n x^n + \sum_{n=0}^{\infty} x^n\right] dx = \int [(1 - x + x^2 - x^3 + x^4 - \cdots) + (1 + x + x^2 + x^3 + x^4 + \cdots)]\, dx$$

$$= \int (2 + 2x^2 + 2x^4 + \cdots)\, dx = \int \sum_{n=0}^{\infty} 2x^{2n}\, dx = C + \sum_{n=0}^{\infty} \frac{2x^{2n+1}}{2n+1}$$

But $f(0) = \ln \frac{1}{1} = 0$, so $C = 0$ and we have $f(x) = \displaystyle\sum_{n=0}^{\infty} \frac{2x^{2n+1}}{2n+1}$ with $R = 1$. If $x = \pm 1$, then $f(x) = \pm 2 \displaystyle\sum_{n=0}^{\infty} \frac{1}{2n+1}$, which both diverge by the Limit Comparison Test with $b_n = \dfrac{1}{n}$.

The partial sums are $s_1 = \dfrac{2x}{1}$, $s_2 = s_1 + \dfrac{2x^3}{3}$, $s_3 = s_2 + \dfrac{2x^5}{5}$, ...



As n increases, $s_n(x)$ approximates f better on the interval of convergence, which is $(-1, 1)$.

24. $f(x) = \tan^{-1}(2x) = 2\displaystyle\int \frac{dx}{1+4x^2} = 2\int \sum_{n=0}^{\infty} (-1)^n \left(4x^2\right)^n\, dx = 2\int \sum_{n=0}^{\infty} (-1)^n 4^n x^{2n}\, dx$

$$= C + 2\sum_{n=0}^{\infty} \frac{(-1)^n 4^n x^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} \frac{(-1)^n 2^{2n+1} x^{2n+1}}{2n+1} \qquad [f(0) = \tan^{-1} 0 = 0, \text{ so } C = 0]$$

The series converges when $|4x^2| < 1 \;\Leftrightarrow\; |x| < \frac{1}{2}$, so $R = \frac{1}{2}$. If $x = \pm\frac{1}{2}$, then $f(x) = \displaystyle\sum_{n=0}^{\infty} (-1)^n \frac{1}{2n+1}$ and $f(x) = \displaystyle\sum_{n=0}^{\infty} (-1)^{n+1} \frac{1}{2n+1}$, respectively. Both series converge by the Alternating Series Test. The partial sums are $s_1 = \dfrac{2x}{1}$, $s_2 = s_1 - \dfrac{2^3 x^3}{3}$, $s_3 = s_2 + \dfrac{2^5 x^5}{5}$, ...



As n increases, $s_n(x)$ approximates f better on the interval of convergence, which is $\left[-\frac{1}{2}, \frac{1}{2}\right]$.

25. $\dfrac{t}{1-t^8} = t \cdot \dfrac{1}{1-t^8} = t \displaystyle\sum_{n=0}^{\infty} (t^8)^n = \sum_{n=0}^{\infty} t^{8n+1} \quad \Rightarrow \quad \int \frac{t}{1-t^8}\, dt = C + \sum_{n=0}^{\infty} \frac{t^{8n+2}}{8n+2}$. The series for $\dfrac{1}{1-t^8}$ converges when $|t^8| < 1 \;\Leftrightarrow\; |t| < 1$, so $R = 1$ for that series and also the series for $t/(1-t^8)$. By Theorem 2, the series for $\displaystyle\int \frac{t}{1-t^8}\, dt$ also has $R = 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

26. $\dfrac{t}{1+t^3} = t \cdot \dfrac{1}{1-(-t^3)} = t \sum_{n=0}^{\infty} (-t^3)^n = \sum_{n=0}^{\infty} (-1)^n t^{3n+1} \quad \Rightarrow \quad \displaystyle\int \frac{t}{1+t^3}\,dt = C + \sum_{n=0}^{\infty} (-1)^n \frac{t^{3n+2}}{3n+2}$. The series for $\dfrac{1}{1+t^3}$ converges when $|-t^3| < 1 \quad \Leftrightarrow \quad |t| < 1$, so $R = 1$ for that series and also for the series $\dfrac{t}{1+t^3}$. By Theorem 2, the series for $\displaystyle\int \frac{t}{1+t^3}\,dt$ also has $R = 1$.

27. From Example 6, $\ln(1+x) = \displaystyle\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$ for $|x| < 1$, so $x^2 \ln(1+x) = \displaystyle\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{n+2}}{n}$ and

$\displaystyle\int x^2 \ln(1+x)\,dx = C + \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{n+3}}{n(n+3)}$. $R = 1$ for the series for $\ln(1+x)$, so $R = 1$ for the series representing $x^2 \ln(1+x)$ as well. By Theorem 2, the series for $\displaystyle\int x^2 \ln(1+x)\,dx$ also has $R = 1$.

28. From Example 7, $\tan^{-1} x = \displaystyle\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$ for $|x| < 1$, so $\dfrac{\tan^{-1} x}{x} = \displaystyle\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{2n+1}$ and

$\displaystyle\int \frac{\tan^{-1} x}{x}\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)^2}$. $R = 1$ for the series for $\tan^{-1} x$, so $R = 1$ for the series representing $\dfrac{\tan^{-1} x}{x}$ as well. By Theorem 2, the series for $\displaystyle\int \frac{\tan^{-1} x}{x}\,dx$ also has $R = 1$.

29. $\dfrac{x}{1+x^3} = x\left[\dfrac{1}{1-(-x^3)}\right] = x \displaystyle\sum_{n=0}^{\infty} (-x^3)^n = \sum_{n=0}^{\infty} (-1)^n x^{3n+1} \quad \Rightarrow$

$\displaystyle\int \frac{x}{1+x^3}\,dx = \int \sum_{n=0}^{\infty} (-1)^n x^{3n+1}\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{3n+2}}{3n+2}$. Thus,

$$I = \int_0^{0.3} \frac{x}{1+x^3}\,dx = \left[\frac{x^2}{2} - \frac{x^5}{5} + \frac{x^8}{8} - \frac{x^{11}}{11} + \cdots\right]_0^{0.3} = \frac{(0.3)^2}{2} - \frac{(0.3)^5}{5} + \frac{(0.3)^8}{8} - \frac{(0.3)^{11}}{11} + \cdots.$$

The series is alternating, so if we use the first three terms, the error is at most $(0.3)^{11}/11 \approx 1.6 \times 10^{-7}$. So $I \approx (0.3)^2/2 - (0.3)^5/5 + (0.3)^8/8 \approx 0.044\,522$ to six decimal places.

30. We substitute $x/2$ for x in Example 7, and find that

$$\begin{aligned}\int \arctan(x/2)\,dx &= \int \sum_{n=0}^{\infty} (-1)^n \frac{(x/2)^{2n+1}}{2n+1}\,dx = \int \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2^{2n+1}(2n+1)}\,dx \\ &= C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+2}}{2^{2n+1}(2n+1)(2n+2)}\end{aligned}$$

Thus,

$$\begin{aligned}I = \int_0^{1/2} \arctan(x/2)\,dx &= \left[\frac{x^2}{2(1)(2)} - \frac{x^4}{2^3(3)(4)} + \frac{x^6}{2^5(5)(6)} - \frac{x^8}{2^7(7)(8)} + \frac{x^{10}}{2^9(9)(10)} - \cdots\right]_0^{1/2} \\ &= \frac{1}{2^3(1)(2)} - \frac{1}{2^7(3)(4)} + \frac{1}{2^{11}(5)(6)} - \frac{1}{2^{15}(7)(8)} + \frac{1}{2^{19}(9)(10)} - \cdots\end{aligned}$$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The series is alternating, so if we use four terms, the error is at most $1/(2^{19} \cdot 90) \approx 2.1 \times 10^{-8}$. So

$I \approx \dfrac{1}{16} - \dfrac{1}{1536} + \dfrac{1}{61{,}440} - \dfrac{1}{1{,}835{,}008} \approx 0.061\,865$ to six decimal places.

Remark: The sum of the first three terms gives us the same answer to six decimal places, but the error is at most $1/1{,}835{,}008 \approx 5.5 \times 10^{-7}$, slightly too large to guarantee the desired accuracy.

31. We substitute x^2 for x in Example 6, and find that

$$\int x \ln(1+x^2)\,dx = \int x \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(x^2)^n}{n}\,dx = \int \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n+1}}{n}\,dx = C + \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n+2}}{n(2n+2)}$$

Thus,

$$I \approx \int_0^{0.2} x \ln(1+x^2)\,dx = \left[\frac{x^4}{1(4)} - \frac{x^6}{2(6)} + \frac{x^8}{3(8)} - \frac{x^{10}}{4(10)} + \cdots\right]_0^{0.2} = \frac{(0.2)^4}{4} - \frac{(0.2)^6}{12} + \frac{(0.2)^8}{24} - \frac{(0.2)^{10}}{40} + \cdots$$

The series is alternating, so if we use two terms, the error is at most $(0.2)^8/24 \approx 1.1 \times 10^{-7}$. So

$I \approx \dfrac{(0.2)^4}{4} - \dfrac{(0.2)^6}{12} \approx 0.000\,395$ to six decimal places.

32.
$$\int_0^{0.3} \frac{x^2}{1+x^4}\,dx = \int_0^{0.3} x^2 \sum_{n=0}^{\infty} (-1)^n x^{4n}\,dx = \sum_{n=0}^{\infty} \left[\frac{(-1)^n x^{4n+3}}{4n+3}\right]_0^{0.3} = \sum_{n=0}^{\infty} \frac{(-1)^n\, 3^{4n+3}}{(4n+3)10^{4n+3}}$$
$$= \frac{3^3}{3 \times 10^3} - \frac{3^7}{7 \times 10^7} + \frac{3^{11}}{11 \times 10^{11}} - \cdots$$

The series is alternating, so if we use only two terms, the error is at most $\dfrac{3^{11}}{11 \times 10^{11}} \approx 0.000\,000\,16$. So, to six decimal places, $\displaystyle\int_0^{0.3} \frac{x^2}{1+x^4}\,dx \approx \frac{3^3}{3 \times 10^3} - \frac{3^7}{7 \times 10^7} \approx 0.008\,969$.

33. By Example 7, $\arctan x = x - \dfrac{x^3}{3} + \dfrac{x^5}{5} - \dfrac{x^7}{7} + \cdots$, so $\arctan 0.2 = 0.2 - \dfrac{(0.2)^3}{3} + \dfrac{(0.2)^5}{5} - \dfrac{(0.2)^7}{7} + \cdots$.

The series is alternating, so if we use three terms, the error is at most $\dfrac{(0.2)^7}{7} \approx 0.000\,002$.

Thus, to five decimal places, $\arctan 0.2 \approx 0.2 - \dfrac{(0.2)^3}{3} + \dfrac{(0.2)^5}{5} \approx 0.197\,40$.

34. $f(x) = \displaystyle\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} \quad \Rightarrow \quad f'(x) = \sum_{n=1}^{\infty} \frac{(-1)^n 2n x^{2n-1}}{(2n)!}$ [the first term disappears], so

$$f''(x) = \sum_{n=1}^{\infty} \frac{(-1)^n (2n)(2n-1) x^{2n-2}}{(2n)!} = \sum_{n=1}^{\infty} \frac{(-1)^n x^{2(n-1)}}{[2(n-1)]!} = -\sum_{n=0}^{\infty} \frac{(-1)^{n+1} x^{2n}}{(2n)!} \quad \text{[substituting } n+1 \text{ for } n\text{]}$$
$$= -\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} = -f(x) \quad \Rightarrow \quad f''(x) + f(x) = 0.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

35. (a) $J_0(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n}(n!)^2}$, $J_0'(x) = \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n-1}}{2^{2n}(n!)^2}$, and $J_0''(x) = \sum_{n=1}^{\infty} \frac{(-1)^n 2n(2n-1)x^{2n-2}}{2^{2n}(n!)^2}$, so

$$\begin{aligned} x^2J_0''(x) + xJ_0'(x) + x^2J_0(x) &= \sum_{n=1}^{\infty} \frac{(-1)^n 2n(2n-1)x^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n}}{2^{2n}(n!)^2} + \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+2}}{2^{2n}(n!)^2} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^n 2n(2n-1)x^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^{2n}}{2^{2n-2}[(n-1)!]^2} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^n 2n(2n-1)x^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^n(-1)^{-1}2^2n^2x^{2n}}{2^{2n}(n!)^2} \\ &= \sum_{n=1}^{\infty} (-1)^n \left[\frac{2n(2n-1) + 2n - 2^2n^2}{2^{2n}(n!)^2}\right] x^{2n} \\ &= \sum_{n=1}^{\infty} (-1)^n \left[\frac{4n^2 - 2n + 2n - 4n^2}{2^{2n}(n!)^2}\right] x^{2n} = 0 \end{aligned}$$

(b) $$\begin{aligned} \int_0^1 J_0(x)\,dx &= \int_0^1 \left[\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n}(n!)^2}\right] dx = \int_0^1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + \cdots\right) dx \\ &= \left[x - \frac{x^3}{3\cdot 4} + \frac{x^5}{5\cdot 64} - \frac{x^7}{7\cdot 2304} + \cdots\right]_0^1 = 1 - \frac{1}{12} + \frac{1}{320} - \frac{1}{16{,}128} + \cdots \end{aligned}$$

Since $\frac{1}{16{,}128} \approx 0.000062$, it follows from The Alternating Series Estimation Theorem that, correct to three decimal places, $\int_0^1 J_0(x)\,dx \approx 1 - \frac{1}{12} + \frac{1}{320} \approx 0.920$.

36. (a) $J_1(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}}$, $J_1'(x) = \sum_{n=0}^{\infty} \frac{(-1)^n (2n+1)\,x^{2n}}{n!\,(n+1)!\,2^{2n+1}}$, and $J_1''(x) = \sum_{n=1}^{\infty} \frac{(-1)^n (2n+1)(2n)\,x^{2n-1}}{n!\,(n+1)!\,2^{2n+1}}$.

$$\begin{aligned} &x^2J_1''(x) + xJ_1'(x) + (x^2-1)J_1(x) \\ &= \sum_{n=1}^{\infty} \frac{(-1)^n (2n+1)(2n)x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} + \sum_{n=0}^{\infty} \frac{(-1)^n (2n+1)x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} \\ &\qquad + \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+3}}{n!\,(n+1)!\,2^{2n+1}} - \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^n (2n+1)(2n)x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} + \sum_{n=0}^{\infty} \frac{(-1)^n (2n+1)x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} \\ &\qquad - \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n+1}}{(n-1)!\,n!\,2^{2n-1}} - \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} \qquad \left[\begin{array}{c}\text{Replace } n \text{ with } n-1 \\ \text{in the third term}\end{array}\right] \\ &= \frac{x}{2} - \frac{x}{2} + \sum_{n=1}^{\infty} (-1)^n \left[\frac{(2n+1)(2n) + (2n+1) - (n)(n+1)2^2 - 1}{n!\,(n+1)!\,2^{2n+1}}\right] x^{2n+1} = 0 \end{aligned}$$

(b) $J_0(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n}(n!)^2} \Rightarrow$

$$\begin{aligned} J_0'(x) &= \sum_{n=1}^{\infty} \frac{(-1)^n (2n)x^{2n-1}}{2^{2n}(n!)^2} = \sum_{n=0}^{\infty} \frac{(-1)^{n+1} 2(n+1)x^{2n+1}}{2^{2n+2}[(n+1)!]^2} \qquad [\text{Replace } n \text{ with } n+1] \\ &= -\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2^{2n+1}(n+1)!\,n!} \quad [\text{cancel 2 and } n+1\text{; take } -1 \text{ outside sum}] \quad = -J_1(x) \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

37. (a) $f(x)=\sum_{n=0}^{\infty}\frac{x^n}{n!} \Rightarrow f'(x)=\sum_{n=1}^{\infty}\frac{nx^{n-1}}{n!}=\sum_{n=1}^{\infty}\frac{x^{n-1}}{(n-1)!}=\sum_{n=0}^{\infty}\frac{x^n}{n!}=f(x)$

(b) By Theorem 9.4.2, the only solution to the differential equation $df(x)/dx=f(x)$ is $f(x)=Ke^x$, but $f(0)=1$, so $K=1$ and $f(x)=e^x$.

Or: We could solve the equation $df(x)/dx=f(x)$ as a separable differential equation.

38. $\frac{|\sin nx|}{n^2}\le\frac{1}{n^2}$, so $\sum_{n=1}^{\infty}\frac{\sin nx}{n^2}$ converges by the Comparison Test. $\frac{d}{dx}\left(\frac{\sin nx}{n^2}\right)=\frac{\cos nx}{n}$, so when $x=2k\pi$ [k an integer], $\sum_{n=1}^{\infty}f_n'(x)=\sum_{n=1}^{\infty}\frac{\cos(2kn\pi)}{n}=\sum_{n=1}^{\infty}\frac{1}{n}$, which diverges [harmonic series]. $f_n''(x)=-\sin nx$, so $\sum_{n=1}^{\infty}f_n''(x)=-\sum_{n=1}^{\infty}\sin nx$, which converges only if $\sin nx=0$, or $x=k\pi$ [k an integer].

39. If $a_n=\frac{x^n}{n^2}$, then by the Ratio Test, $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\frac{x^{n+1}}{(n+1)^2}\cdot\frac{n^2}{x^n}\right|=|x|\lim_{n\to\infty}\left(\frac{n}{n+1}\right)^2=|x|<1$ for convergence, so $R=1$. When $x=\pm1$, $\sum_{n=1}^{\infty}\left|\frac{x^n}{n^2}\right|=\sum_{n=1}^{\infty}\frac{1}{n^2}$ which is a convergent p-series ($p=2>1$), so the interval of convergence for f is $[-1,1]$. By Theorem 2, the radii of convergence of f' and f'' are both 1, so we need only check the endpoints. $f(x)=\sum_{n=1}^{\infty}\frac{x^n}{n^2} \Rightarrow f'(x)=\sum_{n=1}^{\infty}\frac{nx^{n-1}}{n^2}=\sum_{n=0}^{\infty}\frac{x^n}{n+1}$, and this series diverges for $x=1$ (harmonic series) and converges for $x=-1$ (Alternating Series Test), so the interval of convergence is $[-1,1)$. $f''(x)=\sum_{n=1}^{\infty}\frac{nx^{n-1}}{n+1}$ diverges at both 1 and -1 (Test for Divergence) since $\lim_{n\to\infty}\frac{n}{n+1}=1\neq0$, so its interval of convergence is $(-1,1)$.

40. (a) $\sum_{n=1}^{\infty}nx^{n-1}=\sum_{n=0}^{\infty}\frac{d}{dx}x^n=\frac{d}{dx}\left[\sum_{n=0}^{\infty}x^n\right]=\frac{d}{dx}\left[\frac{1}{1-x}\right]=-\frac{1}{(1-x)^2}(-1)=\frac{1}{(1-x)^2}$, $|x|<1$.

(b) (i) $\sum_{n=1}^{\infty}nx^n=x\sum_{n=1}^{\infty}nx^{n-1}=x\left[\frac{1}{(1-x)^2}\right]$ [from part (a)] $=\frac{x}{(1-x)^2}$ for $|x|<1$.

(ii) Put $x=\frac12$ in (i): $\sum_{n=1}^{\infty}\frac{n}{2^n}=\sum_{n=1}^{\infty}n\left(\frac12\right)^n=\frac{1/2}{(1-1/2)^2}=2$.

(c) (i) $\sum_{n=2}^{\infty}n(n-1)x^n=x^2\sum_{n=2}^{\infty}n(n-1)x^{n-2}=x^2\frac{d}{dx}\left[\sum_{n=1}^{\infty}nx^{n-1}\right]=x^2\frac{d}{dx}\frac{1}{(1-x)^2}$

$=x^2\frac{2}{(1-x)^3}=\frac{2x^2}{(1-x)^3}$ for $|x|<1$.

(ii) Put $x=\frac12$ in (i): $\sum_{n=2}^{\infty}\frac{n^2-n}{2^n}=\sum_{n=2}^{\infty}n(n-1)\left(\frac12\right)^n=\frac{2(1/2)^2}{(1-1/2)^3}=4$.

(iii) From (b)(ii) and (c)(ii), we have $\sum_{n=1}^{\infty}\frac{n^2}{2^n}=\sum_{n=1}^{\infty}\frac{n^2-n}{2^n}+\sum_{n=1}^{\infty}\frac{n}{2^n}=4+2=6$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

41. By Example 7, $\tan^{-1}x = \sum_{n=0}^{\infty}(-1)^n \frac{x^{2n+1}}{2n+1}$ for $|x| < 1$. In particular, for $x = \frac{1}{\sqrt{3}}$, we have $\frac{\pi}{6} = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right) = \sum_{n=0}^{\infty}(-1)^n \frac{(1/\sqrt{3})^{2n+1}}{2n+1} = \sum_{n=0}^{\infty}(-1)^n \left(\frac{1}{3}\right)^n \frac{1}{\sqrt{3}}\frac{1}{2n+1}$, so

$$\pi = \frac{6}{\sqrt{3}}\sum_{n=0}^{\infty}\frac{(-1)^n}{(2n+1)3^n} = 2\sqrt{3}\sum_{n=0}^{\infty}\frac{(-1)^n}{(2n+1)3^n}.$$

42. (a) $\displaystyle\int_0^{1/2}\frac{dx}{x^2-x+1} = \int_0^{1/2}\frac{dx}{(x-1/2)^2+3/4} \qquad \left[x-\frac{1}{2} = \frac{\sqrt{3}}{2}u,\ u = \frac{2}{\sqrt{3}}\left(x-\frac{1}{2}\right),\ dx = \frac{\sqrt{3}}{2}\,du\right]$

$$= \int_{-1/\sqrt{3}}^{0}\frac{(\sqrt{3}/2)\,du}{(3/4)(u^2+1)} = \frac{2\sqrt{3}}{3}\Big[\tan^{-1}u\Big]_{-1/\sqrt{3}}^{0} = \frac{2}{\sqrt{3}}\Big[0-\Big(-\frac{\pi}{6}\Big)\Big] = \frac{\pi}{3\sqrt{3}}$$

(b) $\dfrac{1}{x^3+1} = \dfrac{1}{(x+1)(x^2-x+1)} \Rightarrow$

$$\frac{1}{x^2-x+1} = (x+1)\left(\frac{1}{1+x^3}\right) = (x+1)\frac{1}{1-(-x^3)} = (x+1)\sum_{n=0}^{\infty}(-1)^n x^{3n}$$

$$= \sum_{n=0}^{\infty}(-1)^n x^{3n+1} + \sum_{n=0}^{\infty}(-1)^n x^{3n} \quad \text{for } |x|<1 \Rightarrow$$

$$\int\frac{dx}{x^2-x+1} = C + \sum_{n=0}^{\infty}(-1)^n\frac{x^{3n+2}}{3n+2} + \sum_{n=0}^{\infty}(-1)^n\frac{x^{3n+1}}{3n+1} \text{ for } |x|<1 \Rightarrow$$

$$\int_0^{1/2}\frac{dx}{x^2-x+1} = \sum_{n=0}^{\infty}(-1)^n\left[\frac{1}{4\cdot 8^n(3n+2)} + \frac{1}{2\cdot 8^n(3n+1)}\right] = \frac{1}{4}\sum_{n=0}^{\infty}\frac{(-1)^n}{8^n}\left(\frac{2}{3n+1}+\frac{1}{3n+2}\right).$$

By part (a), this equals $\dfrac{\pi}{3\sqrt{3}}$, so $\pi = \dfrac{3\sqrt{3}}{4}\displaystyle\sum_{n=0}^{\infty}\frac{(-1)^n}{8^n}\left(\frac{2}{3n+1}+\frac{1}{3n+2}\right)$.

11.10 Taylor and Maclaurin Series

1. Using Theorem 5 with $\displaystyle\sum_{n=0}^{\infty}b_n(x-5)^n$, $b_n = \dfrac{f^{(n)}(a)}{n!}$, so $b_8 = \dfrac{f^{(8)}(5)}{8!}$.

2. (a) Using Equation 6, a power series expansion of f at 1 must have the form $f(1) + f'(1)(x-1) + \cdots$. Comparing to the given series, $1.6 - 0.8(x-1) + \cdots$, we must have $f'(1) = -0.8$. But from the graph, $f'(1)$ is positive. Hence, the given series is *not* the Taylor series of f centered at 1.

(b) A power series expansion of f at 2 must have the form $f(2) + f'(2)(x-2) + \frac{1}{2}f''(2)(x-2)^2 + \cdots$. Comparing to the given series, $2.8 + 0.5(x-2) + 1.5(x-2)^2 - 0.1(x-2)^3 + \cdots$, we must have $\frac{1}{2}f''(2) = 1.5$; that is, $f''(2)$ is positive. But from the graph, f is concave downward near $x = 2$, so $f''(2)$ must be negative. Hence, the given series is *not* the Taylor series of f centered at 2.

3. Since $f^{(n)}(0) = (n+1)!$, Equation 7 gives the Maclaurin series

$\displaystyle\sum_{n=0}^{\infty}\frac{f^{(n)}(0)}{n!}x^n = \sum_{n=0}^{\infty}\frac{(n+1)!}{n!}x^n = \sum_{n=0}^{\infty}(n+1)x^n$. Applying the Ratio Test with $a_n = (n+1)x^n$ gives us

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(n+2)x^{n+1}}{(n+1)x^n}\right| = |x|\lim_{n\to\infty}\frac{n+2}{n+1} = |x|\cdot 1 = |x|$. For convergence, we must have $|x|<1$, so the radius of convergence $R=1$.

4. Since $f^{(n)}(4) = \dfrac{(-1)^n\, n!}{3^n(n+1)}$, Equation 6 gives the Taylor series

$\sum_{n=0}^{\infty}\dfrac{f^{(n)}(4)}{n!}(x-4)^n = \sum_{n=0}^{\infty}\dfrac{(-1)^n\, n!}{3^n(n+1)\, n!}(x-4)^n = \sum_{n=0}^{\infty}\dfrac{(-1)^n}{3^n(n+1)}(x-4)^n$, which is the Taylor series for f centered at 4. Apply the Ratio Test to find the radius of convergence R.

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^{n+1}(x-4)^{n+1}}{3^{n+1}(n+2)}\cdot\frac{3^n(n+1)}{(-1)^n(x-4)^n}\right| = \lim_{n\to\infty}\left|\frac{(-1)(x-4)(n+1)}{3(n+2)}\right|$$

$$= \frac{1}{3}|x-4|\lim_{n\to\infty}\frac{n+1}{n+2} = \frac{1}{3}|x-4|$$

For convergence, $\frac{1}{3}|x-4|<1 \quad\Leftrightarrow\quad |x-4|<3$, so $R=3$.

5.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	xe^x	0
1	$(x+1)e^x$	1
2	$(x+2)e^x$	2
3	$(x+3)e^x$	3
4	$(x+4)e^x$	4

Using Equation 6 with $n=0$ to 4 and $a=0$, we get

$$\sum_{n=0}^{4}\frac{f^{(n)}(0)}{n!}(x-0)^n = \frac{0}{0!}x^0 + \frac{1}{1!}x^1 + \frac{2}{2!}x^2 + \frac{3}{3!}x^3 + \frac{4}{4!}x^4$$

$$= x + x^2 + \tfrac{1}{2}x^3 + \tfrac{1}{6}x^4$$

6.

n	$f^{(n)}(x)$	$f^{(n)}(2)$
0	$\dfrac{1}{1+x}$	$\frac{1}{3}$
1	$-\dfrac{1}{(1+x)^2}$	$-\frac{1}{9}$
2	$\dfrac{2}{(1+x)^3}$	$\frac{2}{27}$
3	$-\dfrac{6}{(1+x)^4}$	$-\frac{6}{81}$

$$\sum_{n=0}^{3}\frac{f^{(n)}(2)}{n!}(x-2)^n = \frac{\frac{1}{3}}{0!}(x-2)^0 - \frac{\frac{1}{9}}{1!}(x-2)^1 + \frac{\frac{2}{27}}{2!}(x-2)^2 - \frac{\frac{6}{81}}{3!}(x-2)^3$$

$$= \tfrac{1}{3} - \tfrac{1}{9}(x-2) + \tfrac{1}{27}(x-2)^2 - \tfrac{1}{81}(x-2)^3$$

7.

n	$f^{(n)}(x)$	$f^{(n)}(8)$
0	$\sqrt[3]{x}$	2
1	$\dfrac{1}{3x^{2/3}}$	$\frac{1}{12}$
2	$-\dfrac{2}{9x^{5/3}}$	$-\frac{2}{288}$
3	$\dfrac{10}{27x^{8/3}}$	$\frac{10}{6912}$

$$\sum_{n=0}^{3}\frac{f^{(n)}(8)}{n!}(x-8)^n = \frac{2}{0!}(x-8)^0 + \frac{\frac{1}{12}}{1!}(x-8)^1 - \frac{\frac{2}{288}}{2!}(x-8)^2 + \frac{\frac{10}{6912}}{3!}(x-8)^3$$

$$= 2 + \tfrac{1}{12}(x-8) - \tfrac{1}{288}(x-8)^2 + \tfrac{5}{20{,}736}(x-8)^3$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

8.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$\ln x$	0
1	$1/x$	1
2	$-1/x^2$	-1
3	$2/x^3$	2
4	$-6/x^4$	-6

$$\sum_{n=0}^{4} \frac{f^{(n)}(1)}{n!}(x-1)^n = \frac{0}{0!}(x-1)^0 + \frac{1}{1!}(x-1)^1 - \frac{1}{2!}(x-1)^2 + \frac{2}{3!}(x-1)^3 - \frac{6}{4!}(x-1)^4$$
$$= (x-1) - \tfrac{1}{2}(x-1)^2 + \tfrac{1}{3}(x-1)^3 - \tfrac{1}{4}(x-1)^4$$

9.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/6)$
0	$\sin x$	$1/2$
1	$\cos x$	$\sqrt{3}/2$
2	$-\sin x$	$-1/2$
3	$-\cos x$	$-\sqrt{3}/2$

$$\sum_{n=0}^{3} \frac{f^{(n)}(\pi/6)}{n!}\left(x-\frac{\pi}{6}\right)^n = \frac{1/2}{0!}\left(x-\frac{\pi}{6}\right)^0 + \frac{\sqrt{3}/2}{1!}\left(x-\frac{\pi}{6}\right)^1 - \frac{1/2}{2!}\left(x-\frac{\pi}{6}\right)^2 - \frac{\sqrt{3}/2}{3!}\left(x-\frac{\pi}{6}\right)^3$$
$$= \frac{1}{2} + \frac{\sqrt{3}}{2}\left(x-\frac{\pi}{6}\right) - \frac{1}{4}\left(x-\frac{\pi}{6}\right)^2 - \frac{\sqrt{3}}{12}\left(x-\frac{\pi}{6}\right)^3$$

10.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\cos^2 x$	1
1	$-2\cos x \sin x = -\sin 2x$	0
2	$-2\cos 2x$	-2
3	$4\sin 2x$	0
4	$8\cos 2x$	8
5	$-16\sin 2x$	0
6	$-32\cos 2x$	-32

$$\sum_{n=0}^{6} \frac{f^{(n)}(0)}{n!}(x-0)^n = \frac{1}{0!}x^0 - \frac{2}{2!}x^2 + \frac{8}{4!}x^4 - \frac{32}{6!}x^6$$
$$= 1 - x^2 + \tfrac{1}{3}x^4 - \tfrac{2}{45}x^6$$

11.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$(1-x)^{-2}$	1
1	$2(1-x)^{-3}$	2
2	$6(1-x)^{-4}$	6
3	$24(1-x)^{-5}$	24
4	$120(1-x)^{-6}$	120
⋮	⋮	⋮

$$(1-x)^{-2} = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \cdots$$
$$= 1 + 2x + \tfrac{6}{2}x^2 + \tfrac{24}{6}x^3 + \tfrac{120}{24}x^4 + \cdots$$
$$= 1 + 2x + 3x^2 + 4x^3 + 5x^4 + \cdots = \sum_{n=0}^{\infty}(n+1)x^n$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(n+2)x^{n+1}}{(n+1)x^n}\right| = |x|\lim_{n\to\infty}\frac{n+2}{n+1} = |x|\,(1) = |x| < 1$$

for convergence, so $R = 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\ln(1+x)$	0
1	$(1+x)^{-1}$	1
2	$-(1+x)^{-2}$	-1
3	$2(1+x)^{-3}$	2
4	$-6(1+x)^{-4}$	-6
5	$24(1+x)^{-5}$	24
⋮	⋮	⋮

$$\ln(1+x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \frac{f^{(5)}(0)}{5!}x^5 + \cdots$$
$$= 0 + x - \tfrac{1}{2}x^2 + \tfrac{2}{6}x^3 - \tfrac{6}{24}x^4 + \tfrac{24}{120}x^5 - \cdots$$
$$= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \frac{x^5}{5} - \cdots = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} x^n$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{x^{n+1}}{n+1}\cdot\frac{n}{x^n}\right| = \lim_{n\to\infty}\frac{|x|}{1+1/n} = |x| < 1 \text{ for convergence,}$$

so $R = 1$.

Notice that the answer agrees with the entry for $\ln(1+x)$ in Table 1, but we obtained it by a different method. (Compare with Example 11.9.6.)

13.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\cos x$	1
1	$-\sin x$	0
2	$-\cos x$	-1
3	$\sin x$	0
4	$\cos x$	1
⋮	⋮	⋮

$$\cos x = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \cdots$$
$$= 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \cdots$$
$$= \sum_{n=0}^{\infty}(-1)^n\frac{x^{2n}}{(2n)!} \qquad \text{[Agrees with (16).]}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{x^{2n+2}}{(2n+2)!}\cdot\frac{(2n)!}{x^{2n}}\right| = \lim_{n\to\infty}\frac{x^2}{(2n+2)(2n+1)} = 0 < 1$$

for all x, so $R = \infty$.

14.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	e^{-2x}	1
1	$-2e^{-2x}$	-2
2	$4e^{-2x}$	4
3	$-8e^{-2x}$	-8
4	$16e^{-2x}$	16
⋮	⋮	⋮

$$e^{-2x} = \sum_{n=0}^{\infty}\frac{f^{(n)}(0)}{n!}x^n = \sum_{n=0}^{\infty}\frac{(-2)^n}{n!}x^n.$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-2)^{n+1}x^{n+1}}{(n+1)!}\cdot\frac{n!}{(-2)^n x^n}\right| = \lim_{n\to\infty}\frac{2|x|}{n+1}$$
$$= 0 < 1 \quad \text{for all } x, \text{ so } R = \infty.$$

15.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	2^x	1
1	$2^x(\ln 2)$	$\ln 2$
2	$2^x(\ln 2)^2$	$(\ln 2)^2$
3	$2^x(\ln 2)^3$	$(\ln 2)^3$
4	$2^x(\ln 2)^4$	$(\ln 2)^4$
⋮	⋮	⋮

$$2^x = \sum_{n=0}^{\infty}\frac{f^{(n)}(0)}{n!}x^n = \sum_{n=0}^{\infty}\frac{(\ln 2)^n}{n!}x^n.$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(\ln 2)^{n+1}x^{n+1}}{(n+1)!}\cdot\frac{n!}{(\ln 2)^n x^n}\right|$$
$$= \lim_{n\to\infty}\frac{(\ln 2)|x|}{n+1} = 0 < 1 \quad \text{for all } x, \text{ so } R = \infty.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

16.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$x\cos x$	0
1	$-x\sin x+\cos x$	1
2	$-x\cos x-2\sin x$	0
3	$x\sin x-3\cos x$	-3
4	$x\cos x+4\sin x$	0
5	$-x\sin x+5\cos x$	5
6	$-x\cos x-6\sin x$	0
7	$x\sin x-7\cos x$	7
$\vdots$	$\vdots$	$\vdots$

$$\begin{aligned} x\cos x &= f(0)+f'(0)x+\frac{f''(0)}{2!}x^2+\frac{f'''(0)}{3!}x^3+\frac{f^{(4)}(0)}{4!}x^4+\cdots \\ &= 0+1x+0-\frac{3}{3!}x^3+0+\frac{5}{5!}x^5+0-\frac{7}{7!}x^7+\cdots \\ &= x-\frac{1}{2!}x^3+\frac{1}{4!}x^5-\frac{1}{6!}x^7+\cdots \\ &= \sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n)!}x^{2n+1} \end{aligned}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^{n+1}x^{2n+3}}{(2n+2)!}\cdot\frac{(2n)!}{(-1)^n x^{2n+1}}\right|$$

$$= \lim_{n\to\infty}\frac{x^2}{(2n+2)(2n+1)} = 0<1 \quad \text{for all } x, \text{ so } R=\infty.$$

17.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sinh x$	0
1	$\cosh x$	1
2	$\sinh x$	0
3	$\cosh x$	1
4	$\sinh x$	0
$\vdots$	$\vdots$	$\vdots$

$$f^{(n)}(0)=\begin{cases} 0 & \text{if } n \text{ is even} \\ 1 & \text{if } n \text{ is odd} \end{cases} \qquad \text{so } \sinh x=\sum_{n=0}^{\infty}\frac{x^{2n+1}}{(2n+1)!}.$$

Use the Ratio Test to find R. If $a_n=\dfrac{x^{2n+1}}{(2n+1)!}$, then

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{x^{2n+3}}{(2n+3)!}\cdot\frac{(2n+1)!}{x^{2n+1}}\right| = x^2\cdot\lim_{n\to\infty}\frac{1}{(2n+3)(2n+2)}$$

$$= 0<1 \quad \text{for all } x, \text{ so } R=\infty.$$

18.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\cosh x$	1
1	$\sinh x$	0
2	$\cosh x$	1
3	$\sinh x$	0
$\vdots$	$\vdots$	$\vdots$

$$f^{(n)}(0)=\begin{cases} 1 & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases} \qquad \text{so } \cosh x=\sum_{n=0}^{\infty}\frac{x^{2n}}{(2n)!}.$$

Use the Ratio Test to find R. If $a_n=\dfrac{x^{2n}}{(2n)!}$, then

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{x^{2n+2}}{(2n+2)!}\cdot\frac{(2n)!}{x^{2n}}\right| = x^2\cdot\lim_{n\to\infty}\frac{1}{(2n+2)(2n+1)}$$

$$= 0<1 \quad \text{for all } x, \text{ so } R=\infty.$$

19.

n	$f^{(n)}(x)$	$f^{(n)}(2)$
0	x^5+2x^3+x	50
1	$5x^4+6x^2+1$	105
2	$20x^3+12x$	184
3	$60x^2+12$	252
4	$120x$	240
5	120	120
6	0	0
7	0	0
$\vdots$	$\vdots$	$\vdots$

$f^{(n)}(x)=0$ for $n\geq 6$, so f has a finite expansion about $a=2$.

$$\begin{aligned} f(x) &= x^5+2x^3+x=\sum_{n=0}^{5}\frac{f^{(n)}(2)}{n!}(x-2)^n \\ &= \frac{50}{0!}(x-2)^0+\frac{105}{1!}(x-2)^1+\frac{184}{2!}(x-2)^2+\frac{252}{3!}(x-2)^3 \\ &\quad +\frac{240}{4!}(x-2)^4+\frac{120}{5!}(x-2)^5 \\ &= 50+105(x-2)+92(x-2)^2+42(x-2)^3 \\ &\quad +10(x-2)^4+(x-2)^5 \end{aligned}$$

A finite series converges for all x, so $R=\infty$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20.

n	$f^{(n)}(x)$	$f^{(n)}(-2)$
0	$x^6 - x^4 + 2$	50
1	$6x^5 - 4x^3$	-160
2	$30x^4 - 12x^2$	432
3	$120x^3 - 24x$	-912
4	$360x^2 - 24$	1416
5	$720x$	-1440
6	720	720
7	0	0
8	0	0
⋮	⋮	⋮

$f^{(n)}(x) = 0$ for $n \geq 7$, so f has a finite expansion about $a = -2$.

$$\begin{aligned} f(x) = x^6 - x^4 + 2 &= \sum_{n=0}^{6} \frac{f^{(n)}(-2)}{n!}(x+2)^n \\ &= \frac{50}{0!}(x+2)^0 - \frac{160}{1!}(x+2)^1 + \frac{432}{2!}(x+2)^2 - \frac{912}{3!}(x+2)^3 \\ &\quad + \frac{1416}{4!}(x+2)^4 - \frac{1440}{5!}(x+2)^5 + \frac{720}{6!}(x+2)^6 \\ &= 50 - 160(x+2) + 216(x+2)^2 - 152(x+2)^3 + 59(x+2)^4 - 12(x+2)^5 + (x+2)^6 \end{aligned}$$

A finite series converges for all x, so $R = \infty$.

21.

n	$f^{(n)}(x)$	$f^{(n)}(2)$
0	$\ln x$	$\ln 2$
1	$1/x$	$1/2$
2	$-1/x^2$	$-1/2^2$
3	$2/x^3$	$2/2^3$
4	$-6/x^4$	$-6/2^4$
5	$24/x^5$	$24/2^5$
⋮	⋮	⋮

$$\begin{aligned} f(x) = \ln x &= \sum_{n=0}^{\infty} \frac{f^{(n)}(2)}{n!}(x-2)^n \\ &= \frac{\ln 2}{0!}(x-2)^0 + \frac{1}{1!\,2^1}(x-2)^1 + \frac{-1}{2!\,2^2}(x-2)^2 + \frac{2}{3!\,2^3}(x-2)^3 \\ &\quad + \frac{-6}{4!\,2^4}(x-2)^4 + \frac{24}{5!\,2^5}(x-2)^5 + \cdots \\ &= \ln 2 + \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(n-1)!}{n!\,2^n}(x-2)^n \\ &= \ln 2 + \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n\,2^n}(x-2)^n \end{aligned}$$

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^{n+2}(x-2)^{n+1}}{(n+1)\,2^{n+1}} \cdot \frac{n\,2^n}{(-1)^{n+1}(x-2)^n}\right| = \lim_{n\to\infty} \left|\frac{(-1)(x-2)n}{(n+1)2}\right| = \lim_{n\to\infty} \left(\frac{n}{n+1}\right)\frac{|x-2|}{2}$$

$$= \frac{|x-2|}{2} < 1 \quad \text{for convergence, so } |x-2| < 2 \text{ and } R = 2.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

22.

n	$f^{(n)}(x)$	$f^{(n)}(-3)$
0	$1/x$	$-1/3$
1	$-1/x^2$	$-1/3^2$
2	$2/x^3$	$-2/3^3$
3	$-6/x^4$	$-6/3^4$
4	$24/x^5$	$-24/3^5$
⋮	⋮	⋮

$$f(x)=\frac{1}{x}=\sum_{n=0}^{\infty}\frac{f^{(n)}(-3)}{n!}(x+3)^n$$
$$=\frac{-1/3}{0!}(x+3)^0+\frac{-1/3^2}{1!}(x+3)^1+\frac{-2/3^3}{2!}(x+3)^2$$
$$+\frac{-6/3^4}{3!}(x+3)^3+\frac{-24/3^5}{4!}(x+3)^4+\cdots$$
$$=\sum_{n=0}^{\infty}\frac{-n!/3^{n+1}}{n!}(x+3)^n=-\sum_{n=0}^{\infty}\frac{(x+3)^n}{3^{n+1}}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\frac{(x+3)^{n+1}}{3^{n+2}}\cdot\frac{3^{n+1}}{(x+3)^n}\right|=\lim_{n\to\infty}\frac{|x+3|}{3}=\frac{|x+3|}{3}<1 \quad \text{for convergence,}$$

so $|x+3|<3$ and $R=3$.

23.

n	$f^{(n)}(x)$	$f^{(n)}(3)$
0	e^{2x}	e^6
1	$2e^{2x}$	$2e^6$
2	2^2e^{2x}	$4e^6$
3	2^3e^{2x}	$8e^6$
4	2^4e^{2x}	$16e^6$
⋮	⋮	⋮

$$f(x)=e^{2x}=\sum_{n=0}^{\infty}\frac{f^{(n)}(3)}{n!}(x-3)^n$$
$$=\frac{e^6}{0!}(x-3)^0+\frac{2e^6}{1!}(x-3)^1+\frac{4e^6}{2!}(x-3)^2$$
$$+\frac{8e^6}{3!}(x-3)^3+\frac{16e^6}{4!}(x-3)^4+\cdots$$
$$=\sum_{n=0}^{\infty}\frac{2^ne^6}{n!}(x-3)^n$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\frac{2^{n+1}e^6(x-3)^{n+1}}{(n+1)!}\cdot\frac{n!}{2^ne^6(x-3)^n}\right|=\lim_{n\to\infty}\frac{2|x-3|}{n+1}=0<1 \quad \text{for all } x, \text{ so } R=\infty.$$

24.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/2)$
0	$\cos x$	0
1	$-\sin x$	-1
2	$-\cos x$	0
3	$\sin x$	1
4	$\cos x$	0
5	$-\sin x$	-1
6	$\cos x$	0
7	$\sin x$	1
⋮	⋮	⋮

$$f(x)=\cos x=\sum_{n=0}^{\infty}\frac{f^{(n)}(\pi/2)}{n!}\left(x-\frac{\pi}{2}\right)^n$$
$$=\frac{-1}{1!}\left(x-\frac{\pi}{2}\right)^1+\frac{1}{3!}\left(x-\frac{\pi}{2}\right)^3+\frac{-1}{5!}\left(x-\frac{\pi}{2}\right)^5+\frac{1}{7!}\left(x-\frac{\pi}{2}\right)^7+\cdots$$
$$=\sum_{n=0}^{\infty}\frac{(-1)^{n+1}}{(2n+1)!}\left(x-\frac{\pi}{2}\right)^{2n+1}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\frac{(-1)^{n+2}\left(x-\frac{\pi}{2}\right)^{2n+3}}{(2n+3)!}\cdot\frac{(2n+1)!}{(-1)^{n+1}\left(x-\frac{\pi}{2}\right)^{2n+1}}\right|$$
$$=\lim_{n\to\infty}\frac{\left(x-\frac{\pi}{2}\right)^2}{(2n+3)(2n+2)}=0<1 \quad \text{for all } x, \text{ so } R=\infty.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25.

n	$f^{(n)}(x)$	$f^{(n)}(\pi)$
0	$\sin x$	0
1	$\cos x$	-1
2	$-\sin x$	0
3	$-\cos x$	1
4	$\sin x$	0
5	$\cos x$	-1
6	$-\sin x$	0
7	$-\cos x$	1
⋮	⋮	⋮

$$f(x) = \sin x = \sum_{n=0}^{\infty} \frac{f^{(n)}(\pi)}{n!}(x-\pi)^n$$

$$= \frac{-1}{1!}(x-\pi)^1 + \frac{1}{3!}(x-\pi)^3 + \frac{-1}{5!}(x-\pi)^5 + \frac{1}{7!}(x-\pi)^7 + \cdots$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n+1}}{(2n+1)!}(x-\pi)^{2n+1}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^{n+2}(x-\pi)^{2n+3}}{(2n+3)!}\cdot\frac{(2n+1)!}{(-1)^{n+1}(x-\pi)^{2n+1}}\right|$$

$$= \lim_{n\to\infty}\frac{(x-\pi)^2}{(2n+3)(2n+2)} = 0 < 1 \quad \text{for all } x, \text{ so } R = \infty.$$

26.

n	$f^{(n)}(x)$	$f^{(n)}(16)$
0	$\sqrt{x}$	4
1	$\frac{1}{2}x^{-1/2}$	$\frac{1}{2}\cdot\frac{1}{4}$
2	$-\frac{1}{4}x^{-3/2}$	$-\frac{1}{4}\cdot\frac{1}{4^3}$
3	$\frac{3}{8}x^{-5/2}$	$\frac{3}{8}\cdot\frac{1}{4^5}$
4	$-\frac{15}{16}x^{-7/2}$	$-\frac{15}{16}\cdot\frac{1}{4^7}$
⋮	⋮	⋮

$$f(x) = \sqrt{x} = \sum_{n=0}^{\infty} \frac{f^{(n)}(16)}{n!}(x-16)^n$$

$$= \frac{4}{0!}(x-16)^0 + \frac{1}{2}\cdot\frac{1}{4}\cdot\frac{1}{1!}(x-16)^1 - \frac{1}{4}\cdot\frac{1}{4^3}\cdot\frac{1}{2!}(x-16)^2$$

$$+ \frac{3}{8}\cdot\frac{1}{4^5}\cdot\frac{1}{3!}(x-16)^3 - \frac{15}{16}\cdot\frac{1}{4^7}\cdot\frac{1}{4!}(x-16)^4 + \cdots$$

$$= 4 + \frac{1}{8}(x-16) + \sum_{n=2}^{\infty}(-1)^{n-1}\frac{1\cdot3\cdot5\cdot\cdots\cdot(2n-3)}{2^n 4^{2n-1}\,n!}(x-16)^n$$

$$= 4 + \frac{1}{8}(x-16) + \sum_{n=2}^{\infty}(-1)^{n-1}\frac{1\cdot3\cdot5\cdot\cdots\cdot(2n-3)}{2^{5n-2}\,n!}(x-16)^n$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^n\,1\cdot3\cdot5\cdot\cdots\cdot(2n-1)(x-16)^{n+1}}{2^{5n+3}(n+1)!}\cdot\frac{2^{5n-2}n!}{(-1)^{n-1}\,1\cdot3\cdot5\cdot\cdots\cdot(2n-3)(x-16)^n}\right|$$

$$= \lim_{n\to\infty}\frac{(2n-1)\,|x-16|}{2^5(n+1)} = \frac{|x-16|}{32}\lim_{n\to\infty}\frac{2-1/n}{1+1/n} = \frac{|x-16|}{32}\cdot 2$$

$$= \frac{|x-16|}{16} < 1 \quad \text{for convergence, so } |x-16| < 16 \text{ and } R = 16.$$

27. If $f(x) = \cos x$, then $f^{(n+1)}(x) = \pm\sin x$ or $\pm\cos x$. In each case, $\left|f^{(n+1)}(x)\right| \le 1$, so by Formula 9 with $a = 0$ and $M = 1$, $|R_n(x)| \le \dfrac{1}{(n+1)!}|x|^{n+1}$. Thus, $|R_n(x)| \to 0$ as $n \to \infty$ by Equation 10. So $\lim\limits_{n\to\infty} R_n(x) = 0$ and, by Theorem 8, the series in Exercise 13 represents $\cos x$ for all x.

28. If $f(x) = \sin x$, then $f^{(n+1)}(x) = \pm\sin x$ or $\pm\cos x$. In each case, $\left|f^{(n+1)}(x)\right| \le 1$, so by Formula 9 with $a = 0$ and $M = 1$, $|R_n(x)| \le \dfrac{1}{(n+1)!}|x-\pi|^{n+1}$. Thus, $|R_n(x)| \to 0$ as $n \to \infty$ by Equation 10. So $\lim\limits_{n\to\infty} R_n(x) \to 0$ and, by Theorem 8, the series in Exercise 25 represents $\sin x$ for all x.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

29. If $f(x) = \sinh x$, then for all n, $f^{(n+1)}(x) = \cosh x$ or $\sinh x$. Since $|\sinh x| < |\cosh x| = \cosh x$ for all x, we have $\left|f^{(n+1)}(x)\right| \le \cosh x$ for all n. If d is any positive number and $|x| \le d$, then $\left|f^{(n+1)}(x)\right| \le \cosh x \le \cosh d$, so by Formula 9 with $a = 0$ and $M = \cosh d$, we have $|R_n(x)| \le \dfrac{\cosh d}{(n+1)!}\,|x|^{n+1}$. It follows that $|R_n(x)| \to 0$ as $n \to \infty$ for $|x| \le d$ (by Equation 10). But d was an arbitrary positive number. So by Theorem 8, the series represents $\sinh x$ for all x.

30. If $f(x) = \cosh x$, then for all n, $f^{(n+1)}(x) = \cosh x$ or $\sinh x$. Since $|\sinh x| < |\cosh x| = \cosh x$ for all x, we have $\left|f^{(n+1)}(x)\right| \le \cosh x$ for all n. If d is any positive number and $|x| \le d$, then $\left|f^{(n+1)}(x)\right| \le \cosh x \le \cosh d$, so by Formula 9 with $a = 0$ and $M = \cosh d$, we have $|R_n(x)| \le \dfrac{\cosh d}{(n+1)!}\,|x|^{n+1}$. It follows that $|R_n(x)| \to 0$ as $n \to \infty$ for $|x| \le d$ (by Equation 10). But d was an arbitrary positive number. So by Theorem 8, the series represents $\cosh x$ for all x.

31.
$$\sqrt[4]{1-x} = [1+(-x)]^{1/4} = \sum_{n=0}^{\infty} \binom{1/4}{n}(-x)^n = 1 + \tfrac{1}{4}(-x) + \frac{\frac{1}{4}\left(-\frac{3}{4}\right)}{2!}(-x)^2 + \frac{\frac{1}{4}\left(-\frac{3}{4}\right)\left(-\frac{7}{4}\right)}{3!}(-x)^3 + \cdots$$
$$= 1 - \frac{1}{4}x + \sum_{n=2}^{\infty} \frac{(-1)^{n-1}(-1)^n \cdot [3 \cdot 7 \cdot \cdots \cdot (4n-5)]}{4^n \cdot n!}x^n$$
$$= 1 - \frac{1}{4}x - \sum_{n=2}^{\infty} \frac{3 \cdot 7 \cdot \cdots \cdot (4n-5)}{4^n \cdot n!}x^n$$

and $|-x| < 1 \quad \Leftrightarrow \quad |x| < 1$, so $R = 1$.

32.
$$\sqrt[3]{8+x} = \sqrt[3]{8\left(1+\frac{x}{8}\right)} = 2\left(1+\frac{x}{8}\right)^{1/3} = 2\sum_{n=0}^{\infty}\binom{1/3}{n}\left(\frac{x}{8}\right)^n$$
$$= 2\left[1 + \frac{1}{3}\left(\frac{x}{8}\right) + \frac{\frac{1}{3}\left(-\frac{2}{3}\right)}{2!}\left(\frac{x}{8}\right)^2 + \frac{\frac{1}{3}\left(-\frac{2}{3}\right)\left(-\frac{5}{3}\right)}{3!}\left(\frac{x}{8}\right)^3 + \cdots\right]$$
$$= 2\left[1 + \frac{1}{24}x + \sum_{n=2}^{\infty}\frac{(-1)^{n-1}\cdot[2\cdot 5\cdot\cdots\cdot(3n-4)]}{3^n\cdot 8^n\cdot n!}x^n\right]$$
$$= 2 + \frac{1}{12}x + 2\sum_{n=2}^{\infty}\frac{(-1)^{n-1}[2\cdot 5\cdot\cdots\cdot(3n-4)]}{24^n\cdot n!}x^n$$

and $\left|\dfrac{x}{8}\right| < 1 \quad \Leftrightarrow \quad |x| < 8$, so $R = 8$.

33. $\dfrac{1}{(2+x)^3} = \dfrac{1}{[2(1+x/2)]^3} = \dfrac{1}{8}\left(1+\dfrac{x}{2}\right)^{-3} = \dfrac{1}{8}\displaystyle\sum_{n=0}^{\infty}\binom{-3}{n}\left(\frac{x}{2}\right)^n$. The binomial coefficient is

$$\binom{-3}{n} = \frac{(-3)(-4)(-5)\cdot\cdots\cdot(-3-n+1)}{n!} = \frac{(-3)(-4)(-5)\cdot\cdots\cdot[-(n+2)]}{n!}$$
$$= \frac{(-1)^n\cdot 2\cdot 3\cdot 4\cdot 5\cdot\cdots\cdot(n+1)(n+2)}{2\cdot n!} = \frac{(-1)^n(n+1)(n+2)}{2}$$

Thus, $\dfrac{1}{(2+x)^3} = \dfrac{1}{8}\displaystyle\sum_{n=0}^{\infty}\frac{(-1)^n(n+1)(n+2)}{2}\frac{x^n}{2^n} = \sum_{n=0}^{\infty}\frac{(-1)^n(n+1)(n+2)x^n}{2^{n+1}}$ for $\left|\dfrac{x}{2}\right| < 1 \quad \Leftrightarrow \quad |x| < 2$, so $R = 2$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. $(1+x)^{3/4} = \sum_{n=0}^{\infty} \binom{\frac{3}{4}}{n} x^n = 1 + \frac{3}{4}x + \frac{\frac{3}{4}\left(-\frac{1}{4}\right)}{2!}x^2 + \frac{\frac{3}{4}\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)}{3!}x^3 + \cdots$

$$= 1 + \frac{3}{4}x + \sum_{n=2}^{\infty} \frac{(-1)^{n-1} \cdot 3 \cdot [1 \cdot 5 \cdot 9 \cdot \cdots \cdot (4n-7)]}{4^n \cdot n!} x^n$$

for $|x| < 1$, so $R = 1$.

35. $\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$, so $f(x) = \arctan(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{\left(x^2\right)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \frac{1}{2n+1} x^{4n+2}$, $R = 1$.

36. $\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$, so $f(x) = \sin\left(\frac{\pi}{4}x\right) = \sum_{n=0}^{\infty} (-1)^n \frac{\left(\frac{\pi}{4}x\right)^{2n+1}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{\pi^{2n+1}}{4^{2n+1}(2n+1)!} x^{2n+1}$, $R = \infty$.

37. $\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad \Rightarrow \quad \cos 2x = \sum_{n=0}^{\infty} (-1)^n \frac{(2x)^{2n}}{(2n)!} = \sum_{n=0}^{\infty} (-1)^n \frac{2^{2n}x^{2n}}{(2n)!}$, so

$f(x) = x\cos 2x = \sum_{n=0}^{\infty} (-1)^n \frac{2^{2n}}{(2n)!} x^{2n+1}$, $R = \infty$.

38. $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$, so $f(x) = e^{3x} - e^{2x} = \sum_{n=0}^{\infty} \frac{(3x)^n}{n!} - \sum_{n=0}^{\infty} \frac{(2x)^n}{n!} = \sum_{n=0}^{\infty} \frac{3^n x^n}{n!} - \sum_{n=0}^{\infty} \frac{2^n x^n}{n!} = \sum_{n=0}^{\infty} \frac{3^n - 2^n}{n!} x^n$, $R = \infty$.

39. $\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad \Rightarrow \quad \cos\left(\frac{1}{2}x^2\right) = \sum_{n=0}^{\infty} (-1)^n \frac{\left(\frac{1}{2}x^2\right)^{2n}}{(2n)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n}}{2^{2n}\,(2n)!}$, so

$f(x) = x\cos\left(\frac{1}{2}x^2\right) = \sum_{n=0}^{\infty} (-1)^n \frac{1}{2^{2n}(2n)!} x^{4n+1}$, $R = \infty$.

40. $\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} \quad \Rightarrow \quad \ln(1+x^3) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{3n}}{n}$, so $f(x) = x^2 \ln(1+x^3) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{3n+2}}{n}$,

$R = 1$.

41. We must write the binomial in the form (1+ expression), so we'll factor out a 4.

$$\frac{x}{\sqrt{4+x^2}} = \frac{x}{\sqrt{4(1+x^2/4)}} = \frac{x}{2\sqrt{1+x^2/4}} = \frac{x}{2}\left(1+\frac{x^2}{4}\right)^{-1/2} = \frac{x}{2}\sum_{n=0}^{\infty} \binom{-\frac{1}{2}}{n} \left(\frac{x^2}{4}\right)^n$$

$$= \frac{x}{2}\left[1 + \left(-\tfrac{1}{2}\right)\frac{x^2}{4} + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}\left(\frac{x^2}{4}\right)^2 + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!}\left(\frac{x^2}{4}\right)^3 + \cdots\right]$$

$$= \frac{x}{2} + \frac{x}{2}\sum_{n=1}^{\infty} (-1)^n \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{2^n \cdot 4^n \cdot n!} x^{2n}$$

$$= \frac{x}{2} + \sum_{n=1}^{\infty} (-1)^n \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{n!\, 2^{3n+1}} x^{2n+1} \text{ and } \frac{x^2}{4} < 1 \quad \Leftrightarrow \quad \frac{|x|}{2} < 1 \quad \Leftrightarrow \quad |x| < 2, \quad \text{so } R = 2.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $\dfrac{x^2}{\sqrt{2+x}} = \dfrac{x^2}{\sqrt{2\,(1+x/2)}} = \dfrac{x^2}{\sqrt{2}}\left(1+\dfrac{x}{2}\right)^{-1/2} = \dfrac{x^2}{\sqrt{2}}\sum_{n=0}^{\infty}\binom{-\frac{1}{2}}{n}\left(\dfrac{x}{2}\right)^n$

$$= \frac{x^2}{\sqrt{2}}\left[1+\left(-\tfrac{1}{2}\right)\left(\frac{x}{2}\right)+\frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}\left(\frac{x}{2}\right)^2+\frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!}\left(\frac{x}{2}\right)^3+\cdots\right]$$

$$= \frac{x^2}{\sqrt{2}}+\frac{x^2}{\sqrt{2}}\sum_{n=1}^{\infty}(-1)^n\frac{1\cdot3\cdot5\cdot\cdots\cdot(2n-1)}{n!\,2^{2n}}x^n$$

$$= \frac{x^2}{\sqrt{2}}+\sum_{n=1}^{\infty}(-1)^n\frac{1\cdot3\cdot5\cdot\cdots\cdot(2n-1)}{n!\,2^{2n+1/2}}x^{n+2} \text{ and } \left|\frac{x}{2}\right|<1 \iff |x|<2, \text{ so } R=2.$$

43. $\sin^2 x = \dfrac{1}{2}(1 - \cos 2x) = \dfrac{1}{2}\left[1-\sum_{n=0}^{\infty}\dfrac{(-1)^n(2x)^{2n}}{(2n)!}\right] = \dfrac{1}{2}\left[1-1-\sum_{n=1}^{\infty}\dfrac{(-1)^n(2x)^{2n}}{(2n)!}\right] = \sum_{n=1}^{\infty}\dfrac{(-1)^{n+1}2^{2n-1}x^{2n}}{(2n)!}$,

$R=\infty$

44. $\dfrac{x-\sin x}{x^3} = \dfrac{1}{x^3}\left[x-\sum_{n=0}^{\infty}\dfrac{(-1)^n x^{2n+1}}{(2n+1)!}\right] = \dfrac{1}{x^3}\left[x-x-\sum_{n=1}^{\infty}\dfrac{(-1)^n x^{2n+1}}{(2n+1)!}\right] = \dfrac{1}{x^3}\left[-\sum_{n=0}^{\infty}\dfrac{(-1)^{n+1}x^{2n+3}}{(2n+3)!}\right]$

$$= \frac{1}{x^3}\sum_{n=0}^{\infty}\frac{(-1)^n x^{2n+3}}{(2n+3)!} = \sum_{n=0}^{\infty}\frac{(-1)^n x^{2n}}{(2n+3)!}$$

and this series also gives the required value at $x=0$ (namely $1/6$); $R=\infty$.

45. $\cos x \overset{(16)}{=} \sum_{n=0}^{\infty}(-1)^n\dfrac{x^{2n}}{(2n)!} \quad \Rightarrow$

$$f(x)=\cos(x^2)=\sum_{n=0}^{\infty}\frac{(-1)^n(x^2)^{2n}}{(2n)!}=\sum_{n=0}^{\infty}\frac{(-1)^n x^{4n}}{(2n)!}$$
$$= 1-\tfrac{1}{2}x^4+\tfrac{1}{24}x^8-\tfrac{1}{720}x^{12}+\cdots$$

The series for $\cos x$ converges for all x, so the same is true of the series for $f(x)$, that is, $R=\infty$. Notice that, as n increases, $T_n(x)$ becomes a better approximation to $f(x)$.



46. $\ln(1+x) = \sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{x^n}{n} \quad \Rightarrow$

$$f(x)=\ln(1+x^2)=\sum_{n=1}^{\infty}\frac{(-1)^{n-1}(x^2)^n}{n}=\sum_{n=1}^{\infty}\frac{(-1)^{n-1}x^{2n}}{n}$$
$$= x^2-\tfrac{1}{2}x^4+\tfrac{1}{3}x^6-\tfrac{1}{4}x^8+\cdots$$

The series for $\ln(1+x)$ has $R=1$ and $\left|x^2\right|<1 \iff |x|<1$, so the series for $f(x)$ also has $R=1$. From the graphs of f and the first few Taylor polynomials, we see that $T_n(x)$ provides a closer fit to $f(x)$ near 0 as n increases.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $e^x \overset{(11)}{=} \sum_{n=0}^{\infty} \frac{x^n}{n!}$, so $e^{-x} = \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^n}{n!}$, so

$$f(x) = xe^{-x} = \sum_{n=0}^{\infty} (-1)^n \frac{1}{n!} x^{n+1}$$

$$= x - x^2 + \tfrac{1}{2}x^3 - \tfrac{1}{6}x^4 + \tfrac{1}{24}x^5 - \tfrac{1}{120}x^6 + \cdots$$

$$= \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{(n-1)!}$$



The series for e^x converges for all x, so the same is true of the series for $f(x)$; that is, $R = \infty$. From the graphs of f and the first few Taylor polynomials, we see that $T_n(x)$ provides a closer fit to $f(x)$ near 0 as n increases.

48. From Table 1, $\tan^{-1} x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$, so

$$f(x) = \tan^{-1}(x^3) = \sum_{n=0}^{\infty} (-1)^n \frac{(x^3)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{6n+3}}{2n+1}$$

$$= x^3 - \tfrac{1}{3}x^9 + \tfrac{1}{5}x^{15} - \tfrac{1}{7}x^{21} + \cdots$$

The series for $\tan^{-1} x$ has $R = 1$ and $|x^3| < 1 \quad \Leftrightarrow \quad |x| < 1$, so the series for $f(x)$ also has $R = 1$. From the graphs of f and the first few Taylor polynomials, we see that $T_n(x)$ provides a closer fit to $f(x)$ near 0 as n increases.



49. $5° = 5°\left(\frac{\pi}{180°}\right) = \frac{\pi}{36}$ radians and $\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots$, so

$\cos\frac{\pi}{36} = 1 - \frac{(\pi/36)^2}{2!} + \frac{(\pi/36)^4}{4!} - \frac{(\pi/36)^6}{6!} + \cdots$. Now $1 - \frac{(\pi/36)^2}{2!} \approx 0.99619$ and adding $\frac{(\pi/36)^4}{4!} \approx 2.4 \times 10^{-6}$ does not affect the fifth decimal place, so $\cos 5° \approx 0.99619$ by the Alternating Series Estimation Theorem.

50. $1/\sqrt[10]{e} = e^{-1/10}$ and $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$, so

$e^{-1/10} = 1 - \frac{1}{10} + \frac{(1/10)^2}{2!} - \frac{(1/10)^3}{3!} + \frac{(1/10)^4}{4!} - \frac{(1/10)^5}{5!} + \cdots$. Now

$1 - \frac{1}{10} + \frac{(1/10)^2}{2!} - \frac{(1/10)^3}{3!} + \frac{(1/10)^4}{4!} \approx 0.90484$ and subtracting $\frac{(1/10)^5}{5!} \approx 8.3 \times 10^{-8}$ does not affect the fifth decimal place, so $e^{-1/10} \approx 0.90484$ by the Alternating Series Estimation Theorem.

51. (a) $1/\sqrt{1-x^2} = \left[1 + (-x^2)\right]^{-1/2} = 1 + \left(-\frac{1}{2}\right)(-x^2) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}(-x^2)^2 + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!}(-x^2)^3 + \cdots$

$$= 1 + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{2^n \cdot n!} x^{2n}$$

(b) $\sin^{-1} x = \int \frac{1}{\sqrt{1-x^2}}\, dx = C + x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{(2n+1)2^n \cdot n!} x^{2n+1}$

$$= x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{(2n+1)2^n \cdot n!} x^{2n+1} \quad \text{since } 0 = \sin^{-1} 0 = C.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

52. (a) $1/\sqrt[4]{1+x} = (1+x)^{-1/4} = \sum_{n=0}^{\infty} \binom{\frac{1}{4}}{n} x^n = 1 - \frac{1}{4}x + \frac{\left(\frac{1}{4}\right)\left(-\frac{5}{4}\right)}{2!}x^2 + \frac{\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)\left(-\frac{9}{4}\right)}{3!}x^3 + \cdots$

$$= 1 - \frac{1}{4}x + \sum_{n=2}^{\infty} (-1)^n \frac{1 \cdot 5 \cdot 9 \cdot \cdots \cdot (4n-3)}{4^n \cdot n!} x^n$$

(b) $1/\sqrt[4]{1+x} = 1 - \frac{1}{4}x + \frac{5}{32}x^2 - \frac{15}{128}x^3 + \frac{195}{2048}x^4 - \cdots$. $1/\sqrt[4]{1.1} = 1/\sqrt[4]{1+0.1}$, so let $x = 0.1$. The sum of the first four terms is then $1 - \frac{1}{4}(0.1) + \frac{5}{32}(0.1)^2 - \frac{15}{128}(0.1)^3 \approx 0.976$. The fifth term is $\frac{195}{2048}(0.1)^4 \approx 0.000\,009\,5$, which does not affect the third decimal place of the sum, so we have $1/\sqrt[4]{1.1} \approx 0.976$. (Note that the third decimal place of the sum of the first three terms is affected by the fourth term, so we need to use more than three terms for the sum.)

53. $\sqrt{1+x^3} = (1+x^3)^{1/2} = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} (x^3)^n = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} x^{3n} \quad \Rightarrow \quad \int \sqrt{1+x^3}\,dx = C + \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} \frac{x^{3n+1}}{3n+1}$,

with $R = 1$.

54. $\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} \quad \Rightarrow \quad \sin(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n+1}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{(2n+1)!} \quad \Rightarrow$

$x^2 \sin(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+4}}{(2n+1)!} \quad \Rightarrow \quad \int x^2 \sin(x^2)\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+5}}{(2n+1)!(4n+5)}$, with $R = \infty$.

55. $\cos x \overset{(16)}{=} \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad \Rightarrow \quad \cos x - 1 = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad \Rightarrow \quad \frac{\cos x - 1}{x} = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n-1}}{(2n)!} \quad \Rightarrow$

$\int \frac{\cos x - 1}{x}\,dx = C + \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n}}{2n \cdot (2n)!}$, with $R = \infty$.

56. $\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \quad \Rightarrow \quad \arctan(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{2n+1} \quad \rightarrow$

$\int \arctan(x^2)\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+3}}{(2n+1)(4n+3)}$, with $R = 1$.

57. $\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$ for $|x| < 1$, so $x^3 \arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+4}}{2n+1}$ for $|x| < 1$ and

$\int x^3 \arctan x\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+5}}{(2n+1)(2n+5)}$. Since $\frac{1}{2} < 1$, we have

$\int_0^{1/2} x^3 \arctan x\,dx = \sum_{n=0}^{\infty} (-1)^n \frac{(1/2)^{2n+5}}{(2n+1)(2n+5)} = \frac{(1/2)^5}{1 \cdot 5} - \frac{(1/2)^7}{3 \cdot 7} + \frac{(1/2)^9}{5 \cdot 9} - \frac{(1/2)^{11}}{7 \cdot 11} + \cdots$. Now

$\frac{(1/2)^5}{1 \cdot 5} - \frac{(1/2)^7}{3 \cdot 7} + \frac{(1/2)^9}{5 \cdot 9} \approx 0.0059$ and subtracting $\frac{(1/2)^{11}}{7 \cdot 11} \approx 6.3 \times 10^{-6}$ does not affect the fourth decimal place, so $\int_0^{1/2} x^3 \arctan x\,dx \approx 0.0059$ by the Alternating Series Estimation Theorem.

58. $\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$ for all x, so $\sin(x^4) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{8n+4}}{(2n+1)!}$ for all x and

$\int \sin(x^4)\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{8n+5}}{(2n+1)!\,(8n+5)}$. Thus,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\int_0^1 \sin(x^4)\,dx = \sum_{n=0}^{\infty} (-1)^n \dfrac{1}{(2n+1)!\,(8n+5)} = \dfrac{1}{1!\cdot 5} - \dfrac{1}{3!\cdot 13} + \dfrac{1}{5!\cdot 21} - \dfrac{1}{7!\cdot 29} + \cdots$. Now

$\dfrac{1}{1!\cdot 5} - \dfrac{1}{3!\cdot 13} + \dfrac{1}{5!\cdot 21} \approx 0.1876$ and subtracting $\dfrac{1}{7!\cdot 29} \approx 6.84 \times 10^{-6}$ does not affect the fourth decimal place, so $\int_0^1 \sin(x^4)\,dx \approx 0.1876$ by the Alternating Series Estimation Theorem.

59. $\sqrt{1+x^4} = (1+x^4)^{1/2} = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} (x^4)^n$, so $\int \sqrt{1+x^4}\,dx = C + \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} \dfrac{x^{4n+1}}{4n+1}$ and hence, since $0.4 < 1$, we have

$$
\begin{aligned}
I &= \int_0^{0.4} \sqrt{1+x^4}\,dx = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} \frac{(0.4)^{4n+1}}{4n+1} \\
&= (1)\frac{(0.4)^1}{0!} + \frac{\frac{1}{2}}{1!}\frac{(0.4)^5}{5} + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)}{2!}\frac{(0.4)^9}{9} + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3!}\frac{(0.4)^{13}}{13} + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{4!}\frac{(0.4)^{17}}{17} + \cdots \\
&= 0.4 + \frac{(0.4)^5}{10} - \frac{(0.4)^9}{72} + \frac{(0.4)^{13}}{208} - \frac{5(0.4)^{17}}{2176} + \cdots
\end{aligned}
$$

Now $\dfrac{(0.4)^9}{72} \approx 3.6 \times 10^{-6} < 5 \times 10^{-6}$, so by the Alternating Series Estimation Theorem, $I \approx 0.4 + \dfrac{(0.4)^5}{10} \approx 0.40102$ (correct to five decimal places).

60. $\int_0^{0.5} x^2 e^{-x^2}\,dx = \int_0^{0.5} \sum_{n=0}^{\infty} \dfrac{(-1)^n x^{2n+2}}{n!}\,dx = \sum_{n=0}^{\infty} \left[\dfrac{(-1)^n x^{2n+3}}{n!(2n+3)}\right]_0^{1/2} = \sum_{n=0}^{\infty} \dfrac{(-1)^n}{n!(2n+3)2^{2n+3}}$ and since the term with $n = 2$ is $\dfrac{1}{1792} < 0.001$, we use $\sum_{n=0}^{1} \dfrac{(-1)^n}{n!(2n+3)2^{2n+3}} = \dfrac{1}{24} - \dfrac{1}{160} \approx 0.0354$.

61. $\lim_{x\to 0} \dfrac{x - \ln(1+x)}{x^2} = \lim_{x\to 0} \dfrac{x - (x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5 - \cdots)}{x^2} = \lim_{x\to 0} \dfrac{\frac{1}{2}x^2 - \frac{1}{3}x^3 + \frac{1}{4}x^4 - \frac{1}{5}x^5 + \cdots}{x^2}$

$= \lim_{x\to 0} (\frac{1}{2} - \frac{1}{3}x + \frac{1}{4}x^2 - \frac{1}{5}x^3 + \cdots) = \frac{1}{2}$

since power series are continuous functions.

62. $\lim_{x\to 0} \dfrac{1-\cos x}{1+x-e^x} = \lim_{x\to 0} \dfrac{1 - \left(1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \cdots\right)}{1 + x - \left(1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \frac{1}{5!}x^5 + \frac{1}{6!}x^6 + \cdots\right)}$

$= \lim_{x\to 0} \dfrac{\frac{1}{2!}x^2 - \frac{1}{4!}x^4 + \frac{1}{6!}x^6 - \cdots}{-\frac{1}{2!}x^2 - \frac{1}{3!}x^3 - \frac{1}{4!}x^4 - \frac{1}{5!}x^5 - \frac{1}{6!}x^6 - \cdots}$

$= \lim_{x\to 0} \dfrac{\frac{1}{2!} - \frac{1}{4!}x^2 + \frac{1}{6!}x^4 - \cdots}{-\frac{1}{2!} - \frac{1}{3!}x - \frac{1}{4!}x^2 - \frac{1}{5!}x^3 - \frac{1}{6!}x^4 - \cdots} = \dfrac{\frac{1}{2} - 0}{-\frac{1}{2} - 0} = -1$

since power series are continuous functions.

63. $\lim_{x\to 0} \dfrac{\sin x - x + \frac{1}{6}x^3}{x^5} = \lim_{x\to 0} \dfrac{\left(x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \cdots\right) - x + \frac{1}{6}x^3}{x^5}$

$= \lim_{x\to 0} \dfrac{\frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \cdots}{x^5} = \lim_{x\to 0} \left(\dfrac{1}{5!} - \dfrac{x^2}{7!} + \dfrac{x^4}{9!} - \cdots\right) = \dfrac{1}{5!} = \dfrac{1}{120}$

since power series are continuous functions.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

64. $\lim\limits_{x\to 0}\dfrac{\sqrt{1+x}\quad 1-\frac{1}{2}x}{x^2}=\lim\limits_{x\to 0}\dfrac{\left(1+\frac{1}{2}x-\frac{1}{8}x^2+\frac{1}{16}x^3-\cdots\right)-1-\frac{1}{2}x}{x^2}=\lim\limits_{x\to 0}\dfrac{-\frac{1}{8}x^2+\frac{1}{16}x^3-\cdots}{x^2}$

$=\lim\limits_{x\to 0}\left(-\frac{1}{8}+\frac{1}{16}x-\cdots\right)=-\frac{1}{8}$ since power series are continuous functions.

65. $\lim\limits_{x\to 0}\dfrac{x^3-3x+3\tan^{-1}x}{x^5}=\lim\limits_{x\to 0}\dfrac{x^3-3x+3\left(x-\frac{1}{3}x^3+\frac{1}{5}x^5-\frac{1}{7}x^7+\cdots\right)}{x^5}$

$=\lim\limits_{x\to 0}\dfrac{x^3-3x+3x-x^3+\frac{3}{5}x^5-\frac{3}{7}x^7+\cdots}{x^5}=\lim\limits_{x\to 0}\dfrac{\frac{3}{5}x^5-\frac{3}{7}x^7+\cdots}{x^5}$

$=\lim\limits_{x\to 0}\left(\frac{3}{5}-\frac{3}{7}x^2+\cdots\right)=\frac{3}{5}$ since power series are continuous functions.

66. $\lim\limits_{x\to 0}\dfrac{\tan x-x}{x^3}=\lim\limits_{x\to 0}\dfrac{\left(x+\frac{1}{3}x^3+\frac{2}{15}x^5+\cdots\right)-x}{x^3}=\lim\limits_{x\to 0}\dfrac{\frac{1}{3}x^3+\frac{2}{15}x^5+\cdots}{x^3}=\lim\limits_{x\to 0}\left(\frac{1}{3}+\frac{2}{15}x^2+\cdots\right)=\frac{1}{3}$

since power series are continuous functions.

67. From Equation 11, we have $e^{-x^2}=1-\dfrac{x^2}{1!}+\dfrac{x^4}{2!}-\dfrac{x^6}{3!}+\cdots$ and we know that $\cos x=1-\dfrac{x^2}{2!}+\dfrac{x^4}{4!}-\cdots$ from Equation 16. Therefore, $e^{-x^2}\cos x=\left(1-x^2+\frac{1}{2}x^4-\cdots\right)\left(1-\frac{1}{2}x^2+\frac{1}{24}x^4-\cdots\right)$. Writing only the terms with degree ≤ 4, we get $e^{-x^2}\cos x=1-\frac{1}{2}x^2+\frac{1}{24}x^4-x^2+\frac{1}{2}x^4+\frac{1}{2}x^4+\cdots=1-\frac{3}{2}x^2+\frac{25}{24}x^4+\cdots$.

68. $\sec x=\dfrac{1}{\cos x}\overset{(16)}{=}\dfrac{1}{1-\frac{1}{2}x^2+\frac{1}{24}x^4-\cdots}$.

$$
\begin{array}{r}
1+\frac{1}{2}x^2+\frac{5}{24}x^4+\cdots \\
1-\frac{1}{2}x^2+\frac{1}{24}x^4-\cdots \;\big)\overline{\;1\;} \\
\underline{1-\frac{1}{2}x^2+\frac{1}{24}x^4-\cdots} \\
\frac{1}{2}x^2-\frac{1}{24}x^4+\cdots \\
\underline{\frac{1}{2}x^2-\frac{1}{4}x^4+\cdots} \\
\frac{5}{24}x^4+\cdots \\
\underline{\frac{5}{24}x^4+\cdots} \\
\cdots
\end{array}
$$

From the long division above, $\sec x=1+\frac{1}{2}x^2+\frac{5}{24}x^4+\cdots$.

69. $\dfrac{x}{\sin x}\overset{(15)}{=}\dfrac{x}{x-\frac{1}{6}x^3+\frac{1}{120}x^5-\cdots}$.

$$
\begin{array}{r}
1+\frac{1}{6}x^2+\frac{7}{360}x^4+\cdots \\
x-\frac{1}{6}x^3+\frac{1}{120}x^5-\cdots \;\big)\overline{\;x\;} \\
\underline{x-\frac{1}{6}x^3+\frac{1}{120}x^5-\cdots} \\
\frac{1}{6}x^3-\frac{1}{120}x^5+\cdots \\
\underline{\frac{1}{6}x^3-\frac{1}{36}x^5+\cdots} \\
\frac{7}{360}x^5+\cdots \\
\underline{\frac{7}{360}x^5+\cdots} \\
\cdots
\end{array}
$$

From the long division above, $\dfrac{x}{\sin x}=1+\frac{1}{6}x^2+\frac{7}{360}x^4+\cdots$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

70. From Table 1, we have $e^x = 1 + \dfrac{x}{1!} + \dfrac{x^2}{2!} + \dfrac{x^3}{3!} + \cdots$ and that $\ln(1+x) = x - \dfrac{x^2}{2} + \dfrac{x^3}{3} - \dfrac{x^4}{4} + \cdots$. Therefore,

$y = e^x \ln(1+x) = \left(1 + \dfrac{x}{1!} + \dfrac{x^2}{2!} + \dfrac{x^3}{3!} + \cdots\right)\left(x - \dfrac{x^2}{2} + \dfrac{x^3}{3} - \dfrac{x^4}{4} + \cdots\right)$. Writing only terms with degree ≤ 3, we get $e^x \ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 + x^2 - \frac{1}{2}x^3 + \frac{1}{2}x^3 + \cdots = x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \cdots$.

71. $y = (\arctan x)^2 = \left(x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \cdots\right)\left(x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \cdots\right)$. Writing only the terms with degree ≤ 6, we get $(\arctan x)^2 = x^2 - \frac{1}{3}x^4 + \frac{1}{5}x^6 - \frac{1}{3}x^4 + \frac{1}{9}x^6 + \frac{1}{5}x^6 + \cdots = x^2 - \frac{2}{3}x^4 + \frac{23}{45}x^6 + \cdots$.

72. $y = e^x \sin^2 x = (e^x \sin x)\sin x = \left(x + x^2 + \frac{1}{3}x^3 + \cdots\right)\left(x - \frac{1}{6}x^3 + \cdots\right)$ [from Example 13]. Writing only the terms with degree ≤ 4, we get $e^x \sin^2 x = x^2 - \frac{1}{6}x^4 + x^3 + \frac{1}{3}x^4 + \cdots = x^2 + x^3 + \frac{1}{6}x^4 + \cdots$.

73. $\displaystyle\sum_{n=0}^{\infty} (-1)^n \frac{x^{4n}}{n!} = \sum_{n=0}^{\infty} \frac{\left(-x^4\right)^n}{n!} = e^{-x^4}$, by (11).

74. $\displaystyle\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n}}{6^{2n}(2n)!} = \sum_{n=0}^{\infty} (-1)^n \frac{\left(\frac{\pi}{6}\right)^{2n}}{(2n)!} = \cos\frac{\pi}{6} = \frac{\sqrt{3}}{2}$, by (16).

75. $\displaystyle\sum_{n=1}^{\infty} (-1)^{n-1} \frac{3^n}{n5^n} = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(3/5)^n}{n} = \ln\left(1 + \frac{3}{5}\right)$ [from Table 1] $= \ln\dfrac{8}{5}$

76. $\displaystyle\sum_{n=0}^{\infty} \frac{3^n}{5^n\, n!} = \sum_{n=0}^{\infty} \frac{(3/5)^n}{n!} = e^{3/5}$, by (11).

77. $\displaystyle\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n+1}}{4^{2n+1}(2n+1)!} = \sum_{n=0}^{\infty} \frac{(-1)^n \left(\frac{\pi}{4}\right)^{2n+1}}{(2n+1)!} = \sin\frac{\pi}{4} = \frac{1}{\sqrt{2}}$, by (15).

78. $1 - \ln 2 + \dfrac{(\ln 2)^2}{2!} - \dfrac{(\ln 2)^3}{3!} + \cdots = \displaystyle\sum_{n=0}^{\infty} \frac{(-\ln 2)^n}{n!} = e^{-\ln 2} = \left(e^{\ln 2}\right)^{-1} = 2^{-1} = \frac{1}{2}$, by (11).

79. $3 + \dfrac{9}{2!} + \dfrac{27}{3!} + \dfrac{81}{4!} + \cdots = \dfrac{3^1}{1!} + \dfrac{3^2}{2!} + \dfrac{3^3}{3!} + \dfrac{3^4}{4!} + \cdots = \displaystyle\sum_{n=1}^{\infty} \frac{3^n}{n!} = \sum_{n=0}^{\infty} \frac{3^n}{n!} \quad 1 = e^3 - 1$, by (11).

80. $\dfrac{1}{1\cdot 2} - \dfrac{1}{3\cdot 2^3} + \dfrac{1}{5\cdot 2^5} - \dfrac{1}{7\cdot 2^7} + \cdots = \displaystyle\sum_{n-0}^{\infty} (-1)^n \frac{1}{(2n+1)2^{2n+1}} = \sum_{n=0}^{\infty} (-1)^n \frac{(1/2)^{2n+1}}{2n+1} = \tan^{-1}\left(\frac{1}{2}\right)$ [from Table 1]

81. If p is an nth-degree polynomial, then $p^{(i)}(x) = 0$ for $i > n$, so its Taylor series at a is $p(x) = \displaystyle\sum_{i=0}^{n} \frac{p^{(i)}(a)}{i!}(x-a)^i$.

Put $x - a = 1$, so that $x = a + 1$. Then $p(a+1) = \displaystyle\sum_{i=0}^{n} \frac{p^{(i)}(a)}{i!}$.

This is true for any a, so replace a by x: $p(x+1) = \displaystyle\sum_{i=0}^{n} \frac{p^{(i)}(x)}{i!}$

82. The coefficient of x^{58} in the Maclaurin series of $f(x) = (1+x^3)^{30}$ is $\dfrac{f^{(58)}(0)}{58!}$. But the binomial series for $f(x)$ is

$(1+x^3)^{30} = \displaystyle\sum_{n=0}^{\infty} \binom{30}{n} x^{3n}$, so it involves only powers of x that are multiples of 3 and therefore the coefficient of x^{58} is 0.

So $f^{(58)}(0) = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

83. Assume that $|f'''(x)| \le M$, so $f'''(x) \le M$ for $a \le x \le a+d$. Now $\int_a^x f'''(t)\,dt \le \int_a^x M\,dt \;\Rightarrow$

$f''(x) - f''(a) \le M(x-a) \;\Rightarrow\; f''(x) \le f''(a) + M(x-a)$. Thus, $\int_a^x f''(t)\,dt \le \int_a^x [f''(a) + M(t-a)]\,dt \;\Rightarrow$

$f'(x) - f'(a) \le f''(a)(x-a) + \frac{1}{2}M(x-a)^2 \;\Rightarrow\; f'(x) \le f'(a) + f''(a)(x-a) + \frac{1}{2}M(x-a)^2 \;\Rightarrow$

$\int_a^x f'(t)\,dt \le \int_a^x \left[f'(a) + f''(a)(t-a) + \frac{1}{2}M(t-a)^2\right] dt \;\Rightarrow$

$f(x) - f(a) \le f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2 + \frac{1}{6}M(x-a)^3$. So

$f(x) - f(a) - f'(a)(x-a) - \frac{1}{2}f''(a)(x-a)^2 \le \frac{1}{6}M(x-a)^3$. But

$R_2(x) = f(x) - T_2(x) = f(x) - f(a) - f'(a)(x-a) - \frac{1}{2}f''(a)(x-a)^2$, so $R_2(x) \le \frac{1}{6}M(x-a)^3$.

A similar argument using $f'''(x) \ge -M$ shows that $R_2(x) \ge -\frac{1}{6}M(x-a)^3$. So $|R_2(x_2)| \le \frac{1}{6}M\,|x-a|^3$.

Although we have assumed that $x > a$, a similar calculation shows that this inequality is also true if $x < a$.

84. (a) $f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \ne 0 \\ 0 & \text{if } x = 0 \end{cases}$ so $f'(0) = \lim\limits_{x\to 0} \dfrac{f(x)-f(0)}{x-0} = \lim\limits_{x\to 0} \dfrac{e^{-1/x^2}}{x} = \lim\limits_{x\to 0} \dfrac{1/x}{e^{1/x^2}} = \lim\limits_{x\to 0} \dfrac{x}{2e^{1/x^2}} = 0$

(using l'Hospital's Rule and simplifying in the penultimate step). Similarly, we can use the definition of the derivative and l'Hospital's Rule to show that $f''(0) = 0$, $f^{(3)}(0) = 0, \ldots, f^{(n)}(0) = 0$, so that the Maclaurin series for f consists entirely of zero terms. But since $f(x) \ne 0$ except for $x = 0$, we see that f cannot equal its Maclaurin series except at $x = 0$.

(b)



From the graph, it seems that the function is extremely flat at the origin. In fact, it could be said to be "infinitely flat" at $x = 0$, since all of its derivatives are 0 there.

85. (a) $g(x) = \sum\limits_{n=0}^{\infty} \binom{k}{n} x^n \;\Rightarrow\; g'(x) = \sum\limits_{n=1}^{\infty} \binom{k}{n} n x^{n-1}$, so

$$(1+x)g'(x) = (1+x)\sum_{n=1}^{\infty}\binom{k}{n}nx^{n-1} = \sum_{n=1}^{\infty}\binom{k}{n}nx^{n-1} + \sum_{n=1}^{\infty}\binom{k}{n}nx^n$$

$$= \sum_{n=0}^{\infty}\binom{k}{n+1}(n+1)x^n + \sum_{n=0}^{\infty}\binom{k}{n}nx^n \qquad \left[\begin{array}{c}\text{Replace } n \text{ with } n+1\\ \text{in the first series}\end{array}\right]$$

$$\sum_{n=0}^{\infty}(n+1)\frac{k(k-1)(k-2)\cdots(k-n+1)(k-n)}{(n+1)!}x^n + \sum_{n=0}^{\infty}\left[(n)\frac{k(k-1)(k-2)\cdots(k-n+1)}{n!}\right]x^n$$

$$= \sum_{n=0}^{\infty}\frac{(n+1)k(k-1)(k-2)\cdots(k-n+1)}{(n+1)!}\left[(k-n)+n\right]x^n$$

$$= k\sum_{n=0}^{\infty}\frac{k(k-1)(k-2)\cdots(k-n+1)}{n!}x^n = k\sum_{n=0}^{\infty}\binom{k}{n}x^n = kg(x)$$

Thus, $g'(x) = \dfrac{kg(x)}{1+x}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $h(x) = (1+x)^{-k} g(x) \Rightarrow$

$$\begin{aligned} h'(x) &= -k(1+x)^{-k-1} g(x) + (1+x)^{-k} g'(x) && \text{[Product Rule]} \\ &= -k(1+x)^{-k-1} g(x) + (1+x)^{-k} \frac{k g(x)}{1+x} && \text{[from part (a)]} \\ &= -k(1+x)^{-k-1} g(x) + k(1+x)^{-k-1} g(x) = 0 \end{aligned}$$

(c) From part (b) we see that $h(x)$ must be constant for $x \in (-1, 1)$, so $h(x) = h(0) = 1$ for $x \in (-1, 1)$.

Thus, $h(x) = 1 = (1+x)^{-k} g(x) \Leftrightarrow g(x) = (1+x)^k$ for $x \in (-1, 1)$.

86. Using the binomial series to expand $\sqrt{1+x}$ as a power series as in Example 9, we get

$\sqrt{1+x} = (1+x)^{1/2} = 1 + \dfrac{x}{2} + \displaystyle\sum_{n=2}^{\infty} \frac{(-1)^{n-1} 1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-3) x^n}{2^n \cdot n!}$, so

$\left(1 - x^2\right)^{1/2} = 1 - \dfrac{1}{2}x^2 - \displaystyle\sum_{n=2}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-3)}{2^n \cdot n!} x^{2n}$ and

$\sqrt{1 - e^2 \sin^2\theta} = 1 - \dfrac{1}{2}e^2 \sin^2\theta - \displaystyle\sum_{n=2}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-3)}{2^n \cdot n!} e^{2n} \sin^{2n}\theta$. Thus,

$$L = 4a \int_0^{\pi/2} \sqrt{1 - e^2 \sin^2\theta}\, d\theta = 4a \int_0^{\pi/2} \left(1 - \frac{1}{2}e^2 \sin^2\theta - \sum_{n=2}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-3)}{2^n \cdot n!} e^{2n} \sin^{2n}\theta\right) d\theta$$

$$= 4a \left[\frac{\pi}{2} - \frac{e^2}{2} S_1 - \sum_{n=2}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-3)}{n!} \left(\frac{e^2}{2}\right)^n S_n\right]$$

where $S_n = \displaystyle\int_0^{\pi/2} \sin^{2n}\theta\, d\theta = \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{2 \cdot 4 \cdot 6 \cdot \cdots \cdot 2n} \frac{\pi}{2}$ by Exercise 7.1.50.

$$L = 4a\left(\frac{\pi}{2}\right)\left[1 - \frac{e^2}{2} \cdot \frac{1}{2} - \sum_{n=2}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-3)}{n!} \left(\frac{e^2}{2}\right)^n \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{2 \cdot 4 \cdot 6 \cdot \cdots \cdot 2n}\right]$$

$$= 2\pi a\left[1 - \frac{e^2}{4} - \sum_{n=2}^{\infty} \frac{e^{2n}}{2^n} \cdot \frac{1^2 \cdot 3^2 \cdot 5^2 \cdot \cdots \cdot (2n-3)^2 (2n-1)}{n! \cdot 2^n \cdot n!}\right]$$

$$= 2\pi a\left[1 - \frac{e^2}{4} - \sum_{n=2}^{\infty} \frac{e^{2n}}{4^n} \left(\frac{1 \cdot 3 \cdot \cdots \cdot (2n-3)}{n!}\right)^2 (2n-1)\right]$$

$$= 2\pi a\left[1 - \frac{e^2}{4} - \frac{3e^4}{64} - \frac{5e^6}{256} - \cdots\right] = \frac{\pi a}{128}(256 - 64e^2 - 12e^4 - 5e^6 - \cdots)$$

LABORATORY PROJECT An Elusive Limit

1. $f(x) = \dfrac{n(x)}{d(x)} = \dfrac{\sin(\tan x) - \tan(\sin x)}{\arcsin(\arctan x) - \arctan(\arcsin x)}$

The table of function values were obtained using Maple with 10 digits of precision. The results of this project will vary depending on the CAS and precision level. It appears that as $x \to 0^+$, $f(x) \to \frac{10}{3}$. Since f is an even function, we have $f(x) \to \frac{10}{3}$ as $x \to 0$.

x	$f(x)$
1	1.1838
0.1	0.9821
0.01	2.0000
0.001	3.3333
0.0001	3.3333

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

2. The graph is inconclusive about the limit of f as $x \to 0$.



3. The limit has the indeterminate form $\frac{0}{0}$. Applying l'Hospital's Rule, we obtain the form $\frac{0}{0}$ six times. Finally, on the seventh application we obtain $\lim_{x\to 0} \dfrac{n^{(7)}(x)}{d^{(7)}(x)} = \dfrac{-168}{-168} = 1$.

4. $\lim_{x\to 0} f(x) = \lim_{x\to 0} \dfrac{n(x)}{d(x)} \overset{\text{CAS}}{=} \lim_{x\to 0} \dfrac{-\frac{1}{30}x^7 - \frac{29}{756}x^9 + \cdots}{-\frac{1}{30}x^7 + \frac{13}{756}x^9 + \cdots}$

$$= \lim_{x\to 0} \frac{\left(-\frac{1}{30}x^7 - \frac{29}{756}x^9 + \cdots\right)/x^7}{\left(-\frac{1}{30}x^7 + \frac{13}{756}x^9 + \cdots\right)/x^7} = \lim_{x\to 0} \frac{-\frac{1}{30} - \frac{29}{756}x^2 + \cdots}{-\frac{1}{30} + \frac{13}{756}x^2 + \cdots} = \frac{-\frac{1}{30}}{-\frac{1}{30}} = 1$$

Note that $n^{(7)}(x) = d^{(7)}(x) = -\frac{7!}{30} = -\frac{5040}{30} = -168$, which agrees with the result in Problem 3.

5. The limit command gives the result that $\lim_{x\to 0} f(x) = 1$.

6. The strange results (with only 10 digits of precision) must be due to the fact that the terms being subtracted in the numerator and denominator are very close in value when $|x|$ is small. Thus, the differences are imprecise (have few correct digits).

11.11 Applications of Taylor Polynomials

1. (a)

n	$f^{(n)}(x)$	$f^{(n)}(0)$	$T_n(x)$
0	$\sin x$	0	0
1	$\cos x$	1	x
2	$-\sin x$	0	x
3	$\cos x$	−1	$x - \frac{1}{6}x^3$
4	$\sin x$	0	$x - \frac{1}{6}x^3$
5	$\cos x$	1	$x - \frac{1}{6}x^3 + \frac{1}{120}x^5$



Note: $T_n(x) = \sum_{k=0}^{n} \dfrac{f^{(k)}(0)}{k!}x^k$

(b)

x	f	$T_0(x)$	$T_1(x)$ $T_2(x)$	$T_3(x)$ $T_4(x)$	$T_5(x)$
$\frac{\pi}{4}$	0.7071	0	0.7854	0.7047	0.7071
$\frac{\pi}{2}$	1	0	1.5708	0.9248	1.0045
π	0	0	3.1416	−2.0261	0.5240

(c) As n increases, $T_n(x)$ is a good approximation to $f(x)$ on a larger and larger interval.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. (a)

n	$f^{(n)}(x)$	$f^{(n)}(0)$	$T_n(x)$
0	$\tan x$	0	0
1	$\sec^2 x$	1	x
2	$2\sec^2 x \tan x$	0	x
3	$4\sec^2 x \tan^2 x + 2\sec^4 x$	2	$x + \frac{1}{3}x^3$



(b)

x	f	$T_0(x)$	$T_1(x) = T_2(x)$	$T_3(x)$
$\frac{\pi}{6}$	0.5774	0	0.5236	0.5714
$\frac{\pi}{4}$	1	0	0.7854	0.9469
$\frac{\pi}{3}$	1.7321	0	1.0472	1.4300

Note: $T_n(x) = \sum_{k=0}^{n} \dfrac{f^{(k)}(0)}{k!}\, x^k$

(c) As n increases, $T_n(x)$ is a good approximation to $f(x)$ on a larger and larger interval. Because the Taylor polynomials are continuous, they cannot approximate the infinite discontinuities at $x = \pm\pi/2$. They can only approximate $\tan x$ on $(-\pi/2, \pi/2)$.

3.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	e^x	e
1	e^x	e
2	e^x	e
3	e^x	e



$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(1)}{n!}\,(x-1)^n$$

$$= \frac{e}{0!}(x-1)^0 + \frac{e}{1!}(x-1)^1 + \frac{e}{2!}(x-1)^2 + \frac{e}{3!}(x-1)^3$$

$$= e + e(x-1) + \tfrac{1}{2}e(x-1)^2 + \tfrac{1}{6}e(x-1)^3$$

4.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/6)$
0	$\sin x$	$1/2$
1	$\cos x$	$\sqrt{3}/2$
2	$-\sin x$	$-1/2$
3	$-\cos x$	$-\sqrt{3}/2$



$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(\pi/6)}{n!}\left(x - \frac{\pi}{6}\right)^n$$

$$= \frac{1/2}{0!}\left(x-\frac{\pi}{6}\right)^0 + \frac{\sqrt{3}/2}{1!}\left(x-\frac{\pi}{6}\right)^1 - \frac{1/2}{2!}\left(x-\frac{\pi}{6}\right)^2 + \frac{\sqrt{3}/2}{3!}\left(x-\frac{\pi}{6}\right)^3$$

$$= \frac{1}{2} + \frac{\sqrt{3}}{2}\left(x-\frac{\pi}{6}\right) - \frac{1}{4}\left(x-\frac{\pi}{6}\right)^2 - \frac{\sqrt{3}}{12}\left(x-\frac{\pi}{6}\right)^3$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/2)$
0	$\cos x$	0
1	$-\sin x$	-1
2	$-\cos x$	0
3	$\sin x$	1

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(\pi/2)}{n!}\left(x - \tfrac{\pi}{2}\right)^n$$

$$= -\left(x - \tfrac{\pi}{2}\right) + \tfrac{1}{6}\left(x - \tfrac{\pi}{2}\right)^3$$



6.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$e^{-x}\sin x$	0
1	$e^{-x}(\cos x - \sin x)$	1
2	$-2e^{-x}\cos x$	-2
3	$2e^{-x}(\cos x + \sin x)$	2

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(0)}{n!}\,x^n = x - x^2 + \tfrac{1}{3}x^3$$



7.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$\ln x$	0
1	$1/x$	1
2	$-1/x^2$	-1
3	$2/x^3$	2

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(1)}{n!}(x-1)^n$$

$$= 0 + \frac{1}{1!}(x-1) + \frac{-1}{2!}(x-1)^2 + \frac{2}{3!}(x-1)^3$$

$$= (x-1) - \tfrac{1}{2}(x-1)^2 + \tfrac{1}{3}(x-1)^3$$



8.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$x\cos x$	0
1	$-x\sin x + \cos x$	1
2	$-x\cos x - 2\sin x$	0
3	$x\sin x - 3\cos x$	-3

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(0)}{n!}\,x^n$$

$$= 0 + \frac{1}{1!}x + 0 + \frac{-3}{3!}x^3 = x - \frac{1}{2}x^3$$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	xe^{-2x}	0
1	$(1-2x)e^{-2x}$	1
2	$4(x-1)e^{-2x}$	-4
3	$4(3-2x)e^{-2x}$	12



$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(0)}{n!}x^n = \tfrac{0}{1}\cdot 1 + \tfrac{1}{1}x^1 + \tfrac{-4}{2}x^2 + \tfrac{12}{6}x^3 = x - 2x^2 + 2x^3$$

10.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$\tan^{-1} x$	$\frac{\pi}{4}$
1	$\dfrac{1}{1+x^2}$	$\frac{1}{2}$
2	$\dfrac{-2x}{(1+x^2)^2}$	$-\frac{1}{2}$
3	$\dfrac{6x^2-2}{(1+x^2)^3}$	$\frac{1}{2}$



$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(1)}{n!}(x-1)^n = \frac{\pi}{4} + \frac{1/2}{1}(x-1)^1 + \frac{-1/2}{2}(x-1)^2 + \frac{1/2}{6}(x-1)^3$$

$$= \tfrac{\pi}{4} + \tfrac{1}{2}(x-1) - \tfrac{1}{4}(x-1)^2 + \tfrac{1}{12}(x-1)^3$$

11. You may be able to simply find the Taylor polynomials for $f(x) = \cot x$ using your CAS. We will list the values of $f^{(n)}(\pi/4)$ for $n = 0$ to $n = 5$.



n	0	1	2	3	4	5
$f^{(n)}(\pi/4)$	1	-2	4	-16	80	-512

$$T_5(x) = \sum_{n=0}^{5} \frac{f^{(n)}(\pi/4)}{n!}\left(x - \tfrac{\pi}{4}\right)^n$$

$$= 1 - 2\left(x - \tfrac{\pi}{4}\right) + 2\left(x - \tfrac{\pi}{4}\right)^2 - \tfrac{8}{3}\left(x - \tfrac{\pi}{4}\right)^3 + \tfrac{10}{3}\left(x - \tfrac{\pi}{4}\right)^4 - \tfrac{64}{15}\left(x - \tfrac{\pi}{4}\right)^5$$

For $n = 2$ to $n = 5$, $T_n(x)$ is the polynomial consisting of all the terms up to and including the $\left(x - \frac{\pi}{4}\right)^n$ term.

12. You may be able to simply find the Taylor polynomials for $f(x) = \sqrt[3]{1+x^2}$ using your CAS. We will list the values of $f^{(n)}(0)$ for $n = 0$ to $n = 5$.



n	0	1	2	3	4	5
$f^{(n)}(0)$	1	0	$\frac{2}{3}$	0	$-\frac{8}{3}$	0

$$T_5(x) = \sum_{n=0}^{5} \frac{f^{(n)}(0)}{n!}x^n = 1 + \tfrac{1}{3}x^2 - \tfrac{1}{9}x^4$$

For $n = 2$ to $n = 5$, $T_n(x)$ is the polynomial consisting of all the terms up to and including the x^n term. Note that $T_2 = T_3$ and $T_4 = T_5$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

13. (a)

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$1/x$	1
1	$-1/x^2$	-1
2	$2/x^3$	2
3	$-6/x^4$	

$$\begin{aligned} f(x) = 1/x &\approx T_2(x) \\ &= \frac{1}{0!}(x-1)^0 - \frac{1}{1!}(x-1)^1 + \frac{2}{2!}(x-1)^2 \\ &= 1 - (x-1) + (x-1)^2 \end{aligned}$$

(b) $|R_2(x)| \le \dfrac{M}{3!}|x-1|^3$, where $|f'''(x)| \le M$. Now $0.7 \le x \le 1.3 \;\Rightarrow\; |x-1| \le 0.3 \;\Rightarrow\; |x-1|^3 \le 0.027$.

Since $|f'''(x)|$ is decreasing on $[0.7, 1.3]$, we can take $M = |f'''(0.7)| = 6/(0.7)^4$, so

$$|R_2(x)| \le \frac{6/(0.7)^4}{6}(0.027) = 0.112\,453\,1.$$

(c)



From the graph of $|R_2(x)| = \left|\dfrac{1}{x} - T_2(x)\right|$, it seems that the error is less than 0.038 571 on $[0.7, 1.3]$.

14. (a)

n	$f^{(n)}(x)$	$f^{(n)}(4)$
0	$x^{-1/2}$	$\frac{1}{2}$
1	$-\frac{1}{2}x^{-3/2}$	$-\frac{1}{16}$
2	$\frac{3}{4}x^{-5/2}$	$\frac{3}{128}$
3	$-\frac{15}{8}x^{-7/2}$	

$$\begin{aligned} f(x) = x^{-1/2} &\approx T_2(x) \\ &= \frac{1/2}{0!}(x-4)^0 - \frac{1/16}{1!}(x-4)^1 + \frac{3/128}{2!}(x-4)^2 \\ &= \tfrac{1}{2} - \tfrac{1}{16}(x-4) + \tfrac{3}{256}(x-4)^2 \end{aligned}$$

(b) $|R_2(x)| \le \dfrac{M}{3!}|x-4|^3$, where $|f'''(x)| \le M$. Now $3.5 < x < 4.5 \;\Rightarrow\; |x-4| < 0.5 \;\Rightarrow\; |x-4|^3 < 0.125$.

Since $|f'''(x)|$ is decreasing on $[3.5, 4.5]$, we can take $M = |f'''(3.5)| = \dfrac{15}{8(3.5)^{7/2}}$, so

$$|R_2(x)| \le \frac{15}{6 \cdot 8(3.5)^{7/2}}(0.125) \approx 0.000\,487.$$

(c)



From the graph of $|R_2(x)| = \left|x^{-1/2} - T_2(x)\right|$, it seems that the error is less than 0.000 343 on $[3.5, 4.5]$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$x^{2/3}$	1
1	$\frac{2}{3}x^{-1/3}$	$\frac{2}{3}$
2	$-\frac{2}{9}x^{-4/3}$	$-\frac{2}{9}$
3	$\frac{8}{27}x^{-7/3}$	$\frac{8}{27}$
4	$-\frac{56}{81}x^{-10/3}$	

(a) $f(x) = x^{2/3} \approx T_3(x) = 1 + \frac{2}{3}(x-1) - \frac{2/9}{2!}(x-1)^2 + \frac{8/27}{3!}(x-1)^3$

$= 1 + \frac{2}{3}(x-1) - \frac{1}{9}(x-1)^2 + \frac{4}{81}(x-1)^3$

(b) $|R_3(x)| \le \frac{M}{4!}|x-1|^4$, where $\left|f^{(4)}(x)\right| \le M$. Now $0.8 \le x \le 1.2 \Rightarrow$ $|x-1| \le 0.2 \Rightarrow |x-1|^4 \le 0.0016$. Since $\left|f^{(4)}(x)\right|$ is decreasing on $[0.8, 1.2]$, we can take $M = \left|f^{(4)}(0.8)\right| = \frac{56}{81}(0.8)^{-10/3}$, so $|R_3(x)| \le \dfrac{\frac{56}{81}(0.8)^{-10/3}}{24}(0.0016) \approx 0.000\,096\,97$.

(c)



From the graph of $|R_3(x)| = \left|x^{2/3} - T_3(x)\right|$, it seems that the error is less than $0.000\,053\,3$ on $[0.8, 1.2]$.

16.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/6)$
0	$\sin x$	$1/2$
1	$\cos x$	$\sqrt{3}/2$
2	$-\sin x$	$-1/2$
3	$-\cos x$	$-\sqrt{3}/2$
4	$\sin x$	$1/2$
5	$\cos x$	

(a) $f(x) = \sin x \approx T_4(x)$

$= \frac{1}{2} + \frac{\sqrt{3}}{2}\left(x - \frac{\pi}{6}\right) - \frac{1}{4}\left(x - \frac{\pi}{6}\right)^2 - \frac{\sqrt{3}}{12}\left(x - \frac{\pi}{6}\right)^3 + \frac{1}{48}\left(x - \frac{\pi}{6}\right)^4$

(b) $|R_4(x)| \le \frac{M}{5!}\left|x - \frac{\pi}{6}\right|^5$, where $\left|f^{(5)}(x)\right| \le M$. Now $0 \le x \le \frac{\pi}{3} \Rightarrow -\frac{\pi}{6} \le x - \frac{\pi}{6} \le \frac{\pi}{6} \Rightarrow \left|x - \frac{\pi}{6}\right| \le \frac{\pi}{6} \Rightarrow$ $\left|x - \frac{\pi}{6}\right|^5 \le \left(\frac{\pi}{6}\right)^5$. Since $\left|f^{(5)}(x)\right|$ is decreasing on $\left[0, \frac{\pi}{3}\right]$, we can take $M = \left|f^{(5)}(0)\right| = \cos 0 = 1$, so $|R_4(x)| \le \dfrac{1}{5!}\left(\dfrac{\pi}{6}\right)^5 \approx 0.000\,328$.

(c)



From the graph of $|R_4(x)| = |\sin x - T_4(x)|$, it seems that the error is less than $0.000\,297$ on $\left[0, \frac{\pi}{3}\right]$.

17.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sec x$	1
1	$\sec x \tan x$	0
2	$\sec x\,(2\sec^2 x - 1)$	1
3	$\sec x \tan x\,(6\sec^2 x - 1)$	

(a) $f(x) = \sec x \approx T_2(x) = 1 + \frac{1}{2}x^2$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) $|R_2(x)| \le \dfrac{M}{3!}|x|^3$, where $\left|f^{(3)}(x)\right| \le M$. Now $-0.2 \le x \le 0.2 \;\Rightarrow\; |x| \le 0.2 \;\Rightarrow\; |x|^3 \le (0.2)^3$.

$f^{(3)}(x)$ is an odd function and it is increasing on $[0, 0.2]$ since $\sec x$ and $\tan x$ are increasing on $[0, 0.2]$,

so $\left|f^{(3)}(x)\right| \le f^{(3)}(0.2) \approx 1.085\,158\,892$. Thus, $|R_2(x)| \le \dfrac{f^{(3)}(0.2)}{3!}(0.2)^3 \approx 0.001\,447$.

(c)



From the graph of $|R_2(x)| = |\sec x - T_2(x)|$, it seems that the error is less than $0.000\,339$ on $[-0.2, 0.2]$.

18.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$\ln(1+2x)$	$\ln 3$
1	$2/(1+2x)$	$\frac{2}{3}$
2	$-4/(1+2x)^2$	$-\frac{4}{9}$
3	$16/(1+2x)^3$	$\frac{16}{27}$
4	$-96/(1+2x)^4$	

(a) $f(x) = \ln(1+2x) \approx T_3(x)$

$= \ln 3 + \frac{2}{3}(x-1) - \dfrac{4/9}{2!}(x-1)^2 + \dfrac{16/27}{3!}(x-1)^3$

(b) $|R_3(x)| \le \dfrac{M}{4!}|x-1|^4$, where $\left|f^{(4)}(x)\right| \le M$. Now $0.5 \le x \le 1.5 \;\Rightarrow$

$-0.5 \le x - 1 \le 0.5 \;\Rightarrow\; |x-1| \le 0.5 \;\Rightarrow\; |x-1|^4 \le \frac{1}{16}$, and

letting $x = 0.5$ gives $M = 6$, so $|R_3(x)| \le \dfrac{6}{4!} \cdot \dfrac{1}{16} = \dfrac{1}{64} = 0.015\,625$.

(c)



From the graph of $|R_3(x)| = |\ln(1+2x) - T_3(x)|$, it seems that the error is less than 0.005 on $[0.5, 1.5]$.

19.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	e^{x^2}	1
1	$e^{x^2}(2x)$	0
2	$e^{x^2}(2+4x^2)$	2
3	$e^{x^2}(12x+8x^3)$	0
4	$e^{x^2}(12+48x^2+16x^4)$	

(a) $f(x) = e^{x^2} \approx T_3(x) = 1 + \dfrac{2}{2!}x^2 = 1 + x^2$

(b) $|R_3(x)| \le \dfrac{M}{4!}|x|^4$, where $\left|f^{(4)}(x)\right| \le M$. Now $0 \le x \le 0.1 \;\Rightarrow$

$x^4 \le (0.1)^4$, and letting $x = 0.1$ gives

$|R_3(x)| \le \dfrac{e^{0.01}(12 + 0.48 + 0.0016)}{24}(0.1)^4 \approx 0.00006$.

(c)



From the graph of $|R_3(x)| = \left|e^{x^2} - T_3(x)\right|$, it appears that the error is less than $0.000\,051$ on $[0, 0.1]$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$x \ln x$	0
1	$\ln x + 1$	1
2	$1/x$	1
3	$-1/x^2$	-1
4	$2/x^3$	

(a) $f(x) = x \ln x \approx T_3(x) = (x-1) + \frac{1}{2}(x-1)^2 - \frac{1}{6}(x-1)^3$

(b) $|R_3(x)| \le \dfrac{M}{4!}|x-1|^4$, where $\left|f^{(4)}(x)\right| \le M$. Now $0.5 \le x \le 1.5 \Rightarrow |x-1| \le \frac{1}{2} \Rightarrow |x-1|^4 \le \frac{1}{16}$. Since $\left|f^{(4)}(x)\right|$ is decreasing on $[0.5, 1.5]$, we can take $M = \left|f^{(4)}(0.5)\right| = 2/(0.5)^3 = 16$, so $|R_3(x)| \le \frac{16}{24}(1/16) = \frac{1}{24} = 0.041\overline{6}$.

(c)



From the graph of $|R_3(x)| = |x \ln x - T_3(x)|$, it seems that the error is less than 0.0076 on $[0.5, 1.5]$.

21.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$x \sin x$	0
1	$\sin x + x \cos x$	0
2	$2\cos x - x \sin x$	2
3	$-3\sin x - x \cos x$	0
4	$-4\cos x + x \sin x$	-4
5	$5\sin x + x \cos x$	

(a) $f(x) = x \sin x \approx T_4(x) = \dfrac{2}{2!}(x-0)^2 + \dfrac{-4}{4!}(x-0)^4 = x^2 - \dfrac{1}{6}x^4$

(b) $|R_4(x)| \le \dfrac{M}{5!}|x|^5$, where $\left|f^{(5)}(x)\right| \le M$. Now $-1 \le x \le 1 \Rightarrow |x| \le 1$, and a graph of $f^{(5)}(x)$ shows that $\left|f^{(5)}(x)\right| \le 5$ for $-1 \le x \le 1$. Thus, we can take $M = 5$ and get $|R_4(x)| \le \dfrac{5}{5!} \cdot 1^5 = \dfrac{1}{24} = 0.041\overline{6}$.

(c)



From the graph of $|R_4(x)| = |x \sin x - T_4(x)|$, it seems that the error is less than 0.0082 on $[-1, 1]$.

22.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sinh 2x$	0
1	$2\cosh 2x$	2
2	$4\sinh 2x$	0
3	$8\cosh 2x$	8
4	$16\sinh 2x$	0
5	$32\cosh 2x$	32
6	$64\sinh 2x$	

(a) $f(x) = \sinh 2x \approx T_5(x) = 2x + \frac{8}{3!}x^3 + \frac{32}{5!}x^5 = 2x + \frac{4}{3}x^3 + \frac{4}{15}x^5$

(b) $|R_5(x)| \le \frac{M}{6!}|x|^6$, where $\left|f^{(6)}(x)\right| \le M$. For x in $[-1, 1]$, we have $|x| \le 1$. Since $f^{(6)}(x)$ is an increasing odd function on $[-1, 1]$, we see that $\left|f^{(6)}(x)\right| \le f^{(6)}(1) = 64\sinh 2 = 32(e^2 - e^{-2}) \approx 232.119$, so we can take $M = 232.12$ and get $|R_5(x)| \le \frac{232.12}{720} \cdot 1^6 \approx 0.3224$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c)



From the graph of $|R_5(x)| = |\sinh 2x - T_5(x)|$, it seems that the error is less than 0.027 on $[-1, 1]$.

23. From Exercise 5, $\cos x = -\left(x - \frac{\pi}{2}\right) + \frac{1}{6}\left(x - \frac{\pi}{2}\right)^3 + R_3(x)$, where $|R_3(x)| \le \dfrac{M}{4!}\left|x - \frac{\pi}{2}\right|^4$ with $\left|f^{(4)}(x)\right| = |\cos x| \le M = 1$. Now $x = 80° = (90° - 10°) = \left(\frac{\pi}{2} - \frac{\pi}{18}\right) = \frac{4\pi}{9}$ radians, so the error is $\left|R_3\left(\frac{4\pi}{9}\right)\right| \le \frac{1}{24}\left(\frac{\pi}{18}\right)^4 \approx 0.000\,039$, which means our estimate would *not* be accurate to five decimal places. However, $T_3 = T_4$, so we can use $\left|R_4\left(\frac{4\pi}{9}\right)\right| \le \frac{1}{120}\left(\frac{\pi}{18}\right)^5 \approx 0.000\,001$. Therefore, to five decimal places, $\cos 80° \approx -\left(-\frac{\pi}{18}\right) + \frac{1}{6}\left(-\frac{\pi}{18}\right)^3 \approx 0.17365$.

24. From Exercise 16, $\sin x = \frac{1}{2} + \frac{\sqrt{3}}{2}\left(x - \frac{\pi}{6}\right) - \frac{1}{4}\left(x - \frac{\pi}{6}\right)^2 - \frac{\sqrt{3}}{12}\left(x - \frac{\pi}{6}\right)^3 + \frac{1}{48}\left(x - \frac{\pi}{6}\right)^4 + R_4(x)$, where $|R_4(x)| \le \dfrac{M}{5!}\left|x - \frac{\pi}{6}\right|^5$ with $\left|f^{(5)}(x)\right| = |\cos x| \le M = 1$. Now $x = 38° = (30° + 8°) = \left(\frac{\pi}{6} + \frac{2\pi}{45}\right)$ radians, so the error is $\left|R_4\left(\frac{38\pi}{180}\right)\right| \le \frac{1}{120}\left(\frac{2\pi}{45}\right)^5 \approx 0.000\,000\,44$, which means our estimate will be accurate to five decimal places. Therefore, to five decimal places, $\sin 38° = \frac{1}{2} + \frac{\sqrt{3}}{2}\left(\frac{2\pi}{45}\right) - \frac{1}{4}\left(\frac{2\pi}{45}\right)^2 - \frac{\sqrt{3}}{12}\left(\frac{2\pi}{45}\right)^3 + \frac{1}{48}\left(\frac{2\pi}{45}\right)^4 \approx 0.61566$.

25. All derivatives of e^x are e^x, so $|R_n(x)| \le \dfrac{e^x}{(n+1)!}|x|^{n+1}$, where $0 < x < 0.1$. Letting $x = 0.1$, $R_n(0.1) \le \dfrac{e^{0.1}}{(n+1)!}(0.1)^{n+1} < 0.00001$, and by trial and error we find that $n = 3$ satisfies this inequality since $R_3(0.1) < 0.0000046$. Thus, by adding the four terms of the Maclaurin series for e^x corresponding to $n = 0, 1, 2$, and 3, we can estimate $e^{0.1}$ to within 0.00001. (In fact, this sum is $1.10516\overline{6}$ and $e^{0.1} \approx 1.10517$.)

26. From Table 1 in Section 11.10, $\ln(1 + x) = \sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{x^n}{n}$ for $|x| < 1$. Thus, $\ln 1.4 = \ln(1 + 0.4) = \sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{(0.4)^n}{n}$. Since this is an alternating series, the error is less than the first neglected term by the Alternating Series Estimation Theorem, and we find that $|a_6| = (0.4)^6/6 \approx 0.0007 < 0.001$. So we need the first five (nonzero) terms of the Maclaurin series for the desired accuracy. (In fact, this sum is approximately 0.33698 and $\ln 1.4 \approx 0.33647$.)

27. $\sin x = x - \dfrac{1}{3!}x^3 + \dfrac{1}{5!}x^5 - \cdots$. By the Alternating Series Estimation Theorem, the error in the approximation $\sin x = x - \dfrac{1}{3!}x^3$ is less than $\left|\dfrac{1}{5!}x^5\right| < 0.01 \iff$ $|x^5| < 120(0.01) \iff |x| < (1.2)^{1/5} \approx 1.037$. The curves $y = x - \frac{1}{6}x^3$ and $y = \sin x - 0.01$ intersect at $x \approx 1.043$, so the graph confirms our estimate. Since both the sine function



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and the given approximation are odd functions, we need to check the estimate only for $x > 0$. Thus, the desired range of values for x is $-1.037 < x < 1.037$.

28. $\cos x = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \cdots$. By the Alternating Series Estimation Theorem, the error is less than $\left|-\frac{1}{6!}x^6\right| < 0.005 \Leftrightarrow$ $x^6 < 720(0.005) \Leftrightarrow |x| < (3.6)^{1/6} \approx 1.238$. The curves $y = 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4$ and $y = \cos x + 0.005$ intersect at $x \approx 1.244$, so the graph confirms our estimate. Since both the cosine function and the given approximation are even functions, we need to check the estimate only for $x > 0$. Thus, the desired range of values for x is $-1.238 < x < 1.238$.



29. $\arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots$. By the Alternating Series Estimation Theorem, the error is less than $\left|-\frac{1}{7}x^7\right| < 0.05 \Leftrightarrow$ $|x^7| < 0.35 \Leftrightarrow |x| < (0.35)^{1/7} \approx 0.8607$. The curves $y = x - \frac{1}{3}x^3 + \frac{1}{5}x^5$ and $y = \arctan x + 0.05$ intersect at $x \approx 0.9245$, so the graph confirms our estimate. Since both the arctangent function and the given approximation are odd functions, we need to check the estimate only for $x > 0$. Thus, the desired range of values for x is $-0.86 < x < 0.86$.



30. $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(4)}{n!}(x-4)^n = \sum_{n=0}^{\infty} \frac{(-1)^n\, n!}{3^n(n+1)\, n!}(x-4)^n = \sum_{n=0}^{\infty} \frac{(-1)^n}{3^n(n+1)}(x-4)^n$. Now $f(5) = \sum_{n=0}^{\infty} \frac{(-1)^n}{3^n(n+1)} = \sum_{n=0}^{\infty} (-1)^n b_n$ is the sum of an alternating series that satisfies (i) $b_{n+1} \le b_n$ and (ii) $\lim_{n\to\infty} b_n = 0$, so by the Alternating Series Estimation Theorem, $|R_5(5)| = |f(5) - T_5(5)| \le b_6$, and $b_6 = \frac{1}{3^6(7)} = \frac{1}{5103} \approx 0.000196 < 0.0002$; that is, the fifth-degree Taylor polynomial approximates $f(5)$ with error less than 0.0002.

31. Let $s(t)$ be the position function of the car, and for convenience set $s(0) = 0$. The velocity of the car is $v(t) = s'(t)$ and the acceleration is $a(t) = s''(t)$, so the second degree Taylor polynomial is $T_2(t) = s(0) + v(0)t + \frac{a(0)}{2}t^2 = 20t + t^2$. We estimate the distance traveled during the next second to be $s(1) \approx T_2(1) = 20 + 1 = 21$ m. The function $T_2(t)$ would not be accurate over a full minute, since the car could not possibly maintain an acceleration of $2\ \text{m/s}^2$ for that long (if it did, its final speed would be $140\ \text{m/s} \approx 313\ \text{mi/h}$!).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

32. (a)

n	$\rho^{(n)}(t)$	$\rho^{(n)}(20)$
0	$\rho_{20}e^{\alpha(t-20)}$	ρ_{20}
1	$\alpha\rho_{20}e^{\alpha(t-20)}$	$\alpha\rho_{20}$
2	$\alpha^2\rho_{20}e^{\alpha(t-20)}$	$\alpha^2\rho_{20}$

The linear approximation is

$$T_1(t) = \rho(20) + \rho'(20)(t-20) = \rho_{20}[1 + \alpha(t-20)]$$

The quadratic approximation is

$$T_2(t) = \rho(20) + \rho'(20)(t-20) + \frac{\rho''(20)}{2}(t-20)^2$$
$$= \rho_{20}\left[1 + \alpha(t-20) + \tfrac{1}{2}\alpha^2(t-20)^2\right]$$

(b)



(c)



From the graph, it seems that $T_1(t)$ is within 1% of $\rho(t)$, that is, $0.99\rho(t) \le T_1(t) \le 1.01\rho(t)$, for $-14°\text{C} \le t \le 58°\text{C}$.

33. $E = \dfrac{q}{D^2} - \dfrac{q}{(D+d)^2} = \dfrac{q}{D^2} - \dfrac{q}{D^2(1+d/D)^2} = \dfrac{q}{D^2}\left[1 - \left(1 + \dfrac{d}{D}\right)^{-2}\right]$.

We use the Binomial Series to expand $(1 + d/D)^{-2}$:

$$E = \frac{q}{D^2}\left[1 - \left(1 - 2\left(\frac{d}{D}\right) + \frac{2\cdot 3}{2!}\left(\frac{d}{D}\right)^2 - \frac{2\cdot 3\cdot 4}{3!}\left(\frac{d}{D}\right)^3 + \cdots\right)\right] = \frac{q}{D^2}\left[2\left(\frac{d}{D}\right) - 3\left(\frac{d}{D}\right)^2 + 4\left(\frac{d}{D}\right)^3 - \cdots\right]$$
$$\approx \frac{q}{D^2}\cdot 2\left(\frac{d}{D}\right) = 2qd\cdot\frac{1}{D^3}$$

when D is much larger than d; that is, when P is far away from the dipole.

34. (a) $\dfrac{n_1}{\ell_o} + \dfrac{n_2}{\ell_i} = \dfrac{1}{R}\left(\dfrac{n_2 s_i}{\ell_i} - \dfrac{n_1 s_o}{\ell_o}\right)$ [Equation 1] where

$$\ell_o = \sqrt{R^2 + (s_o + R)^2 - 2R(s_o + R)\cos\phi} \quad \text{and} \quad \ell_i = \sqrt{R^2 + (s_i - R)^2 + 2R(s_i - R)\cos\phi} \quad \textbf{(2)}$$

Using $\cos\phi \approx 1$ gives

$$\ell_o = \sqrt{R^2 + (s_o + R)^2 - 2R(s_o + R)} = \sqrt{R^2 + s_o^2 + 2Rs_o + R^2 - 2Rs_o - 2R^2} = \sqrt{s_o^2} = s_o$$

and similarly, $\ell_i = s_i$. Thus, Equation 1 becomes $\dfrac{n_1}{s_o} + \dfrac{n_2}{s_i} = \dfrac{1}{R}\left(\dfrac{n_2 s_i}{s_i} - \dfrac{n_1 s_o}{s_o}\right) \Rightarrow \dfrac{n_1}{s_o} + \dfrac{n_2}{s_i} = \dfrac{n_2 - n_1}{R}$.

(b) Using $\cos\phi \approx 1 - \frac{1}{2}\phi^2$ in **(2)** gives us

$$\ell_o = \sqrt{R^2 + (s_o + R)^2 - 2R(s_o + R)\left(1 - \tfrac{1}{2}\phi^2\right)}$$
$$= \sqrt{R^2 + s_o^2 + 2Rs_o + R^2 - 2Rs_o + Rs_o\phi^2 - 2R^2 + R^2\phi^2} = \sqrt{s_o^2 + Rs_o\phi^2 + R^2\phi^2}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Anticipating that we will use the binomial series expansion $(1+x)^k \approx 1+kx$, we can write the last expression for ℓ_o as $s_o\sqrt{1+\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)}$ and similarly, $\ell_i = s_i\sqrt{1-\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)}$. Thus, from Equation 1,

$$\frac{n_1}{\ell_o}+\frac{n_2}{\ell_i}=\frac{1}{R}\left(\frac{n_2 s_i}{\ell_i}-\frac{n_1 s_o}{\ell_o}\right) \quad \Leftrightarrow \quad n_1\ell_o^{-1}+n_2\ell_i^{-1}=\frac{n_2}{R}\cdot\frac{s_i}{\ell_i}-\frac{n_1}{R}\cdot\frac{s_o}{\ell_o} \quad \Leftrightarrow$$

$$\frac{n_1}{s_o}\left[1+\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\right]^{-1/2}+\frac{n_2}{s_i}\left[1-\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\right]^{-1/2}$$

$$=\frac{n_2}{R}\left[1-\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\right]^{-1/2}-\frac{n_1}{R}\left[1+\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\right]^{-1/2}$$

Approximating the expressions for ℓ_o^{-1} and ℓ_i^{-1} by the first two terms in their binomial series, we get

$$\frac{n_1}{s_o}\left[1-\tfrac{1}{2}\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\right]+\frac{n_2}{s_i}\left[1+\tfrac{1}{2}\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\right]$$

$$=\frac{n_2}{R}\left[1+\tfrac{1}{2}\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\right]-\frac{n_1}{R}\left[1-\tfrac{1}{2}\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\right] \quad \Leftrightarrow$$

$$\frac{n_1}{s_o}-\frac{n_1\phi^2}{2s_o}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)+\frac{n_2}{s_i}+\frac{n_2\phi^2}{2s_i}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)=\frac{n_2}{R}+\frac{n_2\phi^2}{2R}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)-\frac{n_1}{R}+\frac{n_1\phi^2}{2R}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right) \quad \Leftrightarrow$$

$$\frac{n_1}{s_o}+\frac{n_2}{s_i}=\frac{n_2}{R}-\frac{n_1}{R}+\frac{n_1\phi^2}{2s_o}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)+\frac{n_1\phi^2}{2R}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)+\frac{n_2\phi^2}{2R}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)-\frac{n_2\phi^2}{2s_i}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)$$

$$=\frac{n_2-n_1}{R}+\frac{n_1\phi^2}{2}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\left(\frac{1}{s_o}+\frac{1}{R}\right)+\frac{n_2\phi^2}{2}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\left(\frac{1}{R}-\frac{1}{s_i}\right)$$

$$=\frac{n_2-n_1}{R}+\frac{n_1\phi^2R^2}{2s_o}\left(\frac{1}{R}+\frac{1}{s_o}\right)\left(\frac{1}{R}+\frac{1}{s_o}\right)+\frac{n_2\phi^2R^2}{2s_i}\left(\frac{1}{R}-\frac{1}{s_i}\right)\left(\frac{1}{R}-\frac{1}{s_i}\right)$$

$$=\frac{n_2-n_1}{R}+\phi^2R^2\left[\frac{n_1}{2s_o}\left(\frac{1}{R}+\frac{1}{s_o}\right)^2+\frac{n_2}{2s_i}\left(\frac{1}{R}-\frac{1}{s_i}\right)^2\right]$$

From Figure 8, we see that $\sin\phi = h/R$. So if we approximate $\sin\phi$ with ϕ, we get $h = R\phi$ and $h^2 = \phi^2R^2$ and hence, Equation 4, as desired.

35. (a) If the water is deep, then $2\pi d/L$ is large, and we know that $\tanh x \to 1$ as $x \to \infty$. So we can approximate $\tanh(2\pi d/L) \approx 1$, and so $v^2 \approx gL/(2\pi) \quad \Leftrightarrow \quad v \approx \sqrt{gL/(2\pi)}$.

(b) From the table, the first term in the Maclaurin series of $\tanh x$ is x, so if the water is shallow, we can approximate $\tanh\dfrac{2\pi d}{L} \approx \dfrac{2\pi d}{L}$, and so $v^2 \approx \dfrac{gL}{2\pi}\cdot\dfrac{2\pi d}{L} \quad \Leftrightarrow \quad v \approx \sqrt{gd}$.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\tanh x$	0
1	$\operatorname{sech}^2 x$	1
2	$-2\operatorname{sech}^2 x \tanh x$	0
3	$2\operatorname{sech}^2 x\,(3\tanh^2 x - 1)$	−2

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c) Since $\tanh x$ is an odd function, its Maclaurin series is alternating, so the error in the approximation $\tanh\dfrac{2\pi d}{L} \approx \dfrac{2\pi d}{L}$ is less than the first neglected term, which is $\dfrac{|f'''(0)|}{3!}\left(\dfrac{2\pi d}{L}\right)^3 = \dfrac{1}{3}\left(\dfrac{2\pi d}{L}\right)^3$.

If $L \geq 10d$, then $\dfrac{1}{3}\left(\dfrac{2\pi d}{L}\right)^3 \leq \dfrac{1}{3}\left(2\pi \cdot \dfrac{1}{10}\right)^3 = \dfrac{\pi^3}{375}$, so the error in the approximation $v^2 = gd$ is less than $\dfrac{gL}{2\pi} \cdot \dfrac{\pi^3}{375} \approx 0.0132gL$.

36. First note that

$$2\pi\sqrt{R^2+d^2} - 4\pi R = 2\pi\left[\sqrt{R^2}\sqrt{1+\frac{d^2}{R^2}} - R\right]$$

$$\approx 2\pi\left[R\left(1+\frac{d^2}{R^2}\cdot\frac{1}{2}+\cdots\right) - R\right] \quad \text{[use the binomial series } 1+\tfrac{1}{2}x+\cdots \text{ for } \sqrt{1+x}\text{]}$$

$$= 2\pi\left[\left(R+\frac{d^2}{2R}+\cdots\right) - R\right] \approx \frac{\pi d^2}{R}$$

since for large R the other terms are comparatively small. Now $U = \frac{1}{2}mv^2 = \frac{1}{2}m\left(\sqrt{k_e}\sqrt{R^2+d^2}\right)^2 = \frac{1}{2}mk_e(R^2+d^2) \approx \dfrac{\pi k_e R^2 m}{R}$ by the preceding approximation.

37. (a) L is the length of the arc subtended by the angle θ, so $L = R\theta \Rightarrow \theta = L/R$. Now $\sec\theta = (R+C)/R \Rightarrow R\sec\theta = R+C \Rightarrow C = R\sec\theta - R = R\sec(L/R) - R$.



(b) First we'll find a Taylor polynomial $T_4(x)$ for $f(x) = \sec x$ at $x = 0$.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sec x$	1
1	$\sec x\tan x$	0
2	$\sec x(2\tan^2 x + 1)$	1
3	$\sec x\tan x(6\tan^2 x + 5)$	0
4	$\sec x(24\tan^4 x + 28\tan^2 x + 5)$	5

Thus, $f(x) = \sec x \approx T_4(x) = 1 - \frac{1}{2}(x-0)^2 + \frac{5}{4!}(x-0)^4 = 1 - \frac{1}{2}x^2 + \frac{5}{24}x^4$. By part (a),

$$C \approx R\left[1 - \frac{1}{2}\left(\frac{L}{R}\right)^2 + \frac{5}{24}\left(\frac{L}{R}\right)^4\right] - R = R - \frac{1}{2}R\cdot\frac{L^2}{R^2} + \frac{5}{24}R\cdot\frac{L^4}{R^4} - R = \frac{L^2}{2R} + \frac{5L^4}{24R^3}$$

(c) Taking $L = 100$ km and $R = 6370$ km, the formula in part (a) says that

$$C = R\sec(L/R) - R = 6370\sec(100/6370) - 6370 \approx 0.785\,009\,965\,44 \text{ km}.$$

The formula in part (b) says that $C \approx \dfrac{L^2}{2R} + \dfrac{5L^4}{24R^3} = \dfrac{100^2}{2\cdot 6370} + \dfrac{5\cdot 100^4}{24\cdot 6370^3} \approx 0.785\,009\,957\,36$ km.

The difference between these two results is only $0.000\,000\,008\,08$ km, or $0.000\,008\,08$ m!

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. (a) $4\sqrt{\frac{L}{g}}\int_0^{\pi/2}\frac{dx}{\sqrt{1-k^2\sin^2 x}} = 4\sqrt{\frac{L}{g}}\int_0^{\pi/2}\left[1+(-k^2\sin^2 x)\right]^{-1/2}\,dx$

$$= 4\sqrt{\frac{L}{g}}\int_0^{\pi/2}\left[1-\frac{1}{2}(-k^2\sin^2 x)+\frac{\frac{1}{2}\cdot\frac{3}{2}}{2!}(-k^2\sin^2 x)^2-\frac{\frac{1}{2}\cdot\frac{3}{2}\cdot\frac{5}{2}}{3!}(-k^2\sin^2 x)^3+\cdots\right]dx$$

$$= 4\sqrt{\frac{L}{g}}\int_0^{\pi/2}\left[1+\left(\frac{1}{2}\right)k^2\sin^2 x+\left(\frac{1\cdot 3}{2\cdot 4}\right)k^4\sin^4 x+\left(\frac{1\cdot 3\cdot 5}{2\cdot 4\cdot 6}\right)k^6\sin^6 x+\cdots\right]dx$$

$$= 4\sqrt{\frac{L}{g}}\left[\frac{\pi}{2}+\left(\frac{1}{2}\right)\left(\frac{1}{2}\cdot\frac{\pi}{2}\right)k^2+\left(\frac{1\cdot 3}{2\cdot 4}\right)\left(\frac{1\cdot 3}{2\cdot 4}\cdot\frac{\pi}{2}\right)k^4+\left(\frac{1\cdot 3\cdot 5}{2\cdot 4\cdot 6}\right)\left(\frac{1\cdot 3\cdot 5}{2\cdot 4\cdot 6}\cdot\frac{\pi}{2}\right)k^6+\cdots\right]$$

[split up the integral and use the result from Exercise 7.1.50]

$$= 2\pi\sqrt{\frac{L}{g}}\left[1+\frac{1^2}{2^2}k^2+\frac{1^2\cdot 3^2}{2^2\cdot 4^2}k^4+\frac{1^2\cdot 3^2\cdot 5^2}{2^2\cdot 4^2\cdot 6^2}k^6+\cdots\right]$$

(b) The first of the two inequalities is true because all of the terms in the series are positive. For the second,

$$T = 2\pi\sqrt{\frac{L}{g}}\left[1+\frac{1^2}{2^2}k^2+\frac{1^2\cdot 3^2}{2^2\cdot 4^2}k^4+\frac{1^2\cdot 3^2\cdot 5^2}{2^2\cdot 4^2\cdot 6^2}k^6+\frac{1^2\cdot 3^2\cdot 5^2\cdot 7^2}{2^2\cdot 4^2\cdot 6^2\cdot 8^2}k^8+\cdots\right]$$

$$\le 2\pi\sqrt{\frac{L}{g}}\left[1+\tfrac{1}{4}k^2+\tfrac{1}{4}k^4+\tfrac{1}{4}k^6+\tfrac{1}{4}k^8+\cdots\right]$$

The terms in brackets (after the first) form a geometric series with $a=\frac{1}{4}k^2$ and $r=k^2=\sin^2\left(\frac{1}{2}\theta_0\right)<1$.

So $T \le 2\pi\sqrt{\frac{L}{g}}\left[1+\frac{k^2/4}{1-k^2}\right] = 2\pi\sqrt{\frac{L}{g}}\,\frac{4-3k^2}{4-4k^2}$.

(c) We substitute $L=1$, $g=9.8$, and $k=\sin(10^\circ/2)\approx 0.08716$, and the inequality from part (b) becomes $2.01090 \le T \le 2.01093$, so $T\approx 2.0109$. The estimate $T\approx 2\pi\sqrt{L/g}\approx 2.0071$ differs by about 0.2%.

If $\theta_0 = 42^\circ$, then $k\approx 0.35837$ and the inequality becomes $2.07153 \le T \le 2.08103$, so $T\approx 2.0763$.

The one-term estimate is the same, and the discrepancy between the two estimates increases to about 3.4%.

39. Using $f(x)=T_n(x)+R_n(x)$ with $n=1$ and $x=r$, we have $f(r)=T_1(r)+R_1(r)$, where T_1 is the first-degree Taylor polynomial of f at a. Because $a=x_n$, $f(r)=f(x_n)+f'(x_n)(r-x_n)+R_1(r)$. But r is a root of f, so $f(r)=0$ and we have $0=f(x_n)+f'(x_n)(r-x_n)+R_1(r)$. Taking the first two terms to the left side gives us $f'(x_n)(x_n-r)-f(x_n)=R_1(r)$. Dividing by $f'(x_n)$, we get $x_n-r-\dfrac{f(x_n)}{f'(x_n)}=\dfrac{R_1(r)}{f'(x_n)}$. By the formula for Newton's method, the left side of the preceding equation is $x_{n+1}-r$, so $|x_{n+1}-r|=\left|\dfrac{R_1(r)}{f'(x_n)}\right|$. Taylor's Inequality gives us $|R_1(r)|\le\dfrac{|f''(r)|}{2!}|r-x_n|^2$. Combining this inequality with the facts $|f''(x)|\le M$ and $|f'(x)|\ge K$ gives us $|x_{n+1}-r|\le\dfrac{M}{2K}|x_n-r|^2$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

APPLIED PROJECT Radiation from the Stars

1. If we write $f(\lambda) = \dfrac{8\pi hc\lambda^{-5}}{e^{hc/(\lambda kT)} - 1} = \dfrac{a\lambda^{-5}}{e^{b/(\lambda T)} - 1}$, then as $\lambda \to 0^+$, it is of the form ∞/∞, and as $\lambda \to \infty$ it is of the form $0/0$, so in either case we can use l'Hospital's Rule. First of all,

$$\lim_{\lambda\to\infty} f(\lambda) \overset{\text{H}}{=} \lim_{\lambda\to\infty} \frac{a\left(-5\lambda^{-6}\right)}{-\dfrac{bT}{(\lambda T)^2}e^{b/(\lambda T)}} = 5\frac{aT}{b}\lim_{\lambda\to\infty}\frac{\lambda^2\lambda^{-6}}{e^{b/(\lambda T)}} = 5\frac{aT}{b}\lim_{\lambda\to\infty}\frac{\lambda^{-4}}{e^{b/(\lambda T)}} = 0$$

Also,
$$\lim_{\lambda\to 0^+} f(\lambda) \overset{\text{H}}{=} 5\frac{aT}{b}\lim_{\lambda\to 0^+}\frac{\lambda^{-4}}{e^{b/(\lambda T)}} \overset{\text{H}}{=} 5\frac{aT}{b}\lim_{\lambda\to 0^+}\frac{-4\lambda^{-5}}{-\dfrac{bT}{(\lambda T)^2}e^{b/(\lambda T)}} = 20\frac{aT}{b^2}^2\lim_{\lambda\to 0^+}\frac{\lambda^{-3}}{e^{b/(\lambda T)}}$$

This is still indeterminate, but note that each time we use l'Hospital's Rule, we gain a factor of λ in the numerator, as well as a constant factor, and the denominator is unchanged. So if we use l'Hospital's Rule three more times, the exponent of λ in the numerator will become 0. That is, for some $\{k_i\}$, all constant,

$$\lim_{\lambda\to 0^+} f(\lambda) \overset{\text{H}}{=} k_1\lim_{\lambda\to 0^+}\frac{\lambda^{-3}}{e^{b/(\lambda T)}} \overset{\text{H}}{=} k_2\lim_{\lambda\to 0^+}\frac{\lambda^{-2}}{e^{b/(\lambda T)}} \overset{\text{H}}{=} k_3\lim_{\lambda\to 0^-}\frac{\lambda^{-1}}{e^{b/(\lambda T)}} \overset{\text{H}}{=} k_4\lim_{\lambda\to 0^-}\frac{1}{e^{b/(\lambda T)}} = 0$$

2. We expand the denominator of Planck's Law using the Taylor series $e^x = 1 + x + \dfrac{x^2}{2!} + \dfrac{x^3}{3!} + \cdots$ with $x = \dfrac{hc}{\lambda kT}$, and use the fact that if λ is large, then all subsequent terms in the Taylor expansion are very small compared to the first one, so we can approximate using the Taylor polynomial T_1:

$$f(\lambda) = \frac{8\pi hc\lambda^{-5}}{e^{hc/(\lambda kT)} - 1} = \frac{8\pi hc\lambda^{-5}}{\left[1 + \dfrac{hc}{\lambda kT} + \dfrac{1}{2!}\left(\dfrac{hc}{\lambda kT}\right)^2 + \dfrac{1}{3!}\left(\dfrac{hc}{\lambda kT}\right)^3 + \cdots\right] - 1} \approx \frac{8\pi hc\lambda^{-5}}{\left(1 + \dfrac{hc}{\lambda kT}\right) - 1} = \frac{8\pi kT}{\lambda^4}$$

which is the Rayleigh-Jeans Law.

3. To convert to μm, we substitute $\lambda/10^6$ for λ in both laws. The first figure shows that the two laws are similar for large λ. The second figure shows that the two laws are very different for short wavelengths (Planck's Law gives a maximum at $\lambda \approx 0.51\ \mu$m; the Rayleigh-Jeans Law gives no minimum or maximum.).





4. From the graph in Problem 3, $f(\lambda)$ has a maximum under Planck's Law at $\lambda \approx 0.51\ \mu$m.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5.



As T gets larger, the total area under the curve increases, as we would expect: the hotter the star, the more energy it emits. Also, as T increases, the x-value of the maximum decreases, so the higher the temperature, the shorter the peak wavelength (and consequently the average wavelength) of light emitted. This is why Sirius is a blue star and Betelgeuse is a red star: most of Sirius's light is of a fairly short wavelength; that is, a higher frequency, toward the blue end of the spectrum, whereas most of Betelgeuse's light is of a lower frequency, toward the red end of the spectrum.

11 Review

TRUE-FALSE QUIZ

1. False. See Note 2 after Theorem 11.2.6.

2. False. The series $\sum_{n=1}^{\infty} n^{-\sin 1} = \sum_{n=1}^{\infty} \frac{1}{n^{\sin 1}}$ is a p-series with $p = \sin 1 \approx 0.84 \le 1$, so the series diverges.

3. True. If $\lim_{n\to\infty} a_n = L$, then as $n \to \infty$, $2n+1 \to \infty$, so $a_{2n+1} \to L$.

4. True by Theorem 11.8.4.

   Or: Use the Comparison Test to show that $\sum c_n(-2)^n$ converges absolutely.

5. False. For example, take $c_n = (-1)^n/(n6^n)$.

6. True by Theorem 11.8.4.

7. False, since $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{1}{(n+1)^3} \cdot \frac{n^3}{1}\right| = \lim_{n\to\infty} \left|\frac{n^3}{(n+1)^3} \cdot \frac{1/n^3}{1/n^3}\right| = \lim_{n\to\infty} \frac{1}{(1+1/n)^3} \quad 1.$

8. True, since $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{1}{(n+1)!} \cdot \frac{n!}{1}\right| = \lim_{n\to\infty} \frac{1}{n+1} = 0 < 1.$

9. False. See the note after Example 11.4.2.

10. True, since $\frac{1}{e} = e^{-1}$ and $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$, so $e^{-1} = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!}$.

11. True. See (9) in Section 11.1.

12. True, because if $\sum |a_n|$ is convergent, then so is $\sum a_n$ by Theorem 11.6.3.

13. True. By Theorem 11.10.5 the coefficient of x^3 is $\frac{f'''(0)}{3!} = \frac{1}{3} \quad \Rightarrow \quad f'''(0) = 2.$

    Or: Use Theorem 11.9.2 to differentiate f three times.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. False. Let $a_n = n$ and $b_n = -n$. Then $\{a_n\}$ and $\{b_n\}$ are divergent, but $a_n + b_n = 0$, so $\{a_n + b_n\}$ is convergent.

15. False. For example, let $a_n = b_n = (-1)^n$. Then $\{a_n\}$ and $\{b_n\}$ are divergent, but $a_n b_n = 1$, so $\{a_n b_n\}$ is convergent.

16. True by the Monotonic Sequence Theorem, since $\{a_n\}$ is decreasing and $0 < a_n \le a_1$ for all $n \;\Rightarrow\; \{a_n\}$ is bounded.

17. True by Theorem 11.6.3. $[\sum (-1)^n a_n$ is absolutely convergent and hence convergent.]

18. True. $\lim_{n\to\infty} \dfrac{a_{n+1}}{a_n} < 1 \;\Rightarrow\; \sum a_n$ converges (Ratio Test) $\;\Rightarrow\; \lim_{n\to\infty} a_n = 0$ [Theorem 11.2.6].

19. True. $0.99999\ldots = 0.9 + 0.9(0.1)^1 + 0.9(0.1)^2 + 0.9(0.1)^3 + \cdots = \sum_{n=1}^{\infty} (0.9)(0.1)^{n-1} = \dfrac{0.9}{1-0.1} = 1$ by the formula for the sum of a geometric series $[S = a_1/(1-r)]$ with ratio r satisfying $|r| < 1$.

20. True. Since $\lim_{n\to\infty} a_n = 2$, we know that $\lim_{n\to\infty} a_{n+3} = 2$. Thus, $\lim_{n\to\infty} (a_{n+3} - a_n) = \lim_{n\to\infty} a_{n+3} - \lim_{n\to\infty} a_n = 2 - 2 = 0$.

21. True. A finite number of terms doesn't affect convergence or divergence of a series.

22. False. Let $a_n = (0.1)^n$ and $b_n = (0.2)^n$. Then $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (0.1)^n = \dfrac{0.1}{1-0.1} = \dfrac{1}{9} = A$, $\sum_{n=1}^{\infty} b_n = \sum_{n=1}^{\infty} (0.2)^n = \dfrac{0.2}{1-0.2} = \dfrac{1}{4} = B$, and $\sum_{n=1}^{\infty} a_n b_n = \sum_{n=1}^{\infty} (0.02)^n = \dfrac{0.02}{1-0.02} = \dfrac{1}{49}$, but $AB = \frac{1}{9} \cdot \frac{1}{4} = \frac{1}{36}$.

EXERCISES

1. $\left\{ \dfrac{2+n^3}{1+2n^3} \right\}$ converges since $\lim_{n\to\infty} \dfrac{2+n^3}{1+2n^3} = \lim_{n\to\infty} \dfrac{2/n^3+1}{1/n^3+2} = \dfrac{1}{2}$.

2. $a_n = \dfrac{9^{n+1}}{10^n} = 9 \cdot \left(\frac{9}{10}\right)^n$, so $\lim_{n\to\infty} a_n = 9 \lim_{n\to\infty} \left(\frac{9}{10}\right)^n = 9 \cdot 0 = 0$ by (11.1.9).

3. $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \dfrac{n^3}{1+n^2} = \lim_{n\to\infty} \dfrac{n}{1/n^2+1} = \infty$, so the sequence diverges.

4. $a_n = \cos(n\pi/2)$, so $a_n = 0$ if n is odd and $a_n = \pm 1$ if n is even. As n increases, a_n keeps cycling through the values $0, 1, 0, -1$, so the sequence $\{a_n\}$ is divergent.

5. $|a_n| = \left| \dfrac{n \sin n}{n^2+1} \right| \le \dfrac{n}{n^2+1} < \dfrac{1}{n}$, so $|a_n| \to 0$ as $n \to \infty$. Thus, $\lim_{n\to\infty} a_n = 0$. The sequence $\{a_n\}$ is convergent.

6. $a_n = \dfrac{\ln n}{\sqrt{n}}$. Let $f(x) = \dfrac{\ln x}{\sqrt{x}}$ for $x > 0$. Then $\lim_{x\to\infty} f(x) = \lim_{x\to\infty} \dfrac{\ln x}{\sqrt{x}} \overset{\text{H}}{=} \lim_{x\to\infty} \dfrac{1/x}{1/(2\sqrt{x})} = \lim_{x\to\infty} \dfrac{2}{\sqrt{x}} = 0$.

Thus, by Theorem 11.1.3, $\{a_n\}$ converges and $\lim_{n\to\infty} a_n = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

7. $\left\{\left(1+\dfrac{3}{n}\right)^{4n}\right\}$ is convergent. Let $y=\left(1+\dfrac{3}{x}\right)^{4x}$. Then

$\lim\limits_{x\to\infty}\ln y=\lim\limits_{x\to\infty}4x\ln(1+3/x)=\lim\limits_{x\to\infty}\dfrac{\ln(1+3/x)}{1/(4x)}\overset{\text{H}}{=}\lim\limits_{x\to\infty}\dfrac{\frac{1}{1+3/x}\left(-\frac{3}{x^2}\right)}{-1/(4x^2)}=\lim\limits_{x\to\infty}\dfrac{12}{1+3/x}=12$, so

$\lim\limits_{x\to\infty}y=\lim\limits_{n\to\infty}\left(1+\dfrac{3}{n}\right)^{4n}=e^{12}$.

8. $\left\{\dfrac{(-10)^n}{n!}\right\}$ converges, since $\dfrac{10^n}{n!}=\dfrac{10\cdot10\cdot10\cdot\cdots\cdot10}{1\cdot2\cdot3\cdot\cdots\cdot10}\cdot\dfrac{10\cdot10\cdot\cdots\cdot10}{11\cdot12\cdot\cdots\cdot n}\le10^{10}\left(\dfrac{10}{11}\right)^{n-10}\to0$ as $n\to\infty$, so

$\lim\limits_{n\to\infty}\dfrac{(-10)^n}{n!}=0$ [Squeeze Theorem]. *Or:* Use (11.10.10).

9. We use induction, hypothesizing that $a_{n-1}<a_n<2$. Note first that $1<a_2=\frac{1}{3}(1+4)=\frac{5}{3}<2$, so the hypothesis holds for $n=2$. Now assume that $a_{k-1}<a_k<2$. Then $a_k=\frac{1}{3}(a_{k-1}+4)<\frac{1}{3}(a_k+4)<\frac{1}{3}(2+4)=2$. So $a_k<a_{k+1}<2$, and the induction is complete. To find the limit of the sequence, we note that $L=\lim\limits_{n\to\infty}a_n=\lim\limits_{n\to\infty}a_{n+1}\ \Rightarrow$ $L=\frac{1}{3}(L+4)\ \Rightarrow\ L=2$.

10. $\lim\limits_{x\to\infty}\dfrac{x^4}{e^x}\overset{\text{H}}{=}\lim\limits_{x\to\infty}\dfrac{4x^3}{e^x}\overset{\text{H}}{=}\lim\limits_{x\to\infty}\dfrac{12x^2}{e^x}\overset{\text{H}}{=}\lim\limits_{x\to\infty}\dfrac{24x}{e^x}\overset{\text{H}}{=}\lim\limits_{x\to\infty}\dfrac{24}{e^x}=0$

Then we conclude from Theorem 11.1.3 that $\lim\limits_{n\to\infty}n^4e^{-n}=0$.

From the graph, it seems that $12^4e^{-12}>0.1$, but $n^4e^{-n}<0.1$ whenever $n>12$. So the smallest value of N corresponding to $\varepsilon=0.1$ in the definition of the limit is $N=12$.



11. $\dfrac{n}{n^3+1}<\dfrac{n}{n^3}=\dfrac{1}{n^2}$, so $\sum\limits_{n=1}^{\infty}\dfrac{n}{n^3+1}$ converges by the Comparison Test with the convergent p-series $\sum\limits_{n=1}^{\infty}\dfrac{1}{n^2}$ $[p=2>1]$.

12. Let $a_n=\dfrac{n^2+1}{n^3+1}$ and $b_n=\dfrac{1}{n}$, so $\lim\limits_{n\to\infty}\dfrac{a_n}{b_n}=\lim\limits_{n\to\infty}\dfrac{n^3+n}{n^3+1}=\lim\limits_{n\to\infty}\dfrac{1+1/n^2}{1+1/n^3}=1>0$.

Since $\sum\limits_{n=1}^{\infty}b_n$ is the divergent harmonic series, $\sum\limits_{n=1}^{\infty}a_n$ also diverges by the Limit Comparison Test.

13. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim\limits_{n\to\infty}\left[\dfrac{(n+1)^3}{5^{n+1}}\cdot\dfrac{5^n}{n^3}\right]=\lim\limits_{n\to\infty}\left(1+\dfrac{1}{n}\right)^3\cdot\dfrac{1}{5}=\dfrac{1}{5}<1$, so $\sum\limits_{n=1}^{\infty}\dfrac{n^3}{5^n}$ converges by the Ratio Test.

14. Let $b_n=\dfrac{1}{\sqrt{n+1}}$. Then b_n is positive for $n\ge1$, the sequence $\{b_n\}$ is decreasing, and $\lim\limits_{n\to\infty}b_n=0$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{\sqrt{n+1}}$ converges by the Alternating Series Test.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

15. Let $f(x) = \dfrac{1}{x\sqrt{\ln x}}$. Then f is continuous, positive, and decreasing on $[2,\infty)$, so the Integral Test applies.

$$\int_2^\infty f(x)\,dx = \lim_{t\to\infty}\int_2^t \frac{1}{x\sqrt{\ln x}}\,dx \quad \left[u=\ln x,\, du=\frac{1}{x}\,dx\right] \quad = \lim_{t\to\infty}\int_{\ln 2}^{\ln t} u^{-1/2}\,du = \lim_{t\to\infty}\left[2\sqrt{u}\right]_{\ln 2}^{\ln t}$$
$$= \lim_{t\to\infty}\left(2\sqrt{\ln t} - 2\sqrt{\ln 2}\right) = \infty,$$

so the series $\sum_{n=2}^{\infty} \dfrac{1}{n\sqrt{\ln n}}$ diverges.

16. $\lim_{n\to\infty} \dfrac{n}{3n+1} = \dfrac{1}{3}$, so $\lim_{n\to\infty} \ln\left(\dfrac{n}{3n+1}\right) = \ln\frac{1}{3} \neq 0$. Thus, the series $\sum_{n=1}^{\infty} \ln\left(\dfrac{n}{3n+1}\right)$ diverges by the Test for Divergence.

17. $|a_n| = \left|\dfrac{\cos 3n}{1+(1.2)^n}\right| \le \dfrac{1}{1+(1.2)^n} < \dfrac{1}{(1.2)^n} = \left(\dfrac{5}{6}\right)^n$, so $\sum_{n=1}^{\infty} |a_n|$ converges by comparison with the convergent geometric series $\sum_{n=1}^{\infty} \left(\frac{5}{6}\right)^n \ \left[r=\frac{5}{6}<1\right]$. It follows that $\sum_{n=1}^{\infty} a_n$ converges (by Theorem 11.6.3).

18. $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} \sqrt[n]{\left|\dfrac{n^{2n}}{(1+2n^2)^n}\right|} = \lim_{n\to\infty} \dfrac{n^2}{1+2n^2} = \lim_{n\to\infty} \dfrac{1}{1/n^2+2} = \dfrac{1}{2} < 1$, so $\sum_{n=1}^{\infty} \dfrac{n^{2n}}{(1+2n^2)^n}$ converges by the Root Test.

19. $\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)(2n+1)}{5^{n+1}\,(n+1)!} \cdot \dfrac{5^n\,n!}{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)} = \lim_{n\to\infty} \dfrac{2n+1}{5(n+1)} = \dfrac{2}{5} < 1$, so the series converges by the Ratio Test.

20. $\sum_{n=1}^{\infty} \dfrac{(-5)^{2n}}{n^2\,9^n} = \sum_{n=1}^{\infty} \dfrac{1}{n^2}\left(\dfrac{25}{9}\right)^n$. Now $\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \dfrac{25^{n+1}}{(n+1)^2\cdot 9^{n+1}} \cdot \dfrac{n^2\cdot 9^n}{25^n} = \lim_{n\to\infty} \dfrac{25n^2}{9(n+1)^2} = \dfrac{25}{9} > 1$, so the series diverges by the Ratio Test.

21. $b_n = \dfrac{\sqrt{n}}{n+1} > 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series $\sum_{n=1}^{\infty} (-1)^{n-1}\dfrac{\sqrt{n}}{n+1}$ converges by the Alternating Series Test.

22. Use the Limit Comparison Test with $a_n = \dfrac{\sqrt{n+1}-\sqrt{n-1}}{n} = \dfrac{2}{n\left(\sqrt{n+1}+\sqrt{n-1}\right)}$ (rationalizing the numerator) and $b_n = \dfrac{1}{n^{3/2}}$. $\lim_{n\to\infty} \dfrac{a_n}{b_n} = \lim_{n\to\infty} \dfrac{2\sqrt{n}}{\sqrt{n+1}+\sqrt{n-1}} = 1$, so since $\sum_{n=1}^{\infty} b_n$ converges $\left[p=\frac{3}{2}>1\right]$, $\sum_{n=1}^{\infty} a_n$ converges also.

23. Consider the series of absolute values: $\sum_{n=1}^{\infty} n^{-1/3}$ is a p-series with $p = \frac{1}{3} < 1$ and is therefore divergent. But if we apply the Alternating Series Test, we see that $b_n = \dfrac{1}{\sqrt[3]{n}} > 0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n = 0$, so the series $\sum_{n=1}^{\infty} (-1)^{n-1} n^{-1/3}$ converges. Thus, $\sum_{n=1}^{\infty} (-1)^{n-1} n^{-1/3}$ is conditionally convergent.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $\sum_{n=1}^{\infty} \left|(-1)^{n-1} n^{-3}\right| = \sum_{n=1}^{\infty} n^{-3}$ is a convergent p-series $[p = 3 > 1]$. Therefore, $\sum_{n=1}^{\infty} (-1)^{n-1} n^{-3}$ is absolutely convergent.

25. $\left|\dfrac{a_{n+1}}{a_n}\right| = \left|\dfrac{(-1)^{n+1}(n+2)3^{n+1}}{2^{2n+3}} \cdot \dfrac{2^{2n+1}}{(-1)^n (n+1)3^n}\right| = \dfrac{n+2}{n+1} \cdot \dfrac{3}{4} = \dfrac{1+(2/n)}{1+(1/n)} \cdot \dfrac{3}{4} \to \dfrac{3}{4} < 1$ as $n \to \infty$, so by the Ratio Test, $\sum_{n=1}^{\infty} \dfrac{(-1)^n (n+1)3^n}{2^{2n+1}}$ is absolutely convergent.

26. $\lim_{x\to\infty} \dfrac{\sqrt{x}}{\ln x} \overset{\text{H}}{=} \lim_{x\to\infty} \dfrac{1/(2\sqrt{x})}{1/x} = \lim_{x\to\infty} \dfrac{\sqrt{x}}{2} = \infty$. Therefore, $\lim_{n\to\infty} \dfrac{(-1)^n \sqrt{n}}{\ln n} \neq 0$, so the given series is divergent by the Test for Divergence.

27. $\sum_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{2^{3n}} = \sum_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{(2^3)^n} = \sum_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{8^n} = \dfrac{1}{8} \sum_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{8^{n-1}} = \dfrac{1}{8} \sum_{n=1}^{\infty} \left(-\dfrac{3}{8}\right)^{n-1} = \dfrac{1}{8}\left(\dfrac{1}{1-(-3/8)}\right)$

$= \dfrac{1}{8} \cdot \dfrac{8}{11} = \dfrac{1}{11}$

28. $\sum_{n=1}^{\infty} \dfrac{1}{n(n+3)} = \sum_{n=1}^{\infty} \left[\dfrac{1}{3n} - \dfrac{1}{3(n+3)}\right]$ [partial fractions].

$s_n = \sum_{i=1}^{n} \left[\dfrac{1}{3i} - \dfrac{1}{3(i+3)}\right] = \dfrac{1}{3} + \dfrac{1}{6} + \dfrac{1}{9} - \dfrac{1}{3(n+1)} - \dfrac{1}{3(n+2)} - \dfrac{1}{3(n+3)}$ (telescoping sum), so

$\sum_{n=1}^{\infty} \dfrac{1}{n(n+3)} = \lim_{n\to\infty} s_n = \dfrac{1}{3} + \dfrac{1}{6} + \dfrac{1}{9} = \dfrac{11}{18}$.

29. $\sum_{n=1}^{\infty} [\tan^{-1}(n+1) - \tan^{-1} n] = \lim_{n\to\infty} s_n$

$= \lim_{n\to\infty} [(\tan^{-1} 2 - \tan^{-1} 1) + (\tan^{-1} 3 - \tan^{-1} 2) + \cdots + (\tan^{-1}(n+1) - \tan^{-1} n)]$

$= \lim_{n\to\infty} [\tan^{-1}(n+1) - \tan^{-1} 1] = \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}$

30. $\sum_{n=0}^{\infty} \dfrac{(-1)^n \pi^n}{3^{2n}\,(2n)!} = \sum_{n=0}^{\infty} (-1)^n \dfrac{1}{(2n)!} \cdot \dfrac{\pi^n}{3^{2n}} = \sum_{n=0}^{\infty} (-1)^n \dfrac{1}{(2n)!} \cdot \left(\dfrac{\sqrt{\pi}}{3}\right)^{2n} = \cos\left(\dfrac{\sqrt{\pi}}{3}\right)$ since $\cos x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n}}{(2n)!}$ for all x.

31. $1 - e + \dfrac{e^2}{2!} - \dfrac{e^3}{3!} + \dfrac{e^4}{4!} - \cdots = \sum_{n=0}^{\infty} (-1)^n \dfrac{e^n}{n!} = \sum_{n=0}^{\infty} \dfrac{(-e)^n}{n!} = e^{-e}$ since $e^x = \sum_{n=0}^{\infty} \dfrac{x^n}{n!}$ for all x.

32. $4.17\overline{326} = 4.17 + \dfrac{326}{10^5} + \dfrac{326}{10^8} + \cdots = 4.17 + \dfrac{326/10^5}{1 - 1/10^3} = \dfrac{417}{100} + \dfrac{326}{99{,}900} = \dfrac{416{,}909}{99{,}900}$

33. $\cosh x = \dfrac{1}{2}(e^x + e^{-x}) = \dfrac{1}{2}\left(\sum_{n=0}^{\infty} \dfrac{x^n}{n!} + \sum_{n=0}^{\infty} \dfrac{(-x)^n}{n!}\right)$

$= \dfrac{1}{2}\left[\left(1 + x + \dfrac{x^2}{2!} + \dfrac{x^3}{3!} + \dfrac{x^4}{4!} + \cdots\right) + \left(1 - x + \dfrac{x^2}{2!} - \dfrac{x^3}{3!} + \dfrac{x^4}{4!} - \cdots\right)\right]$

$= \dfrac{1}{2}\left(2 + 2 \cdot \dfrac{x^2}{2!} + 2 \cdot \dfrac{x^4}{4!} + \cdots\right) = 1 + \dfrac{1}{2}x^2 + \sum_{n=2}^{\infty} \dfrac{x^{2n}}{(2n)!} \geq 1 + \dfrac{1}{2}x^2$ for all x

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

34. $\sum_{n=1}^{\infty} (\ln x)^n$ is a geometric series which converges whenever $|\ln x| < 1 \;\Rightarrow\; -1 < \ln x < 1 \;\Rightarrow\; e^{-1} < x < e$.

35. $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^5} = 1 - \frac{1}{32} + \frac{1}{243} - \frac{1}{1024} + \frac{1}{3125} - \frac{1}{7776} + \frac{1}{16{,}807} - \frac{1}{32{,}768} + \cdots$.

Since $b_8 = \frac{1}{8^5} = \frac{1}{32{,}768} < 0.000031$, $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^5} \approx \sum_{n=1}^{7} \frac{(-1)^{n+1}}{n^5} \approx 0.9721$.

36. (a) $s_5 = \sum_{n=1}^{5} \frac{1}{n^6} = 1 + \frac{1}{2^6} + \cdots + \frac{1}{5^6} \approx 1.017305$. The series $\sum_{n=1}^{\infty} \frac{1}{n^6}$ converges by the Integral Test, so we estimate the remainder R_5 with (11.3.2): $R_5 \le \int_5^{\infty} \frac{dx}{x^6} = \left[-\frac{x^{-5}}{5}\right]_5^{\infty} = \frac{5^{-5}}{5} = 0.000064$. So the error is at most 0.000064.

(b) In general, $R_n \le \int_n^{\infty} \frac{dx}{x^6} = \frac{1}{5n^5}$. If we take $n = 9$, then $s_9 \approx 1.01734$ and $R_9 \le \frac{1}{5 \cdot 9^5} \approx 3.4 \times 10^{-6}$.

So to five decimal places, $\sum_{n=1}^{\infty} \frac{1}{n^5} \approx \sum_{n=1}^{9} \frac{1}{n^5} \approx 1.01734$.

Another method: Use (11.3.3) instead of (11.3.2).

37. $\sum_{n=1}^{\infty} \frac{1}{2+5^n} \approx \sum_{n=1}^{8} \frac{1}{2+5^n} \approx 0.18976224$. To estimate the error, note that $\frac{1}{2+5^n} < \frac{1}{5^n}$, so the remainder term is

$R_8 = \sum_{n=9}^{\infty} \frac{1}{2+5^n} < \sum_{n=9}^{\infty} \frac{1}{5^n} = \frac{1/5^9}{1-1/5} = 6.4 \times 10^{-7}$ $\left[\text{geometric series with } a = \frac{1}{5^9} \text{ and } r = \frac{1}{5}\right]$.

38. (a) $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(n+1)^{n+1}}{[2(n+1)]!} \cdot \frac{(2n)!}{n^n}\right| = \lim_{n\to\infty} \frac{(n+1)^n (n+1)^1}{(2n+2)(2n+1)n^n} = \lim_{n\to\infty} \left(\frac{n+1}{n}\right)^n \frac{1}{2(2n+1)}$

$= \lim_{n\to\infty} \left(1 + \frac{1}{n}\right)^n \frac{1}{2(2n+1)} = e \cdot 0 = 0 < 1$

so the series converges by the Ratio Test.

(b) The series in part (a) is convergent, so $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{n^n}{(2n)!} = 0$ by Theorem 11.2.6.

39. Use the Limit Comparison Test. $\lim_{n\to\infty} \left|\frac{\left(\frac{n+1}{n}\right)a_n}{a_n}\right| = \lim_{n\to\infty} \frac{n+1}{n} = \lim_{n\to\infty} \left(1 + \frac{1}{n}\right) = 1 > 0$.

Since $\sum |a_n|$ is convergent, so is $\sum \left|\left(\frac{n+1}{n}\right)a_n\right|$, by the Limit Comparison Test.

40. $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{n+1}}{(n+1)^2\, 5^{n+1}} \cdot \frac{n^2 5^n}{x^n}\right| = \lim_{n\to\infty} \frac{1}{(1+1/n)^2} \frac{|x|}{5} = \frac{|x|}{5}$, so by the Ratio Test, $\sum_{n=1}^{\infty} (-1)^n \frac{x^n}{n^2\, 5^n}$ converges when $\frac{|x|}{5} < 1 \quad \Leftrightarrow \quad |x| < 5$, so $R = 5$. When $x = -5$, the series becomes the convergent p-series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ with $p = 2 > 1$. When $x = 5$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$, which converges by the Alternating Series Test. Thus, $I = [-5, 5]$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left[\dfrac{|x+2|^{n+1}}{(n+1)\,4^{n+1}}\cdot\dfrac{n\,4^n}{|x+2|^n}\right] = \lim\limits_{n\to\infty}\left[\dfrac{n}{n+1}\dfrac{|x+2|}{4}\right] = \dfrac{|x+2|}{4} < 1 \quad\Leftrightarrow\quad |x+2|<4$, so $R=4$.

$|x+2|<4 \quad\Leftrightarrow\quad -4<x+2<4 \quad\Leftrightarrow\quad -6<x<2$. If $x=-6$, then the series $\sum\limits_{n=1}^{\infty}\dfrac{(x+2)^n}{n\,4^n}$ becomes $\sum\limits_{n=1}^{\infty}\dfrac{(-4)^n}{n4^n} = \sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{n}$, the alternating harmonic series, which converges by the Alternating Series Test. When $x=2$, the series becomes the harmonic series $\sum\limits_{n=1}^{\infty}\dfrac{1}{n}$, which diverges. Thus, $I=[-6,2)$.

42. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{2^{n+1}\,(x-2)^{n+1}}{(n+3)!}\cdot\dfrac{(n+2)!}{2^n(x-2)^n}\right| = \lim\limits_{n\to\infty}\dfrac{2}{n+3}\,|x-2| = 0 < 1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{2^n\,(x-2)^n}{(n+2)!}$ converges for all x. $R=\infty$ and $I=(-\infty,\infty)$.

43. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{2^{n+1}(x-3)^{n+1}}{\sqrt{n+4}}\cdot\dfrac{\sqrt{n+3}}{2^n(x-3)^n}\right| = 2\,|x-3|\lim\limits_{n\to\infty}\sqrt{\dfrac{n+3}{n+4}} = 2\,|x-3| < 1 \quad\Leftrightarrow\quad |x-3|<\tfrac{1}{2}$,

so $R=\frac{1}{2}$. $|x-3|<\frac{1}{2} \quad\Leftrightarrow\quad -\frac{1}{2}<x-3<\frac{1}{2} \quad\Leftrightarrow\quad \frac{5}{2}<x<\frac{7}{2}$. For $x=\frac{7}{2}$, the series $\sum\limits_{n=1}^{\infty}\dfrac{2^n(x-3)^n}{\sqrt{n+3}}$ becomes $\sum\limits_{n=0}^{\infty}\dfrac{1}{\sqrt{n+3}} = \sum\limits_{n=3}^{\infty}\dfrac{1}{n^{1/2}}$, which diverges $\left[p=\frac{1}{2}\le 1\right]$, but for $x=\frac{5}{2}$, we get $\sum\limits_{n=0}^{\infty}\dfrac{(-1)^n}{\sqrt{n+3}}$, which is a convergent alternating series, so $I=\left[\frac{5}{2},\frac{7}{2}\right)$.

44. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(2n+2)!\,x^{n+1}}{[(n+1)!]^2}\cdot\dfrac{(n!)^2}{(2n)!\,x^n}\right| = \lim\limits_{n\to\infty}\dfrac{(2n+2)(2n+1)}{(n+1)(n+1)}\,|x| = 4\,|x|$.

To converge, we must have $4\,|x|<1 \quad\Leftrightarrow\quad |x|<\frac{1}{4}$, so $R=\frac{1}{4}$.

45.

n	$f^{(n)}(x)$	$f^{(n)}\left(\frac{\pi}{6}\right)$
0	$\sin x$	$\frac{1}{2}$
1	$\cos x$	$\frac{\sqrt{3}}{2}$
2	$-\sin x$	$-\frac{1}{2}$
3	$-\cos x$	$-\frac{\sqrt{3}}{2}$
4	$\sin x$	$\frac{1}{2}$
$\vdots$	$\vdots$	$\vdots$

$$\sin x = f\left(\frac{\pi}{6}\right) + f'\left(\frac{\pi}{6}\right)\left(x-\frac{\pi}{6}\right) + \frac{f''\left(\frac{\pi}{6}\right)}{2!}\left(x-\frac{\pi}{6}\right)^2 + \frac{f^{(3)}\left(\frac{\pi}{6}\right)}{3!}\left(x-\frac{\pi}{6}\right)^3 + \frac{f^{(4)}\left(\frac{\pi}{6}\right)}{4!}\left(x-\frac{\pi}{6}\right)^4 + \cdots$$

$$= \frac{1}{2}\left[1 - \frac{1}{2!}\left(x-\frac{\pi}{6}\right)^2 + \frac{1}{4!}\left(x-\frac{\pi}{6}\right)^4 - \cdots\right] + \frac{\sqrt{3}}{2}\left[\left(x-\frac{\pi}{6}\right) - \frac{1}{3!}\left(x-\frac{\pi}{6}\right)^3 + \cdots\right]$$

$$= \frac{1}{2}\sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n)!}\left(x-\frac{\pi}{6}\right)^{2n} + \frac{\sqrt{3}}{2}\sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n+1)!}\left(x-\frac{\pi}{6}\right)^{2n+1}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

46.

n	$f^{(n)}(x)$	$f^{(n)}\left(\frac{\pi}{3}\right)$
0	$\cos x$	$\frac{1}{2}$
1	$-\sin x$	$\frac{\sqrt{3}}{2}$
2	$-\cos x$	$-\frac{1}{2}$
3	$\sin x$	$\frac{\sqrt{3}}{2}$
4	$\cos x$	$\frac{1}{2}$
$\vdots$	$\vdots$	$\vdots$

$$\cos x = f\left(\frac{\pi}{3}\right) + f'\left(\frac{\pi}{3}\right)\left(x-\frac{\pi}{3}\right) + \frac{f''\left(\frac{\pi}{3}\right)}{2!}\left(x-\frac{\pi}{3}\right)^2 + \frac{f^{(3)}\left(\frac{\pi}{3}\right)}{3!}\left(x-\frac{\pi}{3}\right)^3 + \frac{f^{(4)}\left(\frac{\pi}{3}\right)}{4!}\left(x-\frac{\pi}{3}\right)^4 + \cdots$$

$$= \frac{1}{2}\left[1 - \frac{1}{2!}\left(x-\frac{\pi}{3}\right)^2 + \frac{1}{4!}\left(x-\frac{\pi}{3}\right)^4 - \cdots\right] + \frac{\sqrt{3}}{2}\left[-\left(x-\frac{\pi}{3}\right) + \frac{1}{3!}\left(x-\frac{\pi}{3}\right)^3 - \cdots\right]$$

$$= \frac{1}{2}\sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n)!}\left(x-\frac{\pi}{3}\right)^{2n} + \frac{\sqrt{3}}{2}\sum_{n=0}^{\infty}(-1)^{n+1}\frac{1}{(2n+1)!}\left(x-\frac{\pi}{3}\right)^{2n+1}$$

47. $\dfrac{1}{1+x} = \dfrac{1}{1-(-x)} = \sum_{n=0}^{\infty}(-x)^n = \sum_{n=0}^{\infty}(-1)^n x^n$ for $|x|<1 \quad\Rightarrow\quad \dfrac{x^2}{1+x} = \sum_{n=0}^{\infty}(-1)^n x^{n+2}$ with $R=1$.

48. $\tan^{-1}x = \sum_{n=0}^{\infty}(-1)^n\dfrac{x^{2n+1}}{2n+1}$ with interval of convergence $[-1,1]$, so

$\tan^{-1}(x^2) = \sum_{n=0}^{\infty}(-1)^n\dfrac{(x^2)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty}(-1)^n\dfrac{x^{4n+2}}{2n+1}$, which converges when $x^2\in[-1,1] \quad\Leftrightarrow\quad x\in[-1,1]$.

Therefore, $R=1$.

49. $\displaystyle\int\frac{1}{4-x}\,dx = -\ln(4-x)+C$ and

$\displaystyle\int\frac{1}{4-x}\,dx = \frac{1}{4}\int\frac{1}{1-x/4}\,dx = \frac{1}{4}\int\sum_{n=0}^{\infty}\left(\frac{x}{4}\right)^n dx = \frac{1}{4}\int\sum_{n=0}^{\infty}\frac{x^n}{4^n}\,dx = \frac{1}{4}\sum_{n=0}^{\infty}\frac{x^{n+1}}{4^n(n+1)}+C$. So

$\ln(4-x) = -\dfrac{1}{4}\sum_{n=0}^{\infty}\dfrac{x^{n+1}}{4^n(n+1)}+C = -\sum_{n=0}^{\infty}\dfrac{x^{n+1}}{4^{n+1}(n+1)}+C = -\sum_{n=1}^{\infty}\dfrac{x^n}{n4^n}+C$. Putting $x=0$, we get $C=\ln 4$.

Thus, $f(x)=\ln(4-x)=\ln 4-\sum_{n=1}^{\infty}\dfrac{x^n}{n4^n}$. The series converges for $|x/4|<1 \quad\Leftrightarrow\quad |x|<4$, so $R=4$.

Another solution:

$$\ln(4-x) = \ln[4(1-x/4)] = \ln 4 + \ln(1-x/4) = \ln 4 + \ln[1+(-x/4)]$$

$$= \ln 4 + \sum_{n=1}^{\infty}(-1)^{n+1}\frac{(-x/4)^n}{n} \quad \text{[from Table 1 in Section 11.10]}$$

$$= \ln 4 + \sum_{n=1}^{\infty}(-1)^{2n+1}\frac{x^n}{n4^n} = \ln 4 - \sum_{n=1}^{\infty}\frac{x^n}{n4^n}.$$

50. $e^x = \sum_{n=0}^{\infty}\dfrac{x^n}{n!} \quad\Rightarrow\quad e^{2x} = \sum_{n=0}^{\infty}\dfrac{(2x)^n}{n!} \quad\Rightarrow\quad xe^{2x} = x\sum_{n=0}^{\infty}\dfrac{2^n x^n}{n!} = \sum_{n=0}^{\infty}\dfrac{2^n x^{n+1}}{n!}$, $R=\infty$

51. $\sin x = \sum_{n=0}^{\infty}\dfrac{(-1)^n x^{2n+1}}{(2n+1)!} \quad\Rightarrow\quad \sin(x^4) = \sum_{n=0}^{\infty}\dfrac{(-1)^n (x^4)^{2n+1}}{(2n+1)!} = \sum_{n=0}^{\infty}\dfrac{(-1)^n x^{8n+4}}{(2n+1)!}$ for all x, so the radius of convergence is ∞.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad \Rightarrow \quad 10^x = e^{(\ln 10)x} = \sum_{n=0}^{\infty} \frac{[(\ln 10)x]^n}{n!} = \sum_{n=0}^{\infty} \frac{(\ln 10)^n x^n}{n!}, \ R = \infty$

53. $f(x) = \dfrac{1}{\sqrt[4]{16 - x}} = \dfrac{1}{\sqrt[4]{16(1 - x/16)}} = \dfrac{1}{\sqrt[4]{16}\,(1 - \frac{1}{16}x)^{1/4}} = \frac{1}{2}(1 - \frac{1}{16}x)^{-1/4}$

$$= \frac{1}{2}\left[1 + \left(-\frac{1}{4}\right)\left(-\frac{x}{16}\right) + \frac{\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)}{2!}\left(-\frac{x}{16}\right)^2 + \frac{\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)\left(-\frac{9}{4}\right)}{3!}\left(-\frac{x}{16}\right)^3 + \cdots\right]$$

$$= \frac{1}{2} + \sum_{n=1}^{\infty} \frac{1 \cdot 5 \cdot 9 \cdot \cdots \cdot (4n-3)}{2 \cdot 4^n \cdot n! \cdot 16^n} x^n = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{1 \cdot 5 \cdot 9 \cdot \cdots \cdot (4n-3)}{2^{6n+1}\, n!} x^n$$

for $\left|-\dfrac{x}{16}\right| < 1 \quad \Leftrightarrow \quad |x| < 16$, so $R = 16$.

54. $(1 - 3x)^{-5} = \sum_{n=0}^{\infty} \binom{-5}{n} (-3x)^n = 1 + (-5)(-3x) + \dfrac{(-5)(-6)}{2!}(-3x)^2 + \dfrac{(-5)(-6)(-7)}{3!}(-3x)^3 + \cdots$

$$= 1 + \sum_{n=1}^{\infty} \frac{5 \cdot 6 \cdot 7 \cdot \cdots \cdot (n+4) \cdot 3^n x^n}{n!} \quad \text{for } |-3x| < 1 \quad \Leftrightarrow \quad |x| < \tfrac{1}{3}, \text{ so } R = \tfrac{1}{3}.$$

55. $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$, so $\dfrac{e^x}{x} = \dfrac{1}{x}\sum_{n=0}^{\infty} \dfrac{x^n}{n!} = \sum_{n=0}^{\infty} \dfrac{x^{n-1}}{n!} = x^{-1} + \sum_{n=1}^{\infty} \dfrac{x^{n-1}}{n!} = \dfrac{1}{x} + \sum_{n=1}^{\infty} \dfrac{x^{n-1}}{n!}$ and

$$\int \frac{e^x}{x}\,dx = C + \ln|x| + \sum_{n=1}^{\infty} \frac{x^n}{n \cdot n!}.$$

56. $(1 + x^4)^{1/2} = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} (x^4)^n = 1 + (\tfrac{1}{2})x^4 + \dfrac{(\frac{1}{2})(-\frac{1}{2})}{2!}(x^4)^2 + \dfrac{(\frac{1}{2})(-\frac{1}{2})(-\frac{3}{2})}{3!}(x^4)^3 + \cdots$

$$= 1 + \tfrac{1}{2}x^4 - \tfrac{1}{8}x^8 + \tfrac{1}{16}x^{12} - \cdots$$

so $\int_0^1 (1 + x^4)^{1/2}\,dx = \left[x + \frac{1}{10}x^5 - \frac{1}{72}x^9 + \frac{1}{208}x^{13} - \cdots\right]_0^1 = 1 + \frac{1}{10} - \frac{1}{72} + \frac{1}{208} - \cdots$.

This is an alternating series, so by the Alternating Series Test, the error in the approximation

$\int_0^1 (1 + x^4)^{1/2}\,dx \approx 1 + \frac{1}{10} - \frac{1}{72} \approx 1.086$ is less than $\frac{1}{208}$, sufficient for the desired accuracy.

Thus, correct to two decimal places, $\int_0^1 (1 + x^4)^{1/2}\,dx \approx 1.09$.

57. (a)

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$x^{1/2}$	1
1	$\frac{1}{2}x^{-1/2}$	$\frac{1}{2}$
2	$-\frac{1}{4}x^{-3/2}$	$-\frac{1}{4}$
3	$\frac{3}{8}x^{-5/2}$	$\frac{3}{8}$
4	$-\frac{15}{16}x^{-7/2}$	$-\frac{15}{16}$
⋮	⋮	⋮

$$\sqrt{x} \approx T_3(x) = 1 + \frac{1/2}{1!}(x-1) - \frac{1/4}{2!}(x-1)^2 + \frac{3/8}{3!}(x-1)^3$$

$$= 1 + \tfrac{1}{2}(x-1) - \tfrac{1}{8}(x-1)^2 + \tfrac{1}{16}(x-1)^3$$

(b)



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c) $|R_3(x)| \le \dfrac{M}{4!}|x-1|^4$, where $\left|f^{(4)}(x)\right| \le M$ with $f^{(4)}(x) = -\frac{15}{16}x^{-7/2}$. Now $0.9 \le x \le 1.1 \quad \Rightarrow$

$-0.1 \le x-1 \le 0.1 \quad \Rightarrow \quad (x-1)^4 \le (0.1)^4$, and letting $x = 0.9$ gives $M = \dfrac{15}{16(0.9)^{7/2}}$, so

$$|R_3(x)| \le \frac{15}{16(0.9)^{7/2}\,4!}(0.1)^4 \approx 0.000\,005\,648 \approx 0.000\,006 = 6 \times 10^{-6}.$$

(d)



From the graph of $|R_3(x)| = |\sqrt{x} - T_3(x)|$, it appears that the error is less than 5×10^{-6} on $[0.9, 1.1]$.

58. (a) $\sec x \approx T_2(x) = 1 + \frac{1}{2}x^2$

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sec x$	1
1	$\sec x \tan x$	0
2	$\sec x \tan^2 x + \sec^3 x$	1
3	$\sec x \tan^3 x + 5\sec^3 x \tan x$	0
⋮	⋮	⋮

(b)



(c) $|R_2(x)| \le \dfrac{M}{3!}|x|^3$, where $\left|f^{(3)}(x)\right| \le M$ with $f^{(3)}(x) = \sec x \tan^3 x + 5\sec^3 x \tan x$.

Now $0 \le x \le \frac{\pi}{6} \quad \Rightarrow \quad x^3 \le \left(\frac{\pi}{6}\right)^3$, and letting $x = \frac{\pi}{6}$ gives $M = \frac{14}{3}$, so $|R_2(x)| \le \dfrac{14}{3 \cdot 6}\left(\dfrac{\pi}{6}\right)^3 \approx 0.111648$.

(d)



From the graph of $|R_2(x)| = |\sec x - T_2(x)|$, it appears that the error is less than 0.02 on $\left[0, \frac{\pi}{6}\right]$.

59. $\sin x = \displaystyle\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$, so $\sin x - x = -\dfrac{x^3}{3!} + \dfrac{x^5}{5!} - \dfrac{x^7}{7!} + \cdots$ and

$\dfrac{\sin x - x}{x^3} = -\dfrac{1}{3!} + \dfrac{x^2}{5!} - \dfrac{x^4}{7!} + \cdots$. Thus, $\displaystyle\lim_{x \to 0} \frac{\sin x - x}{x^3} = \lim_{x \to 0}\left(-\frac{1}{6} + \frac{x^2}{120} - \frac{x^4}{5040} + \cdots\right) = -\frac{1}{6}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. (a) $F = \dfrac{mgR^2}{(R+h)^2} = \dfrac{mg}{(1+h/R)^2} = mg \sum_{n=0}^{\infty} \binom{-2}{n} \left(\dfrac{h}{R}\right)^n$ [binomial series]

(b) We expand $F = mg\left[1 - 2\,(h/R) + 3\,(h/R)^2 - \cdots\right]$.

This is an alternating series, so by the Alternating Series Estimation Theorem, the error in the approximation $F = mg$ is less than $2mgh/R$, so for accuracy within 1% we want

$$\left|\frac{2mgh/R}{mgR^2/(R+h)^2}\right| < 0.01 \quad \Leftrightarrow \quad \frac{2h(R+h)^2}{R^3} < 0.01.$$



This inequality would be difficult to solve for h, so we substitute $R = 6{,}400$ km and plot both sides of the inequality. It appears that the approximation is accurate to within 1% for $h < 31$ km.

61. $f(x) = \sum_{n=0}^{\infty} c_n x^n \quad \Rightarrow \quad f(-x) = \sum_{n=0}^{\infty} c_n(-x)^n = \sum_{n=0}^{\infty} (-1)^n c_n x^n$

(a) If f is an odd function, then $f(-x) = -f(x) \quad \Rightarrow \quad \sum_{n=0}^{\infty} (-1)^n c_n x^n = \sum_{n=0}^{\infty} -c_n x^n$. The coefficients of any power series are uniquely determined (by Theorem 11.10.5), so $(-1)^n c_n = -c_n$.

If n is even, then $(-1)^n = 1$, so $c_n = -c_n \quad \Rightarrow \quad 2c_n = 0 \quad \Rightarrow \quad c_n = 0$. Thus, all even coefficients are 0, that is, $c_0 = c_2 = c_4 = \cdots = 0$.

(b) If f is even, then $f(-x) = f(x) \quad \Rightarrow \quad \sum_{n=0}^{\infty} (-1)^n c_n x^n = \sum_{n=0}^{\infty} c_n x^n \quad \Rightarrow \quad (-1)^n c_n = c_n$.

If n is odd, then $(-1)^n = -1$, so $-c_n = c_n \quad \Rightarrow \quad 2c_n = 0 \quad \Rightarrow \quad c_n = 0$. Thus, all odd coefficients are 0, that is, $c_1 = c_3 = c_5 = \cdots = 0$.

62. $e^x = \sum_{n=0}^{\infty} \dfrac{x^n}{n!} \quad \Rightarrow \quad f(x) = e^{x^2} = \sum_{n=0}^{\infty} \dfrac{(x^2)^n}{n!} = \sum_{n=0}^{\infty} \dfrac{x^{2n}}{n!} = \sum_{n=0}^{\infty} \dfrac{1}{n!} x^{2n}$. By Theorem 11.10.6 with $a = 0$, we also have $f(x) = \sum_{k=0}^{\infty} \dfrac{f^{(k)}(0)}{k!} x^k$. Comparing coefficients for $k = 2n$, we have $\dfrac{f^{(2n)}(0)}{(2n)!} = \dfrac{1}{n!} \quad \Rightarrow \quad f^{(2n)}(0) = \dfrac{(2n)!}{n!}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

PROBLEMS PLUS

1. It would be far too much work to compute 15 derivatives of f. The key idea is to remember that $f^{(n)}(0)$ occurs in the coefficient of x^n in the Maclaurin series of f. We start with the Maclaurin series for sin: $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots$. Then $\sin(x^3) = x^3 - \frac{x^9}{3!} + \frac{x^{15}}{5!} - \cdots$, and so the coefficient of x^{15} is $\frac{f^{(15)}(0)}{15!} = \frac{1}{5!}$. Therefore,

$$f^{(15)}(0) = \frac{15!}{5!} = 6 \cdot 7 \cdot 8 \cdot 9 \cdot 10 \cdot 11 \cdot 12 \cdot 13 \cdot 14 \cdot 15 = 10{,}897{,}286{,}400.$$

2. We use the problem-solving strategy of taking cases:

Case (i): If $|x| < 1$, then $0 \le x^2 < 1$, so $\lim_{n\to\infty} x^{2n} = 0$ [see Example 11.1.11]

and $f(x) = \lim_{n\to\infty} \frac{x^{2n} - 1}{x^{2n} + 1} = \frac{0 - 1}{0 + 1} = -1$.

Case (ii): If $|x| = 1$, that is, $x = \pm 1$, then $x^2 = 1$, so $f(x) = \lim_{n\to\infty} \frac{x^{2n} - 1}{x^{2n} + 1} = \lim_{n\to\infty} \frac{1 - 1}{1 + 1} = 0$.

Case (iii): If $|x| > 1$, then $x^2 > 1$, so $\lim_{n\to\infty} x^{2n} = \infty$ and $f(x) = \lim_{n\to\infty} \frac{x^{2n} - 1}{x^{2n} + 1} = \lim_{n\to\infty} \frac{1 - (1/x^{2n})}{1 - (1/x^{2n})} = \frac{1 - 0}{1 + 0} = 1$.

Thus, $f(x) = \begin{cases} 1 & \text{if } x < -1 \\ 0 & \text{if } x = -1 \\ -1 & \text{if } -1 < x < 1 \\ 0 & \text{if } x = 1 \\ 1 & \text{if } x > 1 \end{cases}$



The graph shows that f is continuous everywhere except at $x = \pm 1$.

3. (a) From Formula 14a in Appendix D, with $x = y = \theta$, we get $\tan 2\theta = \frac{2\tan\theta}{1 - \tan^2\theta}$, so $\cot 2\theta = \frac{1 - \tan^2\theta}{2\tan\theta} \Rightarrow$

$2\cot 2\theta = \frac{1 - \tan^2\theta}{\tan\theta} = \cot\theta - \tan\theta$. Replacing θ by $\frac{1}{2}x$, we get $2\cot x = \cot\frac{1}{2}x - \tan\frac{1}{2}x$, or

$\tan\frac{1}{2}x = \cot\frac{1}{2}x - 2\cot x$.

(b) From part (a) with $\frac{x}{2^{n-1}}$ in place of x, $\tan\frac{x}{2^n} = \cot\frac{x}{2^n} - 2\cot\frac{x}{2^{n-1}}$, so the nth partial sum of $\sum_{n=1}^{\infty} \frac{1}{2^n}\tan\frac{x}{2^n}$ is

$$\begin{aligned} s_n &= \frac{\tan(x/2)}{2} + \frac{\tan(x/4)}{4} + \frac{\tan(x/8)}{8} + \cdots + \frac{\tan(x/2^n)}{2^n} \\ &= \left[\frac{\cot(x/2)}{2} - \cot x\right] + \left[\frac{\cot(x/4)}{4} - \frac{\cot(x/2)}{2}\right] + \left[\frac{\cot(x/8)}{8} - \frac{\cot(x/4)}{4}\right] + \cdots \\ &\quad + \left[\frac{\cot(x/2^n)}{2^n} - \frac{\cot(x/2^{n-1})}{2^{n-1}}\right] = -\cot x + \frac{\cot(x/2^n)}{2^n} \quad \text{[telescoping sum]} \end{aligned}$$

Now $\frac{\cot(x/2^n)}{2^n} = \frac{\cos(x/2^n)}{2^n \sin(x/2^n)} = \frac{\cos(x/2^n)}{x} \cdot \frac{x/2^n}{\sin(x/2^n)} \to \frac{1}{x} \cdot 1 = \frac{1}{x}$ as $n \to \infty$ since $x/2^n \to 0$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

for $x \neq 0$. Therefore, if $x \neq 0$ and $x \neq k\pi$ where k is any integer, then

$$\sum_{n=1}^{\infty} \frac{1}{2^n} \tan \frac{x}{2^n} = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \left(-\cot x + \frac{1}{2^n} \cot \frac{x}{2^n} \right) = -\cot x + \frac{1}{x}$$

If $x = 0$, then all terms in the series are 0, so the sum is 0.

4. $|AP_2|^2 = 2$, $|AP_3|^2 = 2 + 2^2$, $|AP_4|^2 = 2 + 2^2 + (2^2)^2$, $|AP_5|^2 = 2 + 2^2 + (2^2)^2 + (2^3)^2, \ldots,$

$|AP_n|^2 = 2 + 2^2 + (2^2)^2 + \cdots + (2^{n-2})^2$ [for $n \geq 3$] $= 2 + (4 + 4^2 + 4^3 + \cdots + 4^{n-2})$

$= 2 + \dfrac{4(4^{n-2} - 1)}{4 - 1}$ [finite geometric sum with $a = 4$, $r = 4$] $= \dfrac{6}{3} + \dfrac{4^{n-1} - 4}{3} = \dfrac{2}{3} + \dfrac{4^{n-1}}{3}$

So $\tan \angle P_n A P_{n+1} = \dfrac{|P_n P_{n+1}|}{|AP_n|} = \dfrac{2^{n-1}}{\sqrt{\frac{2}{3} + \frac{4^{n-1}}{3}}} = \dfrac{\sqrt{4^{n-1}}}{\sqrt{\frac{2}{3} + \frac{4^{n-1}}{3}}} = \dfrac{1}{\sqrt{\frac{2}{3 \cdot 4^{n-1}} - \frac{1}{3}}} \to \sqrt{3}$ as $n \to \infty$.

Thus, $\angle P_n A P_{n+1} \to \frac{\pi}{3}$ as $n \to \infty$.

5. (a) At each stage, each side is replaced by four shorter sides, each of length $\frac{1}{3}$ of the side length at the preceding stage. Writing s_0 and ℓ_0 for the number of sides and the length of the side of the initial triangle, we generate the table at right. In general, we have $s_n = 3 \cdot 4^n$ and $\ell_n = \left(\frac{1}{3}\right)^n$, so the length of the perimeter at the nth stage of construction is $p_n = s_n \ell_n = 3 \cdot 4^n \cdot \left(\frac{1}{3}\right)^n = 3 \cdot \left(\frac{4}{3}\right)^n$.

$s_0 = 3$	$\ell_0 = 1$
$s_1 = 3 \cdot 4$	$\ell_1 = 1/3$
$s_2 = 3 \cdot 4^2$	$\ell_2 = 1/3^2$
$s_3 = 3 \cdot 4^3$	$\ell_3 = 1/3^3$
⋮	⋮

(b) $p_n = \dfrac{4^n}{3^{n-1}} = 4\left(\dfrac{4}{3}\right)^{n-1}$. Since $\frac{4}{3} > 1$, $p_n \to \infty$ as $n \to \infty$.

(c) The area of each of the small triangles added at a given stage is one-ninth of the area of the triangle added at the preceding stage. Let a be the area of the original triangle. Then the area a_n of each of the small triangles added at stage n is $a_n = a \cdot \dfrac{1}{9^n} = \dfrac{a}{9^n}$. Since a small triangle is added to each side at every stage, it follows that the total area A_n added to the figure at the nth stage is $A_n = s_{n-1} \cdot a_n = 3 \cdot 4^{n-1} \cdot \dfrac{a}{9^n} = a \cdot \dfrac{4^{n-1}}{3^{2n-1}}$. Then the total area enclosed by the snowflake curve is $A = a + A_1 + A_2 + A_3 + \cdots = a + a \cdot \dfrac{1}{3} + a \cdot \dfrac{4}{3^3} + a \cdot \dfrac{4^2}{3^5} + a \cdot \dfrac{4^3}{3^7} + \cdots$. After the first term, this is a geometric series with common ratio $\dfrac{4}{9}$, so $A = a + \dfrac{a/3}{1 - \frac{4}{9}} = a + \dfrac{a}{3} \cdot \dfrac{9}{5} = \dfrac{8a}{5}$. But the area of the original equilateral triangle with side 1 is $a = \dfrac{1}{2} \cdot 1 \cdot \sin \dfrac{\pi}{3} = \dfrac{\sqrt{3}}{4}$. So the area enclosed by the snowflake curve is $\dfrac{8}{5} \cdot \dfrac{\sqrt{3}}{4} = \dfrac{2\sqrt{3}}{5}$.

6. Let the series $S = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8} + \frac{1}{9} + \frac{1}{12} + \cdots$. Then every term in S is of the form $\dfrac{1}{2^m 3^n}$, $m, n \geq 0$, and furthermore each term occurs only once. So we can write

$$S = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{2^m 3^n} = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{2^m} \frac{1}{3^n} = \sum_{m=0}^{\infty} \frac{1}{2^m} \sum_{n=0}^{\infty} \frac{1}{3^n} = \frac{1}{1 - \frac{1}{2}} \cdot \frac{1}{1 - \frac{1}{3}} = 2 \cdot \frac{3}{2} = 3$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. (a) Let $a = \arctan x$ and $b = \arctan y$. Then, from Formula 14b in Appendix D,

$$\tan(a-b) = \frac{\tan a - \tan b}{1 + \tan a \tan b} = \frac{\tan(\arctan x) - \tan(\arctan y)}{1 + \tan(\arctan x)\tan(\arctan y)} = \frac{x-y}{1+xy}$$

Now $\arctan x - \arctan y = a - b = \arctan(\tan(a-b)) = \arctan \dfrac{x-y}{1+xy}$ since $-\frac{\pi}{2} < a - b < \frac{\pi}{2}$.

(b) From part (a) we have

$$\arctan \tfrac{120}{119} - \arctan \tfrac{1}{239} = \arctan \frac{\frac{120}{119} - \frac{1}{239}}{1 + \frac{120}{119}\cdot\frac{1}{239}} = \arctan \frac{\frac{28{,}561}{28{,}441}}{\frac{28{,}561}{28{,}441}} = \arctan 1 = \tfrac{\pi}{4}$$

(c) Replacing y by $-y$ in the formula of part (a), we get $\arctan x + \arctan y = \arctan \dfrac{x+y}{1-xy}$. So

$$4\arctan \tfrac{1}{5} = 2\big(\arctan \tfrac{1}{5} + \arctan \tfrac{1}{5}\big) = 2\arctan \frac{\frac{1}{5} + \frac{1}{5}}{1 - \frac{1}{5}\cdot\frac{1}{5}} = 2\arctan \tfrac{5}{12} = \arctan \tfrac{5}{12} + \arctan \tfrac{5}{12}$$

$$= \arctan \frac{\frac{5}{12} + \frac{5}{12}}{1 - \frac{5}{12}\cdot\frac{5}{12}} = \arctan \tfrac{120}{119}$$

Thus, from part (b), we have $4\arctan \frac{1}{5} - \arctan \frac{1}{239} = \arctan \frac{120}{119} - \arctan \frac{1}{239} = \frac{\pi}{4}$.

(d) From Example 11.9.7 we have $\arctan x = x - \dfrac{x^3}{3} + \dfrac{x^5}{5} - \dfrac{x^7}{7} + \dfrac{x^9}{9} - \dfrac{x^{11}}{11} + \cdots$, so

$$\arctan \frac{1}{5} = \frac{1}{5} - \frac{1}{3\cdot 5^3} + \frac{1}{5\cdot 5^5} - \frac{1}{7\cdot 5^7} + \frac{1}{9\cdot 5^9} - \frac{1}{11\cdot 5^{11}} + \cdots$$

This is an alternating series and the size of the terms decreases to 0, so by the Alternating Series Estimation Theorem, the sum lies between s_5 and s_6, that is, $0.197395560 < \arctan \frac{1}{5} < 0.197395562$.

(e) From the series in part (d) we get $\arctan \dfrac{1}{239} = \dfrac{1}{239} - \dfrac{1}{3\cdot 239^3} + \dfrac{1}{5\cdot 239^5} - \cdots$. The third term is less than 2.6×10^{-13}, so by the Alternating Series Estimation Theorem, we have, to nine decimal places, $\arctan \frac{1}{239} \approx s_2 \approx 0.004184076$. Thus, $0.004184075 < \arctan \frac{1}{239} < 0.004184077$.

(f) From part (c) we have $\pi = 16\arctan \frac{1}{5} - 4\arctan \frac{1}{239}$, so from parts (d) and (e) we have

$16(0.197395560) - 4(0.004184077) < \pi < 16(0.197395562) - 4(0.004184075) \;\Rightarrow$

$3.141592652 < \pi < 3.141592692$. So, to 7 decimal places, $\pi \approx 3.1415927$.

8. (a) Let $a = \operatorname{arccot} x$ and $b = \operatorname{arccot} y$ where $0 < a - b < \pi$. Then

$$\cot(a-b) = \frac{1}{\tan(a-b)} = \frac{1 + \tan a \tan b}{\tan a - \tan b} = \frac{\frac{1}{\cot a}\cdot\frac{1}{\cot b} + 1}{\frac{1}{\cot a} - \frac{1}{\cot b}} \cdot \frac{\cot a \cot b}{\cot a \cot b}$$

$$= \frac{1 + \cot a \cot b}{\cot b - \cot a} = \frac{1 + \cot(\operatorname{arccot} x)\cot(\operatorname{arccot} y)}{\cot(\operatorname{arccot} y) - \cot(\operatorname{arccot} x)} = \frac{1+xy}{y-x}$$

Now $\operatorname{arccot} x - \operatorname{arccot} y = a - b = \operatorname{arccot}(\cot(a-b)) = \operatorname{arccot} \dfrac{1+xy}{y-x}$ since $0 < a - b < \pi$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) From part (a), we want $\operatorname{arccot}(n^2+n+1)$ to equal $\operatorname{arccot}\dfrac{1+xy}{y-x}$. Note that $1+xy=n^2+n+1 \quad \Leftrightarrow$

$xy=n^2+n=(n+1)n$, so if we let $x=n+1$ and $y=n$, then $y-x=1$. Therefore,

$$\operatorname{arccot}(n^2+n+1)=\operatorname{arccot}(1+n(n+1))=\operatorname{arccot}\frac{1+n(n+1)}{(n+1)-n}=\operatorname{arccot}n-\operatorname{arccot}(n+1)$$

Thus, we have a telescoping series with nth partial sum

$$s_n=[\operatorname{arccot}0-\operatorname{arccot}1]+[\operatorname{arccot}1-\operatorname{arccot}2]+\cdots+[\operatorname{arccot}n-\operatorname{arccot}(n+1)]=\operatorname{arccot}0-\operatorname{arccot}(n+1).$$

Thus, $\displaystyle\sum_{n=0}^{\infty}\operatorname{arccot}(n^2+n+1)=\lim_{n\to\infty}s_n=\lim_{n\to\infty}[\operatorname{arccot}0-\operatorname{arccot}(n+1)]=\frac{\pi}{2}-0=\frac{\pi}{2}$.

9. We want $\arctan\left(\dfrac{2}{n^2}\right)$ to equal $\arctan\dfrac{x-y}{1+xy}$. Note that $1+xy=n^2 \quad \Leftrightarrow \quad xy=n^2-1=(n+1)(n-1)$, so if we let $x=n+1$ and $y=n-1$, then $x-y=2$ and $xy\neq -1$. Thus, from Problem 7(a),

$\arctan\left(\dfrac{2}{n^2}\right)=\arctan\dfrac{x-y}{1+xy}=\arctan x-\arctan y=\arctan(n+1)-\arctan(n-1)$. Therefore,

$$\begin{aligned}\sum_{n=1}^{k}\arctan\left(\frac{2}{n^2}\right)&=\sum_{n=1}^{k}[\arctan(n+1)-\arctan(n-1)]\\&=\sum_{n=1}^{k}[\arctan(n+1)-\arctan n+\arctan n-\arctan(n-1)]\\&=\sum_{n=1}^{k}[\arctan(n+1)-\arctan n]+\sum_{n=1}^{k}[\arctan n-\arctan(n-1)]\\&=[\arctan(k+1)-\arctan 1]+[\arctan k-\arctan 0]\quad\text{[since both sums are telescoping]}\\&=\arctan(k+1)-\tfrac{\pi}{4}+\arctan k-0\end{aligned}$$

Now $\displaystyle\sum_{n=1}^{k}\arctan\left(\frac{2}{n^2}\right)=\lim_{k\to 0}\sum_{n=1}^{k}\arctan\left(\frac{2}{n^2}\right)=\lim_{k\to\infty}\left[\arctan(k+1)-\frac{\pi}{4}+\arctan k\right]=\frac{\pi}{2}-\frac{\pi}{4}+\frac{\pi}{2}=\frac{3\pi}{4}$.

Note: For all $n\geq 1$, $0\leq\arctan(n-1)<\arctan(n+1)<\frac{\pi}{2}$, so $-\frac{\pi}{2}<\arctan(n+1)-\arctan(n-1)<\frac{\pi}{2}$, and the identity in Problem 7(a) holds.

10. Let's first try the case $k=1$: $a_0+a_1=0 \quad \Rightarrow \quad a_1=-a_0 \quad \Rightarrow$

$$\begin{aligned}\lim_{n\to\infty}\left(a_0\sqrt{n}+a_1\sqrt{n+1}\right)&=\lim_{n\to\infty}\left(a_0\sqrt{n}-a_0\sqrt{n+1}\right)=a_0\lim_{n\to\infty}\left(\sqrt{n}-\sqrt{n+1}\right)\frac{\sqrt{n}+\sqrt{n+1}}{\sqrt{n}+\sqrt{n+1}}\\&=a_0\lim_{n\to\infty}\frac{-1}{\sqrt{n}+\sqrt{n+1}}=0\end{aligned}$$

In general we have $a_0+a_1+\cdots+a_k=0 \quad \Rightarrow \quad a_k=-a_0-a_1-\cdots-a_{k-1} \quad \Rightarrow$

$$\begin{aligned}&\lim_{n\to\infty}\left(a_0\sqrt{n}+a_1\sqrt{n+1}+a_2\sqrt{n+2}+\cdots+a_k\sqrt{n+k}\right)\\&=\lim_{n\to\infty}\left(a_0\sqrt{n}+a_1\sqrt{n+1}+\cdots+a_{k-1}\sqrt{n+k-1}-a_0\sqrt{n+k}-a_1\sqrt{n+k}-\cdots-a_{k-1}\sqrt{n+k}\right)\\&=a_0\lim_{n\to\infty}\left(\sqrt{n}-\sqrt{n+k}\right)+a_1\lim_{n\to\infty}\left(\sqrt{n+1}-\sqrt{n+k}\right)+\cdots+a_{k-1}\lim_{n\to\infty}\left(\sqrt{n+k-1}-\sqrt{n+k}\right)\end{aligned}$$

Each of these limits is 0 by the same type of simplification as in the case $k=1$. So we have

$$\lim_{n\to\infty}\left(a_0\sqrt{n}+a_1\sqrt{n+1}+a_2\sqrt{n+2}+\cdots+a_k\sqrt{n+k}\right)=a_0(0)+a_1(0)+\cdots+a_{k-1}(0)=0$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. We start with the geometric series $\sum_{n=0}^{\infty} x^n = \dfrac{1}{1-x}$, $|x| < 1$, and differentiate:

$$\sum_{n=1}^{\infty} nx^{n-1} = \frac{d}{dx}\left(\sum_{n=0}^{\infty} x^n\right) = \frac{d}{dx}\left(\frac{1}{1-x}\right) = \frac{1}{(1-x)^2} \text{ for } |x| < 1 \quad \Rightarrow \quad \sum_{n=1}^{\infty} nx^n = x\sum_{n=1}^{\infty} nx^{n-1} = \frac{x}{(1-x)^2}$$

for $|x| < 1$. Differentiate again:

$$\sum_{n=1}^{\infty} n^2x^{n-1} = \frac{d}{dx}\frac{x}{(1-x)^2} = \frac{(1-x)^2 - x\cdot 2(1-x)(-1)}{(1-x)^4} = \frac{x+1}{(1-x)^3} \quad \Rightarrow \quad \sum_{n=1}^{\infty} n^2\,x^n = \frac{x^2+x}{(1-x)^3} \quad \Rightarrow$$

$$\sum_{n=1}^{\infty} n^3x^{n-1} = \frac{d}{dx}\frac{x^2+x}{(1-x)^3} = \frac{(1-x)^3(2x+1) - (x^2+x)3(1-x)^2(-1)}{(1-x)^6} = \frac{x^2+4x+1}{(1-x)^4} \quad \Rightarrow$$

$\sum_{n=1}^{\infty} n^3x^n = \dfrac{x^3+4x^2+x}{(1-x)^4}$, $|x| < 1$. The radius of convergence is 1 because that is the radius of convergence for the geometric series we started with. If $x = \pm 1$, the series is $\sum n^3(\pm 1)^n$, which diverges by the Test For Divergence, so the interval of convergence is $(-1, 1)$.

12. Place the y-axis as shown and let the length of each book be L. We want to show that the center of mass of the system of n books lies above the table, that is, $\overline{x} < L$. The x-coordinates of the centers of mass of the books are $x_1 = \dfrac{L}{2}$, $x_2 = \dfrac{L}{2(n-1)} + \dfrac{L}{2}$, $x_3 = \dfrac{L}{2(n-1)} + \dfrac{L}{2(n-2)} + \dfrac{L}{2}$, and so on.



Each book has the same mass m, so if there are n books, then

$$\begin{aligned}
\overline{x} &= \frac{mx_1 + mx_2 + \cdots + mx_n}{mn} = \frac{x_1 + x_2 + \cdots + x_n}{n} \\
&= \frac{1}{n}\left[\frac{L}{2} + \left(\frac{L}{2(n-1)} + \frac{L}{2}\right) + \left(\frac{L}{2(n-1)} + \frac{L}{2(n-2)} + \frac{L}{2}\right) + \cdots \right. \\
&\qquad \left. + \left(\frac{L}{2(n-1)} + \frac{L}{2(n-2)} + \cdots + \frac{L}{4} + \frac{L}{2} + \frac{L}{2}\right)\right] \\
&= \frac{L}{n}\left[\frac{n-1}{2(n-1)} + \frac{n-2}{2(n-2)} + \cdots + \frac{2}{4} + \frac{1}{2} + \frac{n}{2}\right] = \frac{L}{n}\left[(n-1)\frac{1}{2} + \frac{n}{2}\right] = \frac{2n-1}{2n}L < L
\end{aligned}$$

This shows that, no matter how many books are added according to the given scheme, the center of mass lies above the table. It remains to observe that the series $\frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8} + \cdots = \frac{1}{2}\sum(1/n)$ is divergent (harmonic series), so we can make the top book extend as far as we like beyond the edge of the table if we add enough books.

13. $$\begin{aligned}
\ln\left(1 - \frac{1}{n^2}\right) &= \ln\left(\frac{n^2-1}{n^2}\right) = \ln\frac{(n+1)(n-1)}{n^2} = \ln[(n+1)(n-1)] - \ln n^2 \\
&= \ln(n+1) + \ln(n-1) - 2\ln n = \ln(n-1) - \ln n - \ln n + \ln(n+1) \\
&= \ln\frac{n-1}{n} - [\ln n - \ln(n+1)] = \ln\frac{n-1}{n} - \ln\frac{n}{n+1}.
\end{aligned}$$

Let $s_k = \sum_{n=2}^{k} \ln\left(1 - \dfrac{1}{n^2}\right) = \sum_{n=2}^{k}\left(\ln\dfrac{n-1}{n} - \ln\dfrac{n}{n+1}\right)$ for $k \geq 2$. Then

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$s_k = \left(\ln\frac{1}{2} - \ln\frac{2}{3}\right) + \left(\ln\frac{2}{3} - \ln\frac{3}{4}\right) + \cdots + \left(\ln\frac{k-1}{k} - \ln\frac{k}{k+1}\right) = \ln\frac{1}{2} - \ln\frac{k}{k+1}$, so

$\sum_{n=2}^{\infty} \ln\left(1 - \frac{1}{n^2}\right) = \lim_{k\to\infty} s_k = \lim_{k\to\infty}\left(\ln\frac{1}{2} - \ln\frac{k}{k+1}\right) = \ln\frac{1}{2} - \ln 1 = \ln 1 - \ln 2 - \ln 1 = -\ln 2$ (or $\ln\frac{1}{2}$).

14. First notice that both series are absolutely convergent (p-series with $p > 1$.) Let the given expression be called x. Then

$$x = \frac{1 + \frac{1}{2^p} + \frac{1}{3^p} + \frac{1}{4^p} + \cdots}{1 \quad \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots} = \frac{1 + \left(2\cdot\frac{1}{2^p} - \frac{1}{2^p}\right) + \frac{1}{3^p} + \left(2\cdot\frac{1}{4^p} - \frac{1}{4^p}\right) + \cdots}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots}$$

$$= \frac{\left(1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots\right) + \left(2\cdot\frac{1}{2^p} + 2\cdot\frac{1}{4^p} + 2\cdot\frac{1}{6^p} + \cdots\right)}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots}$$

$$= 1 + \frac{2\left(\frac{1}{2^p} + \frac{1}{4^p} + \frac{1}{6^p} + \frac{1}{8^p} + \cdots\right)}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots} = 1 + \frac{\frac{1}{2^{p-1}}\left(1 + \frac{1}{2^p} + \frac{1}{3^p} + \frac{1}{4^p} + \cdots\right)}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots} = 1 + 2^{1-p}x$$

Therefore, $x = 1 + 2^{1-p}x \;\Leftrightarrow\; x - 2^{1-p}x = 1 \;\Leftrightarrow\; x(1 - 2^{1-p}) = 1 \;\Leftrightarrow\; x = \frac{1}{1-2^{1-p}}$.

15. If L is the length of a side of the equilateral triangle, then the area is $A = \frac{1}{2}L\cdot\frac{\sqrt{3}}{2}L = \frac{\sqrt{3}}{4}L^2$ and so $L^2 = \frac{4}{\sqrt{3}}A$.

Let r be the radius of one of the circles. When there are n rows of circles, the figure shows that

$L = \sqrt{3}r + r + (n-2)(2r) + r + \sqrt{3}r = r\left(2n - 2 + 2\sqrt{3}\right)$, so $r = \dfrac{L}{2\left(n + \sqrt{3} - 1\right)}$.

The number of circles is $1 + 2 + \cdots + n = \dfrac{n(n+1)}{2}$, and so the total area of the circles is

$$A_n = \frac{n(n+1)}{2}\pi r^2 = \frac{n(n+1)}{2}\,\pi\,\frac{L^2}{4\left(n+\sqrt{3}-1\right)^2}$$

$$= \frac{n(n+1)}{2}\,\pi\,\frac{4A/\sqrt{3}}{4\left(n+\sqrt{3}-1\right)^2} = \frac{n(n+1)}{\left(n+\sqrt{3}-1\right)^2}\,\frac{\pi A}{2\sqrt{3}} \quad\Rightarrow$$

$$\frac{A_n}{A} = \frac{n(n+1)}{\left(n+\sqrt{3}-1\right)^2}\,\frac{\pi}{2\sqrt{3}}$$

$$= \frac{1 + 1/n}{\left[1 + (\sqrt{3} \quad 1)/n\right]^2}\,\frac{\pi}{2\sqrt{3}} \to \frac{\pi}{2\sqrt{3}} \text{ as } n \to \infty$$



16. Given $a_0 = a_1 = 1$ and $a_n = \dfrac{(n-1)(n-2)a_{n-1} - (n-3)a_{n-2}}{n(n \quad 1)}$, we calculate the next few terms of the sequence:

$a_2 = \dfrac{1\cdot 0\cdot a_1 - (-1)a_0}{2\cdot 1} = \dfrac{1}{2}$, $a_3 = \dfrac{2\cdot 1\cdot a_2 - 0\cdot a_1}{3\cdot 2} = \dfrac{1}{6}$, $a_4 = \dfrac{3\cdot 2\cdot a_3 - 1\cdot a_2}{4\cdot 3} = \dfrac{1}{24}$. It seems that $a_n = \dfrac{1}{n!}$,

so we try to prove this by induction. The first step is done, so assume $a_k = \dfrac{1}{k!}$ and $a_{k-1} = \dfrac{1}{(k-1)!}$. Then

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$a_{k+1} = \dfrac{k(k-1)a_k - (k-2)a_{k-1}}{(k+1)k} = \dfrac{\dfrac{k(k-1)}{k!} - \dfrac{k-2}{(k-1)!}}{(k+1)k} = \dfrac{(k-1)-(k-2)}{[(k+1)(k)](k-1)!} = \dfrac{1}{(k+1)!}$ and the induction is complete. Therefore, $\sum_{n=0}^{\infty} a_n = \sum_{n=0}^{\infty} \dfrac{1}{n!} = e$.

17. (a)



The x-intercepts of the curve occur where $\sin x = 0 \;\Leftrightarrow\; x = n\pi$, n an integer. So using the formula for disks (and either a CAS or $\sin^2 x = \frac{1}{2}(1-\cos 2x)$ and Formula 99 to evaluate the integral), the volume of the nth bead is

$$V_n = \pi \int_{(n-1)\pi}^{n\pi} (e^{-x/10}\sin x)^2\,dx = \pi \int_{(n-1)\pi}^{n\pi} e^{-x/5}\sin^2 x\,dx$$
$$= \tfrac{250\pi}{101}\left(e^{-(n-1)\pi/5} - e^{-n\pi/5}\right)$$

(b) The total volume is

$$\pi \int_0^{\infty} e^{-x/5}\sin^2 x\,dx = \sum_{n=1}^{\infty} V_n = \tfrac{250\pi}{101}\sum_{n=1}^{\infty}\left[e^{-(n-1)\pi/5} - e^{-n\pi/5}\right] = \tfrac{250\pi}{101} \quad \text{[telescoping sum]}.$$

Another method: If the volume in part (a) has been written as $V_n = \frac{250\pi}{101}e^{-n\pi/5}(e^{\pi/5}-1)$, then we recognize $\sum_{n=1}^{\infty} V_n$ as a geometric series with $a = \frac{250\pi}{101}(1-e^{-\pi/5})$ and $r = e^{-\pi/5}$.

18. (a) Since P_n is defined as the midpoint of $P_{n-4}P_{n-3}$, $x_n = \frac{1}{2}(x_{n-4} + x_{n-3})$ for $n \geq 5$. So we prove by induction that $\frac{1}{2}x_n + x_{n+1} + x_{n+2} + x_{n+3} = 2$. The case $n = 1$ is immediate, since $\frac{1}{2}\cdot 0 + 1 + 1 + 0 = 2$. Assume that the result holds for $n = k-1$, that is, $\frac{1}{2}x_{k-1} + x_k + x_{k+1} + x_{k+2} = 2$. Then for $n = k$,

$$\tfrac{1}{2}x_k + x_{k+1} + x_{k+2} + x_{k+3} = \tfrac{1}{2}x_k + x_{k+1} + x_{k+2} + \tfrac{1}{2}(x_{k+3-4} + x_{k+3-3}) \quad \text{[by above]}$$
$$= \tfrac{1}{2}x_{k-1} + x_k + x_{k+1} + x_{k+2} = 2 \quad \text{[by the induction hypothesis]}$$

Similarly, for $n \geq 5$, $y_n = \frac{1}{2}(y_{n-4} + y_{n-3})$, so the same argument as above holds for y, with 2 replaced by $\frac{1}{2}y_1 + y_2 + y_3 + y_4 = \frac{1}{2}\cdot 1 + 1 + 0 + 0 = \frac{3}{2}$. So $\frac{1}{2}y_n + y_{n+1} + y_{n+2} + y_{n+3} = \frac{3}{2}$ for all n.

(b) $\lim_{n\to\infty}\left(\frac{1}{2}x_n + x_{n+1} + x_{n+2} + x_{n+3}\right) = \frac{1}{2}\lim_{n\to\infty} x_n + \lim_{n\to\infty} x_{n+1} + \lim_{n\to\infty} x_{n+2} + \lim_{n\to\infty} x_{n+3} = 2$. Since all the limits on the left hand side are the same, we get $\frac{7}{2}\lim_{n\to\infty} x_n = 2 \;\Rightarrow\; \lim_{n\to\infty} x_n = \frac{4}{7}$. In the same way, $\frac{7}{2}\lim_{n\to\infty} y_n = \frac{3}{2} \;\Rightarrow\; \lim_{n\to\infty} y_n = \frac{3}{7}$, so $P = \left(\frac{4}{7}, \frac{3}{7}\right)$.

19. By Table 1 in Section 11.10, $\tan^{-1} x = \sum_{n=0}^{\infty}(-1)^n \dfrac{x^{2n+1}}{2n+1}$ for $|x| < 1$. In particular, for $x = \dfrac{1}{\sqrt{3}}$, we have $\dfrac{\pi}{6} = \tan^{-1}\left(\dfrac{1}{\sqrt{3}}\right) = \sum_{n=0}^{\infty}(-1)^n \dfrac{\left(1/\sqrt{3}\right)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty}(-1)^n\left(\dfrac{1}{3}\right)^n \dfrac{1}{\sqrt{3}}\dfrac{1}{2n+1}$, so

$$\pi = \frac{6}{\sqrt{3}}\sum_{n=0}^{\infty}\frac{(-1)^n}{(2n+1)3^n} = 2\sqrt{3}\sum_{n=0}^{\infty}\frac{(-1)^n}{(2n+1)3^n} = 2\sqrt{3}\left(1 + \sum_{n=1}^{\infty}\frac{(-1)^n}{(2n+1)3^n}\right) \;\Rightarrow\; \sum_{n=1}^{\infty}\frac{(-1)^n}{(2n+1)3^n} = \frac{\pi}{2\sqrt{3}} - 1.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

20. (a) Using $s_n = a + ar + ar^2 + \cdots + ar^{n-1} = \dfrac{a(1-r^n)}{1-r}$,

$$1 - x + x^2 - x^3 + \cdots + x^{2n-2} - x^{2n-1} = \frac{1\left[1-(-x)^{2n}\right]}{1-(-x)} = \frac{1-x^{2n}}{1+x}.$$

(b) $\displaystyle\int_0^1 (1 - x + x^2 - x^3 + \cdots + x^{2n-2} - x^{2n-1})\,dx = \int_0^1 \frac{1-x^{2n}}{1+x}\,dx \quad \Rightarrow$

$$\left[x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots + \frac{x^{2n-1}}{2n-1} - \frac{x^{2n}}{2n}\right]_0^1 = \int_0^1 \frac{dx}{1+x} - \int_0^1 \frac{x^{2n}}{1+x}\,dx \quad \Rightarrow$$

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots + \frac{1}{2n-1} - \frac{1}{2n} = \int_0^1 \frac{dx}{1+x} - \int_0^1 \frac{x^{2n}}{1+x}\,dx$$

(c) Since $1 - \dfrac{1}{2} = \dfrac{1}{1\cdot 2}, \dfrac{1}{3} - \dfrac{1}{4} = \dfrac{1}{3\cdot 4}, \cdots, \dfrac{1}{2n-1} - \dfrac{1}{2n} = \dfrac{1}{(2n-1)(2n)}$, we see from part (b) that

$\dfrac{1}{1\cdot 2} + \dfrac{1}{3\cdot 4} + \cdots + \dfrac{1}{(2n-1)(2n)} - \displaystyle\int_0^1 \frac{dx}{1+x} = -\int_0^1 \frac{x^{2n}}{1+x}\,dx$. Thus,

$$\left|\frac{1}{1\cdot 2} + \frac{1}{3\cdot 4} + \cdots + \frac{1}{(2n-1)(2n)} - \int_0^1 \frac{dx}{1+x}\right| = \int_0^1 \frac{x^{2n}}{1+x}\,dx < \int_0^1 x^{2n}\,dx$$

$\left[\text{since } \dfrac{x^{2n}}{1+x} < x^{2n} \text{ for } 0 < x \le 1\right]$.

(d) Note that $\displaystyle\int_0^1 \frac{dx}{1+x} = \Big[\ln(1+x)\Big]_0^1 = \ln 2$ and $\displaystyle\int_0^1 x^{2n}\,dx = \left[\frac{x^{2n+1}}{2n+1}\right]_0^1 = \frac{1}{2n+1}$. So part (c) becomes

$\left|\dfrac{1}{1\cdot 2} + \dfrac{1}{3\cdot 4} + \cdots + \dfrac{1}{(2n-1)(2n)} - \ln 2\right| < \dfrac{1}{2n+1}$. In other words, the nth partial sum s_n of the given series satisfies $|s_n - \ln 2| < \dfrac{1}{2n+1}$. Thus, $\lim\limits_{n\to\infty} s_n = \ln 2$, that is, $\dfrac{1}{1\cdot 2} + \dfrac{1}{3\cdot 4} + \dfrac{1}{5\cdot 6} + \dfrac{1}{7\cdot 8} + \cdots = \ln 2$.

21. Let $f(x)$ denote the left-hand side of the equation $1 + \dfrac{x}{2!} + \dfrac{x^2}{4!} + \dfrac{x^3}{6!} + \dfrac{x^4}{8!} + \cdots = 0$. If $x \ge 0$, then $f(x) \ge 1$ and there are no solutions of the equation. Note that $f(-x^2) = 1 - \dfrac{x^2}{2!} + \dfrac{x^4}{4!} - \dfrac{x^6}{6!} + \dfrac{x^8}{8!} - \cdots = \cos x$. The solutions of $\cos x = 0$ for $x < 0$ are given by $x = \dfrac{\pi}{2} - \pi k$, where k is a positive integer. Thus, the solutions of $f(x) = 0$ are $x = -\left(\dfrac{\pi}{2} - \pi k\right)^2$, where k is a positive integer.

22. Suppose the base of the first right triangle has length a. Then by repeated use of the Pythagorean theorem, we find that the base of the second right triangle has length $\sqrt{1+a^2}$, the base of the third right triangle has length $\sqrt{2+a^2}$, and in general, the nth right triangle has base of length $\sqrt{n-1+a^2}$ and hypotenuse of length $\sqrt{n+a^2}$. Thus, $\theta_n = \tan^{-1}\left(1/\sqrt{n-1+a^2}\right)$ and $\displaystyle\sum_{n=1}^{\infty} \theta_n = \sum_{n=1}^{\infty} \tan^{-1}\left(\frac{1}{\sqrt{n-1+a^2}}\right) = \sum_{n=0}^{\infty} \tan^{-1}\left(\frac{1}{\sqrt{n+a^2}}\right)$. We wish to show that this series diverges.

First notice that the series $\displaystyle\sum_{n=1}^{\infty} \frac{1}{\sqrt{n+a^2}}$ diverges by the Limit Comparison Test with the divergent p-series $\displaystyle\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$

$\left[p = \frac{1}{2} \le 1\right]$ since $\displaystyle\lim_{n\to\infty} \frac{1/\sqrt{n+a^2}}{1/\sqrt{n}} = \lim_{n\to\infty} \frac{\sqrt{n}}{\sqrt{n+a^2}} = \lim_{n\to\infty} \sqrt{\frac{n}{n+a^2}} = \lim_{n\to\infty} \sqrt{\frac{1}{1+a^2/n}} = 1 > 0$. Thus,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\sum_{n=0}^{\infty} \dfrac{1}{\sqrt{n+a^2}}$ also diverges. Now $\sum_{n=0}^{\infty} \tan^{-1}\left(\dfrac{1}{\sqrt{n+a^2}}\right)$ diverges by the Limit Comparison Test with $\sum_{n=0}^{\infty} \dfrac{1}{\sqrt{n+a^2}}$ since

$$\lim_{n\to\infty} \frac{\tan^{-1}\left(1/\sqrt{n+a^2}\right)}{1/\sqrt{n+a^2}} = \lim_{x\to\infty} \frac{\tan^{-1}\left(1/\sqrt{x+a^2}\right)}{1/\sqrt{x+a^2}} = \lim_{y\to\infty} \frac{\tan^{-1}(1/y)}{1/y} \qquad \left[y=\sqrt{x+a^2}\right]$$

$$= \lim_{z\to 0^+} \frac{\tan^{-1} z}{z} \quad [z=1/y] \quad \overset{\text{H}}{=} \lim_{z\to 0^+} \frac{1/(1+z^2)}{1} = 1 > 0$$

Thus, $\sum_{n=1}^{\infty} \theta_n$ is a divergent series.

23. Call the series S. We group the terms according to the number of digits in their denominators:

$$S = \underbrace{\left(\tfrac{1}{1}+\tfrac{1}{2}+\cdots+\tfrac{1}{8}+\tfrac{1}{9}\right)}_{g_1} + \underbrace{\left(\tfrac{1}{11}+\cdots+\tfrac{1}{99}\right)}_{g_2} + \underbrace{\left(\tfrac{1}{111}+\cdots+\tfrac{1}{999}\right)}_{g_3} + \cdots$$

Now in the group g_n, since we have 9 choices for each of the n digits in the denominator, there are 9^n terms.

Furthermore, each term in g_n is less than $\frac{1}{10^{n-1}}$ [except for the first term in g_1]. So $g_n < 9^n \cdot \frac{1}{10^{n-1}} = 9\left(\frac{9}{10}\right)^{n-1}$.

Now $\sum_{n=1}^{\infty} 9\left(\frac{9}{10}\right)^{n-1}$ is a geometric series with $a = 9$ and $r = \frac{9}{10} < 1$. Therefore, by the Comparison Test,

$S = \sum_{n=1}^{\infty} g_n < \sum_{n=1}^{\infty} 9\left(\frac{9}{10}\right)^{n-1} = \frac{9}{1-9/10} = 90.$

24. (a) Let $f(x) = \dfrac{x}{1-x-x^2} = \sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots$. Then

$$x = (1-x-x^2)(c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots)$$

$$\begin{aligned} x = {} & c_0 + c_1 x + c_2 x^2 + c_3 x^3 + c_4 x^4 + c_5 x^5 + \cdots \\ & - c_0 x - c_1 x^2 - c_2 x^3 - c_3 x^4 - c_4 x^5 - \cdots \\ & - c_0 x^2 - c_1 x^3 - c_2 x^4 - c_3 x^5 - \cdots \end{aligned}$$

$$x = c_0 + (c_1 - c_0)x + (c_2 - c_1 - c_0)x^2 + (c_3 - c_2 - c_1)x^3 + \cdots$$

Comparing coefficients of powers of x gives us $c_0 = 0$ and

$$\begin{aligned} c_1 - c_0 = 1 \quad &\Rightarrow \quad c_1 = c_0 + 1 = 1 \\ c_2 - c_1 - c_0 = 0 \quad &\Rightarrow \quad c_2 = c_1 + c_0 = 1 + 0 = 1 \\ c_3 - c_2 - c_1 = 0 \quad &\Rightarrow \quad c_3 = c_2 + c_1 = 1 + 1 = 2 \end{aligned}$$

In general, we have $c_n = c_{n-1} + c_{n-2}$ for $n \geq 3$. Each c_n is equal to the nth Fibonacci number, that is,

$$\sum_{n=0}^{\infty} c_n x^n = \sum_{n=1}^{\infty} c_n x^n = \sum_{n=1}^{\infty} f_n x^n$$

(b) Completing the square on $x^2 + x - 1$ gives us

$$\left(x^2 + x + \frac{1}{4}\right) - 1 - \frac{1}{4} = \left(x + \frac{1}{2}\right)^2 - \frac{5}{4} = \left(x + \frac{1}{2}\right)^2 - \left(\frac{\sqrt{5}}{2}\right)^2$$

$$= \left(x + \frac{1}{2} + \frac{\sqrt{5}}{2}\right)\left(x + \frac{1}{2} - \frac{\sqrt{5}}{2}\right) = \left(x + \frac{1+\sqrt{5}}{2}\right)\left(x + \frac{1-\sqrt{5}}{2}\right)$$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

So $\dfrac{x}{1-x-x^2} = \dfrac{-x}{x^2+x-1} = \dfrac{-x}{\left(x+\frac{1+\sqrt{5}}{2}\right)\left(x+\frac{1-\sqrt{5}}{2}\right)}$. The factors in the denominator are linear, so the partial fraction decomposition is

$$\frac{-x}{\left(x+\frac{1+\sqrt{5}}{2}\right)\left(x+\frac{1-\sqrt{5}}{2}\right)} = \frac{A}{x+\frac{1+\sqrt{5}}{2}} + \frac{B}{x+\frac{1-\sqrt{5}}{2}} \quad -x = A\left(x+\tfrac{1-\sqrt{5}}{2}\right) + B\left(x+\tfrac{1+\sqrt{5}}{2}\right)$$

If $x = \frac{-1+\sqrt{5}}{2}$, then $-\frac{-1+\sqrt{5}}{2} = B\sqrt{5} \quad \Rightarrow \quad B = \frac{1-\sqrt{5}}{2\sqrt{5}}$.

If $x = \frac{-1-\sqrt{5}}{2}$, then $-\frac{-1-\sqrt{5}}{2} = A(-\sqrt{5}) \quad \Rightarrow \quad A = \frac{1+\sqrt{5}}{-2\sqrt{5}}$. Thus,

$$\frac{x}{1-x-x^2} = \frac{\frac{1+\sqrt{5}}{-2\sqrt{5}}}{x+\frac{1+\sqrt{5}}{2}} + \frac{\frac{1-\sqrt{5}}{2\sqrt{5}}}{x+\frac{1-\sqrt{5}}{2}} = \frac{\frac{1+\sqrt{5}}{-2\sqrt{5}}}{x+\frac{1+\sqrt{5}}{2}} \cdot \frac{\frac{2}{1+\sqrt{5}}}{\frac{2}{1+\sqrt{5}}} + \frac{\frac{1-\sqrt{5}}{2\sqrt{5}}}{x+\frac{1-\sqrt{5}}{2}} \cdot \frac{\frac{2}{1-\sqrt{5}}}{\frac{2}{1-\sqrt{5}}}$$

$$= \frac{-1/\sqrt{5}}{1+\frac{2}{1+\sqrt{5}}x} + \frac{1/\sqrt{5}}{1+\frac{2}{1-\sqrt{5}}x} = -\frac{1}{\sqrt{5}}\sum_{n=0}^{\infty}\left(-\frac{2}{1+\sqrt{5}}x\right)^n + \frac{1}{\sqrt{5}}\sum_{n=0}^{\infty}\left(-\frac{2}{1-\sqrt{5}}x\right)^n$$

$$= \frac{1}{\sqrt{5}}\sum_{n=0}^{\infty}\left[\left(\frac{-2}{1-\sqrt{5}}\right)^n - \left(\frac{-2}{1+\sqrt{5}}\right)^n\right]x^n$$

$$= \frac{1}{\sqrt{5}}\sum_{n=1}^{\infty}\left[\frac{(-2)^n(1+\sqrt{5})^n - (-2)^n(1-\sqrt{5})^n}{(1-\sqrt{5})^n(1+\sqrt{5})^n}\right]x^n \qquad [\text{the } n=0 \text{ term is } 0]$$

$$= \frac{1}{\sqrt{5}}\sum_{n=1}^{\infty}\left[\frac{(-2)^n\left((1+\sqrt{5})^n - (1-\sqrt{5})^n\right)}{(1-5)^n}\right]x^n$$

$$= \frac{1}{\sqrt{5}}\sum_{n=1}^{\infty}\left[\frac{(1+\sqrt{5})^n - (1-\sqrt{5})^n}{2^n}\right]x^n \qquad [(-4)^n = (-2)^n \cdot 2^n]$$

From part (a), this series must equal $\sum_{n=1}^{\infty} f_n x^n$, so $f_n = \dfrac{(1+\sqrt{5})^n - (1-\sqrt{5})^n}{2^n\sqrt{5}}$, which is an explicit formula for the nth Fibonacci number.

25. $u = 1 + \dfrac{x^3}{3!} + \dfrac{x^6}{6!} + \dfrac{x^9}{9!} + \cdots$, $v = x + \dfrac{x^4}{4!} + \dfrac{x^7}{7!} + \dfrac{x^{10}}{10!} + \cdots$, $w = \dfrac{x^2}{2!} + \dfrac{x^5}{5!} + \dfrac{x^8}{8!} + \cdots$.

Use the Ratio Test to show that the series for u, v, and w have positive radii of convergence (∞ in each case), so Theorem 11.9.2 applies, and hence, we may differentiate each of these series:

$$\frac{du}{dx} = \frac{3x^2}{3!} + \frac{6x^5}{6!} + \frac{9x^8}{9!} + \cdots = \frac{x^2}{2!} + \frac{x^5}{5!} + \frac{x^8}{8!} + \cdots = w$$

Similarly, $\dfrac{dv}{dx} = 1 + \dfrac{x^3}{3!} + \dfrac{x^6}{6!} + \dfrac{x^9}{9!} + \cdots = u$, and $\dfrac{dw}{dx} = x + \dfrac{x^4}{4!} + \dfrac{x^7}{7!} + \dfrac{x^{10}}{10!} + \cdots = v$.

So $u' = w$, $v' = u$, and $w' = v$. Now differentiate the left-hand side of the desired equation:

$$\frac{d}{dx}(u^3 + v^3 + w^3 - 3uvw) = 3u^2u' + 3v^2v' + 3w^2w' - 3(u'vw + uv'w + uvw')$$

$$= 3u^2w + 3v^2u + 3w^2v - 3(vw^2 + u^2w + uv^2) = 0 \quad \Rightarrow$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$u^3 + v^3 + w^3 - 3uvw = C$. To find the value of the constant C, we put $x = 0$ in the last equation and get $1^3 + 0^3 + 0^3 - 3(1 \cdot 0 \cdot 0) = C \quad \Rightarrow \quad C = 1$, so $u^3 + v^3 + w^3 - 3uvw = 1$.

26. To prove: If $n > 1$, then the nth partial sum $s_n = \sum_{i=1}^{n} \frac{1}{i}$ of the harmonic series is not an integer.

Proof: Let 2^k be the largest power of 2 that is less than or equal to n and let M be the product of all the odd positive integers that are less than or equal to n. Suppose that $s_n = m$, an integer. Then $M2^k s_n = M2^k m$. Since $n \geq 2$, we have $k \geq 1$, and hence, $M2^k m$ is an even integer. We will show that $M2^k s_n$ is an odd integer, contradicting the equality $M2^k s_n = M2^k m$ and showing that the supposition that s_n is an integer must have been wrong.

$M2^k s_n = M2^k \sum_{i=1}^{n} \frac{1}{i} = \sum_{i=1}^{n} \frac{M2^k}{i}$. If $1 \leq i \leq n$ and i is odd, then $\frac{M}{i}$ is an odd integer since i is one of the odd integers that were multiplied together to form M. Thus, $\frac{M2^k}{i}$ is an even integer in this case. If $1 \leq i \leq n$ and i is even, then we can write $i = 2^r l$, where 2^r is the largest power of 2 dividing i and l is odd. If $r < k$, then $\frac{M2^k}{i} = \frac{2^k}{2^r} \cdot \frac{M}{l} = 2^{k-r} \frac{M}{l}$, which is an even integer, the product of the even integer 2^{k-r} and the odd integer $\frac{M}{l}$. If $r = k$, then $l = 1$, since $l > 1 = l \geq 2 \quad \Rightarrow$ $i = 2^k l \geq 2^k \cdot 2 = 2^{k+1}$, contrary to the choice of 2^k as the largest power of 2 that is less than or equal to n. This shows that $r = k$ only when $i = 2^k$. In that case, $\frac{M2^k}{i} = M$, an *odd* integer. Since $\frac{M2^k}{i}$ is an even integer for every i except 2^k and $\frac{M2^k}{i}$ is an odd integer when $i = 2^k$, we see that $M2^k s_n$ is an odd integer. This concludes the proof.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

12 □ VECTORS AND THE GEOMETRY OF SPACE

12.1 Three-Dimensional Coordinate Systems

1. We start at the origin, which has coordinates $(0, 0, 0)$. First we move 4 units along the positive x-axis, affecting only the x-coordinate, bringing us to the point $(4, 0, 0)$. We then move 3 units straight downward, in the negative z-direction. Thus only the z-coordinate is affected, and we arrive at $(4, 0, -3)$.

2.



3. The distance from a point to the yz-plane is the absolute value of the x-coordinate of the point. $C(2, 4, 6)$ has the x-coordinate with the smallest absolute value, so C is the point closest to the yz-plane. $A(-4, 0, -1)$ must lie in the xz-plane since the distance from A to the xz-plane, given by the y-coordinate of A, is 0.

4. The projection of $(2, 3, 5)$ onto the xy-plane is $(2, 3, 0)$; onto the yz-plane, $(0, 3, 5)$; onto the xz-plane, $(2, 0, 5)$.

The length of the diagonal of the box is the distance between the origin and $(2, 3, 5)$, given by

$$\sqrt{(2-0)^2 + (3-0)^2 + (5-0)^2} = \sqrt{38} \approx 6.16$$



5. In $\mathbb{R}^2$, the equation $x = 4$ represents a line parallel to the y-axis and 4 units to the right of it. In $\mathbb{R}^3$, the equation $x = 4$ represents the set $\{(x, y, z) \mid x = 4\}$, the set of all points whose x-coordinate is 4. This is the vertical plane that is parallel to the yz-plane and 4 units in front of it.





6. In $\mathbb{R}^3$, the equation $y = 3$ represents a vertical plane that is parallel to the xz-plane and 3 units to the right of it. The equation $z = 5$ represents a horizontal plane parallel to the xy-plane and 5 units above it. The pair of equations $y = 3$, $z = 5$ represents the set of points that are simultaneously on both planes, or in other words, the line of intersection of the planes $y = 3$, $z = 5$.

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

This line can also be described as the set $\{(x, 3, 5) \mid x \in \mathbb{R}\}$, which is the set of all points in $\mathbb{R}^3$ whose x-coordinate may vary but whose y- and z-coordinates are fixed at 3 and 5, respectively. Thus the line is parallel to the x-axis and intersects the yz-plane in the point $(0, 3, 5)$.



7. The equation $x + y = 2$ represents the set of all points in $\mathbb{R}^3$ whose x- and y-coordinates have a sum of 2, or equivalently where $y = 2 - x$. This is the set $\{(x, 2 - x, z) \mid x \in \mathbb{R}, z \in \mathbb{R}\}$ which is a vertical plane that intersects the xy-plane in the line $y = 2 - x$, $z = 0$.



8. The equation $x^2 + z^2 = 9$ has no restrictions on y, and the x- and z-coordinates satisfy the equation for a circle of radius 3 with center the origin. Thus the surface $x^2 + z^2 = 9$ in $\mathbb{R}^3$ consists of all possible vertical circles (parallel to the xz-plane) $x^2 + z^2 = 9$, $y = k$, and is therefore a circular cylinder with radius 3 whose axis is the y-axis.



9. We can find the lengths of the sides of the triangle by using the distance formula between pairs of vertices:

$$|PQ| = \sqrt{(7-3)^2 + [0-(-2)]^2 + [1-(-3)]^2} = \sqrt{16+4+16} = 6$$

$$|QR| = \sqrt{(1-7)^2 + (2-0)^2 + (1-1)^2} = \sqrt{36+4+0} = \sqrt{40} = 2\sqrt{10}$$

$$|RP| = \sqrt{(3-1)^2 + (-2-2)^2 + (-3-1)^2} = \sqrt{4+16+16} = 6$$

The longest side is QR, but the Pythagorean Theorem is not satisfied: $|PQ|^2 + |RP|^2 \neq |QR|^2$. Thus PQR is not a right triangle. PQR is isosceles, as two sides have the same length.

10. Compute the lengths of the sides of the triangle by using the distance formula between pairs of vertices:

$$|PQ| = \sqrt{(4-2)^2 + [1-(-1)]^2 + (1-0)^2} = \sqrt{4+4+1} = 3$$

$$|QR| = \sqrt{(4-4)^2 + (-5-1)^2 + (4-1)^2} = \sqrt{0+36+9} = \sqrt{45} = 3\sqrt{5}$$

$$|RP| = \sqrt{(2-4)^2 + [-1-(-5)]^2 + (0-4)^2} = \sqrt{4+16+16} = 6$$

Since the Pythagorean Theorem is satisfied by $|PQ|^2 + |RP|^2 = |QR|^2$, PQR is a right triangle. PQR is not isosceles, as no two sides have the same length.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. (a) First we find the distances between points:

$$|AB| = \sqrt{(3-2)^2 + (7-4)^2 + (-2-2)^2} = \sqrt{26}$$
$$|BC| = \sqrt{(1-3)^2 + (3-7)^2 + [3-(-2)]^2} = \sqrt{45} = 3\sqrt{5}$$
$$|AC| = \sqrt{(1-2)^2 + (3-4)^2 + (3-2)^2} = \sqrt{3}$$

In order for the points to lie on a straight line, the sum of the two shortest distances must be equal to the longest distance. Since $\sqrt{26} + \sqrt{3} \neq 3\sqrt{5}$, the three points do not lie on a straight line.

(b) First we find the distances between points:

$$|DE| = \sqrt{(1-0)^2 + [-2-(-5)]^2 + (4-5)^2} = \sqrt{11}$$
$$|EF| = \sqrt{(3-1)^2 + [4-(-2)]^2 + (2-4)^2} = \sqrt{44} = 2\sqrt{11}$$
$$|DF| = \sqrt{(3-0)^2 + [4-(-5)]^2 + (2-5)^2} = \sqrt{99} = 3\sqrt{11}$$

Since $|DE| + |EF| = |DF|$, the three points lie on a straight line.

12. (a) The distance from a point to the xy-plane is the absolute value of the z-coordinate of the point. Thus, the distance is $|6| = 6$.

(b) Similarly, the distance to the yz-plane is the absolute value of the x-coordinate of the point: $|4| = 4$.

(c) The distance to the xz-plane is the absolute value of the y-coordinate of the point: $|-2| = 2$.

(d) The point on the x-axis closest to $(4, -2, 6)$ is the point $(4, 0, 0)$. (Approach the x-axis perpendicularly.) The distance from $(4, -2, 6)$ to the x-axis is the distance between these two points: $\sqrt{(4-4)^2 + (-2-0)^2 + (6-0)^2} = \sqrt{40} = 2\sqrt{10} \approx 6.32$.

(e) The point on the y-axis closest to $(4, -2, 6)$ is $(0, -2, 0)$. The distance between these points is $\sqrt{(4-0)^2 + [-2-(-2)]^2 + (6-0)^2} = \sqrt{52} = 2\sqrt{13} \approx 7.21$.

(f) The point on the z-axis closest to $(4, -2, 6)$ is $(0, 0, 6)$. The distance between these points is $\sqrt{(4-0)^2 + (-2-0)^2 + (6-6)^2} = \sqrt{20} = 2\sqrt{5} \approx 4.47$.

13. An equation of the sphere with center $(-3, 2, 5)$ and radius 4 is $[x-(-3)]^2 + (y-2)^2 + (z-5)^2 = 4^2$ or $(x+3)^2 + (y-2)^2 + (z-5)^2 = 16$. The intersection of this sphere with the yz-plane is the set of points on the sphere whose x-coordinate is 0. Putting $x = 0$ into the equation, we have $9 + (y-2)^2 + (z-5)^2 = 16$, $x = 0$ or $(y-2)^2 + (z-5)^2 = 7$, $x = 0$, which represents a circle in the yz-plane with center $(0, 2, 5)$ and radius $\sqrt{7}$.

14. An equation of the sphere with center $(2, -6, 4)$ and radius 5 is $(x-2)^2 + [y-(-6)]^2 + (z-4)^2 = 5^2$ or $(x-2)^2 + (y+6)^2 + (z-4)^2 = 25$. The intersection of this sphere with the xy-plane is the set of points on the sphere whose z-coordinate is 0. Putting $z = 0$ into the equation, we have $(x-2)^2 + (y+6)^2 = 9$, $z = 0$ which represents a circle in the xy-plane with center $(2, -6, 0)$ and radius 3. To find the intersection with the xz-plane, we set $y = 0$: $(x-2)^2 + (z-4)^2 = -11$. Since no points satisfy this equation, the sphere does not intersect the xz-plane. (Also note that the distance from the center of the sphere to the xz-plane is greater than the radius of the sphere.) To find the intersection with the yz-plane, we set $x = 0$: $(y+6)^2 + (z-4)^2 = 21$, $x = 0$, a circle in the yz-plane with center $(0, -6, 4)$ and radius $\sqrt{21}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

15. The radius of the sphere is the distance between $(4, 3, -1)$ and $(3, 8, 1)$: $r = \sqrt{(3-4)^2 + (8-3)^2 + [1-(-1)]^2} = \sqrt{30}$.

Thus, an equation of the sphere is $(x-3)^2 + (y-8)^2 + (z-1)^2 = 30$.

16. If the sphere passes through the origin, the radius of the sphere must be the distance from the origin to the point $(1, 2, 3)$:

$r = \sqrt{(1-0)^2 + (2-0)^2 + (3-0)^2} = \sqrt{14}$. Then an equation of the sphere is $(x-1)^2 + (y-2)^2 + (z-3)^2 = 14$.

17. Completing squares in the equation $x^2 + y^2 + z^2 - 2x - 4y + 8z = 15$ gives

$(x^2 - 2x + 1) + (y^2 - 4y + 4) + (z^2 + 8z + 16) = 15 + 1 + 4 + 16 \;\Rightarrow\; (x-1)^2 + (y-2)^2 + (z+4)^2 = 36$, which we recognize as an equation of a sphere with center $(1, 2, -4)$ and radius 6.

18. Completing squares in the equation gives $(x^2 + 8x + 16) + (y^2 - 6y + 9) + (z^2 + 2z + 1) = -17 + 16 + 9 + 1 \;\Rightarrow$

$(x+4)^2 + (y-3)^2 + (z+1)^2 = 9$, which we recognize as an equation of a sphere with center $(-4, 3, -1)$ and radius 3.

19. Completing squares in the equation $2x^2 - 8x + 2y^2 + 2z^2 + 24z = 1$ gives

$2(x^2 - 4x + 4) + 2y^2 + 2(z^2 + 12z + 36) = 1 + 8 + 72 \;\Rightarrow\; 2(x-2)^2 + 2y^2 + 2(z+6)^2 = 81 \;\Rightarrow$

$(x-2)^2 + y^2 + (z+6)^2 = \frac{81}{2}$, which we recognize as an equation of a sphere with center $(2, 0, -6)$ and radius $\sqrt{\frac{81}{2}} = 9/\sqrt{2}$.

20. Completing squares in the equation $3x^2 + 3y^2 - 6y + 3z^2 - 12z = 10$ gives

$3x^2 + 3(y^2 - 2y + 1) + 3(z^2 - 4z + 4) = 10 + 3 + 12 \;\Rightarrow\; 3x^2 + 3(y-1)^2 + 3(z-2)^2 = 25 \;\Rightarrow$

$x^2 + (y-1)^2 + (z-2)^2 = \frac{25}{3}$, which we recognize as an equation of a sphere with center $(0, 1, 2)$ and radius $\sqrt{\frac{25}{3}} = 5/\sqrt{3}$.

21. (a) If the midpoint of the line segment from $P_1(x_1, y_1, z_1)$ to $P_2(x_2, y_2, z_2)$ is $Q = \left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}, \frac{z_1+z_2}{2}\right)$,

then the distances $|P_1Q|$ and $|QP_2|$ are equal, and each is half of $|P_1P_2|$. We verify that this is the case:

$$|P_1P_2| = \sqrt{(x_2-x_1)^2 + (y_2-y_1)^2 + (z_2-z_1)^2}$$

$$\begin{aligned}
|P_1Q| &= \sqrt{\left[\tfrac{1}{2}(x_1+x_2) - x_1\right]^2 + \left[\tfrac{1}{2}(y_1+y_2) - y_1\right]^2 + \left[\tfrac{1}{2}(z_1+z_2) - z_1\right]^2} \\
&= \sqrt{\left(\tfrac{1}{2}x_2 - \tfrac{1}{2}x_1\right)^2 + \left(\tfrac{1}{2}y_2 - \tfrac{1}{2}y_1\right)^2 + \left(\tfrac{1}{2}z_2 - \tfrac{1}{2}z_1\right)^2} \\
&= \sqrt{\left(\tfrac{1}{2}\right)^2\left[(x_2-x_1)^2 + (y_2-y_1)^2 + (z_2-z_1)^2\right]} = \tfrac{1}{2}\sqrt{(x_2-x_1)^2 + (y_2-y_1)^2 + (z_2-z_1)^2} \\
&= \tfrac{1}{2}\,|P_1P_2|
\end{aligned}$$

$$\begin{aligned}
|QP_2| &= \sqrt{\left[x_2 - \tfrac{1}{2}(x_1+x_2)\right]^2 + \left[y_2 - \tfrac{1}{2}(y_1+y_2)\right]^2 + \left[z_2 - \tfrac{1}{2}(z_1+z_2)\right]^2} \\
&= \sqrt{\left(\tfrac{1}{2}x_2 - \tfrac{1}{2}x_1\right)^2 + \left(\tfrac{1}{2}y_2 - \tfrac{1}{2}y_1\right)^2 + \left(\tfrac{1}{2}z_2 - \tfrac{1}{2}z_1\right)^2} = \sqrt{\left(\tfrac{1}{2}\right)^2\left[(x_2-x_1)^2 + (y_2-y_1)^2 + (z_2-z_1)^2\right]} \\
&= \tfrac{1}{2}\sqrt{(x_2-x_1)^2 + (y_2-y_1)^2 + (z_2-z_1)^2} = \tfrac{1}{2}\,|P_1P_2|
\end{aligned}$$

So Q is indeed the midpoint of P_1P_2.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) By part (a), the midpoints of sides AB, BC and CA are $P_1\left(-\frac{1}{2}, 1, 4\right)$, $P_2\left(1, \frac{1}{2}, 5\right)$ and $P_3\left(\frac{5}{2}, \frac{3}{2}, 4\right)$. Then the lengths of the medians are:

$$|AP_2| = \sqrt{0^2 + \left(\tfrac{1}{2} - 2\right)^2 + (5-3)^2} = \sqrt{\tfrac{9}{4} + 4} = \sqrt{\tfrac{25}{4}} = \tfrac{5}{2}$$

$$|BP_3| = \sqrt{\left(\tfrac{5}{2} + 2\right)^2 + \left(\tfrac{3}{2}\right)^2 + (4-5)^2} = \sqrt{\tfrac{81}{4} + \tfrac{9}{4} + 1} = \sqrt{\tfrac{94}{4}} = \tfrac{1}{2}\sqrt{94}$$

$$|CP_1| = \sqrt{\left(-\tfrac{1}{2} - 4\right)^2 + (1-1)^2 + (4-5)^2} = \sqrt{\tfrac{81}{4} + 1} = \tfrac{1}{2}\sqrt{85}$$

22. By Exercise 21(a), the midpoint of the diameter (and thus the center of the sphere) is $\left(\frac{5+1}{2}, \frac{4+6}{2}, \frac{3+(-9)}{2}\right) = (3, 5, -3)$. The radius is half the diameter, so $r = \frac{1}{2}\sqrt{(1-5)^2 + (6-4)^2 + (-9-3)^2} = \frac{1}{2}\sqrt{164} = \sqrt{41}$. Therefore an equation of the sphere is $(x-3)^2 + (y-5)^2 + (z+3)^2 = 41$.

23. (a) Since the sphere touches the xy-plane, its radius is the distance from its center, $(2, -3, 6)$, to the xy-plane, namely 6. Therefore $r = 6$ and an equation of the sphere is $(x-2)^2 + (y+3)^2 + (z-6)^2 = 6^2 = 36$.

(b) The radius of this sphere is the distance from its center $(2, -3, 6)$ to the yz-plane, which is 2. Therefore, an equation is $(x-2)^2 + (y+3)^2 + (z-6)^2 = 4$.

(c) Here the radius is the distance from the center $(2, -3, 6)$ to the xz-plane, which is 3. Therefore, an equation is $(x-2)^2 + (y+3)^2 + (z-6)^2 = 9$.

24. The largest sphere contained in the first octant must have a radius equal to the minimum distance from the center $(5, 4, 9)$ to any of the three coordinate planes. The shortest such distance is to the xz-plane, a distance of 4. Thus an equation of the sphere is $(x-5)^2 + (y-4)^2 + (z-9)^2 = 16$.

25. The equation $x = 5$ represents a plane parallel to the yz-plane and 5 units in front of it.

26. The equation $y = -2$ represents a plane parallel to the xz-plane and 2 units to the left of it.

27. The inequality $y < 8$ represents a half-space consisting of all points to the left of the plane $y = 8$.

28. The inequality $z \geq -1$ represents a half-space consisting of all points on or above the plane $z = -1$.

29. The inequality $0 \leq z \leq 6$ represents all points on or between the horizontal planes $z = 0$ (the xy-plane) and $z = 6$.

30. The equation $y^2 = 4 \quad \Leftrightarrow \quad y = \pm 2$ represents two vertical planes; $y = 2$ is parallel to the xz-plane, two units to the right of it, and $y = -2$ is two units to the left of it.

31. Because $z = -1$, all points in the region must lie in the horizontal plane $z = -1$. In addition, $x^2 + y^2 = 4$, so the region consists of all points that lie on a circle with radius 2 and center on the z-axis that is contained in the plane $z = -1$.

32. Here $x^2 + y^2 = 4$ with no restrictions on z, so a point in the region must lie on a circle of radius 2, center on the z-axis, but it could be in any horizontal plane $z = k$ (parallel to the xy-plane). Thus the region consists of all possible circles $x^2 + y^2 = 4$, $z = k$ and is therefore a circular cylinder with radius 2 whose axis is the z-axis.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

33. The equation $x^2 + y^2 + z^2 = 4$ is equivalent to $\sqrt{x^2 + y^2 + z^2} = 2$, so the region consists of those points whose distance from the origin is 2. This is the set of all points on a sphere with radius 2 and center $(0, 0, 0)$.

34. The inequality $x^2 + y^2 + z^2 \leq 4$ is equivalent to $\sqrt{x^2 + y^2 + z^2} \leq 2$, so the region consists of those points whose distance from the origin is at most 2. This is the set of all points on or inside a sphere with radius 2 and center $(0, 0, 0)$.

35. The inequalities $1 \leq x^2 + y^2 + z^2 \leq 5$ are equivalent to $1 \leq \sqrt{x^2 + y^2 + z^2} \leq \sqrt{5}$, so the region consists of those points whose distance from the origin is at least 1 and at most $\sqrt{5}$. This is the set of all points on or between spheres with radii 1 and $\sqrt{5}$ and centers $(0, 0, 0)$.

36. The equation $x = z$ represents a plane perpendicular to the xz-plane and intersecting the xz-plane in the line $x = z$, $y = 0$.

37. Here $x^2 + z^2 \leq 9$ or equivalently $\sqrt{x^2 + z^2} \leq 3$ which describes the set of all points in $\mathbb{R}^3$ whose distance from the y-axis is at most 3. Thus the inequality represents the region consisting of all points on or inside a circular cylinder of radius 3 with axis the y-axis.

38. The inequality $x^2 + y^2 + z^2 > 2z \quad \Leftrightarrow \quad x^2 + y^2 + (z - 1)^2 > 1$ is equivalent to $\sqrt{x^2 + y^2 + (z - 1)^2} > 1$, so the region consists of those points whose distance from the point $(0, 0, 1)$ is greater than 1. This is the set of all points outside the sphere with radius 1 and center $(0, 0, 1)$.

39. This describes all points whose x-coordinate is between 0 and 5, that is, $0 < x < 5$.

40. For any point on or above the disk in the xy-plane with center the origin and radius 2 we have $x^2 + y^2 \leq 4$. Also each point lies on or between the planes $z = 0$ and $z = 8$, so the region is described by $x^2 + y^2 \leq 4,\ 0 \leq z \leq 8$.

41. This describes a region all of whose points have a distance to the origin which is greater than r, but smaller than R. So inequalities describing the region are $r < \sqrt{x^2 + y^2 + z^2} < R$, or $r^2 < x^2 + y^2 + z^2 < R^2$.

42. The solid sphere itself is represented by $\sqrt{x^2 + y^2 + z^2} \leq 2$. Since we want only the upper hemisphere, we restrict the z-coordinate to nonnegative values. Then inequalities describing the region are $\sqrt{x^2 + y^2 + z^2} \leq 2$, $z \geq 0$, or $x^2 + y^2 + z^2 \leq 4$, $z \geq 0$.

43. (a) To find the x- and y-coordinates of the point P, we project it onto L_2 and project the resulting point Q onto the x- and y-axes. To find the z-coordinate, we project P onto either the xz-plane or the yz-plane (using our knowledge of its x- or y-coordinate) and then project the resulting point onto the z-axis. (Or, we could draw a line parallel to QO from P to the z-axis.) The coordinates of P are $(2, 1, 4)$.

(b) A is the intersection of L_1 and L_2, B is directly below the y-intercept of L_2, and C is directly above the x-intercept of L_2.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. Let $P = (x, y, z)$. Then $2\,|PB| = |PA| \;\Leftrightarrow\; 4\,|PB|^2 = |PA|^2 \;\Leftrightarrow$

$4\big((x-6)^2 + (y-2)^2 + (z+2)^2\big) = (x+1)^2 + (y-5)^2 + (z-3)^2 \;\Leftrightarrow$

$4\big(x^2 - 12x + 36\big) - x^2 - 2x + 4\big(y^2 - 4y + 4\big) - y^2 + 10y + 4\big(z^2 + 4z + 4\big) - z^2 + 6z = 35 \;\Leftrightarrow$

$3x^2 - 50x + 3y^2 - 6y + 3z^2 + 22z = 35 - 144 - 16 - 16 \;\Leftrightarrow\; x^2 - \frac{50}{3}x + y^2 - 2y + z^2 + \frac{22}{3}z = -\frac{141}{3}$.

By completing the square three times we get $\big(x - \frac{25}{3}\big)^2 + (y-1)^2 + \big(z + \frac{11}{3}\big)^2 = \frac{332}{9}$, which is an equation of a sphere with center $\big(\frac{25}{3}, 1, -\frac{11}{3}\big)$ and radius $\frac{\sqrt{332}}{3}$.

45. We need to find a set of points $\big\{P(x, y, z) \;\big|\; |AP| = |BP|\big\}$.

$\sqrt{(x+1)^2 + (y-5)^2 + (z-3)^2} = \sqrt{(x-6)^2 + (y-2)^2 + (z+2)^2} \;\Rightarrow$

$(x+1)^2 + (y-5) + (z-3)^2 = (x-6)^2 + (y-2)^2 + (z+2)^2 \;\Rightarrow$

$x^2 + 2x + 1 + y^2 - 10y + 25 + z^2 - 6z + 9 = x^2 - 12x + 36 + y^2 - 4y + 4 + z^2 + 4z + 4 \;\Rightarrow\; 14x - 6y - 10z = 9.$

Thus the set of points is a plane perpendicular to the line segment joining A and B (since this plane must contain the perpendicular bisector of the line segment AB).

46. Completing the square three times in the first equation gives $(x+2)^2 + (y-1)^2 + (z+2)^2 = 2^2$, a sphere with center $(-2, 1, 2)$ and radius 2. The second equation is that of a sphere with center $(0, 0, 0)$ and radius 2. The distance between the centers of the spheres is $\sqrt{(-2-0)^2 + (1-0)^2 + (-2-0)^2} = \sqrt{4+1+4} = 3$. Since the spheres have the same radius, the volume inside both spheres is symmetrical about the plane containing the circle of intersection of the spheres. The distance from this plane to the center of the circles is $\frac{3}{2}$. So the region inside both spheres consists of two caps of spheres of height $h = 2 - \frac{3}{2} = \frac{1}{2}$. From Exercise 5.2.49 [ET 6.2.49], the volume of a cap of a sphere is $V = \pi h^2\,(r - \frac{1}{3}h) = \pi\big(\frac{1}{2}\big)^2\big(2 - \frac{1}{3}\cdot\frac{1}{2}\big) = \frac{11\pi}{24}$. So the total volume is $2 \cdot \frac{11\pi}{24} = \frac{11\pi}{12}$.



47. The sphere $x^2 + y^2 + z^2 = 4$ has center $(0, 0, 0)$ and radius 2. Completing squares in $x^2 - 4x + y^2 - 4y + z^2 - 4z = -11$ gives $(x^2 - 4x + 4) + (y^2 - 4y + 4) + (z^2 - 4z + 4) = -11 + 4 + 4 + 4 \;\Rightarrow\; (x-2)^2 + (y-2)^2 + (z-2)^2 = 1$, so this is the sphere with center $(2, 2, 2)$ and radius 1. The (shortest) distance between the spheres is measured along the line segment connecting their centers. The distance between $(0, 0, 0)$ and $(2, 2, 2)$ is $\sqrt{(2-0)^2 + (2-0)^2 + (2-0)^2} = \sqrt{12} = 2\sqrt{3}$, and subtracting the radius of each circle, the distance between the spheres is $2\sqrt{3} - 2 - 1 = 2\sqrt{3} - 3$.

48. There are many different solids that fit the given description. However, any possible solid must have a circular horizontal cross-section at its top or at its base. Here we illustrate a solid with a circular base in the xy-plane. (A circular cross-section at the top results in an inverted version of the solid described below.) The vertical cross-section through the center of the base that is parallel to the xz-plane must be a square, and the vertical cross-section parallel to the yz-plane (perpendicular to the square) through the center of the base must be a triangle with two vertices on the circle and the third vertex at the center of the top side of the square. (See the figure.)



[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

The solid can include any additional points that do not extend beyond these three "silhouettes" when viewed from directions parallel to the coordinate axes. One possibility shown here is to draw the circular base and the vertical square first. Then draw a surface formed by line segments parallel to the yz-plane that connect the top of the square to the circle.



Problem 8 in the Problems Plus section at the end of the chapter illustrates another possible solid.

12.2 Vectors

1. (a) The cost of a theater ticket is a scalar, because it has only magnitude.

   (b) The current in a river is a vector, because it has both magnitude (the speed of the current) and direction at any given location.

   (c) If we assume that the initial path is linear, the initial flight path from Houston to Dallas is a vector, because it has both magnitude (distance) and direction.

   (d) The population of the world is a scalar, because it has only magnitude.

2. If the initial point of the vector $\langle 4, 7\rangle$ is placed at the origin, then $\langle 4, 7\rangle$ is the position vector of the point $(4, 7)$.



3. Vectors are equal when they share the same length and direction (but not necessarily location). Using the symmetry of the parallelogram as a guide, we see that $\overrightarrow{AB} = \overrightarrow{DC}$, $\overrightarrow{DA} = \overrightarrow{CB}$, $\overrightarrow{DE} = \overrightarrow{EB}$, and $\overrightarrow{EA} = \overrightarrow{CE}$.

4. (a) The initial point of $\overrightarrow{BC}$ is positioned at the terminal point of $\overrightarrow{AB}$, so by the Triangle Law the sum $\overrightarrow{AB} + \overrightarrow{BC}$ is the vector with initial point A and terminal point C, namely $\overrightarrow{AC}$.

   (b) By the Triangle Law, $\overrightarrow{CD} + \overrightarrow{DB}$ is the vector with initial point C and terminal point B, namely $\overrightarrow{CB}$.

   (c) First we consider $\overrightarrow{DB} - \overrightarrow{AB}$ as $\overrightarrow{DB} + \left(-\overrightarrow{AB}\right)$. Then since $-\overrightarrow{AB}$ has the same length as $\overrightarrow{AB}$ but points in the opposite direction, we have $-\overrightarrow{AB} = \overrightarrow{BA}$ and so $\overrightarrow{DB} - \overrightarrow{AB} = \overrightarrow{DB} + \overrightarrow{BA} = \overrightarrow{DA}$.

   (d) We use the Triangle Law twice: $\overrightarrow{DC} + \overrightarrow{CA} + \overrightarrow{AB} = \left(\overrightarrow{DC} + \overrightarrow{CA}\right) + \overrightarrow{AB} = \overrightarrow{DA} + \overrightarrow{AB} = \overrightarrow{DB}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. (a)



(b)



(c)

(d)



(e)



(f)



6. (a)



(b)



(c)



(d)



(e)



(f)



7. Because the tail of $\mathbf{d}$ is the midpoint of QR we have $\overrightarrow{QR} = 2\mathbf{d}$, and by the Triangle Law, $\mathbf{a} + 2\mathbf{d} = \mathbf{b} \quad \Rightarrow \quad 2\mathbf{d} = \mathbf{b} - \mathbf{a} \quad \Rightarrow \quad \mathbf{d} = \frac{1}{2}(\mathbf{b} - \mathbf{a}) = \frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a}$. Again by the Triangle Law we have $\mathbf{c} + \mathbf{d} = \mathbf{b}$ so $\mathbf{c} = \mathbf{b} - \mathbf{d} = \mathbf{b} - \left(\frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a}\right) = \frac{1}{2}\mathbf{a} + \frac{1}{2}\mathbf{b}$.

8. We are given $\mathbf{u} + \mathbf{v} + \mathbf{w} = \mathbf{0}$, so $\mathbf{w} = (-\mathbf{u}) + (-\mathbf{v})$. (See the figure.)



Vectors $-\mathbf{u}$, $-\mathbf{v}$, and $\mathbf{w}$ form a right triangle, so from the Pythagorean Theorem we have $|-\mathbf{u}|^2 + |-\mathbf{v}|^2 = |\mathbf{w}|^2$. But $|-\mathbf{u}| = |\mathbf{u}| = 1$ and $|-\mathbf{v}| = |\mathbf{v}| = 1$ so $|\mathbf{w}| = \sqrt{|-\mathbf{u}|^2 + |-\mathbf{v}|^2} = \sqrt{2}$.

9. $\mathbf{a} = \langle 1 - (-2), 2 - 1 \rangle = \langle 3, 1 \rangle$



10. $\mathbf{a} = \langle -3 - (-5), 3 - (-1) \rangle = \langle 2, 4 \rangle$



11. $\mathbf{a} = \langle 2 - 3, 3 - (-1) \rangle = \langle -1, 4 \rangle$



12. $\mathbf{a} = \langle 1 - 3, 0 - 2 \rangle = \langle -2, -2 \rangle$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

13. $\mathbf{a} = \langle 2-0, 3-3, -1-1\rangle = \langle 2, 0, -2\rangle$



14. $\mathbf{a} = \langle 3-0, 4-6, 4-(-1)\rangle = \langle 3, -2, 5\rangle$



15. $\langle -1, 4\rangle + \langle 6, -2\rangle = \langle -1+6, 4+(-2)\rangle = \langle 5, 2\rangle$



16. $\langle 3, -1\rangle + \langle -1, 5\rangle = \langle 3+(-1), -1+5\rangle = \langle 2, 4\rangle$



17. $\langle 3, 0, 1\rangle + \langle 0, 8, 0\rangle = \langle 3+0, 0+8, 1+0\rangle$
$= \langle 3, 8, 1\rangle$



18. $\langle 1, 3, -2\rangle + \langle 0, 0, 6\rangle = \langle 1+0, 3+0, -2+6\rangle$
$= \langle 1, 3, 4\rangle$



19. $\mathbf{a}+\mathbf{b} = \langle -3, 4\rangle + \langle 9, -1\rangle = \langle -3+9, 4+(-1)\rangle = \langle 6, 3\rangle$

$4\mathbf{a}+2\mathbf{b} = 4\langle -3, 4\rangle + 2\langle 9, -1\rangle = \langle -12, 16\rangle + \langle 18, -2\rangle = \langle 6, 14\rangle$

$|\mathbf{a}| = \sqrt{(-3)^2+4^2} = \sqrt{25} = 5$

$|\mathbf{a}-\mathbf{b}| = |\langle -3-9, 4-(-1)\rangle| = |\langle -12, 5\rangle| = \sqrt{(-12)^2+5^2} = \sqrt{169} = 13$

20. $\mathbf{a}+\mathbf{b} = (5\mathbf{i}+3\mathbf{j}) + (-\mathbf{i}-2\mathbf{j}) = 4\mathbf{i}+\mathbf{j}$

$4\mathbf{a}+2\mathbf{b} = 4(5\mathbf{i}+3\mathbf{j}) + 2(-\mathbf{i}-2\mathbf{j}) = 20\mathbf{i}+12\mathbf{j}-2\mathbf{i}-4\mathbf{j} = 18\mathbf{i}+8\mathbf{j}$

$|\mathbf{a}| = \sqrt{5^2+3^2} = \sqrt{34}$

$|\mathbf{a}-\mathbf{b}| = |(5\mathbf{i}+3\mathbf{j}) - (-\mathbf{i}-2\mathbf{j})| = |6\mathbf{i}+5\mathbf{j}| = \sqrt{6^2+5^2} = \sqrt{61}$

21. $\mathbf{a}+\mathbf{b} = (4\mathbf{i}-3\mathbf{j}+2\mathbf{k}) + (2\mathbf{i}-4\mathbf{k}) = 6\mathbf{i}-3\mathbf{j}-2\mathbf{k}$

$4\mathbf{a}+2\mathbf{b} = 4(4\mathbf{i}-3\mathbf{j}+2\mathbf{k}) + 2(2\mathbf{i}-4\mathbf{k}) = 16\mathbf{i}-12\mathbf{j}+8\mathbf{k}+4\mathbf{i}-8\mathbf{k} = 20\mathbf{i}-12\mathbf{j}$

$|\mathbf{a}| = \sqrt{4^2+(-3)^2+2^2} = \sqrt{29}$

$|\mathbf{a}-\mathbf{b}| = |(4\mathbf{i}-3\mathbf{j}+2\mathbf{k}) - (2\mathbf{i}-4\mathbf{k})| = |2\mathbf{i}-3\mathbf{j}+6\mathbf{k}| = \sqrt{2^2+(-3)^2+6^2} = \sqrt{49} = 7$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

22. $\mathbf{a}+\mathbf{b}=\langle 8+5,1+(-2),-4+1\rangle=\langle 13,-1,-3\rangle$

$4\mathbf{a}+2\mathbf{b}=4\langle 8,1,-4\rangle+2\langle 5,-2,1\rangle=\langle 32,4,-16\rangle+\langle 10,-4,2\rangle=\langle 42,0,-14\rangle$

$|\mathbf{a}|=\sqrt{8^2+1^2+(-4)^2}=\sqrt{81}=9$

$|\mathbf{a}-\mathbf{b}|=|\langle 8-5,1-(-2),-4-1\rangle|=|\langle 3,3,-5\rangle|=\sqrt{3^2+3^2+(-5)^2}=\sqrt{43}$

23. The vector $\langle 6,-2\rangle$ has length $|\langle 6,-2\rangle|=\sqrt{6^2+(-2)^2}=\sqrt{40}=2\sqrt{10}$, so by Equation 4 the unit vector with the same direction is $\dfrac{1}{2\sqrt{10}}\langle 6,-2\rangle=\left\langle \dfrac{3}{\sqrt{10}},-\dfrac{1}{\sqrt{10}}\right\rangle$.

24. The vector $-5\,\mathbf{i}+3\,\mathbf{j}-\mathbf{k}$ has length $|-5\,\mathbf{i}+3\,\mathbf{j}-\mathbf{k}|=\sqrt{(-5)^2+3^2+(-1)^2}=\sqrt{35}$, so by Equation 4 the unit vector with the same direction is $\dfrac{1}{\sqrt{35}}(-5\,\mathbf{i}+3\,\mathbf{j}-\mathbf{k})=-\dfrac{5}{\sqrt{35}}\,\mathbf{i}+\dfrac{3}{\sqrt{35}}\,\mathbf{j}-\dfrac{1}{\sqrt{35}}\,\mathbf{k}$.

25. The vector $8\,\mathbf{i}-\mathbf{j}+4\,\mathbf{k}$ has length $|8\,\mathbf{i}-\mathbf{j}+4\,\mathbf{k}|=\sqrt{8^2+(-1)^2+4^2}=\sqrt{81}=9$, so by Equation 4 the unit vector with the same direction is $\frac{1}{9}(8\,\mathbf{i}-\mathbf{j}+4\,\mathbf{k})=\frac{8}{9}\,\mathbf{i}-\frac{1}{9}\,\mathbf{j}+\frac{4}{9}\,\mathbf{k}$.

26. $|\langle 6,2,-3\rangle|=\sqrt{6^2+2^2+(-3)^2}=\sqrt{49}=7$, so a unit vector in the direction of $\langle 6,2,-3\rangle$ is $\mathbf{u}=\frac{1}{7}\langle 6,2,-3\rangle$.
A vector in the same direction but with length 4 is $4\mathbf{u}=4\cdot\frac{1}{7}\langle 6,2,-3\rangle=\left\langle \frac{24}{7},\frac{8}{7},-\frac{12}{7}\right\rangle$.

27.



From the figure, we see that $\tan\theta=\dfrac{\sqrt{3}}{1}=\sqrt{3}\quad\Rightarrow\quad\theta=60°$.

28.



From the figure we see that $\tan\theta=\frac{6}{8}=\frac{3}{4}$, so $\theta=\tan^{-1}\left(\frac{3}{4}\right)\approx 36.9°$.

29. From the figure, we see that the x-component of $\mathbf{v}$ is $v_1=|\mathbf{v}|\cos(\pi/3)=4\cdot\frac{1}{2}=2$ and the y-component is $v_2=|\mathbf{v}|\sin(\pi/3)=4\cdot\frac{\sqrt{3}}{2}=2\sqrt{3}$. Thus $\mathbf{v}=\langle v_1,v_2\rangle=\langle 2,2\sqrt{3}\,\rangle$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

30. From the figure, we see that the horizontal component of the force $\mathbf{F}$ is $|\mathbf{F}|\cos 38^\circ = 50\cos 38^\circ \approx 39.4$ N, and the vertical component is $|\mathbf{F}|\sin 38^\circ = 50\sin 38^\circ \approx 30.8$ N.



31. The velocity vector $\mathbf{v}$ makes an angle of 40° with the horizontal and has magnitude equal to the speed at which the football was thrown. From the figure, we see that the horizontal component of $\mathbf{v}$ is $|\mathbf{v}|\cos 40^\circ = 60\cos 40^\circ \approx 45.96$ ft/s and the vertical component is $|\mathbf{v}|\sin 40^\circ = 60\sin 40^\circ \approx 38.57$ ft/s.



32. The given force vectors can be expressed in terms of their horizontal and vertical components as $20\cos 45^\circ\,\mathbf{i} + 20\sin 45^\circ\,\mathbf{j} = 10\sqrt{2}\,\mathbf{i} + 10\sqrt{2}\,\mathbf{j}$ and $16\cos 30^\circ\,\mathbf{i} - 16\sin 30^\circ\,\mathbf{j} = 8\sqrt{3}\,\mathbf{i} - 8\,\mathbf{j}$. The resultant force $\mathbf{F}$ is the sum of these two vectors: $\mathbf{F} = \left(10\sqrt{2} + 8\sqrt{3}\right)\mathbf{i} + \left(10\sqrt{2} - 8\right)\mathbf{j} \approx 28.00\,\mathbf{i} + 6.14\,\mathbf{j}$. Then we have $|\mathbf{F}| \approx \sqrt{(28.00)^2 + (6.14)^2} \approx 28.7$ lb and, letting θ be the angle $\mathbf{F}$ makes with the positive x-axis,

$$\tan\theta = \frac{10\sqrt{2} - 8}{10\sqrt{2} + 8\sqrt{3}} \quad \Rightarrow \quad \theta = \tan^{-1}\left(\frac{10\sqrt{2} - 8}{10\sqrt{2} + 8\sqrt{3}}\right) \approx 12.4^\circ.$$

33. The given force vectors can be expressed in terms of their horizontal and vertical components as $-300\,\mathbf{i}$ and $200\cos 60^\circ\,\mathbf{i} + 200\sin 60^\circ\,\mathbf{j} = 200\left(\frac{1}{2}\right)\mathbf{i} + 200\left(\frac{\sqrt{3}}{2}\right)\mathbf{j} = 100\,\mathbf{i} + 100\sqrt{3}\,\mathbf{j}$. The resultant force $\mathbf{F}$ is the sum of these two vectors: $\mathbf{F} = (-300 + 100)\,\mathbf{i} + \left(0 + 100\sqrt{3}\right)\mathbf{j} = -200\,\mathbf{i} + 100\sqrt{3}\,\mathbf{j}$. Then we have $|\mathbf{F}| \approx \sqrt{(-200)^2 + \left(100\sqrt{3}\right)^2} = \sqrt{70{,}000} = 100\sqrt{7} \approx 264.6$ N. Let θ be the angle $\mathbf{F}$ makes with the positive x-axis. Then $\tan\theta = \dfrac{100\sqrt{3}}{-200} = -\dfrac{\sqrt{3}}{2}$ and the terminal point of $\mathbf{F}$ lies in the second quadrant, so

$$\theta = \tan^{-1}\left(-\frac{\sqrt{3}}{2}\right) + 180^\circ \approx -40.9^\circ + 180^\circ = 139.1^\circ.$$

34. Set up the coordinate axes so that north is the positive y-direction, and east is the positive x-direction. The wind is blowing at 50 km/h from the direction N45°W, so that its velocity vector is 50 km/h S45°E, which can be written as $\mathbf{v}_{\text{wind}} = 50(\cos 45^\circ\,\mathbf{i} - \sin 45^\circ\,\mathbf{j})$. With respect to the still air, the velocity vector of the plane is 250 km/h N 60°E, or equivalently $\mathbf{v}_{\text{plane}} = 250(\cos 30^\circ\,\mathbf{i} + \sin 30^\circ\,\mathbf{j})$. The velocity of the plane relative to the ground is

$$\begin{aligned}\mathbf{v} &= \mathbf{v}_{\text{plane}} + \mathbf{v}_{\text{wind}} \\ &= (250\cos 30^\circ + 50\cos 45^\circ)\,\mathbf{i} + (250\sin 30^\circ - 50\sin 45^\circ)\,\mathbf{j} \\ &= \left(125\sqrt{3} + 25\sqrt{2}\right)\mathbf{i} + \left(125 - 25\sqrt{2}\right)\mathbf{j} \\ &\approx 251.9\,\mathbf{i} + 89.6\,\mathbf{j}\end{aligned}$$



(See the figure.) The ground speed is $|\mathbf{v}| \approx \sqrt{(251.9)^2 + (89.6)^2} \approx 267$ km/h. The angle the velocity vector makes with the x-axis is $\theta \approx \tan^{-1}\left(\frac{89.6}{251.9}\right) \approx 20^\circ$. Therefore, the true course of the plane is about N$(90 - 20)^\circ$E = N 70°E.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. With respect to the water's surface, the woman's velocity is the vector sum of the velocity of the ship with respect to the water, and the woman's velocity with respect to the ship. If we let north be the positive y-direction, then $\mathbf{v} = \langle 0, 22\rangle + \langle -3, 0\rangle = \langle -3, 22\rangle$. The woman's speed is $|\mathbf{v}| = \sqrt{9 + 484} \approx 22.2$ mi/h. The vector $\mathbf{v}$ makes an angle θ with the east, where $\theta = \tan^{-1}\left(\frac{22}{-3}\right) \approx 98°$. Therefore, the woman's direction is about $\mathrm{N}(98 - 90)°\mathrm{W} = \mathrm{N}8°\mathrm{W}$.

36. Let $\mathbf{T}_1$ and $\mathbf{T}_2$ be the tension vectors corresponding to the support cables as shown in the figure. In terms of vertical and horizontal components,



$$\mathbf{T}_1 = |\mathbf{T}_1|\cos 60°\,\mathbf{i} + |\mathbf{T}_1|\sin 60°\,\mathbf{j} = \frac{1}{2}|\mathbf{T}_1|\,\mathbf{i} + \frac{\sqrt{3}}{2}|\mathbf{T}_1|\,\mathbf{j}$$

$$\mathbf{T}_2 = -|\mathbf{T}_2|\cos 60°\,\mathbf{i} + |\mathbf{T}_2|\sin 60°\,\mathbf{j} = -\frac{1}{2}|\mathbf{T}_2|\,\mathbf{i} + \frac{\sqrt{3}}{2}|\mathbf{T}_2|\,\mathbf{j}$$

The resultant of these tensions, $\mathbf{T}_1 + \mathbf{T}_2$, counterbalances the weight $\mathbf{w} = -500\,\mathbf{j}$. So $\mathbf{T}_1 + \mathbf{T}_2 = -\mathbf{w} = 500\,\mathbf{j} \;\Rightarrow$

$$\left(\frac{1}{2}|\mathbf{T}_1|\,\mathbf{i} + \frac{\sqrt{3}}{2}|\mathbf{T}_1|\,\mathbf{j}\right) + \left(-\frac{1}{2}|\mathbf{T}_2|\,\mathbf{i} + \frac{\sqrt{3}}{2}|\mathbf{T}_2|\,\mathbf{j}\right) = 500\,\mathbf{j}.$$

Equating x-components gives $\frac{1}{2}|\mathbf{T}_1|\,\mathbf{i} - \frac{1}{2}|\mathbf{T}_2|\,\mathbf{i} = 0$, so $|\mathbf{T}_1| = |\mathbf{T}_2|$ (as we would expect from the symmetry of the problem). Equating y-components, we have $\frac{\sqrt{3}}{2}|\mathbf{T}_1|\,\mathbf{j} + \frac{\sqrt{3}}{2}|\mathbf{T}_2|\,\mathbf{j} = \sqrt{3}\,|\mathbf{T}_1|\,\mathbf{j} = 500\,\mathbf{j} \;\Rightarrow\; |\mathbf{T}_1| = \frac{500}{\sqrt{3}}$. Thus the magnitude of each tension is $|\mathbf{T}_1| = |\mathbf{T}_2| = \frac{500}{\sqrt{3}} \approx 288.68$ lb. The tension vectors are

$$\mathbf{T}_1 = \frac{1}{2}|\mathbf{T}_1|\,\mathbf{i} + \frac{\sqrt{3}}{2}|\mathbf{T}_1|\,\mathbf{j} = \frac{250}{\sqrt{3}}\,\mathbf{i} + 250\,\mathbf{j} \approx 144.34\,\mathbf{i} + 250\,\mathbf{j} \text{ and } \mathbf{T}_2 = -\frac{250}{\sqrt{3}}\,\mathbf{i} + 250\,\mathbf{j} \approx -144.34\,\mathbf{i} + 250\,\mathbf{j}.$$

37. Call the two tension vectors $\mathbf{T}_2$ and $\mathbf{T}_3$, corresponding to the ropes of length 2 m and 3 m. In terms of vertical and horizontal components,

$$\mathbf{T}_2 = -|\mathbf{T}_2|\cos 50°\,\mathbf{i} + |\mathbf{T}_2|\sin 50°\,\mathbf{j} \quad (1) \qquad \text{and} \qquad \mathbf{T}_3 = |\mathbf{T}_3|\cos 38°\,\mathbf{i} + |\mathbf{T}_3|\sin 38°\,\mathbf{j} \quad (2)$$

The resultant of these forces, $\mathbf{T}_2 + \mathbf{T}_3$, counterbalances the weight of the hoist (which is $-350\,\mathbf{j}$), so

$\mathbf{T}_2 + \mathbf{T}_3 = 350\,\mathbf{j} \;\Rightarrow$

$(-|\mathbf{T}_2|\cos 50° + |\mathbf{T}_3|\cos 38°)\,\mathbf{i} + (|\mathbf{T}_2|\sin 50° + |\mathbf{T}_3|\sin 38°)\,\mathbf{j} = 350\,\mathbf{j}$. Equating components, we have

$-|\mathbf{T}_2|\cos 50° + |\mathbf{T}_3|\cos 38° = 0 \;\Rightarrow\; |\mathbf{T}_2| = |\mathbf{T}_3|\dfrac{\cos 38°}{\cos 50°}$ and

$|\mathbf{T}_2|\sin 50° + |\mathbf{T}_3|\sin 38° = 350$. Substituting the first equation into the second gives

$|\mathbf{T}_3|\dfrac{\cos 38°}{\cos 50°}\sin 50° + |\mathbf{T}_3|\sin 38° = 350 \;\Rightarrow\; |\mathbf{T}_3|(\cos 38°\tan 50° + \sin 38°) = 350$, so the magnitudes of the tensions are $|\mathbf{T}_3| = \dfrac{350}{\cos 38°\tan 50° + \sin 38°} \approx 225.11$ N and $|\mathbf{T}_2| = |\mathbf{T}_3|\dfrac{\cos 38°}{\cos 50°} \approx 275.97$ N. Finally, from (1) and (2), the tension vectors are $\mathbf{T}_2 \approx -177.39\,\mathbf{i} + 211.41\,\mathbf{j}$ and $\mathbf{T}_3 \approx 177.39\,\mathbf{i} + 138.59\,\mathbf{j}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

38. We can consider the weight of the chain to be concentrated at its midpoint. The forces acting on the chain then are the tension vectors $\mathbf{T}_1$, $\mathbf{T}_2$ in each end of the chain and the weight $\mathbf{w}$, as shown in the figure. We know $|\mathbf{T}_1| = |\mathbf{T}_2| = 25$ N so, in terms of vertical and horizontal components, we have



$$\mathbf{T}_1 = -25\cos 37°\,\mathbf{i} + 25\sin 37°\,\mathbf{j} \qquad \mathbf{T}_2 = 25\cos 37°\,\mathbf{i} + 25\sin 37°\,\mathbf{j}$$

The resultant vector $\mathbf{T}_1 + \mathbf{T}_2$ of the tensions counterbalances the weight $\mathbf{w}$, giving $\mathbf{T}_1 + \mathbf{T}_2 = -\mathbf{w}$. Since $\mathbf{w} = -|\mathbf{w}|\,\mathbf{j}$, we have $(-25\cos 37°\,\mathbf{i} + 25\sin 37°\,\mathbf{j}) + (25\cos 37°\,\mathbf{i} + 25\sin 37°\,\mathbf{j}) = |\mathbf{w}|\,\mathbf{j} \;\Rightarrow\; 50\sin 37°\,\mathbf{j} = |\mathbf{w}|\,\mathbf{j} \;\Rightarrow$ $|\mathbf{w}| = 50\sin 37° \approx 30.1$. So the weight is 30.1 N, and since $w = mg$, the mass is $\frac{30.1}{9.8} \approx 3.07$ kg.

39. (a) Set up coordinate axes so that the boatman is at the origin, the canal is bordered by the y-axis and the line $x = 3$, and the current flows in the negative y-direction. The boatman wants to reach the point $(3, 2)$. Let θ be the angle, measured from the positive y-axis, in the direction he should steer. (See the figure.)



In still water, the boat has velocity $\mathbf{v}_b = \langle 13\sin\theta, 13\cos\theta\rangle$ and the velocity of the current is $\mathbf{v}_c = \langle 0, -3.5\rangle$, so the true path of the boat is determined by the velocity vector $\mathbf{v} = \mathbf{v}_b + \mathbf{v}_c = \langle 13\sin\theta, 13\cos\theta - 3.5\rangle$. Let t be the time (in hours) after the boat departs; then the position of the boat at time t is given by $t\mathbf{v}$ and the boat crosses the canal when $t\mathbf{v} = \langle 13\sin\theta, 13\cos\theta - 3.5\rangle\, t = \langle 3, 2\rangle$. Thus $13(\sin\theta)t = 3 \;\Rightarrow\; t = \dfrac{3}{13\sin\theta}$ and $(13\cos\theta - 3.5)\,t = 2$.

Substituting gives $(13\cos\theta - 3.5)\,\dfrac{3}{13\sin\theta} = 2 \;\Rightarrow\; 39\cos\theta - 10.5 = 26\sin\theta$ **(1)**. Squaring both sides, we have

$$1521\cos^2\theta - 819\cos\theta + 110.25 = 676\sin^2\theta = 676\left(1 - \cos^2\theta\right)$$

$$2197\cos^2\theta - 819\cos\theta - 565.75 = 0$$

The quadratic formula gives

$$\cos\theta = \frac{819 \pm \sqrt{(-819)^2 - 4(2197)(-565.75)}}{2(2197)}$$

$$= \frac{819 \pm \sqrt{5{,}612{,}572}}{4394} \approx 0.72699 \text{ or } -0.35421$$

The acute value for θ is approximately $\cos^{-1}(0.72699) \approx 43.4°$. Thus the boatman should steer in the direction that is $43.4°$ from the bank, toward upstream.

Alternate solution: We could solve **(1)** graphically by plotting $y = 39\cos\theta - 10.5$ and $y = 26\sin\theta$ on a graphing device and finding the appoximate intersection point $(0.757, 17.85)$. Thus $\theta \approx 0.757$ radians or equivalently $43.4°$.

(b) From part (a) we know the trip is completed when $t = \dfrac{3}{13\sin\theta}$. But $\theta \approx 43.4°$, so the time required is approximately $\dfrac{3}{13\sin 43.4°} \approx 0.336$ hours or 20.2 minutes.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. Let $\mathbf{v}_1$, $\mathbf{v}_2$, and $\mathbf{v}_3$ be the force vectors where $|\mathbf{v}_1| = 25$, $|\mathbf{v}_2| = 12$, and $|\mathbf{v}_3| = 4$. Set up coordinate axes so that the object is at the origin and $\mathbf{v}_1$, $\mathbf{v}_2$ lie in the xy-plane. We can position the vectors so that $\mathbf{v}_1 = 25\,\mathbf{i}$, $\mathbf{v}_2 = 12\cos 100^\circ\,\mathbf{i} + 12\sin 100^\circ\,\mathbf{j}$, and $\mathbf{v}_3 = 4\,\mathbf{k}$. The magnitude of a force that counterbalances the three given forces must match the magnitude of the resultant force. We have $\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3 = (25 + 12\cos 100^\circ)\,\mathbf{i} + 12\sin 100^\circ\,\mathbf{j} + 4\,\mathbf{k}$, so the counterbalancing force must have magnitude $|\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3| = \sqrt{(25 + 12\cos 100^\circ)^2 + (12\sin 100^\circ)^2 + 4^2} \approx 26.1$ N.

41. The slope of the tangent line to the graph of $y = x^2$ at the point $(2, 4)$ is

$$\left.\frac{dy}{dx}\right|_{x=2} = \left.2x\right|_{x=2} = 4$$

and a parallel vector is $\mathbf{i} + 4\,\mathbf{j}$ which has length $|\mathbf{i} + 4\,\mathbf{j}| = \sqrt{1^2 + 4^2} = \sqrt{17}$, so unit vectors parallel to the tangent line are $\pm\frac{1}{\sqrt{17}}(\mathbf{i} + 4\,\mathbf{j})$.

42. (a) The slope of the tangent line to the graph of $y = 2\sin x$ at the point $(\pi/6, 1)$ is

$$\left.\frac{dy}{dx}\right|_{x=\pi/6} = \left.2\cos x\right|_{x=\pi/6} = 2\cdot\frac{\sqrt{3}}{2} = \sqrt{3}$$

and a parallel vector is $\mathbf{i} + \sqrt{3}\,\mathbf{j}$ which has length $\left|\mathbf{i} + \sqrt{3}\,\mathbf{j}\right| = \sqrt{1^2 + \left(\sqrt{3}\right)^2} = \sqrt{4} = 2$, so unit vectors parallel to the tangent line are $\pm\frac{1}{2}\left(\mathbf{i} + \sqrt{3}\,\mathbf{j}\right)$.

(b) The slope of the tangent line is $\sqrt{3}$, so the slope of a line perpendicular to the tangent line is $-\frac{1}{\sqrt{3}}$ and a vector in this direction is $\sqrt{3}\,\mathbf{i} - \mathbf{j}$. Since $\left|\sqrt{3}\,\mathbf{i} - \mathbf{j}\right| = \sqrt{\left(\sqrt{3}\right)^2 + (-1)^2} = 2$, unit vectors perpendicular to the tangent line are $\pm\frac{1}{2}\left(\sqrt{3}\,\mathbf{i} - \mathbf{j}\right)$.

(c)



43. By the Triangle Law, $\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC}$. Then $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = \overrightarrow{AC} + \overrightarrow{CA}$, but $\overrightarrow{AC} + \overrightarrow{CA} = \overrightarrow{AC} + \left(-\overrightarrow{AC}\right) = \mathbf{0}$. So $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = \mathbf{0}$.

44. $\overrightarrow{AC} = \frac{1}{3}\overrightarrow{AB}$ and $\overrightarrow{BC} = \frac{2}{3}\overrightarrow{BA}$. $\mathbf{c} = \overrightarrow{OA} + \overrightarrow{AC} = \mathbf{a} + \frac{1}{3}\overrightarrow{AB} \;\Rightarrow\; \overrightarrow{AB} = 3\,\mathbf{c} - 3\,\mathbf{a}$. $\mathbf{c} = \overrightarrow{OB} + \overrightarrow{BC} = \overrightarrow{OA} + \frac{2}{3}\overrightarrow{BA} \;\Rightarrow\; \overrightarrow{BA} = \frac{3}{2}\,\mathbf{c} - \frac{3}{2}\,\mathbf{b}$. $\overrightarrow{BA} = \overrightarrow{AB}$, so $\frac{3}{2}\,\mathbf{c} - \frac{3}{2}\,\mathbf{b} = 3\,\mathbf{a} - 3\,\mathbf{c} \;\Leftrightarrow\; \mathbf{c} + 2\,\mathbf{c} = 2\,\mathbf{a} + \mathbf{b} \;\Leftrightarrow\; \mathbf{c} = \frac{2}{3}\,\mathbf{a} + \frac{1}{3}\,\mathbf{b}$.

45. (a), (b)



(c) From the sketch, we estimate that $s \approx 1.3$ and $t \approx 1.6$.

(d) $\mathbf{c} = s\,\mathbf{a} + t\,\mathbf{b} \;\Leftrightarrow\; 7 = 3s + 2t$ and $1 = 2s - t$.
Solving these equations gives $s = \frac{9}{7}$ and $t = \frac{11}{7}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

46. Draw $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ emanating from the origin. Extend $\mathbf{a}$ and $\mathbf{b}$ to form lines A and B, and draw lines A' and B' parallel to these two lines through the terminal point of $\mathbf{c}$. Since $\mathbf{a}$ and $\mathbf{b}$ are not parallel, A and B' must meet (at P), and A' and B must also meet (at Q). Now we see that $\overrightarrow{OP} + \overrightarrow{OQ} = \mathbf{c}$, so if



$s = \dfrac{|\overrightarrow{OP}|}{|\mathbf{a}|}$ $\left(\text{or its negative, if } \mathbf{a} \text{ points in the direction opposite } \overrightarrow{OP}\right)$ and $t = \dfrac{|\overrightarrow{OQ}|}{|\mathbf{b}|}$ (or its negative, as in the diagram), then $\mathbf{c} = s\mathbf{a} + t\mathbf{b}$, as required.

Argument using components: Since $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ all lie in the same plane, we can consider them to be vectors in two dimensions. Let $\mathbf{a} = \langle a_1, a_2 \rangle$, $\mathbf{b} = \langle b_1, b_2 \rangle$, and $\mathbf{c} = \langle c_1, c_2 \rangle$. We need $sa_1 + tb_1 = c_1$ and $sa_2 + tb_2 = c_2$. Multiplying the first equation by a_2 and the second by a_1 and subtracting, we get $t = \dfrac{c_2a_1 - c_1a_2}{b_2a_1 - b_1a_2}$. Similarly $s = \dfrac{b_2c_1 - b_1c_2}{b_2a_1 - b_1a_2}$. Since $\mathbf{a} \neq \mathbf{0}$ and $\mathbf{b} \neq \mathbf{0}$ and $\mathbf{a}$ is not a scalar multiple of $\mathbf{b}$, the denominator is not zero.

47. $|\mathbf{r} - \mathbf{r}_0|$ is the distance between the points (x, y, z) and (x_0, y_0, z_0), so the set of points is a sphere with radius 1 and center (x_0, y_0, z_0).

Alternate method: $|\mathbf{r} - \mathbf{r}_0| = 1 \quad \Leftrightarrow \quad \sqrt{(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2} = 1 \quad \Leftrightarrow$
$(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = 1$, which is the equation of a sphere with radius 1 and center (x_0, y_0, z_0).

48. Let P_1 and P_2 be the points with position vectors $\mathbf{r}_1$ and $\mathbf{r}_2$ respectively. Then $|\mathbf{r} - \mathbf{r}_1| + |\mathbf{r} - \mathbf{r}_2|$ is the sum of the distances from (x, y) to P_1 and P_2. Since this sum is constant, the set of points (x, y) represents an ellipse with foci P_1 and P_2. The condition $k > |\mathbf{r}_1 - \mathbf{r}_2|$ assures us that the ellipse is not degenerate.

49.
$$\begin{aligned}
\mathbf{a} + (\mathbf{b} + \mathbf{c}) &= \langle a_1, a_2 \rangle + (\langle b_1, b_2 \rangle + \langle c_1, c_2 \rangle) = \langle a_1, a_2 \rangle + \langle b_1 + c_1, b_2 + c_2 \rangle \\
&= \langle a_1 + b_1 + c_1, a_2 + b_2 + c_2 \rangle = \langle (a_1 + b_1) + c_1, (a_2 + b_2) + c_2 \rangle \\
&= \langle a_1 + b_1, a_2 + b_2 \rangle + \langle c_1, c_2 \rangle = (\langle a_1, a_2 \rangle + \langle b_1, b_2 \rangle) + \langle c_1, c_2 \rangle \\
&= (\mathbf{a} + \mathbf{b}) + \mathbf{c}
\end{aligned}$$

50. *Algebraically:*
$$\begin{aligned}
c(\mathbf{a} + \mathbf{b}) &= c\left(\langle a_1, a_2, a_3 \rangle + \langle b_1, b_2, b_3 \rangle\right) = c\left\langle a_1 + b_1, a_2 + b_2, a_3 + b_3 \right\rangle \\
&= \left\langle c\left(a_1 + b_1\right), c\left(a_2 + b_2\right), c\left(a_3 + b_3\right) \right\rangle = \langle ca_1 + cb_1, ca_2 + cb_2, ca_3 + cb_3 \rangle \\
&= \langle ca_1, ca_2, ca_3 \rangle + \langle cb_1, cb_2, cb_3 \rangle = c\,\mathbf{a} + c\,\mathbf{b}
\end{aligned}$$

Geometrically:

According to the Triangle Law, if $\mathbf{a} = \overrightarrow{PQ}$ and $\mathbf{b} = \overrightarrow{QR}$, then $\mathbf{a} + \mathbf{b} = \overrightarrow{PR}$. Construct triangle PST as shown so that $\overrightarrow{PS} = c\,\mathbf{a}$ and $\overrightarrow{ST} = c\,\mathbf{b}$. (We have drawn the case where $c > 1$.) By the Triangle Law, $\overrightarrow{PT} = c\,\mathbf{a} + c\,\mathbf{b}$. But triangle PQR and triangle PST are similar triangles because $c\,\mathbf{b}$ is parallel to $\mathbf{b}$. Therefore, $\overrightarrow{PR}$ and $\overrightarrow{PT}$ are parallel and, in fact, $\overrightarrow{PT} = c\overrightarrow{PR}$. Thus, $c\,\mathbf{a} + c\,\mathbf{b} = c(\mathbf{a} + \mathbf{b})$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. Consider triangle ABC, where D and E are the midpoints of AB and BC. We know that $\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC}$ **(1)** and $\overrightarrow{DB} + \overrightarrow{BE} = \overrightarrow{DE}$ **(2)**. However, $\overrightarrow{DB} = \frac{1}{2}\overrightarrow{AB}$, and $\overrightarrow{BE} = \frac{1}{2}\overrightarrow{BC}$. Substituting these expressions for $\overrightarrow{DB}$ and $\overrightarrow{BE}$ into **(2)** gives $\frac{1}{2}\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC} = \overrightarrow{DE}$. Comparing this with **(1)** gives $\overrightarrow{DE} = \frac{1}{2}\overrightarrow{AC}$. Therefore $\overrightarrow{AC}$ and $\overrightarrow{DE}$ are parallel and $\left|\overrightarrow{DE}\right| = \frac{1}{2}\left|\overrightarrow{AC}\right|$.

52. The question states that the light ray strikes all three mirrors, so it is not parallel to any of them and $a_1 \neq 0$, $a_2 \neq 0$ and $a_3 \neq 0$. Let $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$, as in the diagram. We can let $|\mathbf{b}| = |\mathbf{a}|$, since only its direction is important. Then

$$\frac{|b_2|}{|\mathbf{b}|} = \sin\theta = \frac{|a_2|}{|\mathbf{a}|} \quad \Rightarrow \quad |b_2| = |a_2|.$$



From the diagram $b_2\,\mathbf{j}$ and $a_2\,\mathbf{j}$ point in opposite directions, so $b_2 = -a_2$. $|AB| = |BC|$, so $|b_3| = \sin\phi\ |BC| = \sin\phi\ |AB| = |a_3|$, and $|b_1| = \cos\phi\ |BC| = \cos\phi\ |AB| = |a_1|$. $b_3\,\mathbf{k}$ and $a_3\,\mathbf{k}$ have the same direction, as do $b_1\,\mathbf{i}$ and $a_1\,\mathbf{i}$, so $\mathbf{b} = \langle a_1, -a_2, a_3 \rangle$. When the ray hits the other mirrors, similar arguments show that these reflections will reverse the signs of the other two coordinates, so the final reflected ray will be $\langle -a_1, -a_2, -a_3 \rangle = -\mathbf{a}$, which is parallel to $\mathbf{a}$.

12.3 The Dot Product

1. (a) $\mathbf{a} \cdot \mathbf{b}$ is a scalar, and the dot product is defined only for vectors, so $(\mathbf{a} \cdot \mathbf{b}) \cdot \mathbf{c}$ has no meaning.

(b) $(\mathbf{a} \cdot \mathbf{b})\,\mathbf{c}$ is a scalar multiple of a vector, so it does have meaning.

(c) Both $|\mathbf{a}|$ and $\mathbf{b} \cdot \mathbf{c}$ are scalars, so $|\mathbf{a}|\,(\mathbf{b} \cdot \mathbf{c})$ is an ordinary product of real numbers, and has meaning.

(d) Both $\mathbf{a}$ and $\mathbf{b} + \mathbf{c}$ are vectors, so the dot product $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c})$ has meaning.

(e) $\mathbf{a} \cdot \mathbf{b}$ is a scalar, but $\mathbf{c}$ is a vector, and so the two quantities cannot be added and $\mathbf{a} \cdot \mathbf{b} + \mathbf{c}$ has no meaning.

(f) $|\mathbf{a}|$ is a scalar, and the dot product is defined only for vectors, so $|\mathbf{a}| \cdot (\mathbf{b} + \mathbf{c})$ has no meaning.

2. $\mathbf{a} \cdot \mathbf{b} = \langle 5, -2 \rangle \cdot \langle 3, 4 \rangle = (5)(3) + (-2)(4) = 15 - 8 = 7$

3. $\mathbf{a} \cdot \mathbf{b} = \langle 1.5, 0.4 \rangle \cdot \langle -4, 6 \rangle = (1.5)(-4) + (0.4)(6) = -6 + 2.4 = -3.6$

4. $\mathbf{a} \cdot \mathbf{b} = \langle 6, -2, 3 \rangle \cdot \langle 2, 5, -1 \rangle = (6)(2) + (-2)\,(5) + (3)(-1) = 12 - 10 - 3 = -1$

5. $\mathbf{a} \cdot \mathbf{b} = \left\langle 4, 1, \frac{1}{4} \right\rangle \cdot \langle 6, -3, -8 \rangle = (4)(6) + (1)(-3) + \left(\frac{1}{4}\right)(-8) = 19$

6. $\mathbf{a} \cdot \mathbf{b} = \langle p, -p, 2p \rangle \cdot \langle 2q, q, -q \rangle = (p)(2q) + (-p)(q) + (2p)(-q) = 2pq - pq - 2pq = -pq$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

7. $\mathbf{a} \cdot \mathbf{b} = (2\mathbf{i} + \mathbf{j}) \cdot (\mathbf{i} - \mathbf{j} + \mathbf{k}) = (2)(1) + (1)(-1) + (0)(1) = 1$

8. $\mathbf{a} \cdot \mathbf{b} = (3\mathbf{i} + 2\mathbf{j} - \mathbf{k}) \cdot (4\mathbf{i} + 5\mathbf{k}) = (3)(4) + (2)(0) + (-1)(5) = 7$

9. By Theorem 3, $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}|\,|\mathbf{b}| \cos\theta = (7)(4)\cos 30^\circ = 28\left(\frac{\sqrt{3}}{2}\right) = 14\sqrt{3} \approx 24.25$.

10. By Theorem 3, $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}|\,|\mathbf{b}| \cos\theta = (80)(50)\cos\frac{3\pi}{4} = 4000\left(-\frac{\sqrt{2}}{2}\right) = -2000\sqrt{2} \approx -2828.43$.

11. $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are all unit vectors, so the triangle is an equilateral triangle. Thus the angle between $\mathbf{u}$ and $\mathbf{v}$ is 60° and $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}|\,|\mathbf{v}| \cos 60^\circ = (1)(1)\left(\frac{1}{2}\right) = \frac{1}{2}$. If $\mathbf{w}$ is moved so it has the same initial point as $\mathbf{u}$, we can see that the angle between them is 120° and we have $\mathbf{u} \cdot \mathbf{w} = |\mathbf{u}|\,|\mathbf{w}| \cos 120^\circ = (1)(1)\left(-\frac{1}{2}\right) = -\frac{1}{2}$.

12. $\mathbf{u}$ is a unit vector, so $\mathbf{w}$ is also a unit vector, and $|\mathbf{v}|$ can be determined by examining the right triangle formed by $\mathbf{u}$ and $\mathbf{v}$. Since the angle between $\mathbf{u}$ and $\mathbf{v}$ is 45°, we have $|\mathbf{v}| = |\mathbf{u}| \cos 45^\circ = \frac{\sqrt{2}}{2}$. Then $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}|\,|\mathbf{v}| \cos 45^\circ = (1)\left(\frac{\sqrt{2}}{2}\right)\frac{\sqrt{2}}{2} = \frac{1}{2}$. Since $\mathbf{u}$ and $\mathbf{w}$ are orthogonal, $\mathbf{u} \cdot \mathbf{w} = 0$.

13. (a) $\mathbf{i} \cdot \mathbf{j} = \langle 1, 0, 0\rangle \cdot \langle 0, 1, 0\rangle = (1)(0) + (0)(1) + (0)(0) = 0$. Similarly, $\mathbf{j} \cdot \mathbf{k} = (0)(0) + (1)(0) + (0)(1) = 0$ and $\mathbf{k} \cdot \mathbf{i} = (0)(1) + (0)(0) + (1)(0) = 0$.

Another method: Because $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ are mutually perpendicular, the cosine factor in each dot product (see Theorem 3) is $\cos\frac{\pi}{2} = 0$.

(b) By Property 1 of the dot product, $\mathbf{i} \cdot \mathbf{i} = |\mathbf{i}|^2 = 1^2 = 1$ since $\mathbf{i}$ is a unit vector. Similarly, $\mathbf{j} \cdot \mathbf{j} = |\mathbf{j}|^2 = 1$ and $\mathbf{k} \cdot \mathbf{k} = |\mathbf{k}|^2 = 1$.

14. The dot product $\mathbf{A} \cdot \mathbf{P}$ is

$$\begin{aligned}\langle a, b, c\rangle \cdot \langle 4, 2.5, 1\rangle &= a(4) + b(2.5) + c(1)\\ &= (\text{number of hamburgers sold})(\text{price per hamburger})\\ &\quad + (\text{number of hot dogs sold})(\text{price per hot dog})\\ &\quad + (\text{number of soft drinks sold})(\text{price per soft drink})\end{aligned}$$

so it is equal to the vendor's total revenue for that day.

15. $|\mathbf{a}| = \sqrt{4^2 + 3^2} = 5$, $|\mathbf{b}| = \sqrt{2^2 + (-1)^2} = \sqrt{5}$, and $\mathbf{a} \cdot \mathbf{b} = (4)(2) + (3)(-1) = 5$. From Corollary 6, we have $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{5}{5 \cdot \sqrt{5}} = \dfrac{1}{\sqrt{5}}$. So the angle between $\mathbf{a}$ and $\mathbf{b}$ is $\theta = \cos^{-1}\left(\dfrac{1}{\sqrt{5}}\right) \approx 63^\circ$.

16. $|\mathbf{a}| = \sqrt{(-2)^2 + 5^2} = \sqrt{29}$, $|\mathbf{b}| = \sqrt{5^2 + 12^2} = 13$, and $\mathbf{a} \cdot \mathbf{b} = (-2)(5) + (5)(12) = 50$. Using Corollary 6, we have $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{50}{\sqrt{29} \cdot 13} = \dfrac{50}{13\sqrt{29}}$ and the angle between $\mathbf{a}$ and $\mathbf{b}$ is $\theta = \cos^{-1}\left(\dfrac{50}{13\sqrt{29}}\right) \approx 44^\circ$.

17. $|\mathbf{a}| = \sqrt{1^2 + (-4)^2 + 1^2} = \sqrt{18} = 3\sqrt{2}$, $|\mathbf{b}| = \sqrt{0^2 + 2^2 + (-2)^2} = \sqrt{8} = 2\sqrt{2}$, and $\mathbf{a} \cdot \mathbf{b} = (1)(0) + (-4)(2) + (1)(-2) = -10$. From Corollary 6, we have $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{-10}{3\sqrt{2} \cdot 2\sqrt{2}} = -\dfrac{10}{12} = -\dfrac{5}{6}$ and the angle between $\mathbf{a}$ and $\mathbf{b}$ is $\theta = \cos^{-1}\left(-\frac{5}{6}\right) \approx 146^\circ$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $|\mathbf{a}| = \sqrt{(-1)^2 + 3^2 + 4^2} = \sqrt{26}$, $|\mathbf{b}| = \sqrt{5^2 + 2^2 + 1^2} = \sqrt{30}$, and $\mathbf{a} \cdot \mathbf{b} = (-1)(5) + (3)(2) + (4)(1) = 5$.

Then $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{5}{\sqrt{26} \cdot \sqrt{30}} = \dfrac{5}{\sqrt{780}} = \dfrac{5}{2\sqrt{195}}$ and $\theta = \cos^{-1}\left(\dfrac{5}{2\sqrt{195}}\right) \approx 80°$.

19. $|\mathbf{a}| = \sqrt{4^2 + (-3)^2 + 1^2} = \sqrt{26}$, $|\mathbf{b}| = \sqrt{2^2 + 0^2 + (-1)^2} = \sqrt{5}$, and $\mathbf{a} \cdot \mathbf{b} = (4)(2) + (-3)(0) + (1)(-1) = 7$.

Then $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{7}{\sqrt{26} \cdot \sqrt{5}} = \dfrac{7}{\sqrt{130}}$ and $\theta = \cos^{-1}\left(\dfrac{7}{\sqrt{130}}\right) \approx 52°$.

20. $|\mathbf{a}| = \sqrt{8^2 + (-1)^2 + 4^2} = \sqrt{81} = 9$, $|\mathbf{b}| = \sqrt{0^2 + 4^2 + 2^2} = \sqrt{20} = 2\sqrt{5}$, and $\mathbf{a} \cdot \mathbf{b} = (8)(0) + (-1)(4) + (4)(2) = 4$.

Then $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{4}{9 \cdot 2\sqrt{5}} = \dfrac{2}{9\sqrt{5}}$ and $\theta = \cos^{-1}\left(\dfrac{2}{9\sqrt{5}}\right) \approx 84°$.

21. Let p, q, and r be the angles at vertices P, Q, and R respectively.



Then p is the angle between vectors $\overrightarrow{PQ}$ and $\overrightarrow{PR}$, q is the angle between vectors $\overrightarrow{QP}$ and $\overrightarrow{QR}$, and r is the angle between vectors $\overrightarrow{RP}$ and $\overrightarrow{RQ}$.

Thus $\cos p = \dfrac{\overrightarrow{PQ} \cdot \overrightarrow{PR}}{\left|\overrightarrow{PQ}\right|\left|\overrightarrow{PR}\right|} = \dfrac{\langle -2, 3\rangle \cdot \langle 1, 4\rangle}{\sqrt{(-2)^2 + 3^2}\sqrt{1^2 + 4^2}} = \dfrac{-2 + 12}{\sqrt{13}\sqrt{17}} = \dfrac{10}{\sqrt{221}}$ and $p = \cos^{-1}\left(\dfrac{10}{\sqrt{221}}\right) \approx 48°$. Similarly,

$\cos q = \dfrac{\overrightarrow{QP} \cdot \overrightarrow{QR}}{\left|\overrightarrow{QP}\right|\left|\overrightarrow{QR}\right|} = \dfrac{\langle 2, -3\rangle \cdot \langle 3, 1\rangle}{\sqrt{4 + 9}\sqrt{9 + 1}} = \dfrac{6 - 3}{\sqrt{13}\sqrt{10}} = \dfrac{3}{\sqrt{130}}$ so $q = \cos^{-1}\left(\dfrac{3}{\sqrt{130}}\right) \approx 75°$ and

$r \approx 180° - (48° + 75°) = 57°$.

Alternate solution: Apply the Law of Cosines three times as follows: $\cos p = \dfrac{\left|\overrightarrow{QR}\right|^2 - \left|\overrightarrow{PQ}\right|^2 - \left|\overrightarrow{PR}\right|^2}{2\left|\overrightarrow{PQ}\right|\left|\overrightarrow{PR}\right|}$,

$\cos q = \dfrac{\left|\overrightarrow{PR}\right|^2 - \left|\overrightarrow{PQ}\right|^2 - \left|\overrightarrow{QR}\right|^2}{2\left|\overrightarrow{PQ}\right|\left|\overrightarrow{QR}\right|}$, and $\cos r = \dfrac{\left|\overrightarrow{PQ}\right|^2 - \left|\overrightarrow{PR}\right|^2 - \left|\overrightarrow{QR}\right|^2}{2\left|\overrightarrow{PR}\right|\left|\overrightarrow{QR}\right|}$.

22. Let a, b, and c be the angles at vertices A, B, and C. Then a is the angle between vectors $\overrightarrow{AB}$ and $\overrightarrow{AC}$, b is the angle between vectors $\overrightarrow{BA}$ and $\overrightarrow{BC}$, and c is the angle between vectors $\overrightarrow{CA}$ and $\overrightarrow{CB}$.



Thus $\cos a = \dfrac{\overrightarrow{AB} \cdot \overrightarrow{AC}}{\left|\overrightarrow{AB}\right|\left|\overrightarrow{AC}\right|} = \dfrac{\langle 2, -2, 1\rangle \cdot \langle 0, 3, 4\rangle}{\sqrt{2^2 + (-2)^2 + 1^2}\sqrt{0^2 + 3^2 + 4^2}} = \dfrac{0 - 6 + 4}{3 \cdot 5} = -\dfrac{2}{15}$ and $a = \cos^{-1}\left(-\frac{2}{15}\right) \approx 98°$.

Similarly, $\cos b = \dfrac{\overrightarrow{BA} \cdot \overrightarrow{BC}}{\left|\overrightarrow{BA}\right|\left|\overrightarrow{BC}\right|} = \dfrac{\langle -2, 2, -1\rangle \cdot \langle -2, 5, 3\rangle}{\sqrt{4 + 4 + 1}\sqrt{4 + 25 + 9}} = \dfrac{4 + 10 - 3}{3 \cdot \sqrt{38}} = \dfrac{11}{3\sqrt{38}}$ so $b = \cos^{-1}\left(\dfrac{11}{3\sqrt{38}}\right) \approx 54°$ and

$c \approx 180° - (98° + 54°) = 28°$.

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Alternate solution: Apply the Law of Cosines three times as follows:

$$\cos a = \frac{\left|\overrightarrow{BC}\right|^2 - \left|\overrightarrow{AB}\right|^2 - \left|\overrightarrow{AC}\right|^2}{2\left|\overrightarrow{AB}\right|\left|\overrightarrow{AC}\right|} \qquad \cos b = \frac{\left|\overrightarrow{AC}\right|^2 - \left|\overrightarrow{AB}\right|^2 - \left|\overrightarrow{BC}\right|^2}{2\left|\overrightarrow{AB}\right|\left|\overrightarrow{BC}\right|} \qquad \cos c = \frac{\left|\overrightarrow{AB}\right|^2 - \left|\overrightarrow{AC}\right|^2 - \left|\overrightarrow{BC}\right|^2}{2\left|\overrightarrow{AC}\right|\left|\overrightarrow{BC}\right|}$$

23. (a) $\mathbf{a}\cdot\mathbf{b} = (9)(-2) + (3)(6) = 0$, so $\mathbf{a}$ and $\mathbf{b}$ are orthogonal (and not parallel).

(b) $\mathbf{a}\cdot\mathbf{b} = (4)(3) + (5)(-1) + (-2)(5) = -3 \neq 0$, so $\mathbf{a}$ and $\mathbf{b}$ are not orthogonal. Also, since $\mathbf{a}$ is not a scalar multiple of $\mathbf{b}$, $\mathbf{a}$ and $\mathbf{b}$ are not parallel.

(c) $\mathbf{a}\cdot\mathbf{b} = (-8)(6) + (12)(-9) + (4)(-3) = -168 \neq 0$, so $\mathbf{a}$ and $\mathbf{b}$ are not orthogonal. Because $\mathbf{a} = -\frac{4}{3}\,\mathbf{b}$, $\mathbf{a}$ and $\mathbf{b}$ are parallel.

(d) $\mathbf{a}\cdot\mathbf{b} = (3)(5) + (-1)(9) + (3)(-2) = 0$, so $\mathbf{a}$ and $\mathbf{b}$ are orthogonal (and not parallel).

24. (a) $\mathbf{u}\cdot\mathbf{v} = (-5)(3) + (4)(4) + (-2)(-1) = 3 \neq 0$, so $\mathbf{u}$ and $\mathbf{v}$ are not orthogonal. Also, $\mathbf{u}$ is not a scalar multiple of $\mathbf{v}$, so $\mathbf{u}$ and $\mathbf{v}$ are not parallel.

(b) $\mathbf{u}\cdot\mathbf{v} = (9)(-6) + (-6)(4) + (3)(-2) = -84 \neq 0$, so $\mathbf{u}$ and $\mathbf{v}$ are not orthogonal. Because $\mathbf{u} = -\frac{3}{2}\,\mathbf{v}$, $\mathbf{u}$ and $\mathbf{v}$ are parallel.

(c) $\mathbf{u}\cdot\mathbf{v} = (c)(c) + (c)(0) + (c)(-c) = c^2 + 0 - c^2 = 0$, so $\mathbf{u}$ and $\mathbf{v}$ are orthogonal (and not parallel). (Note that if $c = 0$ then $\mathbf{u} = \mathbf{v} = \mathbf{0}$, and the zero vector is considered orthogonal to all vectors. Although in this case $\mathbf{u}$ and $\mathbf{v}$ are identical, they are not considered parallel, as only nonzero vectors can be parallel.)

25. $\overrightarrow{QP} = \langle -1, -3, 2\rangle$, $\overrightarrow{QR} = \langle 4, -2, -1\rangle$, and $\overrightarrow{QP}\cdot\overrightarrow{QR} = -4 + 6 - 2 = 0$. Thus $\overrightarrow{QP}$ and $\overrightarrow{QR}$ are orthogonal, so the angle of the triangle at vertex Q is a right angle.

26. By Theorem 3, vectors $\langle 2, 1, -1\rangle$ and $\langle 1, x, 0\rangle$ meet at an angle of 45° when

$\langle 2, 1, -1\rangle \cdot \langle 1, x, 0\rangle = \sqrt{4+1+1}\,\sqrt{1+x^2+0}\,\cos 45^\circ$ or $2 + x - 0 = \sqrt{6}\,\sqrt{1+x^2}\cdot\frac{\sqrt{2}}{2} \quad\Leftrightarrow\quad 2 + x = \sqrt{3}\sqrt{1+x^2}$.

Squaring both sides gives $4 + 4x + x^2 = 3 + 3x^2 \quad\Leftrightarrow\quad 2x^2 - 4x - 1 = 0$. By the quadratic formula,

$x = \dfrac{-(-4) \pm \sqrt{(-4)^2 - 4(2)(-1)}}{2(2)} = \dfrac{4 \pm \sqrt{24}}{4} = \dfrac{4 \pm 2\sqrt{6}}{4} = 1 \pm \dfrac{\sqrt{6}}{2}$. (You can verify that both values are valid.)

27. Let $\mathbf{a} = a_1\,\mathbf{i} + a_2\,\mathbf{j} + a_3\,\mathbf{k}$ be a vector orthogonal to both $\mathbf{i} + \mathbf{j}$ and $\mathbf{i} + \mathbf{k}$. Then $\mathbf{a}\cdot(\mathbf{i} + \mathbf{j}) = 0 \quad\Leftrightarrow\quad a_1 + a_2 = 0$ and $\mathbf{a}\cdot(\mathbf{i} + \mathbf{k}) = 0 \quad\Leftrightarrow\quad a_1 + a_3 = 0$, so $a_1 = -a_2 = -a_3$. Furthermore $\mathbf{a}$ is to be a unit vector, so $1 = a_1^2 + a_2^2 + a_3^2 = 3a_1^2$ implies $a_1 = \pm\frac{1}{\sqrt{3}}$. Thus $\mathbf{a} = \frac{1}{\sqrt{3}}\,\mathbf{i} - \frac{1}{\sqrt{3}}\,\mathbf{j} - \frac{1}{\sqrt{3}}\,\mathbf{k}$ and $\mathbf{a} = -\frac{1}{\sqrt{3}}\,\mathbf{i} + \frac{1}{\sqrt{3}}\,\mathbf{j} + \frac{1}{\sqrt{3}}\,\mathbf{k}$ are two such unit vectors.

28. Let $\mathbf{u} = \langle a, b\rangle$ be a unit vector. By Theorem 3 we need $\mathbf{u}\cdot\mathbf{v} = |\mathbf{u}|\,|\mathbf{v}|\cos 60^\circ \quad\Leftrightarrow\quad 3a + 4b = (1)(5)\frac{1}{2} \quad\Leftrightarrow$

$b = \frac{5}{8} - \frac{3}{4}a$. Since $\mathbf{u}$ is a unit vector, $|\mathbf{u}| = \sqrt{a^2 + b^2} = 1 \quad\Leftrightarrow\quad a^2 + b^2 = 1 \quad\Leftrightarrow\quad a^2 + \left(\frac{5}{8} - \frac{3}{4}a\right)^2 = 1 \quad\Leftrightarrow$

$\frac{25}{16}a^2 - \frac{15}{16}a + \frac{25}{64} = 1 \quad\Leftrightarrow\quad 100a^2 - 60a - 39 = 0$. By the quadratic formula,

$a = \dfrac{-(-60) \pm \sqrt{(-60)^2 - 4(100)(-39)}}{2(100)} = \dfrac{60 \pm \sqrt{19{,}200}}{200} = \dfrac{3 \pm 4\sqrt{3}}{10}$. If $a = \dfrac{3 + 4\sqrt{3}}{10}$ then

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$b = \frac{5}{8} - \frac{3}{4}\left(\frac{3+4\sqrt{3}}{10}\right) = \frac{4-3\sqrt{3}}{10}$, and if $a = \frac{3-4\sqrt{3}}{10}$ then $b = \frac{5}{8} - \frac{3}{4}\left(\frac{3-4\sqrt{3}}{10}\right) = \frac{4+3\sqrt{3}}{10}$. Thus the two unit vectors are $\left\langle \frac{3+4\sqrt{3}}{10}, \frac{4-3\sqrt{3}}{10} \right\rangle \approx \langle 0.9928, -0.1196\rangle$ and $\left\langle \frac{3-4\sqrt{3}}{10}, \frac{4+3\sqrt{3}}{10} \right\rangle \approx \langle -0.3928, 0.9196\rangle$.

29. The line $2x - y = 3 \Leftrightarrow y = 2x - 3$ has slope 2, so a vector parallel to the line is $\mathbf{a} = \langle 1, 2\rangle$. The line $3x + y = 7 \Leftrightarrow y = -3x + 7$ has slope -3, so a vector parallel to the line is $\mathbf{b} = \langle 1, -3\rangle$. The angle between the lines is the same as the angle θ between the vectors. Here we have $\mathbf{a} \cdot \mathbf{b} = (1)(1) + (2)(-3) = -5$, $|\mathbf{a}| = \sqrt{1^2 + 2^2} = \sqrt{5}$, and $|\mathbf{b}| = \sqrt{1^2 + (-3)^2} = \sqrt{10}$, so $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{-5}{\sqrt{5} \cdot \sqrt{10}} = \dfrac{-5}{5\sqrt{2}} = -\dfrac{1}{\sqrt{2}}$ or $-\dfrac{\sqrt{2}}{2}$. Thus $\theta = 135°$, and the acute angle between the lines is $180° - 135° = 45°$.

30. The line $x + 2y = 7 \Leftrightarrow y = -\frac{1}{2}x + \frac{7}{2}$ has slope $-\frac{1}{2}$, so a vector parallel to the line is $\mathbf{a} = \langle 2, -1\rangle$. The line $5x - y = 2 \Leftrightarrow y = 5x - 2$ has slope 5, so a vector parallel to the line is $\mathbf{b} = \langle 1, 5\rangle$. The lines meet at the same angle θ that the vectors meet at. Here we have $\mathbf{a} \cdot \mathbf{b} = (2)(1) + (-1)(5) = -3$, $|\mathbf{a}| = \sqrt{2^2 + (-1)^2} = \sqrt{5}$, and $|\mathbf{b}| = \sqrt{1^2 + 5^2} = \sqrt{26}$, so $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{-3}{\sqrt{5} \cdot \sqrt{26}} = \dfrac{-3}{\sqrt{130}}$ and $\theta = \cos^{-1}\left(\dfrac{-3}{\sqrt{130}}\right) \approx 105.3°$. The acute angle between the lines is approximately $180° - 105.3° = 74.7°$.

31. The curves $y = x^2$ and $y = x^3$ meet when $x^2 = x^3 \Leftrightarrow x^3 - x^2 = 0 \Leftrightarrow x^2(x-1) = 0 \Leftrightarrow x = 0, x = 1$. We have $\frac{d}{dx}x^2 = 2x$ and $\frac{d}{dx}x^3 = 3x^2$, so the tangent lines of both curves have slope 0 at $x = 0$. Thus the angle between the curves is 0° at the point $(0, 0)$. For $x = 1$, $\left.\dfrac{d}{dx}\,x^2\right|_{x=1} = 2$ and $\left.\dfrac{d}{dx}\,x^3\right|_{x=1} = 3$ so the tangent lines at the point $(1, 1)$ have slopes 2 and 3. Vectors parallel to the tangent lines are $\langle 1, 2\rangle$ and $\langle 1, 3\rangle$, and the angle θ between them is given by

$$\cos\theta = \frac{\langle 1,2\rangle \cdot \langle 1,3\rangle}{|\langle 1,2\rangle|\,|\langle 1,3\rangle|} = \frac{1+6}{\sqrt{5}\sqrt{10}} = \frac{7}{5\sqrt{2}}$$

Thus $\theta = \cos^{-1}\left(\dfrac{7}{5\sqrt{2}}\right) \approx 8.1°$.

32. The curves $y = \sin x$ and $y = \cos x$ meet when $\sin x = \cos x \Leftrightarrow \tan x = 1 \Leftrightarrow x = \pi/4$ $[0 \le x \le \pi/2]$. Thus the point of intersection is $(\pi/4, \sqrt{2}/2)$. We have $\left.\dfrac{d}{dx}\sin x\right|_{x=\pi/4} = \left.\cos x\right|_{x=\pi/4} = \dfrac{\sqrt{2}}{2}$ and $\left.\dfrac{d}{dx}\cos x\right|_{x=\pi/4} = -\left.\sin x\right|_{x=\pi/4} = -\dfrac{\sqrt{2}}{2}$, so the tangent lines at that point have slopes $\dfrac{\sqrt{2}}{2}$ and $-\dfrac{\sqrt{2}}{2}$. Vectors parallel to the tangent lines are $\left\langle 1, \dfrac{\sqrt{2}}{2}\right\rangle$ and $\left\langle 1, -\dfrac{\sqrt{2}}{2}\right\rangle$, and the angle θ between them is given by

$$\cos\theta = \frac{\langle 1, \sqrt{2}/2\rangle \cdot \langle 1, -\sqrt{2}/2\rangle}{|\langle 1, \sqrt{2}/2\rangle|\,|\langle 1, -\sqrt{2}/2\rangle|} = \frac{1 - \frac{1}{2}}{\sqrt{\frac{3}{2}}\sqrt{\frac{3}{2}}} = \frac{1/2}{3/2} = \frac{1}{3}$$

Thus $\theta = \cos^{-1}\left(\frac{1}{3}\right) \approx 70.5°$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

33. Since $|\langle 2,1,2\rangle| = \sqrt{4+1+4} = \sqrt{9} = 3$, using Equations 8 and 9 we have $\cos\alpha = \frac{2}{3}$, $\cos\beta = \frac{1}{3}$, and $\cos\gamma = \frac{2}{3}$. The direction angles are given by $\alpha = \cos^{-1}\left(\frac{2}{3}\right) \approx 48°$, $\beta = \cos^{-1}\left(\frac{1}{3}\right) \approx 71°$, and $\gamma = \cos^{-1}\left(\frac{2}{3}\right) = 48°$.

34. Since $|\langle 6,3,-2\rangle| = \sqrt{36+9+4} = \sqrt{49} = 7$, using Equations 8 and 9 we have $\cos\alpha = \frac{6}{7}$, $\cos\beta = \frac{3}{7}$, and $\cos\gamma = \frac{-2}{7}$. The direction angles are given by $\alpha = \cos^{-1}\left(\frac{6}{7}\right) \approx 31°$, $\beta = \cos^{-1}\left(\frac{3}{7}\right) \approx 65°$, and $\gamma = \cos^{-1}\left(-\frac{2}{7}\right) = 107°$.

35. Since $|\mathbf{i} - 2\mathbf{j} - 3\mathbf{k}| = \sqrt{1+4+9} = \sqrt{14}$, Equations 8 and 9 give $\cos\alpha = \frac{1}{\sqrt{14}}$, $\cos\beta = \frac{-2}{\sqrt{14}}$, and $\cos\gamma = \frac{-3}{\sqrt{14}}$, while $\alpha = \cos^{-1}\left(\frac{1}{\sqrt{14}}\right) \approx 74°$, $\beta = \cos^{-1}\left(-\frac{2}{\sqrt{14}}\right) \approx 122°$, and $\gamma = \cos^{-1}\left(-\frac{3}{\sqrt{14}}\right) \approx 143°$.

36. Since $\left|\frac{1}{2}\mathbf{i} + \mathbf{j} + \mathbf{k}\right| = \sqrt{\frac{1}{4}+1+1} = \sqrt{\frac{9}{4}} = \frac{3}{2}$, Equations 8 and 9 give $\cos\alpha = \frac{1/2}{3/2} = \frac{1}{3}$, $\cos\beta = \cos\gamma = \frac{1}{3/2} = \frac{2}{3}$, while $\alpha = \cos^{-1}\left(\frac{1}{3}\right) \approx 71°$ and $\beta = \gamma = \cos^{-1}\left(\frac{2}{3}\right) \approx 48°$.

37. $|\langle c,c,c\rangle| = \sqrt{c^2+c^2+c^2} = \sqrt{3}c$ [since $c > 0$], so $\cos\alpha = \cos\beta = \cos\gamma = \dfrac{c}{\sqrt{3}c} = \dfrac{1}{\sqrt{3}}$ and $\alpha = \beta = \gamma = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right) \approx 55°$.

38. Since $\cos^2\alpha + \cos^2\beta + \cos^2\gamma = 1$, $\cos^2\gamma = 1 - \cos^2\alpha - \cos^2\beta = 1 - \cos^2\left(\frac{\pi}{4}\right) - \cos^2\left(\frac{\pi}{3}\right) = 1 - \left(\frac{\sqrt{2}}{2}\right)^2 - \left(\frac{1}{2}\right)^2 = \frac{1}{4}$. Thus $\cos\gamma = \pm\frac{1}{2}$ and $\gamma = \frac{\pi}{3}$ or $\gamma = \frac{2\pi}{3}$.

39. $|\mathbf{a}| = \sqrt{(-5)^2+12^2} = \sqrt{169} = 13$. The scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{-5\cdot 4 + 12\cdot 6}{13} = 4$ and the vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{proj}_{\mathbf{a}}\,\mathbf{b} = \left(\dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|}\right)\dfrac{\mathbf{a}}{|\mathbf{a}|} = 4\cdot\frac{1}{13}\langle -5,12\rangle = \left\langle -\frac{20}{13}, \frac{48}{13}\right\rangle$.

40. $|\mathbf{a}| = \sqrt{1^2+4^2} = \sqrt{17}$. The scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{1\cdot 2 + 4\cdot 3}{\sqrt{17}} = \dfrac{14}{\sqrt{17}}$ and the vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{proj}_{\mathbf{a}}\,\mathbf{b} = \left(\dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|}\right)\dfrac{\mathbf{a}}{|\mathbf{a}|} = \frac{14}{\sqrt{17}}\cdot\frac{1}{\sqrt{17}}\langle 1,4\rangle = \left\langle \frac{14}{17}, \frac{56}{17}\right\rangle$.

41. $|\mathbf{a}| = \sqrt{4^2+7^2+(-4)^2} = \sqrt{81} = 9$ so the scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{comp}_{\mathbf{a}}\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{(4)(3)+(7)(-1)+(-4)(1)}{9} = \dfrac{1}{9}$. The vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{proj}_{\mathbf{a}}\mathbf{b} = \left(\dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|}\right)\dfrac{\mathbf{a}}{|\mathbf{a}|} = \frac{1}{9}\cdot\frac{1}{9}\langle 4,7,-4\rangle = \frac{1}{81}\langle 4,7,-4\rangle = \left\langle \frac{4}{81}, \frac{7}{81}, -\frac{4}{81}\right\rangle$.

42. $|\mathbf{a}| = \sqrt{1+16+64} = \sqrt{81} = 9$ so the scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \frac{1}{9}(-12+4+16) = \frac{8}{9}$, while the vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{proj}_{\mathbf{a}}\,\mathbf{b} = \left(\dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|}\right)\dfrac{\mathbf{a}}{|\mathbf{a}|} = \frac{8}{9}\cdot\frac{1}{9}\langle -1,4,8\rangle = \frac{8}{81}\langle -1,4,8\rangle = \left\langle -\frac{8}{81}, \frac{32}{81}, \frac{64}{81}\right\rangle$.

43. $|\mathbf{a}| = \sqrt{9+9+1} = \sqrt{19}$ so the scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{6-12-1}{\sqrt{19}} = -\dfrac{7}{\sqrt{19}}$ while the vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\mathrm{proj}_{\mathbf{a}}\,\mathbf{b} = -\dfrac{7}{\sqrt{19}}\dfrac{\mathbf{a}}{|\mathbf{a}|} = -\frac{7}{\sqrt{19}}\cdot\frac{1}{\sqrt{19}}(3\mathbf{i}-3\mathbf{j}+\mathbf{k}) = -\frac{7}{19}(3\mathbf{i}-3\mathbf{j}+\mathbf{k}) = -\frac{21}{19}\mathbf{i} + \frac{21}{19}\mathbf{j} - \frac{7}{19}\mathbf{k}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. $|\mathbf{a}| = \sqrt{1+4+9} = \sqrt{14}$ so the scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{5+0-3}{\sqrt{14}} = \dfrac{2}{\sqrt{14}}$ while the vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \dfrac{2}{\sqrt{14}}\dfrac{\mathbf{a}}{|\mathbf{a}|} = \frac{2}{\sqrt{14}}\cdot\frac{1}{\sqrt{14}}\,(\mathbf{i}+2\mathbf{j}+3\mathbf{k}) = \frac{1}{7}(\mathbf{i}+2\mathbf{j}+3\mathbf{k}) = \frac{1}{7}\mathbf{i}+\frac{2}{7}\mathbf{j}+\frac{3}{7}\mathbf{k}$.

45. $(\text{orth}_{\mathbf{a}}\,\mathbf{b})\cdot\mathbf{a} = (\mathbf{b}-\text{proj}_{\mathbf{a}}\,\mathbf{b})\cdot\mathbf{a} = \mathbf{b}\cdot\mathbf{a} - (\text{proj}_{\mathbf{a}}\,\mathbf{b})\cdot\mathbf{a} = \mathbf{b}\cdot\mathbf{a} - \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\,\mathbf{a}\cdot\mathbf{a} = \mathbf{b}\cdot\mathbf{a} - \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\,|\mathbf{a}|^2 = \mathbf{b}\cdot\mathbf{a} - \mathbf{a}\cdot\mathbf{b} = 0$.

So they are orthogonal by (7).

46. Using the formula in Exercise 45 and the result of Exercise 40, we have $\text{orth}_{\mathbf{a}}\,\mathbf{b} = \mathbf{b} - \text{proj}_{\mathbf{a}}\,\mathbf{b} = \langle 2,3\rangle - \left\langle \frac{14}{17}, \frac{56}{17}\right\rangle = \left\langle \frac{20}{17}, -\frac{5}{17}\right\rangle$.



47. $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = 2 \iff \mathbf{a}\cdot\mathbf{b} = 2\,|\mathbf{a}| = 2\sqrt{10}$. If $\mathbf{b} = \langle b_1, b_2, b_3\rangle$, then we need $3b_1 + 0b_2 - 1b_3 = 2\sqrt{10}$.

One possible solution is obtained by taking $b_1 = 0$, $b_2 = 0$, $b_3 = -2\sqrt{10}$. In general, $\mathbf{b} = \langle s, t, 3s - 2\sqrt{10}\,\rangle$, $s, t \in \mathbb{R}$.

48. (a) $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \text{comp}_{\mathbf{b}}\,\mathbf{a} \iff \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{\mathbf{b}\cdot\mathbf{a}}{|\mathbf{b}|} \iff \dfrac{1}{|\mathbf{a}|} = \dfrac{1}{|\mathbf{b}|}$ or $\mathbf{a}\cdot\mathbf{b} = 0 \iff |\mathbf{b}| = |\mathbf{a}|$ or $\mathbf{a}\cdot\mathbf{b} = 0$.

That is, if $\mathbf{a}$ and $\mathbf{b}$ are orthogonal or if they have the same length.

(b) $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \text{proj}_{\mathbf{b}}\,\mathbf{a} \iff \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\,\mathbf{a} = \dfrac{\mathbf{b}\cdot\mathbf{a}}{|\mathbf{b}|^2}\,\mathbf{b} \iff \mathbf{a}\cdot\mathbf{b} = 0$ or $\dfrac{\mathbf{a}}{|\mathbf{a}|^2} = \dfrac{\mathbf{b}}{|\mathbf{b}|^2}$.

But $\dfrac{\mathbf{a}}{|\mathbf{a}|^2} = \dfrac{\mathbf{b}}{|\mathbf{b}|^2} \Rightarrow \dfrac{|\mathbf{a}|}{|\mathbf{a}|^2} = \dfrac{|\mathbf{b}|}{|\mathbf{b}|^2} \Rightarrow |\mathbf{a}| = |\mathbf{b}|$. Substituting this into the previous equation gives $\mathbf{a} = \mathbf{b}$.

So $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \text{proj}_{\mathbf{b}}\,\mathbf{a} \iff \mathbf{a}$ and $\mathbf{b}$ are orthogonal, or they are equal.

49. The displacement vector is $\mathbf{D} = (6-0)\,\mathbf{i} + (12-10)\,\mathbf{j} + (20-8)\,\mathbf{k} = 6\,\mathbf{i} + 2\,\mathbf{j} + 12\,\mathbf{k}$ so, by Equation 12, the work done is $W = \mathbf{F}\cdot\mathbf{D} = (8\,\mathbf{i} - 6\,\mathbf{j} + 9\,\mathbf{k})\cdot(6\,\mathbf{i} + 2\,\mathbf{j} + 12\,\mathbf{k}) = 48 - 12 + 108 = 144$ joules.

50. Here $|\mathbf{D}| = 1000$ m, $|\mathbf{F}| = 1500$ N, and $\theta = 30°$. Thus

$W = \mathbf{F}\cdot\mathbf{D} = |\mathbf{F}|\,|\mathbf{D}|\cos\theta = (1500)(1000)\left(\frac{\sqrt{3}}{2}\right) = 750{,}000\sqrt{3}$ joules.

51. Here $|\mathbf{D}| = 80$ ft, $|\mathbf{F}| = 30$ lb, and $\theta = 40°$. Thus

$W = \mathbf{F}\cdot\mathbf{D} = |\mathbf{F}|\,|\mathbf{D}|\cos\theta = (30)(80)\cos 40° = 2400\cos 40° \approx 1839$ ft-lb.

52. $W = \mathbf{F}\cdot\mathbf{D} = |\mathbf{F}|\,|\mathbf{D}|\cos\theta = (400)(120)\cos 36° \approx 38{,}833$ ft-lb



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

53. First note that $\mathbf{n} = \langle a, b\rangle$ is perpendicular to the line, because if $Q_1 = (a_1, b_1)$ and $Q_2 = (a_2, b_2)$ lie on the line, then $\mathbf{n} \cdot \overrightarrow{Q_1Q_2} = aa_2 - aa_1 + bb_2 - bb_1 = 0$, since $aa_2 + bb_2 = -c = aa_1 + bb_1$ from the equation of the line.

Let $P_2 = (x_2, y_2)$ lie on the line. Then the distance from P_1 to the line is the absolute value of the scalar projection of $\overrightarrow{P_1P_2}$ onto $\mathbf{n}$. $\operatorname{comp}_{\mathbf{n}}\left(\overrightarrow{P_1P_2}\right) = \dfrac{|\mathbf{n} \cdot \langle x_2 - x_1, y_2 - y_1\rangle|}{|\mathbf{n}|} = \dfrac{|ax_2 - ax_1 + by_2 - by_1|}{\sqrt{a^2 + b^2}} = \dfrac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$

since $ax_2 + by_2 = -c$. The required distance is $\dfrac{|(3)(-2) + (-4)(3) + 5|}{\sqrt{3^2 + (-4)^2}} = \dfrac{13}{5}$.

54. $(\mathbf{r} - \mathbf{a}) \cdot (\mathbf{r} - \mathbf{b}) = 0$ implies that the vectors $\mathbf{r} - \mathbf{a}$ and $\mathbf{r} - \mathbf{b}$ are orthogonal. From the diagram (in which A, B and R are the terminal points of the vectors), we see that this implies that R lies on a sphere whose diameter is the line from A to B. The center of this circle is the midpoint of AB, that is, $\frac{1}{2}(\mathbf{a} + \mathbf{b}) = \left\langle \frac{1}{2}(a_1 + b_1), \frac{1}{2}(a_2 + b_2), \frac{1}{2}(a_3 + b_3)\right\rangle$, and its radius is $\frac{1}{2}|\mathbf{a} - \mathbf{b}| = \frac{1}{2}\sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2 + (a_3 - b_3)^2}$.



Or: Expand the given equation, substitute $\mathbf{r} \cdot \mathbf{r} = x^2 + y^2 + z^2$ and complete the squares.

55. For convenience, consider the unit cube positioned so that its back left corner is at the origin, and its edges lie along the coordinate axes. The diagonal of the cube that begins at the origin and ends at $(1, 1, 1)$ has vector representation $\langle 1, 1, 1\rangle$. The angle θ between this vector and the vector of the edge which also begins at the origin and runs along the x-axis [that is, $\langle 1, 0, 0\rangle$] is given by $\cos\theta = \dfrac{\langle 1, 1, 1\rangle \cdot \langle 1, 0, 0\rangle}{|\langle 1, 1, 1\rangle|\,|\langle 1, 0, 0\rangle|} = \dfrac{1}{\sqrt{3}} \quad \Rightarrow \quad \theta = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right) \approx 55°$.

56. Consider a cube with sides of unit length, wholly within the first octant and with edges along each of the three coordinate axes. $\mathbf{i} + \mathbf{j} + \mathbf{k}$ and $\mathbf{i} + \mathbf{j}$ are vector representations of a diagonal of the cube and a diagonal of one of its faces. If θ is the angle between these diagonals, then $\cos\theta = \dfrac{(\mathbf{i} + \mathbf{j} + \mathbf{k}) \cdot (\mathbf{i} + \mathbf{j})}{|\mathbf{i} + \mathbf{j} + \mathbf{k}|\,|\mathbf{i} + \mathbf{j}|} = \dfrac{1 + 1}{\sqrt{3}\sqrt{2}} = \sqrt{\dfrac{2}{3}} \quad \Rightarrow \quad \theta = \cos^{-1}\sqrt{\tfrac{2}{3}} \approx 35°$.

57. Consider the H — C — H combination consisting of the sole carbon atom and the two hydrogen atoms that are at $(1, 0, 0)$ and $(0, 1, 0)$ (or any H—C—H combination, for that matter). Vector representations of the line segments emanating from the carbon atom and extending to these two hydrogen atoms are $\left\langle 1 - \frac{1}{2}, 0 - \frac{1}{2}, 0 - \frac{1}{2}\right\rangle = \left\langle \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right\rangle$ and $\left\langle 0 - \frac{1}{2}, 1 - \frac{1}{2}, 0 - \frac{1}{2}\right\rangle = \left\langle -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\right\rangle$. The bond angle, θ, is therefore given by

$$\cos\theta = \frac{\left\langle \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right\rangle \cdot \left\langle -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\right\rangle}{\left|\left\langle \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right\rangle\right|\left|\left\langle -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\right\rangle\right|} = \frac{-\frac{1}{4} - \frac{1}{4} + \frac{1}{4}}{\sqrt{\frac{3}{4}}\sqrt{\frac{3}{4}}} = -\frac{1}{3} \quad \Rightarrow \quad \theta = \cos^{-1}\left(-\tfrac{1}{3}\right) \approx 109.5°.$$

58. Let α be the angle between $\mathbf{a}$ and $\mathbf{c}$ and β be the angle between $\mathbf{c}$ and $\mathbf{b}$. We need to show that $\alpha = \beta$. Now $\cos\alpha = \dfrac{\mathbf{a} \cdot \mathbf{c}}{|\mathbf{a}|\,|\mathbf{c}|} = \dfrac{\mathbf{a} \cdot |\mathbf{a}|\,\mathbf{b} + \mathbf{a} \cdot |\mathbf{b}|\,\mathbf{a}}{|\mathbf{a}|\,|\mathbf{c}|} = \dfrac{|\mathbf{a}|\,\mathbf{a} \cdot \mathbf{b} + |\mathbf{a}|^2\,|\mathbf{b}|}{|\mathbf{a}|\,|\mathbf{c}|} = \dfrac{\mathbf{a} \cdot \mathbf{b} + |\mathbf{a}|\,|\mathbf{b}|}{|\mathbf{c}|}$. Similarly,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\cos\beta = \dfrac{\mathbf{b}\cdot\mathbf{c}}{|\mathbf{b}|\,|\mathbf{c}|} = \dfrac{|\mathbf{a}|\,|\mathbf{b}| + \mathbf{b}\cdot\mathbf{a}}{|\mathbf{c}|}$. Thus $\cos\alpha = \cos\beta$. However $0° \le \alpha \le 180°$ and $0° \le \beta \le 180°$, so $\alpha = \beta$ and $\mathbf{c}$ bisects the angle between $\mathbf{a}$ and $\mathbf{b}$.

59. Let $\mathbf{a} = \langle a_1, a_2, a_3\rangle$ and $\mathbf{b} = \langle b_1, b_2, b_3\rangle$.

Property 2: $\mathbf{a}\cdot\mathbf{b} = \langle a_1, a_2, a_3\rangle \cdot \langle b_1, b_2, b_3\rangle = a_1b_1 + a_2b_2 + a_3b_3$

$= b_1a_1 + b_2a_2 + b_3a_3 = \langle b_1, b_2, b_3\rangle \cdot \langle a_1, a_2, a_3\rangle = \mathbf{b}\cdot\mathbf{a}$

Property 4: $(c\,\mathbf{a})\cdot\mathbf{b} = \langle ca_1, ca_2, ca_3\rangle \cdot \langle b_1, b_2, b_3\rangle = (ca_1)b_1 + (ca_2)b_2 + (ca_3)b_3$

$= c(a_1b_1 + a_2b_2 + a_3b_3) = c(\mathbf{a}\cdot\mathbf{b}) = a_1(cb_1) + a_2(cb_2) + a_3(cb_3)$

$= \langle a_1, a_2, a_3\rangle \cdot \langle cb_1, cb_2, cb_3\rangle = \mathbf{a}\cdot(c\,\mathbf{b})$

Property 5: $\mathbf{0}\cdot\mathbf{a} = \langle 0,0,0\rangle \cdot \langle a_1, a_2, a_3\rangle = (0)(a_1) + (0)(a_2) + (0)(a_3) = 0$

60. Let the figure be called quadrilateral $ABCD$. The diagonals can be represented by $\overrightarrow{AC}$ and $\overrightarrow{BD}$. $\overrightarrow{AC} = \overrightarrow{AB} + \overrightarrow{BC}$ and $\overrightarrow{BD} = \overrightarrow{BC} + \overrightarrow{CD} = \overrightarrow{BC} - \overrightarrow{DC} = \overrightarrow{BC} - \overrightarrow{AB}$ (Since opposite sides of the object are of the same length and parallel, $\overrightarrow{AB} = \overrightarrow{DC}$.) Thus

$$\overrightarrow{AC}\cdot\overrightarrow{BD} = \left(\overrightarrow{AB} + \overrightarrow{BC}\right)\cdot\left(\overrightarrow{BC} - \overrightarrow{AB}\right) = \overrightarrow{AB}\cdot\left(\overrightarrow{BC} - \overrightarrow{AB}\right) + \overrightarrow{BC}\cdot\left(\overrightarrow{BC} - \overrightarrow{AB}\right)$$

$$= \overrightarrow{AB}\cdot\overrightarrow{BC} - \left|\overrightarrow{AB}\right|^2 + \left|\overrightarrow{BC}\right|^2 - \overrightarrow{AB}\cdot\overrightarrow{BC} = \left|\overrightarrow{BC}\right|^2 - \left|\overrightarrow{AB}\right|^2$$

But $\left|\overrightarrow{AB}\right|^2 = \left|\overrightarrow{BC}\right|^2$ because all sides of the quadrilateral are equal in length. Therefore $\overrightarrow{AC}\cdot\overrightarrow{BD} = 0$, and since both of these vectors are nonzero this tells us that the diagonals of the quadrilateral are perpendicular.

61. $|\mathbf{a}\cdot\mathbf{b}| = \big||\mathbf{a}|\,|\mathbf{b}|\cos\theta\big| = |\mathbf{a}|\,|\mathbf{b}|\,|\cos\theta|$. Since $|\cos\theta| \le 1$, $|\mathbf{a}\cdot\mathbf{b}| = |\mathbf{a}|\,|\mathbf{b}|\,|\cos\theta| \le |\mathbf{a}|\,|\mathbf{b}|$.

Note: We have equality in the case of $\cos\theta = \pm 1$, so $\theta = 0$ or $\theta = \pi$, thus equality when $\mathbf{a}$ and $\mathbf{b}$ are parallel.

62. (a)



The Triangle Inequality states that the length of the longest side of a triangle is less than or equal to the sum of the lengths of the two shortest sides.

(b) $|\mathbf{a}+\mathbf{b}|^2 = (\mathbf{a}+\mathbf{b})\cdot(\mathbf{a}+\mathbf{b}) = (\mathbf{a}\cdot\mathbf{a}) + 2(\mathbf{a}\cdot\mathbf{b}) + (\mathbf{b}\cdot\mathbf{b}) = |\mathbf{a}|^2 + 2(\mathbf{a}\cdot\mathbf{b}) + |\mathbf{b}|^2$

$\le |\mathbf{a}|^2 + 2\,|\mathbf{a}|\,|\mathbf{b}| + |\mathbf{b}|^2$ [by the Cauchy-Schwartz Inequality]

$= (|\mathbf{a}| + |\mathbf{b}|)^2$

Thus, taking the square root of both sides, $|\mathbf{a}+\mathbf{b}| \le |\mathbf{a}| + |\mathbf{b}|$.

63. (a)



The Parallelogram Law states that the sum of the squares of the lengths of the diagonals of a parallelogram equals the sum of the squares of its (four) sides.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) $|\mathbf{a}+\mathbf{b}|^2 = (\mathbf{a}+\mathbf{b})\cdot(\mathbf{a}+\mathbf{b}) = |\mathbf{a}|^2 + 2(\mathbf{a}\cdot\mathbf{b}) + |\mathbf{b}|^2$ and $|\mathbf{a}-\mathbf{b}|^2 = (\mathbf{a}-\mathbf{b})\cdot(\mathbf{a}-\mathbf{b}) = |\mathbf{a}|^2 - 2(\mathbf{a}\cdot\mathbf{b}) + |\mathbf{b}|^2$.

Adding these two equations gives $|\mathbf{a}+\mathbf{b}|^2 + |\mathbf{a}-\mathbf{b}|^2 = 2|\mathbf{a}|^2 + 2|\mathbf{b}|^2$.

64. If the vectors $\mathbf{u}+\mathbf{v}$ and $\mathbf{u}-\mathbf{v}$ are orthogonal then $(\mathbf{u}+\mathbf{v})\cdot(\mathbf{u}-\mathbf{v}) = 0$. But

$$\begin{aligned}(\mathbf{u}+\mathbf{v})\cdot(\mathbf{u}-\mathbf{v}) &= (\mathbf{u}+\mathbf{v})\cdot\mathbf{u} - (\mathbf{u}+\mathbf{v})\cdot\mathbf{v} && \text{by Property 3 of the dot product}\\ &= \mathbf{u}\cdot\mathbf{u}+\mathbf{v}\cdot\mathbf{u}-\mathbf{u}\cdot\mathbf{v}-\mathbf{v}\cdot\mathbf{v} && \text{by Property 3}\\ &= |\mathbf{u}|^2 + \mathbf{u}\cdot\mathbf{v} - \mathbf{u}\cdot\mathbf{v} - |\mathbf{v}|^2 && \text{by Properties 1 and 2}\\ &= |\mathbf{u}|^2 - |\mathbf{v}|^2\end{aligned}$$

Thus $|\mathbf{u}|^2 - |\mathbf{v}|^2 = 0 \;\Rightarrow\; |\mathbf{u}|^2 = |\mathbf{v}|^2 \;\Rightarrow\; |\mathbf{u}| = |\mathbf{v}|$ [since $|\mathbf{u}|, |\mathbf{v}| \geq 0$].

65.
$$\begin{aligned}\text{proj}_{\mathbf{a}}\,\mathbf{b}\cdot\text{proj}_{\mathbf{b}}\,\mathbf{a} &= \frac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\mathbf{a}\cdot\frac{\mathbf{b}\cdot\mathbf{a}}{|\mathbf{b}|^2}\mathbf{b} = \frac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\cdot\frac{\mathbf{b}\cdot\mathbf{a}}{|\mathbf{b}|^2}\,(\mathbf{a}\cdot\mathbf{b}) && \text{by Property 4 of the dot product}\\ &= \frac{(\mathbf{a}\cdot\mathbf{b})^2}{|\mathbf{a}|^2\,|\mathbf{b}|^2}\,(\mathbf{a}\cdot\mathbf{b}) = \left(\frac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|}\right)^2(\mathbf{a}\cdot\mathbf{b}) && \text{by Property 2}\\ &= (\cos\theta)^2\,(\mathbf{a}\cdot\mathbf{b}) = (\mathbf{a}\cdot\mathbf{b})\cos^2\theta && \text{by Corollary 6}\end{aligned}$$

12.4 The Cross Product

1. $\mathbf{a}\times\mathbf{b} = \begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\2&3&0\\1&0&5\end{vmatrix} = \begin{vmatrix}3&0\\0&5\end{vmatrix}\mathbf{i} - \begin{vmatrix}2&0\\1&5\end{vmatrix}\mathbf{j} + \begin{vmatrix}2&3\\1&0\end{vmatrix}\mathbf{k}$

$= (15-0)\,\mathbf{i} - (10-0)\,\mathbf{j} + (0-3)\,\mathbf{k} = 15\,\mathbf{i} - 10\,\mathbf{j} - 3\,\mathbf{k}$

Now $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a} = \langle 15,-10,-3\rangle\cdot\langle 2,3,0\rangle = 30-30+0 = 0$ and

$(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b} = \langle 15,-10,-3\rangle\cdot\langle 1,0,5\rangle = 15+0-15 = 0$, so $\mathbf{a}\times\mathbf{b}$ is orthogonal to both $\mathbf{a}$ and $\mathbf{b}$.

2. $\mathbf{a}\times\mathbf{b} = \begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\4&3&-2\\2&-1&1\end{vmatrix} = \begin{vmatrix}3&-2\\-1&1\end{vmatrix}\mathbf{i} - \begin{vmatrix}4&-2\\2&1\end{vmatrix}\mathbf{j} + \begin{vmatrix}4&3\\2&-1\end{vmatrix}\mathbf{k}$

$= (3-2)\,\mathbf{i} - [4-(-4)]\,\mathbf{j} + (-4-6)\,\mathbf{k} = \mathbf{i} - 8\,\mathbf{j} - 10\,\mathbf{k}$

Now $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a} = \langle 1,-8,-10\rangle\cdot\langle 4,3,-2\rangle = 4-24+20 = 0$ and

$(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b} = \langle 1,-8,-10\rangle\cdot\langle 2,-1,1\rangle = 2+8-10 = 0$, so $\mathbf{a}\times\mathbf{b}$ is orthogonal to both $\mathbf{a}$ and $\mathbf{b}$.

3. $\mathbf{a}\times\mathbf{b} = \begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\0&2&-4\\-1&3&1\end{vmatrix} = \begin{vmatrix}2&-4\\3&1\end{vmatrix}\mathbf{i} - \begin{vmatrix}0&-4\\-1&1\end{vmatrix}\mathbf{j} + \begin{vmatrix}0&2\\-1&3\end{vmatrix}\mathbf{k}$

$= [2-(-12)]\,\mathbf{i} - (0-4)\,\mathbf{j} + [0-(-2)]\,\mathbf{k} = 14\,\mathbf{i} + 4\,\mathbf{j} + 2\,\mathbf{k}$

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a} = (14\,\mathbf{i}+4\,\mathbf{j}+2\,\mathbf{k})\cdot(2\,\mathbf{j}-4\,\mathbf{k}) = 0+8-8 = 0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b} = (14\,\mathbf{i}+4\,\mathbf{j}+2\,\mathbf{k})\cdot(-\mathbf{i}+3\,\mathbf{j}+\mathbf{k}) = -14+12+2 = 0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\3&3&-3\\3&-3&3\end{vmatrix}=\begin{vmatrix}3&-3\\-3&3\end{vmatrix}\mathbf{i}-\begin{vmatrix}3&-3\\3&3\end{vmatrix}\mathbf{j}+\begin{vmatrix}3&3\\3&-3\end{vmatrix}\mathbf{k}$

$=(9-9)\,\mathbf{i}-[9-(-9)]\,\mathbf{j}+(-9-9)\,\mathbf{k}=-18\,\mathbf{j}-18\,\mathbf{k}$

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}=(-18\,\mathbf{j}-18\,\mathbf{k})\cdot(3\,\mathbf{i}+3\,\mathbf{j}-3\,\mathbf{k})=0-54+54=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}=(-18\,\mathbf{j}-18\,\mathbf{k})\cdot(3\,\mathbf{i}-3\,\mathbf{j}+3\,\mathbf{k})=0+54-54=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

5. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\\frac{1}{2}&\frac{1}{3}&\frac{1}{4}\\1&2&-3\end{vmatrix}=\begin{vmatrix}\frac{1}{3}&\frac{1}{4}\\2&-3\end{vmatrix}\mathbf{i}-\begin{vmatrix}\frac{1}{2}&\frac{1}{4}\\1&-3\end{vmatrix}\mathbf{j}+\begin{vmatrix}\frac{1}{2}&\frac{1}{3}\\1&2\end{vmatrix}\mathbf{k}$

$=\left(-1-\frac{1}{2}\right)\mathbf{i}-\left(-\frac{3}{2}-\frac{1}{4}\right)\mathbf{j}+\left(1-\frac{1}{3}\right)\mathbf{k}=-\frac{3}{2}\,\mathbf{i}+\frac{7}{4}\,\mathbf{j}+\frac{2}{3}\,\mathbf{k}$

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}=\left(-\frac{3}{2}\,\mathbf{i}+\frac{7}{4}\,\mathbf{j}+\frac{2}{3}\,\mathbf{k}\right)\cdot\left(\frac{1}{2}\,\mathbf{i}+\frac{1}{3}\,\mathbf{j}+\frac{1}{4}\,\mathbf{k}\right)=-\frac{3}{4}+\frac{7}{12}+\frac{1}{6}=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}=\left(-\frac{3}{2}\,\mathbf{i}+\frac{7}{4}\,\mathbf{j}+\frac{2}{3}\,\mathbf{k}\right)\cdot(\mathbf{i}+2\,\mathbf{j}-3\,\mathbf{k})=-\frac{3}{2}+\frac{7}{2}-2=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

6. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\t&\cos t&\sin t\\1&-\sin t&\cos t\end{vmatrix}=\begin{vmatrix}\cos t&\sin t\\-\sin t&\cos t\end{vmatrix}\mathbf{i}-\begin{vmatrix}t&\sin t\\1&\cos t\end{vmatrix}\mathbf{j}+\begin{vmatrix}t&\cos t\\1&-\sin t\end{vmatrix}\mathbf{k}$

$=[\cos^2 t-(-\sin^2 t)]\,\mathbf{i}-(t\cos t-\sin t)\,\mathbf{j}+(-t\sin t-\cos t)\,\mathbf{k}=\mathbf{i}+(\sin t-t\cos t)\,\mathbf{j}+(-t\sin t-\cos t)\,\mathbf{k}$

Since

$$\begin{aligned}(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}&=[\mathbf{i}+(\sin t-t\cos t)\,\mathbf{j}+(-t\sin t-\cos t)\,\mathbf{k}]\cdot(t\,\mathbf{i}+\cos t\,\mathbf{j}+\sin t\,\mathbf{k})\\&=t+\sin t\cos t-t\cos^2 t-t\sin^2 t-\sin t\cos t\\&=t-t\left(\cos^2 t+\sin^2 t\right)=0\end{aligned}$$

$\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since

$$\begin{aligned}(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}&=[\mathbf{i}+(\sin t-t\cos t)\,\mathbf{j}+(-t\sin t-\cos t)\,\mathbf{k}]\cdot(\mathbf{i}-\sin t\,\mathbf{j}+\cos t\,\mathbf{k})\\&=1-\sin^2 t+t\sin t\cos t-t\sin t\cos t-\cos^2 t\\&=1-\left(\sin^2 t+\cos^2 t\right)=0\end{aligned}$$

$\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

7. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\t&1&1/t\\t^2&t^2&1\end{vmatrix}=\begin{vmatrix}1&1/t\\t^2&1\end{vmatrix}\mathbf{i}-\begin{vmatrix}t&1/t\\t^2&1\end{vmatrix}\mathbf{j}+\begin{vmatrix}t&1\\t^2&t^2\end{vmatrix}\mathbf{k}$

$=(1-t)\,\mathbf{i}-(t-t)\,\mathbf{j}+(t^3-t^2)\,\mathbf{k}=(1-t)\,\mathbf{i}+(t^3-t^2)\,\mathbf{k}$

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}=\langle 1-t,0,t^3-t^2\rangle\cdot\langle t,1,1/t\rangle=t-t^2+0+t^2-t=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}=\langle 1-t,0,t^3-t^2\rangle\cdot\langle t^2,t^2,1\rangle=t^2-t^3+0+t^3-t^2=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

8. $\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & -2 \\ 0 & 1 & 1 \end{vmatrix}$

$= \begin{vmatrix} 0 & -2 \\ 1 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & -2 \\ 0 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} \mathbf{k}$

$= 2\mathbf{i} - \mathbf{j} + \mathbf{k}$



9. According to the discussion following Example 4, $\mathbf{i} \times \mathbf{j} = \mathbf{k}$, so $(\mathbf{i} \times \mathbf{j}) \times \mathbf{k} = \mathbf{k} \times \mathbf{k} = \mathbf{0}$ [by Example 2].

10. $\mathbf{k} \times (\mathbf{i} - 2\mathbf{j}) = \mathbf{k} \times \mathbf{i} + \mathbf{k} \times (-2\mathbf{j})$ by Property 3 of the cross product

$= \mathbf{k} \times \mathbf{i} + (-2)(\mathbf{k} \times \mathbf{j})$ by Property 2

$= \mathbf{j} + (-2)(-\mathbf{i}) = 2\mathbf{i} + \mathbf{j}$ by the discussion following Example 4

11. $(\mathbf{j} - \mathbf{k}) \times (\mathbf{k} - \mathbf{i}) = (\mathbf{j} - \mathbf{k}) \times \mathbf{k} + (\mathbf{j} - \mathbf{k}) \times (-\mathbf{i})$ by Property 3 of the cross product

$= \mathbf{j} \times \mathbf{k} + (-\mathbf{k}) \times \mathbf{k} + \mathbf{j} \times (-\mathbf{i}) + (-\mathbf{k}) \times (-\mathbf{i})$ by Property 4

$= (\mathbf{j} \times \mathbf{k}) + (-1)(\mathbf{k} \times \mathbf{k}) + (-1)(\mathbf{j} \times \mathbf{i}) + (-1)^2(\mathbf{k} \times \mathbf{i})$ by Property 2

$= \mathbf{i} + (-1)\,\mathbf{0} + (-1)(-\mathbf{k}) + \mathbf{j} = \mathbf{i} + \mathbf{j} + \mathbf{k}$ by Example 2 and the discussion following Example 4

12. $(\mathbf{i} + \mathbf{j}) \times (\mathbf{i} - \mathbf{j}) = (\mathbf{i} + \mathbf{j}) \times \mathbf{i} + (\mathbf{i} + \mathbf{j}) \times (-\mathbf{j})$ by Property 3 of the cross product

$= \mathbf{i} \times \mathbf{i} + \mathbf{j} \times \mathbf{i} + \mathbf{i} \times (-\mathbf{j}) + \mathbf{j} \times (-\mathbf{j})$ by Property 4

$= (\mathbf{i} \times \mathbf{i}) + (\mathbf{j} \times \mathbf{i}) + (-1)(\mathbf{i} \times \mathbf{j}) + (-1)(\mathbf{j} \times \mathbf{j})$ by Property 2

$= \mathbf{0} + (-\mathbf{k}) + (-1)\,\mathbf{k} + (-1)\,\mathbf{0} = -2\,\mathbf{k}$ by Example 2 and the discussion following Example 4

13. (a) Since $\mathbf{b} \times \mathbf{c}$ is a vector, the dot product $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$ is meaningful and is a scalar.

(b) $\mathbf{b} \cdot \mathbf{c}$ is a scalar, so $\mathbf{a} \times (\mathbf{b} \cdot \mathbf{c})$ is meaningless, as the cross product is defined only for two *vectors*.

(c) Since $\mathbf{b} \times \mathbf{c}$ is a vector, the cross product $\mathbf{a} \times (\mathbf{b} \times \mathbf{c})$ is meaningful and results in another vector.

(d) $\mathbf{b} \cdot \mathbf{c}$ is a scalar, so the dot product $\mathbf{a} \cdot (\mathbf{b} \cdot \mathbf{c})$ is meaningless, as the dot product is defined only for two vectors.

(e) Since $(\mathbf{a} \cdot \mathbf{b})$ and $(\mathbf{c} \cdot \mathbf{d})$ are both scalars, the cross product $(\mathbf{a} \cdot \mathbf{b}) \times (\mathbf{c} \cdot \mathbf{d})$ is meaningless.

(f) $\mathbf{a} \times \mathbf{b}$ and $\mathbf{c} \times \mathbf{d}$ are both vectors, so the dot product $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d})$ is meaningful and is a scalar.

14. Using Theorem 9, we have $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| \sin\theta = (4)(5) \sin 45^\circ = 20 \cdot \dfrac{\sqrt{2}}{2} = 10\sqrt{2}$. By the right-hand rule, $\mathbf{u} \times \mathbf{v}$ is directed out of the page.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. If we sketch $\mathbf{u}$ and $\mathbf{v}$ starting from the same initial point, we see that the angle between them is $60°$. Using Theorem 9, we have

$$|\mathbf{u}\times\mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}|\sin\theta = (12)(16)\sin 60° = 192\cdot\frac{\sqrt{3}}{2} = 96\sqrt{3}.$$

By the right-hand rule, $\mathbf{u}\times\mathbf{v}$ is directed into the page.



16. (a) $|\mathbf{a}\times\mathbf{b}| = |\mathbf{a}|\,|\mathbf{b}|\sin\theta = 3\cdot 2\cdot\sin\frac{\pi}{2} = 6$

(b) $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{k}$, so it lies in the xy-plane, and its z-coordinate is 0. By the right-hand rule, its y-component is negative and its x-component is positive.



17. $\mathbf{a}\times\mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -1 & 3 \\ 4 & 2 & 1 \end{vmatrix} = \begin{vmatrix} -1 & 3 \\ 2 & 1 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 2 & 3 \\ 4 & 1 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 2 & -1 \\ 4 & 2 \end{vmatrix}\mathbf{k} = (-1-6)\,\mathbf{i} - (2-12)\,\mathbf{j} + [4-(-4)]\,\mathbf{k} = -7\mathbf{i}+10\mathbf{j}+8\mathbf{k}$

$\mathbf{b}\times\mathbf{a} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4 & 2 & 1 \\ 2 & -1 & 3 \end{vmatrix} = \begin{vmatrix} 2 & 1 \\ -1 & 3 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 4 & 1 \\ 2 & 3 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 4 & 2 \\ 2 & -1 \end{vmatrix}\mathbf{k} = [6-(-1)]\,\mathbf{i} - (12-2)\,\mathbf{j} + (-4-4)\,\mathbf{k} = 7\mathbf{i}-10\mathbf{j}-8\mathbf{k}$

Notice $\mathbf{a}\times\mathbf{b} = -\mathbf{b}\times\mathbf{a}$ here, as we know is always true by Property 1 of the cross product.

18. $\mathbf{b}\times\mathbf{c} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 1 & -1 \\ 0 & 1 & 3 \end{vmatrix} = \begin{vmatrix} 1 & -1 \\ 1 & 3 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 2 & -1 \\ 0 & 3 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}\mathbf{k} = 4\mathbf{i}-6\mathbf{j}+2\mathbf{k}$ so

$\mathbf{a}\times(\mathbf{b}\times\mathbf{c}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 1 \\ 4 & -6 & 2 \end{vmatrix} = \begin{vmatrix} 0 & 1 \\ -6 & 2 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 1 & 1 \\ 4 & 2 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 1 & 0 \\ 4 & -6 \end{vmatrix}\mathbf{k} = 6\mathbf{i}+2\mathbf{j}-6\mathbf{k}.$

$\mathbf{a}\times\mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 1 \\ 2 & 1 & -1 \end{vmatrix} = \begin{vmatrix} 0 & 1 \\ 1 & -1 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 1 & 1 \\ 2 & -1 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 1 & 0 \\ 2 & 1 \end{vmatrix}\mathbf{k} = -\mathbf{i}+3\mathbf{j}+\mathbf{k}$ so

$(\mathbf{a}\times\mathbf{b})\times\mathbf{c} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 3 & 1 \\ 0 & 1 & 3 \end{vmatrix} = \begin{vmatrix} 3 & 1 \\ 1 & 3 \end{vmatrix}\mathbf{i} - \begin{vmatrix} -1 & 1 \\ 0 & 3 \end{vmatrix}\mathbf{j} + \begin{vmatrix} -1 & 3 \\ 0 & 1 \end{vmatrix}\mathbf{k} = 8\mathbf{i}+3\mathbf{j}-\mathbf{k}.$

Thus $\mathbf{a}\times(\mathbf{b}\times\mathbf{c}) \neq (\mathbf{a}\times\mathbf{b})\times\mathbf{c}$.

19. By Theorem 8, the cross product of two vectors is orthogonal to both vectors. So we calculate

$$\langle 3,2,1\rangle\times\langle -1,1,0\rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 2 & 1 \\ -1 & 1 & 0 \end{vmatrix} = \begin{vmatrix} 2 & 1 \\ 1 & 0 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 3 & 1 \\ -1 & 0 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 3 & 2 \\ -1 & 1 \end{vmatrix}\mathbf{k} = -\mathbf{i}-\mathbf{j}+5\mathbf{k}.$$

So two unit vectors orthogonal to both given vectors are $\pm\dfrac{\langle -1,-1,5\rangle}{\sqrt{1+1+25}} = \pm\dfrac{\langle -1,-1,5\rangle}{3\sqrt{3}}$, that is, $\left\langle -\frac{1}{3\sqrt{3}}, -\frac{1}{3\sqrt{3}}, \frac{5}{3\sqrt{3}}\right\rangle$ and $\left\langle \frac{1}{3\sqrt{3}}, \frac{1}{3\sqrt{3}}, -\frac{5}{3\sqrt{3}}\right\rangle$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

20. By Theorem 8, the cross product of two vectors is orthogonal to both vectors. So we calculate

$$(\mathbf{j}-\mathbf{k})\times(\mathbf{i}+\mathbf{j}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & -1 \\ 1 & 1 & 0 \end{vmatrix} = \begin{vmatrix} 1 & -1 \\ 1 & 0 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 0 & -1 \\ 1 & 0 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix}\mathbf{k} = \mathbf{i}-\mathbf{j}-\mathbf{k}$$

Thus two unit vectors orthogonal to both given vectors are $\pm\frac{1}{\sqrt{3}}(\mathbf{i}-\mathbf{j}-\mathbf{k})$, that is, $\frac{1}{\sqrt{3}}\,\mathbf{i} - \frac{1}{\sqrt{3}}\,\mathbf{j} - \frac{1}{\sqrt{3}}\,\mathbf{k}$ and $-\frac{1}{\sqrt{3}}\,\mathbf{i} + \frac{1}{\sqrt{3}}\,\mathbf{j} + \frac{1}{\sqrt{3}}\,\mathbf{k}$.

21. Let $\mathbf{a} = \langle a_1, a_2, a_3\rangle$. Then

$$\mathbf{0}\times\mathbf{a} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & 0 \\ a_1 & a_2 & a_3 \end{vmatrix} = \begin{vmatrix} 0 & 0 \\ a_2 & a_3 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 0 & 0 \\ a_1 & a_3 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 0 & 0 \\ a_1 & a_2 \end{vmatrix}\mathbf{k} = \mathbf{0},$$

$$\mathbf{a}\times\mathbf{0} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ 0 & 0 & 0 \end{vmatrix} = \begin{vmatrix} a_2 & a_3 \\ 0 & 0 \end{vmatrix}\mathbf{i} - \begin{vmatrix} a_1 & a_3 \\ 0 & 0 \end{vmatrix}\mathbf{j} + \begin{vmatrix} a_1 & a_2 \\ 0 & 0 \end{vmatrix}\mathbf{k} = \mathbf{0}.$$

22. Let $\mathbf{a} = \langle a_1, a_2, a_3\rangle$ and $\mathbf{b} = \langle b_1, b_2, b_3\rangle$.

$$(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b} = \left\langle \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix}, \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix}, \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \right\rangle \cdot \langle b_1, b_2, b_3\rangle = \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} b_1 - \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} b_2 + \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} b_3$$

$$= (a_2b_3b_1 - a_3b_2b_1) - (a_1b_3b_2 - a_3b_1b_2) + (a_1b_2b_3 - a_2b_1b_3) = 0$$

23. $\mathbf{a}\times\mathbf{b} = \langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1\rangle$

$$= \langle (-1)(b_2a_3 - b_3a_2), (-1)(b_3a_1 - b_1a_3), (-1)(b_1a_2 - b_2a_1)\rangle$$

$$= -\langle b_2a_3 - b_3a_2, b_3a_1 - b_1a_3, b_1a_2 - b_2a_1\rangle = -\mathbf{b}\times\mathbf{a}$$

24. $c\mathbf{a} = \langle ca_1, ca_2, ca_3\rangle$, so

$$(c\,\mathbf{a})\times\mathbf{b} = \langle ca_2b_3 - ca_3b_2, ca_3b_1 - ca_1b_3, ca_1b_2 - ca_2b_1\rangle$$

$$= c\langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1\rangle = c(\mathbf{a}\times\mathbf{b})$$

$$= \langle ca_2b_3 - ca_3b_2, ca_3b_1 - ca_1b_3, ca_1b_2 - ca_2b_1\rangle$$

$$= \langle a_2(cb_3) - a_3(cb_2), a_3(cb_1) - a_1(cb_3), a_1(cb_2) - a_2(cb_1)\rangle$$

$$= \mathbf{a}\times c\,\mathbf{b}$$

25. $\mathbf{a}\times(\mathbf{b}+\mathbf{c}) = \mathbf{a}\times\langle b_1+c_1, b_2+c_2, b_3+c_3\rangle$

$$= \langle a_2(b_3+c_3) - a_3(b_2+c_2), a_3(b_1+c_1) - a_1(b_3+c_3), a_1(b_2+c_2) - a_2(b_1+c_1)\rangle$$

$$= \langle a_2b_3 + a_2c_3 - a_3b_2 - a_3c_2, a_3b_1 + a_3c_1 - a_1b_3 - a_1c_3, a_1b_2 + a_1c_2 - a_2b_1 - a_2c_1\rangle$$

$$= \langle (a_2b_3 - a_3b_2) + (a_2c_3 - a_3c_2),\ (a_3b_1 - a_1b_3) + (a_3c_1 - a_1c_3),\ (a_1b_2 - a_2b_1) + (a_1c_2 - a_2c_1)\rangle$$

$$= \langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1\rangle + \langle a_2c_3 - a_3c_2, a_3c_1 - a_1c_3, a_1c_2 - a_2c_1\rangle$$

$$= (\mathbf{a}\times\mathbf{b}) + (\mathbf{a}\times\mathbf{c})$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

26. $(\mathbf{a}+\mathbf{b})\times\mathbf{c} = -\mathbf{c}\times(\mathbf{a}+\mathbf{b})$ by Property 1 of the cross product

$= -(\mathbf{c}\times\mathbf{a}+\mathbf{c}\times\mathbf{b})$ by Property 3

$= -(-\mathbf{a}\times\mathbf{c}+(-\mathbf{b}\times\mathbf{c}))$ by Property 1

$= \mathbf{a}\times\mathbf{c}+\mathbf{b}\times\mathbf{c}$ by Property 2

27. By plotting the vertices, we can see that the parallelogram is determined by the vectors $\overrightarrow{AB} = \langle 2,3\rangle$ and $\overrightarrow{AD} = \langle 6,-1\rangle$. We know that the area of the parallelogram determined by two vectors is equal to the length of the cross product of these vectors. In order to compute the cross product, we consider the vector $\overrightarrow{AB}$ as the three-dimensional vector $\langle 2,3,0\rangle$ (and similarly for $\overrightarrow{AD}$), and then the area of parallelogram $ABCD$ is



$$\left|\overrightarrow{AB}\times\overrightarrow{AD}\right| = \left\|\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 2 & 3 & 0\\ 6 & -1 & 0\end{vmatrix}\right\| = |(0-0)\,\mathbf{i}-(0-0)\,\mathbf{j}+(-2-18)\,\mathbf{k}| = |-20\,\mathbf{k}| = 20$$

28. By plotting the vertices, we can see that the parallelogram is determined by the vectors $\overrightarrow{PQ} = \langle 2,3,1\rangle$ and $\overrightarrow{PS} = \langle 4,2,5\rangle$. Thus the area of parallelogram $PQRS$ is



$$\left|\overrightarrow{PQ}\times\overrightarrow{PS}\right| = \left\|\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 2 & 3 & 1\\ 4 & 2 & 5\end{vmatrix}\right\| = |(15-2)\,\mathbf{i}-(10-4)\,\mathbf{j}+(4-12)\,\mathbf{k}|$$

$$= |13\,\mathbf{i}-6\,\mathbf{j}-8\,\mathbf{k}| = \sqrt{169+36+64} = \sqrt{269} \approx 16.40$$

29. (a) Because the plane through P, Q, and R contains the vectors $\overrightarrow{PQ}$ and $\overrightarrow{PR}$, a vector orthogonal to both of these vectors (such as their cross product) is also orthogonal to the plane. Here $\overrightarrow{PQ} = \langle -3,1,2\rangle$ and $\overrightarrow{PR} = \langle 3,2,4\rangle$, so

$$\overrightarrow{PQ}\times\overrightarrow{PR} = \langle (1)(4)-(2)(2),(2)(3)-(-3)(4),(-3)(2)-(1)(3)\rangle = \langle 0,18,-9\rangle$$

Therefore, $\langle 0,18,-9\rangle$ (or any nonzero scalar multiple thereof, such as $\langle 0,2,-1\rangle$) is orthogonal to the plane through P, Q, and R.

(b) Note that the area of the triangle determined by P, Q, and R is equal to half of the area of the parallelogram determined by the three points. From part (a), the area of the parallelogram is $\left|\overrightarrow{PQ}\times\overrightarrow{PR}\right| = |\langle 0,18,-9\rangle| = \sqrt{0+324+81} = \sqrt{405} = 9\sqrt{5}$, so the area of the triangle is $\frac{1}{2}\cdot 9\sqrt{5} = \frac{9}{2}\sqrt{5}$.

30. (a) $\overrightarrow{PQ} = \langle 4,2,3\rangle$ and $\overrightarrow{PR} = \langle 3,3,4\rangle$, so a vector orthogonal to the plane through P, Q, and R is $\overrightarrow{PQ}\times\overrightarrow{PR} = \langle (2)(4)-(3)(3),(3)(3)-(4)(4),(4)(3)-(2)(3)\rangle = \langle -1,-7,6\rangle$ (or any nonzero scalar mutiple thereof).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) The area of the parallelogram determined by $\overrightarrow{PQ}$ and $\overrightarrow{PR}$ is $\left|\overrightarrow{PQ} \times \overrightarrow{PR}\right| = |\langle -1, -7, 6\rangle| = \sqrt{1 + 49 + 36} = \sqrt{86}$, so the area of triangle PQR is $\frac{1}{2}\sqrt{86}$.

31. (a) $\overrightarrow{PQ} = \langle 4, 3, -2\rangle$ and $\overrightarrow{PR} = \langle 5, 5, 1\rangle$, so a vector orthogonal to the plane through P, Q, and R is

$\overrightarrow{PQ} \times \overrightarrow{PR} = \langle (3)(1) - (-2)(5), (-2)(5) - (4)(1), (4)(5) - (3)(5)\rangle = \langle 13, -14, 5\rangle$ [or any scalar mutiple thereof].

(b) The area of the parallelogram determined by $\overrightarrow{PQ}$ and $\overrightarrow{PR}$ is

$\left|\overrightarrow{PQ} \times \overrightarrow{PR}\right| = |\langle 13, -14, 5\rangle| = \sqrt{13^2 + (-14)^2 + 5^2} = \sqrt{390}$, so the area of triangle PQR is $\frac{1}{2}\sqrt{390}$.

32. (a) $\overrightarrow{PQ} = \langle -3, 1, -2\rangle$ and $\overrightarrow{PR} = \langle 1, 4, -7\rangle$, so a vector orthogonal to the plane through P, Q, and R is

$\overrightarrow{PQ} \times \overrightarrow{PR} = \langle (1)(-7) - (-2)(4), (-2)(1) - (-3)(-7), (-3)(4) - (1)(1)\rangle = \langle 1, -23, -13\rangle$ [or any scalar multiple thereof].

(b) The area of the parallelogram determined by $\overrightarrow{PQ}$ and $\overrightarrow{PR}$ is

$\left|\overrightarrow{PQ} \times \overrightarrow{PR}\right| = |\langle 1, -23, -13\rangle| = \sqrt{1 + 529 + 169} = \sqrt{699}$, so the area of triangle PQR is $\frac{1}{2}\sqrt{699}$.

33. By Equation 14, the volume of the parallelepiped determined by $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ is the magnitude of their scalar triple product,

which is $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} 1 & 2 & 3 \\ -1 & 1 & 2 \\ 2 & 1 & 4 \end{vmatrix} = 1\begin{vmatrix} 1 & 2 \\ 1 & 4 \end{vmatrix} - 2\begin{vmatrix} -1 & 2 \\ 2 & 4 \end{vmatrix} + 3\begin{vmatrix} -1 & 1 \\ 2 & 1 \end{vmatrix} = 1(4-2) - 2(-4-4) + 3(-1-2) = 9.$

Thus the volume of the parallelepiped is 9 cubic units.

34. $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{vmatrix} = 1\begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} - 1\begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} + 0\begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} = 0 + 1 + 0 = 1.$

So the volume of the parallelepiped determined by $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ is 1 cubic unit.

35. $\mathbf{a} = \overrightarrow{PQ} = \langle 4, 2, 2\rangle$, $\mathbf{b} = \overrightarrow{PR} = \langle 3, 3, -1\rangle$, and $\mathbf{c} = \overrightarrow{PS} = \langle 5, 5, 1\rangle$.

$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} 4 & 2 & 2 \\ 3 & 3 & -1 \\ 5 & 5 & 1 \end{vmatrix} = 4\begin{vmatrix} 3 & -1 \\ 5 & 1 \end{vmatrix} - 2\begin{vmatrix} 3 & -1 \\ 5 & 1 \end{vmatrix} + 2\begin{vmatrix} 3 & 3 \\ 5 & 5 \end{vmatrix} = 32 - 16 + 0 = 16,$

so the volume of the parallelepiped is 16 cubic units.

36. $\mathbf{a} = \overrightarrow{PQ} = \langle -4, 2, 4\rangle$, $\mathbf{b} = \overrightarrow{PR} = \langle 2, 1, -2\rangle$ and $\mathbf{c} = \overrightarrow{PS} = \langle -3, 4, 1\rangle$.

$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} -4 & 2 & 4 \\ 2 & 1 & -2 \\ -3 & 4 & 1 \end{vmatrix} = -4\begin{vmatrix} 1 & -2 \\ 4 & 1 \end{vmatrix} - 2\begin{vmatrix} 2 & -2 \\ -3 & 1 \end{vmatrix} + 4\begin{vmatrix} 2 & 1 \\ -3 & 4 \end{vmatrix} = -36 + 8 + 44 = 16$, so the volume of the parallelepiped is 16 cubic units.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $\mathbf{u}\cdot(\mathbf{v}\times\mathbf{w}) = \begin{vmatrix} 1 & 5 & -2 \\ 3 & -1 & 0 \\ 5 & 9 & -4 \end{vmatrix} = 1\begin{vmatrix} -1 & 0 \\ 9 & -4 \end{vmatrix} - 5\begin{vmatrix} 3 & 0 \\ 5 & -4 \end{vmatrix} + (-2)\begin{vmatrix} 3 & -1 \\ 5 & 9 \end{vmatrix} = 4 + 60 - 64 = 0$, which says that the volume of the parallelepiped determined by $\mathbf{u}$, $\mathbf{v}$ and $\mathbf{w}$ is 0, and thus these three vectors are coplanar.

38. $\mathbf{u} = \overrightarrow{AB} = \langle 2, -4, 4\rangle$, $\mathbf{v} = \overrightarrow{AC} = \langle 4, -1, -2\rangle$ and $\mathbf{w} = \overrightarrow{AD} = \langle 2, 3, -6\rangle$.

$\mathbf{u}\cdot(\mathbf{v}\times\mathbf{w}) = \begin{vmatrix} 2 & -4 & 4 \\ 4 & -1 & -2 \\ 2 & 3 & -6 \end{vmatrix} = 2\begin{vmatrix} -1 & -2 \\ 3 & -6 \end{vmatrix} - (-4)\begin{vmatrix} 4 & -2 \\ 2 & -6 \end{vmatrix} + 4\begin{vmatrix} 4 & -1 \\ 2 & 3 \end{vmatrix} = 24 - 80 + 56 = 0$, so the volume of the parallelepiped determined by $\mathbf{u}$, $\mathbf{v}$ and $\mathbf{w}$ is 0, which says these vectors lie in the same plane. Therefore, their initial and terminal points A, B, C and D also lie in the same plane.

39. Using the notation of the text, $|\mathbf{r}| = 0.18$ m, $|\mathbf{F}| = 60$ N, and the angle between $\mathbf{r}$ and $\mathbf{F}$ is $\theta = 70° + 10° = 80°$. (Move $\mathbf{F}$ so that both vectors start from the same point.) Then the magnitude of the torque is $|\boldsymbol{\tau}| = |\mathbf{r}\times\mathbf{F}| = |\mathbf{r}|\,|\mathbf{F}|\sin\theta = (0.18)(60)\sin 80° = 10.8\sin 80° \approx 10.6$ N·m.

40. (a) The position vector from the point P to the handle is $\mathbf{r} = \langle 1, 2\rangle$ and has magnitude $|\mathbf{r}| = \sqrt{1^2 + 2^2} = \sqrt{5}$ ft. Since the force vector $\mathbf{F}$ is parallel to the x-axis, the angle between $\mathbf{r}$ and $\mathbf{F}$ is $\theta = \tan^{-1}\left(\frac{2}{1}\right) \approx 63.43°$ and the magnitude of the torque is $|\boldsymbol{\tau}| = |\mathbf{r}\times\mathbf{F}| = |\mathbf{r}|\,|\mathbf{F}|\sin\theta \approx \left(\sqrt{5}\right)(20)\sin 63.43° \approx 40.0$ ft-lb. (Alternatively, we can observe that $\sin\theta = \frac{2}{\sqrt{5}}$, so $|\mathbf{r}|\,|\mathbf{F}|\sin\theta = \sqrt{5}\cdot 20\cdot\frac{2}{\sqrt{5}} = 40$.)

(b) In this case $\mathbf{r} = \overrightarrow{PQ} = \langle 0.6, 0.6\rangle$, so $|\mathbf{r}| = \sqrt{(0.6)^2 + (0.6)^2} = \sqrt{0.72}$ and $\theta = 45°$. The magnitude of the torque is $|\boldsymbol{\tau}| = |\mathbf{r}|\,|\mathbf{F}|\sin\theta = \left(\sqrt{0.72}\right)(20)\sin 45° = \left(\sqrt{0.72}\right)(20)\cdot\frac{\sqrt{2}}{2} = 10\sqrt{1.44} = 12$ ft-lb.

41. Using the notation of the text, $\mathbf{r} = \langle 0, 0.3, 0\rangle$ (measuring in meters) and $\mathbf{F}$ has direction $\langle 0, 3, -4\rangle$. The angle θ between them can be determined by $\cos\theta = \dfrac{\langle 0, 0.3, 0\rangle\cdot\langle 0, 3, -4\rangle}{|\langle 0, 0.3, 0\rangle|\,|\langle 0, 3, -4\rangle|} \;\Rightarrow\; \cos\theta = \dfrac{0.9}{(0.3)(5)} \;\Rightarrow\; \cos\theta = 0.6 \;\Rightarrow$

$\theta = \cos^{-1}(0.6) \approx 53.1°$. Then $|\boldsymbol{\tau}| = |\mathbf{r}|\,|\mathbf{F}|\sin\theta \;\Rightarrow\; 100 \approx 0.3\,|\mathbf{F}|\sin 53.1° \;\Rightarrow\; |\mathbf{F}| \approx \dfrac{100}{0.3\sin 53.1°} \approx 417$ N.

42. Since $|\mathbf{u}\times\mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}|\sin\theta$, $0 \le \theta \le \pi$, $|\mathbf{u}\times\mathbf{v}|$ achieves its maximum value for $\sin\theta = 1 \;\Rightarrow\; \theta = \frac{\pi}{2}$, in which case $|\mathbf{u}\times\mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| = 15$. The minimum value is zero, which occurs when $\sin\theta = 0 \;\Rightarrow\; \theta = 0$ or π, so when $\mathbf{u}$, $\mathbf{v}$ are parallel. Thus, when $\mathbf{u}$ points in the same direction as $\mathbf{v}$, so $\mathbf{u} = 3\mathbf{j}$, $|\mathbf{u}\times\mathbf{v}| = 0$. As $\mathbf{u}$ rotates counterclockwise, $\mathbf{u}\times\mathbf{v}$ is directed in the negative z-direction (by the right-hand rule) and the length increases until $\theta = \frac{\pi}{2}$, in which case $\mathbf{u} = -3\mathbf{i}$ and $|\mathbf{u}\times\mathbf{v}| = 15$. As $\mathbf{u}$ rotates to the negative y-axis, $\mathbf{u}\times\mathbf{v}$ remains pointed in the negative z-direction and the length of $\mathbf{u}\times\mathbf{v}$ decreases to 0, after which the direction of $\mathbf{u}\times\mathbf{v}$ reverses to point in the positive z-direction and $|\mathbf{u}\times\mathbf{v}|$ increases. When $\mathbf{u} = 3\mathbf{i}$ (so $\theta = \frac{\pi}{2}$), $|\mathbf{u}\times\mathbf{v}|$ again reaches its maximum of 15, after which $|\mathbf{u}\times\mathbf{v}|$ decreases to 0 as $\mathbf{u}$ rotates to the positive y-axis.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

43. From Theorem 9 we have $|\mathbf{a}\times\mathbf{b}| = |\mathbf{a}|\,|\mathbf{b}|\sin\theta$, where θ is the angle between $\mathbf{a}$ and $\mathbf{b}$, and from Theorem 12.3.3 we have $\mathbf{a}\cdot\mathbf{b} = |\mathbf{a}|\,|\mathbf{b}|\cos\theta \quad\Rightarrow\quad |\mathbf{a}|\,|\mathbf{b}| = \dfrac{\mathbf{a}\cdot\mathbf{b}}{\cos\theta}$. Substituting the second equation into the first gives $|\mathbf{a}\times\mathbf{b}| = \dfrac{\mathbf{a}\cdot\mathbf{b}}{\cos\theta}\sin\theta$, so $\dfrac{|\mathbf{a}\times\mathbf{b}|}{\mathbf{a}\cdot\mathbf{b}} = \tan\theta$. Here $|\mathbf{a}\times\mathbf{b}| = |\langle 1,2,2\rangle| = \sqrt{1+4+4} = 3$, so $\tan\theta = \dfrac{|\mathbf{a}\times\mathbf{b}|}{\mathbf{a}\cdot\mathbf{b}} = \dfrac{3}{\sqrt{3}} = \sqrt{3} \quad\Rightarrow\quad \theta = 60°$.

44. (a) Let $\mathbf{v} = \langle v_1, v_2, v_3\rangle$. Then

$$\langle 1,2,1\rangle \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 1 \\ v_1 & v_2 & v_3 \end{vmatrix} = \begin{vmatrix} 2 & 1 \\ v_2 & v_3 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 1 & 1 \\ v_1 & v_3 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 1 & 2 \\ v_1 & v_2 \end{vmatrix}\mathbf{k} = (2v_3 - v_2)\,\mathbf{i} - (v_3 - v_1)\,\mathbf{j} + (v_2 - 2v_1)\,\mathbf{k}.$$

If $\langle 1,2,1\rangle \times \mathbf{v} = \langle 3,1,-5\rangle$ then $\langle 2v_3 - v_2, v_1 - v_3, v_2 - 2v_1\rangle = \langle 3,1,-5\rangle \quad\Leftrightarrow\quad 2v_3 - v_2 = 3$ **(1)**, $v_1 - v_3 = 1$ **(2)**, and $v_2 - 2v_1 = -5$ **(3)**. From **(3)** we have $v_2 = 2v_1 - 5$ and from **(2)** we have $v_3 = v_1 - 1$; substitution into **(1)** gives $2(v_1 - 1) - (2v_1 - 5) = 3 \quad\Rightarrow\quad 3 = 3$, so this is a dependent system. If we let $v_1 = a$ then $v_2 = 2a - 5$ and $v_3 = a - 1$, so $\mathbf{v}$ is any vector of the form $\langle a, 2a-5, a-1\rangle$.

(b) If $\langle 1,2,1\rangle \times \mathbf{v} = \langle 3,1,5\rangle$ then $2v_3 - v_2 = 3$ **(1)**, $v_1 - v_3 = 1$ **(2)**, and $v_2 - 2v_1 = 5$ **(3)**. From **(3)** we have $v_2 = 2v_1 + 5$ and from **(2)** we have $v_3 = v_1 - 1$; substitution into **(1)** gives $2(v_1 - 1) - (2v_1 + 5) = 3 \quad\Rightarrow\quad -7 = 3$, so this is an inconsistent system and has no solution.

Alternatively, if we use matrices to solve the system we could show that the determinant is 0 (and hence the system has no solution).

45. (a)



The distance between a point and a line is the length of the perpendicular from the point to the line, here $\left|\overrightarrow{PS}\right| = d$. But referring to triangle PQS, $d = \left|\overrightarrow{PS}\right| = \left|\overrightarrow{QP}\right|\sin\theta = |\mathbf{b}|\sin\theta$. But θ is the angle between $\overrightarrow{QP} = \mathbf{b}$ and $\overrightarrow{QR} = \mathbf{a}$. Thus by Theorem 9, $\sin\theta = \dfrac{|\mathbf{a}\times\mathbf{b}|}{|\mathbf{a}|\,|\mathbf{b}|}$ and so $d = |\mathbf{b}|\sin\theta = \dfrac{|\mathbf{b}|\,|\mathbf{a}\times\mathbf{b}|}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{|\mathbf{a}\times\mathbf{b}|}{|\mathbf{a}|}$.

(b) $\mathbf{a} = \overrightarrow{QR} = \langle -1,\ 2,\ 1\rangle$ and $\mathbf{b} = \overrightarrow{QP} = \langle 1,\ -5,\ -7\rangle$. Then

$\mathbf{a}\times\mathbf{b} = \langle (-2)(-7) - (-1)(-5), (-1)(1) - (-1)(-7), (-1)(-5) - (-2)(1)\rangle = \langle 9, -8, 7\rangle$.

Thus the distance is $d = \dfrac{|\mathbf{a}\times\mathbf{b}|}{|\mathbf{a}|} = \frac{1}{\sqrt{6}}\sqrt{81 + 64 + 49} \quad \sqrt{\frac{194}{6}} \quad \sqrt{\frac{97}{3}}$.

46. (a) The distance between a point and a plane is the length of the perpendicular from the point to the plane, here $\left|\overrightarrow{TP}\right| = d$. But $\overrightarrow{TP}$ is parallel to $\mathbf{b}\times\mathbf{a}$ (because $\mathbf{b}\times\mathbf{a}$ is perpendicular to $\mathbf{b}$ and $\mathbf{a}$) and $d = \left|\overrightarrow{TP}\right| =$ the absolute value of the scalar projection of $\mathbf{c}$ along $\mathbf{b}\times\mathbf{a}$, which is $|\mathbf{c}|\,|\cos\theta|$. (Notice that this is the same



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

setup as the development of the volume of a parallelepiped with $h = |\mathbf{c}|\,|\cos\theta|$). Thus $d = |\mathbf{c}|\,|\cos\theta| = h = V/A$ where $A = |\mathbf{a} \times \mathbf{b}|$, the area of the base. So finally $d = \dfrac{V}{A} = \dfrac{|\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|}{|\mathbf{a} \times \mathbf{b}|}$.

(b) $\mathbf{a} = \overrightarrow{QR} = \langle -1, 2, 0\rangle$, $\mathbf{b} = \overrightarrow{QS} = \langle -1, 0, 3\rangle$ and $\mathbf{c} = \overrightarrow{QP} = \langle 1, 1, 4\rangle$. Then

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} -1 & 2 & 0 \\ -1 & 0 & 3 \\ 1 & 1 & 4 \end{vmatrix} = (-1)\begin{vmatrix} 0 & 3 \\ 1 & 4 \end{vmatrix} - 2\begin{vmatrix} -1 & 3 \\ 1 & 4 \end{vmatrix} + 0 = 17$$

and
$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 2 & 0 \\ -1 & 0 & 3 \end{vmatrix} = \begin{vmatrix} 2 & 0 \\ 0 & 3 \end{vmatrix}\mathbf{i} - \begin{vmatrix} -1 & 0 \\ -1 & 3 \end{vmatrix}\mathbf{j} + \begin{vmatrix} -1 & 2 \\ -1 & 0 \end{vmatrix}\mathbf{k} = 6\,\mathbf{i} + 3\,\mathbf{j} + 2\,\mathbf{k}$$

Thus $d = \dfrac{|\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|}{|\mathbf{a} \times \mathbf{b}|} = \dfrac{17}{\sqrt{36 + 9 + 4}} = \dfrac{17}{7}$.

47. From Theorem 9 we have $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}|\,|\mathbf{b}| \sin\theta$ so

$$\begin{aligned} |\mathbf{a} \times \mathbf{b}|^2 &= |\mathbf{a}|^2\,|\mathbf{b}|^2 \sin^2\theta = |\mathbf{a}|^2\,|\mathbf{b}|^2\left(1 - \cos^2\theta\right) \\ &= |\mathbf{a}|^2\,|\mathbf{b}|^2 - (|\mathbf{a}|\,|\mathbf{b}|\cos\theta)^2 = |\mathbf{a}|^2\,|\mathbf{b}|^2 - (\mathbf{a} \cdot \mathbf{b})^2 \end{aligned}$$

by Theorem 12.3.3.

48. If $\mathbf{a} + \mathbf{b} + \mathbf{c} = \mathbf{0}$ then $\mathbf{b} = -(\mathbf{a} + \mathbf{c})$, so

$$\begin{aligned} \mathbf{a} \times \mathbf{b} &= \mathbf{a} \times [-(\mathbf{a} + \mathbf{c})] = -[\mathbf{a} \times (\mathbf{a} + \mathbf{c})] && \text{by Property 2 of the cross product (with } c = -1) \\ &= -[(\mathbf{a} \times \mathbf{a}) + (\mathbf{a} \times \mathbf{c})] && \text{by Property 3} \\ &= -[\mathbf{0} + (\mathbf{a} \times \mathbf{c})] = -\mathbf{a} \times \mathbf{c} && \text{by Example 2} \\ &= \mathbf{c} \times \mathbf{a} && \text{by Property 1} \end{aligned}$$

Similarly, $\mathbf{a} = -(\mathbf{b} + \mathbf{c})$ so

$$\begin{aligned} \mathbf{c} \times \mathbf{a} &= \mathbf{c} \times [-(\mathbf{b} + \mathbf{c})] = -[\mathbf{c} \times (\mathbf{b} + \mathbf{c})] \\ &= -[(\mathbf{c} \times \mathbf{b}) + (\mathbf{c} \times \mathbf{c})] = -[(\mathbf{c} \times \mathbf{b}) + \mathbf{0}] \\ &= -\mathbf{c} \times \mathbf{b} = \mathbf{b} \times \mathbf{c} \end{aligned}$$

Thus $\mathbf{a} \times \mathbf{b} = \mathbf{b} \times \mathbf{c} = \mathbf{c} \times \mathbf{a}$.

49.
$$\begin{aligned} (\mathbf{a} - \mathbf{b}) \times (\mathbf{a} + \mathbf{b}) &= (\mathbf{a} - \mathbf{b}) \times \mathbf{a} + (\mathbf{a} - \mathbf{b}) \times \mathbf{b} && \text{by Property 3 of the cross product} \\ &= \mathbf{a} \times \mathbf{a} + (-\mathbf{b}) \times \mathbf{a} + \mathbf{a} \times \mathbf{b} + (-\mathbf{b}) \times \mathbf{b} && \text{by Property 4} \\ &= (\mathbf{a} \times \mathbf{a}) - (\mathbf{b} \times \mathbf{a}) + (\mathbf{a} \times \mathbf{b}) - (\mathbf{b} \times \mathbf{b}) && \text{by Property 2 (with } c = -1) \\ &= \mathbf{0} - (\mathbf{b} \times \mathbf{a}) + (\mathbf{a} \times \mathbf{b}) - \mathbf{0} && \text{by Example 2} \\ &= (\mathbf{a} \times \mathbf{b}) + (\mathbf{a} \times \mathbf{b}) && \text{by Property 1} \\ &= 2(\mathbf{a} \times \mathbf{b}) \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

50. Let $\mathbf{a} = \langle a_1, a_2, a_3\rangle$, $\mathbf{b} = \langle b_1, b_2, b_3\rangle$ and $\mathbf{c} = \langle c_1, c_2, c_3\rangle$, so $\mathbf{b} \times \mathbf{c} = \langle b_2c_3 - b_3c_2, b_3c_1 - b_1c_3, b_1c_2 - b_2c_1\rangle$ and

$$\begin{aligned}
\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) &= \langle a_2(b_1c_2 - b_2c_1) - a_3(b_3c_1 - b_1c_3),\ a_3(b_2c_3 - b_3c_2) - a_1(b_1c_2 - b_2c_1),\\
&\qquad a_1(b_3c_1 - b_1c_3) - a_2(b_2c_3 - b_3c_2)\rangle\\
&= \langle a_2b_1c_2 - a_2b_2c_1 - a_3b_3c_1 + a_3b_1c_3,\ a_3b_2c_3 - a_3b_3c_2 - a_1b_1c_2 + a_1b_2c_1,\\
&\qquad a_1b_3c_1 - a_1b_1c_3 - a_2b_2c_3 + a_2b_3c_2\rangle\\
&= \langle (a_2c_2 + a_3c_3)b_1 - (a_2b_2 + a_3b_3)c_1,\ (a_1c_1 + a_3c_3)b_2 - (a_1b_1 + a_3b_3)c_2,\\
&\qquad (a_1c_1 + a_2c_2)b_3 - (a_1b_1 + a_2b_2)c_3\rangle\\
(\star)\quad &= \langle (a_2c_2 + a_3c_3)b_1 - (a_2b_2 + a_3b_3)c_1 + a_1b_1c_1 - a_1b_1c_1,\\
&\qquad (a_1c_1 + a_3c_3)b_2 - (a_1b_1 + a_3b_3)c_2 + a_2b_2c_2 - a_2b_2c_2,\\
&\qquad (a_1c_1 + a_2c_2)b_3 - (a_1b_1 + a_2b_2)c_3 + a_3b_3c_3 - a_3b_3c_3\rangle\\
&= \langle (a_1c_1 + a_2c_2 + a_3c_3)b_1 - (a_1b_1 + a_2b_2 + a_3b_3)c_1,\\
&\qquad (a_1c_1 + a_2c_2 + a_3c_3)b_2 - (a_1b_1 + a_2b_2 + a_3b_3)c_2,\\
&\qquad (a_1c_1 + a_2c_2 + a_3c_3)b_3 - (a_1b_1 + a_2b_2 + a_3b_3)c_3\rangle\\
&= (a_1c_1 + a_2c_2 + a_3c_3)\,\langle b_1, b_2, b_3\rangle - (a_1b_1 + a_2b_2 + a_3b_3)\,\langle c_1, c_2, c_3\rangle\\
&= (\mathbf{a}\cdot\mathbf{c})\mathbf{b} - (\mathbf{a}\cdot\mathbf{b})\mathbf{c}
\end{aligned}$$

$(\star)$ Here we look ahead to see what terms are still needed to arrive at the desired equation. By adding and subtracting the same terms, we don't change the value of the component.

51. $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) + \mathbf{b} \times (\mathbf{c} \times \mathbf{a}) + \mathbf{c} \times (\mathbf{a} \times \mathbf{b})$

$$\begin{aligned}
&= [(\mathbf{a}\cdot\mathbf{c})\mathbf{b} - (\mathbf{a}\cdot\mathbf{b})\mathbf{c}] + [(\mathbf{b}\cdot\mathbf{a})\mathbf{c} - (\mathbf{b}\cdot\mathbf{c})\mathbf{a}] + [(\mathbf{c}\cdot\mathbf{b})\mathbf{a} - (\mathbf{c}\cdot\mathbf{a})\mathbf{b}] \qquad \text{by Exercise 50}\\
&= (\mathbf{a}\cdot\mathbf{c})\mathbf{b} - (\mathbf{a}\cdot\mathbf{b})\mathbf{c} + (\mathbf{a}\cdot\mathbf{b})\mathbf{c} - (\mathbf{b}\cdot\mathbf{c})\mathbf{a} + (\mathbf{b}\cdot\mathbf{c})\mathbf{a} - (\mathbf{a}\cdot\mathbf{c})\mathbf{b} = \mathbf{0}
\end{aligned}$$

52. Let $\mathbf{c} \times \mathbf{d} = \mathbf{v}$. Then

$$\begin{aligned}
(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) &= (\mathbf{a} \times \mathbf{b}) \cdot \mathbf{v} = \mathbf{a} \cdot (\mathbf{b} \times \mathbf{v}) && \text{by Property 5 of the cross product}\\
&= \mathbf{a} \cdot [\mathbf{b} \times (\mathbf{c} \times \mathbf{d})] = \mathbf{a} \cdot [(\mathbf{b}\cdot\mathbf{d})\mathbf{c} - (\mathbf{b}\cdot\mathbf{c})\mathbf{d}] && \text{by Exercise 50}\\
&= (\mathbf{b}\cdot\mathbf{d})(\mathbf{a}\cdot\mathbf{c}) - (\mathbf{b}\cdot\mathbf{c})(\mathbf{a}\cdot\mathbf{d}) && \text{by Properties 3 and 4 of the dot product}\\
&= \begin{vmatrix} \mathbf{a}\cdot\mathbf{c} & \mathbf{b}\cdot\mathbf{c} \\ \mathbf{a}\cdot\mathbf{d} & \mathbf{b}\cdot\mathbf{d} \end{vmatrix}
\end{aligned}$$

53. (a) No. If $\mathbf{a}\cdot\mathbf{b} = \mathbf{a}\cdot\mathbf{c}$, then $\mathbf{a}\cdot(\mathbf{b} - \mathbf{c}) = 0$, so $\mathbf{a}$ is perpendicular to $\mathbf{b} - \mathbf{c}$, which can happen if $\mathbf{b} \neq \mathbf{c}$. For example, let $\mathbf{a} = \langle 1, 1, 1\rangle$, $\mathbf{b} = \langle 1, 0, 0\rangle$ and $\mathbf{c} = \langle 0, 1, 0\rangle$.

(b) No. If $\mathbf{a} \times \mathbf{b} = \mathbf{a} \times \mathbf{c}$ then $\mathbf{a} \times (\mathbf{b} - \mathbf{c}) = \mathbf{0}$, which implies that $\mathbf{a}$ is parallel to $\mathbf{b} - \mathbf{c}$, which of course can happen if $\mathbf{b} \neq \mathbf{c}$.

(c) Yes. Since $\mathbf{a}\cdot\mathbf{c} = \mathbf{a}\cdot\mathbf{b}$, $\mathbf{a}$ is perpendicular to $\mathbf{b} - \mathbf{c}$, by part (a). From part (b), $\mathbf{a}$ is also parallel to $\mathbf{b} - \mathbf{c}$. Thus since $\mathbf{a} \neq \mathbf{0}$ but is both parallel and perpendicular to $\mathbf{b} - \mathbf{c}$, we have $\mathbf{b} - \mathbf{c} = \mathbf{0}$, so $\mathbf{b} = \mathbf{c}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. (a) $\mathbf{k}_i$ is perpendicular to $\mathbf{v}_i$ if $i \neq j$ by the definition of $\mathbf{k}_i$ and Theorem 8.

(b) $\mathbf{k}_1 \cdot \mathbf{v}_1 = \dfrac{\mathbf{v}_2 \times \mathbf{v}_3}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} \cdot \mathbf{v}_1 = \dfrac{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} = 1$

$\mathbf{k}_2 \cdot \mathbf{v}_2 = \dfrac{\mathbf{v}_3 \times \mathbf{v}_1}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} \cdot \mathbf{v}_2 = \dfrac{\mathbf{v}_2 \cdot (\mathbf{v}_3 \times \mathbf{v}_1)}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} = \dfrac{(\mathbf{v}_2 \times \mathbf{v}_3) \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} = 1$ [by Property 5 of the cross product]

$\mathbf{k}_3 \cdot \mathbf{v}_3 = \dfrac{(\mathbf{v}_1 \times \mathbf{v}_2) \cdot \mathbf{v}_3}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} = \dfrac{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} = 1$ [by Property 5]

(c) $\mathbf{k}_1 \cdot (\mathbf{k}_2 \times \mathbf{k}_3) = \mathbf{k}_1 \cdot \left(\dfrac{\mathbf{v}_3 \times \mathbf{v}_1}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} \times \dfrac{\mathbf{v}_1 \times \mathbf{v}_2}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} \right) = \dfrac{\mathbf{k}_1}{[\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)]^2} \cdot [(\mathbf{v}_3 \times \mathbf{v}_1) \times (\mathbf{v}_1 \times \mathbf{v}_2)]$

$= \dfrac{\mathbf{k}_1}{[\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)]^2} \cdot ([(\mathbf{v}_3 \times \mathbf{v}_1) \cdot \mathbf{v}_2]\,\mathbf{v}_1 - [(\mathbf{v}_3 \times \mathbf{v}_1) \cdot \mathbf{v}_1]\,\mathbf{v}_2)$ [by Exercise 50]

But $(\mathbf{v}_3 \times \mathbf{v}_1) \cdot \mathbf{v}_1 = 0$ since $\mathbf{v}_3 \times \mathbf{v}_1$ is orthogonal to $\mathbf{v}_1$, and

$(\mathbf{v}_3 \times \mathbf{v}_1) \cdot \mathbf{v}_2 = \mathbf{v}_2 \cdot (\mathbf{v}_3 \times \mathbf{v}_1) = (\mathbf{v}_2 \times \mathbf{v}_3) \cdot \mathbf{v}_1 = \mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)$. Thus

$$\mathbf{k}_1 \cdot (\mathbf{k}_2 \times \mathbf{k}_3) = \frac{\mathbf{k}_1}{[\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)]^2} \cdot [\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)]\,\mathbf{v}_1 = \frac{\mathbf{k}_1 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} = \frac{1}{\mathbf{v}_1 \cdot (\mathbf{v}_2 \times \mathbf{v}_3)} \quad \text{[by part (b)]}$$

DISCOVERY PROJECT The Geometry of a Tetrahedron

1. Set up a coordinate system so that vertex S is at the origin, $R = (0, y_1, 0)$, $Q = (x_2, y_2, 0)$, $P = (x_3, y_3, z_3)$.

Then $\overrightarrow{SR} = \langle 0, y_1, 0 \rangle$, $\overrightarrow{SQ} = \langle x_2, y_2, 0 \rangle$, $\overrightarrow{SP} = \langle x_3, y_3, z_3 \rangle$, $\overrightarrow{QR} = \langle -x_2, y_1 - y_2, 0 \rangle$, and $\overrightarrow{QP} = \langle x_3 - x_2, y_3 - y_2, z_3 \rangle$.

Let

$$\mathbf{v}_S = \overrightarrow{QR} \times \overrightarrow{QP} = (y_1 z_3 - y_2 z_3)\,\mathbf{i} + x_2 z_3\,\mathbf{j} + (-x_2 y_3 - x_3 y_1 + x_3 y_2 + x_2 y_1)\,\mathbf{k}$$

Then $\mathbf{v}_S$ is an outward normal to the face opposite vertex S. Similarly,

$\mathbf{v}_R = \overrightarrow{SQ} \times \overrightarrow{SP} = y_2 z_3\,\mathbf{i} - x_2 z_3\,\mathbf{j} + (x_2 y_3 - x_3 y_2)\,\mathbf{k}$, $\mathbf{v}_Q = \overrightarrow{SP} \times \overrightarrow{SR} = -y_1 z_3\,\mathbf{i} + x_3 y_1\,\mathbf{k}$, and

$\mathbf{v}_P = \overrightarrow{SR} \times \overrightarrow{SQ} = -x_2 y_1\,\mathbf{k} \quad \Rightarrow \quad \mathbf{v}_S + \mathbf{v}_R + \mathbf{v}_Q + \mathbf{v}_P = \mathbf{0}$. Now

$$\begin{aligned} |\mathbf{v}_S| &= \text{ area of the parallelogram determined by } \overrightarrow{QR} \text{ and } \overrightarrow{QP} \\ &= 2\,(\text{area of triangle } RQP) = 2|\mathbf{v}_1| \end{aligned}$$

So $\mathbf{v}_S = 2\mathbf{v}_1$, and similarly $\mathbf{v}_R = 2\mathbf{v}_2$, $\mathbf{v}_Q = 2\mathbf{v}_3$, $\mathbf{v}_P = 2\mathbf{v}_4$. Thus $\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3 + \mathbf{v}_4 = \mathbf{0}$.

2. (a) Let $S = (x_0, y_0, z_0)$, $R = (x_1, y_1, z_1)$, $Q = (x_2, y_2, z_2)$, $P = (x_3, y_3, z_3)$ be the four vertices. Then

$$\begin{aligned} \text{Volume} &= \tfrac{1}{3}(\text{distance from } S \text{ to plane } RQP) \times (\text{area of triangle } RQP) \\ &= \frac{1}{3} \frac{\left| \mathbf{N} \cdot \overrightarrow{SR} \right|}{|\mathbf{N}|} \cdot \frac{1}{2} \left| \overrightarrow{RQ} \times \overrightarrow{RP} \right| \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

where $\mathbf{N}$ is a vector which is normal to the face RQP. Thus $\mathbf{N} = \overrightarrow{RQ} \times \overrightarrow{RP}$. Therefore

$$V = \left|\tfrac{1}{6}\left(\overrightarrow{RQ} \times \overrightarrow{RP}\right) \cdot \overrightarrow{SR}\right| = \frac{1}{6}\left|\begin{vmatrix} x_0 - x & y_0 - y_1 & z_0 - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix}\right|$$

(b) Using the formula from part (a), $V = \dfrac{1}{6}\left|\begin{vmatrix} 1-1 & 1-2 & 1-3 \\ 1-1 & 1-2 & 2-3 \\ 3-1 & -1-2 & 2-3 \end{vmatrix}\right| = \dfrac{1}{6}\left|2(1-2)\right| = \dfrac{1}{3}$.

3. We define a vector $\mathbf{v}_1$ to have length equal to the area of the face opposite vertex P, so we can say $|\mathbf{v}_1| = A$, and direction perpendicular to the face and pointing outward, as in Problem 1. Similarly, we define $\mathbf{v}_2$, $\mathbf{v}_3$, and $\mathbf{v}_4$ so that $|\mathbf{v}_2| = B$, $|\mathbf{v}_3| = C$, and $|\mathbf{v}_4| = D$ and with the analogous directions. From Problem 1, we know $\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3 + \mathbf{v}_4 = \mathbf{0} \;\Rightarrow$

$\mathbf{v}_4 = -(\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3) \;\Rightarrow\; |\mathbf{v}_4| = |-(\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3)| = |\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3| \;\Rightarrow\; |\mathbf{v}_4|^2 = |\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3|^2 \;\Rightarrow$

$$\begin{aligned}\mathbf{v}_4 \cdot \mathbf{v}_4 &= (\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3) \cdot (\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3) \\ &= \mathbf{v}_1 \cdot \mathbf{v}_1 + \mathbf{v}_1 \cdot \mathbf{v}_2 + \mathbf{v}_1 \cdot \mathbf{v}_3 + \mathbf{v}_2 \cdot \mathbf{v}_1 + \mathbf{v}_2 \cdot \mathbf{v}_2 + \mathbf{v}_2 \cdot \mathbf{v}_3 + \mathbf{v}_3 \cdot \mathbf{v}_1 + \mathbf{v}_3 \cdot \mathbf{v}_2 + \mathbf{v}_3 \cdot \mathbf{v}_3\end{aligned}$$

Since the vertex S is trirectangular, we know the three faces meeting at S are mutually perpendicular, so the vectors $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ are also mutually perpendicular. Therefore, $\mathbf{v}_i \cdot \mathbf{v}_j = 0$ for $i \neq j$ and $i, j \in \{1, 2, 3\}$. Thus we have

$\mathbf{v}_4 \cdot \mathbf{v}_4 = \mathbf{v}_1 \cdot \mathbf{v}_1 + \mathbf{v}_2 \cdot \mathbf{v}_2 + \mathbf{v}_3 \cdot \mathbf{v}_3 \;\Rightarrow\; |\mathbf{v}_4|^2 = |\mathbf{v}_1|^2 + |\mathbf{v}_2|^2 + |\mathbf{v}_3|^2 \;\Rightarrow\; D^2 = A^2 + B^2 + C^2$.

Another method: We introduce a coordinate system, as shown. Recall that the area of the parallelogram spanned by two vectors is equal to the length of their cross product, so since $\mathbf{u} \times \mathbf{v} = \langle -q, r, 0\rangle \times \langle -q, 0, p\rangle = \langle pr, pq, qr\rangle$, we have $|\mathbf{u} \times \mathbf{v}| = \sqrt{(pr)^2 + (pq)^2 + (qr)^2}$, and therefore



$$\begin{aligned}D^2 &= \left(\tfrac{1}{2}|\mathbf{u} \times \mathbf{v}|\right)^2 = \tfrac{1}{4}[(pr)^2 + (pq)^2 + (qr)^2] \\ &= \left(\tfrac{1}{2}pr\right)^2 + \left(\tfrac{1}{2}pq\right)^2 + \left(\tfrac{1}{2}qr\right)^2 = A^2 + B^2 + C^2.\end{aligned}$$

A third method: We draw a line from S perpendicular to QR, as shown. Now $D = \frac{1}{2}ch$, so $D^2 = \frac{1}{4}c^2h^2$. Substituting $h^2 = p^2 + k^2$, we get $D^2 = \frac{1}{4}c^2(p^2 + k^2) = \frac{1}{4}c^2p^2 + \frac{1}{4}c^2k^2$. But $C = \frac{1}{2}ck$, so $D^2 = \frac{1}{4}c^2p^2 + C^2$. Now substituting $c^2 = q^2 + r^2$ gives $D^2 = \frac{1}{4}p^2q^2 + \frac{1}{4}q^2r^2 + C^2 = A^2 + B^2 + C^2$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12.5 Equations of Lines and Planes

1. (a) True; each of the first two lines has a direction vector parallel to the direction vector of the third line, so these vectors are each scalar multiples of the third direction vector. Then the first two direction vectors are also scalar multiples of each other, so these vectors, and hence the two lines, are parallel.

   (b) False; for example, the x- and y-axes are both perpendicular to the z-axis, yet the x- and y-axes are not parallel.

   (c) True; each of the first two planes has a normal vector parallel to the normal vector of the third plane, so these two normal vectors are parallel to each other and the planes are parallel.

   (d) False; for example, the xy- and yz-planes are not parallel, yet they are both perpendicular to the xz-plane.

   (e) False; the x- and y-axes are not parallel, yet they are both parallel to the plane $z = 1$.

   (f) True; if each line is perpendicular to a plane, then the lines' direction vectors are both parallel to a normal vector for the plane. Thus, the direction vectors are parallel to each other and the lines are parallel.

   (g) False; the planes $y = 1$ and $z = 1$ are not parallel, yet they are both parallel to the x-axis.

   (h) True; if each plane is perpendicular to a line, then any normal vector for each plane is parallel to a direction vector for the line. Thus, the normal vectors are parallel to each other and the planes are parallel.

   (i) True; see Figure 9 and the accompanying discussion.

   (j) False; they can be skew, as in Example 3.

   (k) True. Consider any normal vector for the plane and any direction vector for the line. If the normal vector is perpendicular to the direction vector, the line and plane are parallel. Otherwise, the vectors meet at an angle θ, $0^\circ \le \theta < 90^\circ$, and the line will intersect the plane at an angle $90^\circ - \theta$.

2. For this line, we have $\mathbf{r}_0 = 6\,\mathbf{i} - 5\,\mathbf{j} + 2\,\mathbf{k}$ and $\mathbf{v} = \mathbf{i} + 3\,\mathbf{j} - \frac{2}{3}\,\mathbf{k}$, so a vector equation is $\mathbf{r} = \mathbf{r}_0 + t\,\mathbf{v} = (6\,\mathbf{i} - 5\,\mathbf{j} + 2\,\mathbf{k}) + t\left(\mathbf{i} + 3\,\mathbf{j} - \frac{2}{3}\,\mathbf{k}\right) = (6 + t)\,\mathbf{i} + (-5 + 3t)\,\mathbf{j} + \left(2 - \frac{2}{3}t\right)\mathbf{k}$ and parametric equations are $x = 6 + t$, $y = -5 + 3t$, $z = 2 - \frac{2}{3}t$.

3. For this line, we have $\mathbf{r}_0 = 2\,\mathbf{i} + 2.4\,\mathbf{j} + 3.5\,\mathbf{k}$ and $\mathbf{v} = 3\,\mathbf{i} + 2\,\mathbf{j} - \mathbf{k}$, so a vector equation is $\mathbf{r} = \mathbf{r}_0 + t\,\mathbf{v} = (2\,\mathbf{i} + 2.4\,\mathbf{j} + 3.5\,\mathbf{k}) + t(3\,\mathbf{i} + 2\,\mathbf{j} - \mathbf{k}) = (2 + 3t)\,\mathbf{i} + (2.4 + 2t)\,\mathbf{j} + (3.5 - t)\,\mathbf{k}$ and parametric equations are $x = 2 + 3t$, $y = 2.4 + 2t$, $z = 3.5 - t$.

4. This line has the same direction as the given line, $\mathbf{v} = 2\,\mathbf{i} - 3\,\mathbf{j} + 9\,\mathbf{k}$. Here $\mathbf{r}_0 = 14\,\mathbf{j} - 10\,\mathbf{k}$, so a vector equation is $\mathbf{r} = (14\,\mathbf{j} - 10\,\mathbf{k}) + t(2\,\mathbf{i} - 3\,\mathbf{j} + 9\,\mathbf{k}) = 2t\,\mathbf{i} + (14 - 3t)\,\mathbf{j} + (-10 + 9t)\,\mathbf{k}$ and parametric equations are $x = 2t$, $y = 14 - 3t$, $z = -10 + 9t$.

5. A line perpendicular to the given plane has the same direction as a normal vector to the plane, such as $\mathbf{n} = \langle 1, 3, 1\rangle$. So $\mathbf{r}_0 = \mathbf{i} + 6\,\mathbf{k}$, and we can take $\mathbf{v} = \mathbf{i} + 3\,\mathbf{j} + \mathbf{k}$. Then a vector equation is $\mathbf{r} = (\mathbf{i} + 6\,\mathbf{k}) + t(\mathbf{i} + 3\,\mathbf{j} + \mathbf{k}) = (1 + t)\,\mathbf{i} + 3t\,\mathbf{j} + (6 + t)\,\mathbf{k}$, and parametric equations are $x = 1 + t$, $y = 3t$, $z = 6 + t$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

6. The vector $\mathbf{v} = \langle 4-0, 3-0, -1-0\rangle = \langle 4, 3, -1\rangle$ is parallel to the line. Letting $P_0 = (0, 0, 0)$, parametric equations are $x = 0 + 4 \cdot t = 4t$, $y = 0 + 3 \cdot t = 3t$, $z = 0 + (-1) \cdot t = -t$, while symmetric equations are $\dfrac{x}{4} = \dfrac{y}{3} = \dfrac{z}{-1}$ or $\dfrac{x}{4} = \dfrac{y}{3} = -z$.

7. The vector $\mathbf{v} = \langle 2-0, 1-\frac{1}{2}, -3-1\rangle = \langle 2, \frac{1}{2}, -4\rangle$ is parallel to the line. Letting $P_0 = (2, 1, -3)$, parametric equations are $x = 2 + 2t$, $y = 1 + \frac{1}{2}t$, $z = -3 - 4t$, while symmetric equations are $\dfrac{x-2}{2} = \dfrac{y-1}{1/2} = \dfrac{z+3}{-4}$ or $\dfrac{x-2}{2} = 2y - 2 = \dfrac{z+3}{-4}$.

8. $\mathbf{v} = \langle 2.6-1, 1.2-2.4, 0.3-4.6\rangle = \langle 1.6, -1.2, -4.3\rangle$, and letting $P_0 = (1, 2.4, 4.6)$, parametric equations are $x = 1 + 1.6t$, $y = 2.4 - 1.2t$, $z = 4.6 - 4.3t$, while symmetric equations are $\dfrac{x-1}{1.6} = \dfrac{y-2.4}{-1.2} = \dfrac{z-4.6}{-4.3}$.

9. $\mathbf{v} = \langle 3-(-8), -2-1, 4-4\rangle = \langle 11, -3, 0\rangle$, and letting $P_0 = (-8, 1, 4)$, parametric equations are $x = -8 + 11t$, $y = 1 - 3t$, $z = 4 + 0t = 4$, while symmetric equations are $\dfrac{x+8}{11} = \dfrac{y-1}{-3}$, $z = 4$. Notice here that the direction number $c = 0$, so rather than writing $\dfrac{z-4}{0}$ in the symmetric equation we must write the equation $z = 4$ separately.

10. $\mathbf{v} = (\mathbf{i}+\mathbf{j}) \times (\mathbf{j}+\mathbf{k}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix} = \mathbf{i} - \mathbf{j} + \mathbf{k}$ is the direction of the line perpendicular to both $\mathbf{i}+\mathbf{j}$ and $\mathbf{j}+\mathbf{k}$.

With $P_0 = (2, 1, 0)$, parametric equations are $x = 2 + t$, $y = 1 - t$, $z = t$ and symmetric equations are $x - 2 = \dfrac{y-1}{-1} = z$ or $x - 2 = 1 - y = z$.

11. The given line $\dfrac{x}{2} = \dfrac{y}{3} = \dfrac{z+1}{1}$ has direction $\mathbf{v} = \langle 2, 3, 1\rangle$. Taking $(-6, 2, 3)$ as P_0, parametric equations are $x = -6 + 2t$, $y = 2 + 3t$, $z = 3 + t$ and symmetric equations are $\dfrac{x+6}{2} = \dfrac{y-2}{3} = z - 3$.

12. Setting $z = 0$ we see that $(1, 0, 0)$ satisfies the equations of both planes, so they do in fact have a line of intersection. The line is perpendicular to the normal vectors of both planes, so a direction vector for the line is $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, 2, 3\rangle \times \langle 1, -1, 1\rangle = \langle 5, 2, -3\rangle$. Taking the point $(1, 0, 0)$ as P_0, parametric equations are $x = 1 + 5t$, $y = 2t$, $z = -3t$, and symmetric equations are $\dfrac{x-1}{5} = \dfrac{y}{2} = \dfrac{z}{-3}$.

13. Direction vectors of the lines are $\mathbf{v}_1 = \langle -2-(-4), 0-(-6), -3-1\rangle = \langle 2, 6, -4\rangle$ and $\mathbf{v}_2 = \langle 5-10, 3-18, 14-4\rangle = \langle -5, -15, 10\rangle$, and since $\mathbf{v}_2 = -\frac{5}{2}\mathbf{v}_1$, the direction vectors and thus the lines are parallel.

14. Direction vectors of the lines are $\mathbf{v}_1 = \langle 3, -3, 1\rangle$ and $\mathbf{v}_2 = \langle 1, -4, -12\rangle$. Since $\mathbf{v}_1 \cdot \mathbf{v}_2 = 3 + 12 - 12 \neq 0$, the vectors and thus the lines are not perpendicular.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. (a) The line passes through the point $(1, -5, 6)$ and a direction vector for the line is $\langle -1, 2, -3 \rangle$, so symmetric equations for the line are $\dfrac{x-1}{-1} = \dfrac{y+5}{2} = \dfrac{z-6}{-3}$.

(b) The line intersects the xy-plane when $z = 0$, so we need $\dfrac{x-1}{-1} = \dfrac{y+5}{2} = \dfrac{0-6}{-3}$ or $\dfrac{x-1}{-1} = 2 \Rightarrow x = -1$, $\dfrac{y+5}{2} = 2 \Rightarrow y = -1$. Thus the point of intersection with the xy-plane is $(-1, -1, 0)$. Similarly for the yz-plane, we need $x = 0 \Rightarrow 1 = \dfrac{y+5}{2} = \dfrac{z-6}{-3} \Rightarrow y = -3, z = 3$. Thus the line intersects the yz-plane at $(0, -3, 3)$. For the xz-plane, we need $y = 0 \Rightarrow \dfrac{x-1}{-1} = \dfrac{5}{2} = \dfrac{z-6}{-3} \Rightarrow x = -\frac{3}{2}, z = -\frac{3}{2}$. So the line intersects the xz-plane at $\left(-\frac{3}{2}, 0, -\frac{3}{2}\right)$.

16. (a) A vector normal to the plane $x - y + 3z = 7$ is $\mathbf{n} = \langle 1, -1, 3 \rangle$, and since the line is to be perpendicular to the plane, $\mathbf{n}$ is also a direction vector for the line. Thus parametric equations of the line are $x = 2 + t, y = 4 - t, z = 6 + 3t$.

(b) On the xy-plane, $z = 0$. So $z = 6 + 3t = 0 \Rightarrow t = -2$ in the parametric equations of the line, and therefore $x = 0$ and $y = 6$, giving the point of intersection $(0, 6, 0)$. For the yz-plane, $x = 0$ so we get the same point of interesection: $(0, 6, 0)$. For the xz-plane, $y = 0$ which implies $t = 4$, so $x = 6$ and $z = 18$ and the point of intersection is $(6, 0, 18)$.

17. From Equation 4, the line segment from $\mathbf{r}_0 = 6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}$ to $\mathbf{r}_1 = 7\,\mathbf{i} + 6\,\mathbf{j}$ has vector equation

$$\begin{aligned}\mathbf{r}(t) &= (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)(6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}) + t(7\,\mathbf{i} + 6\,\mathbf{j}) \\ &= (6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}) - t(6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}) + t(7\,\mathbf{i} + 6\,\mathbf{j}) \\ &= (6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}) + t(\mathbf{i} + 7\,\mathbf{j} - 9\,\mathbf{k}), \quad 0 \le t \le 1.\end{aligned}$$

18. From Equation 4, the line segment from $\mathbf{r}_0 = -2\,\mathbf{i} + 18\,\mathbf{j} + 31\,\mathbf{k}$ to $\mathbf{r}_1 = 11\,\mathbf{i} - 4\,\mathbf{j} + 48\,\mathbf{k}$ has vector equation

$$\begin{aligned}\mathbf{r}(t) &= (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)(-2\,\mathbf{i} + 18\,\mathbf{j} + 31\,\mathbf{k}) + t(11\,\mathbf{i} - 4\,\mathbf{j} + 48\,\mathbf{k}) \\ &= (-2\,\mathbf{i} + 18\,\mathbf{j} + 31\,\mathbf{k}) + t(13\,\mathbf{i} - 22\,\mathbf{j} + 17\,\mathbf{k}), \quad 0 \le t \le 1.\end{aligned}$$

The corresponding parametric equations are $x = -2 + 13t, y = 18 - 22t, z = 31 + 17t, 0 \le t \le 1$.

19. Since the direction vectors $\langle 2, -1, 3 \rangle$ and $\langle 4, -2, 5 \rangle$ are not scalar multiples of each other, the lines aren't parallel. For the lines to intersect, we must be able to find one value of t and one value of s that produce the same point from the respective parametric equations. Thus we need to satisfy the following three equations: $3 + 2t = 1 + 4s, 4 - t = 3 - 2s$, $1 + 3t = 4 + 5s$. Solving the last two equations we get $t = 1, s = 0$ and checking, we see that these values don't satisfy the first equation. Thus the lines aren't parallel and don't intersect, so they must be skew lines.

20. Since the direction vectors are $\mathbf{v}_1 = \langle -12, 9, -3 \rangle$ and $\mathbf{v}_2 = \langle 8, -6, 2 \rangle$, we have $\mathbf{v}_1 = -\frac{3}{2}\mathbf{v}_2$ so the lines are parallel.

21. Since the direction vectors $\langle 1, -2, -3 \rangle$ and $\langle 1, 3, -7 \rangle$ aren't scalar multiples of each other, the lines aren't parallel. Parametric equations of the lines are L_1: $x = 2 + t, y = 3 - 2t, z = 1 - 3t$ and L_2: $x = 3 + s, y = -4 + 3s, z = 2 - 7s$. Thus, for the lines to intersect, the three equations $2 + t = 3 + s, 3 - 2t = -4 + 3s$, and $1 - 3t = 2 - 7s$ must be satisfied simultaneously. Solving the first two equations gives $t = 2, s = 1$ and checking, we see that these values do satisfy the third equation, so the lines intersect when $t = 2$ and $s = 1$, that is, at the point $(4, -1, -5)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

22. The direction vectors $\langle 1, -1, 3\rangle$ and $\langle 2, -2, 7\rangle$ are not parallel, so neither are the lines. Parametric equations for the lines are L_1: $x = t$, $y = 1 - t$, $z = 2 + 3t$ and L_2: $x = 2 + 2s$, $y = 3 - 2s$, $z = 7s$. Thus, for the lines to interesect, the three equations $t = 2 + 2s$, $1 - t = 3 - 2s$, and $2 + 3t = 7s$ must be satisfied simultaneously. Solving the last two equations gives $t = -10$, $s = -4$ and checking, we see that these values don't satisfy the first equation. Thus the lines aren't parallel and don't intersect, so they must be skew.

23. Since the plane is perpendicular to the vector $\langle 1, -2, 5\rangle$, we can take $\langle 1, -2, 5\rangle$ as a normal vector to the plane. $(0, 0, 0)$ is a point on the plane, so setting $a = 1$, $b = -2$, $c = 5$ and $x_0 = 0$, $y_0 = 0$, $z_0 = 0$ in Equation 7 gives $1(x - 0) + (-2)(y - 0) + 5(z - 0) = 0$ or $x - 2y + 5z = 0$ as an equation of the plane.

24. $2\,\mathbf{i} + \mathbf{j} - \mathbf{k} = \langle 2, 1, -1\rangle$ is a normal vector to the plane and $(5, 3, 5)$ is a point on the plane, so setting $a = 2$, $b = 1$, $c = -1$, $x_0 = 5$, $y_0 = 3$, $z_0 = 5$ in Equation 7 gives $2(x - 5) + 1(y - 3) + (-1)(z - 5) = 0$ or $2x + y - z = 8$ as an equation of the plane.

25. $\mathbf{i} + 4\,\mathbf{j} + \mathbf{k} = \langle 1, 4, 1\rangle$ is a normal vector to the plane and $\left(-1, \frac{1}{2}, 3\right)$ is a point on the plane, so setting $a = 1$, $b = 4$, $c = 1$, $x_0 = -1$, $y_0 = \frac{1}{2}$, $z_0 = 3$ in Equation 7 gives $1[x - (-1)] + 4\left(y - \frac{1}{2}\right) + 1(z - 3) = 0$ or $x + 4y + z = 4$ as an equation of the plane.

26. Since the line is perpendicular to the plane, its direction vector $\langle 3, -1, 4\rangle$ is a normal vector to the plane. The point $(2, 0, 1)$ is on the plane, so an equation of the plane is $3(x - 2) + (-1)(y - 0) + 4(z - 1) = 0$ or $3x - y + 4z = 10$.

27. Since the two planes are parallel, they will have the same normal vectors. So we can take $\mathbf{n} = \langle 5, -1, -1\rangle$, and an equation of the plane is $5(x - 1) - 1[y - (-1)] - 1[z - (-1)] = 0$ or $5x - y - z = 7$.

28. Since the two planes are parallel, they will have the same normal vectors. A normal vector for the plane $z = x + y$ or $x + y - z = 0$ is $\mathbf{n} = \langle 1, 1, -1\rangle$, and an equation of the desired plane is $1(x - 3) + 1[y - (-2)] - 1(z - 8) = 0$ or $x + y - z = -7$.

29. Since the two planes are parallel, they will have the same normal vectors. So we can take $\mathbf{n} = \langle 1, 1, 1\rangle$, and an equation of the plane is $1(x - 1) + 1\left(y - \frac{1}{2}\right) + 1\left(z - \frac{1}{3}\right) = 0$ or $x + y + z = \frac{11}{6}$ or $6x + 6y + 6z = 11$.

30. First, a normal vector for the plane $5x + 2y + z = 1$ is $\mathbf{n} = \langle 5, 2, 1\rangle$. A direction vector for the line is $\mathbf{v} = \langle 1, -1, -3\rangle$, and since $\mathbf{n} \cdot \mathbf{v} = 0$ we know the line is perpendicular to $\mathbf{n}$ and hence parallel to the plane. Thus, there is a parallel plane which contains the line. By putting $t = 0$, we know that the point $(1, 2, 4)$ is on the line and hence the new plane. We can use the same normal vector $\mathbf{n} = \langle 5, 2, 1\rangle$, so an equation of the plane is $5(x - 1) + 2(y - 2) + 1(z - 4) = 0$ or $5x + 2y + z = 13$.

31. The vector from $(0, 1, 1)$ to $(1, 0, 1)$, namely $\mathbf{a} = \langle 1 - 0, 0 - 1, 1 - 1\rangle = \langle 1, -1, 0\rangle$, and the vector from $(0, 1, 1)$ to $(1, 1, 0)$, $\mathbf{b} = \langle 1 - 0, 1 - 1, 0 - 1\rangle = \langle 1, 0, -1\rangle$, both lie in the plane, so $\mathbf{a} \times \mathbf{b}$ is a normal vector to the plane. Thus, we can take $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle (-1)((-1) - (0)(0), (0)(1) - (1)(-1), (1)(0) - (-1)(1)\rangle = \langle 1, 1, 1\rangle$. If P_0 is the point $(0, 1, 1)$, an equation of the plane is $1(x - 0) + 1(y - 1) + 1(z - 1) = 0$ or $x + y + z = 2$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. Here the vectors $\mathbf{a} = \langle 3, -2, 1\rangle$ and $\mathbf{b} = \langle 1, 1, 1\rangle$ lie in the plane, so $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle (-2)(1) - (1)(1), (1)(1) - (3)(1), (3)(1) - (-2)(1)\rangle = \langle -3, -2, 5\rangle$ is a normal vector to the plane. We can take the origin as P_0, so an equation of the plane is $-3(x - 0) - 2(y - 0) + 5(z - 0) = 0$ or $-3x - 2y + 5z = 0$ or $3x + 2y - 5z = 0$.

33. Here the vectors $\mathbf{a} = \langle 3 - 2, -8 - 1, 6 - 2\rangle = \langle 1, -9, 4\rangle$ and $\mathbf{b} = \langle -2 - 2, -3 - 1, 1 - 2\rangle = \langle -4, -4, -1\rangle$ lie in the plane, so a normal vector to the plane is $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle 9 + 16, -16 + 1, -4 - 36\rangle = \langle 25, -15, -40\rangle$ and an equation of the plane is $25(x - 2) - 15(y - 1) - 40(z - 2) = 0$ or $25x - 15y - 40z = -45$ or $5x - 3y - 8z = -9$.

34. The vectors $\mathbf{a} = \langle -2 - 3, -2 - 0, 3 - (-1)\rangle = \langle -5, -2, 4\rangle$ and $\mathbf{b} = \langle 7 - 3, 1 - 0, -4 - (-1)\rangle = \langle 4, 1, -3\rangle$ lie in the plane, so a normal vector to the plane is $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle 6 - 4, 16 - 15, -5 + 8\rangle = \langle 2, 1, 3\rangle$ and an equation of the plane is $2(x - 3) + 1(y - 0) + 3[z - (-1)] = 0$ or $2x + y + 3z = 3$.

35. If we first find two nonparallel vectors in the plane, their cross product will be a normal vector to the plane. Since the given line lies in the plane, its direction vector $\mathbf{a} = \langle -1, 2, -3\rangle$ is one vector in the plane. We can verify that the given point $(3, 5, -1)$ does not lie on this line, so to find another nonparallel vector $\mathbf{b}$ which lies in the plane, we can pick any point on the line and find a vector connecting the points. If we put $t = 0$, we see that $(4, -1, 0)$ is on the line, so $\mathbf{b} = \langle 4 - 3, -1 - 5, 0 - (-1)\rangle = \langle 1, -6, 1\rangle$ and $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle 2 - 18, -3 + 1, 6 - 2\rangle = \langle -16, -2, 4\rangle$. Thus, an equation of the plane is $-16(x - 3) - 2(y - 5) + 4[z - (-1)] = 0$ or $-16x - 2y + 4z = -62$ or $8x + y - 2z = 31$.

36. Since the line $\dfrac{x}{3} = \dfrac{y + 4}{1} = \dfrac{z}{2}$ lies in the plane, its direction vector $\mathbf{a} = \langle 3, 1, 2\rangle$ is parallel to the plane. The point $(0, -4, 0)$ is on the line (put $t = 0$ in the corresponding parametric equations), and we can verify that the given point $(6, -1, 3)$ in the plane is not on the line. The vector connecting these two points, $\mathbf{b} = \langle 6, 3, 3\rangle$, is therefore parallel to the plane, but not parallel to $\mathbf{a}$. Then $\mathbf{a} \times \mathbf{b} = \langle 3 - 6, 12 - 9, 9 - 6\rangle = \langle -3, 3, 3\rangle$ is a normal vector to the plane, and an equation of the plane is $-3(x - 0) + 3[y - (-4)] + 3(z - 0) = 0$ or $-3x + 3y + 3z = -12$ or $x - y - z = 4$.

37. Normal vectors for the given planes are $\mathbf{n}_1 = \langle 1, 2, 3\rangle$ and $\mathbf{n}_2 = \langle 2, -1, 1\rangle$. A direction vector, then, for the line of intersection is $\mathbf{a} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 2 + 3, 6 - 1, -1 - 4\rangle = \langle 5, 5, -5\rangle$, and $\mathbf{a}$ is parallel to the desired plane. Another vector parallel to the plane is the vector connecting any point on the line of intersection to the given point $(3, 1, 4)$ in the plane. Setting $z = 0$, the equations of the planes reduce to $x + 2y = 1$ and $2x - y = -3$ with simultaneous solution $x = -1$ and $y = 1$. So a point on the line is $(-1, 1, 0)$ and another vector parallel to the plane is $\mathbf{b} = \langle 3 - (-1), 1 - 1, 4 - 0\rangle = \langle 4, 0, 4\rangle$. Then a normal vector to the plane is $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle 20 - 0, -20 - 20, 0 - 20\rangle = \langle 20, -40, -20\rangle$. Equivalently, we can take $\langle 1, -2, -1\rangle$ as a normal vector, and an equation of the plane is $1(x - 3) - 2(y - 1) - 1(z - 4) = 0$ or $x - 2y - z = -3$.

38. The points $(0, -2, 5)$ and $(-1, 3, 1)$ lie in the desired plane, so the vector $\mathbf{v}_1 = \langle -1, 5, -4\rangle$ connecting them is parallel to the plane. The desired plane is perpendicular to the plane $2z - 5x + 4y$ or $5x + 4y - 2z = 0$ and for perpendicular planes.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

a normal vector for one plane is parallel to the other plane, so $\mathbf{v}_2 = \langle 5, 4, -2\rangle$ is also parallel to the desired plane. A normal vector to the desired plane is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle -10 + 16, -20 - 2, -4 - 25\rangle = \langle 6, -22, -29\rangle$. Taking $(x_0, y_0, z_0) = (0, -2, 5)$, the equation we are looking for is $6(x - 0) - 22(y + 2) - 29(z - 5) = 0$ or $6x - 22y - 29z = -101$.

39. If a plane is perpendicular to two other planes, its normal vector is perpendicular to the normal vectors of the other two planes. Thus $\langle 2, 1, -2\rangle \times \langle 1, 0, 3\rangle = \langle 3 - 0, -2 - 6, 0 - 1\rangle = \langle 3, -8, -1\rangle$ is a normal vector to the desired plane. The point $(1, 5, 1)$ lies on the plane, so an equation is $3(x - 1) - 8(y - 5) - (z - 1) = 0$ or $3x - 8y - z = -38$.

40. $\mathbf{n}_1 = \langle 1, 0, -1\rangle$ and $\mathbf{n}_2 = \langle 0, 1, 2\rangle$. Setting $z = 0$, it is easy to see that $(1, 3, 0)$ is a point on the line of intersection of $x - z = 1$ and $y + 2z = 3$. The direction of this line is $\mathbf{v}_1 = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, -2, 1\rangle$. A second vector parallel to the desired plane is $\mathbf{v}_2 = \langle 1, 1, -2\rangle$, since it is perpendicular to $x + y - 2z = 1$. Therefore, a normal of the plane in question is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 4 - 1, 1 + 2, 1 + 2\rangle = \langle 3, 3, 3\rangle$, or we can use $\langle 1, 1, 1\rangle$. Taking $(x_0, y_0, z_0) = (1, 3, 0)$, the equation we are looking for is $(x - 1) + (y - 3) + z = 0 \quad \Leftrightarrow \quad x + y + z = 4$.

41. To find the x-intercept we set $y = z = 0$ in the equation $2x + 5y + z = 10$ and obtain $2x = 10 \quad \Rightarrow \quad x = 5$ so the x-intercept is $(5, 0, 0)$. When $x = z = 0$ we get $5y = 10 \quad \Rightarrow \quad y = 2$, so the y-intercept is $(0, 2, 0)$. Setting $x = y = 0$ gives $z = 10$, so the z-intercept is $(0, 0, 10)$ and we graph the portion of the plane that lies in the first octant.



42. To find the x-intercept we set $y = z = 0$ in the equation $3x + y + 2z = 6$ and obtain $3x = 6 \quad \Rightarrow \quad x = 2$ so the x-intercept is $(2, 0, 0)$. When $x = z = 0$ we get $y = 6$ so the y-intercept is $(0, 6, 0)$. Setting $x = y = 0$ gives $2z = 6 \quad \Rightarrow \quad z = 3$, so the z-intercept is $(0, 0, 3)$. The figure shows the portion of the plane that lies in the first octant.



43. Setting $y = z = 0$ in the equation $6x - 3y + 4z = 6$ gives $6x = 6 \quad \Rightarrow \quad x = 1$, when $x = z = 0$ we have $-3y = 6 \quad \Rightarrow \quad y = -2$, and $x = y = 0$ implies $4z = 6 \quad \Rightarrow \quad z = \frac{3}{2}$, so the intercepts are $(1, 0, 0)$, $(0, -2, 0)$, and $(0, 0, \frac{3}{2})$. The figure shows the portion of the plane cut off by the coordinate planes.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. Setting $y = z = 0$ in the equation $6x + 5y - 3z = 15$ gives $6x = 15 \Rightarrow x = \frac{5}{2}$, when $x = z = 0$ we have $5y = 15 \Rightarrow y = 3$, and $x = y = 0$ implies $-3z = 15 \Rightarrow z = -5$, so the intercepts are $(\frac{5}{2}, 0, 0)$, $(0, 3, 0)$, and $(0, 0, -5)$. The figure shows the portion of the plane cut off by the coordinate planes.



45. Substitute the parametric equations of the line into the equation of the plane: $x + 2y - z = 7 \Rightarrow (2 - 2t) + 2(3t) - (1 + t) = 7 \Rightarrow 3t + 1 = 7 \Rightarrow t = 2$. Therefore, the point of intersection of the line and the plane is given by $x = 2 - 2(2) = -2$, $y = 3(2) = 6$, and $z = 1 + 2 = 3$, that is, the point $(-2, 6, 3)$.

46. Substitute the parametric equations of the line into the equation of the plane: $3(t - 1) - (1 + 2t) + 2(3 - t) = 5 \Rightarrow -t + 2 = 5 \Rightarrow t = -3$. Therefore, the point of intersection of the line and the plane is given by $x = -3 - 1 = -4$, $y = 1 + 2(-3) = -5$, and $z = 3 - (-3) = 6$, that is, the point $(-4, -5, 6)$.

47. Parametric equations for the line are $x = \frac{1}{5}t$, $y = 2t$, $z = t - 2$ and substitution into the equation of the plane gives $10(\frac{1}{5}t) - 7(2t) + 3(t - 2) + 24 = 0 \Rightarrow -9t + 18 = 0 \Rightarrow t = 2$. Thus $x = \frac{1}{5}(2) = \frac{2}{5}$, $y = 2(2) = 4$, $z = 2 - 2 = 0$ and the point of intersection is $(\frac{2}{5}, 4, 0)$.

48. A direction vector for the line through $(-3, 1, 0)$ and $(-1, 5, 6)$ is $\mathbf{v} = \langle 2, 4, 6 \rangle$ and, taking $P_0 = (-3, 1, 0)$, parametric equations for the line are $x = -3 + 2t$, $y = 1 + 4t$, $z = 6t$. Substitution of the parametric equations into the equation of the plane gives $2(-3 + 2t) + (1 + 4t) - (6t) = -2 \Rightarrow 2t - 5 = -2 \Rightarrow t = \frac{3}{2}$. Then $x = -3 + 2(\frac{3}{2}) = 0$, $y = 1 + 4(\frac{3}{2}) = 7$, and $z = 6(\frac{3}{2}) = 9$, and the point of intersection is $(0, 7, 9)$.

49. Setting $x = 0$, we see that $(0, 1, 0)$ satisfies the equations of both planes, so that they do in fact have a line of intersection. $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, 1, 1 \rangle \times \langle 1, 0, 1 \rangle = \langle 1, 0, -1 \rangle$ is the direction of this line. Therefore, direction numbers of the intersecting line are 1, 0, -1.

50. The angle between the two planes is the same as the angle between their normal vectors. The normal vectors of the two planes are $\langle 1, 1, 1 \rangle$ and $\langle 1, 2, 3 \rangle$. The cosine of the angle θ between these two planes is

$$\cos\theta = \frac{\langle 1,1,1\rangle \cdot \langle 1,2,3\rangle}{|\langle 1,1,1\rangle||\langle 1,2,3\rangle|} = \frac{1+2+3}{\sqrt{1+1+1}\sqrt{1+4+9}} = \frac{6}{\sqrt{42}} = \sqrt{\frac{6}{7}}.$$

51. Normal vectors for the planes are $\mathbf{n}_1 = \langle 1, 4, -3 \rangle$ and $\mathbf{n}_2 = \langle -3, 6, 7 \rangle$. The normals aren't parallel (they are not scalar multiples of each other), so neither are the planes. But $\mathbf{n}_1 \cdot \mathbf{n}_2 = -3 + 24 - 21 = 0$, so the normals, and thus the planes, are perpendicular.

52. Normal vectors for the planes are $\mathbf{n}_1 = \langle 9, -3, 6 \rangle$ and $\mathbf{n}_2 = \langle 6, -2, 4 \rangle$ (the plane's equation is $6x - 2y + 4z = 0$). Since $\mathbf{n}_1 = \frac{3}{2}\mathbf{n}_2$, the normals, and thus the planes, are parallel.

53. Normal vectors for the planes are $\mathbf{n}_1 = \langle 1, 2, -1 \rangle$ and $\mathbf{n}_2 = \langle 2, -2, 1 \rangle$. The normals are not parallel (they are not scalar multiples of each other), so neither are the planes. Furthermore, $\mathbf{n}_1 \cdot \mathbf{n}_2 = 2 - 4 - 1 = -3 \neq 0$, so the planes aren't

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

perpendicular. The angle between the planes is the same as the angle between the normals, given by

$$\cos\theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\,|\mathbf{n}_2|} = \frac{-3}{\sqrt{6}\sqrt{9}} = -\frac{1}{\sqrt{6}} \quad \Rightarrow \quad \theta = \cos^{-1}\left(-\frac{1}{\sqrt{6}}\right) \approx 114.1°.$$

54. Normal vectors for the planes are $\mathbf{n}_1 = \langle 1, -1, 3\rangle$ and $\mathbf{n}_2 = \langle 3, 1, -1\rangle$. The normals are not parallel, so neither are the planes. Since $\mathbf{n}_1 \cdot \mathbf{n}_2 = 3 - 1 - 3 = -1 \neq 0$, the planes aren't perpendicular. The angle between the planes is given by

$$\cos\theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\,|\mathbf{n}_2|} = \frac{-1}{\sqrt{11}\sqrt{11}} = -\frac{1}{11} \quad \Rightarrow \quad \theta = \cos^{-1}\left(-\tfrac{1}{11}\right) \approx 95.2°.$$

55. The planes are $2x - 3y - z = 0$ and $4x - 6y - 2z = 3$ with normal vectors $\mathbf{n}_1 = \langle 2, -3, -1\rangle$ and $\mathbf{n}_2 = \langle 4, -6, -2\rangle$. Since $\mathbf{n}_2 = 2\mathbf{n}_1$, the normals, and thus the planes, are parallel.

56. The normals are $\mathbf{n}_1 = \langle 5, 2, 3\rangle$ and $\mathbf{n}_2 = \langle 4, -1, -6\rangle$ which are not scalar multiples of each other, so the planes aren't parallel. Since $\mathbf{n}_1 \cdot \mathbf{n}_2 = 20 - 2 - 18 = 0$, the normals, and thus the planes, are perpendicular.

57. (a) To find a point on the line of intersection, set one of the variables equal to a constant, say $z = 0$. (This will fail if the line of intersection does not cross the xy-plane; in that case, try setting x or y equal to 0.) The equations of the two planes reduce to $x + y = 1$ and $x + 2y = 1$. Solving these two equations gives $x = 1$, $y = 0$. Thus a point on the line is $(1, 0, 0)$. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes, so we can take $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, 1, 1\rangle \times \langle 1, 2, 2\rangle = \langle 2 - 2, 1 - 2, 2 - 1\rangle = \langle 0, -1, 1\rangle$. By Equations 2, parametric equations for the line are $x = 1$, $y = -t$, $z = t$.

(b) The angle between the planes satisfies $\cos\theta = \dfrac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\,|\mathbf{n}_2|} = \dfrac{1 + 2 + 2}{\sqrt{3}\sqrt{9}} = \dfrac{5}{3\sqrt{3}}$. Therefore $\theta = \cos^{-1}\left(\dfrac{5}{3\sqrt{3}}\right) \approx 15.8°$.

58. (a) If we set $z = 0$ then the equations of the planes reduce to $3x - 2y = 1$ and $2x + y = 3$ and solving these two equations gives $x = 1$, $y = 1$. Thus a point on the line of intersection is $(1, 1, 0)$. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes, so let $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 3, -2, 1\rangle \times \langle 2, 1, -3\rangle = \langle 5, 11, 7\rangle$. By Equations 2, parametric equations for the line are $x = 1 + 5t$, $y = 1 + 11t$, $z = 7t$.

(b) $\cos\theta = \dfrac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\,|\mathbf{n}_2|} = \dfrac{6 - 2 - 3}{\sqrt{14}\sqrt{14}} = \dfrac{1}{14} \quad \Rightarrow \quad \theta = \cos^{-1}\left(\frac{1}{14}\right) \approx 85.9°$.

59. Setting $z = 0$, the equations of the two planes become $5x - 2y = 1$ and $4x + y = 6$. Solving these two equations gives $x = 1$, $y = 2$ so a point on the line of intersection is $(1, 2, 0)$. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes. So we can use $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 5, -2, -2\rangle \times \langle 4, 1, 1\rangle = \langle 0, -13, 13\rangle$ or equivalently we can take $\mathbf{v} = \langle 0, -1, 1\rangle$, and symmetric equations for the line are $x = 1$, $\dfrac{y - 2}{-1} = \dfrac{z}{1}$ or $x = 1$, $y - 2 = -z$.

60. If we set $z = 0$ then the equations of the planes reduce to $2x - y - 5 = 0$ and $4x + 3y - 5 = 0$ and solving these two equations gives $x = 2$, $y = -1$. Thus a point on the line of intersection is $(2, -1, 0)$. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes, so take $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 2, -1, -1\rangle \times \langle 4, 3, -1\rangle = \langle 4, -2, 10\rangle$ or equivalently we can take $\mathbf{v} = \langle 2, -1, 5\rangle$. Symmetric equations for the line are $\dfrac{x - 2}{2} = \dfrac{y + 1}{-1} = \dfrac{z}{5}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. The distance from a point (x, y, z) to $(1, 0, -2)$ is $d_1 = \sqrt{(x-1)^2 + y^2 + (z+2)^2}$ and the distance from (x, y, z) to $(3, 4, 0)$ is $d_2 = \sqrt{(x-3)^2 + (y-4)^2 + z^2}$. The plane consists of all points (x, y, z) where $d_1 = d_2 \Rightarrow d_1^2 = d_2^2 \Leftrightarrow (x-1)^2 + y^2 + (z+2)^2 = (x-3)^2 + (y-4)^2 + z^2 \Leftrightarrow$

$x^2 - 2x + y^2 + z^2 + 4z + 5 = x^2 - 6x + y^2 - 8y + z^2 + 25 \Leftrightarrow 4x + 8y + 4z = 20$ so an equation for the plane is $4x + 8y + 4z = 20$ or equivalently $x + 2y + z = 5$.

Alternatively, you can argue that the segment joining points $(1, 0, -2)$ and $(3, 4, 0)$ is perpendicular to the plane and the plane includes the midpoint of the segment.

62. The distance from a point (x, y, z) to $(2, 5, 5)$ is $d_1 = \sqrt{(x-2)^2 + (y-5)^2 + (z-5)^2}$ and the distance from (x, y, z) to $(-6, 3, 1)$ is $d_2 = \sqrt{(x+6)^2 + (y-3)^2 + (z-1)^2}$. The plane consists of all points (x, y, z) where $d_1 = d_2 \Rightarrow d_1^2 = d_2^2 \Leftrightarrow (x-2)^2 + (y-5)^2 + (z-5)^2 = (x+6)^2 + (y-3)^2 + (z-1)^2 \Leftrightarrow$

$x^2 - 4x + y^2 - 10y + z^2 - 10z + 54 = x^2 + 12x + y^2 - 6y + z^2 - 2z + 46 \Leftrightarrow 16x + 4y + 8z = 8$ so an equation for the plane is $16x + 4y + 8z = 8$ or equivalently $4x + y + 2z = 2$.

63. The plane contains the points $(a, 0, 0)$, $(0, b, 0)$ and $(0, 0, c)$. Thus the vectors $\mathbf{a} = \langle -a, b, 0\rangle$ and $\mathbf{b} = \langle -a, 0, c\rangle$ lie in the plane, and $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle bc - 0, 0 + ac, 0 + ab\rangle = \langle bc, ac, ab\rangle$ is a normal vector to the plane. The equation of the plane is therefore $bcx + acy + abz = abc + 0 + 0$ or $bcx + acy + abz = abc$. Notice that if $a \neq 0$, $b \neq 0$ and $c \neq 0$ then we can rewrite the equation as $\dfrac{x}{a} + \dfrac{y}{b} + \dfrac{z}{c} = 1$. This is a good equation to remember!

64. (a) For the lines to intersect, we must be able to find one value of t and one value of s satisfying the three equations $1 + t = 2 - s$, $1 - t = s$ and $2t = 2$. From the third we get $t = 1$, and putting this in the second gives $s = 0$. These values of s and t do satisfy the first equation, so the lines intersect at the point $P_0 = (1 + 1, 1 - 1, 2(1)) = (2, 0, 2)$.

(b) The direction vectors of the lines are $\langle 1, -1, 2\rangle$ and $\langle -1, 1, 0\rangle$, so a normal vector for the plane is $\langle -1, 1, 0\rangle \times \langle 1, -1, 2\rangle = \langle 2, 2, 0\rangle$ and it contains the point $(2, 0, 2)$. Then an equation of the plane is $2(x - 2) + 2(y - 0) + 0(z - 2) = 0 \Leftrightarrow x + y = 2$.

65. Two vectors which are perpendicular to the required line are the normal of the given plane, $\langle 1, 1, 1\rangle$, and a direction vector for the given line, $\langle 1, -1, 2\rangle$. So a direction vector for the required line is $\langle 1, 1, 1\rangle \times \langle 1, -1, 2\rangle = \langle 3, -1, -2\rangle$. Thus L is given by $\langle x, y, z\rangle = \langle 0, 1, 2\rangle + t\langle 3, -1, -2\rangle$, or in parametric form, $x = 3t$, $y = 1 - t$, $z = 2 - 2t$.

66. Let L be the given line. Then $(1, 1, 0)$ is the point on L corresponding to $t = 0$. L is in the direction of $\mathbf{a} = \langle 1, -1, 2\rangle$ and $\mathbf{b} = \langle -1, 0, 2\rangle$ is the vector joining $(1, 1, 0)$ and $(0, 1, 2)$. Then

$\mathbf{b} - \text{proj}_{\mathbf{a}}\,\mathbf{b} = \langle -1, 0, 2\rangle - \dfrac{\langle 1, -1, 2\rangle \cdot \langle -1, 0, 2\rangle}{1^2 + (-1)^2 + 2^2}\langle 1, -1, 2\rangle = \langle -1, 0, 2\rangle - \frac{1}{2}\langle 1, -1, 2\rangle = \langle -\frac{3}{2}, \frac{1}{2}, 1\rangle$ is a direction vector for the required line. Thus $2\langle -\frac{3}{2}, \frac{1}{2}, 1\rangle = \langle -3, 1, 2\rangle$ is also a direction vector, and the line has parametric equations $x = -3t$, $y = 1 + t$, $z = 2 + 2t$. (Notice that this is the same line as in Exercise 65.)

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

67. Let P_i have normal vector $\mathbf{n}_i$. Then $\mathbf{n}_1 = \langle 3, 6, -3\rangle$, $\mathbf{n}_2 = \langle 4, -12, 8\rangle$, $\mathbf{n}_3 = \langle 3, -9, 6\rangle$, $\mathbf{n}_4 = \langle 1, 2, -1\rangle$. Now $\mathbf{n}_1 = 3\mathbf{n}_4$, so $\mathbf{n}_1$ and $\mathbf{n}_4$ are parallel, and hence P_1 and P_4 are parallel; similarly P_2 and P_3 are parallel because $\mathbf{n}_2 = \frac{4}{3}\mathbf{n}_3$. However, $\mathbf{n}_1$ and $\mathbf{n}_2$ are not parallel (so not all four planes are parallel). Notice that the point $(2, 0, 0)$ lies on both P_1 and P_4, so these two planes are identical. The point $\left(\frac{5}{4}, 0, 0\right)$ lies on P_2 but not on P_3, so these are different planes.

68. Let L_i have direction vector $\mathbf{v}_i$. Rewrite the symmetric equations for L_3 as $\dfrac{x-1}{1/2} = \dfrac{y-1}{-1/4} = \dfrac{z+1}{1}$; then $\mathbf{v}_1 = \langle 6, -3, 12\rangle$, $\mathbf{v}_2 = \langle 2, 1, 4\rangle$, $\mathbf{v}_3 = \left\langle \frac{1}{2}, -\frac{1}{4}, 1\right\rangle$, and $\mathbf{v}_4 = \langle 4, 2, 8\rangle$. $\mathbf{v}_1 = 12\mathbf{v}_3$, so L_1 and L_3 are parallel. $\mathbf{v}_4 = 2\mathbf{v}_2$, so L_2 and L_4 are parallel. (Note that L_1 and L_2 are not parallel.) L_1 contains the point $(1, 1, 5)$, but this point does not lie on L_3, so they're not identical. $(3, 1, 5)$ lies on L_4 and also on L_2 (for $t = 1$), so L_2 and L_4 are the same line.

69. Let $Q = (1, 3, 4)$ and $R = (2, 1, 1)$, points on the line corresponding to $t = 0$ and $t = 1$. Let $P = (4, 1, -2)$. Then $\mathbf{a} = \overrightarrow{QR} = \langle 1, -2, -3\rangle$, $\mathbf{b} = \overrightarrow{QP} = \langle 3, -2, -6\rangle$. The distance is

$$d = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \frac{|\langle 1, -2, -3\rangle \times \langle 3, -2, -6\rangle|}{|\langle 1, -2, -3\rangle|} = \frac{|\langle 6, -3, 4\rangle|}{|\langle 1, -2, -3\rangle|} = \frac{\sqrt{6^2 + (-3)^2 + 4^2}}{\sqrt{1^2 + (-2)^2 + (-3)^2}} = \frac{\sqrt{61}}{\sqrt{14}} = \sqrt{\frac{61}{14}}.$$

70. Let $Q = (0, 6, 3)$ and $R = (2, 4, 4)$, points on the line corresponding to $t = 0$ and $t = 1$. Let $P = (0, 1, 3)$. Then $\mathbf{a} = \overrightarrow{QR} = \langle 2, -2, 1\rangle$ and $\mathbf{b} = \overrightarrow{QP} = \langle 0, -5, 0\rangle$. The distance is

$$d = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \frac{|\langle 2, -2, 1\rangle \times \langle 0, -5, 0\rangle|}{|\langle 2, -2, 1\rangle|} = \frac{|\langle 5, 0, -10\rangle|}{|\langle 2, -2, 1\rangle|} = \frac{\sqrt{5^2 + 0^2 + (-10)^2}}{\sqrt{2^2 + (-2)^2 + 1^2}} = \frac{\sqrt{125}}{\sqrt{9}} = \frac{5\sqrt{5}}{3}.$$

71. By Equation 9, the distance is $D = \dfrac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}} = \dfrac{|3(1) + 2(-2) + 6(4) - 5|}{\sqrt{3^2 + 2^2 + 6^2}} = \dfrac{|18|}{\sqrt{49}} = \dfrac{18}{7}$.

72. By Equation 9, the distance is $D = \dfrac{|1(-6) - 2(3) - 4(5) - 8|}{\sqrt{1^2 + (-2)^2 + (-4)^2}} = \dfrac{|-40|}{\sqrt{21}} = \dfrac{40}{\sqrt{21}}$.

73. Put $y = z = 0$ in the equation of the first plane to get the point $(2, 0, 0)$ on the plane. Because the planes are parallel, the distance D between them is the distance from $(2, 0, 0)$ to the second plane. By Equation 9,

$$D = \frac{|4(2) - 6(0) + 2(0) - 3|}{\sqrt{4^2 + (-6)^2 + (2)^2}} = \frac{5}{\sqrt{56}} = \frac{5}{2\sqrt{14}} \text{ or } \frac{5\sqrt{14}}{28}.$$

74. Put $x = y = 0$ in the equation of the first plane to get the point $(0, 0, 0)$ on the plane. Because the planes are parallel the distance D between them is the distance from $(0, 0, 0)$ to the second plane $3x - 6y + 9z - 1 = 0$. By Equation 9,

$$D = \frac{|3(0) - 6(0) + 9(0) - 1|}{\sqrt{3^2 + (-6)^2 + 9^2}} = \frac{1}{\sqrt{126}} = \frac{1}{3\sqrt{14}}.$$

75. The distance between two parallel planes is the same as the distance between a point on one of the planes and the other plane. Let $P_0 = (x_0, y_0, z_0)$ be a point on the plane given by $ax + by + cz + d_1 = 0$. Then $ax_0 + by_0 + cz_0 + d_1 = 0$ and the

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

distance between P_0 and the plane given by $ax + by + cz + d_2 = 0$ is, from Equation 9,

$$D = \frac{|ax_0 + by_0 + cz_0 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|-d_1 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}.$$

76. The planes must have parallel normal vectors, so if $ax + by + cz + d = 0$ is such a plane, then for some $t \neq 0$, $\langle a, b, c\rangle = t\langle 1, 2, -2\rangle = \langle t, 2t, -2t\rangle$. So this plane is given by the equation $x + 2y - 2z + k = 0$, where $k = d/t$. By Exercise 75, the distance between the planes is $2 = \dfrac{|1 - k|}{\sqrt{1^2 + 2^2 + (-2)^2}} \quad \Leftrightarrow \quad 6 = |1 - k| \quad \Leftrightarrow \quad k = 7 \text{ or } -5$. So the desired planes have equations $x + 2y - 2z = 7$ and $x + 2y - 2z = -5$.

77. L_1: $x = y = z \quad \Rightarrow \quad x = y$ **(1)**. L_2: $x + 1 = y/2 = z/3 \quad \Rightarrow \quad x + 1 = y/2$ **(2)**. The solution of **(1)** and **(2)** is $x = y = -2$. However, when $x = -2$, $x = z \quad \Rightarrow \quad z = -2$, but $x + 1 = z/3 \quad \Rightarrow \quad z = -3$, a contradiction. Hence the lines do not intersect. For L_1, $\mathbf{v}_1 = \langle 1, 1, 1\rangle$, and for L_2, $\mathbf{v}_2 = \langle 1, 2, 3\rangle$, so the lines are not parallel. Thus the lines are skew lines. If two lines are skew, they can be viewed as lying in two parallel planes and so the distance between the skew lines would be the same as the distance between these parallel planes. The common normal vector to the planes must be perpendicular to both $\langle 1, 1, 1\rangle$ and $\langle 1, 2, 3\rangle$, the direction vectors of the two lines. So set $\mathbf{n} = \langle 1, 1, 1\rangle \times \langle 1, 2, 3\rangle = \langle 3 - 2, -3 + 1, 2 - 1\rangle = \langle 1, -2, 1\rangle$. From above, we know that $(-2, -2, -2)$ and $(-2, -2, -3)$ are points of L_1 and L_2 respectively. So in the notation of Equation 8, $1(-2) - 2(-2) + 1(-2) + d_1 = 0 \quad \Rightarrow \quad d_1 = 0$ and $1(-2) - 2(-2) + 1(-3) + d_2 = 0 \quad \Rightarrow \quad d_2 = 1$.

By Exercise 75, the distance between these two skew lines is $D = \dfrac{|0 - 1|}{\sqrt{1 + 4 + 1}} = \dfrac{1}{\sqrt{6}}$.

Alternate solution (without reference to planes): A vector which is perpendicular to both of the lines is $\mathbf{n} = \langle 1, 1, 1\rangle \times \langle 1, 2, 3\rangle = \langle 1, -2, 1\rangle$. Pick any point on each of the lines, say $(-2, -2, -2)$ and $(-2, -2, -3)$, and form the vector $\mathbf{b} = \langle 0, 0, 1\rangle$ connecting the two points. The distance between the two skew lines is the absolute value of the scalar projection of $\mathbf{b}$ along $\mathbf{n}$, that is, $D = \dfrac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \dfrac{|1 \cdot 0 - 2 \cdot 0 + 1 \cdot 1|}{\sqrt{1 + 4 + 1}} = \dfrac{1}{\sqrt{6}}$.

78. First notice that if two lines are skew, they can be viewed as lying in two parallel planes and so the distance between the skew lines would be the same as the distance between these parallel planes. The common normal vector to the planes must be perpendicular to both $\mathbf{v}_1 = \langle 1, 6, 2\rangle$ and $\mathbf{v}_2 = \langle 2, 15, 6\rangle$, the direction vectors of the two lines respectively. Thus set $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 36 - 30, 4 - 6, 15 - 12\rangle = \langle 6, -2, 3\rangle$. Setting $t = 0$ and $s = 0$ gives the points $(1, 1, 0)$ and $(1, 5, -2)$. So in the notation of Equation 8, $6 - 2 + 0 + d_1 = 0 \quad \Rightarrow \quad d_1 = -4$ and $6 - 10 - 6 + d_2 = 0 \quad \Rightarrow \quad d_2 = 10$.

Then by Exercise 75, the distance between the two skew lines is given by $D = \dfrac{|-4 - 10|}{\sqrt{36 + 4 + 9}} = \dfrac{14}{7} = 2$.

Alternate solution (without reference to planes): We already know that the direction vectors of the two lines are $\mathbf{v}_1 = \langle 1, 6, 2\rangle$ and $\mathbf{v}_2 = \langle 2, 15, 6\rangle$. Then $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 6, -2, 3\rangle$ is perpendicular to both lines. Pick any point on each of the lines, say $(1, 1, 0)$ and $(1, 5, -2)$, and form the vector $\mathbf{b} = \langle 0, 4, -2\rangle$ connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of $\mathbf{b}$ along $\mathbf{n}$, that is,

$$D = \frac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \frac{1}{\sqrt{36 + 4 + 9}}\,|0 - 8 - 6| = \frac{14}{7} = 2.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

79. A direction vector for L_1 is $\mathbf{v}_1 = \langle 2, 0, -1 \rangle$ and a direction vector for L_2 is $\mathbf{v}_2 = \langle 3, 2, 2 \rangle$. These vectors are not parallel so neither are the lines. Parametric equations for the lines are L_1: $x = 2t$, $y = 0$, $z = -t$, and L_2: $x = 1 + 3s$, $y = -1 + 2s$, $z = 1 + 2s$. No values of t and s satisfy these equations simultaneously, so the lines don't intersect and hence are skew. We can view the lines as lying in two parallel planes; a common normal vector to the planes is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 2, -7, 4 \rangle$. Line L_1 passes through the origin, so $(0, 0, 0)$ lies on one of the planes, and $(1, -1, 1)$ is a point on L_2 and therefore on the other plane. Equations of the planes then are $2x - 7y + 4z = 0$ and $2x - 7y + 4z - 13 = 0$, and by Exercise 75, the distance between the two skew lines is $D = \dfrac{|0 - (-13)|}{\sqrt{4 + 49 + 16}} = \dfrac{13}{\sqrt{69}}$.

Alternate solution (without reference to planes): Direction vectors of the two lines are $\mathbf{v}_1 = \langle 2, 0, -1 \rangle$ and $\mathbf{v}_2 = \langle 3, 2, 2 \rangle$. Then $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 2, -7, 4 \rangle$ is perpendicular to both lines. Pick any point on each of the lines, say $(0, 0, 0)$ and $(1, -1, 1)$, and form the vector $\mathbf{b} = \langle 1, -1, 1 \rangle$ connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of $\mathbf{b}$ along $\mathbf{n}$, that is, $D = \dfrac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \dfrac{|2 + 7 + 4|}{\sqrt{4 + 49 + 16}} = \dfrac{13}{\sqrt{69}}$.

80. A direction vector for the line L_1 is $\mathbf{v}_1 = \langle 1, 2, 2 \rangle$. A normal vector for the plane P_1 is $\mathbf{n}_1 = \langle 1, -1, 2 \rangle$. The vector from the point $(0, 0, 1)$ to $(3, 2, -1)$, $\langle 3, 2, -2 \rangle$, is parallel to the plane P_2, as is the vector from $(0, 0, 1)$ to $(1, 2, 1)$, namely $\langle 1, 2, 0 \rangle$. Thus a normal vector for P_2 is $\langle 3, 2, -2 \rangle \times \langle 1, 2, 0 \rangle = \langle 4, -2, 4 \rangle$, or we can use $\mathbf{n}_2 = \langle 2, -1, 2 \rangle$, and a direction vector for the line L_2 of intersection of these planes is $\mathbf{v}_2 = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, -1, 2 \rangle \times \langle 2, -1, 2 \rangle = \langle 0, 2, 1 \rangle$. Notice that the point $(3, 2, -1)$ lies on both planes, so it also lies on L_2. The lines are skew, so we can view them as lying in two parallel planes; a common normal vector to the planes is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle -2, -1, 2 \rangle$. Line L_1 passes through the point $(1, 2, 6)$, so $(1, 2, 6)$ lies on one of the planes, and $(3, 2, -1)$ is a point on L_2 and therefore on the other plane. Equations of the planes then are $-2x - y + 2z - 8 = 0$ and $-2x - y + 2z + 10 = 0$, and by Exercise 75, the distance between the lines is $D = \dfrac{|-8 - 10|}{\sqrt{4 + 1 + 4}} = \dfrac{18}{3} = 6$.

Alternatively, direction vectors for the lines are $\mathbf{v}_1 = \langle 1, 2, 2 \rangle$ and $\mathbf{v}_2 = \langle 0, 2, 1 \rangle$, so $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle -2, -1, 2 \rangle$ is perpendicular to both lines. Pick any point on each of the lines, say $(1, 2, 6)$ and $(3, 2, -1)$, and form the vector $\mathbf{b} = \langle 2, 0, -7 \rangle$ connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of $\mathbf{b}$ along $\mathbf{n}$, that is, $D = \dfrac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \dfrac{|-4 + 0 - 14|}{\sqrt{4 + 1 + 4}} = \dfrac{18}{3} = 6$.

81. (a) A direction vector from tank A to tank B is $\langle 765 - 325, 675 - 810, 599 - 561 \rangle = \langle 440, -135, 38 \rangle$. Taking tank A's position $(325, 810, 561)$ as the initial point, parametric equations for the line of sight are $x = 325 + 440t$, $y = 810 - 135t$, $z = 561 + 38t$ for $0 \le t \le 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) We divide the line of sight into 5 equal segments, corresponding to $\Delta t = 0.2$, and compute the elevation from the z-component of the parametric equations in part (a):

t	$z = 561 + 38t$	terrain elevation
0	561.0	
0.2	568.6	549
0.4	576.2	566
0.6	583.8	586
0.8	591.4	589
1.0	599.0	

Since the terrain is higher than the line of sight when $t = 0.6$, the tanks can't see each other.

82. (a) The planes $x + y + z = c$ have normal vector $\langle 1, 1, 1 \rangle$, so they are all parallel. Their x-, y-, and z-intercepts are all c. When $c > 0$ their intersection with the first octant is an equilateral triangle and when $c < 0$ their intersection with the octant diagonally opposite the first is an equilateral triangle.



(b) The planes $x + y + cz = 1$ have x-intercept 1, y-intercept 1, and z-intercept $1/c$. The plane with $c = 0$ is parallel to the z-axis. As c gets larger, the planes get closer to the xy-plane.

(c) The planes $y \cos\theta + z \cos\theta = 1$ have normal vectors $\langle 0, \cos\theta, \sin\theta \rangle$, which are perpendicular to the x-axis, and so the planes are parallel to the x-axis. We look at their intersection with the yz-plane. These are lines that are perpendicular to $\langle \cos\theta, \sin\theta \rangle$ and pass through $(\cos\theta, \sin\theta)$, since $\cos^2\theta + \sin^2\theta = 1$. So these are the tangent lines to the unit circle. Thus the family consists of all planes tangent to the circular cylinder with radius 1 and axis the x-axis.

83. If $a \neq 0$, then $ax + by + cz + d = 0 \quad \Rightarrow \quad a(x + d/a) + b(y - 0) + c(z - 0) = 0$ which by (7) is the scalar equation of the plane through the point $(-d/a, 0, 0)$ with normal vector $\langle a, b, c \rangle$. Similarly, if $b \neq 0$ (or if $c \neq 0$) the equation of the plane can be rewritten as $a(x - 0) + b(y + d/b) + c(z - 0) = 0$ [or as $a(x - 0) + b(y - 0) + c(z + d/c) = 0$] which by (7) is the scalar equation of a plane through the point $(0, -d/b, 0)$ [or the point $(0, 0, -d/c)$] with normal vector $\langle a, b, c \rangle$.

LABORATORY PROJECT Putting 3D in Perspective

1. If we view the screen from the camera's location, the vertical clipping plane on the left passes through the points $(1000, 0, 0)$, $(0, -400, 0)$, and $(0, -400, 600)$. A vector from the first point to the second is $\mathbf{v}_1 = \langle -1000, -400, 0 \rangle$ and a vector from the first point to the third is $\mathbf{v}_2 = \langle -1000, -400, 600 \rangle$. A normal vector for the clipping plane is $\mathbf{v}_1 \times \mathbf{v}_2 = -240{,}000\,\mathbf{i} + 600{,}000\,\mathbf{j}$ or $-2\,\mathbf{i} + 5\,\mathbf{j}$, and an equation for the plane is $-2(x - 1000) + 5(y - 0) + 0(z - 0) = 0 \quad \Rightarrow \quad 2x - 5y = 2000$. By symmetry, the vertical clipping plane on the right is given by $2x + 5y = 2000$. The lower clipping plane is $z = 0$. The upper clipping plane passes through the points $(1000, 0, 0)$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$(0, -400, 600)$, and $(0, 400, 600)$. Vectors from the first point to the second and third points are $\mathbf{v}_1 = \langle -1000, -400, 600\rangle$ and $\mathbf{v}_2 = \langle -1000, 400, 600\rangle$, and a normal vector for the plane is $\mathbf{v}_1 \times \mathbf{v}_2 = -480{,}000\,\mathbf{i} - 800{,}000\,\mathbf{k}$ or $3\,\mathbf{i} + 5\,\mathbf{k}$. An equation for the plane is $3(x - 1000) + 0(y - 0) + 5(z - 0) = 0 \;\Rightarrow\; 3x + 5z = 3000$.

A direction vector for the line L is $\mathbf{v} = \langle 630, 390, 162\rangle$ and taking $P_0 = (230, -285, 102)$, parametric equations are $x = 230 + 630t$, $y = -285 + 390t$, $z = 102 + 162t$. L intersects the left clipping plane when $2(230 + 630t) - 5(-285 + 390t) = 2000 \;\Rightarrow\; t = -\frac{1}{6}$. The corresponding point is $(125, -350, 75)$. L intersects the right clipping plane when $2(230 + 630t) + 5(-285 + 390t) = 2000 \;\Rightarrow\; t = \frac{593}{642}$. The corresponding point is approximately $(811.9, 75.2, 251.6)$, but this point is not contained within the viewing volume. L intersects the upper clipping plane when $3(230 + 630t) + 5(102 + 162t) = 3000 \;\Rightarrow\; t = \frac{2}{3}$, corresponding to the point $(650, -25, 210)$, and L intersects the lower clipping plane when $z = 0 \;\Rightarrow\; 102 + 162t = 0 \;\Rightarrow\; t = -\frac{17}{27}$. The corresponding point is approximately $(-166.7, -530.6, 0)$, which is not contained within the viewing volume. Thus L should be clipped at the points $(125, -350, 75)$ and $(650, -25, 210)$.

2. A sight line from the camera at $(1000, 0, 0)$ to the left endpoint $(125, -350, 75)$ of the clipped line has direction $\mathbf{v} = \langle -875, -350, 75\rangle$. Parametric equations are $x = 1000 - 875t$, $y = -350t$, $z = 75t$. This line intersects the screen when $x = 0 \;\Rightarrow\; 1000 - 875t = 0 \;\Rightarrow\; t = \frac{8}{7}$, corresponding to the point $\left(0, -400, \frac{600}{7}\right)$. Similarly, a sight line from the camera to the right endpoint $(650, -25, 210)$ of the clipped line has direction $\langle -350, -25, 210\rangle$ and parametric equations are $x = 1000 - 350t$, $y = -25t$, $z = 210t$. $x = 0 \;\Rightarrow\; 1000 - 350t = 0 \;\Rightarrow\; t = \frac{20}{7}$, corresponding to the point $\left(0, -\frac{500}{7}, 600\right)$. Thus the projection of the clipped line is the line segment between the points $\left(0, -400, \frac{600}{7}\right)$ and $\left(0, -\frac{500}{7}, 600\right)$.

3. From Equation 12.5.4, equations for the four sides of the screen are $\mathbf{r}_1(t) = (1 - t)\langle 0, -400, 0\rangle + t\,\langle 0, -400, 600\rangle$,
$\mathbf{r}_2(t) = (1 - t)\langle 0, -400, 600\rangle + t\,\langle 0, 400, 600\rangle$,
$\mathbf{r}_3(t) = (1 - t)\langle 0, 400, 0\rangle + t\,\langle 0, 400, 600\rangle$, and
$\mathbf{r}_4(t) = (1 - t)\langle 0, -400, 0\rangle + t\,\langle 0, 400, 0\rangle$. The clipped line segment connects the points $(125, -350, 75)$ and $(650, -25, 210)$, so an equation for the segment is $\mathbf{r}_5(t) = (1 - t)\langle 125, -350, 75\rangle + t\,\langle 650, -25, 210\rangle$.



The projection of the clipped segment connects the points $\left(0, -400, \frac{600}{7}\right)$ and $\left(0, -\frac{500}{7}, 600\right)$, so an equation is $\mathbf{r}_6(t) = (1 - t)\left\langle 0, -400, \frac{600}{7}\right\rangle + t\,\left\langle 0, -\frac{500}{7}, 600\right\rangle$.
The sight line on the left connects the points $(1000, 0, 0)$ and $\left(0, -400, \frac{600}{7}\right)$, so an equation is $\mathbf{r}_7(t) = (1 - t)\langle 1000, 0, 0\rangle + t\left\langle 0, -400, \frac{600}{7}\right\rangle$. The other sight line connects $(1000, 0, 0)$ to $\left(0, -\frac{500}{7}, 600\right)$, so an equation is $\mathbf{r}_8(t) = (1 - t)\langle 1000, 0, 0\rangle + t\left\langle 0, -\frac{500}{7}, 600\right\rangle$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. The vector from $(621, -147, 206)$ to $(563, 31, 242)$, $\mathbf{v}_1 = \langle -58, 178, 36\rangle$, lies in the plane of the rectangle, as does the vector from $(621, -147, 206)$ to $(657, -111, 86)$, $\mathbf{v}_2 = \langle 36, 36, -120\rangle$. A normal vector for the plane is $\mathbf{v}_1 \times \mathbf{v}_2 = \langle -1888, -142, -708\rangle$ or $\langle 8, 2, 3\rangle$, and an equation of the plane is $8x + 2y + 3z = 5292$. The line L intersects this plane when $8(230 + 630t) + 2(-285 + 390t) + 3(102 + 162t) = 5292 \Rightarrow t = \frac{1858}{3153} \approx 0.589$. The corresponding point is approximately $(601.25, -55.18, 197.46)$. Starting at this point, a portion of the line is hidden behind the rectangle. The line becomes visible again at the left edge of the rectangle, specifically the edge between the points $(621, -147, 206)$ and $(657, -111, 86)$. (This is most easily determined by graphing the rectangle and the line.) A plane through these two points and the camera's location, $(1000, 0, 0)$, will clip the line at the point it becomes visible. Two vectors in this plane are $\mathbf{v}_1 = \langle -379, -147, 206\rangle$ and $\mathbf{v}_2 = \langle -343, -111, 86\rangle$. A normal vector for the plane is $\mathbf{v}_1 \times \mathbf{v}_2 = \langle 10224, -38064, -8352\rangle$ and an equation of the plane is $213x - 793y - 174z = 213{,}000$. L intersects this plane when $213(230 + 630t) - 793(-285 + 390t) - 174(102 + 162t) = 213{,}000 \Rightarrow t = \frac{44{,}247}{203{,}268} \approx 0.2177$. The corresponding point is approximately $(367.14, -200.11, 137.26)$. Thus the portion of L that should be removed is the segment between the points $(601.25, -55.18, 197.46)$ and $(367.14, -200.11, 137.26)$.

12.6 Cylinders and Quadric Surfaces

1. (a) In $\mathbb{R}^2$, the equation $y = x^2$ represents a parabola.



(b) In $\mathbb{R}^3$, the equation $y = x^2$ doesn't involve z, so any horizontal plane with equation $z = k$ intersects the graph in a curve with equation $y = x^2$. Thus, the surface is a parabolic cylinder, made up of infinitely many shifted copies of the same parabola. The rulings are parallel to the z-axis.



(c) In $\mathbb{R}^3$, the equation $z = y^2$ also represents a parabolic cylinder. Since x doesn't appear, the graph is formed by moving the parabola $z = y^2$ in the direction of the x-axis. Thus, the rulings of the cylinder are parallel to the x-axis.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

2. (a)



(b) Since the equation $y = e^x$ doesn't involve z, horizontal traces are copies of the curve $y = e^x$. The rulings are parallel to the z-axis.



(c) The equation $z = e^y$ doesn't involve x, so vertical traces in $x = k$ (parallel to the yz-plane) are copies of the curve $z = e^y$. The rulings are parallel to the x-axis.



3. Since y is missing from the equation, the vertical traces $x^2 + z^2 = 1$, $y = k$, are copies of the same circle in the plane $y = k$. Thus the surface $x^2 + z^2 = 1$ is a circular cylinder with rulings parallel to the y-axis.



4. Since z is missing from the equation, the horizontal traces $4x^2 + y^2 = 4$, $z = k$, are copies of the same ellipse in the plane $z = k$. Thus the surface $4x^2 + y^2 = 4$ is an elliptic cylinder with rulings parallel to the z-axis.



5. Since x is missing, each vertical trace $z = 1 - y^2$, $x = k$, is a copy of the same parabola in the plane $x = k$. Thus the surface $z = 1 - y^2$ is a parabolic cylinder with rulings parallel to the x-axis.



6. Since x is missing, each vertical trace $y = z^2$, $x = k$, is a copy of the same parabola in the plane $x = k$. Thus the surface $y = z^2$ is a parabolic cylinder with rulings parallel to the x-axis.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. Since z is missing, each horizontal trace $xy = 1$, $z = k$, is a copy of the same hyperbola in the plane $z = k$. Thus the surface $xy = 1$ is a hyperbolic cylinder with rulings parallel to the z-axis.



8. Since x is missing, each vertical trace $z = \sin y$, $x = k$, is a copy of a sine curve in the plane $x = k$. Thus the surface $z = \sin y$ is a cylindrical surface with rulings parallel to the x-axis.



9. (a) The traces of $x^2 + y^2 - z^2 = 1$ in $x = k$ are $y^2 - z^2 = 1 - k^2$, a family of hyperbolas. (Note that the hyperbolas are oriented differently for $-1 < k < 1$ than for $k < -1$ or $k > 1$.) The traces in $y = k$ are $x^2 - z^2 = 1 - k^2$, a similar family of hyperbolas. The traces in $z = k$ are $x^2 + y^2 = 1 + k^2$, a family of circles. For $k = 0$, the trace in the xy-plane, the circle is of radius 1. As $|k|$ increases, so does the radius of the circle. This behavior, combined with the hyperbolic vertical traces, gives the graph of the hyperboloid of one sheet in Table 1.

(b) The shape of the surface is unchanged, but the hyperboloid is rotated so that its axis is the y-axis. Traces in $y = k$ are circles, while traces in $x = k$ and $z = k$ are hyperbolas.



(c) Completing the square in y gives $x^2 + (y + 1)^2 - z^2 = 1$. The surface is a hyperboloid identical to the one in part (a) but shifted one unit in the negative y-direction.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

10. (a) The traces of $-x^2 - y^2 + z^2 = 1$ in $x = k$ are $-y^2 + z^2 = 1 + k^2$, a family of hyperbolas, as are the traces in $y = k$, $-x^2 + z^2 = 1 + k^2$. The traces in $z = k$ are $x^2 + y^2 = k^2 - 1$, a family of circles for $|k| > 1$. As $|k|$ increases, the radii of the circles increase; the traces are empty for $|k| < 1$. This behavior, combined with the vertical traces, gives the graph of the hyperboloid of two sheets in Table 1.

(b) The graph has the same shape as the hyperboloid in part (a) but is rotated so that its axis is the x-axis. Traces in $x = k$, $|k| > 1$, are circles, while traces in $y = k$ and $z = k$ are hyperbolas.



11. For $x = y^2 + 4z^2$, the traces in $x = k$ are $y^2 + 4z^2 = k$. When $k > 0$ we have a family of ellipses. When $k = 0$ we have just a point at the origin, and the trace is empty for $k < 0$. The traces in $y = k$ are $x = 4z^2 + k^2$, a family of parabolas opening in the positive x-direction. Similarly, the traces in $z = k$ are $x = y^2 + 4k^2$, a family of parabolas opening in the positive x-direction. We recognize the graph as an elliptic paraboloid with axis the x-axis and vertex the origin.



12. $4x^2 + 9y^2 + 9z^2 = 36$. The traces in $x = k$ are $9y^2 + 9z^2 = 36 - 4k^2 \Leftrightarrow y^2 + z^2 = 4 - \frac{4}{9}k^2$, a family of circles for $|k| < 3$. (The traces are a single point for $|k| = 3$ and are empty for $|k| > 3$.) The traces in $y = k$ are $4x^2 + 9z^2 = 36 - 9k^2$, a family of ellipses for $|k| < 2$. Similarly, the traces in $z = k$ are the ellipses $4x^2 + 9y^2 = 36 - 9k^2$, $|k| < 2$. The graph is an ellipsoid centered at the origin with intercepts $x = \pm 3$, $y = \pm 2$, $z = \pm 2$.



13. $x^2 = 4y^2 + z^2$. The traces in $x = k$ are the ellipses $4y^2 + z^2 = k^2$. The traces in $y = k$ are $x^2 - z^2 = 4k^2$, hyperbolas for $k \neq 0$ and two intersecting lines if $k = 0$. Similarly, the traces in $z = k$ are $x^2 - 4y^2 = k^2$, hyperbolas for $k \neq 0$ and two intersecting lines if $k = 0$. We recognize the graph as an elliptic cone with axis the x-axis and vertex the origin.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $z^2 - 4x^2 - y^2 = 4$. The traces in $x = k$ are the hyperbolas $z^2 - y^2 = 4 + 4k^2$, and the traces in $y = k$ are the hyperbolas $z^2 - 4x^2 = 4 + k^2$. The traces in $z = k$ are $4x^2 + y^2 = k^2 - 4$, a family of ellipses for $|k| > 2$. (The traces are a single point for $|k| = 2$ and are empty for $|k| < 2$.) The surface is a hyperboloid of two sheets with axis the z-axis.



15. $9y^2 + 4z^2 = x^2 + 36$. The traces in $x = k$ are $9y^2 + 4z^2 = k^2 + 36$, a family of ellipses. The traces in $y = k$ are $4z^2 - x^2 = 9(4 - k^2)$, a family of hyperbolas for $|k| \neq 2$ and two intersecting lines when $|k| = 2$. (Note that the hyperbolas are oriented differently for $|k| < 2$ than for $|k| > 2$.) The traces in $z = k$ are $9y^2 - x^2 = 4(9 - k^2)$, a family of hyperbolas when $|k| \neq 3$ (oriented differently for $|k| < 3$ than for $|k| > 3$) and two intersecting lines when $|k| = 3$. We recognize the graph as a hyperboloid of one sheet with axis the x-axis.



16. $3x^2 + y + 3z^2 = 0$. The traces in $x = k$ are the parabolas $y = -3z^2 - 3k^2$ which open to the left (in the negative y-direction). Traces in $y = k$ are $3x^2 + 3z^2 = -k \quad \Leftrightarrow \quad x^2 + z^2 = -\dfrac{k}{3}$, a family of circles for $k < 0$. (Traces are empty for $k > 0$ and a single point for $k = 0$.) Traces in $z = k$ are the parabolas $y = -3x^2 - 3k^2$ which open in the negative y-direction. The graph is a circular paraboloid with axis the y-axis, opening in the negative y-direction, and vertex the origin.



17. $\dfrac{x^2}{9} + \dfrac{y^2}{25} + \dfrac{z^2}{4} = 1$. The traces in $x = k$ are $\dfrac{y^2}{25} + \dfrac{z^2}{4} = 1 - \dfrac{k^2}{9}$, a family of ellipses for $|k| < 3$. (The traces are a single point for $|k| = 3$ and are empty for $|k| > 3$.) The traces in $y = k$ are the ellipses $\dfrac{x^2}{9} + \dfrac{z^2}{4} = 1 - \dfrac{k^2}{25}$, $|k| < 5$, and the traces in $z = k$ are the ellipses $\dfrac{x^2}{9} + \dfrac{y^2}{25} = 1 - \dfrac{k^2}{4}$, $|k| < 2$. The surface is an ellipsoid centered at the origin with intercepts $x = \pm 3$, $y = \pm 5$, $z = \pm 2$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

18. $3x^2 - y^2 + 3z^2 = 0$. The traces in $x = k$ are $y^2 - 3z^2 = 3k^2$, a family of hyperbolas for $k \neq 0$ and two intersecting lines if $k = 0$. Traces in $y = k$ are the circles $3x^2 + 3z^2 = k^2 \quad \Leftrightarrow \quad x^2 + z^2 = \frac{1}{3}k^2$. The traces in $z = k$ are $y^2 - 3x^2 = 3k^2$, hyperbolas for $k \neq 0$ and two intersecting lines if $k = 0$. We recognize the surface as a circular cone with axis the y-axis and vertex the origin.



19. $y = z^2 - x^2$. The traces in $x = k$ are the parabolas $y = z^2 - k^2$, opening in the positive y-direction. The traces in $y = k$ are $k = z^2 - x^2$, two intersecting lines when $k = 0$ and a family of hyperbolas for $k \neq 0$ (note that the hyperbolas are oriented differently for $k > 0$ than for $k < 0$). The traces in $z = k$ are the parabolas $y = k^2 - x^2$ which open in the negative y-direction. Thus the surface is a hyperbolic paraboloid centered at $(0, 0, 0)$.



20. $x = y^2 - z^2$. The traces in $x = k$ are $y^2 - z^2 = k$, two intersecting lines when $k = 0$ and a family of hyperbolas for $k \neq 0$ (oriented differently for $k > 0$ than for $k < 0$). The traces in $y = k$ are the parabolas $x = -z^2 + k^2$, opening in the negative x-direction, and the traces in $z = k$ are the parabolas $x = y^2 - k^2$ which open in the positive x-direction. The graph is a hyperbolic paraboloid centered at $(0, 0, 0)$.



21. This is the equation of an ellipsoid: $x^2 + 4y^2 + 9z^2 = x^2 + \dfrac{y^2}{(1/2)^2} + \dfrac{z^2}{(1/3)^2} = 1$, with x-intercepts ± 1, y-intercepts $\pm\frac{1}{2}$ and z-intercepts $\pm\frac{1}{3}$. So the major axis is the x-axis and the only possible graph is VII.

22. This is the equation of an ellipsoid: $9x^2 + 4y^2 + z^2 = \dfrac{x^2}{(1/3)^2} + \dfrac{y^2}{(1/2)^2} + z^2 = 1$, with x-intercepts $\pm\frac{1}{3}$, y-intercepts $\pm\frac{1}{2}$ and z-intercepts ± 1. So the major axis is the z-axis and the only possible graph is IV.

23. This is the equation of a hyperboloid of one sheet, with $a = b = c = 1$. Since the coefficient of y^2 is negative, the axis of the hyperboloid is the y-axis, hence the correct graph is II.

24. This is a hyperboloid of two sheets, with $a = b = c = 1$. This surface does not intersect the xz-plane at all, so the axis of the hyperboloid is the y-axis and the graph is III.

25. There are no real values of x and z that satisfy this equation for $y < 0$, so this surface does not extend to the left of the xz-plane. The surface intersects the plane $y = k > 0$ in an ellipse. Notice that y occurs to the first power whereas x and z occur to the second power. So the surface is an elliptic paraboloid with axis the y-axis. Its graph is VI.

26. This is the equation of a cone with axis the y-axis, so the graph is I.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. This surface is a cylinder because the variable y is missing from the equation. The intersection of the surface and the xz-plane is an ellipse. So the graph is VIII.

28. This is the equation of a hyperbolic paraboloid. The trace in the xy-plane is the parabola $y = x^2$. So the correct graph is V.

29. Vertical traces parallel to the xz-plane are circles centered at the origin whose radii increase as y decreases. (The trace in $y = 1$ is just a single point and the graph suggests that traces in $y = k$ are empty for $k > 1$.) The traces in vertical planes parallel to the yz-plane are parabolas opening to the left that shift to the left as $|x|$ increases. One surface that fits this description is a circular paraboloid, opening to the left, with vertex $(0, 1, 0)$.



30. The vertical traces parallel to the yz-plane are ellipses that are smallest in the yz-plane and increase in size as $|x|$ increases. One surface that fits this description is a hyperboloid of one sheet with axis the x-axis. The horizontal traces in $z = k$ (hyperbolas and intersecting lines) also fit this surface, as shown in the figure.



31. $y^2 = x^2 + \frac{1}{9}z^2$ or $y^2 = x^2 + \dfrac{z^2}{9}$ represents an elliptic cone with vertex $(0, 0, 0)$ and axis the y-axis.



32. $4x^2 - y + 2z^2 = 0$ or $y = \dfrac{x^2}{1/4} + \dfrac{z^2}{1/2}$ or $\dfrac{y}{4} = x^2 + \dfrac{z^2}{2}$ represents an elliptic paraboloid with vertex $(0, 0, 0)$ and axis the y-axis.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

33. $x^2 + 2y - 2z^2 = 0$ or $2y = 2z^2 - x^2$ or $y = z^2 - \dfrac{x^2}{2}$ represents a hyperbolic paraboloid with center $(0, 0, 0)$.

34. $y^2 = x^2 + 4z^2 + 4$ or $-x^2 + y^2 - 4z^2 = 4$ or $-\dfrac{x^2}{4} + \dfrac{y^2}{4} - z^2 = 1$ represents a hyperboloid of two sheets with axis the y-axis.



35. Completing squares in x and y gives

$(x^2 - 2x + 1) + (y^2 - 6y + 9) - z = 0 \quad \Leftrightarrow$

$(x - 1)^2 + (y - 3)^2 - z = 0$ or $z = (x - 1)^2 + (y - 3)^2$, a circular paraboloid opening upward with vertex $(1, 3, 0)$ and axis the vertical line $x = 1, y = 3$.



36. Completing squares in x and z gives $(x^2 - 4x + 4) - y^2 - (z^2 + 2z + 1) + 3 = 0 + 4 - 1 \quad \Leftrightarrow$

$(x - 2)^2 - y^2 - (z + 1)^2 = 0$ or $(x - 2)^2 = y^2 + (z + 1)^2$, a circular cone with vertex $(2, 0, -1)$ and axis the horizontal line $y = 0, z = -1$.



37. Completing squares in x and z gives

$(x^2 - 4x + 4) - y^2 + (z^2 - 2z + 1) = 0 + 4 + 1 \quad \Leftrightarrow$

$(x - 2)^2 - y^2 + (z - 1)^2 = 5$ or $\dfrac{(x - 2)^2}{5} - \dfrac{y^2}{5} + \dfrac{(z - 1)^2}{5}$ 1, a hyperboloid of one sheet with center $(2, 0, 1)$ and axis the horizontal line $x = 2, z = 1$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. Completing squares in all three variables gives

$4(x^2 - 6x + 9) + (y^2 - 8y + 16) + (z^2 + 4z + 4) = -55 + 36 + 16 + 4 \quad \Leftrightarrow$

$4(x-3)^2 + (y-4)^2 + (z+2)^2 = 1$ or

$\dfrac{(x-3)^2}{1/4} + (y-4)^2 + (z+2)^2 = 1$, an ellipsoid with center $(3, 4, -2)$.



39. Solving the equation for z we get $z = \pm\sqrt{1 + 4x^2 + y^2}$, so we plot separately $z = \sqrt{1 + 4x^2 + y^2}$ and $z = -\sqrt{1 + 4x^2 + y^2}$.





To restrict the z-range as in the second graph, we can use the option `view=-4..4` in Maple's `plot3d` command, or `PlotRange->{-4,4}` in Mathematica's `Plot3D` command.

40. We plot the surface $z = x^2 - y^2$.



41. Solving the equation for z we get $z = \pm\sqrt{4x^2 + y^2}$, so we plot separately $z = \sqrt{4x^2 + y^2}$ and $z = -\sqrt{4x^2 + y^2}$.





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

42. We plot the surface $z = x^2 - 6x + 4y^2$.





43.



44.



45. The curve $y = \sqrt{x}$ is equivalent to $x = y^2$, $y \geq 0$. Rotating the curve about the x-axis creates a circular paraboloid with vertex at the origin, axis the x-axis, opening in the positive x-direction. The trace in the xy-plane is $x = y^2$, $z = 0$, and the trace in the xz-plane is a parabola of the same shape: $x = z^2$, $y = 0$. An equation for the surface is $x = y^2 + z^2$.



46. Rotating the line $z = 2y$ about the z-axis creates a (right) circular cone with vertex at the origin and axis the z-axis. Traces in $z = k$ ($k \neq 0$) are circles with center $(0, 0, k)$ and radius $y = z/2 = k/2$, so an equation for the trace is $x^2 + y^2 = (k/2)^2$, $z = k$. Thus an equation for the surface is $x^2 + y^2 = (z/2)^2$ or $4x^2 + 4y^2 = z^2$.



47. Let $P = (x, y, z)$ be an arbitrary point equidistant from $(-1, 0, 0)$ and the plane $x = 1$. Then the distance from P to $(-1, 0, 0)$ is $\sqrt{(x+1)^2 + y^2 + z^2}$ and the distance from P to the plane $x = 1$ is $|x - 1|\,/\sqrt{1^2} = |x - 1|$ (by Equation 12.5.9). So $|x - 1| = \sqrt{(x+1)^2 + y^2 + z^2} \quad \Leftrightarrow \quad (x-1)^2 = (x+1)^2 + y^2 + z^2 \quad \Leftrightarrow$ $x^2 - 2x + 1 = x^2 + 2x + 1 + y^2 + z^2 \quad \Leftrightarrow \quad -4x = y^2 + z^2$. Thus the collection of all such points P is a circular paraboloid with vertex at the origin, axis the x-axis, which opens in the negative x-direction.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. Let $P = (x, y, z)$ be an arbitrary point whose distance from the x-axis is twice its distance from the yz-plane. The distance from P to the x-axis is $\sqrt{(x-x)^2 + y^2 + z^2} = \sqrt{y^2 + z^2}$ and the distance from P to the yz-plane ($x = 0$) is $|x|\,/1 = |x|$. Thus $\sqrt{y^2 + z^2} = 2\,|x| \quad \Leftrightarrow \quad y^2 + z^2 = 4x^2 \quad \Leftrightarrow \quad x^2 = (y^2/2^2) + (z^2/2^2)$. So the surface is a right circular cone with vertex the origin and axis the x-axis.

49. (a) An equation for an ellipsoid centered at the origin with intercepts $x = \pm a$, $y = \pm b$, and $z = \pm c$ is $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} + \dfrac{z^2}{c^2} = 1$.

Here the poles of the model intersect the z-axis at $z = \pm 6356.523$ and the equator intersects the x- and y-axes at $x = \pm 6378.137$, $y = \pm 6378.137$, so an equation is

$$\frac{x^2}{(6378.137)^2} + \frac{y^2}{(6378.137)^2} + \frac{z^2}{(6356.523)^2} = 1$$

(b) Traces in $z = k$ are the circles $\dfrac{x^2}{(6378.137)^2} + \dfrac{y^2}{(6378.137)^2} = 1 - \dfrac{k^2}{(6356.523)^2} \quad \Leftrightarrow$

$$x^2 + y^2 = (6378.137)^2 - \left(\frac{6378.137}{6356.523}\right)^2 k^2.$$

(c) To identify the traces in $y = mx$ we substitute $y = mx$ into the equation of the ellipsoid:

$$\frac{x^2}{(6378.137)^2} + \frac{(mx)^2}{(6378.137)^2} + \frac{z^2}{(6356.523)^2} = 1$$

$$\frac{(1+m^2)x^2}{(6378.137)^2} + \frac{z^2}{(6356.523)^2} = 1$$

$$\frac{x^2}{(6378.137)^2/(1+m^2)} + \frac{z^2}{(6356.523)^2} = 1$$

As expected, this is a family of ellipses.

50. If we position the hyperboloid on coordinate axes so that it is centered at the origin with axis the z-axis then its equation is given by $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} - \dfrac{z^2}{c^2} = 1$. Horizontal traces in $z = k$ are $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 + \dfrac{k^2}{c^2}$, a family of ellipses, but we know that the traces are circles so we must have $a = b$. The trace in $z = 0$ is $\dfrac{x^2}{a^2} + \dfrac{y^2}{a^2} = 1 \quad \Leftrightarrow \quad x^2 + y^2 = a^2$ and since the minimum radius of 100 m occurs there, we must have $a = 100$. The base of the tower is the trace in $z = -500$ given by $\dfrac{x^2}{a^2} + \dfrac{y^2}{a^2} = 1 + \dfrac{(-500)^2}{c^2}$ but $a = 100$ so the trace is $x^2 + y^2 = 100^2 + 50{,}000^2\dfrac{1}{c^2}$. We know the base is a circle of radius 140, so we must have $100^2 + 50{,}000^2\dfrac{1}{c^2} = 140^2 \quad \Rightarrow \quad c^2 = \dfrac{50{,}000^2}{140^2 - 100^2} = \dfrac{781{,}250}{3}$ and an equation for the tower is $\dfrac{x^2}{100^2} + \dfrac{y^2}{100^2} - \dfrac{z^2}{(781{,}250)/3} = 1$ or $\dfrac{x^2}{10{,}000} + \dfrac{y^2}{10{,}000} - \dfrac{3z^2}{781{,}250} = 1, \ -500 \le z \le 500.$

51. If (a, b, c) satisfies $z = y^2 - x^2$, then $c = b^2 - a^2$. L_1: $x = a + t$, $y = b + t$, $z = c + 2(b - a)t$, L_2: $x = a + t$, $y = b - t$, $z = c - 2(b + a)t$. Substitute the parametric equations of L_1 into the equation of the hyperbolic paraboloid in order to find the points of intersection: $z = y^2 - x^2 \quad \Rightarrow$

$c + 2(b - a)t = (b + t)^2 - (a + t)^2 = b^2 - a^2 + 2(b - a)t \quad \Rightarrow \quad c = b^2 - a^2$. As this is true for all values of t,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

L_1 lies on $z = y^2 - x^2$. Performing similar operations with L_2 gives: $z = y^2 - x^2 \Rightarrow$ $c - 2(b+a)t = (b-t)^2 - (a+t)^2 = b^2 - a^2 - 2(b+a)t \Rightarrow c = b^2 - a^2$. This tells us that all of L_2 also lies on $z = y^2 - x^2$.

52. Any point on the curve of intersection must satisfy both $2x^2 + 4y^2 - 2z^2 + 6x = 2$ and $2x^2 + 4y^2 - 2z^2 - 5y = 0$. Subtracting, we get $6x + 5y = 2$, which is linear and therefore the equation of a plane. Thus the curve of intersection lies in this plane.

53.



The curve of intersection looks like a bent ellipse. The projection of this curve onto the xy-plane is the set of points $(x, y, 0)$ which satisfy $x^2 + y^2 = 1 - y^2 \Leftrightarrow x^2 + 2y^2 = 1 \Leftrightarrow$ $x^2 + \dfrac{y^2}{\left(1/\sqrt{2}\right)^2} = 1$. This is an equation of an ellipse.

12 Review

TRUE-FALSE QUIZ

1. This is false, as the dot product of two vectors is a scalar, not a vector.
2. False. For example, if $\mathbf{u} = \mathbf{i}$ and $\mathbf{v} = -\mathbf{i}$ then $|\mathbf{u} + \mathbf{v}| = |\mathbf{0}| = 0$ but $|\mathbf{u}| + |\mathbf{v}| = 1 + 1 = 2$.
3. False. For example, if $\mathbf{u} = \mathbf{i}$ and $\mathbf{v} = \mathbf{j}$ then $|\mathbf{u} \cdot \mathbf{v}| = |0| = 0$ but $|\mathbf{u}|\,|\mathbf{v}| = 1 \cdot 1 = 1$. In fact, by Theorem 12.3.3, $|\mathbf{u} \cdot \mathbf{v}| = \big||\mathbf{u}|\,|\mathbf{v}| \cos\theta\big|$.
4. False. For example, $|\mathbf{i} \times \mathbf{i}| = |\mathbf{0}| = 0$ (see Example 12.4.2) but $|\mathbf{i}|\,|\mathbf{i}| = 1 \cdot 1 = 1$. In fact, by Theorem 12.4.9, $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| \sin\theta$.
5. True, by Theorem 12.3.2, property 2.
6. False. Property 1 of Theorem 12.4.11 says that $\mathbf{u} \times \mathbf{v} = -\mathbf{v} \times \mathbf{u}$.
7. True. If θ is the angle between $\mathbf{u}$ and $\mathbf{v}$, then by Theorem 12.4.9, $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| \sin\theta = |\mathbf{v}|\,|\mathbf{u}| \sin\theta = |\mathbf{v} \times \mathbf{u}|$. (Or, by Theorem 12.4.11, $|\mathbf{u} \times \mathbf{v}| = |-\mathbf{v} \times \mathbf{u}| = |-1|\,|\mathbf{v} \times \mathbf{u}| = |\mathbf{v} \times \mathbf{u}|$.)
8. This is true by Theorem 12.3.2, property 4.
9. Theorem 12.4.11, property 2 tells us that this is true.
10. This is true by Theorem 12.4.11, property 4.
11. This is true by Theorem 12.4.11, property 5.
12. In general, this assertion is false; a counterexample is $\mathbf{i} \times (\mathbf{i} \times \mathbf{j}) \neq (\mathbf{i} \times \mathbf{i}) \times \mathbf{j}$. (See the paragraph preceding Theorem 12.4.11.)
13. This is true because $\mathbf{u} \times \mathbf{v}$ is orthogonal to $\mathbf{u}$ (see Theorem 12.4.8), and the dot product of two orthogonal vectors is 0.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $(\mathbf{u}+\mathbf{v})\times\mathbf{v}=\mathbf{u}\times\mathbf{v}+\mathbf{v}\times\mathbf{v}$ [by Theorem 12.4.11, property 4]

$=\mathbf{u}\times\mathbf{v}+\mathbf{0}$ [by Example 12.4.2]

$=\mathbf{u}\times\mathbf{v}$, so this is true.

15. This is false. A normal vector to the plane is $\mathbf{n}=\langle 6,-2,4\rangle$. Because $\langle 3,-1,2\rangle=\frac{1}{2}\mathbf{n}$, the vector is parallel to $\mathbf{n}$ and hence perpendicular to the plane.

16. This is false, because according to Equation 12.5.8, $ax+by+cz+d=0$ is the general equation of a plane.

17. This is false. In $\mathbb{R}^2$, $x^2+y^2=1$ represents a circle, but $\{(x,y,z)\mid x^2+y^2=1\}$ represents a *three-dimensional surface*, namely, a circular cylinder with axis the z-axis.

18. This is false. In $\mathbb{R}^3$ the graph of $y=x^2$ is a parabolic cylinder (see Example 12.6.1). A paraboloid has an equation such as $z=x^2+y^2$.

19. False. For example, $\mathbf{i}\cdot\mathbf{j}=\mathbf{0}$ but $\mathbf{i}\neq\mathbf{0}$ and $\mathbf{j}\neq\mathbf{0}$.

20. This is false. By Corollary 12.4.10, $\mathbf{u}\times\mathbf{v}=\mathbf{0}$ for any nonzero parallel vectors $\mathbf{u}$, $\mathbf{v}$. For instance, $\mathbf{i}\times\mathbf{i}=\mathbf{0}$.

21. This is true. If $\mathbf{u}$ and $\mathbf{v}$ are both nonzero, then by (7) in Section 12.3, $\mathbf{u}\cdot\mathbf{v}=0$ implies that $\mathbf{u}$ and $\mathbf{v}$ are orthogonal. But $\mathbf{u}\times\mathbf{v}=\mathbf{0}$ implies that $\mathbf{u}$ and $\mathbf{v}$ are parallel (see Corollary 12.4.10). Two nonzero vectors can't be both parallel and orthogonal, so at least one of $\mathbf{u}$, $\mathbf{v}$ must be $\mathbf{0}$.

22. This is true. We know $\mathbf{u}\cdot\mathbf{v}=|\mathbf{u}|\,|\mathbf{v}|\cos\theta$ where $|\mathbf{u}|\geq 0$, $|\mathbf{v}|\geq 0$, and $|\cos\theta|\leq 1$, so $|\mathbf{u}\cdot\mathbf{v}|=|\mathbf{u}|\,|\mathbf{v}|\,|\cos\theta|\leq|\mathbf{u}|\,|\mathbf{v}|$.

EXERCISES

1. (a) The radius of the sphere is the distance between the points $(-1,2,1)$ and $(6,-2,3)$, namely, $\sqrt{[6-(-1)]^2+(-2-2)^2+(3-1)^2}=\sqrt{69}$. By the formula for an equation of a sphere (see page 835 [ET 795]), an equation of the sphere with center $(-1,2,1)$ and radius $\sqrt{69}$ is $(x+1)^2+(y-2)^2+(z-1)^2=69$.

(b) The intersection of this sphere with the yz-plane is the set of points on the sphere whose x-coordinate is 0. Putting $x=0$ into the equation, we have $(y-2)^2+(z-1)^2=68, x=0$ which represents a circle in the yz-plane with center $(0,2,1)$ and radius $\sqrt{68}$.

(c) Completing squares gives $(x-4)^2+(y+1)^2+(z+3)^2=-1+16+1+9=25$. Thus the sphere is centered at $(4,-1,-3)$ and has radius 5.

2. (a)



(b)



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c)



(d)



3. $\mathbf{u}\cdot\mathbf{v} = |\mathbf{u}|\,|\mathbf{v}|\cos 45° = (2)(3)\frac{\sqrt{2}}{2} = 3\sqrt{2}$. $|\mathbf{u}\times\mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}|\sin 45° = (2)(3)\frac{\sqrt{2}}{2} = 3\sqrt{2}$.
By the right-hand rule, $\mathbf{u}\times\mathbf{v}$ is directed out of the page.

4. (a) $2\mathbf{a}+3\mathbf{b} = 2\mathbf{i}+2\mathbf{j}-4\mathbf{k}+9\mathbf{i}-6\mathbf{j}+3\mathbf{k} = 11\mathbf{i}-4\mathbf{j}-\mathbf{k}$

(b) $|\mathbf{b}| = \sqrt{9+4+1} = \sqrt{14}$

(c) $\mathbf{a}\cdot\mathbf{b} = (1)(3)+(1)(-2)+(-2)(1) = -1$

(d) $\mathbf{a}\times\mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & -2 \\ 3 & -2 & 1 \end{vmatrix} = (1-4)\,\mathbf{i}-(1+6)\,\mathbf{j}+(-2-3)\,\mathbf{k} = -3\mathbf{i}-7\mathbf{j}-5\mathbf{k}$

(e) $\mathbf{b}\times\mathbf{c} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -2 & 1 \\ 0 & 1 & -5 \end{vmatrix} = 9\mathbf{i}+15\mathbf{j}+3\mathbf{k}, \quad |\mathbf{b}\times\mathbf{c}| = 3\sqrt{9+25+1} = 3\sqrt{35}$

(f) $\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c}) = \begin{vmatrix} 1 & 1 & -2 \\ 3 & -2 & 1 \\ 0 & 1 & -5 \end{vmatrix} = \begin{vmatrix} -2 & 1 \\ 1 & -5 \end{vmatrix} - \begin{vmatrix} 3 & 1 \\ 0 & -5 \end{vmatrix} - 2\begin{vmatrix} 3 & -2 \\ 0 & 1 \end{vmatrix} = 9+15-6 = 18$

(g) $\mathbf{c}\times\mathbf{c} = \mathbf{0}$ for any $\mathbf{c}$.

(h) From part (e),

$$\mathbf{a}\times(\mathbf{b}\times\mathbf{c}) = \mathbf{a}\times(9\mathbf{i}+15\mathbf{j}+3\mathbf{k}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & -2 \\ 9 & 15 & 3 \end{vmatrix}$$

$$= (3+30)\,\mathbf{i}-(3+18)\,\mathbf{j}+(15-9)\,\mathbf{k} = 33\mathbf{i}-21\mathbf{j}+6\mathbf{k}$$

(i) The scalar projection is $\text{comp}_{\mathbf{a}}\,\mathbf{b} = |\mathbf{b}|\cos\theta = \mathbf{a}\cdot\mathbf{b}/|\mathbf{a}| = -\frac{1}{\sqrt{6}}$.

(j) The vector projection is $\text{proj}_{\mathbf{a}}\,\mathbf{b} = -\dfrac{1}{\sqrt{6}}\left(\dfrac{\mathbf{a}}{|\mathbf{a}|}\right) = -\frac{1}{6}(\mathbf{i}+\mathbf{j}-2\mathbf{k})$.

(k) $\cos\theta = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{-1}{\sqrt{6}\sqrt{14}} = \dfrac{-1}{2\sqrt{21}}$ and $\theta = \cos^{-1}\left(\dfrac{-1}{2\sqrt{21}}\right) \approx 96°$.

5. For the two vectors to be orthogonal, we need $\langle 3,2,x\rangle\cdot\langle 2x,4,x\rangle = 0 \quad\Leftrightarrow\quad (3)(2x)+(2)(4)+(x)(x) = 0 \quad\Leftrightarrow$
$x^2+6x+8 = 0 \quad\Leftrightarrow\quad (x+2)(x+4) = 0 \quad\Leftrightarrow\quad x = -2$ or $x = -4$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. We know that the cross product of two vectors is orthogonal to both given vectors. So we calculate

$(\mathbf{j}+2\mathbf{k})\times(\mathbf{i}-2\mathbf{j}+3\mathbf{k}) = [3-(-4)]\,\mathbf{i}-(0-2)\,\mathbf{j}+(0-1)\,\mathbf{k} = 7\mathbf{i}+2\mathbf{j}-\mathbf{k}.$

Then two unit vectors orthogonal to both given vectors are $\pm\dfrac{7\mathbf{i}+2\mathbf{j}-\mathbf{k}}{\sqrt{7^2+2^2+(-1)^2}} = \pm\dfrac{1}{3\sqrt{6}}(7\mathbf{i}+2\mathbf{j}-\mathbf{k})$,

that is, $\frac{7}{3\sqrt{6}}\mathbf{i}+\frac{2}{3\sqrt{6}}\mathbf{j}-\frac{1}{3\sqrt{6}}\mathbf{k}$ and $-\frac{7}{3\sqrt{6}}\mathbf{i}-\frac{2}{3\sqrt{6}}\mathbf{j}+\frac{1}{3\sqrt{6}}\mathbf{k}$.

7. (a) $(\mathbf{u}\times\mathbf{v})\cdot\mathbf{w} = \mathbf{u}\cdot(\mathbf{v}\times\mathbf{w}) = 2$

(b) $\mathbf{u}\cdot(\mathbf{w}\times\mathbf{v}) = \mathbf{u}\cdot[-(\mathbf{v}\times\mathbf{w})] = -\mathbf{u}\cdot(\mathbf{v}\times\mathbf{w}) = -2$

(c) $\mathbf{v}\cdot(\mathbf{u}\times\mathbf{w}) = (\mathbf{v}\times\mathbf{u})\cdot\mathbf{w} = -(\mathbf{u}\times\mathbf{v})\cdot\mathbf{w} = -2$

(d) $(\mathbf{u}\times\mathbf{v})\cdot\mathbf{v} = \mathbf{u}\cdot(\mathbf{v}\times\mathbf{v}) = \mathbf{u}\cdot\mathbf{0} = 0$

8. $(\mathbf{a}\times\mathbf{b})\cdot[(\mathbf{b}\times\mathbf{c})\times(\mathbf{c}\times\mathbf{a})] = (\mathbf{a}\times\mathbf{b})\cdot([(\mathbf{b}\times\mathbf{c})\cdot\mathbf{a}]\,\mathbf{c}-[(\mathbf{b}\times\mathbf{c})\cdot\mathbf{c}]\,\mathbf{a})$

[by Property 6 of the cross product]

$= (\mathbf{a}\times\mathbf{b})\cdot[(\mathbf{b}\times\mathbf{c})\cdot\mathbf{a}]\,\mathbf{c} = [\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})]\,(\mathbf{a}\times\mathbf{b})\cdot\mathbf{c}$

$= [\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})]\,[\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})] = [\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})]^2$

9. For simplicity, consider a unit cube positioned with its back left corner at the origin. Vector representations of the diagonals joining the points $(0,0,0)$ to $(1,1,1)$ and $(1,0,0)$ to $(0,1,1)$ are $\langle 1,1,1\rangle$ and $\langle -1,1,1\rangle$. Let θ be the angle between these two vectors. $\langle 1,1,1\rangle\cdot\langle -1,1,1\rangle = -1+1+1 = 1 = |\langle 1,1,1\rangle|\,|\langle -1,1,1\rangle|\cos\theta = 3\cos\theta \;\Rightarrow\; \cos\theta = \frac{1}{3} \;\Rightarrow$ $\theta = \cos^{-1}\left(\frac{1}{3}\right) \approx 71°$.

10. $\overrightarrow{AB} = \langle 1,3,-1\rangle$, $\overrightarrow{AC} = \langle -2,1,3\rangle$ and $\overrightarrow{AD} = \langle -1,3,1\rangle$. By Equation 12.4.13,

$$\overrightarrow{AB}\cdot\left(\overrightarrow{AC}\times\overrightarrow{AD}\right) = \begin{vmatrix} 1 & 3 & -1 \\ -2 & 1 & 3 \\ -1 & 3 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 3 \\ 3 & 1 \end{vmatrix} - 3\begin{vmatrix} -2 & 3 \\ -1 & 1 \end{vmatrix} - \begin{vmatrix} -2 & 1 \\ -1 & 3 \end{vmatrix} = -8-3+5 = -6.$$

The volume is $\left|\overrightarrow{AB}\cdot\left(\overrightarrow{AC}\times\overrightarrow{AD}\right)\right| = 6$ cubic units.

11. $\overrightarrow{AB} = \langle 1,0,-1\rangle$, $\overrightarrow{AC} = \langle 0,4,3\rangle$, so

(a) a vector perpendicular to the plane is $\overrightarrow{AB}\times\overrightarrow{AC} = \langle 0+4,-(3+0),4-0\rangle = \langle 4,-3,4\rangle$.

(b) $\frac{1}{2}\left|\overrightarrow{AB}\times\overrightarrow{AC}\right| = \frac{1}{2}\sqrt{16+9+16} = \frac{\sqrt{41}}{2}$.

12. $\mathbf{D} = 4\mathbf{i}+3\mathbf{j}+6\mathbf{k}$, $W = \mathbf{F}\cdot\mathbf{D} = 12+15+60 = 87$ J

13. Let F_1 be the magnitude of the force directed $20°$ away from the direction of shore, and let F_2 be the magnitude of the other force. Separating these forces into components parallel to the direction of the resultant force and perpendicular to it gives

$F_1\cos 20° + F_2\cos 30° = 255$ **(1)**, and $F_1\sin 20° - F_2\sin 30° = 0 \;\Rightarrow\; F_1 = F_2\dfrac{\sin 30°}{\sin 20°}$ **(2)**. Substituting **(2)** into **(1)** gives $F_2(\sin 30°\cot 20° + \cos 30°) = 255 \;\Rightarrow\; F_2 \approx 114$ N. Substituting this into **(2)** gives $F_1 \approx 166$ N.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

14. $|\boldsymbol{\tau}| = |\mathbf{r}|\,|\mathbf{F}| \sin\theta = (0.40)(50)\sin(90^\circ - 30^\circ) \approx 17.3$ N·m.

15. The line has direction $\mathbf{v} = \langle -3, 2, 3\rangle$. Letting $P_0 = (4, -1, 2)$, parametric equations are $x = 4 - 3t,\ y = -1 + 2t,\ z = 2 + 3t$.

16. A direction vector for the line is $\mathbf{v} = \langle 3, 2, 1\rangle$, so parametric equations for the line are $x = 1 + 3t,\ y = 2t,\ z = -1 + t$.

17. A direction vector for the line is a normal vector for the plane, $\mathbf{n} = \langle 2, -1, 5\rangle$, and parametric equations for the line are $x = -2 + 2t,\ y = 2 - t,\ z = 4 + 5t$.

18. Since the two planes are parallel, they will have the same normal vectors. Then we can take $\mathbf{n} = \langle 1, 4, -3\rangle$ and an equation of the plane is $1(x - 2) + 4(y - 1) - 3(z - 0) = 0$ or $x + 4y - 3z = 6$.

19. Here the vectors $\mathbf{a} = \langle 4 - 3, 0 - (-1), 2 - 1\rangle = \langle 1, 1, 1\rangle$ and $\mathbf{b} = \langle 6 - 3, 3 - (-1), 1 - 1\rangle = \langle 3, 4, 0\rangle$ lie in the plane, so $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle -4, 3, 1\rangle$ is a normal vector to the plane and an equation of the plane is $-4(x - 3) + 3(y - (-1)) + 1(z - 1) = 0$ or $-4x + 3y + z = -14$.

20. If we first find two nonparallel vectors in the plane, their cross product will be a normal vector to the plane. Since the given line lies in the plane, its direction vector $\mathbf{a} = \langle 2, -1, 3\rangle$ is one vector in the plane. We can verify that the given point $(1, 2, -2)$ does not lie on this line. The point $(0, 3, 1)$ is on the line (obtained by putting $t = 0$) and hence in the plane, so the vector $\mathbf{b} = \langle 0 - 1, 3 - 2, 1 - (-2)\rangle = \langle -1, 1, 3\rangle$ lies in the plane, and a normal vector is $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle -6, -9, 1\rangle$. Thus an equation of the plane is $-6(x - 1) - 9(y - 2) + (z + 2) = 0$ or $6x + 9y - z = 26$.

21. Substitution of the parametric equations into the equation of the plane gives $2x - y + z = 2(2 - t) - (1 + 3t) + 4t = 2 \quad \Rightarrow \quad -t + 3 = 2 \quad \Rightarrow \quad t = 1$. When $t = 1$, the parametric equations give $x = 2 - 1 = 1$, $y = 1 + 3 = 4$ and $z = 4$. Therefore, the point of intersection is $(1, 4, 4)$.

22. Use the formula proven in Exercise 12.4.45(a). In the notation used in that exercise, $\mathbf{a}$ is just the direction of the line; that is, $\mathbf{a} = \langle 1, -1, 2\rangle$. A point on the line is $(1, 2, -1)$ (setting $t = 0$), and therefore $\mathbf{b} = \langle 1 - 0, 2 - 0, -1 - 0\rangle = \langle 1, 2, -1\rangle$.

Hence $d = \dfrac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \dfrac{|\langle 1, -1, 2\rangle \times \langle 1, 2, -1\rangle|}{\sqrt{1 + 1 + 4}} = \dfrac{|\langle -3, 3, 3\rangle|}{\sqrt{6}} = \sqrt{\dfrac{27}{6}} = \dfrac{3}{\sqrt{2}}$.

23. Since the direction vectors $\langle 2, 3, 4\rangle$ and $\langle 6, -1, 2\rangle$ aren't parallel, neither are the lines. For the lines to intersect, the three equations $1 + 2t = -1 + 6s$, $2 + 3t = 3 - s$, $3 + 4t = -5 + 2s$ must be satisfied simultaneously. Solving the first two equations gives $t = \frac{1}{5}$, $s = \frac{2}{5}$ and checking we see these values don't satisfy the third equation. Thus the lines aren't parallel and they don't intersect, so they must be skew.

24. (a) The normal vectors are $\langle 1, 1, -1\rangle$ and $\langle 2, -3, 4\rangle$. Since these vectors aren't parallel, neither are the planes parallel.

Also $\langle 1, 1, -1\rangle \cdot \langle 2, -3, 4\rangle = 2 - 3 - 4 = -5 \neq 0$ so the normal vectors, and thus the planes, are not perpendicular.

(b) $\cos\theta = \dfrac{\langle 1, 1, -1\rangle \cdot \langle 2, -3, 4\rangle}{\sqrt{3}\sqrt{29}} = -\dfrac{5}{\sqrt{87}}$ and $\theta = \cos^{-1}\left(-\frac{5}{\sqrt{87}}\right) \approx 122^\circ$ [or we can say $\approx 58^\circ$].

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $\mathbf{n}_1 = \langle 1, 0, -1\rangle$ and $\mathbf{n}_2 = \langle 0, 1, 2\rangle$. Setting $z = 0$, it is easy to see that $(1, 3, 0)$ is a point on the line of intersection of $x - z = 1$ and $y + 2z = 3$. The direction of this line is $\mathbf{v}_1 = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, -2, 1\rangle$. A second vector parallel to the desired plane is $\mathbf{v}_2 = \langle 1, 1, -2\rangle$, since it is perpendicular to $x + y - 2z = 1$. Therefore, the normal of the plane in question is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 4 - 1, 1 + 2, 1 + 2\rangle = 3\langle 1, 1, 1\rangle$. Taking $(x_0, y_0, z_0) = (1, 3, 0)$, the equation we are looking for is $(x - 1) + (y - 3) + z = 0 \quad \Leftrightarrow \quad x + y + z = 4$.

26. (a) The vectors $\overrightarrow{AB} = \langle -1 - 2, -1 - 1, 10 - 1\rangle = \langle -3, -2, 9\rangle$ and $\overrightarrow{AC} = \langle 1 - 2, 3 - 1, -4 - 1\rangle = \langle -1, 2, -5\rangle$ lie in the plane, so $\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \langle -3, -2, 9\rangle \times \langle -1, 2, -5\rangle = \langle -8, -24, -8\rangle$ or equivalently $\langle 1, 3, 1\rangle$ is a normal vector to the plane. The point $A(2, 1, 1)$ lies on the plane so an equation of the plane is $1(x - 2) + 3(y - 1) + 1(z - 1) = 0$ or $x + 3y + z = 6$.

(b) The line is perpendicular to the plane so it is parallel to a normal vector for the plane, namely $\langle 1, 3, 1\rangle$. If the line passes through $B(-1, -1, 10)$ then symmetric equations are $\dfrac{x - (-1)}{1} = \dfrac{y - (-1)}{3} = \dfrac{z - 10}{1}$ or $x + 1 = \dfrac{y + 1}{3} = z - 10$.

(c) Normal vectors for the two planes are $\mathbf{n}_1 = \langle 1, 3, 1\rangle$ and $\mathbf{n}_2 = \langle 2, -4, -3\rangle$. The angle θ between the planes is given by

$$\cos\theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\ |\mathbf{n}_2|} = \frac{\langle 1, 3, 1\rangle \cdot \langle 2, -4, -3\rangle}{\sqrt{1^2 + 3^2 + 1^2}\sqrt{2^2 + (-4)^2 + (-3)^2}} = \frac{2 - 12 - 3}{\sqrt{11}\sqrt{29}} = -\frac{13}{\sqrt{319}}$$

Thus $\theta = \cos^{-1}\left(-\dfrac{13}{\sqrt{319}}\right) \approx 137°$ or $180° - 137° = 43°$.

(d) From part (c), the point $(2, 0, 4)$ lies on the second plane, but notice that the point also satisfies the equation of the first plane, so the point lies on the line of intersection of the planes. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes, so take $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, 3, 1\rangle \times \langle 2, -4, -3\rangle = \langle -5, 5, -10\rangle$ or equivalently we can take $\mathbf{v} = \langle 1, -1, 2\rangle$. Parametric equations for the line are $x = 2 + t$, $y = -t$, $z = 4 + 2t$.

27. By Exercise 12.5.75, $D = \dfrac{|-2 - (-24)|}{\sqrt{3^2 + 1^2 + (-4)^2}} = \dfrac{22}{\sqrt{26}}$.

28. The equation $x = 3$ represents a plane parallel to the yz-plane and 3 units in front of it.



29. The equation $x = z$ represents a plane perpendicular to the xz-plane and intersecting the xz-plane in the line $x = z$, $y = 0$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

30. The equation $y = z^2$ represents a parabolic cylinder whose trace in the xz-plane is the x-axis and which opens to the right.



31. The equation $x^2 = y^2 + 4z^2$ represents a (right elliptical) cone with vertex at the origin and axis the x-axis.



32. $4x - y + 2z = 4$ is a plane with intercepts $(1, 0, 0)$, $(0, -4, 0)$, and $(0, 0, 2)$.



33. An equivalent equation is $-x^2 + \frac{y^2}{4} - z^2 = 1$, a hyperboloid of two sheets with axis the y-axis. For $|y| > 2$, traces parallel to the xz-plane are circles.



34. An equivalent equation is $-x^2 + y^2 + z^2 = 1$, a hyperboloid of one sheet with axis the x-axis.



35. Completing the square in y gives $4x^2 + 4(y-1)^2 + z^2 = 4$ or $x^2 + (y-1)^2 + \frac{z^2}{4} = 1$, an ellipsoid centered at $(0, 1, 0)$.



36. Completing the square in y and z gives $x = (y-1)^2 + (z-2)^2$, a circular paraboloid with vertex $(0, 1, 2)$ and axis the horizontal line $y = 1$, $z = 2$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $4x^2 + y^2 = 16 \quad \Leftrightarrow \quad \dfrac{x^2}{4} + \dfrac{y^2}{16} = 1$. The equation of the ellipsoid is $\dfrac{x^2}{4} + \dfrac{y^2}{16} + \dfrac{z^2}{c^2} = 1$, since the horizontal trace in the plane $z = 0$ must be the original ellipse. The traces of the ellipsoid in the yz-plane must be circles since the surface is obtained by rotation about the x-axis. Therefore, $c^2 = 16$ and the equation of the ellipsoid is $\dfrac{x^2}{4} + \dfrac{y^2}{16} + \dfrac{z^2}{16} = 1 \quad \Leftrightarrow$

$4x^2 + y^2 + z^2 = 16$.

38. The distance from a point $P(x, y, z)$ to the plane $y = 1$ is $|y - 1|$, so the given condition becomes

$|y - 1| = 2\sqrt{(x - 0)^2 + (y + 1)^2 + (z - 0)^2} \quad \Rightarrow \quad |y - 1| = 2\sqrt{x^2 + (y + 1)^2 + z^2} \quad \Rightarrow$

$(y - 1)^2 = 4x^2 + 4(y + 1)^2 + 4z^2 \quad \Leftrightarrow \quad -3 = 4x^2 + (3y^2 + 10y) + 4z^2 \quad \Leftrightarrow$

$\frac{16}{3} = 4x^2 + 3\left(y + \frac{5}{3}\right)^2 + 4z^2 \quad \Rightarrow \quad \frac{3}{4}x^2 + \frac{9}{16}\left(y + \frac{5}{3}\right)^2 + \frac{3}{4}z^2 = 1.$

This is the equation of an ellipsoid whose center is $\left(0, -\frac{5}{3}, 0\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

PROBLEMS PLUS

1. Since three-dimensional situations are often difficult to visualize and work with, let us first try to find an analogous problem in two dimensions. The analogue of a cube is a square and the analogue of a sphere is a circle. Thus a similar problem in two dimensions is the following: if five circles with the same radius r are contained in a square of side 1 m so that the circles touch each other and four of the circles touch two sides of the square, find r.



The diagonal of the square is $\sqrt{2}$. The diagonal is also $4r + 2x$. But x is the diagonal of a smaller square of side r. Therefore

$x = \sqrt{2}\,r \quad \Rightarrow \quad \sqrt{2} = 4r + 2x = 4r + 2\sqrt{2}\,r = (4 + 2\sqrt{2})r \quad \Rightarrow \quad r = \frac{\sqrt{2}}{4 + 2\sqrt{2}}$.

Let's use these ideas to solve the original three-dimensional problem. The diagonal of the cube is $\sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}$. The diagonal of the cube is also $4r + 2x$ where x is the diagonal of a smaller cube with edge r. Therefore

$x = \sqrt{r^2 + r^2 + r^2} = \sqrt{3}\,r \quad \Rightarrow \quad \sqrt{3} = 4r + 2x = 4r + 2\sqrt{3}\,r = (4 + 2\sqrt{3})r$. Thus $r = \dfrac{\sqrt{3}}{4 + 2\sqrt{3}} = \dfrac{2\sqrt{3} - 3}{2}$.

The radius of each ball is $\left(\sqrt{3} - \frac{3}{2}\right)$ m.

2. Try an analogous problem in two dimensions. Consider a rectangle with length L and width W and find the area of S in terms of L and W. Since S contains B, it has area



$A(S) = LW +$ the area of two $L \times 1$ rectangles

$+$ the area of two $1 \times W$ rectangles

$+$ the area of four quarter-circles of radius 1

as seen in the diagram. So $A(S) = LW + 2L + 2W + \pi \cdot 1^2$.

Now in three dimensions, the volume of S is

$LWH + 2(L \times W \times 1) + 2(1 \times W \times H) + 2(L \times 1 \times H)$

$+$ the volume of 4 quarter-cylinders with radius 1 and height W

$+$ the volume of 4 quarter-cylinders with radius 1 and height L

$+$ the volume of 4 quarter-cylinders with radius 1 and height H

$+$ the volume of 8 eighths of a sphere of radius 1

So

$$V(S) = LWH + 2LW + 2WH + 2LH + \pi \cdot 1^2 \cdot W + \pi \cdot 1^2 \cdot L + \pi \cdot 1^2 \cdot H + \tfrac{4}{3}\pi \cdot 1^3$$

$$= LWH + 2(LW + WH + LH) + \pi(L + W + H) + \tfrac{4}{3}\pi.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

3. (a) We find the line of intersection L as in Example 12.5.7(b). Observe that the point $(-1, c, c)$ lies on both planes. Now since L lies in both planes, it is perpendicular to both of the normal vectors $\mathbf{n}_1$ and $\mathbf{n}_2$, and thus parallel to their cross product

$$\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ c & 1 & 1 \\ 1 & -c & c \end{vmatrix} = \langle 2c, -c^2+1, -c^2-1 \rangle$$

So symmetric equations of L can be written as $\dfrac{x+1}{-2c} = \dfrac{y-c}{c^2-1} = \dfrac{z-c}{c^2+1}$, provided that $c \neq 0, \pm 1$.

If $c = 0$, then the two planes are given by $y + z = 0$ and $x = -1$, so symmetric equations of L are $x = -1$, $y = -z$. If $c = -1$, then the two planes are given by $-x + y + z = -1$ and $x + y + z = -1$, and they intersect in the line $x = 0$, $y = -z - 1$. If $c = 1$, then the two planes are given by $x + y + z = 1$ and $x - y + z = 1$, and they intersect in the line $y = 0$, $x = 1 - z$.

(b) If we set $z = t$ in the symmetric equations and solve for x and y separately, we get $x + 1 = \dfrac{(t-c)(-2c)}{c^2+1}$, $y - c = \dfrac{(t-c)(c^2-1)}{c^2+1} \quad \Rightarrow \quad x = \dfrac{-2ct + (c^2-1)}{c^2+1}$, $y = \dfrac{(c^2-1)t + 2c}{c^2+1}$. Eliminating c from these equations, we have $x^2 + y^2 = t^2 + 1$. So the curve traced out by L in the plane $z = t$ is a circle with center at $(0, 0, t)$ and radius $\sqrt{t^2+1}$.

(c) The area of a horizontal cross-section of the solid is $A(z) = \pi(z^2+1)$, so $V = \int_0^1 A(z)\,dz = \pi\left[\frac{1}{3}z^3 + z\right]_0^1 = \frac{4\pi}{3}$.

4. (a) We consider velocity vectors for the plane and the wind. Let $\mathbf{v}_i$ be the initial, intended velocity for the plane and $\mathbf{v}_g$ the actual velocity relative to the ground. If $\mathbf{w}$ is the velocity of the wind, $\mathbf{v}_g$ is the resultant, that is, the vector sum $\mathbf{v}_i + \mathbf{w}$ as shown in the figure. We know $\mathbf{v}_i = 180\,\mathbf{j}$, and since the plane actually flew 80 km in $\frac{1}{2}$ hour, $|\mathbf{v}_g| = 160$. Thus $\mathbf{v}_g = (160\cos 85^\circ)\,\mathbf{i} + (160\sin 85^\circ)\,\mathbf{j} \approx 13.9\,\mathbf{i} + 159.4\,\mathbf{j}$. Finally, $\mathbf{v}_i + \mathbf{w} = \mathbf{v}_g$, so $\mathbf{w} = \mathbf{v}_g - \mathbf{v}_i \approx 13.9\,\mathbf{i} - 20.6\,\mathbf{j}$. Thus, the wind velocity is about $13.9\,\mathbf{i} - 20.6\,\mathbf{j}$, and the wind speed is $|\mathbf{w}| \approx \sqrt{(13.9)^2 + (-20.6)^2} \approx 24.9$ km/h.

(b) Let $\mathbf{v}$ be the velocity the pilot should have taken. The pilot would need to head a little west of north to compensate for the wind, so let θ be the angle $\mathbf{v}$ makes with the north direction. Then we can write $\mathbf{v} = \langle 180\cos(\theta + 90^\circ), 180\sin(\theta + 90^\circ) \rangle$. With the effect of the wind, the actual velocity (with respect to the ground) will be $\mathbf{v} + \mathbf{w}$, which we want to be due north. Equating horizontal components of the vectors, we must have $180\cos(\theta + 90^\circ) + 160\cos 85^\circ = 0 \quad \Rightarrow \quad \cos(\theta + 90^\circ) = -\frac{160}{180}\cos 85^\circ \approx -0.0775$, so $\theta \approx \cos^{-1}(-0.0775) - 90^\circ \approx 4.4^\circ$. Thus the pilot should have headed about 4.4° west of north.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\mathbf{v}_3 = \text{proj}_{\mathbf{v}_1}\mathbf{v}_2 = \dfrac{\mathbf{v}_1 \cdot \mathbf{v}_2}{|\mathbf{v}_1|^2}\,\mathbf{v}_1 = \dfrac{5}{2^2}\,\mathbf{v}_1$ so $|\mathbf{v}_3| = \dfrac{5}{2^2}\,|\mathbf{v}_1| = \dfrac{5}{2}$,

$\mathbf{v}_4 = \text{proj}_{\mathbf{v}_2}\mathbf{v}_3 = \dfrac{\mathbf{v}_2 \cdot \mathbf{v}_3}{|\mathbf{v}_2|^2}\,\mathbf{v}_2 = \dfrac{\mathbf{v}_2 \cdot \frac{5}{2^2}\,\mathbf{v}_1}{|\mathbf{v}_2|^2}\,\mathbf{v}_2 = \dfrac{5}{2^2 \cdot 3^2}\,(\mathbf{v}_1 \cdot \mathbf{v}_2)\,\mathbf{v}_2 = \dfrac{5^2}{2^2 \cdot 3^2}\,\mathbf{v}_2 \quad \Rightarrow \quad |\mathbf{v}_4| = \dfrac{5^2}{2^2 \cdot 3^2}\,|\mathbf{v}_2| = \dfrac{5^2}{2^2 \cdot 3}$,

$\mathbf{v}_5 = \text{proj}_{\mathbf{v}_3}\mathbf{v}_4 = \dfrac{\mathbf{v}_3 \cdot \mathbf{v}_4}{|\mathbf{v}_3|^2}\,\mathbf{v}_3 = \dfrac{\frac{5}{2^2}\,\mathbf{v}_1 \cdot \frac{5^2}{2^2 3^2}\,\mathbf{v}_2}{\left(\frac{5}{2}\right)^2}\left(\dfrac{5}{2^2}\,\mathbf{v}_1\right) = \dfrac{5^2}{2^4 \cdot 3^2}\,(\mathbf{v}_1 \cdot \mathbf{v}_2)\,\mathbf{v}_1 = \dfrac{5^3}{2^4 \cdot 3^2}\,\mathbf{v}_1 \quad \Rightarrow$

$|\mathbf{v}_5| = \dfrac{5^3}{2^4 \cdot 3^2}\,|\mathbf{v}_1| = \dfrac{5^3}{2^3 \cdot 3^2}$. Similarly, $|\mathbf{v}_6| = \dfrac{5^4}{2^4 \cdot 3^3}$, $|\mathbf{v}_7| = \dfrac{5^5}{2^5 \cdot 3^4}$, and in general, $|\mathbf{v}_n| = \dfrac{5^{n-2}}{2^{n-2} \cdot 3^{n-3}} = 3\left(\frac{5}{6}\right)^{n-2}$.

Thus

$$\sum_{n=1}^{\infty} |\mathbf{v}_n| = |\mathbf{v}_1| + |\mathbf{v}_2| + \sum_{n=3}^{\infty} 3\left(\tfrac{5}{6}\right)^{n-2} = 2 + 3 + \sum_{n=1}^{\infty} 3\left(\tfrac{5}{6}\right)^{n}$$

$$= 5 + \sum_{n=1}^{\infty} \tfrac{5}{2}\left(\tfrac{5}{6}\right)^{n-1} = 5 + \frac{\frac{5}{2}}{1 - \frac{5}{6}} \quad \text{[sum of a geometric series]} \quad = 5 + 15 = 20$$

6. Completing squares in the inequality $x^2 + y^2 + z^2 < 136 + 2(x + 2y + 3z)$ gives $(x-1)^2 + (y-2)^2 + (z-3)^2 < 150$ which describes the set of all points (x, y, z) whose distance from the point $P\,(1, 2, 3)$ is less than $\sqrt{150} = 5\sqrt{6}$. The distance from P to $Q\,(-1, 1, 4)$ is $\sqrt{4 + 1 + 1} = \sqrt{6}$, so the largest possible sphere that passes through Q and satisfies the stated conditions extends $5\sqrt{6}$ units in the opposite direction, giving a diameter of $6\sqrt{6}$. (See the figure.)



Thus the radius of the desired sphere is $3\sqrt{6}$, and its center is $3\sqrt{6}$ units from Q in the direction of P. A unit vector in this direction is $\mathbf{u} = \frac{1}{\sqrt{6}}\,\langle 2, 1, -1\rangle$, so starting at $Q(-1, 1, 4)$ and following the vector $3\sqrt{6}\,\mathbf{u} = \langle 6, 3, -3\rangle$ we arrive at the center of the sphere, $(5, 4, 1)$. An equation of the sphere then is $(x-5)^2 + (y-4)^2 + (z-1)^2 = \left(3\sqrt{6}\right)^2$ or $(x-5)^2 + (y-4)^2 + (z-1)^2 = 54$.

7. (a) When $\theta = \theta_s$, the block is not moving, so the sum of the forces on the block must be $\mathbf{0}$, thus $\mathbf{N} + \mathbf{F} + \mathbf{W} = \mathbf{0}$. This relationship is illustrated geometrically in the figure. Since the vectors form a right triangle, we have $\tan(\theta_s) = \dfrac{|\mathbf{F}|}{|\mathbf{N}|} = \dfrac{\mu_s n}{n} = \mu_s$.



(b) We place the block at the origin and sketch the force vectors acting on the block, including the additional horizontal force $\mathbf{H}$, with initial points at the origin. We then rotate this system so that $\mathbf{F}$ lies along the positive x-axis and the inclined plane is parallel to the x-axis. (See the following figure.)

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part





$|\mathbf{F}|$ is maximal, so $|\mathbf{F}| = \mu_s n$ for $\theta > \theta_s$. Then the vectors, in terms of components parallel and perpendicular to the inclined plane, are

$$\mathbf{N} = n\,\mathbf{j} \qquad \mathbf{F} = (\mu_s n)\,\mathbf{i}$$

$$\mathbf{W} = (-mg\sin\theta)\,\mathbf{i} + (-mg\cos\theta)\,\mathbf{j} \qquad \mathbf{H} = (h_{\min}\cos\theta)\,\mathbf{i} + (-h_{\min}\sin\theta)\,\mathbf{j}$$

Equating components, we have

$$\mu_s n - mg\sin\theta + h_{\min}\cos\theta = 0 \quad \Rightarrow \quad h_{\min}\cos\theta + \mu_s n = mg\sin\theta \tag{1}$$

$$n - mg\cos\theta - h_{\min}\sin\theta = 0 \quad \Rightarrow \quad h_{\min}\sin\theta + mg\cos\theta = n \tag{2}$$

(c) Since **(2)** is solved for n, we substitute into **(1)**:

$$h_{\min}\cos\theta + \mu_s(h_{\min}\sin\theta + mg\cos\theta) = mg\sin\theta \quad \Rightarrow$$

$$h_{\min}\cos\theta + h_{\min}\mu_s\sin\theta = mg\sin\theta - mg\mu_s\cos\theta \quad \Rightarrow$$

$$h_{\min} = mg\left(\frac{\sin\theta - \mu_s\cos\theta}{\cos\theta + \mu_s\sin\theta}\right) = mg\left(\frac{\tan\theta - \mu_s}{1 + \mu_s\tan\theta}\right)$$

From part (a) we know $\mu_s = \tan\theta_s$, so this becomes $h_{\min} = mg\left(\dfrac{\tan\theta - \tan\theta_s}{1 + \tan\theta_s\tan\theta}\right)$ and using a trigonometric identity, this is $mg\tan(\theta - \theta_s)$ as desired.

Note for $\theta = \theta_s$, $h_{\min} = mg\tan 0 = 0$, which makes sense since the block is at rest for θ_s, thus no additional force $\mathbf{H}$ is necessary to prevent it from moving. As θ increases, the factor $\tan(\theta - \theta_s)$, and hence the value of $h_{\min}$, increases slowly for small values of $\theta - \theta_s$ but much more rapidly as $\theta - \theta_s$ becomes significant. This seems reasonable, as the steeper the inclined plane, the less the horizontal components of the various forces affect the movement of the block, so we would need a much larger magnitude of horizontal force to keep the block motionless. If we allow $\theta \to 90^\circ$, corresponding to the inclined plane being placed vertically, the value of $h_{\min}$ is quite large; this is to be expected, as it takes a great amount of horizontal force to keep an object from moving vertically. In fact, without friction (so $\theta_s = 0$), we would have $\theta \to 90^\circ \quad \Rightarrow \quad h_{\min} \to \infty$, and it would be impossible to keep the block from slipping.

(d) Since $h_{\max}$ is the largest value of h that keeps the block from slipping, the force of friction is keeping the block from moving *up* the inclined plane; thus, $\mathbf{F}$ is directed *down* the plane. Our system of forces is similar to that in part (b), then, except that we have $\mathbf{F} = -(\mu_s n)\,\mathbf{i}$. (Note that $|\mathbf{F}|$ is again maximal.) Following our procedure in parts (b) and (c), we

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

equate components:

$$-\mu_s n - mg\sin\theta + h_{\max}\cos\theta = 0 \quad \Rightarrow \quad h_{\max}\cos\theta - \mu_s n = mg\sin\theta$$

$$n - mg\cos\theta - h_{\max}\sin\theta = 0 \quad \Rightarrow \quad h_{\max}\sin\theta + mg\cos\theta = n$$

Then substituting,

$$h_{\max}\cos\theta - \mu_s(h_{\max}\sin\theta + mg\cos\theta) = mg\sin\theta \quad \Rightarrow$$

$$h_{\max}\cos\theta - h_{\max}\mu_s\sin\theta = mg\sin\theta + mg\mu_s\cos\theta \quad \Rightarrow$$

$$h_{\max} = mg\left(\frac{\sin\theta + \mu_s\cos\theta}{\cos\theta - \mu_s\sin\theta}\right) = mg\left(\frac{\tan\theta + \mu_s}{1 - \mu_s\tan\theta}\right)$$

$$= mg\left(\frac{\tan\theta + \tan\theta_s}{1 - \tan\theta_s\tan\theta}\right) = mg\tan(\theta + \theta_s)$$

We would expect $h_{\max}$ to increase as θ increases, with similar behavior as we established for $h_{\min}$, but with $h_{\max}$ values always larger than $h_{\min}$. We can see that this is the case if we graph $h_{\max}$ as a function of θ, as the curve is the graph of $h_{\min}$ translated $2\theta_s$ to the left, so the equation does seem reasonable. Notice that the equation predicts $h_{\max} \to \infty$ as $\theta \to (90° - \theta_s)$. In fact, as $h_{\max}$ increases, the normal force increases as well. When $(90° - \theta_s) \le \theta \le 90°$, the horizontal force is completely counteracted by the sum of the normal and frictional forces, so no part of the horizontal force contributes to moving the block up the plane no matter how large its magnitude.

8. (a) The largest possible solid is achieved by starting with a circular cylinder of diameter 1 with axis the z-axis and with a height of 1. This is the largest solid that creates a square shadow with side length 1 in the y-direction and a circular disk shadow in the z-direction. For convenience, we place the base of the cylinder on the xy-plane so that it intersects the x- and y-axes at $\pm\frac{1}{2}$. We then remove as little as possible from the solid that leaves an isosceles triangle shadow in the x-direction. This is achieved by cutting the solid with planes parallel to the x-axis that intersect the z-axis at 1 and the y-axis at $\pm\frac{1}{2}$ (see the figure).



To compute the volume of this solid, we take vertical slices parallel to the xz-plane. The equation of the base of the solid is $x^2 + y^2 = \frac{1}{4}$, so a cross-section y units to the right of the origin is a rectangle with base $2\sqrt{\frac{1}{4} - y^2}$. For $0 \le y \le \frac{1}{2}$, the solid is cut off on top by the plane $z = 1 - 2y$, so the height of the rectangular cross-section is $1 - 2y$ and the cross-sectional area is $A(y) = 2\sqrt{\frac{1}{4} - y^2}\,(1 - 2y)$. The volume of the right half of the solid is

$$\int_0^{1/2} 2\sqrt{\tfrac{1}{4} - y^2}\,(1 - 2y)\,dy = 2\int_0^{1/2}\sqrt{\tfrac{1}{4} - y^2}\,dy - 4\int_0^{1/2} y\sqrt{\tfrac{1}{4} - y^2}\,dy$$

$$= 2\left[\tfrac{1}{4}\text{ area of a circle of radius }\tfrac{1}{2}\right] - 4\left[-\tfrac{1}{3}\left(\tfrac{1}{4} - y^2\right)^{3/2}\right]_0^{1/2}$$

$$= 2\left[\tfrac{1}{4}\cdot\pi\left(\tfrac{1}{2}\right)^2\right] + \tfrac{4}{3}\left[0 - \left(\tfrac{1}{4}\right)^{3/2}\right] = \tfrac{\pi}{8} - \tfrac{1}{6}$$

Thus the volume of the solid is $2\left(\frac{\pi}{8} - \frac{1}{6}\right) = \frac{\pi}{4} - \frac{1}{3} \approx 0.45$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) There is not a smallest volume. We can remove portions of the solid from part (a) to create smaller and smaller volumes as long as we leave the "skeleton" shown in the figure intact (which has no volume at all and is not a solid). Thus we can create solids with arbitrarily small volume.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13 □ VECTOR FUNCTIONS

13.1 Vector Functions and Space Curves

1. The component functions $\ln(t+1)$, $\dfrac{t}{\sqrt{9-t^2}}$, and 2^t are all defined when $t+1>0 \;\Rightarrow\; t>-1$ and $9-t^2>0 \;\Rightarrow$ $-3<t<3$, so the domain of $\mathbf{r}$ is $(-1,3)$.

2. The component functions $\cos t$, $\ln t$, and $\dfrac{1}{t-2}$ are all defined when $t>0$ and $t\neq 2$, so the domain of $\mathbf{r}$ is $(0,2)\cup(2,\infty)$.

3. $\lim\limits_{t\to 0} e^{-3t} = e^0 = 1$, $\lim\limits_{t\to 0} \dfrac{t^2}{\sin^2 t} = \lim\limits_{t\to 0} \dfrac{1}{\dfrac{\sin^2 t}{t^2}} = \dfrac{1}{\lim\limits_{t\to 0} \dfrac{\sin^2 t}{t^2}} = \dfrac{1}{\left(\lim\limits_{t\to 0} \dfrac{\sin t}{t}\right)^2} = \dfrac{1}{1^2} = 1$,

and $\lim\limits_{t\to 0} \cos 2t = \cos 0 = 1$. Thus

$$\lim_{t\to 0}\left(e^{-3t}\,\mathbf{i} + \frac{t^2}{\sin^2 t}\,\mathbf{j} + \cos 2t\,\mathbf{k}\right) = \left[\lim_{t\to 0} e^{-3t}\right]\mathbf{i} + \left[\lim_{t\to 0}\frac{t^2}{\sin^2 t}\right]\mathbf{j} + \left[\lim_{t\to 0}\cos 2t\right]\mathbf{k} = \mathbf{i}+\mathbf{j}+\mathbf{k}.$$

4. $\lim\limits_{t\to 1} \dfrac{t^2-t}{t-1} = \lim\limits_{t\to 1} \dfrac{t(t-1)}{t-1} = \lim\limits_{t\to 1} t = 1$, $\lim\limits_{t\to 1} \sqrt{t+8} = 3$, $\lim\limits_{t\to 1} \dfrac{\sin \pi t}{\ln t} = \lim\limits_{t\to 1} \dfrac{\pi \cos \pi t}{1/t} = -\pi$ [by l'Hospital's Rule].

Thus the given limit equals $\mathbf{i} + 3\,\mathbf{j} - \pi\,\mathbf{k}$.

5. $\lim\limits_{t\to\infty} \dfrac{1+t^2}{1-t^2} = \lim\limits_{t\to\infty} \dfrac{(1/t^2)+1}{(1/t^2)-1} = \dfrac{0+1}{0-1} = -1$, $\lim\limits_{t\to\infty} \tan^{-1} t = \dfrac{\pi}{2}$, $\lim\limits_{t\to\infty} \dfrac{1-e^{-2t}}{t} = \lim\limits_{t\to\infty} \dfrac{1}{t} - \dfrac{1}{te^{2t}} = 0-0=0$. Thus

$$\lim_{t\to\infty}\left\langle \frac{1+t^2}{1-t^2}, \tan^{-1} t, \frac{1-e^{-2t}}{t}\right\rangle = \left\langle -1, \tfrac{\pi}{2}, 0\right\rangle.$$

6. $\lim\limits_{t\to\infty} te^{-t} = \lim\limits_{t\to\infty} \dfrac{t}{e^t} = \lim\limits_{t\to\infty} \dfrac{1}{e^t} = 0$ [by l'Hospital's Rule], $\lim\limits_{t\to\infty} \dfrac{t^3+t}{2t^3-1} = \lim\limits_{t\to\infty} \dfrac{1+(1/t^2)}{2-(1/t^3)} = \dfrac{1+0}{2-0} = \dfrac{1}{2}$,

and $\lim\limits_{t\to\infty} t\sin\dfrac{1}{t} = \lim\limits_{t\to\infty} \dfrac{\sin(1/t)}{1/t} = \lim\limits_{t\to\infty} \dfrac{\cos(1/t)(-1/t^2)}{-1/t^2} = \lim\limits_{t\to\infty} \cos\dfrac{1}{t} = \cos 0 = 1$ [again by l'Hospital's Rule].

Thus $\lim\limits_{t\to\infty}\left\langle te^{-t}, \dfrac{t^3+t}{2t^3-1}, t\sin\dfrac{1}{t}\right\rangle = \left\langle 0, \tfrac{1}{2}, 1\right\rangle$.

7. The corresponding parametric equations for this curve are $x=\sin t,\ y=t$.

We can make a table of values, or we can eliminate the parameter: $t=y \;\Rightarrow$ $x=\sin y$, with $y\in\mathbb{R}$. By comparing different values of t, we find the direction in which t increases as indicated in the graph.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. The corresponding parametric equations for this curve are $x = t^2 - 1,\ y = t$. We can make a table of values, or we can eliminate the parameter:

$t = y \quad \Rightarrow \quad x = y^2 - 1$, with $y \in \mathbb{R}$. Thus the curve is a parabola with vertex $(-1, 0)$ that opens to the right. By comparing different values of t, we find the direction in which t increases as indicated in the graph.



9. The corresponding parametric equations are $x = t,\ y = 2 - t,\ z = 2t$, which are parametric equations of a line through the point $(0, 2, 0)$ and with direction vector $\langle 1, -1, 2 \rangle$.



10. The corresponding parametric equations are $x = \sin \pi t,\ y = t,\ z = \cos \pi t$. Note that $x^2 + z^2 = \sin^2 \pi t + \cos^2 \pi t = 1$, so the curve lies on the circular cylinder $x^2 + z^2 = 1$. A point (x, y, z) on the curve lies directly to the left or right of the point $(x, 0, z)$ which moves clockwise (when viewed from the left) along the circle $x^2 + z^2 = 1$ in the xz-plane as t increases. Since $y = t$, the curve is a helix that spirals toward the right around the cylinder.



11. The corresponding parametric equations are $x = 3,\ y = t,\ z = 2 - t^2$. Eliminating the parameter in y and z gives $z = 2 - y^2$. Because $x = 3$, the curve is a parabola in the vertical plane $x = 3$ with vertex $(3, 0, 2)$.



12. The corresponding parametric equations are $x = 2 \cos t,\ y = 2 \sin t$, $z = 1$. Eliminating the parameter in x and y gives $x^2 + y^2 = 4 \cos^2 t + 4 \sin^2 t = 4(\cos^2 t + \sin^2 t) = 4$. Since $z = 1$, the curve is a circle of radius 2 centered at $(0, 0, 1)$ in the horizontal plane $z = 1$.



13. The parametric equations are $x = t^2, y = t^4, z = t^6$. These are positive for $t \neq 0$ and 0 when $t = 0$. So the curve lies entirely in the first octant. The projection of the graph onto the xy-plane is $y = x^2,\ y > 0$, a half parabola. The projection onto the xz-plane is $z = x^3,\ z > 0$, a half cubic, and the projection onto the yz-plane is $y^3 = z^2$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. If $x = \cos t,\ y = -\cos t,\ z = \sin t$, then $x^2 + z^2 = 1$ and $y^2 + z^2 = 1$, so the curve is contained in the intersection of circular cylinders along the x- and y-axes. Furthermore, $y = -x$, so the curve is an ellipse in the plane $y = -x$, centered at the origin.



15. The projection of the curve onto the xy-plane is given by $\mathbf{r}(t) = \langle t, \sin t, 0\rangle$ [we use 0 for the z-component] whose graph is the curve $y = \sin x,\ z = 0$. Similarly, the projection onto the xz-plane is $\mathbf{r}(t) = \langle t, 0, 2\cos t\rangle$, whose graph is the cosine wave $z = 2\cos x,\ y = 0$, and the projection onto the yz-plane is $\mathbf{r}(t) = \langle 0, \sin t, 2\cos t\rangle$ whose graph is the ellipse $y^2 + \frac{1}{4}z^2 = 1,\ x = 0$.



xy-plane



xz-plane



yz-plane

From the projection onto the yz-plane we see that the curve lies on an elliptical cylinder with axis the x-axis. The other two projections show that the curve oscillates both vertically and horizontally as we move in the x-direction, suggesting that the curve is an elliptical helix that spirals along the cylinder.



16. The projection of the curve onto the xy-plane is given by $\mathbf{r}(t) = \langle t, t, 0\rangle$ whose graph is the line $y = x,\ z = 0$.
The projection onto the xz-plane is $\mathbf{r}(t) = \langle t, 0, t^2\rangle$ whose graph is the parabola $z = x^2,\ y = 0$.
The projection onto the yz-plane is $\mathbf{r}(t) = \langle 0, t, t^2\rangle$ whose graph is the parabola $z = y^2,\ x = 0$.



xy-plane



xz-plane



yz-plane

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

From the projection onto the xy-plane we see that the curve lies on the vertical plane $y = x$. The other two projections show that the curve is a parabola contained in this plane.



17. Taking $\mathbf{r}_0 = \langle 2, 0, 0\rangle$ and $\mathbf{r}_1 = \langle 6, 2, -2\rangle$, we have from Equation 12.5.4

$\mathbf{r}(t) = (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)\,\langle 2,0,0\rangle + t\,\langle 6,2,-2\rangle, 0 \le t \le 1$ or $\mathbf{r}(t) = \langle 2+4t, 2t, -2t\rangle, 0 \le t \le 1$.

Parametric equations are $x = 2+4t,\ y = 2t,\ z = -2t,\ 0 \le t \le 1$.

18. Taking $\mathbf{r}_0 = \langle -1, 2, -2\rangle$ and $\mathbf{r}_1 = \langle -3, 5, 1\rangle$, we have from Equation 12.5.4

$\mathbf{r}(t) = (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)\,\langle -1,2,-2\rangle + t\,\langle -3,5,1\rangle, 0 \le t \le 1$ or $\mathbf{r}(t) = \langle -1-2t, 2+3t, -2+3t\rangle, 0 \le t \le 1$.

Parametric equations are $x = -1-2t,\ y = 2+3t,\ z = -2+3t,\ 0 \le t \le 1$.

19. Taking $\mathbf{r}_0 = \langle 0, -1, 1\rangle$ and $\mathbf{r}_1 = \left\langle \frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right\rangle$, we have

$\mathbf{r}(t) = (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)\,\langle 0,-1,1\rangle + t\left\langle \frac{1}{2}, \frac{1}{3}, \frac{1}{4}\right\rangle, 0 \le t \le 1$ or $\mathbf{r}(t) = \left\langle \frac{1}{2}t, -1+\frac{4}{3}t, 1-\frac{3}{4}t\right\rangle, 0 \le t \le 1$.

Parametric equations are $x = \frac{1}{2}t,\ y = -1+\frac{4}{3}t,\ z = 1-\frac{3}{4}t,\ 0 \le t \le 1$.

20. Taking $\mathbf{r}_0 = \langle a, b, c\rangle$ and $\mathbf{r}_1 = \langle u, v, w\rangle$, we have

$\mathbf{r}(t) = (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)\,\langle a,b,c\rangle + t\,\langle u,v,w\rangle, 0 \le t \le 1$ or $\mathbf{r}(t) = \langle a+(u-a)t, b+(v-b)t, c+(w-c)t\rangle$, $0 \le t \le 1$. Parametric equations are $x = a+(u-a)t,\ y = b+(v-b)t,\ z = c+(w-c)t,\ 0 \le t \le 1$.

21. $x = t\cos t,\ y = t,\ z = t\sin t,\ t \ge 0$. At any point (x, y, z) on the curve, $x^2 + z^2 = t^2\cos^2 t + t^2\sin^2 t = t^2 = y^2$ so the curve lies on the circular cone $x^2 + z^2 = y^2$ with axis the y-axis. Also notice that $y \ge 0$; the graph is II.

22. $x = \cos t,\ y = \sin t,\ z = 1/(1+t^2)$. At any point on the curve we have $x^2 + y^2 = \cos^2 t + \sin^2 t = 1$, so the curve lies on the circular cylinder $x^2 + y^2 = 1$ with axis the z-axis. Notice that $0 < z \le 1$ and $z = 1$ only for $t = 0$. A point (x, y, z) on the curve lies directly above the point $(x, y, 0)$, which moves counterclockwise around the unit circle in the xy-plane as t increases, and $z \to 0$ as $t \to \pm\infty$. The graph must be VI.

23. $x = t,\ y = 1/(1+t^2),\ z = t^2$. At any point on the curve we have $z = x^2$, so the curve lies on a parabolic cylinder parallel to the y-axis. Notice that $0 < y \le 1$ and $z \ge 0$. Also the curve passes through $(0, 1, 0)$ when $t = 0$ and $y \to 0$, $z \to \infty$ as $t \to \pm\infty$, so the graph must be V.

24. $x = \cos t,\ y = \sin t,\ z = \cos 2t$. $x^2 + y^2 = \cos^2 t + \sin^2 t = 1$, so the curve lies on a circular cylinder with axis the z-axis. A point (x, y, z) on the curve lies directly above or below $(x, y, 0)$, which moves around the unit circle in the xy-plane with period 2π. At the same time, the z-value of the point (x, y, z) oscillates with a period of π. So the curve repeats itself and the graph is I.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $x = \cos 8t,\ y = \sin 8t,\ z = e^{0.8t},\ t \geq 0.$ $x^2 + y^2 = \cos^2 8t + \sin^2 8t = 1$, so the curve lies on a circular cylinder with axis the z-axis. A point (x, y, z) on the curve lies directly above the point $(x, y, 0)$, which moves counterclockwise around the unit circle in the xy-plane as t increases. The curve starts at $(1, 0, 1)$, when $t = 0$, and $z \to \infty$ (at an increasing rate) as $t \to \infty$, so the graph is IV.

26. $x = \cos^2 t,\ y = \sin^2 t,\ z = t.$ $x + y = \cos^2 t + \sin^2 t = 1$, so the curve lies in the vertical plane $x + y = 1$. x and y are periodic, both with period π, and z increases as t increases, so the graph is III.

27. If $x = t\cos t,\ y = t\sin t,\ z = t$, then $x^2 + y^2 = t^2\cos^2 t + t^2\sin^2 t = t^2 = z^2$, so the curve lies on the cone $z^2 = x^2 + y^2$. Since $z = t$, the curve is a spiral on this cone.



28. Here $x^2 = \sin^2 t = z$ and $x^2 + y^2 = \sin^2 t + \cos^2 t = 1$, so the curve is contained in the intersection of the parabolic cylinder $z = x^2$ with the circular cylinder $x^2 + y^2 = 1$. We get the complete intersection for $0 \leq t \leq 2\pi$.



29. Here $x = 2t,\ y = e^t,\ z = e^{2t}$. Then $t = x/2 \Rightarrow y = e^t = e^{x/2}$, so the curve lies on the cylinder $y = e^{x/2}$. Also $z = e^{2t} = e^x$, so the curve lies on the cylinder $z = e^x$. Since $z = e^{2t} = \left(e^t\right)^2 = y^2$, the curve also lies on the parabolic cylinder $z = y^2$.

30. Here $x = t^2,\ y = \ln t,\ z = 1/t$. The domain of $\mathbf{r}$ is $(0, \infty)$, so $x = t^2 \Rightarrow t = \sqrt{x} \Rightarrow y = \ln\sqrt{x}$. Thus one surface containing the curve is the cylinder $y = \ln\sqrt{x}$ or $y = \ln x^{1/2} = \frac{1}{2}\ln x$. Also $z = 1/t = 1/\sqrt{x}$, so the curve also lies on the cylinder $z = 1/\sqrt{x}$ or $x = 1/z^2,\ z > 0$. Finally $z = 1/t \Rightarrow t = 1/z \Rightarrow y = \ln(1/z)$, so the curve also lies on the cylinder $y = \ln(1/z)$ or $y = \ln z^{-1} = -\ln z$. Note that the surface $y = \ln(xz)$ also contains the curve, since $\ln(xz) = \ln(t^2 \cdot 1/t) = \ln t = y$.

31. Parametric equations for the curve are $x = t,\ y = 0,\ z = 2t - t^2$. Substituting into the equation of the paraboloid gives $2t - t^2 = t^2 \Rightarrow 2t = 2t^2 \Rightarrow t = 0, 1$. Since $\mathbf{r}(0) = \mathbf{0}$ and $\mathbf{r}(1) = \mathbf{i} + \mathbf{k}$, the points of intersection are $(0, 0, 0)$ and $(1, 0, 1)$.

32. Parametric equations for the helix are $x = \sin t,\ y = \cos t,\ z = t$. Substituting into the equation of the sphere gives $\sin^2 t + \cos^2 t + t^2 = 5 \Rightarrow 1 + t^2 = 5 \Rightarrow t = \pm 2$. Since $\mathbf{r}(2) = \langle \sin 2, \cos 2, 2\rangle$ and $\mathbf{r}(-2) = \langle \sin(-2), \cos(-2), -2\rangle$, the points of intersection are $(\sin 2, \cos 2, 2) \approx (0.909, -0.416, 2)$ and $(\sin(-2), \cos(-2), -2) \approx (-0.909, -0.416, -2)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

33. $\mathbf{r}(t) = \langle \cos t \sin 2t, \sin t \sin 2t, \cos 2t \rangle$.

We include both a regular plot and a plot showing a tube of radius 0.08 around the curve.





34. $\mathbf{r}(t) = \langle te^t, e^{-t}, t \rangle$



35. $\mathbf{r}(t) = \langle \sin 3t \cos t, \frac{1}{4}t, \sin 3t \sin t \rangle$



36. $\mathbf{r}(t) = \langle \cos(8 \cos t) \sin t, \sin(8 \cos t) \sin t, \cos t \rangle$



37. $\mathbf{r}(t) = \langle \cos 2t, \cos 3t, \cos 4t \rangle$



38. $x = \sin t$, $y = \sin 2t$, $z = \cos 4t$.



We graph the projections onto the coordinate planes.



xy-plane



xz-plane



yz-plane

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

From the projection onto the xy-plane we see that from above the curve appears to be shaped like a "figure eight." The curve can be visualized as this shape wrapped around an almost parabolic cylindrical surface, the profile of which is visible in the projection onto the yz-plane.

39.



$x = (1 + \cos 16t)\cos t$, $y = (1 + \cos 16t)\sin t$, $z = 1 + \cos 16t$. At any point on the graph,

$$\begin{aligned} x^2 + y^2 &= (1 + \cos 16t)^2 \cos^2 t + (1 + \cos 16t)^2 \sin^2 t \\ &= (1 + \cos 16t)^2 = z^2, \text{ so the graph lies on the cone } x^2 + y^2 = z^2. \end{aligned}$$

From the graph at left, we see that this curve looks like the projection of a leaved two-dimensional curve onto a cone.

40.



$x = \sqrt{1 - 0.25\cos^2 10t}\cos t$, $y = \sqrt{1 - 0.25\cos^2 10t}\sin t$, $z = 0.5\cos 10t$. At any point on the graph,

$$\begin{aligned} x^2 + y^2 + z^2 &= (1 - 0.25\cos^2 10t)\cos^2 t \\ &\quad + (1 - 0.25\cos^2 10t)\sin^2 t + 0.25\cos^2 t \\ &= 1 - 0.25\cos^2 10t + 0.25\cos^2 10t = 1, \end{aligned}$$

so the graph lies on the sphere $x^2 + y^2 + z^2 = 1$, and since $z = 0.5\cos 10t$ the graph resembles a trigonometric curve with ten peaks projected onto the sphere. We get the complete graph for $0 \le t \le 2\pi$.

41. If $t = -1$, then $x = 1$, $y = 4$, $z = 0$, so the curve passes through the point $(1, 4, 0)$. If $t = 3$, then $x = 9$, $y = -8$, $z = 28$, so the curve passes through the point $(9, -8, 28)$. For the point $(4, 7, -6)$ to be on the curve, we require $y = 1 - 3t = 7 \Rightarrow t = -2$. But then $z = 1 + (-2)^3 = -7 \neq -6$, so $(4, 7, -6)$ is not on the curve.

42. The projection of the curve C of intersection onto the xy-plane is the circle $x^2 + y^2 = 4$, $z = 0$. Then we can write $x = 2\cos t$, $y = 2\sin t$, $0 \le t \le 2\pi$. Since C also lies on the surface $z = xy$, we have $z = xy = (2\cos t)(2\sin t) = 4\cos t \sin t$, or $2\sin(2t)$. Then parametric equations for C are $x = 2\cos t$, $y = 2\sin t$, $z = 2\sin(2t)$, $0 \le t \le 2\pi$, and the corresponding vector function is $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} + 2\sin(2t)\,\mathbf{k}$, $0 \le t \le 2\pi$.

43. Both equations are solved for z, so we can substitute to eliminate z: $\sqrt{x^2 + y^2} = 1 + y \Rightarrow x^2 + y^2 = 1 + 2y + y^2 \Rightarrow x^2 = 1 + 2y \Rightarrow y = \frac{1}{2}(x^2 - 1)$. We can form parametric equations for the curve C of intersection by choosing a parameter $x = t$, then $y = \frac{1}{2}(t^2 - 1)$ and $z = 1 + y = 1 + \frac{1}{2}(t^2 - 1) = \frac{1}{2}(t^2 + 1)$. Thus a vector function representing C is $\mathbf{r}(t) = t\,\mathbf{i} + \frac{1}{2}(t^2 - 1)\,\mathbf{j} + \frac{1}{2}(t^2 + 1)\,\mathbf{k}$.

44. The projection of the curve C of intersection onto the xy-plane is the parabola $y = x^2$, $z = 0$. Then we can choose the parameter $x = t \Rightarrow y = t^2$. Since C also lies on the surface $z = 4x^2 + y^2$, we have $z = 4x^2 + y^2 = 4t^2 + (t^2)^2$. Then parametric equations for C are $x = t$, $y = t^2$, $z = 4t^2 + t^4$, and the corresponding vector function is $\mathbf{r}(t) = t\,\mathbf{i} + t^2\,\mathbf{j} + (4t^2 + t^4)\,\mathbf{k}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

45. The projection of the curve C of intersection onto the xy-plane is the circle $x^2 + y^2 = 1$, $z = 0$, so we can write $x = \cos t$, $y = \sin t$, $0 \le t \le 2\pi$. Since C also lies on the surface $z = x^2 - y^2$, we have $z = x^2 - y^2 = \cos^2 t - \sin^2 t$ or $\cos 2t$. Thus parametric equations for C are $x = \cos t$, $y = \sin t$, $z = \cos 2t$, $0 \le t \le 2\pi$, and the corresponding vector function is $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + \cos 2t\,\mathbf{k}$, $0 \le t \le 2\pi$.

46. The projection of the curve C of intersection onto the xz-plane is the circle $x^2 + z^2 = 1$, $y = 0$, so we can write $x = \cos t$, $z = \sin t$, $0 \le t \le 2\pi$. C also lies on the surface $x^2 + y^2 + 4z^2 = 4$, and since $y \ge 0$ we can write

$$y = \sqrt{4 - x^2 - 4z^2} = \sqrt{4 - \cos^2 t - 4\sin^2 t} = \sqrt{4 - \cos^2 t - 4(1 - \cos^2 t)} = \sqrt{3\cos^2 t} = \sqrt{3}\,|\cos t\,|$$

Thus parametric equations for C are $x = \cos t$, $y = \sqrt{3}\,|\cos t\,|$, $z = \sin t$, $0 \le t \le 2\pi$, and the corresponding vector function is $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sqrt{3}\,|\cos t\,|\,\mathbf{j} + \sin t\,\mathbf{k}$, $0 \le t \le 2\pi$.

47.





The projection of the curve C of intersection onto the xy-plane is the circle $x^2 + y^2 = 4$, $z = 0$. Then we can write $x = 2\cos t,\ y = 2\sin t,\ 0 \le t \le 2\pi$. Since C also lies on the surface $z = x^2$, we have $z = x^2 = (2\cos t)^2 = 4\cos^2 t$. Then parametric equations for C are $x = 2\cos t,\ y = 2\sin t,\ z = 4\cos^2 t,\ 0 \le t \le 2\pi$.

48.





$x = t \ \Rightarrow\ y = t^2 \ \Rightarrow\ 4z^2 = 16 - x^2 - 4y^2 = 16 - t^2 - 4t^4 \ \Rightarrow\ z = \sqrt{4 - \left(\frac{1}{2}t\right)^2 - t^4}$.

Note that z is positive because the intersection is with the top half of the ellipsoid. Hence the curve is given by $x = t,\ y = t^2,\ z = \sqrt{4 - \frac{1}{4}t^2 - t^4}$.

49. For the particles to collide, we require $\mathbf{r}_1(t) = \mathbf{r}_2(t) \ \Leftrightarrow\ \langle t^2, 7t - 12, t^2\rangle = \langle 4t - 3, t^2, 5t - 6\rangle$. Equating components gives $t^2 = 4t - 3$, $7t - 12 = t^2$, and $t^2 = 5t - 6$. From the first equation, $t^2 - 4t + 3 = 0 \ \Leftrightarrow\ (t - 3)(t - 1) = 0$ so $t = 1$ or $t = 3$. $t = 1$ does not satisfy the other two equations, but $t = 3$ does. The particles collide when $t = 3$, at the point $(9, 9, 9)$.

50. The particles collide provided $\mathbf{r}_1(t) = \mathbf{r}_2(t) \ \Leftrightarrow\ \langle t, t^2, t^3\rangle = \langle 1 + 2t, 1 + 6t, 1 + 14t\rangle$. Equating components gives $t = 1 + 2t$, $t^2 = 1 + 6t$, and $t^3 = 1 + 14t$. The first equation gives $t = -1$, but this does not satisfy the other equations, so the particles do not collide. For the paths to intersect, we need to find a value for t and a value for s where $\mathbf{r}_1(t) = \mathbf{r}_2(s) \ \Leftrightarrow$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\langle t, t^2, t^3 \rangle = \langle 1+2s, 1+6s, 1+14s \rangle$. Equating components, $t = 1+2s$, $t^2 = 1+6s$, and $t^3 = 1+14s$. Substituting the first equation into the second gives $(1+2s)^2 = 1+6s \;\Rightarrow\; 4s^2 - 2s = 0 \;\Rightarrow\; 2s(2s-1) = 0 \;\Rightarrow\; s = 0$ or $s = \frac{1}{2}$. From the first equation, $s = 0 \;\Rightarrow\; t = 1$ and $s = \frac{1}{2} \;\Rightarrow\; t = 2$. Checking, we see that both pairs of values satisfy the third equation. Thus the paths intersect twice, at the point $(1, 1, 1)$ when $s = 0$ and $t = 1$, and at $(2, 4, 8)$ when $s = \frac{1}{2}$ and $t = 2$.

51. (a) We plot the parametric equations for $0 \le t \le 2\pi$ in the first figure. We get a better idea of the shape of the curve if we plot it simultaneously with the hyperboloid of one sheet from part (b), as shown in the second figure.





(b) Here $x = \frac{27}{26}\sin 8t - \frac{8}{39}\sin 18t$, $y = -\frac{27}{26}\cos 8t + \frac{8}{39}\cos 18t$, $z = \frac{144}{65}\sin 5t$.

For any point on the curve,

$$\begin{aligned} x^2 + y^2 &= \left(\tfrac{27}{26}\sin 8t - \tfrac{8}{39}\sin 18t\right)^2 + \left(-\tfrac{27}{26}\cos 8t + \tfrac{8}{39}\cos 18t\right)^2 \\ &= \tfrac{27^2}{26^2}\sin^2 8t - 2\cdot\tfrac{27\cdot 8}{26\cdot 39}\sin 8t \sin 18t + \tfrac{64}{39^2}\sin^2 18t \\ &\quad + \tfrac{27^2}{26^2}\cos^2 8t - 2\cdot\tfrac{27\cdot 8}{26\cdot 39}\cos 8t\cos 18t + \tfrac{64}{39^2}\cos^2 18t \\ &= \tfrac{27^2}{26^2}\left(\sin^2 8t + \cos^2 8t\right) + \tfrac{64}{39^2}\left(\sin^2 18t + \cos^2 18t\right) - \tfrac{72}{169}\left(\sin 8t \sin 18t + \cos 8t \cos 18t\right) \\ &= \tfrac{27^2}{26^2} + \tfrac{64}{39^2} - \tfrac{72}{169}\cos\left(18t - 8t\right) = \tfrac{27^2}{26^2} + \tfrac{64}{39^2} - \tfrac{72}{169}\cos 10t \end{aligned}$$

using the trigonometric identities $\sin^2\theta + \cos^2\theta = 1$ and $\cos(x-y) = \cos x\cos y + \sin x \sin y$. Also $z^2 = \frac{144^2}{65^2}\sin^2 5t$, and the identity $\sin^2 x = \dfrac{1-\cos 2x}{2}$ gives $z^2 = \frac{144^2}{65^2}\cdot\frac{1}{2}\left[1 - \cos(2\cdot 5t)\right] = \frac{144^2}{2\cdot 65^2} - \frac{144^2}{2\cdot 65^2}\cos 10t$.

Then

$$\begin{aligned} 144(x^2+y^2) - 25z^2 &= 144\left(\tfrac{27^2}{26^2} + \tfrac{64}{39^2} - \tfrac{72}{169}\cos 10t\right) - 25\left(\tfrac{144^2}{2\cdot 65^2} - \tfrac{144^2}{2\cdot 65^2}\cos 10t\right) \\ &= 144\left(\tfrac{27^2}{26^2} + \tfrac{64}{39^2} - \tfrac{25\cdot 144}{2\cdot 65^2} - \tfrac{72}{169}\cos 10t + \tfrac{25\cdot 144}{2\cdot 65^2}\cos 10t\right) \\ &= 144\left(\tfrac{27^2}{26^2} + \tfrac{64}{39^2} - \tfrac{72}{169} - \tfrac{72}{169}\cos 10t + \tfrac{72}{169}\cos 10t\right) = 144\left(\tfrac{25}{36}\right) = 100 \end{aligned}$$

Thus the curve lies on the surface $144(x^2+y^2) - 25z^2 = 100$ or $144x^2 + 144y^2 - 25z^2 = 100$, a hyperboloid of one sheet with axis the z-axis.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

52. The projection of the curve onto the xy-plane is given by the parametric equations $x = (2 + \cos 1.5t)\cos t$, $y = (2 + \cos 1.5t)\sin t$. If we convert to polar coordinates, we have

$r^2 = x^2 + y^2 = [(2 + \cos 1.5t)\cos t]^2 + [(2 + \cos 1.5t)\sin t]^2 = (2 + \cos 1.5t)^2(\cos^2 t + \sin^2 t) = (2 + \cos 1.5t)^2 \quad \Rightarrow$

$r = 2 + \cos 1.5t$. Also, $\tan\theta = \dfrac{y}{x} = \dfrac{(2 + \cos 1.5t)\sin t}{(2 + \cos 1.5t)\cos t} = \tan t \quad \Rightarrow \quad \theta = t$.

Thus the polar equation of the curve is $r = 2 + \cos 1.5\theta$. At $\theta = 0$, we have $r = 3$, and r decreases to 1 as θ increases to $\frac{2\pi}{3}$. For $\frac{2\pi}{3} \le \theta \le \frac{4\pi}{3}$, r increases to 3; r decreases to 1 again at $\theta = 2\pi$, increases to 3 at $\theta = \frac{8\pi}{3}$, decreases to 1 at $\theta = \frac{10\pi}{3}$, and completes the closed curve by increasing to 3 at $\theta = 4\pi$. We sketch an approximate graph as shown in the figure.



We can determine how the curve passes over itself by investigating the maximum and minimum values of z for $0 \le t \le 4\pi$. Since $z = \sin 1.5t$, z is maximized where $\sin 1.5t = 1 \quad \Rightarrow \quad 1.5t = \frac{\pi}{2}, \frac{5\pi}{2}$, or $\frac{9\pi}{2} \quad \Rightarrow$ $t = \frac{\pi}{3}, \frac{5\pi}{3}$, or 3π. z is minimized where $\sin 1.5t = -1 \quad \Rightarrow$ $1.5t = \frac{3\pi}{2}, \frac{7\pi}{2}$, or $\frac{11\pi}{2} \quad \Rightarrow \quad t = \pi, \frac{7\pi}{3}$, or $\frac{11\pi}{3}$. Note that these are precisely the values for which $\cos 1.5t = 0 \quad \Rightarrow \quad r = 2$, and on the graph of the projection, these six points appear to be at the three self-intersections we see. Comparing the maximum and minimum values of z at these intersections, we can determine where the curve passes over itself, as indicated in the figure.



We show a computer-drawn graph of the curve from above, as well as views from the front and from the right side.



Top view



Front view



Side view

The top view graph shows a more accurate representation of the projection of the trefoil knot onto the xy-plane (the axes are rotated 90°). Notice the indentations the graph exhibits at the points corresponding to $r = 1$. Finally, we graph several

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

additional viewpoints of the trefoil knot, along with two plots showing a tube of radius 0.2 around the curve.



53. Let $\mathbf{u}(t) = \langle u_1(t), u_2(t), u_3(t) \rangle$ and $\mathbf{v}(t) = \langle v_1(t), v_2(t), v_3(t) \rangle$. In each part of this problem the basic procedure is to use Equation 1 and then analyze the individual component functions using the limit properties we have already developed for real-valued functions.

(a) $\lim_{t \to a} \mathbf{u}(t) + \lim_{t \to a} \mathbf{v}(t) = \left\langle \lim_{t \to a} u_1(t), \lim_{t \to a} u_2(t), \lim_{t \to a} u_3(t) \right\rangle + \left\langle \lim_{t \to a} v_1(t), \lim_{t \to a} v_2(t), \lim_{t \to a} v_3(t) \right\rangle$ and the limits of these component functions must each exist since the vector functions both possess limits as $t \to a$. Then adding the two vectors and using the addition property of limits for real-valued functions, we have that

$$\begin{aligned} \lim_{t \to a} \mathbf{u}(t) + \lim_{t \to a} \mathbf{v}(t) &= \left\langle \lim_{t \to a} u_1(t) + \lim_{t \to a} v_1(t), \lim_{t \to a} u_2(t) + \lim_{t \to a} v_2(t), \lim_{t \to a} u_3(t) + \lim_{t \to a} v_3(t) \right\rangle \\ &= \left\langle \lim_{t \to a} [u_1(t) + v_1(t)], \lim_{t \to a} [u_2(t) + v_2(t)], \lim_{t \to a} [u_3(t) + v_3(t)] \right\rangle \\ &= \lim_{t \to a} \langle u_1(t) + v_1(t), u_2(t) + v_2(t), u_3(t) + v_3(t) \rangle \qquad \text{[using (1) backward]} \\ &= \lim_{t \to a} [\mathbf{u}(t) + \mathbf{v}(t)] \end{aligned}$$

(b) $\lim_{t \to a} c\mathbf{u}(t) = \lim_{t \to a} \langle cu_1(t), cu_2(t), cu_3(t) \rangle = \left\langle \lim_{t \to a} cu_1(t), \lim_{t \to a} cu_2(t), \lim_{t \to a} cu_3(t) \right\rangle$

$$= \left\langle c \lim_{t \to a} u_1(t), c \lim_{t \to a} u_2(t), c \lim_{t \to a} u_3(t) \right\rangle = c \left\langle \lim_{t \to a} u_1(t), \lim_{t \to a} u_2(t), \lim_{t \to a} u_3(t) \right\rangle$$

$$= c \lim_{t \to a} \langle u_1(t), u_2(t), u_3(t) \rangle = c \lim_{t \to a} \mathbf{u}(t)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c) $\lim_{t\to a}\mathbf{u}(t)\cdot\lim_{t\to a}\mathbf{v}(t)=\left\langle\lim_{t\to a}u_1(t),\lim_{t\to a}u_2(t),\lim_{t\to a}u_3(t)\right\rangle\cdot\left\langle\lim_{t\to a}v_1(t),\lim_{t\to a}v_2(t),\lim_{t\to a}v_3(t)\right\rangle$

$$=\left[\lim_{t\to a}u_1(t)\right]\left[\lim_{t\to a}v_1(t)\right]+\left[\lim_{t\to a}u_2(t)\right]\left[\lim_{t\to a}v_2(t)\right]+\left[\lim_{t\to a}u_3(t)\right]\left[\lim_{t\to a}v_3(t)\right]$$

$$=\lim_{t\to a}u_1(t)v_1(t)+\lim_{t\to a}u_2(t)v_2(t)+\lim_{t\to a}u_3(t)v_3(t)$$

$$=\lim_{t\to a}\left[u_1(t)v_1(t)+u_2(t)v_2(t)+u_3(t)v_3(t)\right]=\lim_{t\to a}\left[\mathbf{u}(t)\cdot\mathbf{v}(t)\right]$$

(d) $\lim_{t\to a}\mathbf{u}(t)\times\lim_{t\to a}\mathbf{v}(t)=\left\langle\lim_{t\to a}u_1(t),\lim_{t\to a}u_2(t),\lim_{t\to a}u_3(t)\right\rangle\times\left\langle\lim_{t\to a}v_1(t),\lim_{t\to a}v_2(t),\lim_{t\to a}v_3(t)\right\rangle$

$$=\Big\langle\left[\lim_{t\to a}u_2(t)\right]\left[\lim_{t\to a}v_3(t)\right]-\left[\lim_{t\to a}u_3(t)\right]\left[\lim_{t\to a}v_2(t)\right],$$
$$\left[\lim_{t\to a}u_3(t)\right]\left[\lim_{t\to a}v_1(t)\right]-\left[\lim_{t\to a}u_1(t)\right]\left[\lim_{t\to a}v_3(t)\right],$$
$$\left[\lim_{t\to a}u_1(t)\right]\left[\lim_{t\to a}v_2(t)\right]-\left[\lim_{t\to a}u_2(t)\right]\left[\lim_{t\to a}v_1(t)\right]\Big\rangle$$

$$=\Big\langle\lim_{t\to a}\left[u_2(t)v_3(t)-u_3(t)v_2(t)\right],\lim_{t\to a}\left[u_3(t)v_1(t)-u_1(t)v_3(t)\right],$$
$$\lim_{t\to a}\left[u_1(t)v_2(t)-u_2(t)v_1(t)\right]\Big\rangle$$

$$=\lim_{t\to a}\left\langle u_2(t)v_3(t)-u_3(t)v_2(t),u_3(t)v_1(t)-u_1(t)v_3(t),u_1(t)v_2(t)-u_2(t)v_1(t)\right\rangle$$

$$=\lim_{t\to a}\left[\mathbf{u}(t)\times\mathbf{v}(t)\right]$$

54. Let $\mathbf{r}(t)=\langle f(t),g(t),h(t)\rangle$ and $\mathbf{b}=\langle b_1,b_2,b_3\rangle$. If $\lim_{t\to a}\mathbf{r}(t)=\mathbf{b}$, then $\lim_{t\to a}\mathbf{r}(t)$ exists, so by (1), $\mathbf{b}=\lim_{t\to a}\mathbf{r}(t)=\left\langle\lim_{t\to a}f(t),\lim_{t\to a}g(t),\lim_{t\to a}h(t)\right\rangle$. By the definition of equal vectors we have $\lim_{t\to a}f(t)=b_1$, $\lim_{t\to a}g(t)=b_2$ and $\lim_{t\to a}h(t)=b_3$. But these are limits of real-valued functions, so by the definition of limits, for every $\varepsilon>0$ there exists $\delta_1>0$, $\delta_2>0$, $\delta_3>0$ so that if $0<|t-a|<\delta_1$ then $|f(t)-b_1|<\varepsilon/3$, if $0<|t-a|<\delta_2$ then $|g(t)-b_2|<\varepsilon/3$, and if $0<|t-a|<\delta_3$ then $|h(t)-b_3|<\varepsilon/3$. Letting $\delta=$ minimum of $\{\delta_1,\delta_2,\delta_3\}$, then if $0<|t-a|<\delta$ we have $|f(t)-b_1|+|g(t)-b_2|+|h(t)-b_3|<\varepsilon/3+\varepsilon/3+\varepsilon/3=\varepsilon$. But

$$|\mathbf{r}(t)-\mathbf{b}|=|\langle f(t)-b_1,g(t)-b_2,h(t)-b_3\rangle|=\sqrt{(f(t)-b_1)^2+(g(t)-b_2)^2+(h(t)-b_3)^2}$$
$$\le\sqrt{[f(t)-b_1]^2}+\sqrt{[g(t)-b_2]^2}+\sqrt{[h(t)-b_3]^2}=|f(t)-b_1|+|g(t)-b_2|+|h(t)-b_3|$$

Thus for every $\varepsilon>0$ there exists $\delta>0$ such that if $0<|t-a|<\delta$ then $|\mathbf{r}(t)-\mathbf{b}|\le|f(t)-b_1|+|g(t)-b_2|+|h(t)-b_3|<\varepsilon$. Conversely, suppose for every $\varepsilon>0$, there exists $\delta>0$ such that if $0<|t-a|<\delta$ then $|\mathbf{r}(t)-\mathbf{b}|<\varepsilon\ \Leftrightarrow\ |\langle f(t)-b_1,g(t)-b_2,h(t)-b_3\rangle|<\varepsilon\ \Leftrightarrow$ $\sqrt{[f(t)-b_1]^2+[g(t)-b_2]^2+[h(t)-b_3]^2}<\varepsilon\ \Leftrightarrow\ [f(t)-b_1]^2+[g(t)-b_2]^2+[h(t)-b_3]^2<\varepsilon^2$. But each term on the left side of the last inequality is positive, so if $0<|t-a|<\delta$, then $[f(t)-b_1]^2<\varepsilon^2$, $[g(t)-b_2]^2<\varepsilon^2$ and $[h(t)-b_3]^2<\varepsilon^2$ or, taking the square root of both sides in each of the above, $|f(t)-b_1|<\varepsilon$, $|g(t)-b_2|<\varepsilon$ and $|h(t)-b_3|<\varepsilon$. And by definition of limits of real-valued functions we have $\lim_{t\to a}f(t)=b_1$, $\lim_{t\to a}g(t)=b_2$ and $\lim_{t\to a}h(t)=b_3$. But by (1), $\lim_{t\to a}\mathbf{r}(t)=\left\langle\lim_{t\to a}f(t),\lim_{t\to a}g(t),\lim_{t\to a}h(t)\right\rangle$, so $\lim_{t\to a}\mathbf{r}(t)=\langle b_1,b_2,b_3\rangle=\mathbf{b}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13.2 Derivatives and Integrals of Vector Functions

1. (a)



(b) $\dfrac{\mathbf{r}(4.5)-\mathbf{r}(4)}{0.5} = 2[\mathbf{r}(4.5)-\mathbf{r}(4)]$, so we draw a vector in the same direction but with twice the length of the vector $\mathbf{r}(4.5)-\mathbf{r}(4)$.

$\dfrac{\mathbf{r}(4.2)-\mathbf{r}(4)}{0.2} = 5[\mathbf{r}(4.2)-\mathbf{r}(4)]$, so we draw a vector in the same direction but with 5 times the length of the vector $\mathbf{r}(4.2)-\mathbf{r}(4)$.



(c) By Definition 1, $\mathbf{r}'(4) = \lim_{h\to 0} \dfrac{\mathbf{r}(4+h)-\mathbf{r}(4)}{h}$. $\mathbf{T}(4) = \dfrac{\mathbf{r}'(4)}{|\mathbf{r}'(4)|}$.

(d) $\mathbf{T}(4)$ is a unit vector in the same direction as $\mathbf{r}'(4)$, that is, parallel to the tangent line to the curve at $\mathbf{r}(4)$ with length 1.



2. (a) The curve can be represented by the parametric equations $x = t^2$, $y = t$, $0 \le t \le 2$. Eliminating the parameter, we have $x = y^2$, $0 \le y \le 2$, a portion of which we graph here, along with the vectors $\mathbf{r}(1)$, $\mathbf{r}(1.1)$, and $\mathbf{r}(1.1)-\mathbf{r}(1)$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) Since $\mathbf{r}(t) = \langle t^2, t \rangle$, we differentiate components, giving $\mathbf{r}'(t) = \langle 2t, 1 \rangle$, so $\mathbf{r}'(1) = \langle 2, 1 \rangle$.

$$\frac{\mathbf{r}(1.1) - \mathbf{r}(1)}{0.1} = \frac{\langle 1.21, 1.1 \rangle - \langle 1, 1 \rangle}{0.1} = 10 \langle 0.21, 0.1 \rangle = \langle 2.1, 1 \rangle.$$



As we can see from the graph, these vectors are very close in length and direction. $\mathbf{r}'(1)$ is defined to be $\lim_{h \to 0} \dfrac{\mathbf{r}(1+h) - \mathbf{r}(1)}{h}$, and we recognize $\dfrac{\mathbf{r}(1.1) - \mathbf{r}(1)}{0.1}$ as the expression after the limit sign with $h = 0.1$. Since h is close to 0, we would expect $\dfrac{\mathbf{r}(1.1) - \mathbf{r}(1)}{0.1}$ to be a vector close to $\mathbf{r}'(1)$.

3. $\mathbf{r}(t) = \langle t - 2, t^2 + 1 \rangle$, $\mathbf{r}(-1) = \langle -3, 2 \rangle$. Since $(x+2)^2 = t^2 = y - 1 \Rightarrow y = (x+2)^2 + 1$, the curve is a parabola.

(a), (c)



(b) $\mathbf{r}'(t) = \langle 1, 2t \rangle$, $\mathbf{r}'(-1) = \langle 1, -2 \rangle$

4. $\mathbf{r}(t) = \langle t^2, t^3 \rangle$, $\mathbf{r}(1) = \langle 1, 1 \rangle$. Since $x = t^2 = (t^3)^{2/3} = y^{2/3}$, the curve is the graph of $x = y^{2/3}$.

(a), (c)

(b) $\mathbf{r}'(t) = \langle 2t, 3t^2 \rangle$, $\mathbf{r}'(1) = \langle 2, 3 \rangle$

5. $\mathbf{r}(t) = e^{2t}\,\mathbf{i} + e^t\,\mathbf{j}$, $\mathbf{r}(0) = \mathbf{i} + \mathbf{j}$. Since $x = e^{2t} = (e^t)^2 = y^2$, the curve is part of a parabola. Note that here $x > 0$, $y > 0$.

(a), (c)



(b) $\mathbf{r}'(t) = 2e^{2t}\,\mathbf{i} + e^t\,\mathbf{j}$, $\mathbf{r}'(0) = 2\,\mathbf{i} + \mathbf{j}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $\mathbf{r}(t) = e^t\,\mathbf{i} + 2t\,\mathbf{j},\ \mathbf{r}(0) = \mathbf{i}$.

Since $x = e^t \Leftrightarrow t = \ln x$ and $y = 2t = 2\ln x$, the curve is the graph of $y = 2\ln x$.

(a), (c)



(b) $\mathbf{r}'(t) = e^t\,\mathbf{i} + 2\,\mathbf{j}$, $\mathbf{r}'(0) = \mathbf{i} + 2\,\mathbf{j}$

7. $\mathbf{r}(t) = 4\sin t\,\mathbf{i} - 2\cos t\,\mathbf{j},\ \mathbf{r}(3\pi/4) = 4(\sqrt{2}/2)\,\mathbf{i} - 2(-\sqrt{2}/2)\,\mathbf{j} = 2\sqrt{2}\,\mathbf{i} + \sqrt{2}\,\mathbf{j}$.

Here $(x/4)^2 + (y/2)^2 = \sin^2 t + \cos^2 t = 1$, so the curve is the ellipse $\dfrac{x^2}{16} + \dfrac{y^2}{4} = 1$.

(a), (c)



(b) $\mathbf{r}'(t) = 4\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j}$, $\mathbf{r}'(3\pi/4) = -2\sqrt{2}\,\mathbf{i} + \sqrt{2}\,\mathbf{j}$.

8. $\mathbf{r}(t) = (\cos t + 1)\,\mathbf{i} + (\sin t - 1)\,\mathbf{j},\ \mathbf{r}(-\pi/3) = \left(\frac{1}{2} + 1\right)\mathbf{i} + \left(-\frac{\sqrt{3}}{2} - 1\right)\mathbf{j} = \frac{3}{2}\,\mathbf{i} + \left(-\frac{\sqrt{3}}{2} - 1\right)\mathbf{j} \approx 1.5\,\mathbf{i} - 1.87\,\mathbf{j}$.

Here $(x-1)^2 + (y+1)^2 = \cos^2 t + \sin^2 t = 1$, so the curve is a circle of radius 1 with center $(1, -1)$.

(a), (c)



(b) $\mathbf{r}'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j}$, $\mathbf{r}'(-\pi/3) = \frac{\sqrt{3}}{2}\,\mathbf{i} + \frac{1}{2}\,\mathbf{j} \approx 0.87\,\mathbf{i} + 0.5\,\mathbf{j}$

9. $\mathbf{r}(t) = \left\langle \sqrt{t-2},\ 3,\ 1/t^2 \right\rangle \Rightarrow$

$$\mathbf{r}'(t) = \left\langle \frac{d}{dt}\left[\sqrt{t-2}\right], \frac{d}{dt}\left[3\right], \frac{d}{dt}\left[1/t^2\right] \right\rangle = \left\langle \tfrac{1}{2}(t-2)^{-1/2}, 0, -2t^{-3} \right\rangle = \left\langle \frac{1}{2\sqrt{t-2}}, 0, -\frac{2}{t^3} \right\rangle$$

10. $\mathbf{r}(t) = \left\langle e^{-t},\ t - t^3,\ \ln t \right\rangle \Rightarrow \mathbf{r}'(t) = \left\langle -e^{-t},\ 1 - 3t^2,\ 1/t \right\rangle$

11. $\mathbf{r}(t) = t^2\,\mathbf{i} + \cos\left(t^2\right)\,\mathbf{j} + \sin^2 t\,\mathbf{k} \Rightarrow$

$$\mathbf{r}'(t) = 2t\,\mathbf{i} + \left[-\sin(t^2)\cdot 2t\right]\mathbf{j} + (2\sin t\cdot\cos t)\,\mathbf{k} = 2t\,\mathbf{i} - 2t\sin(t^2)\,\mathbf{j} + 2\sin t\cos t\,\mathbf{k}$$

12. $\mathbf{r}(t) = \dfrac{1}{1+t}\,\mathbf{i} + \dfrac{t}{1+t}\,\mathbf{j} + \dfrac{t^2}{1+t}\,\mathbf{k} \Rightarrow$

$$\mathbf{r}'(t) = \frac{0 - 1(1)}{(1+t)^2}\,\mathbf{i} + \frac{(1+t)\cdot 1 - t(1)}{(1+t)^2}\,\mathbf{j} + \frac{(1+t)\cdot 2t - t^2(1)}{(1+t)^2}\,\mathbf{k} = -\frac{1}{(1+t)^2}\,\mathbf{i} + \frac{1}{(1+t)^2}\,\mathbf{j} + \frac{t^2+2t}{(1+t)^2}\,\mathbf{k}$$

13. $\mathbf{r}(t) = t\sin t\,\mathbf{i} + e^t\cos t\,\mathbf{j} + \sin t\cos t\,\mathbf{k} \Rightarrow$

$$\begin{aligned}\mathbf{r}'(t) &= \left[t\cdot\cos t + (\sin t)\cdot 1\right]\mathbf{i} + \left[e^t(-\sin t) + (\cos t)e^t\right]\mathbf{j} + \left[(\sin t)(-\sin t) + (\cos t)(\cos t)\right]\mathbf{k} \\ &= (t\cos t + \sin t)\,\mathbf{i} + e^t\,(\cos t - \sin t)\,\mathbf{j} + \left(\cos^2 t - \sin^2 t\right)\mathbf{k}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

14. $\mathbf{r}(t) = \sin^2 at\,\mathbf{i} + te^{bt}\,\mathbf{j} + \cos^2 ct\,\mathbf{k} \;\Rightarrow$

$\mathbf{r}'(t) = [2(\sin at)\cdot(\cos at)(a)]\,\mathbf{i} + \left[t\cdot e^{bt}(b) + e^{bt}\cdot 1\right]\mathbf{j} + [2(\cos ct)\cdot(-\sin ct)(c)]\,\mathbf{k}$

$= 2a\sin at\cos at\,\mathbf{i} + e^{bt}\,(bt+1)\,\mathbf{j} - 2c\sin ct\cos ct\,\mathbf{k}$

15. $\mathbf{r}'(t) = \mathbf{0} + \mathbf{b} + 2t\,\mathbf{c} = \mathbf{b} + 2t\,\mathbf{c}$ by Formulas 1 and 3 of Theorem 3.

16. To find $\mathbf{r}'(t)$, we first expand $\mathbf{r}(t) = t\,\mathbf{a}\times(\mathbf{b}+t\,\mathbf{c}) = t(\mathbf{a}\times\mathbf{b}) + t^2(\mathbf{a}\times\mathbf{c})$, so $\mathbf{r}'(t) = \mathbf{a}\times\mathbf{b} + 2t(\mathbf{a}\times\mathbf{c})$.

17. $\mathbf{r}(t) = \langle t^2 - 2t, 1+3t, \frac{1}{3}t^3 + \frac{1}{2}t^2\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 2t-2, 3, t^2+t\rangle \;\Rightarrow\; \mathbf{r}'(2) = \langle 2,3,6\rangle$.

So $|\mathbf{r}'(2)| = \sqrt{2^2+3^2+6^2} = \sqrt{49} = 7$ and $\mathbf{T}(2) = \dfrac{\mathbf{r}'(2)}{|\mathbf{r}'(2)|} = \frac{1}{7}\langle 2,3,6\rangle = \left\langle \frac{2}{7}, \frac{3}{7}, \frac{6}{7}\right\rangle$.

18. $\mathbf{r}(t) = \langle \tan^{-1} t, 2e^{2t}, 8te^t\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1/(1+t^2), 4e^{2t}, 8te^t + 8e^t\rangle \;\Rightarrow\; \mathbf{r}'(0) = \langle 1,4,8\rangle$.

So $|\mathbf{r}'(0)| = \sqrt{1^2+4^2+8^2} = \sqrt{81} = 9$ and $\mathbf{T}(0) = \dfrac{\mathbf{r}'(0)}{|\mathbf{r}'(0)|} = \frac{1}{9}\langle 1,4,8\rangle = \left\langle \frac{1}{9}, \frac{4}{9}, \frac{8}{9}\right\rangle$.

19. $\mathbf{r}'(t) = -\sin t\,\mathbf{i} + 3\,\mathbf{j} + 4\cos 2t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(0) = 3\,\mathbf{j} + 4\,\mathbf{k}$. Thus

$\mathbf{T}(0) = \dfrac{\mathbf{r}'(0)}{|\mathbf{r}'(0)|} = \dfrac{1}{\sqrt{0^2+3^2+4^2}}\,(3\,\mathbf{j}+4\,\mathbf{k}) = \frac{1}{5}(3\,\mathbf{j}+4\,\mathbf{k}) = \frac{3}{5}\,\mathbf{j} + \frac{4}{5}\,\mathbf{k}$.

20. $\mathbf{r}'(t) = 2\sin t\cos t\,\mathbf{i} - 2\cos t\sin t\,\mathbf{j} + 2\tan t\sec^2 t\,\mathbf{k} \;\Rightarrow$

$\mathbf{r}'\left(\frac{\pi}{4}\right) = 2\cdot\frac{\sqrt{2}}{2}\cdot\frac{\sqrt{2}}{2}\,\mathbf{i} - 2\cdot\frac{\sqrt{2}}{2}\cdot\frac{\sqrt{2}}{2}\,\mathbf{j} + 2\cdot 1\cdot(\sqrt{2})^2\,\mathbf{k} = \mathbf{i} - \mathbf{j} + 4\,\mathbf{k}$ and $\left|\mathbf{r}'\left(\frac{\pi}{4}\right)\right| = \sqrt{1+1+16} = \sqrt{18} = 3\sqrt{2}$. Thus

$\mathbf{T}\left(\frac{\pi}{4}\right) = \dfrac{\mathbf{r}'\left(\frac{\pi}{4}\right)}{\left|\mathbf{r}'\left(\frac{\pi}{4}\right)\right|} = \dfrac{1}{3\sqrt{2}}\,(\mathbf{i}-\mathbf{j}+4\,\mathbf{k}) = \dfrac{1}{3\sqrt{2}}\,\mathbf{i} - \dfrac{1}{3\sqrt{2}}\,\mathbf{j} + \dfrac{4}{3\sqrt{2}}\,\mathbf{k}$.

21. $\mathbf{r}(t) = \langle t, t^2, t^3\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, 2t, 3t^2\rangle$. Then $\mathbf{r}'(1) = \langle 1,2,3\rangle$ and $|\mathbf{r}'(1)| = \sqrt{1^2+2^2+3^2} = \sqrt{14}$, so

$\mathbf{T}(1) = \dfrac{\mathbf{r}'(1)}{|\mathbf{r}'(1)|} = \frac{1}{\sqrt{14}}\langle 1,2,3\rangle = \left\langle \frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}\right\rangle$. $\mathbf{r}''(t) = \langle 0, 2, 6t\rangle$, so

$$\mathbf{r}'(t)\times\mathbf{r}''(t) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2t & 3t^2 \\ 0 & 2 & 6t \end{vmatrix} = \begin{vmatrix} 2t & 3t^2 \\ 2 & 6t \end{vmatrix}\mathbf{i} - \begin{vmatrix} 1 & 3t^2 \\ 0 & 6t \end{vmatrix}\mathbf{j} + \begin{vmatrix} 1 & 2t \\ 0 & 2 \end{vmatrix}\mathbf{k}$$

$$= (12t^2 - 6t^2)\,\mathbf{i} - (6t-0)\,\mathbf{j} + (2-0)\,\mathbf{k} = \langle 6t^2, -6t, 2\rangle$$

22. $\mathbf{r}(t) = \langle e^{2t}, e^{-2t}, te^{2t}\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 2e^{2t}, -2e^{-2t}, (2t+1)e^{2t}\rangle \;\Rightarrow\; \mathbf{r}'(0) = \langle 2e^0, -2e^0, (0+1)e^0\rangle = \langle 2,-2,1\rangle$

and $|\mathbf{r}'(0)| = \sqrt{2^2+(-2)^2+1^2} = 3$. Then $\mathbf{T}(0) = \dfrac{\mathbf{r}'(0)}{|\mathbf{r}'(0)|} = \frac{1}{3}\langle 2,-2,1\rangle = \left\langle \frac{2}{3}, -\frac{2}{3}, \frac{1}{3}\right\rangle$.

$\mathbf{r}''(t) = \langle 4e^{2t}, 4e^{-2t}, (4t+4)e^{2t}\rangle \;\Rightarrow\; \mathbf{r}''(0) = \langle 4e^0, 4e^0, (0+4)e^0\rangle = \langle 4,4,4\rangle$.

$\mathbf{r}'(t)\cdot\mathbf{r}''(t) = \langle 2e^{2t}, -2e^{-2t}, (2t+1)e^{2t}\rangle\cdot\langle 4e^{2t}, 4e^{-2t}, (4t+4)e^{2t}\rangle$

$= (2e^{2t})(4e^{2t}) + (-2e^{-2t})(4e^{-2t}) + ((2t+1)e^{2t})((4t+4)e^{2t})$

$= 8e^{4t} - 8e^{-4t} + (8t^2+12t+4)e^{4t} = (8t^2+12t+12)e^{4t} - 8e^{-4t}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. The vector equation for the curve is $\mathbf{r}(t) = \left\langle t^2+1, 4\sqrt{t}, e^{t^2-t} \right\rangle$, so $\mathbf{r}'(t) = \left\langle 2t, 2/\sqrt{t}, (2t-1)e^{t^2-t} \right\rangle$. The point $(2,4,1)$ corresponds to $t=1$, so the tangent vector there is $\mathbf{r}'(1) = \langle 2,2,1 \rangle$. Thus, the tangent line goes through the point $(2,4,1)$ and is parallel to the vector $\langle 2,2,1 \rangle$. Parametric equations are $x = 2+2t$, $y = 4+2t$, $z = 1+t$.

24. The vector equation for the curve is $\mathbf{r}(t) = \langle \ln(t+1), t\cos 2t, 2^t \rangle$, so $\mathbf{r}'(t) = \langle 1/(t+1), \cos 2t - 2t\sin 2t, 2^t \ln 2 \rangle$. The point $(0,0,1)$ corresponds to $t=0$, so the tangent vector there is $\mathbf{r}'(0) = \langle 1, 1, \ln 2 \rangle$. Thus, the tangent line goes through the point $(0,0,1)$ and is parallel to the vector $\langle 1,1,\ln 2 \rangle$. Parametric equations are $x = 0+1\cdot t = t$, $y = 0+1\cdot t = t$, $z = 1+(\ln 2)t$.

25. The vector equation for the curve is $\mathbf{r}(t) = \langle e^{-t}\cos t, e^{-t}\sin t, e^{-t} \rangle$, so

$$\begin{aligned}\mathbf{r}'(t) &= \langle e^{-t}(-\sin t) + (\cos t)(-e^{-t}),\ e^{-t}\cos t + (\sin t)(-e^{-t}),\ (-e^{-t}) \rangle \\ &= \langle -e^{-t}(\cos t + \sin t),\ e^{-t}(\cos t - \sin t),\ -e^{-t} \rangle\end{aligned}$$

The point $(1,0,1)$ corresponds to $t=0$, so the tangent vector there is $\mathbf{r}'(0) = \langle -e^0(\cos 0 + \sin 0), e^0(\cos 0 - \sin 0), -e^0 \rangle = \langle -1,1,-1 \rangle$. Thus, the tangent line is parallel to the vector $\langle -1,1,-1 \rangle$ and parametric equations are $x = 1+(-1)t = 1-t$, $y = 0+1\cdot t = t$, $z = 1+(-1)t = 1-t$.

26. The vector equation for the curve is $\mathbf{r}(t) = \langle \sqrt{t^2+3}, \ln(t^2+3), t \rangle$, so $\mathbf{r}'(t) = \langle t/\sqrt{t^2+3}, 2t/(t^2+3), 1 \rangle$. At $(2, \ln 4, 1)$, $t=1$ and $\mathbf{r}'(1) = \langle \frac{1}{2}, \frac{1}{2}, 1 \rangle$. Thus, parametric equations of the tangent line are $x = 2+\frac{1}{2}t$, $y = \ln 4 + \frac{1}{2}t$, $z = 1+t$.

27. First we parametrize the curve C of intersection. The projection of C onto the xy-plane is contained in the circle $x^2+y^2 = 25$, $z=0$, so we can write $x = 5\cos t$, $y = 5\sin t$. C also lies on the cylinder $y^2+z^2 = 20$, and $z \geq 0$ near the point $(3,4,2)$, so we can write $z = \sqrt{20-y^2} = \sqrt{20-25\sin^2 t}$. A vector equation then for C is $\mathbf{r}(t) = \left\langle 5\cos t, 5\sin t, \sqrt{20-25\sin^2 t} \right\rangle \Rightarrow \mathbf{r}'(t) = \left\langle -5\sin t, 5\cos t, \frac{1}{2}(20-25\sin^2 t)^{-1/2}(-50\sin t\cos t) \right\rangle$.

The point $(3,4,2)$ corresponds to $t = \cos^{-1}\left(\frac{3}{5}\right)$, so the tangent vector there is

$$\mathbf{r}'\left(\cos^{-1}\left(\tfrac{3}{5}\right)\right) = \left\langle -5\left(\tfrac{4}{5}\right), 5\left(\tfrac{3}{5}\right), \tfrac{1}{2}\left(20-25\left(\tfrac{4}{5}\right)^2\right)^{-1/2}\left(-50\left(\tfrac{4}{5}\right)\left(\tfrac{3}{5}\right)\right) \right\rangle = \langle -4,3,-6 \rangle.$$

The tangent line is parallel to this vector and passes through $(3,4,2)$, so a vector equation for the line is $\mathbf{r}(t) = (3-4t)\mathbf{i} + (4+3t)\mathbf{j} + (2-6t)\mathbf{k}$.

28. $\mathbf{r}(t) = \langle 2\cos t, 2\sin t, e^t \rangle \Rightarrow \mathbf{r}'(t) = \langle -2\sin t, 2\cos t, e^t \rangle$. The tangent line to the curve is parallel to the plane when the curve's tangent vector is orthogonal to the plane's normal vector. Thus we require $\langle -2\sin t, 2\cos t, e^t \rangle \cdot \langle \sqrt{3}, 1, 0 \rangle = 0 \Rightarrow$ $-2\sqrt{3}\sin t + 2\cos t + 0 = 0 \Rightarrow \tan t = \frac{1}{\sqrt{3}} \Rightarrow t = \frac{\pi}{6}$ [since $0 \leq t \leq \pi$].

$\mathbf{r}\left(\frac{\pi}{6}\right) = \left\langle \sqrt{3}, 1, e^{\pi/6} \right\rangle$, so the point is $(\sqrt{3}, 1, e^{\pi/6})$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

29. $\mathbf{r}(t) = \langle t, e^{-t}, 2t - t^2 \rangle \Rightarrow \mathbf{r}'(t) = \langle 1, -e^{-t}, 2 - 2t \rangle$. At $(0, 1, 0)$, $t = 0$ and $\mathbf{r}'(0) = \langle 1, -1, 2 \rangle$. Thus, parametric equations of the tangent line are $x = t$, $y = 1 - t$, $z = 2t$.



30. $\mathbf{r}(t) = \langle 2\cos t, 2\sin t, 4\cos 2t \rangle$,
$\mathbf{r}'(t) = \langle -2\sin t, 2\cos t, -8\sin 2t \rangle$. At $\left(\sqrt{3}, 1, 2\right)$, $t = \frac{\pi}{6}$ and $\mathbf{r}'\left(\frac{\pi}{6}\right) = \left\langle -1, \sqrt{3}, -4\sqrt{3} \right\rangle$. Thus, parametric equations of the tangent line are $x = \sqrt{3} - t$, $y = 1 + \sqrt{3}\,t$, $z = 2 - 4\sqrt{3}\,t$.



31. $\mathbf{r}(t) = \langle t\cos t, t, t\sin t \rangle \Rightarrow \mathbf{r}'(t) = \langle \cos t - t\sin t, 1, t\cos t + \sin t \rangle$.
At $(-\pi, \pi, 0)$, $t = \pi$ and $\mathbf{r}'(\pi) = \langle -1, 1, -\pi \rangle$. Thus, parametric equations of the tangent line are $x = -\pi - t$, $y = \pi + t$, $z = -\pi t$.



32. (a) The tangent line at $t = 0$ is the line through the point with position vector $\mathbf{r}(0) = \langle \sin 0, 2\sin 0, \cos 0 \rangle = \langle 0, 0, 1 \rangle$, and in the direction of the tangent vector, $\mathbf{r}'(0) = \langle \pi\cos 0, 2\pi\cos 0, -\pi\sin 0 \rangle = \langle \pi, 2\pi, 0 \rangle$. So an equation of the line is

$$\langle x, y, z \rangle = \mathbf{r}(0) + u\,\mathbf{r}'(0) = \langle 0 + \pi u, 0 + 2\pi u, 1 \rangle = \langle \pi u, 2\pi u, 1 \rangle.$$

$\mathbf{r}\left(\frac{1}{2}\right) = \left\langle \sin\frac{\pi}{2}, 2\sin\frac{\pi}{2}, \cos\frac{\pi}{2} \right\rangle = \langle 1, 2, 0 \rangle$,

$\mathbf{r}'\left(\frac{1}{2}\right) = \left\langle \pi\cos\frac{\pi}{2}, 2\pi\cos\frac{\pi}{2}, -\pi\sin\frac{\pi}{2} \right\rangle = \langle 0, 0, -\pi \rangle$.

So the equation of the second line is

$\langle x, y, z \rangle = \langle 1, 2, 0 \rangle + v\,\langle 0, 0, -\pi \rangle = \langle 1, 2, -\pi v \rangle$.

The lines intersect where $\langle \pi u, 2\pi u, 1 \rangle = \langle 1, 2, -\pi v \rangle$,

so the point of intersection is $(1, 2, 1)$.

(b)



33. The angle of intersection of the two curves is the angle between the two tangent vectors to the curves at the point of intersection. Since $\mathbf{r}_1'(t) = \langle 1, 2t, 3t^2 \rangle$ and $t = 0$ at $(0, 0, 0)$, $\mathbf{r}_1'(0) = \langle 1, 0, 0 \rangle$ is a tangent vector to $\mathbf{r}_1$ at $(0, 0, 0)$. Similarly,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{r}_2'(t) = \langle \cos t, 2\cos 2t, 1\rangle$ and since $\mathbf{r}_2(0) = \langle 0,0,0\rangle$, $\mathbf{r}_2'(0) = \langle 1,2,1\rangle$ is a tangent vector to $\mathbf{r}_2$ at $(0,0,0)$. If θ is the angle between these two tangent vectors, then $\cos\theta = \frac{1}{\sqrt{1}\sqrt{6}}\langle 1,0,0\rangle \cdot \langle 1,2,1\rangle = \frac{1}{\sqrt{6}}$ and $\theta = \cos^{-1}\left(\frac{1}{\sqrt{6}}\right) \approx 66°$.

34. To find the point of intersection, we must find the values of t and s which satisfy the following three equations simultaneously: $t = 3 - s$, $1 - t = s - 2$, $3 + t^2 = s^2$. Solving the last two equations gives $t = 1$, $s = 2$ (check these in the first equation). Thus the point of intersection is $(1,0,4)$. To find the angle θ of intersection, we proceed as in Exercise 33. The tangent vectors to the respective curves at $(1,0,4)$ are $\mathbf{r}_1'(1) = \langle 1,-1,2\rangle$ and $\mathbf{r}_2'(2) = \langle -1,1,4\rangle$. So

$\cos\theta = \frac{1}{\sqrt{6}\sqrt{18}}(-1-1+8) = \frac{6}{6\sqrt{3}} = \frac{1}{\sqrt{3}}$ and $\theta = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right) \approx 55°$.

Note: In Exercise 33, the curves intersect when the value of both parameters is zero. However, as seen in this exercise, it is not necessary for the parameters to be of equal value at the point of intersection.

35. $\int_0^2 (t\,\mathbf{i} - t^3\,\mathbf{j} + 3t^5\,\mathbf{k})\,dt = \left(\int_0^2 t\,dt\right)\mathbf{i} - \left(\int_0^2 t^3\,dt\right)\mathbf{j} + \left(\int_0^2 3t^5\,dt\right)\mathbf{k}$

$= \left[\frac{1}{2}t^2\right]_0^2\mathbf{i} - \left[\frac{1}{4}t^4\right]_0^2\mathbf{j} + \left[\frac{1}{2}t^6\right]_0^2\mathbf{k}$

$= \frac{1}{2}(4-0)\,\mathbf{i} - \frac{1}{4}(16-0)\,\mathbf{j} + \frac{1}{2}(64-0)\,\mathbf{k} = 2\,\mathbf{i} - 4\,\mathbf{j} + 32\,\mathbf{k}$

36. $\int_1^4 \left(2t^{3/2}\,\mathbf{i} + (t+1)\sqrt{t}\,\mathbf{k}\right)dt = \left(\int_1^4 2t^{3/2}\,dt\right)\mathbf{i} + \left[\int_1^4 (t^{3/2} + t^{1/2})\,dt\right]\mathbf{k}$

$= \left[\frac{4}{5}t^{5/2}\right]_1^4\mathbf{i} + \left[\frac{2}{5}t^{5/2} + \frac{2}{3}t^{3/2}\right]_1^4\mathbf{k}$

$= \frac{4}{5}(4^{5/2} - 1)\,\mathbf{i} + \left(\frac{2}{5}(4)^{5/2} + \frac{2}{3}(4)^{3/2} - \frac{2}{5} - \frac{2}{3}\right)\mathbf{k}$

$= \frac{4}{5}(31)\,\mathbf{i} + \left(\frac{2}{5}(32) + \frac{2}{3}(8) - \frac{2}{5} - \frac{2}{3}\right)\mathbf{k} = \frac{124}{5}\,\mathbf{i} + \frac{256}{15}\,\mathbf{k}$

37. $\int_0^1 \left(\frac{1}{t+1}\,\mathbf{i} + \frac{1}{t^2+1}\,\mathbf{j} + \frac{t}{t^2+1}\,\mathbf{k}\right)dt = \left(\int_0^1 \frac{1}{t+1}\,dt\right)\mathbf{i} + \left(\int_0^1 \frac{1}{t^2+1}\,dt\right)\mathbf{j} + \left(\int_0^1 \frac{t}{t^2+1}\,dt\right)\mathbf{k}$

$= \left[\ln|t+1|\right]_0^1\mathbf{i} + \left[\tan^{-1}t\right]_0^1\mathbf{j} + \left[\frac{1}{2}\ln(t^2+1)\right]_0^1\mathbf{k}$

$= (\ln 2 - \ln 1)\,\mathbf{i} + (\frac{\pi}{4} - 0)\,\mathbf{j} + \frac{1}{2}(\ln 2 - \ln 1)\,\mathbf{k} = \ln 2\,\mathbf{i} + \frac{\pi}{4}\,\mathbf{j} + \frac{1}{2}\ln 2\,\mathbf{k}$

38. $\int_0^{\pi/4}(\sec t\tan t\,\mathbf{i} + t\cos 2t\,\mathbf{j} + \sin^2 2t\cos 2t\,\mathbf{k})\,dt$

$= \left(\int_0^{\pi/4}\sec t\tan t\,dt\right)\mathbf{i} + \left(\int_0^{\pi/4} t\cos 2t\,dt\right)\mathbf{j} + \left(\int_0^{\pi/4}\sin^2 2t\cos 2t\,dt\right)\mathbf{k}$

$= \left[\sec t\right]_0^{\pi/4}\mathbf{i} + \left(\left[\frac{1}{2}t\sin 2t\right]_0^{\pi/4} - \int_0^{\pi/4}\frac{1}{2}\sin 2t\,dt\right)\mathbf{j} + \left[\frac{1}{6}\sin^3 2t\right]_0^{\pi/4}\mathbf{k}$

[For the y-component, integrate by parts with $u = t$, $dv = \cos 2t\,dt$.]

$= (\sec\frac{\pi}{4} - \sec 0)\,\mathbf{i} + \left(\frac{\pi}{8}\sin\frac{\pi}{2} - 0 - \left[-\frac{1}{4}\cos 2t\right]_0^{\pi/4}\right)\mathbf{j} + \frac{1}{6}\left(\sin^3\frac{\pi}{2} - \sin^3 0\right)\mathbf{k}$

$= (\sqrt{2} - 1)\,\mathbf{i} + \left(\frac{\pi}{8} + \frac{1}{4}\cos\frac{\pi}{2} - \frac{1}{4}\cos 0\right)\mathbf{j} + \frac{1}{6}(1-0)\,\mathbf{k} = (\sqrt{2}-1)\,\mathbf{i} + \left(\frac{\pi}{8} - \frac{1}{4}\right)\mathbf{j} + \frac{1}{6}\,\mathbf{k}$

39. $\int(\sec^2 t\,\mathbf{i} + t(t^2+1)^3\,\mathbf{j} + t^2\ln t\,\mathbf{k})\,dt = \left(\int \sec^2 t\,dt\right)\mathbf{i} + \left(\int t(t^2+1)^3\,dt\right)\mathbf{j} + \left(\int t^2\ln t\,dt\right)\mathbf{k}$

$= \tan t\,\mathbf{i} + \frac{1}{8}(t^2+1)^4\,\mathbf{j} + \left(\frac{1}{3}t^3\ln t - \frac{1}{9}t^3\right)\mathbf{k} + \mathbf{C},$

where $\mathbf{C}$ is a vector constant of integration. [For the z-component, integrate by parts with $u = \ln t$, $dv = t^2\,dt$.]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

40. $\displaystyle\int \left(te^{2t}\,\mathbf{i}+\frac{t}{1-t}\,\mathbf{j}+\frac{1}{\sqrt{1-t^2}}\,\mathbf{k}\right)dt=\left(\int te^{2t}\,dt\right)\mathbf{i}+\left(\int \frac{t}{1-t}\,dt\right)\mathbf{j}+\left(\int \frac{1}{\sqrt{1-t^2}}\,dt\right)\mathbf{k}$

$$=\left(\tfrac{1}{2}te^{2t}-\int \tfrac{1}{2}e^{2t}\,dt\right)\mathbf{i}+\left[\int\left(-1+\frac{1}{1-t}\right)dt\right]\mathbf{j}+\left(\int \frac{1}{\sqrt{1-t^2}}\,dt\right)\mathbf{k}$$

$$=\left(\tfrac{1}{2}te^{2t}-\tfrac{1}{4}e^{2t}\right)\mathbf{i}+\left(-t-\ln|1-t|\right)\mathbf{j}+\sin^{-1}t\,\mathbf{k}+\mathbf{C}$$

41. $\mathbf{r}'(t)=2t\,\mathbf{i}+3t^2\,\mathbf{j}+\sqrt{t}\,\mathbf{k} \;\Rightarrow\; \mathbf{r}(t)=t^2\,\mathbf{i}+t^3\,\mathbf{j}+\frac{2}{3}t^{3/2}\,\mathbf{k}+\mathbf{C}$, where $\mathbf{C}$ is a constant vector.

But $\mathbf{i}+\mathbf{j}=\mathbf{r}(1)=\mathbf{i}+\mathbf{j}+\frac{2}{3}\mathbf{k}+\mathbf{C}$. Thus $\mathbf{C}=-\frac{2}{3}\mathbf{k}$ and $\mathbf{r}(t)=t^2\,\mathbf{i}+t^3\,\mathbf{j}+\left(\frac{2}{3}t^{3/2}-\frac{2}{3}\right)\mathbf{k}$.

42. $\mathbf{r}'(t)=t\,\mathbf{i}+e^t\,\mathbf{j}+te^t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}(t)=\frac{1}{2}t^2\,\mathbf{i}+e^t\,\mathbf{j}+\left(te^t-e^t\right)\mathbf{k}+\mathbf{C}$. But $\mathbf{i}+\mathbf{j}+\mathbf{k}=\mathbf{r}(0)=\mathbf{j}-\mathbf{k}+\mathbf{C}$.

Thus $\mathbf{C}=\mathbf{i}+2\,\mathbf{k}$ and $\mathbf{r}(t)=\left(\frac{1}{2}t^2+1\right)\mathbf{i}+e^t\,\mathbf{j}+(te^t-e^t+2)\,\mathbf{k}$.

For Exercises 43–46, let $\mathbf{u}(t)=\langle u_1(t),u_2(t),u_3(t)\rangle$ and $\mathbf{v}(t)=\langle v_1(t),v_2(t),v_3(t)\rangle$. In each of these exercises, the procedure is to apply Theorem 2 so that the corresponding properties of derivatives of real-valued functions can be used.

43. $\dfrac{d}{dt}\left[\mathbf{u}(t)+\mathbf{v}(t)\right]=\dfrac{d}{dt}\left\langle u_1(t)+v_1(t),u_2(t)+v_2(t),u_3(t)+v_3(t)\right\rangle$

$$=\left\langle \frac{d}{dt}\left[u_1(t)+v_1(t)\right],\frac{d}{dt}\left[u_2(t)+v_2(t)\right],\frac{d}{dt}\left[u_3(t)+v_3(t)\right]\right\rangle$$

$$=\langle u_1'(t)+v_1'(t),u_2'(t)+v_2'(t),u_3'(t)+v_3'(t)\rangle$$

$$=\langle u_1'(t),u_2'(t),u_3'(t)\rangle+\langle v_1'(t),v_2'(t),v_3'(t)\rangle=\mathbf{u}'(t)+\mathbf{v}'(t)$$

44. $\dfrac{d}{dt}\left[f(t)\,\mathbf{u}(t)\right]=\dfrac{d}{dt}\left\langle f(t)u_1(t),f(t)u_2(t),f(t)u_3(t)\right\rangle$

$$=\left\langle \frac{d}{dt}\left[f(t)u_1(t)\right],\frac{d}{dt}\left[f(t)u_2(t)\right],\frac{d}{dt}\left[f(t)u_3(t)\right]\right\rangle$$

$$=\langle f'(t)u_1(t)+f(t)u_1'(t),f'(t)u_2(t)+f(t)u_2'(t),f'(t)u_3(t)+f(t)u_3'(t)\rangle$$

$$=f'(t)\left\langle u_1(t),u_2(t),u_3(t)\right\rangle+f(t)\left\langle u_1'(t),u_2'(t),u_3'(t)\right\rangle=f'(t)\,\mathbf{u}(t)+f(t)\,\mathbf{u}'(t)$$

45. $\dfrac{d}{dt}\left[\mathbf{u}(t)\times\mathbf{v}(t)\right]=\dfrac{d}{dt}\left\langle u_2(t)v_3(t)-u_3(t)v_2(t),u_3(t)v_1(t)-u_1(t)v_3(t),u_1(t)v_2(t)-u_2(t)v_1(t)\right\rangle$

$$\begin{aligned}=\langle\, & u_2'v_3(t)+u_2(t)v_3'(t)-u_3'(t)v_2(t)-u_3(t)v_2'(t),\\ & u_3'(t)v_1(t)+u_3(t)v_1'(t)-u_1'(t)v_3(t)-u_1(t)v_3'(t),\\ & u_1'(t)v_2(t)+u_1(t)v_2'(t)-u_2'(t)v_1(t)-u_2(t)v_1'(t)\rangle\end{aligned}$$

$$\begin{aligned}=\,&\langle u_2'(t)v_3(t)-u_3'(t)v_2(t),u_3'(t)v_1(t)-u_1'(t)v_3(t),u_1'(t)v_2(t)-u_2'(t)v_1(t)\rangle\\ &+\langle u_2(t)v_3'(t)-u_3(t)v_2'(t),u_3(t)v_1'(t)-u_1(t)v_3'(t),u_1(t)v_2'(t)-u_2(t)v_1'(t)\rangle\end{aligned}$$

$$=\mathbf{u}'(t)\times\mathbf{v}(t)+\mathbf{u}(t)\times\mathbf{v}'(t)$$

Alternate solution: Let $\mathbf{r}(t)=\mathbf{u}(t)\times\mathbf{v}(t)$. Then

$$\begin{aligned}\mathbf{r}(t+h)-\mathbf{r}(t)&=\left[\mathbf{u}(t+h)\times\mathbf{v}(t+h)\right]-\left[\mathbf{u}(t)\times\mathbf{v}(t)\right]\\ &=\left[\mathbf{u}(t+h)\times\mathbf{v}(t+h)\right]-\left[\mathbf{u}(t)\times\mathbf{v}(t)\right]+\left[\mathbf{u}(t+h)\times\mathbf{v}(t)\right]-\left[\mathbf{u}(t+h)\times\mathbf{v}(t)\right]\\ &=\mathbf{u}(t+h)\times\left[\mathbf{v}(t+h)-\mathbf{v}(t)\right]+\left[\mathbf{u}(t+h)-\mathbf{u}(t)\right]\times\mathbf{v}(t)\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(Be careful of the order of the cross product.) Dividing through by h and taking the limit as $h \to 0$ we have

$$\mathbf{r}'(t) = \lim_{h \to 0} \frac{\mathbf{u}(t+h) \times [\mathbf{v}(t+h) - \mathbf{v}(t)]}{h} + \lim_{h \to 0} \frac{[\mathbf{u}(t+h) - \mathbf{u}(t)] \times \mathbf{v}(t)}{h} = \mathbf{u}(t) \times \mathbf{v}'(t) + \mathbf{u}'(t) \times \mathbf{v}(t)$$

by Exercise 13.1.53(a) and Definition 1.

46. $$\frac{d}{dt}[\mathbf{u}(f(t))] = \frac{d}{dt}\langle u_1(f(t)), u_2(f(t)), u_3(f(t))\rangle = \left\langle \frac{d}{dt}[u_1(f(t))], \frac{d}{dt}[u_2(f(t))], \frac{d}{dt}[u_3(f(t))] \right\rangle$$
$$= \langle f'(t)u_1'(f(t)), f'(t)u_2'(f(t)), f'(t)u_3'(f(t))\rangle = f'(t)\,\mathbf{u}'(t)$$

47. $$\frac{d}{dt}[\mathbf{u}(t) \cdot \mathbf{v}(t)] = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t) \qquad \text{[by Formula 4 of Theorem 3]}$$
$$= \langle \cos t, -\sin t, 1\rangle \cdot \langle t, \cos t, \sin t\rangle + \langle \sin t, \cos t, t\rangle \cdot \langle 1, -\sin t, \cos t\rangle$$
$$= t\cos t - \cos t\,\sin t + \sin t + \sin t - \cos t\,\sin t + t\cos t$$
$$= 2t\cos t + 2\sin t - 2\cos t\,\sin t$$

48. $$\frac{d}{dt}[\mathbf{u}(t) \times \mathbf{v}(t)] = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t) \qquad \text{[by Formula 5 of Theorem 3]}$$
$$= \langle \cos t, -\sin t, 1\rangle \times \langle t, \cos t, \sin t\rangle + \langle \sin t, \cos t, t\rangle \times \langle 1, -\sin t, \cos t\rangle$$
$$= \left\langle -\sin^2 t - \cos t, t - \cos t\,\sin t, \cos^2 t + t\sin t \right\rangle$$
$$+ \left\langle \cos^2 t + t\sin t, t - \cos t\,\sin t, -\sin^2 t - \cos t \right\rangle$$
$$= \left\langle \cos^2 t - \sin^2 t - \cos t + t\sin t, 2t - 2\cos t\,\sin t, \cos^2 t - \sin^2 t - \cos t + t\sin t \right\rangle$$

49. By Formula 4 of Theorem 3, $f'(t) = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t)$, and $\mathbf{v}'(t) = \langle 1, 2t, 3t^2\rangle$, so

$f'(2) = \mathbf{u}'(2) \cdot \mathbf{v}(2) + \mathbf{u}(2) \cdot \mathbf{v}'(2) = \langle 3, 0, 4\rangle \cdot \langle 2, 4, 8\rangle + \langle 1, 2, -1\rangle \cdot \langle 1, 4, 12\rangle = 6 + 0 + 32 + 1 + 8 - 12 = 35.$

50. By Formula 5 of Theorem 3, $\mathbf{r}'(t) = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$, so

$$\mathbf{r}'(2) = \mathbf{u}'(2) \times \mathbf{v}(2) + \mathbf{u}(2) \times \mathbf{v}'(2) = \langle 3, 0, 4\rangle \times \langle 2, 4, 8\rangle + \langle 1, 2, -1\rangle \times \langle 1, 4, 12\rangle$$
$$= \langle -16, -16, 12\rangle + \langle 28, -13, 2\rangle = \langle 12, -29, 14\rangle$$

51. $\mathbf{r}(t) = \mathbf{a}\cos\omega t + \mathbf{b}\sin\omega t \quad \Rightarrow \quad \mathbf{r}'(t) = -\mathbf{a}\,\omega\sin\omega t + \mathbf{b}\,\omega\cos\omega t$ by Formulas 1 and 3 of Theorem 3. Then

$$\mathbf{r}(t) \times \mathbf{r}'(t) = (\mathbf{a}\cos\omega t + \mathbf{b}\sin\omega t) \times (-\mathbf{a}\,\omega\sin\omega t + \mathbf{b}\,\omega\cos\omega t)$$
$$= (\mathbf{a}\cos\omega t + \mathbf{b}\sin\omega t) \times (-\mathbf{a}\,\omega\sin\omega t) + (\mathbf{a}\cos\omega t + \mathbf{b}\sin\omega t) \times (\mathbf{b}\,\omega\cos\omega t)$$
[by Property 3 of Theorem 12.4.11]
$$= \mathbf{a}\cos\omega t \times (-\mathbf{a}\,\omega\sin\omega t) + \mathbf{b}\sin\omega t \times (-\mathbf{a}\,\omega\sin\omega t) + \mathbf{a}\cos\omega t \times \mathbf{b}\,\omega\cos\omega t + \mathbf{b}\sin\omega t \times \mathbf{b}\,\omega\cos\omega t$$
[by Property 4]
$$= (\cos\omega t)\,(-\omega\sin\omega t)\,(\mathbf{a} \times \mathbf{a}) + (\sin\omega t)\,(-\omega\sin\omega t)\,(\mathbf{b} \times \mathbf{a}) + (\cos\omega t)\,(\omega\cos\omega t)\,(\mathbf{a} \times \mathbf{b})$$
$$+ (\sin\omega t)\,(\omega\cos\omega t)\,(\mathbf{b} \times \mathbf{b}) \qquad \text{[by Property 2]}$$
$$= \mathbf{0} + \left(\omega\sin^2\omega t\right)(\mathbf{a} \times \mathbf{b}) + \left(\omega\cos^2\omega t\right)(\mathbf{a} \times \mathbf{b}) + \mathbf{0} \qquad \text{[by Property 1 and Example 12.4.2]}$$
$$= \omega\left(\sin^2\omega t + \cos^2\omega t\right)(\mathbf{a} \times \mathbf{b}) = \omega\,(\mathbf{a} \times \mathbf{b}) = \omega\,\mathbf{a} \times \mathbf{b} \qquad \text{[by Property 2]}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. From Exercise 51, $\mathbf{r}'(t) = -\mathbf{a}\,\omega \sin \omega t + \mathbf{b}\,\omega \cos \omega t \quad \Rightarrow \quad \mathbf{r}''(t) = -\mathbf{a}\,\omega^2 \cos \omega t - \mathbf{b}\,\omega^2 \sin \omega t$. Then

$$\begin{aligned} \mathbf{r}''(t) + \omega^2 \mathbf{r}(t) &= \left(-\mathbf{a}\,\omega^2 \cos \omega t - \mathbf{b}\,\omega^2 \sin \omega t\right) + \omega^2 \left(\mathbf{a} \cos \omega t + \mathbf{b} \sin \omega t\right) \\ &= -\mathbf{a}\,\omega^2 \cos \omega t - \mathbf{b}\,\omega^2 \sin \omega t + \mathbf{a}\,\omega^2 \cos \omega t + \mathbf{b}\,\omega^2 \sin \omega t = \mathbf{0} \end{aligned}$$

53. $\dfrac{d}{dt}\left[\mathbf{r}(t) \times \mathbf{r}'(t)\right] = \mathbf{r}'(t) \times \mathbf{r}'(t) + \mathbf{r}(t) \times \mathbf{r}''(t)$ by Formula 5 of Theorem 3. But $\mathbf{r}'(t) \times \mathbf{r}'(t) = \mathbf{0}$ (by Example 12.4.2).

Thus, $\dfrac{d}{dt}\left[\mathbf{r}(t) \times \mathbf{r}'(t)\right] = \mathbf{r}(t) \times \mathbf{r}''(t)$.

54.
$$\begin{aligned} \frac{d}{dt}\left(\mathbf{u}(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right]\right) &= \mathbf{u}'(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right] + \mathbf{u}(t) \cdot \frac{d}{dt}\left[\mathbf{v}(t) \times \mathbf{w}(t)\right] \\ &= \mathbf{u}'(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right] + \mathbf{u}(t) \cdot \left[\mathbf{v}'(t) \times \mathbf{w}(t) + \mathbf{v}(t) \times \mathbf{w}'(t)\right] \\ &= \mathbf{u}'(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right] + \mathbf{u}(t) \cdot \left[\mathbf{v}'(t) \times \mathbf{w}(t)\right] + \mathbf{u}(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}'(t)\right] \\ &= \mathbf{u}'(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right] - \mathbf{v}'(t) \cdot \left[\mathbf{u}(t) \times \mathbf{w}(t)\right] + \mathbf{w}'(t) \cdot \left[\mathbf{u}(t) \times \mathbf{v}(t)\right] \end{aligned}$$

55. $\dfrac{d}{dt}\left|\mathbf{r}(t)\right| = \dfrac{d}{dt}\left[\mathbf{r}(t) \cdot \mathbf{r}(t)\right]^{1/2} = \frac{1}{2}\left[\mathbf{r}(t) \cdot \mathbf{r}(t)\right]^{-1/2}\left[2\mathbf{r}(t) \cdot \mathbf{r}'(t)\right] = \dfrac{1}{\left|\mathbf{r}(t)\right|}\,\mathbf{r}(t) \cdot \mathbf{r}'(t)$

56. Since $\mathbf{r}(t) \cdot \mathbf{r}'(t) = 0$, we have $0 = 2\mathbf{r}(t) \cdot \mathbf{r}'(t) = \dfrac{d}{dt}\left[\mathbf{r}(t) \cdot \mathbf{r}(t)\right] = \dfrac{d}{dt}\left|\mathbf{r}(t)\right|^2$. Thus $\left|\mathbf{r}(t)\right|^2$, and so $\left|\mathbf{r}(t)\right|$, is a constant, and hence the curve lies on a sphere with center the origin.

57. Since $\mathbf{u}(t) = \mathbf{r}(t) \cdot \left[\mathbf{r}'(t) \times \mathbf{r}''(t)\right]$,

$$\begin{aligned} \mathbf{u}'(t) &= \mathbf{r}'(t) \cdot \left[\mathbf{r}'(t) \times \mathbf{r}''(t)\right] + \mathbf{r}(t) \cdot \frac{d}{dt}\left[\mathbf{r}'(t) \times \mathbf{r}''(t)\right] \\ &= 0 + \mathbf{r}(t) \cdot \left[\mathbf{r}''(t) \times \mathbf{r}''(t) + \mathbf{r}'(t) \times \mathbf{r}'''(t)\right] && \left[\text{since } \mathbf{r}'(t) \perp \mathbf{r}'(t) \times \mathbf{r}''(t)\right] \\ &= \mathbf{r}(t) \cdot \left[\mathbf{r}'(t) \times \mathbf{r}'''(t)\right] && \left[\text{since } \mathbf{r}''(t) \times \mathbf{r}''(t) = \mathbf{0}\right] \end{aligned}$$

58. The tangent vector $\mathbf{r}'(t)$ is defined as $\lim\limits_{h \to 0} \dfrac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$. Here we assume that this limit exists and $\mathbf{r}'(t) \neq \mathbf{0}$; then we know that this vector lies on the tangent line to the curve. As in Figure 1, let points P and Q have position vectors $\mathbf{r}(t)$ and $\mathbf{r}(t+h)$. The vector $\mathbf{r}(t+h) - \mathbf{r}(t)$ points from P to Q, so $\mathbf{r}(t+h) - \mathbf{r}(t) = \overrightarrow{PQ}$. If $h > 0$ then $t < t+h$, so Q lies "ahead" of P on the curve. If h is sufficiently small (we can take h to be as small as we like since $h \to 0$) then $\overrightarrow{PQ}$ approximates the curve from P to Q and hence points approximately in the direction of the curve as t increases. Since h is positive, $\dfrac{1}{h}\overrightarrow{PQ} = \dfrac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$ points in the same direction. If $h < 0$, then $t > t+h$ so Q lies "behind" P on the curve. For h sufficiently small, $\overrightarrow{PQ}$ approximates the curve but points in the direction of decreasing t. However, h is negative, so $\dfrac{1}{h}\overrightarrow{PQ}$ $\dfrac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$ points in the opposite direction, that is, in the direction of increasing t. In both cases, the difference quotient $\dfrac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$ points in the direction of increasing t. The tangent vector $\mathbf{r}'(t)$ is the limit of this difference quotient, so it must also point in the direction of increasing t.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13.3 Arc Length and Curvature

1. $\mathbf{r}(t) = \langle t, 3\cos t, 3\sin t\rangle \Rightarrow \mathbf{r}'(t) = \langle 1, -3\sin t, 3\cos t\rangle \Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{1^2 + (-3\sin t)^2 + (3\cos t)^2} = \sqrt{1 + 9(\sin^2 t + \cos^2 t)} = \sqrt{10}.$

Then using Formula 3, we have $L = \int_{-5}^{5} |\mathbf{r}'(t)|\,dt = \int_{-5}^{5} \sqrt{10}\,dt = \sqrt{10}\,t\big]_{-5}^{5} = 10\sqrt{10}.$

2. $\mathbf{r}(t) = \langle 2t, t^2, \frac{1}{3}t^3\rangle \Rightarrow \mathbf{r}'(t) = \langle 2, 2t, t^2\rangle \Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{2^2 + (2t)^2 + (t^2)^2} = \sqrt{4 + 4t^2 + t^4} = \sqrt{(2 + t^2)^2} = 2 + t^2$ for $0 \le t \le 1$. Then using Formula 3, we have

$L = \int_0^1 |\mathbf{r}'(t)|\,dt = \int_0^1 (2 + t^2)\,dt = 2t + \frac{1}{3}t^3\big]_0^1 = \frac{7}{3}.$

3. $\mathbf{r}(t) = \sqrt{2}\,t\,\mathbf{i} + e^t\mathbf{j} + e^{-t}\mathbf{k} \Rightarrow \mathbf{r}'(t) = \sqrt{2}\,\mathbf{i} + e^t\mathbf{j} - e^{-t}\mathbf{k} \Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{\left(\sqrt{2}\right)^2 + (e^t)^2 + (-e^{-t})^2} = \sqrt{2 + e^{2t} + e^{-2t}} = \sqrt{(e^t + e^{-t})^2} = e^t + e^{-t}$ [since $e^t + e^{-t} > 0$].

Then $L = \int_0^1 |\mathbf{r}'(t)|\,dt = \int_0^1 (e^t + e^{-t})\,dt = \left[e^t - e^{-t}\right]_0^1 = e - e^{-1}.$

4. $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + \ln\cos t\,\mathbf{k} \Rightarrow \mathbf{r}'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j} + \dfrac{-\sin t}{\cos t}\,\mathbf{k} = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j} - \tan t\,\mathbf{k},$

$|\mathbf{r}'(t)| = \sqrt{(-\sin t)^2 + \cos^2 t + (-\tan t)^2} = \sqrt{1 + \tan^2 t} = \sqrt{\sec^2 t} = |\sec t|$. Since $\sec t > 0$ for $0 \le t \le \pi/4$, here we can say $|\mathbf{r}'(t)| = \sec t$. Then

$$L = \int_0^{\pi/4} \sec t\,dt = \left[\ln|\sec t + \tan t|\right]_0^{\pi/4} = \ln\left|\sec\tfrac{\pi}{4} + \tan\tfrac{\pi}{4}\right| - \ln|\sec 0 + \tan 0|$$
$$= \ln\left|\sqrt{2} + 1\right| - \ln|1 + 0| = \ln(\sqrt{2} + 1).$$

5. $\mathbf{r}(t) = \mathbf{i} + t^2\,\mathbf{j} + t^3\,\mathbf{k} \Rightarrow \mathbf{r}'(t) = 2t\,\mathbf{j} + 3t^2\,\mathbf{k} \Rightarrow |\mathbf{r}'(t)| = \sqrt{4t^2 + 9t^4} = t\sqrt{4 + 9t^2}$ [since $t \ge 0$].

Then $L = \int_0^1 |\mathbf{r}'(t)|\,dt = \int_0^1 t\sqrt{4 + 9t^2}\,dt = \frac{1}{18}\cdot\frac{2}{3}(4 + 9t^2)^{3/2}\big]_0^1 = \frac{1}{27}(13^{3/2} - 4^{3/2}) = \frac{1}{27}(13^{3/2} - 8).$

6. $\mathbf{r}(t) = t^2\,\mathbf{i} + 9t\,\mathbf{j} + 4t^{3/2}\,\mathbf{k} \Rightarrow \mathbf{r}'(t) = 2t\,\mathbf{i} + 9\,\mathbf{j} + 6\sqrt{t}\,\mathbf{k} \Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{4t^2 + 81 + 36t} = \sqrt{(2t + 9)^2} = |2t + 9| = 2t + 9$ [since $2t + 9 \ge 0$ for $1 \le t \le 4$]. Then

$L = \int_1^4 |\mathbf{r}'(t)|\,dt = \int_1^4 (2t + 9)\,dt = \left[t^2 + 9t\right]_1^4 = 52 - 10 = 42.$

7. $\mathbf{r}(t) = \langle t^2, t^3, t^4\rangle \Rightarrow \mathbf{r}'(t) = \langle 2t, 3t^2, 4t^3\rangle \Rightarrow |\mathbf{r}'(t)| = \sqrt{(2t)^2 + (3t^2)^2 + (4t^3)^2} = \sqrt{4t^2 + 9t^4 + 16t^6}$, so

$L = \int_0^2 |\mathbf{r}'(t)|\,dt = \int_0^2 \sqrt{4t^2 + 9t^4 + 16t^6}\,dt \approx 18.6833.$

8. $\mathbf{r}(t) = \langle t, e^{-t}, te^{-t}\rangle \Rightarrow \mathbf{r}'(t) = \langle 1, -e^{-t}, (1 - t)e^{-t}\rangle \Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{1^2 + (-e^{-t})^2 + [(1 - t)e^{-t}]^2} = \sqrt{1 + e^{-2t} + (1 - t)^2e^{-2t}} = \sqrt{1 + (2 - 2t + t^2)e^{-2t}}$, so

$L = \int_1^3 |\mathbf{r}'(t)|\,dt = \int_1^3 \sqrt{1 + (2 + 2t + t^2)e^{-2t}}\,dt \approx 2.0454.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

9. $\mathbf{r}(t) = \langle \cos \pi t, 2t, \sin 2\pi t \rangle \Rightarrow \mathbf{r}'(t) = \langle -\pi \sin \pi t, 2, 2\pi \cos 2\pi t \rangle \Rightarrow |\mathbf{r}'(t)| = \sqrt{\pi^2 \sin^2 \pi t + 4 + 4\pi^2 \cos^2 2\pi t}$.

The point $(1, 0, 0)$ corresponds to $t = 0$ and $(1, 4, 0)$ corresponds to $t = 2$, so the length is

$L = \int_0^2 |\mathbf{r}'(t)|\, dt = \int_0^2 \sqrt{\pi^2 \sin^2 \pi t + 4 + 4\pi^2 \cos^2 2\pi t}\, dt \approx 10.3311$.

10. We plot two different views of the curve with parametric equations $x = \sin t$, $y = \sin 2t$, $z = \sin 3t$. To help visualize the curve, we also include a plot showing a tube of radius 0.07 around the curve.







The complete curve is given by the parameter interval $[0, 2\pi]$ and we have $\mathbf{r}'(t) = \langle \cos t, 2\cos 2t, 3\cos 3t \rangle \Rightarrow$ $|\mathbf{r}'(t)| = \sqrt{\cos^2 t + 4\cos^2 2t + 9\cos^2 3t}$, so $L = \int_0^{2\pi} |\mathbf{r}'(t)|\, dt = \int_0^{2\pi} \sqrt{\cos^2 t + 4\cos^2 2t + 9\cos^2 3t}\, dt \approx 16.0264$.

11. The projection of the curve C onto the xy-plane is the curve $x^2 = 2y$ or $y = \frac{1}{2}x^2$, $z = 0$. Then we can choose the parameter $x = t \Rightarrow y = \frac{1}{2}t^2$. Since C also lies on the surface $3z = xy$, we have $z = \frac{1}{3}xy = \frac{1}{3}(t)(\frac{1}{2}t^2) = \frac{1}{6}t^3$. Then parametric equations for C are $x = t$, $y = \frac{1}{2}t^2$, $z = \frac{1}{6}t^3$ and the corresponding vector equation is $\mathbf{r}(t) = \langle t, \frac{1}{2}t^2, \frac{1}{6}t^3 \rangle$. The origin corresponds to $t = 0$ and the point $(6, 18, 36)$ corresponds to $t = 6$, so

$$L = \int_0^6 |\mathbf{r}'(t)|\, dt = \int_0^6 \left|\left\langle 1, t, \tfrac{1}{2}t^2 \right\rangle\right|\, dt = \int_0^6 \sqrt{1^2 + t^2 + \left(\tfrac{1}{2}t^2\right)^2}\, dt = \int_0^6 \sqrt{1 + t^2 + \tfrac{1}{4}t^4}\, dt$$
$$= \int_0^6 \sqrt{(1 + \tfrac{1}{2}t^2)^2}\, dt = \int_0^6 (1 + \tfrac{1}{2}t^2)\, dt = \left[t + \tfrac{1}{6}t^3\right]_0^6 = 6 + 36 = 42$$

12. Let C be the curve of intersection. The projection of C onto the xy-plane is the ellipse $4x^2 + y^2 = 4$ or $x^2 + y^2/4 = 1$, $z = 0$. Then we can write $x = \cos t$, $y = 2\sin t$, $0 \le t \le 2\pi$. Since C also lies on the plane $x + y + z = 2$, we have $z = 2 - x - y = 2 - \cos t - 2\sin t$. Then parametric equations for C are $x = \cos t$, $y = 2\sin t$, $z = 2 - \cos t - 2\sin t$, $0 \le t \le 2\pi$, and the corresponding vector equation is $\mathbf{r}(t) = \langle \cos t, 2\sin t, 2 - \cos t - 2\sin t \rangle$. Differentiating gives $\mathbf{r}'(t) = \langle -\sin t, 2\cos t, \sin t - 2\cos t \rangle \Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{(-\sin t)^2 + (2\cos t)^2 + (\sin t - 2\cos t)^2} = \sqrt{2\sin^2 t + 8\cos^2 t - 4\sin t \cos t}$. The length of C is

$L = \int_0^{2\pi} |\mathbf{r}'(t)|\, dt = \int_0^{2\pi} \sqrt{2\sin^2 t + 8\cos^2 t - 4\sin t \cos t}\, dt \approx 13.5191$.

13. (a) $\mathbf{r}(t) = (5 - t)\,\mathbf{i} + (4t - 3)\,\mathbf{j} + 3t\,\mathbf{k} \Rightarrow \mathbf{r}'(t) = -\mathbf{i} + 4\,\mathbf{j} + 3\,\mathbf{k}$ and $\frac{ds}{dt} = |\mathbf{r}'(t)| = \sqrt{1 + 16 + 9} = \sqrt{26}$. The point $P\,(4, 1, 3)$ corresponds to $t = 1$, so the arc length function from P is

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$s(t) = \int_1^t |\mathbf{r}'(u)|\,du = \int_1^t \sqrt{26}\,du = \sqrt{26}\,u\big|_1^t = \sqrt{26}\,(t-1)$. Since $s = \sqrt{26}\,(t-1)$, we have $t = \dfrac{s}{\sqrt{26}} + 1$.

Substituting for t in the original equation, the reparametrization of the curve with respect to arc length is

$$\begin{aligned}\mathbf{r}(t(s)) &= \left[5 - \left(\frac{s}{\sqrt{26}} + 1\right)\right]\mathbf{i} + \left[4\left(\frac{s}{\sqrt{26}} + 1\right) - 3\right]\mathbf{j} + 3\left(\frac{s}{\sqrt{26}} + 1\right)\mathbf{k} \\ &= \left(4 - \frac{s}{\sqrt{26}}\right)\mathbf{i} + \left(\frac{4s}{\sqrt{26}} + 1\right)\mathbf{j} + \left(\frac{3s}{\sqrt{26}} + 3\right)\mathbf{k}\end{aligned}$$

(b) The point 4 units along the curve from P has position vector

$\mathbf{r}(t(4)) = \left(4 - \dfrac{4}{\sqrt{26}}\right)\mathbf{i} + \left(\dfrac{4(4)}{\sqrt{26}} + 1\right)\mathbf{j} + \left(\dfrac{3(4)}{\sqrt{26}} + 3\right)\mathbf{k}$, so the point is $\left(4 - \dfrac{4}{\sqrt{26}}, \dfrac{16}{\sqrt{26}} + 1, \dfrac{12}{\sqrt{26}} + 3\right)$.

14. (a) $\mathbf{r}(t) = e^t \sin t\,\mathbf{i} + e^t \cos t\,\mathbf{j} + \sqrt{2}\,e^t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = e^t(\cos t + \sin t)\,\mathbf{i} + e^t(\cos t - \sin t)\,\mathbf{j} + \sqrt{2}\,e^t\,\mathbf{k}$ and

$$\begin{aligned}\frac{ds}{dt} = |\mathbf{r}'(t)| &= \sqrt{e^{2t}(\cos t + \sin t)^2 + e^{2t}(\cos t - \sin t)^2 + 2e^{2t}} \\ &= \sqrt{e^{2t}\left[2(\cos^2 t + \sin^2 t) + 2\cos t \sin t - 2\cos t \sin t + 2\right]} = \sqrt{4e^{2t}} = 2e^t\end{aligned}$$

The point $P\left(0, 1, \sqrt{2}\right)$ corresponds to $t = 0$, so the arc length function from P is

$s(t) = \int_0^t |\mathbf{r}'(u)|\,du = \int_0^t 2e^u\,du = 2e^u\big|_0^t = 2(e^t - 1)$. Since $s = 2(e^t - 1)$, we have $e^t = \dfrac{s}{2} + 1 \;\Leftrightarrow$

$t = \ln\left(\frac{1}{2}s + 1\right)$. Substituting for t in the original equation, the reparametrization of the curve with respect to arc length is

$\mathbf{r}(t(s)) = \left(\frac{1}{2}s + 1\right)\sin\left(\ln\left(\frac{1}{2}s + 1\right)\right)\mathbf{i} + \left(\frac{1}{2}s + 1\right)\cos\left(\ln\left(\frac{1}{2}s + 1\right)\right)\mathbf{j} + \left(\frac{\sqrt{2}}{2}s + \sqrt{2}\right)\mathbf{k}$.

(b) The point 4 units along the curve from P has position vector

$\mathbf{r}(t(4)) = \left(\frac{1}{2}(4) + 1\right)\sin\left(\ln\left(\frac{1}{2}(4) + 1\right)\right)\mathbf{i} + \left(\frac{1}{2}(4) + 1\right)\cos\left(\ln\left(\frac{1}{2}(4) + 1\right)\right)\mathbf{j} + \left(\frac{\sqrt{2}}{2}(4) + \sqrt{2}\right)\mathbf{k}$, so the point is

$\left(3\sin(\ln 3), 3\cos(\ln 3), 3\sqrt{2}\right)$.

15. Here $\mathbf{r}(t) = \langle 3\sin t, 4t, 3\cos t\rangle$, so $\mathbf{r}'(t) = \langle 3\cos t, 4, -3\sin t\rangle$ and $|\mathbf{r}'(t)| = \sqrt{9\cos^2 t + 16 + 9\sin^2 t} = \sqrt{25} = 5$.

The point $(0, 0, 3)$ corresponds to $t = 0$, so the arc length function beginning at $(0, 0, 3)$ and measuring in the positive direction is given by $s(t) = \int_0^t |\mathbf{r}'(u)|\,du = \int_0^t 5\,du = 5t$. $s(t) = 5 \;\Rightarrow\; 5t = 5 \;\Rightarrow\; t = 1$, thus your location after moving 5 units along the curve is $(3\sin 1, 4, 3\cos 1)$.

16. $\mathbf{r}(t) = \left(\dfrac{2}{t^2+1} - 1\right)\mathbf{i} + \dfrac{2t}{t^2+1}\,\mathbf{j} \;\Rightarrow\; \mathbf{r}'(t) = \dfrac{-4t}{(t^2+1)^2}\,\mathbf{i} + \dfrac{-2t^2+2}{(t^2+1)^2}\,\mathbf{j}$,

$$\frac{ds}{dt} = |\mathbf{r}'(t)| = \sqrt{\left[\frac{-4t}{(t^2+1)^2}\right]^2 + \left[\frac{-2t^2+2}{(t^2+1)^2}\right]^2} = \sqrt{\frac{4t^4 + 8t^2 + 4}{(t^2+1)^4}} = \sqrt{\frac{4(t^2+1)^2}{(t^2+1)^4}} = \sqrt{\frac{4}{(t^2+1)^2}} = \frac{2}{t^2+1}.$$

Since the initial point $(1, 0)$ corresponds to $t = 0$, the arc length function is

$s(t) = \displaystyle\int_0^t |\mathbf{r}'(u)|\,du = \int_0^t \frac{2}{u^2+1}\,du = 2\arctan t$. Then $\arctan t = \frac{1}{2}s \;\Rightarrow\; t = \tan\frac{1}{2}s$. Substituting, we have

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$$\mathbf{r}(t(s)) = \left[\frac{2}{\tan^2\left(\frac{1}{2}s\right)+1} - 1\right]\mathbf{i} + \frac{2\tan\left(\frac{1}{2}s\right)}{\tan^2\left(\frac{1}{2}s\right)+1}\mathbf{j} = \frac{1-\tan^2\left(\frac{1}{2}s\right)}{1+\tan^2\left(\frac{1}{2}s\right)}\mathbf{i} + \frac{2\tan\left(\frac{1}{2}s\right)}{\sec^2\left(\frac{1}{2}s\right)}\mathbf{j}$$

$$- \frac{1-\tan^2\left(\frac{1}{2}s\right)}{\sec^2\left(\frac{1}{2}s\right)}\mathbf{i} + 2\tan\left(\frac{1}{2}s\right)\cos^2\left(\frac{1}{2}s\right)\mathbf{j} = \left[\cos^2\left(\frac{1}{2}s\right) - \sin^2\left(\frac{1}{2}s\right)\right]\mathbf{i} + 2\sin\left(\frac{1}{2}s\right)\cos\left(\frac{1}{2}s\right)\mathbf{j} = \cos s\,\mathbf{i} + \sin s\,\mathbf{j}$$

With this parametrization, we recognize the function as representing the unit circle. Note here that the curve approaches, but does not include, the point $(-1, 0)$, since $\cos s = -1$ for $s = \pi + 2k\pi$ (k an integer) but then $t = \tan\left(\frac{1}{2}s\right)$ is undefined.

17. (a) $\mathbf{r}(t) = \langle t, 3\cos t, 3\sin t\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, -3\sin t, 3\cos t\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{1 + 9\sin^2 t + 9\cos^2 t} = \sqrt{10}$.

Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \frac{1}{\sqrt{10}}\langle 1, -3\sin t, 3\cos t\rangle$ or $\left\langle \frac{1}{\sqrt{10}}, -\frac{3}{\sqrt{10}}\sin t, \frac{3}{\sqrt{10}}\cos t\right\rangle$.

$\mathbf{T}'(t) = \frac{1}{\sqrt{10}}\langle 0, -3\cos t, -3\sin t\rangle \;\Rightarrow\; |\mathbf{T}'(t)| = \frac{1}{\sqrt{10}}\sqrt{0 + 9\cos^2 t + 9\sin^2 t} = \frac{3}{\sqrt{10}}$. Thus

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \frac{1/\sqrt{10}}{3/\sqrt{10}}\langle 0, -3\cos t, -3\sin t\rangle = \langle 0, -\cos t, -\sin t\rangle.$$

(b) $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{3/\sqrt{10}}{\sqrt{10}} = \dfrac{3}{10}$

18. (a) $\mathbf{r}(t) = \langle t^2, \sin t - t\cos t, \cos t + t\sin t\rangle \;\Rightarrow$

$\mathbf{r}'(t) = \langle 2t, \cos t + t\sin t - \cos t,\ -\sin t + t\cos t + \sin t\rangle = \langle 2t, t\sin t, t\cos t\rangle \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{4t^2 + t^2\sin^2 t + t^2\cos^2 t} = \sqrt{4t^2 + t^2(\cos^2 t + \sin^2 t)} = \sqrt{5t^2} = \sqrt{5}\,t$ [since $t > 0$]. Then

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{5}\,t}\langle 2t, t\sin t, t\cos t\rangle = \frac{1}{\sqrt{5}}\langle 2, \sin t, \cos t\rangle$. $\mathbf{T}'(t) = \frac{1}{\sqrt{5}}\langle 0, \cos t, -\sin t\rangle \;\Rightarrow$

$|\mathbf{T}'(t)| = \frac{1}{\sqrt{5}}\sqrt{0 + \cos^2 t + \sin^2 t} = \frac{1}{\sqrt{5}}$. Thus $\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{1/\sqrt{5}}{1/\sqrt{5}}\langle 0, \cos t, -\sin t\rangle = \langle 0, \cos t, -\sin t\rangle$.

(b) $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{1/\sqrt{5}}{\sqrt{5}\,t} = \dfrac{1}{5t}$

19. (a) $\mathbf{r}(t) = \langle \sqrt{2}\,t, e^t, e^{-t}\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle \sqrt{2}, e^t, -e^{-t}\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{2 + e^{2t} + e^{-2t}} = \sqrt{(e^t + e^{-t})^2} = e^t + e^{-t}$.

Then

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \frac{1}{e^t + e^{-t}}\langle \sqrt{2}, e^t, -e^{-t}\rangle = \frac{1}{e^{2t}+1}\langle \sqrt{2}\,e^t, e^{2t}, -1\rangle \quad \left[\text{after multiplying by } \frac{e^t}{e^t}\right] \quad \text{and}$$

$$\mathbf{T}'(t) = \frac{1}{e^{2t}+1}\langle \sqrt{2}\,e^t, 2e^{2t}, 0\rangle - \frac{2e^{2t}}{(e^{2t}+1)^2}\langle \sqrt{2}\,e^t, e^{2t}, -1\rangle$$

$$= \frac{1}{(e^{2t}+1)^2}\left[(e^{2t}+1)\langle \sqrt{2}\,e^t, 2e^{2t}, 0\rangle - 2e^{2t}\langle \sqrt{2}\,e^t, e^{2t}, -1\rangle\right] = \frac{1}{(e^{2t}+1)^2}\langle \sqrt{2}\,e^t(1 - e^{2t}), 2e^{2t}, 2e^{2t}\rangle$$

Then

$$|\mathbf{T}'(t)| = \frac{1}{(e^{2t}+1)^2}\sqrt{2e^{2t}(1 - 2e^{2t} + e^{4t}) + 4e^{4t} + 4e^{4t}} = \frac{1}{(e^{2t}+1)^2}\sqrt{2e^{2t}(1 + 2e^{2t} + e^{4t})}$$

$$= \frac{1}{(e^{2t}+1)^2}\sqrt{2e^{2t}(1 + e^{2t})^2} = \frac{\sqrt{2}\,e^t(1 + e^{2t})}{(e^{2t}+1)^2} = \frac{\sqrt{2}\,e^t}{e^{2t}+1}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Therefore

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \frac{e^{2t}+1}{\sqrt{2}\,e^t}\,\frac{1}{(e^{2t}+1)^2}\,\langle \sqrt{2}\,e^t(1-e^{2t}), 2e^{2t}, 2e^{2t}\rangle$$

$$= \frac{1}{\sqrt{2}\,e^t(e^{2t}+1)}\,\langle \sqrt{2}\,e^t(1-e^{2t}), 2e^{2t}, 2e^{2t}\rangle = \frac{1}{e^{2t}+1}\,\langle 1-e^{2t}, \sqrt{2}\,e^t, \sqrt{2}\,e^t\rangle$$

(b) $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{\sqrt{2}\,e^t}{e^{2t}+1}\cdot\dfrac{1}{e^t+e^{-t}} = \dfrac{\sqrt{2}\,e^t}{e^{3t}+2e^t+e^{-t}} = \dfrac{\sqrt{2}\,e^{2t}}{e^{4t}+2e^{2t}+1} = \dfrac{\sqrt{2}\,e^{2t}}{(e^{2t}+1)^2}$

20. (a) $\mathbf{r}(t) = \langle t, \frac{1}{2}t^2, t^2\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, t, 2t\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{1+t^2+4t^2} = \sqrt{1+5t^2}$. Then

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \frac{1}{\sqrt{1+5t^2}}\,\langle 1, t, 2t\rangle.$$

$$\mathbf{T}'(t) = \frac{-5t}{(1+5t^2)^{3/2}}\,\langle 1, t, 2t\rangle + \frac{1}{\sqrt{1+5t^2}}\,\langle 0, 1, 2\rangle \qquad \text{[by Formula 3 of Theorem 13.2.3]}$$

$$= \frac{1}{(1+5t^2)^{3/2}}\,\left(\langle -5t, -5t^2, -10t^2\rangle + \langle 0, 1+5t^2, 2+10t^2\rangle\right) = \frac{1}{(1+5t^2)^{3/2}}\,\langle -5t, 1, 2\rangle$$

$$|\mathbf{T}'(t)| = \frac{1}{(1+5t^2)^{3/2}}\,\sqrt{25t^2+1+4} = \frac{1}{(1+5t^2)^{3/2}}\,\sqrt{25t^2+5} = \frac{\sqrt{5}\,\sqrt{5t^2+1}}{(1+5t^2)^{3/2}} = \frac{\sqrt{5}}{1+5t^2}$$

Thus $\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{1+5t^2}{\sqrt{5}}\cdot\dfrac{1}{(1+5t^2)^{3/2}}\,\langle -5t, 1, 2\rangle = \dfrac{1}{\sqrt{5+25t^2}}\,\langle -5t, 1, 2\rangle.$

(b) $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{\sqrt{5}/(1+5t^2)}{\sqrt{1+5t^2}} = \dfrac{\sqrt{5}}{(1+5t^2)^{3/2}}$

21. $\mathbf{r}(t) = t^3\,\mathbf{j} + t^2\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = 3t^2\,\mathbf{j} + 2t\,\mathbf{k},\quad \mathbf{r}''(t) = 6t\,\mathbf{j} + 2\,\mathbf{k},\quad |\mathbf{r}'(t)| = \sqrt{0^2+(3t^2)^2+(2t)^2} = \sqrt{9t^4+4t^2}$,

$\mathbf{r}'(t)\times\mathbf{r}''(t) = -6t^2\,\mathbf{i},\quad |\mathbf{r}'(t)\times\mathbf{r}''(t)| = 6t^2$. Then $\kappa(t) = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{6t^2}{\left(\sqrt{9t^4+4t^2}\,\right)^3} = \dfrac{6t^2}{(9t^4+4t^2)^{3/2}}.$

22. $\mathbf{r}(t) = t\,\mathbf{i} + t^2\,\mathbf{j} + e^t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = \mathbf{i} + 2t\,\mathbf{j} + e^t\,\mathbf{k},\quad \mathbf{r}''(t) = 2\,\mathbf{j} + e^t\,\mathbf{k}$,

$|\mathbf{r}'(t)| = \sqrt{1^2+(2t)^2+(e^t)^2} = \sqrt{1+4t^2+e^{2t}},\quad \mathbf{r}'(t)\times\mathbf{r}''(t) = (2t-2)e^t\,\mathbf{i} - e^t\,\mathbf{j} + 2\,\mathbf{k}$,

$|\mathbf{r}'(t)\times\mathbf{r}''(t)| = \sqrt{[(2t-2)e^t]^2+(-e^t)^2+2^2} = \sqrt{(2t-2)^2e^{2t}+e^{2t}+4} = \sqrt{(4t^2-8t+5)e^{2t}+4}.$

Then $\kappa(t) = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{\sqrt{(4t^2-8t+5)e^{2t}+4}}{\left(\sqrt{1+4t^2+e^{2t}}\,\right)^3} = \dfrac{\sqrt{(4t^2-8t+5)e^{2t}+4}}{(1+4t^2+e^{2t}\,)^{3/2}}.$

23. $\mathbf{r}(t) = \sqrt{6}\,t^2\,\mathbf{i} + 2t\,\mathbf{j} + 2t^3\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = 2\sqrt{6}\,t\,\mathbf{i} + 2\,\mathbf{j} + 6t^2\,\mathbf{k},\quad \mathbf{r}''(t) = 2\sqrt{6}\,\mathbf{i} + 12t\,\mathbf{k}$,

$|\mathbf{r}'(t)| = \sqrt{24t^2+4+36t^4} = \sqrt{4(9t^4+6t^2+1)} = \sqrt{4(3t^2+1)^2} = 2(3t^2+1)$,

$\mathbf{r}'(t)\times\mathbf{r}''(t) = 24t\,\mathbf{i} - 12\sqrt{6}\,t^2\,\mathbf{j} - 4\sqrt{6}\,\mathbf{k}$,

$|\mathbf{r}'(t)\times\mathbf{r}''(t)| = \sqrt{576t^2+864t^4+96} = \sqrt{96(9t^4+6t^2+1)} = \sqrt{96(3t^2+1)^2} = 4\sqrt{6}\,(3t^2+1)$.

Then $\kappa(t) = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{4\sqrt{6}\,(3t^2+1)}{8(3t^2+1)^3} = \dfrac{\sqrt{6}}{2(3t^2+1)^2}.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

24. $\mathbf{r}(t) = \langle t^2, \ln t, t \ln t \rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 2t, 1/t, 1 + \ln t \rangle, \quad \mathbf{r}''(t) = \langle 2, -1/t^2, 1/t \rangle.$ The point $(1, 0, 0)$ corresponds to $t = 1$, and $\mathbf{r}'(1) = \langle 2, 1, 1 \rangle, \quad |\mathbf{r}'(1)| = \sqrt{2^2 + 1^2 + 1^2} = \sqrt{6}, \quad \mathbf{r}''(1) = \langle 2, -1, 1 \rangle, \quad \mathbf{r}'(1) \times \mathbf{r}''(1) = \langle 2, 0, -4 \rangle,$ $|\mathbf{r}'(1) \times \mathbf{r}''(1)| = \sqrt{2^2 + 0^2 + (-4)^2} = \sqrt{20} = 2\sqrt{5}$. Then $\kappa(1) = \dfrac{|\mathbf{r}'(1) \times \mathbf{r}''(1)|}{|\mathbf{r}'(1)|^3} = \dfrac{2\sqrt{5}}{(\sqrt{6})^3} = \dfrac{2\sqrt{5}}{6\sqrt{6}}$ or $\dfrac{\sqrt{30}}{18}$.

25. $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, 2t, 3t^2 \rangle.$ The point $(1, 1, 1)$ corresponds to $t = 1$, and $\mathbf{r}'(1) = \langle 1, 2, 3 \rangle \;\Rightarrow$ $|\mathbf{r}'(1)| = \sqrt{1 + 4 + 9} = \sqrt{14}. \quad \mathbf{r}''(t) = \langle 0, 2, 6t \rangle \;\Rightarrow\; \mathbf{r}''(1) = \langle 0, 2, 6 \rangle. \quad \mathbf{r}'(1) \times \mathbf{r}''(1) = \langle 6, -6, 2 \rangle$, so $|\mathbf{r}'(1) \times \mathbf{r}''(1)| = \sqrt{36 + 36 + 4} = \sqrt{76}$. Then $\kappa(1) = \dfrac{|\mathbf{r}'(1) \times \mathbf{r}''(1)|}{|\mathbf{r}'(1)|^3} = \dfrac{\sqrt{76}}{\sqrt{14}^3} = \dfrac{1}{7}\sqrt{\dfrac{19}{14}}$.

26.



Note that we get the complete curve for $0 \le t < 2\pi$.

$\mathbf{r}(t) = \langle \cos t, \sin t, \sin 5t \rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle -\sin t, \cos t, 5\cos 5t \rangle,$

$\mathbf{r}''(t) = \langle -\cos t, -\sin t, -25\sin 5t \rangle.$ The point $(1, 0, 0)$ corresponds to $t = 0$, and $\mathbf{r}'(0) = \langle 0, 1, 5 \rangle \;\Rightarrow$

$|\mathbf{r}'(0)| = \sqrt{0^2 + 1^2 + 5^2} = \sqrt{26}, \quad \mathbf{r}''(0) = \langle -1, 0, 0 \rangle,$

$\mathbf{r}'(0) \times \mathbf{r}''(0) = \langle 0, -5, 1 \rangle \;\Rightarrow$

$|\mathbf{r}'(0) \times \mathbf{r}''(0)| = \sqrt{0^2 + (-5)^2 + 1^2} = \sqrt{26}.$ The curvature at the point $(1, 0, 0)$ is $\kappa(0) = \dfrac{|\mathbf{r}'(0) \times \mathbf{r}''(0)|}{|\mathbf{r}'(0)|^3} = \dfrac{\sqrt{26}}{(\sqrt{26})^3} = \dfrac{1}{26}$.

27. $f(x) = x^4, \quad f'(x) = 4x^3, \quad f''(x) = 12x^2, \quad \kappa(x) = \dfrac{|f''(x)|}{[1 + (f'(x))^2]^{3/2}} = \dfrac{|12x^2|}{[1 + (4x^3)^2]^{3/2}} = \dfrac{12x^2}{(1 + 16x^6)^{3/2}}$

28. $f(x) = \tan x, \quad f'(x) = \sec^2 x, \quad f''(x) = 2\sec x \cdot \sec x \tan x = 2\sec^2 x \tan x,$

$$\kappa(x) = \frac{|f''(x)|}{[1 + (f'(x))^2]^{3/2}} = \frac{|2\sec^2 x \tan x|}{[1 + (\sec^2 x)^2]^{3/2}} = \frac{2\sec^2 x\,|\tan x|}{(1 + \sec^4 x)^{3/2}}$$

29. $f(x) = xe^x, \quad f'(x) = xe^x + e^x, \quad f''(x) = xe^x + 2e^x,$

$$\kappa(x) = \frac{|f''(x)|}{[1 + (f'(x))^2]^{3/2}} = \frac{|xe^x + 2e^x|}{[1 + (xe^x + e^x)^2]^{3/2}} = \frac{|x + 2|\,e^x}{[1 + (xe^x + e^x)^2]^{3/2}}$$

30. $y' = \dfrac{1}{x}, \quad y'' = -\dfrac{1}{x^2},$

$$\kappa(x) = \frac{|y''(x)|}{\left[1 + (y'(x))^2\right]^{3/2}} = \left|\frac{-1}{x^2}\right| \frac{1}{(1 + 1/x^2)^{3/2}} = \frac{1}{x^2}\frac{(x^2)^{3/2}}{(x^2 + 1)^{3/2}} = \frac{|x|}{(x^2 + 1)^{3/2}} = \frac{x}{(x^2 + 1)^{3/2}} \quad [\text{since } x > 0].$$

To find the maximum curvature, we first find the critical numbers of $\kappa(x)$:

$$\kappa'(x) = \frac{(x^2 + 1)^{3/2} \cdot 1 - x\left(\frac{3}{2}\right)(x^2 + 1)^{1/2}(2x)}{[(x^2 + 1)^{3/2}]^2} = \frac{(x^2 + 1)^{1/2}[(x^2 + 1) - 3x^2]}{(x^2 + 1)^3} = \frac{1 - 2x^2}{(x^2 + 1)^{5/2}},$$

$\kappa'(x) = 0 \;\Rightarrow\; 1 - 2x^2 = 0$, so the only critical number in the domain is $x = \frac{1}{\sqrt{2}}$. Since $\kappa'(x) > 0$ for $0 < x < \frac{1}{\sqrt{2}}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and $\kappa'(x) < 0$ for $x > \frac{1}{\sqrt{2}}$, $\kappa(x)$ attains its maximum at $x = \frac{1}{\sqrt{2}}$. Thus, the maximum curvature occurs at $\left(\frac{1}{\sqrt{2}}, \ln\frac{1}{\sqrt{2}}\right)$.

Since $\lim_{x\to\infty} \dfrac{x}{(x^2+1)^{3/2}} = 0$, $\kappa(x)$ approaches 0 as $x \to \infty$.

31. Since $y' = y'' = e^x$, the curvature is $\kappa(x) = \dfrac{|y''(x)|}{[1+(y'(x))^2]^{3/2}} = \dfrac{e^x}{(1+e^{2x})^{3/2}} = e^x(1+e^{2x})^{-3/2}$.

To find the maximum curvature, we first find the critical numbers of $\kappa(x)$:

$$\kappa'(x) = e^x(1+e^{2x})^{-3/2} + e^x\left(-\tfrac{3}{2}\right)(1+e^{2x})^{-5/2}(2e^{2x}) = e^x\frac{1+e^{2x}-3e^{2x}}{(1+e^{2x})^{5/2}} = e^x\frac{1-2e^{2x}}{(1+e^{2x})^{5/2}}.$$

$\kappa'(x) = 0$ when $1-2e^{2x} = 0$, so $e^{2x} = \frac{1}{2}$ or $x = -\frac{1}{2}\ln 2$. And since $1-2e^{2x} > 0$ for $x < -\frac{1}{2}\ln 2$ and $1-2e^{2x} < 0$ for $x > -\frac{1}{2}\ln 2$, the maximum curvature is attained at the point $\left(-\frac{1}{2}\ln 2, e^{(-\ln 2)/2}\right) = \left(-\frac{1}{2}\ln 2, \frac{1}{\sqrt{2}}\right)$.

Since $\lim_{x\to\infty} e^x(1+e^{2x})^{-3/2} = 0$, $\kappa(x)$ approaches 0 as $x \to \infty$.

32. We can take the parabola as having its vertex at the origin and opening upward, so the equation is $f(x) = ax^2, a > 0$. Then by Equation 11, $\kappa(x) = \dfrac{|f''(x)|}{[1+(f'(x))^2]^{3/2}} = \dfrac{|2a|}{[1+(2ax)^2]^{3/2}} = \dfrac{2a}{(1+4a^2x^2)^{3/2}}$, thus $\kappa(0) = 2a$. We want $\kappa(0) = 4$, so $a = 2$ and the equation is $y = 2x^2$.

33. (a) C appears to be changing direction more quickly at P than Q, so we would expect the curvature to be greater at P.

(b) First we sketch approximate osculating circles at P and Q. Using the axes scale as a guide, we measure the radius of the osculating circle at P to be approximately 0.8 units, thus $\rho = \dfrac{1}{\kappa} \Rightarrow \kappa = \dfrac{1}{\rho} \approx \dfrac{1}{0.8} \approx 1.3$. Similarly, we estimate the radius of the osculating circle at Q to be 1.4 units, so $\kappa = \dfrac{1}{\rho} \approx \dfrac{1}{1.4} \approx 0.7$.



34. $y = x^4 - 2x^2 \Rightarrow y' = 4x^3 - 4x$, $y'' = 12x^2 - 4$, and $\kappa(x) = \dfrac{|y''|}{\left[1+(y')^2\right]^{3/2}} = \dfrac{|12x^2-4|}{\left[1+(4x^3-4x)^2\right]^{3/2}}$. The graph of the curvature here is what we would expect. The graph of $y = x^4 - 2x^2$ appears to be bending most sharply at the origin and near $x = \pm 1$.



35. $y = x^{-2} \Rightarrow y' = -2x^{-3}$, $y'' = 6x^{-4}$, and

$$\kappa(x) = \frac{|y''|}{\left[1+(y')^2\right]^{3/2}} = \frac{|6x^{-4}|}{\left[1+(-2x^{-3})^2\right]^{3/2}} = \frac{6}{x^4\left(1+4x^{-6}\right)^{3/2}}.$$

The appearance of the two humps in this graph is perhaps a little surprising, but it is explained by the fact that $y = x^{-2}$ increases asymptotically at the origin from both directions, and so its graph has very little bend there. [Note that $\kappa(0)$ is undefined.]



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

36. $\mathbf{r}(t) = \langle t - \sin t, 1 - \cos t, 4\cos(t/2)\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1 - \cos t, \sin t, -2\sin(t/2)\rangle, \quad \mathbf{r}''(t) = \langle \sin t, \cos t, -\cos(t/2)\rangle.$

Using a CAS, $\mathbf{r}'(t) \times \mathbf{r}''(t) = \langle -2\sin^3(t/2), -\sin(t/2)\sin t, \cos t - 1\rangle$, $|\mathbf{r}'(t) \times \mathbf{r}''(t)| = \sqrt{3 - 4\cos t + \cos 2t}$ or $2\sqrt{2}\sin^2(t/2)$, and $|\mathbf{r}'(t)| = 2\sqrt{1 - \cos t}$ or $2\sqrt{2}\,|\sin(t/2)|$. (To compute cross products in Maple, use the `VectorCalculus` or `LinearAlgebra` package and the `CrossProduct(a,b)` command. Here loading the `RealDomain` package will give simpler results. In Mathematica, use `Cross[a,b]`.)

Then $\kappa(t) = \dfrac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{\sqrt{3 - 4\cos t + \cos 2t}}{8\,(1 - \cos t)^{3/2}}$ or $\dfrac{1}{4\sqrt{2 - 2\cos t}}$ or $\dfrac{1}{8\,|\sin(t/2)|}$. We plot the space curve and its curvature function for $0 \le t \le 8\pi$ below.





The asymptotes in the graph of $\kappa(t)$ correspond to the sharp cusps we see in the graph of $\mathbf{r}(t)$. The space curve bends most sharply as it approaches these cusps (mostly in the x-direction) and bends most gradually between these, near its intersections with the xy-plane, where $t = \pi + 2n\pi$ (n an integer). (The bending we see in the z-direction on the curve near these points is deceiving; most of the curvature occurs in the x-direction.) The curvature graph has local minima at these values of t.

37. $\mathbf{r}(t) = \langle te^t, e^{-t}, \sqrt{2}t\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle (t+1)e^t, -e^{-t}, \sqrt{2}\rangle, \quad \mathbf{r}''(t) = \langle (t+2)e^t, e^{-t}, 0\rangle.$ Then

$\mathbf{r}'(t) \times \mathbf{r}''(t) = \langle -\sqrt{2}e^{-t}, \sqrt{2}(t+2)e^t, 2t+3\rangle, \quad |\mathbf{r}'(t) \times \mathbf{r}''(t)| = \sqrt{2e^{-2t} + 2(t+2)^2e^{2t} + (2t+3)^2},$

$|\mathbf{r}'(t)| = \sqrt{(t+1)^2e^{2t} + e^{-2t} + 2}$, and $\kappa(t) = \dfrac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{\sqrt{2e^{-2t} + 2(t+2)^2e^{2t} + (2t+3)^2}}{\left[(t+1)^2e^{2t} + e^{-2t} + 2\right]^{3/2}}.$

We plot the space curve and its curvature function for $-5 \le t \le 5$ below.





From the graph of $\kappa(t)$ we see that curvature is maximized for $t = 0$, so the curve bends most sharply at the point $(0, 1, 0)$. The curve bends more gradually as we move away from this point, becoming almost linear. This is reflected in the curvature graph, where $\kappa(t)$ becomes nearly 0 as $|t|$ increases.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. Notice that the curve a is highest for the same x-values at which curve b is turning more sharply, and a is 0 or near 0 where b is nearly straight. So, a must be the graph of $y = \kappa(x)$, and b is the graph of $y = f(x)$.

39. Notice that the curve b has two inflection points at which the graph appears almost straight. We would expect the curvature to be 0 or nearly 0 at these values, but the curve a isn't near 0 there. Thus, a must be the graph of $y = f(x)$ rather than the graph of curvature, and b is the graph of $y = \kappa(x)$.

40. (a) The complete curve is given by $0 \le t \le 2\pi$. Curvature appears to have a local (or absolute) maximum at 6 points. (Look at points where the curve appears to turn more sharply.)



(b) Using a CAS, we find (after simplifying) $\kappa(t) = \dfrac{3\sqrt{2}\sqrt{(5\sin t + \sin 5t)^2}}{(9\cos 6t + 2\cos 4t + 11)^{3/2}}$. (To compute cross products in Maple, use the `VectorCalculus` or `LinearAlgebra` package and the `CrossProduct(a,b)` command; in Mathematica, use `Cross[a,b]`.) The graph shows 6 local (or absolute) maximum points for $0 \le t \le 2\pi$, as observed in part (a).



41. Using a CAS, we find (after simplifying) $\kappa(t) = \dfrac{6\sqrt{4\cos^2 t - 12\cos t + 13}}{(17 - 12\cos t)^{3/2}}$. (To compute cross products in Maple, use the `VectorCalculus` or `LinearAlgebra` package and the `CrossProduct(a,b)` command; in Mathematica, use `Cross[a,b]`.) Curvature is largest at integer multiples of 2π.



42. Here $\mathbf{r}(t) = \langle f(t), g(t)\rangle$, $\mathbf{r}'(t) = \langle f'(t), g'(t)\rangle$, $\mathbf{r}''(t) = \langle f''(t), g''(t)\rangle$,

$|\mathbf{r}'(t)|^3 = \left[\sqrt{(f'(t))^2 + (g'(t))^2}\right]^3 = [(f'(t))^2 + (g'(t))^2]^{3/2} = (\dot{x}^2 + \dot{y}^2)^{3/2}$, and

$|\mathbf{r}'(t) \times \mathbf{r}''(t)| = |\langle 0, 0, f'(t)\, g''(t) - f''(t)\, g'(t)\rangle| = \left[(\dot{x}\ddot{y} - \ddot{x}\dot{y})^2\right]^{1/2} = |\dot{x}\ddot{y} - \dot{y}\ddot{x}|$. Thus $\kappa(t) = \dfrac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}}$.

43. $x = t^2 \;\Rightarrow\; \dot{x} = 2t \;\Rightarrow\; \ddot{x} = 2$, $y = t^3 \;\Rightarrow\; \dot{y} = 3t^2 \;\Rightarrow\; \ddot{y} = 6t$.

Then $\kappa(t) = \dfrac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \dfrac{|(2t)(6t) - (3t^2)(2)|}{[(2t)^2 + (3t^2)^2]^{3/2}} = \dfrac{|12t^2 - 6t^2|}{(4t^2 + 9t^4)^{3/2}} = \dfrac{6t^2}{(4t^2 + 9t^4)^{3/2}}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

44. $x = a\cos\omega t \;\Rightarrow\; \dot{x} = -a\omega\sin\omega t \;\Rightarrow\; \ddot{x} = -a\omega^2\cos\omega t$,

$y = b\sin\omega t \;\Rightarrow\; \dot{y} = b\omega\cos\omega t \;\Rightarrow\; \ddot{y} = -b\omega^2\sin\omega t$. Then

$$\kappa(t) = \frac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \frac{\left|(-a\omega\sin\omega t)(-b\omega^2\sin\omega t) - (b\omega\cos\omega t)(-a\omega^2\cos\omega t)\right|}{[(-a\omega\sin\omega t)^2 + (b\omega\cos\omega t)^2]^{3/2}}$$

$$= \frac{|ab\omega^3\sin^2\omega t + ab\omega^3\cos^2\omega t|}{(a^2\omega^2\sin^2\omega t + b^2\omega^2\cos^2\omega t)^{3/2}} = \frac{|ab\omega^3|}{(a^2\omega^2\sin^2\omega t + b^2\omega^2\cos^2\omega t)^{3/2}}$$

45. $x = e^t\cos t \;\Rightarrow\; \dot{x} = e^t(\cos t - \sin t) \;\Rightarrow\; \ddot{x} = e^t(-\sin t - \cos t) + e^t(\cos t - \sin t) = -2e^t\sin t$,

$y = e^t\sin t \;\Rightarrow\; \dot{y} = e^t(\cos t + \sin t) \;\Rightarrow\; \ddot{y} = e^t(-\sin t + \cos t) + e^t(\cos t + \sin t) = 2e^t\cos t$. Then

$$\kappa(t) = \frac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \frac{\left|e^t(\cos t - \sin t)(2e^t\cos t) - e^t(\cos t + \sin t)(-2e^t\sin t)\right|}{([e^t(\cos t - \sin t)]^2 + [e^t(\cos t + \sin t)]^2)^{3/2}}$$

$$= \frac{\left|2e^{2t}(\cos^2 t - \sin t\cos t + \sin t\cos t + \sin^2 t)\right|}{\left[e^{2t}(\cos^2 t - 2\cos t\sin t + \sin^2 t + \cos^2 t + 2\cos t\sin t + \sin^2 t)\right]^{3/2}} = \frac{|2e^{2t}(1)|}{[e^{2t}(1+1)]^{3/2}} = \frac{2e^{2t}}{e^{3t}(2)^{3/2}} = \frac{1}{\sqrt{2}\,e^t}$$

46. $f(x) = e^{cx}$, $f'(x) = ce^{cx}$, $f''(x) = c^2e^{cx}$. Using Formula 11 we have

$\kappa(x) = \dfrac{|f''(x)|}{[1 + (f'(x))^2]^{3/2}} = \dfrac{|c^2e^{cx}|}{[1 + (ce^{cx})^2]^{3/2}} = \dfrac{c^2e^{cx}}{(1 + c^2e^{2cx})^{3/2}}$ so the curvature at $x = 0$ is

$\kappa(0) = \dfrac{c^2}{(1+c^2)^{3/2}}$. To determine the maximum value for $\kappa(0)$, let $f(c) = \dfrac{c^2}{(1+c^2)^{3/2}}$. Then

$f'(c) = \dfrac{2c\cdot(1+c^2)^{3/2} - c^2\cdot\frac{3}{2}(1+c^2)^{1/2}(2c)}{[(1+c^2)^{3/2}]^2} = \dfrac{(1+c^2)^{1/2}\left[2c(1+c^2) - 3c^3\right]}{(1+c^2)^3} = \dfrac{(2c - c^3)}{(1+c^2)^{5/2}}$. We have a critical number when $2c - c^3 = 0 \;\Rightarrow\; c(2 - c^2) = 0 \;\Rightarrow\; c = 0$ or $c = \pm\sqrt{2}$. $f'(c)$ is positive for $c < -\sqrt{2}$, $0 < c < \sqrt{2}$ and negative elsewhere, so f achieves its maximum value when $c = \sqrt{2}$ or $-\sqrt{2}$. In either case, $\kappa(0) = \dfrac{2}{3^{3/2}}$, so the members of the family with the largest value of $\kappa(0)$ are $f(x) = e^{\sqrt{2}x}$ and $f(x) = e^{-\sqrt{2}x}$.

47. $(1, \frac{2}{3}, 1)$ corresponds to $t = 1$. $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{\langle 2t, 2t^2, 1\rangle}{\sqrt{4t^2 + 4t^4 + 1}} = \dfrac{\langle 2t, 2t^2, 1\rangle}{2t^2 + 1}$, so $\mathbf{T}(1) = \langle \frac{2}{3}, \frac{2}{3}, \frac{1}{3}\rangle$.

$\mathbf{T}'(t) = -4t(2t^2+1)^{-2}\langle 2t, 2t^2, 1\rangle + (2t^2+1)^{-1}\langle 2, 4t, 0\rangle$ [by Formula 3 of Theorem 13.2.3]

$= (2t^2+1)^{-2}\langle -8t^2 + 4t^2 + 2, -8t^3 + 8t^3 + 4t, -4t\rangle = 2(2t^2+1)^{-2}\langle 1 - 2t^2, 2t, -2t\rangle$

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{2(2t^2+1)^{-2}\langle 1 - 2t^2, 2t, -2t\rangle}{2(2t^2+1)^{-2}\sqrt{(1-2t^2)^2 + (2t)^2 + (-2t)^2}} = \dfrac{\langle 1 - 2t^2, 2t, -2t\rangle}{\sqrt{1 - 4t^2 + 4t^4 + 8t^2}} = \dfrac{\langle 1 - 2t^2, 2t, -2t\rangle}{1 + 2t^2}$

$\mathbf{N}(1) = \langle -\frac{1}{3}, \frac{2}{3}, -\frac{2}{3}\rangle$ and $\mathbf{B}(1) = \mathbf{T}(1) \times \mathbf{N}(1) = \langle -\frac{4}{9} - \frac{2}{9}, -\left(-\frac{4}{9} + \frac{1}{9}\right), \frac{4}{9} + \frac{2}{9}\rangle = \langle \frac{2}{3}, \frac{1}{3}, \frac{2}{3}\rangle$.

48. $(1, 0, 0)$ corresponds to $t = 0$. $\mathbf{r}(t) = \langle \cos t, \sin t, \ln\cos t\rangle$, and in Exercise 4 we found that $\mathbf{r}'(t) = \langle -\sin t, \cos t, -\tan t\rangle$ and $|\mathbf{r}'(t)| = |\sec t|$. Here we can assume $-\frac{\pi}{2} < t < \frac{\pi}{2}$ and then $\sec t > 0 \;\Rightarrow\; |\mathbf{r}'(t)| = \sec t$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{\langle -\sin t, \cos t, -\tan t\rangle}{\sec t} = \langle -\sin t\cos t, \cos^2 t, -\sin t\rangle$ and $\mathbf{T}(0) = \langle 0, 1, 0\rangle$.

$\mathbf{T}'(t) = \langle -[(\sin t)(-\sin t) + (\cos t)(\cos t)], 2(\cos t)(-\sin t), -\cos t\rangle = \langle \sin^2 t - \cos^2 t, -2\sin t\cos t, -\cos t\rangle$, so

$\mathbf{N}(0) = \dfrac{\mathbf{T}'(0)}{|\mathbf{T}'(0)|} = \dfrac{\langle -1, 0, -1\rangle}{\sqrt{1+0+1}} = \dfrac{1}{\sqrt{2}}\langle -1, 0, -1\rangle = \left\langle -\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}}\right\rangle$.

Finally, $\mathbf{B}(0) = \mathbf{T}(0) \times \mathbf{N}(0) = \langle 0, 1, 0\rangle \times \left\langle -\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}}\right\rangle = \left\langle -\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}}\right\rangle$.

49. $\mathbf{r}(t) = \langle \sin 2t, -\cos 2t, 4t\rangle \Rightarrow \mathbf{r}'(t) = \langle 2\cos 2t, 2\sin 2t, 4\rangle$. The point $(0, 1, 2\pi)$ corresponds to $t = \pi/2$, and the normal plane there has normal vector $\mathbf{r}'(\pi/2) = \langle -2, 0, 4\rangle$. An equation for the normal plane is $-2(x-0) + 0(y-1) + 4(z-2\pi) = 0$ or $-2x + 4z = 8\pi$ or $x - 2z = -4\pi$.

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{\langle 2\cos 2t, 2\sin 2t, 4\rangle}{\sqrt{4\cos^2 2t + 4\sin^2 2t + 16}} = \dfrac{1}{2\sqrt{5}}\langle 2\cos 2t, 2\sin 2t, 4\rangle = \dfrac{1}{\sqrt{5}}\langle \cos 2t, \sin 2t, 2\rangle \Rightarrow$

$\mathbf{T}'(t) = \frac{1}{\sqrt{5}}\langle -2\sin 2t, 2\cos 2t, 0\rangle \Rightarrow |\mathbf{T}'(t)| = \frac{1}{\sqrt{5}}\sqrt{4\sin^2 2t + 4\cos^2 2t} = \frac{2}{\sqrt{5}}$, and

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \langle -\sin 2t, \cos 2t, 0\rangle$. Then $\mathbf{T}(\pi/2) = \frac{1}{\sqrt{5}}\langle -1, 0, 2\rangle$, $\mathbf{N}(\pi/2) = \langle 0, -1, 0\rangle$, and $\mathbf{B}(\pi/2) = \mathbf{T}(\pi/2) \times \mathbf{N}(\pi/2) = \frac{1}{\sqrt{5}}\langle 2, 0, 1\rangle$. Since $\mathbf{B}(\pi/2)$ is normal to the osculating plane, so is $\langle 2, 0, 1\rangle$, and an equation of the plane is $2(x-0) + 0(y-1) + 1(z-2\pi) = 0$ or $2x + z = 2\pi$.

50. $\mathbf{r}(t) = \langle \ln t, 2t, t^2\rangle \Rightarrow \mathbf{r}'(t) = \langle 1/t, 2, 2t\rangle$. The point $(0, 2, 1)$ corresponds to $t = 1$, and the normal plane there has normal vector $\mathbf{r}'(1) = \langle 1, 2, 2\rangle$. An equation for the normal plane is $1(x-0) + 2(y-2) + 2(z-1) = 0$ or $x + 2y + 2z = 6$.

$|\mathbf{r}'(t)| = \sqrt{1/t^2 + 4 + 4t^2} = \sqrt{[(1/t) + 2t]^2} = (1/t) + 2t$ [since $t > 0$] and then

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{\langle 1/t, 2, 2t\rangle}{(1/t) + 2t} = \dfrac{1}{1+2t^2}\langle 1, 2t, 2t^2\rangle$ $\left[\text{after multiplying by } \dfrac{t}{t}\right]$. By Formula 3 of Theorem 13.2.3,

$$\begin{aligned}\mathbf{T}'(t) &= -\frac{4t}{(1+2t^2)^2}\langle 1, 2t, 2t^2\rangle + \frac{1}{1+2t^2}\langle 0, 2, 4t\rangle \\ &= \frac{1}{(1+2t^2)^2}\langle -4t, -8t^2 + 2(1+2t^2), -8t^3 + 4t(1+2t^2)\rangle = \frac{1}{(1+2t^2)^2}\langle -4t, 2-4t^2, 4t\rangle\end{aligned}$$

Then

$$\begin{aligned}|\mathbf{T}'(t)| &= \frac{1}{(1+2t^2)^2}\sqrt{16t^2 + (2-4t^2)^2 + 16t^2} = \frac{1}{(1+2t^2)^2}\sqrt{16t^2 + 4 + 16t^4} \\ &= \frac{1}{(1+2t^2)^2}\cdot 2\sqrt{(1+2t^2)^2} = \frac{2}{1+2t^2}\end{aligned}$$

and $\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{1}{2(1+2t^2)}\langle -4t, 2-4t^2, 4t\rangle = \dfrac{1}{1+2t^2}\langle -2t, 1-2t^2, 2t\rangle$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Thus $\mathbf{T}(1) = \frac{1}{3}\langle 1,2,2\rangle$, $\mathbf{N}(1) = \frac{1}{3}\langle -2,-1,2\rangle$, and $\mathbf{B}(1) = \mathbf{T}(1) \times \mathbf{N}(1) = \frac{1}{9}\langle 6,-6,3\rangle$ is normal to the osculating plane. We can take the parallel vector $\langle 2,-2,1\rangle$ as a normal vector for the plane, so an equation is

$2(x-0) - 2(y-2) + 1(z-1) = 0$ or $2x - 2y + z = -3$.

Note: Since $\mathbf{r}'(1)$ is parallel to $\mathbf{T}(1)$ and $\mathbf{T}'(1)$ is parallel to $\mathbf{N}(1)$, we could have taken $\mathbf{r}'(1) \times \mathbf{T}'(1)$ as a normal vector for the plane.

51. The ellipse is given by the parametric equations $x = 2\cos t,\ y = 3\sin t$, so using the result from Exercise 42,

$$\kappa(t) = \frac{|\dot{x}\ddot{y} - \ddot{x}\dot{y}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \frac{|(-2\sin t)(-3\sin t) - (3\cos t)(-2\cos t)|}{(4\sin^2 t + 9\cos^2 t)^{3/2}} = \frac{6}{(4\sin^2 t + 9\cos^2 t)^{3/2}}.$$

At $(2,0)$, $t = 0$. Now $\kappa(0) = \frac{6}{27} = \frac{2}{9}$, so the radius of the osculating circle is $1/\kappa(0) = \frac{9}{2}$ and its center is $\left(-\frac{5}{2}, 0\right)$. Its equation is therefore $\left(x + \frac{5}{2}\right)^2 + y^2 = \frac{81}{4}$.

At $(0,3)$, $t = \frac{\pi}{2}$, and $\kappa\left(\frac{\pi}{2}\right) = \frac{6}{8} = \frac{3}{4}$. So the radius of the osculating circle is $\frac{4}{3}$ and its center is $\left(0, \frac{5}{3}\right)$. Hence its equation is $x^2 + \left(y - \frac{5}{3}\right)^2 = \frac{16}{9}$.



52. $y = \frac{1}{2}x^2 \ \Rightarrow\ y' = x$ and $y'' = 1$, so Formula 11 gives $\kappa(x) = \dfrac{1}{(1+x^2)^{3/2}}$. So the curvature at $(0,0)$ is $\kappa(0) = 1$ and the osculating circle has radius 1 and center $(0,1)$, and hence equation $x^2 + (y-1)^2 = 1$. The curvature at $\left(1, \frac{1}{2}\right)$ is $\kappa(1) = \dfrac{1}{(1+1^2)^{3/2}} = \dfrac{1}{2\sqrt{2}}$. The tangent line to the parabola at $\left(1, \frac{1}{2}\right)$ has slope 1, so the normal line has slope -1. Thus the center of the osculating circle lies in the direction of the unit vector $\left\langle -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right\rangle$. The circle has radius $2\sqrt{2}$, so its center has position vector $\left\langle 1, \frac{1}{2}\right\rangle + 2\sqrt{2}\left\langle -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right\rangle = \left\langle -1, \frac{5}{2}\right\rangle$. So the equation of the circle is $(x+1)^2 + \left(y - \frac{5}{2}\right)^2 = 8$.



53. Here $\mathbf{r}(t) = \langle t^3, 3t, t^4\rangle$, and $\mathbf{r}'(t) = \langle 3t^2, 3, 4t^3\rangle$ is normal to the normal plane for any t. The given plane has normal vector $\langle 6,6,-8\rangle$, and the planes are parallel when their normal vectors are parallel. Thus we need to find a value for t where $\langle 3t^2, 3, 4t^3\rangle = k\langle 6,6,-8\rangle$ for some $k \neq 0$. From the y-component we see that $k = \frac{1}{2}$, and $\langle 3t^2, 3, 4t^3\rangle = \frac{1}{2}\langle 6,6,-8\rangle = \langle 3,3,-4\rangle$ for $t = -1$. Thus the planes are parallel at the point $(-1,-3,1)$.

54. To find the osculating plane, we first calculate the unit tangent and normal vectors.

In Maple, we use the `VectorCalculus` package and set `r:=<t^3,3*t,t^4>;`. After differentiating, the `Normalize` command converts the tangent vector to the unit tangent vector: `T:=Normalize(diff(r,t));`. After

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

simplifying, we find that $\mathbf{T}(t) = \dfrac{\langle 3t^2, 3, 4t^3 \rangle}{\sqrt{16t^6 + 9t^4 + 9}}$. We use a similar procedure to compute the unit normal vector, `N:=Normalize(diff(T,t));`. After simplifying, we have $\mathbf{N}(t) = \dfrac{\langle -t(8t^6 - 9), -3t^3(3 + 8t^2), 6t^2(t^4 + 3) \rangle}{\sqrt{t^2(4t^6 + 36t^2 + 9)(16t^6 + 9t^4 + 9)}}$. Then we use the command `B:=CrossProduct(T,N);`. After simplification, we find that $\mathbf{B}(t) = \dfrac{\langle 6t^2, -2t^4, -3t \rangle}{\sqrt{t^2(4t^6 + 36t^2 + 9)}}$.

In Mathematica, we define the vector function `r={t^3, 3*t, t^4}` and use the command `Dt` to differentiate. We find $\mathbf{T}(t)$ by dividing the result by its magnitude, computed using the `Norm` command. (You may wish to include the option `Element[t,Reals]` to obtain simpler expressions.) $\mathbf{N}(t)$ is found similarly, and we use `Cross[T,N]` to find $\mathbf{B}(t)$.

Now $\mathbf{B}(t)$ is parallel to $\langle 6t^2, -2t^4, -3t \rangle$, so if $\mathbf{B}(t)$ is parallel to $\langle 1, 1, 1 \rangle$ for some $t \neq 0$ [since $\mathbf{B}(0) = \mathbf{0}$], then $\langle 6t^2, -2t^4, -3t \rangle = k \langle 1, 1, 1 \rangle$ for some value of k. But then $6t^2 = -2t^4 = -3t$ which has no solution for $t \neq 0$. So there is no such osculating plane.

55. First we parametrize the curve of intersection. We can choose $y = t$; then $x = y^2 = t^2$ and $z = x^2 = t^4$, and the curve is given by $\mathbf{r}(t) = \langle t^2, t, t^4 \rangle$. $\mathbf{r}'(t) = \langle 2t, 1, 4t^3 \rangle$ and the point $(1, 1, 1)$ corresponds to $t = 1$, so $\mathbf{r}'(1) = \langle 2, 1, 4 \rangle$ is a normal vector for the normal plane. Thus an equation of the normal plane is

$2(x - 1) + 1(y - 1) + 4(z - 1) = 0$ or $2x + y + 4z = 7$. $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{4t^2 + 1 + 16t^6}} \langle 2t, 1, 4t^3 \rangle$ and

$\mathbf{T}'(t) = -\frac{1}{2}(4t^2 + 1 + 16t^6)^{-3/2}(8t + 96t^5) \langle 2t, 1, 4t^3 \rangle + (4t^2 + 1 + 16t^6)^{-1/2} \langle 2, 0, 12t^2 \rangle$. A normal vector for the osculating plane is $\mathbf{B}(1) = \mathbf{T}(1) \times \mathbf{N}(1)$, but $\mathbf{r}'(1) = \langle 2, 1, 4 \rangle$ is parallel to $\mathbf{T}(1)$ and $\mathbf{T}'(1) = -\frac{1}{2}(21)^{-3/2}(104)\langle 2, 1, 4 \rangle + (21)^{-1/2}\langle 2, 0, 12 \rangle = \frac{2}{21\sqrt{21}} \langle -31, -26, 22 \rangle$ is parallel to $\mathbf{N}(1)$ as is $\langle -31, -26, 22 \rangle$, so $\langle 2, 1, 4 \rangle \times \langle -31, -26, 22 \rangle = \langle 126, -168, -21 \rangle$ is normal to the osculating plane. Thus an equation for the osculating plane is $126(x - 1) - 168(y - 1) - 21(z - 1) = 0$ or $6x - 8y - z = -3$.

56. $\mathbf{r}(t) = \langle t + 2, 1 - t, \frac{1}{2}t^2 \rangle \Rightarrow \mathbf{r}'(t) = \langle 1, -1, t \rangle$, $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{2 + t^2}} \langle 1, -1, t \rangle$,

$$\begin{aligned} \mathbf{T}'(t) &= -\tfrac{1}{2}(2 + t^2)^{-3/2}(2t)\langle 1, -1, t \rangle + (2 + t^2)^{1/2} \langle 0, 0, 1 \rangle \\ &= -(2 + t^2)^{-3/2} \left[t\langle 1, -1, t \rangle - (2 + t^2)\langle 0, 0, 1 \rangle \right] = \tfrac{-1}{(2+t^2)^{3/2}} \langle t, -t, -2 \rangle \end{aligned}$$

A normal vector for the osculating plane is $\mathbf{B}(t) = \mathbf{T}(t) \times \mathbf{N}(t)$, but $\mathbf{r}'(t) = \langle 1, -1, t \rangle$ is parallel to $\mathbf{T}(t)$ and $\langle t, -t, -2 \rangle$ is parallel to $\mathbf{T}'(t)$ and hence parallel to $\mathbf{N}(t)$, so $\langle 1, -1, t \rangle \times \langle t, -t, -2 \rangle = \langle t^2 + 2, t^2 + 2, 0 \rangle$ is normal to the osculating plane for any t. All such vectors are parallel to $\langle 1, 1, 0 \rangle$, so at any point $\left(t + 2, 1 - t, \frac{1}{2}t^2\right)$ on the curve, an equation for the osculating plane is $1[x - (t + 2)] + 1[y - (1 - t)] + 0\left(z - \frac{1}{2}t^2\right) = 0$ or $x + y = 3$. Because the osculating plane at every point on the curve is the same, we can conclude that the curve itself lies in that same plane. In fact, we can easily verify that the parametric equations of the curve satisfy $x + y = 3$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

57. $\mathbf{r}(t) = \langle e^t \cos t, e^t \sin t, e^t \rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \langle e^t(\cos t - \sin t), e^t(\cos t + \sin t), e^t \rangle$ so

$$|\mathbf{r}'(t)| = \sqrt{e^{2t}(\cos t - \sin t)^2 + e^{2t}(\cos t + \sin t)^2 + e^{2t}}$$
$$= \sqrt{e^{2t}\left[2(\cos^2 t + \sin^2 t) - 2\cos t \sin t + 2\cos t \sin t + 1\right]} = \sqrt{3e^{2t}} = \sqrt{3}\, e^t$$

and $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{3}\, e^t} \langle e^t(\cos t - \sin t), e^t(\cos t + \sin t), e^t \rangle = \dfrac{1}{\sqrt{3}} \langle \cos t - \sin t, \cos t + \sin t, 1 \rangle$. The vector $\mathbf{k} = \langle 0, 0, 1 \rangle$ is parallel to the z-axis, so for any t, the angle α between $\mathbf{T}(t)$ and the z-axis is given by

$\cos\alpha = \dfrac{\mathbf{T}(t) \cdot \mathbf{k}}{|\mathbf{T}(t)|\,|\mathbf{k}|} = \dfrac{\frac{1}{\sqrt{3}} \langle \cos t - \sin t, \cos t + \sin t, 1 \rangle \cdot \langle 0, 0, 1 \rangle}{\frac{1}{\sqrt{3}} \sqrt{(\cos t - \sin t)^2 + (\cos t + \sin t)^2 + 1}\, \sqrt{1}} = \dfrac{1}{\sqrt{2(\cos^2 t + \sin^2 t) + 1}} = \dfrac{1}{\sqrt{3}}$. Thus the angle is constant; specifically, $\alpha = \cos^{-1}(1/\sqrt{3}) \approx 54.7°$.

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{(1/\sqrt{3}) \langle -\sin t - \cos t, -\sin t + \cos t, 0 \rangle}{(1/\sqrt{3}) \sqrt{2\,(\sin^2 t + \cos^2 t)}} = \dfrac{1}{\sqrt{2}} \langle -\sin t - \cos t, -\sin t + \cos t, 0 \rangle$, and the angle β made with the z-axis is given by $\cos\beta = \dfrac{\mathbf{N}(t) \cdot \mathbf{k}}{|\mathbf{N}(t)|\,|\mathbf{k}|} = 0$, so $\beta = 90°$.

$\mathbf{B}(t) = \mathbf{T}(t) \times \mathbf{N}(t) = \frac{1}{\sqrt{6}} \langle \sin t - \cos t, -\sin t - \cos t, 2 \rangle$ and the angle γ made with the z-axis is given by

$\cos\gamma = \dfrac{\mathbf{B}(t) \cdot \mathbf{k}}{|\mathbf{B}(t)|\,|\mathbf{k}|} = \dfrac{\frac{1}{\sqrt{6}} \langle \sin t - \cos t, -\sin t - \cos t, 2 \rangle \cdot \langle 0, 0, 1 \rangle}{\frac{1}{\sqrt{6}} \sqrt{(\sin t - \cos t)^2 + (-\sin t - \cos t)^2 + 4}\, \sqrt{1}} = \dfrac{2}{\sqrt{6}}$ or equivalently $\dfrac{\sqrt{6}}{3}$. Again the angle is constant; specifically, $\gamma = \cos^{-1}(2/\sqrt{6}) \approx 35.3°$.

58. If vectors $\mathbf{T}$ and $\mathbf{B}$ lie in the rectifying plane then $\mathbf{N}$ is a normal vector for the plane, as it is orthogonal to both $\mathbf{T}$ and $\mathbf{B}$. The point $(\sqrt{2}/2, \sqrt{2}/2, 1)$ corresponds to $t = \pi/4$, so we can take $\mathbf{T}'(\pi/4)$ as a normal vector for the plane [since it is parallel to $\mathbf{N}(\pi/4)$]. $\mathbf{r}(t) = \sin t\,\mathbf{i} + \cos t\,\mathbf{j} + \tan t\,\mathbf{k} \quad \Rightarrow \quad \mathbf{r}'(t) = \cos t\,\mathbf{i} - \sin t\,\mathbf{j} + \sec^2 t\,\mathbf{k}$ and $|\mathbf{r}'(t)| = \sqrt{\cos^2 t + \sin^2 t + \sec^4 t} = \sqrt{1 + \sec^4 t}$. Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{1 + \sec^4 t}} \left(\cos t\,\mathbf{i} - \sin t\,\mathbf{j} + \sec^2 t\,\mathbf{k}\right)$. By Formula 3 of Theorem 13.2.3,

$$\mathbf{T}'(t) = -\frac{2\sec^4 t \tan t}{(1 + \sec^4 t)^{3/2}} \left(\cos t\,\mathbf{i} - \sin t\,\mathbf{j} + \sec^2 t\,\mathbf{k}\right) + \frac{1}{\sqrt{1 + \sec^4 t}} \left(-\sin t\,\mathbf{i} - \cos t\,\mathbf{j} + 2\sec^2 t \tan t\,\mathbf{k}\right) \text{ and}$$

$$\mathbf{T}'(\pi/4) = \frac{2(\sqrt{2})^4(1)}{[1 + (\sqrt{2})^4]^{3/2}} \left(\frac{\sqrt{2}}{2}\,\mathbf{i} - \frac{\sqrt{2}}{2}\,\mathbf{j} + (\sqrt{2})^2\,\mathbf{k}\right) + \frac{1}{\sqrt{1 + (\sqrt{2})^4}} \left(-\frac{\sqrt{2}}{2}\,\mathbf{i} - \frac{\sqrt{2}}{2}\,\mathbf{j} + 2(\sqrt{2})^2(1)\,\mathbf{k}\right)$$
$$= -\frac{8}{5\sqrt{5}} \left(\frac{\sqrt{2}}{2}\,\mathbf{i} - \frac{\sqrt{2}}{2}\,\mathbf{j} + 2\,\mathbf{k}\right) + \frac{1}{\sqrt{5}} \left(-\frac{\sqrt{2}}{2}\,\mathbf{i} - \frac{\sqrt{2}}{2}\,\mathbf{j} + 4\,\mathbf{k}\right) = -\frac{13\sqrt{2}}{10\sqrt{5}}\,\mathbf{i} + \frac{3\sqrt{2}}{10\sqrt{5}}\,\mathbf{j} + \frac{4}{5\sqrt{5}}\,\mathbf{k}$$

We can take the parallel vector $-13\sqrt{2}\,\mathbf{i} + 3\sqrt{2}\,\mathbf{j} + 8\,\mathbf{k}$ as a normal for the plane, so an equation for the plane is

$$-13\sqrt{2}\left(x - \tfrac{\sqrt{2}}{2}\right) + 3\sqrt{2}\left(y - \tfrac{\sqrt{2}}{2}\right) + 8\,(z - 1) = 0 \text{ or } -13\sqrt{2}\,x + 3\sqrt{2}\,y + 8z = -2 \text{ or } 13x - 3y - 4\sqrt{2}\,z = \sqrt{2}.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $\kappa = \left|\dfrac{d\mathbf{T}}{ds}\right| = \left|\dfrac{d\mathbf{T}/dt}{ds/dt}\right| = \dfrac{|d\mathbf{T}/dt|}{ds/dt}$ and $\mathbf{N} = \dfrac{d\mathbf{T}/dt}{|d\mathbf{T}/dt|}$, so $\kappa\mathbf{N} = \dfrac{\left|\dfrac{d\mathbf{T}}{dt}\right|\dfrac{d\mathbf{T}}{dt}}{\left|\dfrac{d\mathbf{T}}{dt}\right|\dfrac{ds}{dt}} = \dfrac{d\mathbf{T}/dt}{ds/dt} = \dfrac{d\mathbf{T}}{ds}$ by the Chain Rule.

60. For a plane curve, $\mathbf{T} = |\mathbf{T}|\cos\phi\,\mathbf{i} + |\mathbf{T}|\sin\phi\,\mathbf{j} = \cos\phi\,\mathbf{i} + \sin\phi\,\mathbf{j}$. Then

$\dfrac{d\mathbf{T}}{ds} = \left(\dfrac{d\mathbf{T}}{d\phi}\right)\left(\dfrac{d\phi}{ds}\right) = (-\sin\phi\,\mathbf{i} + \cos\phi\,\mathbf{j})\left(\dfrac{d\phi}{ds}\right)$ and $\left|\dfrac{d\mathbf{T}}{ds}\right| = |-\sin\phi\,\mathbf{i} + \cos\phi\,\mathbf{j}|\left|\dfrac{d\phi}{ds}\right| = \left|\dfrac{d\phi}{ds}\right|$. Hence for a plane curve, the curvature is $\kappa = |d\phi/ds|$.

61. (a) $|\mathbf{B}| = 1 \;\Rightarrow\; \mathbf{B}\cdot\mathbf{B} = 1 \;\Rightarrow\; \dfrac{d}{ds}(\mathbf{B}\cdot\mathbf{B}) = 0 \;\Rightarrow\; 2\dfrac{d\mathbf{B}}{ds}\cdot\mathbf{B} = 0 \;\Rightarrow\; \dfrac{d\mathbf{B}}{ds} \perp \mathbf{B}$

(b) $\mathbf{B} = \mathbf{T}\times\mathbf{N} \;\Rightarrow$

$$\frac{d\mathbf{B}}{ds} = \frac{d}{ds}(\mathbf{T}\times\mathbf{N}) = \frac{d}{dt}(\mathbf{T}\times\mathbf{N})\frac{1}{ds/dt} = \frac{d}{dt}(\mathbf{T}\times\mathbf{N})\frac{1}{|\mathbf{r}'(t)|} = [(\mathbf{T}'\times\mathbf{N}) + (\mathbf{T}\times\mathbf{N}')]\frac{1}{|\mathbf{r}'(t)|}$$

$$= \left[\left(\mathbf{T}'\times\frac{\mathbf{T}'}{|\mathbf{T}'|}\right) + (\mathbf{T}\times\mathbf{N}')\right]\frac{1}{|\mathbf{r}'(t)|} = \frac{\mathbf{T}\times\mathbf{N}'}{|\mathbf{r}'(t)|} \;\Rightarrow\; \frac{d\mathbf{B}}{ds} \perp \mathbf{T}$$

(c) $\mathbf{B} = \mathbf{T}\times\mathbf{N} \;\Rightarrow\; \mathbf{T}\perp\mathbf{N}, \mathbf{B}\perp\mathbf{T}$ and $\mathbf{B}\perp\mathbf{N}$. So $\mathbf{B}$, $\mathbf{T}$ and $\mathbf{N}$ form an orthogonal set of vectors in the three-dimensional space $\mathbb{R}^3$. From parts (a) and (b), $d\mathbf{B}/ds$ is perpendicular to both $\mathbf{B}$ and $\mathbf{T}$, so $d\mathbf{B}/ds$ is parallel to $\mathbf{N}$. Therefore, $d\mathbf{B}/ds = -\tau(s)\mathbf{N}$, where $\tau(s)$ is a scalar.

(d) Since $\mathbf{B} = \mathbf{T}\times\mathbf{N}$, $\mathbf{T}\perp\mathbf{N}$ and both $\mathbf{T}$ and $\mathbf{N}$ are unit vectors, $\mathbf{B}$ is a unit vector mutually perpendicular to both $\mathbf{T}$ and $\mathbf{N}$. For a plane curve, $\mathbf{T}$ and $\mathbf{N}$ always lie in the plane of the curve, so that $\mathbf{B}$ is a constant unit vector always perpendicular to the plane. Thus $d\mathbf{B}/ds = \mathbf{0}$, but $d\mathbf{B}/ds = -\tau(s)\mathbf{N}$ and $\mathbf{N}\neq\mathbf{0}$, so $\tau(s) = 0$.

62. $\mathbf{N} = \mathbf{B}\times\mathbf{T} \;\Rightarrow$

$$\frac{d\mathbf{N}}{ds} = \frac{d}{ds}(\mathbf{B}\times\mathbf{T}) = \frac{d\mathbf{B}}{ds}\times\mathbf{T} + \mathbf{B}\times\frac{d\mathbf{T}}{ds} \qquad \text{[by Formula 5 of Theorem 13.2.3]}$$

$$= -\tau\mathbf{N}\times\mathbf{T} + \mathbf{B}\times\kappa\mathbf{N} \qquad \text{[by Formulas 3 and 1]}$$

$$= -\tau(\mathbf{N}\times\mathbf{T}) + \kappa(\mathbf{B}\times\mathbf{N}) \qquad \text{[by Property 2 of Theorem 12.4.11]}$$

But $\mathbf{B}\times\mathbf{N} = \mathbf{B}\times(\mathbf{B}\times\mathbf{T}) = (\mathbf{B}\cdot\mathbf{T})\mathbf{B} - (\mathbf{B}\cdot\mathbf{B})\mathbf{T}$ [by Property 6 of Theorem 12.4.11] $= -\mathbf{T} \;\Rightarrow$ $d\mathbf{N}/ds = \tau(\mathbf{T}\times\mathbf{N}) - \kappa\mathbf{T} = -\kappa\,\mathbf{T} + \tau\,\mathbf{B}$.

63. (a) $\mathbf{r}' = s'\,\mathbf{T} \;\Rightarrow\; \mathbf{r}'' = s''\,\mathbf{T} + s'\,\mathbf{T}' = s''\,\mathbf{T} + s'\dfrac{d\mathbf{T}}{ds}s' = s''\,\mathbf{T} + \kappa(s')^2\,\mathbf{N}$ by the first Serret-Frenet formula.

(b) Using part (a), we have

$$\mathbf{r}'\times\mathbf{r}'' = (s'\,\mathbf{T})\times[s''\,\mathbf{T} + \kappa(s')^2\,\mathbf{N}]$$

$$= [(s'\,\mathbf{T})\times(s''\,\mathbf{T})] + [(s'\mathbf{T})\times(\kappa(s')^2\,\mathbf{N})] \qquad \text{[by Property 3 of Theorem 12.4.11]}$$

$$= (s's'')(\mathbf{T}\times\mathbf{T}) + \kappa(s')^3(\mathbf{T}\times\mathbf{N}) = \mathbf{0} + \kappa(s')^3\,\mathbf{B} = \kappa(s')^3\,\mathbf{B}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c) Using part (a), we have

$$\begin{aligned}\mathbf{r}''' &= [s''\,\mathbf{T} + \kappa(s')^2\,\mathbf{N}]' = s'''\,\mathbf{T} + s''\,\mathbf{T}' + \kappa'(s')^2\,\mathbf{N} + 2\kappa s's''\,\mathbf{N} + \kappa(s')^2\,\mathbf{N}' \\ &= s'''\,\mathbf{T} + s''\frac{d\mathbf{T}}{ds}\,s' + \kappa'(s')^2\,\mathbf{N} + 2\kappa s's''\,\mathbf{N} + \kappa(s')^2\,\frac{d\mathbf{N}}{ds}\,s' \\ &= s'''\,\mathbf{T} + s''s'\kappa\,\mathbf{N} + \kappa'(s')^2\,\mathbf{N} + 2\kappa s's''\,\mathbf{N} + \kappa(s')^3(-\kappa\,\mathbf{T} + \tau\,\mathbf{B}) \qquad \text{[by the second formula]} \\ &= [s''' - \kappa^2(s')^3]\,\mathbf{T} + [3\kappa s's'' + \kappa'(s')^2]\,\mathbf{N} + \kappa\tau(s')^3\,\mathbf{B}\end{aligned}$$

(d) Using parts (b) and (c) and the facts that $\mathbf{B}\cdot\mathbf{T} = 0$, $\mathbf{B}\cdot\mathbf{N} = 0$, and $\mathbf{B}\cdot\mathbf{B} = 1$, we get

$$\frac{(\mathbf{r}'\times\mathbf{r}'')\cdot\mathbf{r}'''}{|\mathbf{r}'\times\mathbf{r}''|^2} = \frac{\kappa(s')^3\,\mathbf{B}\cdot\{[s''' - \kappa^2(s')^3]\,\mathbf{T} + [3\kappa s's'' + \kappa'(s')^2]\,\mathbf{N} + \kappa\tau(s')^3\,\mathbf{B}\}}{|\kappa(s')^3\,\mathbf{B}|^2} = \frac{\kappa(s')^3\kappa\tau(s')^3}{[\kappa(s')^3]^2} = \tau.$$

64. First we find the quantities required to compute κ:

$$\mathbf{r}'(t) = \langle -a\sin t, a\cos t, b\rangle \;\Rightarrow\; \mathbf{r}''(t) = \langle -a\cos t, -a\sin t, 0\rangle \;\Rightarrow\; \mathbf{r}'''(t) = \langle a\sin t, -a\cos t, 0\rangle$$

$$|\mathbf{r}'(t)| = \sqrt{(-a\sin t)^2 + (a\cos t)^2 + b^2} = \sqrt{a^2 + b^2}$$

$$\mathbf{r}'(t)\times\mathbf{r}''(t) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a\sin t & a\cos t & b \\ -a\cos t & -a\sin t & 0 \end{vmatrix} = ab\sin t\,\mathbf{i} - ab\cos t\,\mathbf{j} + a^2\,\mathbf{k}$$

$$|\mathbf{r}'(t)\times\mathbf{r}''(t)| = \sqrt{(ab\sin t)^2 + (-ab\cos t)^2 + (a^2)^2} = \sqrt{a^2b^2 + a^4}$$

$$(\mathbf{r}'(t)\times\mathbf{r}''(t))\cdot\mathbf{r}'''(t) = (ab\sin t)(a\sin t) + (-ab\cos t)(-a\cos t) + (a^2)(0) = a^2b$$

Then by Theorem 10, $\kappa(t) = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{\sqrt{a^2b^2+a^4}}{\left(\sqrt{a^2+b^2}\right)^3} = \dfrac{a\sqrt{a^2+b^2}}{\left(\sqrt{a^2+b^2}\right)^3} = \dfrac{a}{a^2+b^2}$ which is a constant.

From Exercise 63(d), the torsion τ is given by $\tau = \dfrac{(\mathbf{r}'\times\mathbf{r}'')\cdot\mathbf{r}'''}{|\mathbf{r}'\times\mathbf{r}''|^2} = \dfrac{a^2b}{\left(\sqrt{a^2b^2+a^4}\right)^2} = \dfrac{b}{a^2+b^2}$ which is also a constant.

65. $\mathbf{r} = \left\langle t, \frac{1}{2}t^2, \frac{1}{3}t^3\right\rangle \;\Rightarrow\; \mathbf{r}' = \langle 1, t, t^2\rangle,\ \mathbf{r}'' = \langle 0, 1, 2t\rangle,\ \mathbf{r}''' = \langle 0, 0, 2\rangle \;\Rightarrow\; \mathbf{r}'\times\mathbf{r}'' = \langle t^2, -2t, 1\rangle \;\Rightarrow$

$$\tau = \frac{(\mathbf{r}'\times\mathbf{r}'')\cdot\mathbf{r}'''}{|\mathbf{r}'\times\mathbf{r}''|^2} = \frac{\langle t^2, -2t, 1\rangle\cdot\langle 0, 0, 2\rangle}{t^4 + 4t^2 + 1} = \frac{2}{t^4 + 4t^2 + 1}$$

66. $\mathbf{r} = \langle \sinh t, \cosh t, t\rangle \;\Rightarrow\; \mathbf{r}' = \langle \cosh t, \sinh t, 1\rangle, \mathbf{r}'' = \langle \sinh t, \cosh t, 0\rangle, \mathbf{r}''' = \langle \cosh t, \sinh t, 0\rangle \;\Rightarrow$

$\mathbf{r}'\times\mathbf{r}'' = \langle -\cosh t, \sinh t, \cosh^2 t - \sinh^2 t\rangle = \langle -\cosh t, \sinh t, 1\rangle \;\Rightarrow$

$$\kappa = \frac{|\mathbf{r}'\times\mathbf{r}''|}{|\mathbf{r}'|^3} = \frac{|\langle -\cosh t, \sinh t, 1\rangle|}{|\langle \cosh t, \sinh t, 1\rangle|^3} = \frac{\sqrt{\cosh^2 t + \sinh^2 t + 1}}{(\cosh^2 t + \sinh^2 t + 1)^{3/2}} = \frac{1}{\cosh^2 t + \sinh^2 t + 1} = \frac{1}{2\cosh^2 t},$$

$$\tau = \frac{(\mathbf{r}'\times\mathbf{r}'')\cdot\mathbf{r}'''}{|\mathbf{r}'\times\mathbf{r}''|^2} = \frac{\langle -\cosh t, \sinh t, 1\rangle\cdot\langle \cosh t, \sinh t, 0\rangle}{\cosh^2 t + \sinh^2 t + 1} = \frac{-\cosh^2 t + \sinh^2 t}{2\cosh^2 t} = \frac{-1}{2\cosh^2 t}$$

So at the point $(0, 1, 0)$, $t = 0$, and $\kappa = \frac{1}{2}$ and $\tau = -\frac{1}{2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. For one helix, the vector equation is $\mathbf{r}(t) = \langle 10\cos t, 10\sin t, 34t/(2\pi)\rangle$ (measuring in angstroms), because the radius of each helix is 10 angstroms, and z increases by 34 angstroms for each increase of 2π in t. Using the arc length formula, letting t go from 0 to $2.9\times10^8\times2\pi$, we find the approximate length of each helix to be

$$L = \int_0^{2.9\times10^8\times2\pi} |\mathbf{r}'(t)|\,dt = \int_0^{2.9\times10^8\times2\pi} \sqrt{(-10\sin t)^2 + (10\cos t)^2 + \left(\tfrac{34}{2\pi}\right)^2}\,dt = \sqrt{100+\left(\tfrac{34}{2\pi}\right)^2}\,t\Big]_0^{2.9\times10^8\times2\pi}$$

$$= 2.9\times10^8\times2\pi\sqrt{100+\left(\tfrac{34}{2\pi}\right)^2} \approx 2.07\times10^{10}\ \text{Å} \text{ — more than two meters!}$$

68. (a) For the function $F(x) = \begin{cases} 0 & \text{if } x<0 \\ P(x) & \text{if } 0<x<1 \\ 1 & \text{if } x\ge 1 \end{cases}$ to be continuous, we must have $P(0)=0$ and $P(1)=1$.

For F' to be continuous, we must have $P'(0)=P'(1)=0$. The curvature of the curve $y=F(x)$ at the point $(x,F(x))$ is $\kappa(x) = \dfrac{|F''(x)|}{\left(1+[F'(x)]^2\right)^{3/2}}$. For $\kappa(x)$ to be continuous, we must have $P''(0)=P''(1)=0$.

Write $P(x) = ax^5+bx^4+cx^3+dx^2+ex+f$. Then $P'(x) = 5ax^4+4bx^3+3cx^2+2dx+e$ and $P''(x) = 20ax^3+12bx^2+6cx+2d$. Our six conditions are:

$$P(0)=0 \Rightarrow f=0 \quad \textbf{(1)} \qquad P(1)=1 \Rightarrow a+b+c+d+e+f=1 \quad \textbf{(2)}$$

$$P'(0)=0 \Rightarrow e=0 \quad \textbf{(3)} \qquad P'(1)=0 \Rightarrow 5a+4b+3c+2d+e=0 \quad \textbf{(4)}$$

$$P''(0)=0 \Rightarrow d=0 \quad \textbf{(5)} \qquad P''(1)=0 \Rightarrow 20a+12b+6c+2d=0 \quad \textbf{(6)}$$

From **(1)**, **(3)**, and **(5)**, we have $d=e=f=0$. Thus **(2)**, **(4)** and **(6)** become **(7)** $a+b+c=1$, **(8)** $5a+4b+3c=0$, and **(9)** $10a+6b+3c=0$. Subtracting **(8)** from **(9)** gives **(10)** $5a+2b=0$. Multiplying **(7)** by 3 and subtracting from **(8)** gives **(11)** $2a+b=-3$. Multiplying **(11)** by 2 and subtracting from **(10)** gives $a=6$. By **(10)**, $b=-15$. By **(7)**, $c=10$. Thus, $P(x) = 6x^5-15x^4+10x^3$.

(b)



13.4 Motion in Space: Velocity and Acceleration

1. (a) If $\mathbf{r}(t) = x(t)\,\mathbf{i} + y(t)\,\mathbf{j} + z(t)\,\mathbf{k}$ is the position vector of the particle at time t, then the average velocity over the time interval $[0,1]$ is

$$\mathbf{v}_{\text{ave}} = \frac{\mathbf{r}(1)-\mathbf{r}(0)}{1-0} = \frac{(4.5\,\mathbf{i}+6.0\,\mathbf{j}+3.0\,\mathbf{k})-(2.7\,\mathbf{i}+9.8\,\mathbf{j}+3.7\,\mathbf{k})}{1} = 1.8\,\mathbf{i}-3.8\,\mathbf{j}-0.7\,\mathbf{k}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Similarly, over the other intervals we have

$$[0.5,1]: \quad \mathbf{v}_{\text{ave}} = \frac{\mathbf{r}(1)-\mathbf{r}(0.5)}{1-0.5} = \frac{(4.5\mathbf{i}+6.0\mathbf{j}+3.0\mathbf{k})-(3.5\mathbf{i}+7.2\mathbf{j}+3.3\mathbf{k})}{0.5} = 2.0\mathbf{i}-2.4\mathbf{j}-0.6\mathbf{k}$$

$$[1,2]: \quad \mathbf{v}_{\text{ave}} = \frac{\mathbf{r}(2)-\mathbf{r}(1)}{2-1} = \frac{(7.3\mathbf{i}+7.8\mathbf{j}+2.7\mathbf{k})-(4.5\mathbf{i}+6.0\mathbf{j}+3.0\mathbf{k})}{1} = 2.8\mathbf{i}+1.8\mathbf{j}-0.3\mathbf{k}$$

$$[1,1.5]: \quad \mathbf{v}_{\text{ave}} = \frac{\mathbf{r}(1.5)\quad \mathbf{r}(1)}{1.5-1} = \frac{(5.9\mathbf{i}+6.4\mathbf{j}+2.8\mathbf{k})-(4.5\mathbf{i}+6.0\mathbf{j}+3.0\mathbf{k})}{0.5} = 2.8\mathbf{i}+0.8\mathbf{j}-0.4\mathbf{k}$$

(b) We can estimate the velocity at $t=1$ by averaging the average velocities over the time intervals $[0.5,1]$ and $[1,1.5]$: $\mathbf{v}(1) \approx \frac{1}{2}[(2\mathbf{i}-2.4\mathbf{j}-0.6\mathbf{k})+(2.8\mathbf{i}+0.8\mathbf{j}-0.4\mathbf{k})] = 2.4\mathbf{i}-0.8\mathbf{j}-0.5\mathbf{k}$. Then the speed is $|\mathbf{v}(1)| \approx \sqrt{(2.4)^2+(-0.8)^2+(-0.5)^2} \approx 2.58$.

2. (a) The average velocity over $2 \le t \le 2.4$ is $\dfrac{\mathbf{r}(2.4)-\mathbf{r}(2)}{2.4-2} = 2.5\,[\mathbf{r}(2.4)-\mathbf{r}(2)]$, so we sketch a vector in the same direction but 2.5 times the length of $[\mathbf{r}(2.4)-\mathbf{r}(2)]$.



(b) The average velocity over $1.5 \le t \le 2$ is $\dfrac{\mathbf{r}(2)-\mathbf{r}(1.5)}{2-1.5} = 2[\mathbf{r}(2)-\mathbf{r}(1.5)]$, so we sketch a vector in the same direction but twice the length of $[\mathbf{r}(2)-\mathbf{r}(1.5)]$.

(c) Using Equation 2 we have $\mathbf{v}(2) = \lim\limits_{h\to 0} \dfrac{\mathbf{r}(2+h)-\mathbf{r}(2)}{h}$.

(d) $\mathbf{v}(2)$ is tangent to the curve at $\mathbf{r}(2)$ and points in the direction of increasing t. Its length is the speed of the particle at $t=2$. We can estimate the speed by averaging the lengths of the vectors found in parts (a) and (b) which represent the average speed over $2 \le t \le 2.4$ and $1.5 \le t \le 2$ respectively. Using the axes scale as a guide, we estimate the vectors to have lengths 2.8 and 2.7. Thus, we estimate the speed at $t=2$ to be $|\mathbf{v}(2)| \approx \frac{1}{2}(2.8+2.7) = 2.75$ and we draw the velocity vector $\mathbf{v}(2)$ with this length.



3. $\mathbf{r}(t) = \left\langle -\frac{1}{2}t^2, t \right\rangle \Rightarrow$ At $t=2$:

$\mathbf{v}(t) = \mathbf{r}'(t) = \langle -t, 1 \rangle$ $\quad \mathbf{v}(2) = \langle -2, 1 \rangle$

$\mathbf{a}(t) = \mathbf{r}''(t) = \langle -1, 0 \rangle$ $\quad \mathbf{a}(2) = \langle -1, 0 \rangle$

$|\mathbf{v}(t)| = \sqrt{t^2+1}$



Notice that $x = -\frac{1}{2}y^2$, so the path is a parabola.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $\mathbf{r}(t) = \langle t^2, 1/t^2 \rangle \Rightarrow$ At $t = 1$:

$\mathbf{v}(t) = \mathbf{r}'(t) = \langle 2t, -2/t^3 \rangle$ $\quad \mathbf{v}(1) = \langle 2, -2 \rangle$

$\mathbf{a}(t) = \mathbf{r}''(t) = \langle 2, 6/t^4 \rangle$ $\quad \mathbf{a}(1) = \langle 2, 6 \rangle$

$|\mathbf{v}(t)| = \sqrt{4t^2 + 4/t^6} = 2\sqrt{t^2 + 1/t^6}$

Notice that $y = 1/x$ and $x > 0$, so the path is part of the hyperbola $y = 1/x$.



5. $\mathbf{r}(t) = 3\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} \Rightarrow$ At $t = \pi/3$:

$\mathbf{v}(t) = -3\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$ $\quad \mathbf{v}\left(\frac{\pi}{3}\right) = -\frac{3\sqrt{3}}{2}\,\mathbf{i} + \mathbf{j}$

$\mathbf{a}(t) = -3\cos t\,\mathbf{i} - 2\sin t\,\mathbf{j}$ $\quad \mathbf{a}\left(\frac{\pi}{3}\right) = -\frac{3}{2}\,\mathbf{i} - \sqrt{3}\,\mathbf{j}$

$|\mathbf{v}(t)| = \sqrt{9\sin^2 t + 4\cos^2 t} = \sqrt{5\sin^2 t + 4\sin^2 t + 4\cos^2 t}$

$= \sqrt{4 + 5\sin^2 t}$

Notice that $x^2/9 + y^2/4 = \sin^2 t + \cos^2 t = 1$, so the path is an ellipse.



6. $\mathbf{r}(t) = e^t\,\mathbf{i} + e^{2t}\,\mathbf{j} \Rightarrow$ At $t = 0$:

$\mathbf{v}(t) = e^t\,\mathbf{i} + 2e^{2t}\,\mathbf{j}$ $\quad \mathbf{v}(0) = \mathbf{i} + 2\mathbf{j}$

$\mathbf{a}(t) = e^t\,\mathbf{i} + 4e^{2t}\,\mathbf{j}$ $\quad \mathbf{a}(0) = \mathbf{i} + 4\mathbf{j}$

$|\mathbf{v}(t)| = \sqrt{e^{2t} + 4e^{4t}} = e^t\sqrt{1 + 4e^{2t}}$

Notice that $y = e^{2t} = (e^t)^2 = x^2$, so the particle travels along a parabola, but $x = e^t$, so $x > 0$.



7. $\mathbf{r}(t) = t\,\mathbf{i} + t^2\,\mathbf{j} + 2\,\mathbf{k} \Rightarrow$ At $t = 1$:

$\mathbf{v}(t) = \mathbf{i} + 2t\,\mathbf{j}$ $\quad \mathbf{v}(1) = \mathbf{i} + 2\mathbf{j}$

$\mathbf{a}(t) = 2\,\mathbf{j}$ $\quad \mathbf{a}(1) = 2\mathbf{j}$

$|\mathbf{v}(t)| = \sqrt{1 + 4t^2}$

Here $x = t$, $y = t^2 \Rightarrow y = x^2$ and $z = 2$, so the path of the particle is a parabola in the plane $z = 2$.



8. $\mathbf{r}(t) = t\,\mathbf{i} + 2\cos t\,\mathbf{j} + \sin t\,\mathbf{k} \Rightarrow$ At $t = 0$:

$\mathbf{v}(t) = \mathbf{i} - 2\sin t\,\mathbf{j} + \cos t\,\mathbf{k}$ $\quad \mathbf{v}(0) = \mathbf{i} + \mathbf{k}$

$\mathbf{a}(t) = -2\cos t\,\mathbf{j} - \sin t\,\mathbf{k}$ $\quad \mathbf{a}(0) = -2\mathbf{j}$

$|\mathbf{v}(t)| = \sqrt{1 + 4\sin^2 t + \cos^2 t} = \sqrt{2 + 3\sin^2 t}$

Since $y^2/4 + z^2 = 1$, $x = t$, the path of the particle is an elliptical helix about the x-axis.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

9. $\mathbf{r}(t) = \langle t^2 + t, t^2 - t, t^3 \rangle \;\Rightarrow\; \mathbf{v}(t) = \mathbf{r}'(t) = \langle 2t + 1, 2t - 1, 3t^2 \rangle, \;\; \mathbf{a}(t) = \mathbf{v}'(t) = \langle 2, 2, 6t \rangle,$

$|\mathbf{v}(t)| = \sqrt{(2t+1)^2 + (2t-1)^2 + (3t^2)^2} = \sqrt{9t^4 + 8t^2 + 2}.$

10. $\mathbf{r}(t) = \langle 2\cos t, 3t, 2\sin t \rangle \;\Rightarrow\; \mathbf{v}(t) = \mathbf{r}'(t) = \langle -2\sin t, 3, 2\cos t \rangle, \;\; \mathbf{a}(t) = \mathbf{v}'(t) = \langle -2\cos t, 0, -2\sin t \rangle,$

$|\mathbf{v}(t)| = \sqrt{4\sin^2 t + 9 + 4\cos^2 t} = \sqrt{13}.$

11. $\mathbf{r}(t) = \sqrt{2}\,t\,\mathbf{i} + e^t\,\mathbf{j} + e^{-t}\,\mathbf{k} \;\Rightarrow\; \mathbf{v}(t) = \mathbf{r}'(t) = \sqrt{2}\,\mathbf{i} + e^t\,\mathbf{j} - e^{-t}\,\mathbf{k}, \;\; \mathbf{a}(t) = \mathbf{v}'(t) = e^t\,\mathbf{j} + e^{-t}\,\mathbf{k},$

$|\mathbf{v}(t)| = \sqrt{2 + e^{2t} + e^{-2t}} = \sqrt{(e^t + e^{-t})^2} = e^t + e^{-t}.$

12. $\mathbf{r}(t) = t^2\,\mathbf{i} + 2t\,\mathbf{j} + \ln t\,\mathbf{k} \;\Rightarrow\; \mathbf{v}(t) = \mathbf{r}'(t) = 2t\,\mathbf{i} + 2\,\mathbf{j} + (1/t)\,\mathbf{k}, \;\; \mathbf{a}(t) = \mathbf{v}'(t) = 2\,\mathbf{i} - (1/t^2)\,\mathbf{k},$

$|\mathbf{v}(t)| = \sqrt{4t^2 + 4 + (1/t^2)} = \sqrt{[2t + (1/t)]^2} = |2t + (1/t)|.$

13. $\mathbf{r}(t) = e^t\langle \cos t, \sin t, t \rangle \;\Rightarrow$

$\mathbf{v}(t) = \mathbf{r}'(t) = e^t\langle \cos t, \sin t, t \rangle + e^t\,\langle -\sin t, \cos t, 1 \rangle = e^t\langle \cos t - \sin t, \sin t + \cos t, t + 1 \rangle$

$\mathbf{a}(t) = \mathbf{v}'(t) = e^t\langle \cos t - \sin t - \sin t - \cos t, \sin t + \cos t + \cos t - \sin t, t + 1 + 1 \rangle$

$= e^t\langle -2\sin t, 2\cos t, t + 2 \rangle$

$|\mathbf{v}(t)| = e^t\sqrt{\cos^2 t + \sin^2 t - 2\cos t \sin t + \sin^2 t + \cos^2 t + 2\sin t \cos t + t^2 + 2t + 1}$

$= e^t\sqrt{t^2 + 2t + 3}$

14. $\mathbf{r}(t) = \langle t^2, \sin t - t\cos t, \cos t + t\sin t \rangle \;\Rightarrow$

$\mathbf{v}(t) = \mathbf{r}'(t) = \langle 2t, \cos t - (-t\sin t + \cos t), -\sin t + t\cos t + \sin t \rangle = \langle 2t, t\sin t, t\cos t \rangle,$

$\mathbf{a}(t) = \mathbf{v}'(t) = \langle 2, t\cos t + \sin t, -t\sin t + \cos t \rangle,$

$|\mathbf{v}(t)| = \sqrt{4t^2 + t^2\sin^2 t + t^2\cos^2 t} = \sqrt{4t^2 + t^2} = \sqrt{5t^2} = \sqrt{5}\,t \quad [\text{since } t \geq 0].$

15. $\mathbf{a}(t) = 2\,\mathbf{i} + 2t\,\mathbf{k} \;\Rightarrow\; \mathbf{v}(t) = \int \mathbf{a}(t)\,dt = \int (2\,\mathbf{i} + 2t\,\mathbf{k})\,dt = 2t\,\mathbf{i} + t^2\,\mathbf{k} + \mathbf{C}$. Then $\mathbf{v}(0) = \mathbf{C}$ but we were given that $\mathbf{v}(0) = 3\,\mathbf{i} - \mathbf{j}$, so $\mathbf{C} = 3\,\mathbf{i} - \mathbf{j}$ and $\mathbf{v}(t) = 2t\,\mathbf{i} + t^2\,\mathbf{k} + 3\,\mathbf{i} - \mathbf{j} = (2t + 3)\,\mathbf{i} - \mathbf{j} + t^2\,\mathbf{k}$.

$\mathbf{r}(t) = \int \mathbf{v}(t)\,dt = \int \left[(2t + 3)\,\mathbf{i} - \mathbf{j} + t^2\,\mathbf{k}\right] dt = (t^2 + 3t)\,\mathbf{i} - t\,\mathbf{j} + \frac{1}{3}t^3\,\mathbf{k} + \mathbf{D}$. Here $\mathbf{r}(0) = \mathbf{D}$ and we were given that $\mathbf{r}(0) = \mathbf{j} + \mathbf{k}$, so $\mathbf{D} = \mathbf{j} + \mathbf{k}$ and $\mathbf{r}(t) = (t^2 + 3t)\,\mathbf{i} + (1 - t)\,\mathbf{j} + \left(\frac{1}{3}t^3 + 1\right)\mathbf{k}$.

16. $\mathbf{a}(t) = \sin t\,\mathbf{i} + 2\cos t\,\mathbf{j} + 6t\,\mathbf{k} \;\Rightarrow\; \mathbf{v}(t) = \int \mathbf{a}(t)\,dt = \int (\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j} + 6t\,\mathbf{k})\,dt = -\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} + 3t^2\,\mathbf{k} + \mathbf{C}.$

Then $\mathbf{v}(0) = -\mathbf{i} + \mathbf{C}$ but we were given that $\mathbf{v}(0) = -\mathbf{k}$, so $-\mathbf{i} + \mathbf{C} = -\mathbf{k} \;\Rightarrow\; \mathbf{C} = \mathbf{i} - \mathbf{k}$

and $\mathbf{v}(t) = (1 - \cos t)\,\mathbf{i} + 2\sin t\,\mathbf{j} + (3t^2 - 1)\,\mathbf{k}$.

$\mathbf{r}(t) = \int \mathbf{v}(t)\,dt = \int \left[(1 - \cos t)\,\mathbf{i} + 2\sin t\,\mathbf{j} + (3t^2 - 1)\,\mathbf{k}\right] dt = (t - \sin t)\,\mathbf{i} - 2\cos t\,\mathbf{j} + (t^3 - t)\,\mathbf{k} + \mathbf{D}$. Here

$\mathbf{r}(0) = -2\,\mathbf{j} + \mathbf{D}$ and we were given that $\mathbf{r}(0) = \mathbf{j} - 4\,\mathbf{k}$, so $\mathbf{D} = 3\,\mathbf{j} - 4\,\mathbf{k}$ and

$\mathbf{r}(t) = (t - \sin t)\,\mathbf{i} + (3 - 2\cos t)\,\mathbf{j} + (t^3 - t - 4)\,\mathbf{k}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. (a) $\mathbf{a}(t) = 2t\,\mathbf{i} + \sin t\,\mathbf{j} + \cos 2t\,\mathbf{k} \Rightarrow$

$\mathbf{v}(t) = \int (2t\,\mathbf{i} + \sin t\,\mathbf{j} + \cos 2t\,\mathbf{k})\,dt = t^2\,\mathbf{i} - \cos t\,\mathbf{j} + \frac{1}{2}\sin 2t\,\mathbf{k} + \mathbf{C}$

and $\mathbf{i} = \mathbf{v}(0) = -\mathbf{j} + \mathbf{C}$, so $\mathbf{C} = \mathbf{i} + \mathbf{j}$

and $\mathbf{v}(t) = (t^2 + 1)\,\mathbf{i} + (1 - \cos t)\,\mathbf{j} + \frac{1}{2}\sin 2t\,\mathbf{k}$.

$\mathbf{r}(t) = \int [(t^2 + 1)\,\mathbf{i} + (1 - \cos t)\,\mathbf{j} + \frac{1}{2}\sin 2t\,\mathbf{k}]dt$

$= (\frac{1}{3}t^3 + t)\,\mathbf{i} + (t - \sin t)\,\mathbf{j} - \frac{1}{4}\cos 2t\,\mathbf{k} + \mathbf{D}$

(b)



But $\mathbf{j} = \mathbf{r}(0) = -\frac{1}{4}\,\mathbf{k} + \mathbf{D}$, so $\mathbf{D} = \mathbf{j} + \frac{1}{4}\,\mathbf{k}$ and $\mathbf{r}(t) = (\frac{1}{3}t^3 + t)\,\mathbf{i} + (t - \sin t + 1)\,\mathbf{j} + (\frac{1}{4} - \frac{1}{4}\cos 2t)\,\mathbf{k}$.

18. (a) $\mathbf{a}(t) = t\,\mathbf{i} + e^t\,\mathbf{j} + e^{-t}\,\mathbf{k} \Rightarrow$

$\mathbf{v}(t) = \int (t\,\mathbf{i} + e^t\,\mathbf{j} + e^{-t}\,\mathbf{k})\,dt = \frac{1}{2}t^2\,\mathbf{i} + e^t\,\mathbf{j} - e^{-t}\,\mathbf{k} + \mathbf{C}$

and $\mathbf{k} = \mathbf{v}(0) = \mathbf{j} - \mathbf{k} + \mathbf{C}$, so $\mathbf{C} = -\mathbf{j} + 2\,\mathbf{k}$

and $\mathbf{v}(t) = \frac{1}{2}t^2\,\mathbf{i} + (e^t - 1)\,\mathbf{j} + (2 - e^{-t})\,\mathbf{k}$.

$\mathbf{r}(t) = \int [\frac{1}{2}t^2\,\mathbf{i} + (e^t - 1)\,\mathbf{j} + (2 - e^{-t})\,\mathbf{k}]\,dt$

$= \frac{1}{6}t^3\,\mathbf{i} + (e^t - t)\,\mathbf{j} + (e^{-t} + 2t)\,\mathbf{k} + \mathbf{D}$

(b)



But $\mathbf{j} + \mathbf{k} = \mathbf{r}(0) = \mathbf{j} + \mathbf{k} + \mathbf{D}$, so $\mathbf{D} = \mathbf{0}$ and $\mathbf{r}(t) = \frac{1}{6}t^3\,\mathbf{i} + (e^t - t)\,\mathbf{j} + (e^{-t} + 2t)\,\mathbf{k}$.

19. $\mathbf{r}(t) = \langle t^2, 5t, t^2 - 16t\rangle \Rightarrow \mathbf{v}(t) = \langle 2t, 5, 2t - 16\rangle$, $|\mathbf{v}(t)| = \sqrt{4t^2 + 25 + 4t^2 - 64t + 256} = \sqrt{8t^2 - 64t + 281}$

and $\dfrac{d}{dt}|\mathbf{v}(t)| = \frac{1}{2}(8t^2 - 64t + 281)^{-1/2}(16t - 64)$. This is zero if and only if the numerator is zero, that is, $16t - 64 = 0$ or $t = 4$. Since $\dfrac{d}{dt}|\mathbf{v}(t)| < 0$ for $t < 4$ and $\dfrac{d}{dt}|\mathbf{v}(t)| > 0$ for $t > 4$, the minimum speed of $\sqrt{153}$ is attained at $t = 4$ units of time.

20. Since $\mathbf{r}(t) = t^3\,\mathbf{i} + t^2\,\mathbf{j} + t^3\,\mathbf{k}$, $\mathbf{a}(t) = \mathbf{r}''(t) = 6t\,\mathbf{i} + 2\,\mathbf{j} + 6t\,\mathbf{k}$. By Newton's Second Law, $\mathbf{F}(t) = m\,\mathbf{a}(t) = 6mt\,\mathbf{i} + 2m\,\mathbf{j} + 6mt\,\mathbf{k}$ is the required force.

21. $|\mathbf{F}(t)| = 20$ N in the direction of the positive z-axis, so $\mathbf{F}(t) = 20\,\mathbf{k}$. Also $m = 4$ kg, $\mathbf{r}(0) = \mathbf{0}$ and $\mathbf{v}(0) = \mathbf{i} - \mathbf{j}$. Since $20\mathbf{k} = \mathbf{F}(t) = 4\,\mathbf{a}(t)$, $\mathbf{a}(t) = 5\,\mathbf{k}$. Then $\mathbf{v}(t) = 5t\,\mathbf{k} + \mathbf{c}_1$ where $\mathbf{c}_1 = \mathbf{i} - \mathbf{j}$ so $\mathbf{v}(t) = \mathbf{i} - \mathbf{j} + 5t\,\mathbf{k}$ and the speed is $|\mathbf{v}(t)| = \sqrt{1 + 1 + 25t^2} = \sqrt{25t^2 + 2}$. Also $\mathbf{r}(t) = t\,\mathbf{i} - t\,\mathbf{j} + \frac{5}{2}t^2\,\mathbf{k} + \mathbf{c}_2$ and $\mathbf{0} = \mathbf{r}(0)$, so $\mathbf{c}_2 = \mathbf{0}$ and $\mathbf{r}(t) = t\,\mathbf{i} - t\,\mathbf{j} + \frac{5}{2}t^2\,\mathbf{k}$.

22. The argument here is the same as that in Example 13.2.4 with $\mathbf{r}(t)$ replaced by $\mathbf{v}(t)$ and $\mathbf{r}'(t)$ replaced by $\mathbf{a}(t)$.

23. $|\mathbf{v}(0)| = 200$ m/s and, since the angle of elevation is $60°$, a unit vector in the direction of the velocity is $(\cos 60°)\mathbf{i} + (\sin 60°)\mathbf{j} = \frac{1}{2}\mathbf{i} + \frac{\sqrt{3}}{2}\,\mathbf{j}$. Thus $\mathbf{v}(0) = 200\left(\frac{1}{2}\mathbf{i} + \frac{\sqrt{3}}{2}\,\mathbf{j}\right) = 100\,\mathbf{i} + 100\sqrt{3}\,\mathbf{j}$ and if we set up the axes so that the projectile starts at the origin, then $\mathbf{r}(0) = \mathbf{0}$. Ignoring air resistance, the only force is that due to gravity, so $\mathbf{F}(t) = m\mathbf{a}(t) = -mg\,\mathbf{j}$ where $g \approx 9.8$ m/s^2. Thus $\mathbf{a}(t) = -9.8\,\mathbf{j}$ and, integrating, we have $\mathbf{v}(t) = -9.8t\,\mathbf{j} + \mathbf{C}$. But

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$100\,\mathbf{i} + 100\sqrt{3}\,\mathbf{j} = \mathbf{v}(0) = \mathbf{C}$, so $\mathbf{v}(t) = 100\,\mathbf{i} + \left(100\sqrt{3} - 9.8t\right)\mathbf{j}$ and then (integrating again)

$\mathbf{r}(t) = 100t\,\mathbf{i} + \left(100\sqrt{3}\,t - 4.9t^2\right)\mathbf{j} + \mathbf{D}$ where $\mathbf{0} = \mathbf{r}(0) = \mathbf{D}$. Thus the position function of the projectile is

$\mathbf{r}(t) = 100t\,\mathbf{i} + \left(100\sqrt{3}\,t - 4.9t^2\right)\mathbf{j}$.

(a) Parametric equations for the projectile are $x(t) = 100t$, $y(t) = 100\sqrt{3}\,t - 4.9t^2$. The projectile reaches the ground when $y(t) = 0$ (and $t > 0$) $\Rightarrow$ $100\sqrt{3}\,t - 4.9t^2 = t\left(100\sqrt{3} - 4.9t\right) = 0$ $\Rightarrow$ $t = \frac{100\sqrt{3}}{4.9} \approx 35.3$ s. So the range is $x\left(\frac{100\sqrt{3}}{4.9}\right) = 100\left(\frac{100\sqrt{3}}{4.9}\right) \approx 3535$ m.

(b) The maximum height is reached when $y(t)$ has a critical number (or equivalently, when the vertical component of velocity is 0): $y'(t) = 0$ $\Rightarrow$ $100\sqrt{3} - 9.8t = 0$ $\Rightarrow$ $t = \frac{100\sqrt{3}}{9.8} \approx 17.7$ s. Thus the maximum height is $y\left(\frac{100\sqrt{3}}{9.8}\right) = 100\sqrt{3}\left(\frac{100\sqrt{3}}{9.8}\right) - 4.9\left(\frac{100\sqrt{3}}{9.8}\right)^2 \approx 1531$ m.

(c) From part (a), impact occurs at $t = \frac{100\sqrt{3}}{4.9}$ s. Thus, the velocity at impact is $\mathbf{v}\left(\frac{100\sqrt{3}}{4.9}\right) = 100\,\mathbf{i} + \left[100\sqrt{3} - 9.8\left(\frac{100\sqrt{3}}{4.9}\right)\right]\mathbf{j} = 100\,\mathbf{i} - 100\sqrt{3}\,\mathbf{j}$ and the speed is $\left|\mathbf{v}\left(\frac{100\sqrt{3}}{4.9}\right)\right| = \sqrt{10{,}000 + 30{,}000} = 200$ m/s.

24. As in Exercise 23, $\mathbf{v}(t) = 100\,\mathbf{i} + \left(100\sqrt{3} - 9.8t\right)\mathbf{j}$ and $\mathbf{r}(t) = 100t\,\mathbf{i} + \left(100\sqrt{3}\,t - 4.9t^2\right)\mathbf{j} + \mathbf{D}$.

But $\mathbf{r}(0) = 100\,\mathbf{j}$, so $\mathbf{D} = 100\,\mathbf{j}$ and $\mathbf{r}(t) = 100t\,\mathbf{i} + \left(100 + 100\sqrt{3}\,t - 4.9t^2\right)\mathbf{j}$.

(a) $y = 0$ $\Rightarrow$ $100 + 100\sqrt{3}\,t - 4.9t^2 = 0$ or $4.9t^2 - 100\sqrt{3}\,t - 100 = 0$. From the quadratic formula we have $t = \dfrac{100\sqrt{3} \pm \sqrt{(-100\sqrt{3})^2 - 4(4.9)(-100)}}{2(4.9)} = \dfrac{100\sqrt{3} \pm \sqrt{31{,}960}}{9.8}$. Taking the positive t-value gives $t = \frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8} \approx 35.9$ s. Thus the range is $x = 100 \cdot \frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8} \approx 3592$ m.

(b) The maximum height is attained when $\dfrac{dy}{dt} = 0$ $\Rightarrow$ $100\sqrt{3} - 9.8t = 0$ $\Rightarrow$ $t = \frac{100\sqrt{3}}{9.8} \approx 17.7$ s and the maximum height is $100 + 100\sqrt{3}\left(\frac{100\sqrt{3}}{9.8}\right) - 4.9\left(\frac{100\sqrt{3}}{9.8}\right)^2 \approx 1631$ m.

Alternate solution: Because the projectile is fired in the same direction and with the same velocity as in Exercise 23, but from a point 100 m higher, the maximum height reached is 100 m higher than that found in Exercise 23, that is, 1531 m + 100 m = 1631 m.

(c) From part (a), impact occurs at $t = \frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8}$ s. Thus the velocity at impact is $\mathbf{v}\left(\frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8}\right) = 100\,\mathbf{i} + \left[100\sqrt{3} - 9.8\left(\frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8}\right)\right]\mathbf{j} = 100\,\mathbf{i} - \sqrt{31{,}960}\,\mathbf{j}$ and the speed is $|\mathbf{v}| = \sqrt{10{,}000 + 31{,}960} = \sqrt{41{,}960} \approx 205$ m/s.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. As in Example 5, $\mathbf{r}(t) = (v_0 \cos 45°)t\,\mathbf{i} + \left[(v_0 \sin 45°)t - \frac{1}{2}gt^2\right]\mathbf{j} = \frac{1}{2}\left[v_0\sqrt{2}\,t\,\mathbf{i} + \left(v_0\sqrt{2}\,t - gt^2\right)\mathbf{j}\right]$. The ball lands when $y = 0$ (and $t > 0$) $\Rightarrow$ $t = \dfrac{v_0\sqrt{2}}{g}$ s. Now since it lands 90 m away, $90 = x = \frac{1}{2}v_0\sqrt{2}\,\dfrac{v_0\sqrt{2}}{g}$ or $v_0^2 = 90g$ and the initial velocity is $v_0 = \sqrt{90g} \approx 30$ m/s.

26. Let α be the angle of elevation. Here $v_0 = 400$ m/s and from Example 5, the horizontal distance traveled by the projectile is $d = \dfrac{v_0^2 \sin 2\alpha}{g}$. We want $\dfrac{400^2 \sin 2\alpha}{g} = 3000$ $\Rightarrow$ $\sin 2\alpha = \dfrac{3000g}{400^2} \approx 0.1838$ $\Rightarrow$ $2\alpha \approx \sin^{-1}(0.1838) \approx 10.6°$ or $2\alpha \approx 180° - 10.6° = 169.4°$. Thus two angles of elevation are $\alpha \approx 5.3°$ and $\alpha \approx 84.7°$.

27. As in Example 5, $\mathbf{r}(t) = (v_0 \cos 36°)t\,\mathbf{i} + \left[(v_0 \sin 36°)t - \frac{1}{2}gt^2\right]\mathbf{j}$ and then $\mathbf{v}(t) = \mathbf{r}'(t) = (v_0 \cos 36°)\,\mathbf{i} + [(v_0 \sin 36°) - gt]\,\mathbf{j}$. The shell reaches its maximum height when the vertical component of velocity is zero, so $(v_0 \sin 36°) - gt = 0$ $\Rightarrow$ $t = \dfrac{v_0 \sin 36°}{g}$. The vertical height of the shell at that time is 1600 ft, so

$$(v_0 \sin 36°)\left(\frac{v_0 \sin 36°}{g}\right) - \frac{1}{2}g\left(\frac{v_0 \sin 36°}{g}\right)^2 = 1600 \quad \Rightarrow \quad \left(\frac{v_0^2 \sin^2 36°}{g}\right) - \frac{1}{2}\left(\frac{v_0^2 \sin^2 36°}{g}\right) = 1600 \quad \Rightarrow$$

$$\frac{v_0^2 \sin^2 36°}{2g} = 1600 \quad \Rightarrow \quad v_0^2 = \frac{1600(2g)}{\sin^2 36°} \quad \Rightarrow \quad v_0 = \sqrt{\frac{3200g}{\sin^2 36°}} \approx \frac{\sqrt{3200(32)}}{\sin 36°} \approx 544 \text{ ft/s}.$$

28. Here $v_0 = 115$ ft/s, the angle of elevation is $\alpha = 50°$, and if we place the origin at home plate, then $\mathbf{r}(0) = 3\,\mathbf{j}$. As in Example 5, we have $\mathbf{r}\,(t) = -\frac{1}{2}gt^2\,\mathbf{j} + t\,\mathbf{v}_0 + \mathbf{D}$ where $\mathbf{D} = \mathbf{r}(0) = 3\,\mathbf{j}$ and $\mathbf{v}_0 = v_0 \cos\alpha\,\mathbf{i} + v_0 \sin\alpha\,\mathbf{j}$, so $\mathbf{r}(t) = (v_0 \cos\alpha)t\,\mathbf{i} + \left[(v_0 \sin\alpha)t - \frac{1}{2}gt^2 + 3\right]\mathbf{j}$. Thus, parametric equations for the trajectory of the ball are $x = (v_0 \cos\alpha)t$, $y = (v_0 \sin\alpha)t - \frac{1}{2}gt^2 + 3$. The ball reaches the fence when $x = 400$ $\Rightarrow$ $(v_0 \cos\alpha)t = 400$ $\Rightarrow$ $t = \dfrac{400}{v_0 \cos\alpha} = \dfrac{400}{115 \cos 50°} \approx 5.41$ s. At this time, the height of the ball is $y = (v_0 \sin\alpha)t - \frac{1}{2}gt^2 + 3 \approx (115 \sin 50°)(5.41) - \frac{1}{2}(32)(5.41)^2 + 3 \approx 11.2$ ft. Since the fence is 10 ft high, the ball clears the fence.

29. Place the catapult at the origin and assume the catapult is 100 meters from the city, so the city lies between $(100, 0)$ and $(600, 0)$. The initial speed is $v_0 = 80$ m/s and let θ be the angle the catapult is set at. As in Example 5, the trajectory of the catapulted rock is given by $\mathbf{r}\,(t) = (80 \cos\theta)t\,\mathbf{i} + \left[(80 \sin\theta)t - 4.9t^2\right]\,\mathbf{j}$. The top of the near city wall is at $(100, 15)$, which the rock will hit when $(80 \cos\theta)\,t = 100$ $\Rightarrow$ $t = \dfrac{5}{4 \cos\theta}$ and $(80 \sin\theta)t - 4.9t^2 = 15$ $\Rightarrow$

$$80 \sin\theta \cdot \frac{5}{4 \cos\theta} - 4.9\left(\frac{5}{4 \cos\theta}\right)^2 = 15 \quad \Rightarrow \quad 100 \tan\theta - 7.65625 \sec^2\theta = 15.$$

Replacing $\sec^2\theta$ with $\tan^2\theta + 1$ gives $7.65625 \tan^2\theta - 100 \tan\theta + 22.65625 = 0$. Using the quadratic formula, we have $\tan\theta \approx 0.230635, 12.8306$ $\Rightarrow$ $\theta \approx 13.0°, 85.5°$. So for $13.0° < \theta < 85.5°$, the rock will land beyond the near city wall. The base of the far wall is

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

located at $(600, 0)$ which the rock hits if $(80\cos\theta)t = 600 \Rightarrow t = \dfrac{15}{2\cos\theta}$ and $(80\sin\theta)t - 4.9t^2 = 0 \Rightarrow$

$80\sin\theta \cdot \dfrac{15}{2\cos\theta} - 4.9\left(\dfrac{15}{2\cos\theta}\right)^2 = 0 \Rightarrow 600\tan\theta - 275.625\sec^2\theta = 0 \Rightarrow$

$275.625\tan^2\theta - 600\tan\theta + 275.625 = 0$. Solutions are $\tan\theta \approx 0.658678, 1.51819 \Rightarrow \theta \approx 33.4°, 56.6°$. Thus the rock lands beyond the enclosed city ground for $33.4° < \theta < 56.6°$, and the angles that allow the rock to land on city ground are $13.0° < \theta < 33.4°, 56.6° < \theta < 85.5°$. If you consider that the rock can hit the far wall and bounce back into the city, we calculate the angles that cause the rock to hit the top of the wall at $(600, 15)$: $(80\cos\theta)t = 600 \Rightarrow t = \dfrac{15}{2\cos\theta}$ and $(80\sin\theta)t - 4.9t^2 = 15 \Rightarrow 600\tan\theta - 275.625\sec^2\theta = 15 \Rightarrow 275.625\tan^2\theta - 600\tan\theta + 290.625 = 0$. Solutions are $\tan\theta \approx 0.727506, 1.44936 \Rightarrow \theta \approx 36.0°, 55.4°$, so the catapult should be set with angle θ where $13.0° < \theta < 36.0°, 55.4° < \theta < 85.5°$.

30. If we place the projectile at the origin then, as in Example 5, $\mathbf{r}(t) = (v_0\cos\alpha)t\,\mathbf{i} + \left[(v_0\sin\alpha)t - \frac{1}{2}gt^2\right]\mathbf{j}$ and $\mathbf{v}(t) = (v_0\cos\alpha)\,\mathbf{i} + [(v_0\sin\alpha) - gt]\,\mathbf{j}$. The maximum height is reached when the vertical component of velocity is zero, so $(v_0\sin\alpha) - gt = 0 \Rightarrow t = \dfrac{v_0\sin\alpha}{g}$, and the corresponding height is the vertical component of the position function:

$$(v_0\sin\alpha)t - \tfrac{1}{2}gt^2 = (v_0\sin\alpha)\left(\frac{v_0\sin\alpha}{g}\right) - \tfrac{1}{2}g\left(\frac{v_0\sin\alpha}{g}\right)^2 = \frac{1}{2g}v_0^2\sin^2\alpha$$

Half that time is $t = \dfrac{v_0\sin\alpha}{2g}$, when the height of the projectile is

$$\begin{aligned}(v_0\sin\alpha)t - \tfrac{1}{2}gt^2 &= (v_0\sin\alpha)\left(\frac{v_0\sin\alpha}{2g}\right) - \tfrac{1}{2}g\left(\frac{v_0\sin\alpha}{2g}\right)^2 \\ &= \frac{1}{2g}v_0^2\sin^2\alpha - \frac{1}{8g}v_0^2\sin^2\alpha = \frac{3}{8g}v_0^2\sin^2\alpha = \frac{3}{4}\left(\frac{1}{2g}v_0^2\sin^2\alpha\right)\end{aligned}$$

or three-quarters of the maximum height.

31. Here $\mathbf{a}(t) = -4\,\mathbf{j} - 32\,\mathbf{k}$ so $\mathbf{v}(t) = -4t\,\mathbf{j} - 32t\,\mathbf{k} + \mathbf{v}_0 = -4t\,\mathbf{j} - 32t\,\mathbf{k} + 50\,\mathbf{i} + 80\,\mathbf{k} = 50\,\mathbf{i} - 4t\,\mathbf{j} + (80 - 32t)\,\mathbf{k}$ and $\mathbf{r}(t) = 50t\,\mathbf{i} - 2t^2\,\mathbf{j} + (80t - 16t^2)\,\mathbf{k}$ (note that $\mathbf{r}_0 = \mathbf{0}$). The ball lands when the z-component of $\mathbf{r}(t)$ is zero and $t > 0$: $80t - 16t^2 = 16t(5 - t) = 0 \Rightarrow t = 5$. The position of the ball then is $\mathbf{r}(5) = 50(5)\,\mathbf{i} - 2(5)^2\,\mathbf{j} + [80(5) - 16(5)^2]\,\mathbf{k} = 250\,\mathbf{i} - 50\,\mathbf{j}$ or equivalently the point $(250, -50, 0)$. This is a distance of $\sqrt{250^2 + (-50)^2 + 0^2} = \sqrt{65{,}000} \approx 255$ ft from the origin at an angle of $\tan^{-1}\left(\frac{50}{250}\right) \approx 11.3°$ from the eastern direction toward the south. The speed of the ball is $|\mathbf{v}(5)| = |50\,\mathbf{i} - 20\,\mathbf{j} - 80\,\mathbf{k}| = \sqrt{50^2 + (-20)^2 + (-80)^2} = \sqrt{9300} \approx 96.4$ ft/s.

32. Place the ball at the origin and consider $\mathbf{j}$ to be pointing in the northward direction with $\mathbf{i}$ pointing east and $\mathbf{k}$ pointing upward. Force = mass × acceleration $\Rightarrow$ acceleration = force/mass, so the wind applies a constant acceleration of $4\text{ N}/0.8\text{ kg} = 5\text{ m/s}^2$ in the easterly direction. Combined with the acceleration due to gravity, the acceleration acting on the

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

ball is $\mathbf{a}(t) = 5\,\mathbf{i} - 9.8\,\mathbf{k}$. Then $\mathbf{v}(t) = \int \mathbf{a}(t)\,dt = 5t\,\mathbf{i} - 9.8t\,\mathbf{k} + \mathbf{C}$ where $\mathbf{C}$ is a constant vector.

We know $\mathbf{v}(0) = \mathbf{C} = -30\cos 30^\circ\,\mathbf{j} + 30\sin 30^\circ\,\mathbf{k} = -15\sqrt{3}\,\mathbf{j} + 15\,\mathbf{k} \;\Rightarrow\; \mathbf{C} = -15\sqrt{3}\,\mathbf{j} + 15\,\mathbf{k}$ and $\mathbf{v}(t) = 5t\,\mathbf{i} - 15\sqrt{3}\,\mathbf{j} + (15 - 9.8t)\,\mathbf{k}$. $\mathbf{r}(t) = \int \mathbf{v}(t)\,dt = 2.5t^2\,\mathbf{i} - 15\sqrt{3}\,t\,\mathbf{j} + \left(15t - 4.9t^2\right)\mathbf{k} + \mathbf{D}$ but $\mathbf{r}(0) = \mathbf{D} = \mathbf{0}$ so $\mathbf{r}(t) = 2.5t^2\,\mathbf{i} - 15\sqrt{3}\,t\,\mathbf{j} + \left(15t - 4.9t^2\right)\mathbf{k}$. The ball lands when $15t - 4.9t^2 = 0 \;\Rightarrow\; t = 0, t = 15/4.9 \approx 3.0612$ s, so the ball lands at approximately $\mathbf{r}(3.0612) \approx 23.43\,\mathbf{i} - 79.53\,\mathbf{j}$ which is 82.9 m away in the direction S 16.4°E. Its speed is approximately $|\mathbf{v}(3.0612)| \approx \left|15.306\,\mathbf{i} - 15\sqrt{3}\,\mathbf{j} - 15\,\mathbf{k}\right| \approx 33.68$ m/s.

33. (a) After t seconds, the boat will be $5t$ meters west of point A. The velocity of the water at that location is $\frac{3}{400}(5t)(40 - 5t)\,\mathbf{j}$. The velocity of the boat in still water is $5\,\mathbf{i}$, so the resultant velocity of the boat is $\mathbf{v}(t) = 5\,\mathbf{i} + \frac{3}{400}(5t)(40 - 5t)\,\mathbf{j} = 5\mathbf{i} + \left(\frac{3}{2}t - \frac{3}{16}t^2\right)\mathbf{j}$. Integrating, we obtain $\mathbf{r}(t) = 5t\,\mathbf{i} + \left(\frac{3}{4}t^2 - \frac{1}{16}t^3\right)\mathbf{j} + \mathbf{C}$. If we place the origin at A (and consider $\mathbf{j}$ to coincide with the northern direction) then $\mathbf{r}(0) = \mathbf{0} \;\Rightarrow\; \mathbf{C} = \mathbf{0}$ and we have $\mathbf{r}(t) = 5t\,\mathbf{i} + \left(\frac{3}{4}t^2 - \frac{1}{16}t^3\right)\mathbf{j}$. The boat reaches the east bank after 8 s, and it is located at $\mathbf{r}(8) = 5(8)\mathbf{i} + \left(\frac{3}{4}(8)^2 - \frac{1}{16}(8)^3\right)\mathbf{j} = 40\,\mathbf{i} + 16\,\mathbf{j}$. Thus the boat is 16 m downstream.



(b) Let α be the angle north of east that the boat heads. Then the velocity of the boat in still water is given by $5(\cos\alpha)\,\mathbf{i} + 5(\sin\alpha)\,\mathbf{j}$. At t seconds, the boat is $5(\cos\alpha)t$ meters from the west bank, at which point the velocity of the water is $\frac{3}{400}[5(\cos\alpha)t][40 - 5(\cos\alpha)t]\,\mathbf{j}$. The resultant velocity of the boat is given by $\mathbf{v}(t) = 5(\cos\alpha)\,\mathbf{i} + \left[5\sin\alpha + \frac{3}{400}(5t\cos\alpha)(40 - 5t\cos\alpha)\right]\mathbf{j} = (5\cos\alpha)\,\mathbf{i} + \left(5\sin\alpha + \frac{3}{2}t\cos\alpha - \frac{3}{16}t^2\cos^2\alpha\right)\mathbf{j}$. Integrating, $\mathbf{r}(t) = (5t\cos\alpha)\,\mathbf{i} + \left(5t\sin\alpha + \frac{3}{4}t^2\cos\alpha - \frac{1}{16}t^3\cos^2\alpha\right)\mathbf{j}$ (where we have again placed the origin at A). The boat will reach the east bank when $5t\cos\alpha = 40 \;\Rightarrow\; t = \dfrac{40}{5\cos\alpha} = \dfrac{8}{\cos\alpha}$.

In order to land at point $B(40, 0)$ we need $5t\sin\alpha + \frac{3}{4}t^2\cos\alpha - \frac{1}{16}t^3\cos^2\alpha = 0 \;\Rightarrow$

$$5\left(\frac{8}{\cos\alpha}\right)\sin\alpha + \frac{3}{4}\left(\frac{8}{\cos\alpha}\right)^2\cos\alpha - \frac{1}{16}\left(\frac{8}{\cos\alpha}\right)^3\cos^2\alpha = 0 \;\Rightarrow\; \frac{1}{\cos\alpha}\left(40\sin\alpha + 48 - 32\right) = 0 \;\Rightarrow$$

$40\sin\alpha + 16 = 0 \;\Rightarrow\; \sin\alpha = -\frac{2}{5}$. Thus $\alpha = \sin^{-1}\left(-\frac{2}{5}\right) \approx -23.6^\circ$, so the boat should head 23.6° south of east (upstream). The path does seem realistic. The boat initially heads upstream to counteract the effect of the current. Near the center of the river, the current is stronger and the boat is pushed downstream. When the boat nears the eastern bank, the current is slower and the boat is able to progress upstream to arrive at point B.



34. As in Exercise 33(b), let α be the angle north of east that the boat heads, so the velocity of the boat in still water is given by $5(\cos\alpha)\,\mathbf{i} + 5(\sin\alpha)\,\mathbf{j}$. At t seconds, the boat is $5(\cos\alpha)t$ meters from the west bank, at which point the velocity

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

of the water is $3\sin(\pi x/40)\,\mathbf{j} = 3\sin[\pi\cdot 5(\cos\alpha)t/40]\,\mathbf{j} = 3\sin\left(\frac{\pi}{8}t\cos\alpha\right)\mathbf{j}$. The resultant velocity of the boat then is given by $\mathbf{v}(t) = 5(\cos\alpha)\,\mathbf{i} + \left[5\sin\alpha + 3\sin\left(\frac{\pi}{8}t\cos\alpha\right)\right]\mathbf{j}$. Integrating,

$$\mathbf{r}(t) = (5t\cos\alpha)\,\mathbf{i} + \left[5t\sin\alpha - \frac{24}{\pi\cos\alpha}\cos\left(\tfrac{\pi}{8}t\cos\alpha\right)\right]\mathbf{j} + \mathbf{C}.$$

If we place the origin at A then $\mathbf{r}(0) = \mathbf{0} \;\Rightarrow\; -\frac{24}{\pi\cos\alpha}\,\mathbf{j} + \mathbf{C} = \mathbf{0} \;\Rightarrow\; \mathbf{C} = \frac{24}{\pi\cos\alpha}\,\mathbf{j}$ and

$\mathbf{r}(t) = (5t\cos\alpha)\,\mathbf{i} + \left[5t\sin\alpha - \frac{24}{\pi\cos\alpha}\cos\left(\frac{\pi}{8}t\cos\alpha\right) + \frac{24}{\pi\cos\alpha}\right]\mathbf{j}$. The boat will reach the east bank when

$5t\cos\alpha = 40 \;\Rightarrow\; t = \frac{8}{\cos\alpha}$. In order to land at point $B(40,0)$ we need

$$5t\sin\alpha - \frac{24}{\pi\cos\alpha}\cos\left(\tfrac{\pi}{8}t\cos\alpha\right) + \frac{24}{\pi\cos\alpha} = 0 \;\Rightarrow$$

$$5\left(\frac{8}{\cos\alpha}\right)\sin\alpha - \frac{24}{\pi\cos\alpha}\cos\left[\frac{\pi}{8}\left(\frac{8}{\cos\alpha}\right)\cos\alpha\right] + \frac{24}{\pi\cos\alpha} - 0 \;\Rightarrow\; \frac{1}{\cos\alpha}\left(40\sin\alpha - \frac{24}{\pi}\cos\pi + \frac{24}{\pi}\right) = 0 \;\Rightarrow$$

$40\sin\alpha + \frac{48}{\pi} = 0 \;\Rightarrow\; \sin\alpha = -\frac{6}{5\pi}$. Thus $\alpha = \sin^{-1}\left(-\frac{6}{5\pi}\right) \approx -22.5°$, so the boat should head $22.5°$ south of east.

35. If $\mathbf{r}'(t) = \mathbf{c}\times\mathbf{r}(t)$ then $\mathbf{r}'(t)$ is perpendicular to both $\mathbf{c}$ and $\mathbf{r}(t)$. Remember that $\mathbf{r}'(t)$ points in the direction of motion, so if $\mathbf{r}'(t)$ is always perpendicular to $\mathbf{c}$, the path of the particle must lie in a plane perpendicular to $\mathbf{c}$. But $\mathbf{r}'(t)$ is also perpendicular to the position vector $\mathbf{r}(t)$ which confines the path to a sphere centered at the origin. Considering both restrictions, the path must be contained in a circle that lies in a plane perpendicular to $\mathbf{c}$, and the circle is centered on a line through the origin in the direction of $\mathbf{c}$.

36. (a) From Equation 7 we have $\mathbf{a} = v'\mathbf{T} + \kappa v^2\mathbf{N}$. If a particle moves along a straight line, then $\kappa = 0$ [see Section 13.3], so the acceleration vector becomes $\mathbf{a} = v'\mathbf{T}$. Because the acceleration vector is a scalar multiple of the unit tangent vector, it is parallel to the tangent vector.

(b) If the speed of the particle is constant, then $v' = 0$ and Equation 7 gives $\mathbf{a} = \kappa v^2\mathbf{N}$. Thus the acceleration vector is parallel to the unit normal vector (which is perpendicular to the tangent vector and points in the direction that the curve is turning).

37. $\mathbf{r}(t) = (t^2+1)\,\mathbf{i} + t^3\,\mathbf{j} \;\Rightarrow\; \mathbf{r}'(t) = 2t\,\mathbf{i} + 3t^2\,\mathbf{j}$,

$|\mathbf{r}'(t)| = \sqrt{(2t)^2 + (3t^2)^2} = \sqrt{4t^2+9t^4} = t\sqrt{4+9t^2}$ [since $t \ge 0$], $\mathbf{r}''(t) = 2\,\mathbf{i} + 6t\,\mathbf{j}$, $\mathbf{r}'(t)\times\mathbf{r}''(t) = 6t^2\,\mathbf{k}$.

Then Equation 9 gives $a_T = \dfrac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \dfrac{(2t)(2)+(3t^2)(6t)}{t\sqrt{4+9t^2}} = \dfrac{4t+18t^3}{t\sqrt{4+9t^2}} = \dfrac{4+18t^2}{\sqrt{4+9t^2}}$

$$\left[\text{or by Equation 8,}\quad a_T = v' = \frac{d}{dt}\left[t\sqrt{4+9t^2}\right] = t\cdot\tfrac{1}{2}\left(4+9t^2\right)^{-1/2}(18t) + \left(4+9t^2\right)^{1/2}\cdot 1\right.$$
$$\left. = \left(4+9t^2\right)^{-1/2}\left(9t^2+4+9t^2\right) = (4+18t^2)/\sqrt{4+9t^2}\right]$$

and Equation 10 gives $a_N = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{6t^2}{t\sqrt{4+9t^2}} = \dfrac{6t}{\sqrt{4+9t^2}}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. $\mathbf{r}(t) = 2t^2\,\mathbf{i} + \left(\frac{2}{3}t^3 - 2t\right)\mathbf{j} \;\Rightarrow\; \mathbf{r}'(t) = 4t\,\mathbf{i} + (2t^2 - 2)\,\mathbf{j},$

$|\mathbf{r}'(t)| = \sqrt{16t^2 + (2t^2-2)^2} = \sqrt{4t^4 + 8t^2 + 4} = \sqrt{4\,(t^2+1)^2} = 2(t^2+1),$

$\mathbf{r}''(t) = 4\,\mathbf{i} + 4t\,\mathbf{j}, \quad \mathbf{r}'(t) \times \mathbf{r}''(t) = (8t^2+8)\,\mathbf{k}.$ Then Equation 9 gives

$$a_T = \frac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \frac{(4t)(4) + (2t^2-2)(4t)}{2(t^2+1)} = \frac{8t(t^2+1)}{2(t^2+1)} = 4t \quad \left[\text{or by Equation 8, } a_T = v' = \frac{d}{dt}\left[2(t^2+1)\right] = 4t\right]$$

and Equation 10 gives $a_N = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{8(t^2+1)}{2(t^2+1)} = 4.$

39. $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j} + \mathbf{k}, \quad |\mathbf{r}'(t)| = \sqrt{\sin^2 t + \cos^2 t + 1} = \sqrt{2},$

$\mathbf{r}''(t) = -\cos t\,\mathbf{i} - \sin t\,\mathbf{j}, \quad \mathbf{r}'(t)\times\mathbf{r}''(t) = \sin t\,\mathbf{i} - \cos t\,\mathbf{j} + \mathbf{k}.$

Then $a_T = \dfrac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \dfrac{\sin t\cos t - \sin t\cos t}{\sqrt{2}} = 0$ and $a_N = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{\sqrt{\sin^2 t + \cos^2 t + 1}}{\sqrt{2}} = \dfrac{\sqrt{2}}{\sqrt{2}} = 1.$

40. $\mathbf{r}(t) = t\,\mathbf{i} + 2e^t\,\mathbf{j} + e^{2t}\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = \mathbf{i} + 2e^t\,\mathbf{j} + 2e^{2t}\,\mathbf{k}, \quad |\mathbf{r}'(t)| = \sqrt{1 + 4e^{2t} + 4e^{4t}} = \sqrt{(1+2e^{2t})^2} = 1 + 2e^{2t},$

$\mathbf{r}''(t) = 2e^t\,\mathbf{j} + 4e^{2t}\,\mathbf{k}, \quad \mathbf{r}'(t)\times\mathbf{r}''(t) = 4e^{3t}\,\mathbf{i} - 4e^{2t}\,\mathbf{j} + 2e^t\,\mathbf{k},$

$|\mathbf{r}'(t)\times\mathbf{r}''(t)| = \sqrt{16e^{6t} + 16e^{4t} + 4e^{2t}} = \sqrt{4e^{2t}(2e^{2t}+1)^2} = 2e^t(2e^{2t}+1).$ Then

$$a_T = \frac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \frac{4e^{2t} + 8e^{4t}}{1 + 2e^{2t}} = \frac{4e^{2t}(1+2e^{2t})}{1+2e^{2t}} = 4e^{2t} \text{ and } a_N = \frac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \frac{2e^t(2e^{2t}+1)}{1+2e^{2t}} = 2e^t.$$

41. $\mathbf{r}(t) = \ln t\,\mathbf{i} + (t^2+3t)\,\mathbf{j} + 4\sqrt{t}\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = (1/t)\,\mathbf{i} + (2t+3)\,\mathbf{j} + (2/\sqrt{t}\,)\,\mathbf{k} \;\Rightarrow$

$\mathbf{r}''(t) = (-1/t^2)\,\mathbf{i} + 2\,\mathbf{j} - (1/t^{3/2})\,\mathbf{k}.$ The point $(0,4,4)$ corresponds to $t = 1$, where

$\mathbf{r}'(1) = \mathbf{i} + 5\,\mathbf{j} + 2\,\mathbf{k}, \quad \mathbf{r}''(1) = -\mathbf{i} + 2\,\mathbf{j} - \mathbf{k},$ and $\mathbf{r}'(1)\times\mathbf{r}''(1) = -9\,\mathbf{i} - \mathbf{j} + 7\,\mathbf{k}.$ Thus at the point $(0,4,4)$,

$$a_T = \frac{\mathbf{r}'(1)\cdot\mathbf{r}''(1)}{|\mathbf{r}'(1)|} = \frac{-1+10-2}{\sqrt{1+25+4}} = \frac{7}{\sqrt{30}} \text{ and } a_N = \frac{|\mathbf{r}'(1)\times\mathbf{r}''(1)|}{|\mathbf{r}'(1)|} = \frac{\sqrt{81+1+49}}{\sqrt{30}} = \sqrt{\frac{131}{30}}.$$

42. $\mathbf{r}(t) = t^{-1}\,\mathbf{i} + t^{-2}\,\mathbf{j} + t^{-3}\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = -t^{-2}\,\mathbf{i} - 2t^{-3}\,\mathbf{j} - 3t^{-4}\,\mathbf{k} \;\Rightarrow\; \mathbf{r}''(t) = 2t^{-3}\,\mathbf{i} + 6t^{-4}\,\mathbf{j} + 12t^{-5}\,\mathbf{k}.$ The point $(1,1,1)$ corresponds to $t = 1$, where $\mathbf{r}'(1) = -\mathbf{i} - 2\,\mathbf{j} - 3\,\mathbf{k}, \quad \mathbf{r}''(1) = 2\,\mathbf{i} + 6\,\mathbf{j} + 12\,\mathbf{k},$ and

$\mathbf{r}'(1)\times\mathbf{r}''(1) = -6\,\mathbf{i} + 6\,\mathbf{j} - 2\,\mathbf{k}.$ Thus at the point $(1,1,1)$, $a_T = \dfrac{\mathbf{r}'(1)\cdot\mathbf{r}''(1)}{|\mathbf{r}'(1)|} = \dfrac{-2-12-36}{\sqrt{1+4+9}} = -\dfrac{50}{\sqrt{14}}$ and

$$a_N = \frac{|\mathbf{r}'(1)\times\mathbf{r}''(1)|}{|\mathbf{r}'(1)|} = \frac{\sqrt{36+36+4}}{\sqrt{14}} = \sqrt{\frac{76}{14}} = \sqrt{\frac{38}{7}}.$$

43. The tangential component of $\mathbf{a}$ is the length of the projection of $\mathbf{a}$ onto $\mathbf{T}$, so we sketch the scalar projection of $\mathbf{a}$ in the tangential direction to the curve and estimate its length to be 4.5 (using the fact that $\mathbf{a}$ has length 10 as a guide). Similarly, the normal component of $\mathbf{a}$ is the length of the projection of $\mathbf{a}$ onto $\mathbf{N}$, so we sketch the scalar projection of $\mathbf{a}$ in the normal direction to the curve and estimate its length to be 9.0. Thus $a_T \approx 4.5$ cm/s^2 and $a_N \approx 9.0$ cm/s^2.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

44. $\mathbf{L}(t) = m\,\mathbf{r}(t) \times \mathbf{v}(t) \;\Rightarrow$

$$\begin{aligned}\mathbf{L}'(t) &= m[\mathbf{r}'(t) \times \mathbf{v}(t) + \mathbf{r}\,(t) \times \mathbf{v}'(t)] \qquad \text{[by Formula 5 of Theorem 13.2.3]}\\ &= m[\mathbf{v}(t) \times \mathbf{v}(t) + \mathbf{r}(t) \times \mathbf{v}'(t)] = m[\mathbf{0} + \mathbf{r}(t) \times \mathbf{a}\,(t)] = \boldsymbol{\tau}(t)\end{aligned}$$

So if the torque is always $\mathbf{0}$, then $\mathbf{L}'(t) = \mathbf{0}$ for all t, and so $\mathbf{L}(t)$ is constant.

45. If the engines are turned off at time t, then the spacecraft will continue to travel in the direction of $\mathbf{v}(t)$, so we need a t such that for some scalar $s > 0$, $\mathbf{r}(t) + s\,\mathbf{v}(t) = \langle 6, 4, 9\rangle$. $\mathbf{v}(t) = \mathbf{r}'(t) = \mathbf{i} + \dfrac{1}{t}\,\mathbf{j} + \dfrac{8t}{(t^2+1)^2}\,\mathbf{k} \;\Rightarrow$

$$\mathbf{r}(t) + s\,\mathbf{v}(t) = \left\langle 3 + t + s,\, 2 + \ln t + \frac{s}{t},\, 7 - \frac{4}{t^2+1} + \frac{8st}{(t^2+1)^2} \right\rangle \;\Rightarrow\; 3 + t + s = 6 \;\Rightarrow\; s = 3 - t,$$

$$\text{so } 7 - \frac{4}{t^2+1} + \frac{8(3-t)t}{(t^2+1)^2} = 9 \;\Leftrightarrow\; \frac{24t - 12t^2 - 4}{(t^2+1)^2} = 2 \;\Leftrightarrow\; t^4 + 8t^2 - 12t + 3 = 0.$$

It is easily seen that $t = 1$ is a root of this polynomial. Also $2 + \ln 1 + \dfrac{3-1}{1} = 4$, so $t = 1$ is the desired solution.

46. (a) $m\,\dfrac{d\mathbf{v}}{dt} = \dfrac{dm}{dt}\,\mathbf{v}_e \;\Leftrightarrow\; \dfrac{d\mathbf{v}}{dt} = \dfrac{1}{m}\dfrac{dm}{dt}\,\mathbf{v}_e$. Integrating both sides of this equation with respect to t gives

$$\int_0^t \frac{d\mathbf{v}}{du}\,du = \mathbf{v}_e \int_0^t \frac{1}{m}\frac{dm}{du}\,du \;\Rightarrow\; \int_{\mathbf{v}(0)}^{\mathbf{v}(t)} d\mathbf{v} = \mathbf{v}_e \int_{m(0)}^{m(t)} \frac{dm}{m} \quad \text{[Substitution Rule]} \;\Rightarrow$$

$$\mathbf{v}(t) - \mathbf{v}(0) = \ln\left(\frac{m(t)}{m(0)}\right)\mathbf{v}_e \;\Rightarrow\; \mathbf{v}(t) = \mathbf{v}(0) - \ln\left(\frac{m(0)}{m(t)}\right)\mathbf{v}_e.$$

(b) $|\mathbf{v}(t)| = 2\,|\mathbf{v}_e|$, and $|\mathbf{v}(0)| = 0$. Therefore, by part (a), $2\,|\mathbf{v}_e| = \left|-\ln\left(\dfrac{m(0)}{m(t)}\right)\mathbf{v}_e\right| \;\Rightarrow$

$$2\,|\mathbf{v}_e| = \ln\left(\frac{m(0)}{m\,(t)}\right)|\mathbf{v}_e|. \quad \left[\textit{Note: } m(0) > m(t) \text{ so that } \ln\left(\frac{m(0)}{m(t)}\right) > 0\right] \;\Rightarrow\; m(t) = e^{-2}m(0).$$

Thus $\dfrac{m(0) - e^{-2}m(0)}{m(0)} = 1 - e^{-2}$ is the fraction of the initial mass that is burned as fuel.

APPLIED PROJECT Kepler's Laws

1. With $\mathbf{r} = (r\cos\theta)\,\mathbf{i} + (r\sin\theta)\,\mathbf{j}$ and $\mathbf{h} = \alpha\,\mathbf{k}$ where $\alpha > 0$,

(a) $\mathbf{h} = \mathbf{r} \times \mathbf{r}' = [(r\cos\theta)\,\mathbf{i} + (r\sin\theta)\,\mathbf{j}] \times \left[\left(r'\cos\theta - r\sin\theta\,\dfrac{d\theta}{dt}\right)\mathbf{i} + \left(r'\sin\theta + r\cos\theta\,\dfrac{d\theta}{dt}\right)\mathbf{j}\right]$

$$= \left[rr'\cos\theta\sin\theta + r^2\cos^2\theta\,\frac{d\theta}{dt} - rr'\cos\theta\sin\theta + r^2\sin^2\theta\,\frac{d\theta}{dt}\right]\mathbf{k} = r^2\frac{d\theta}{dt}\,\mathbf{k}$$

(b) Since $\mathbf{h} = \alpha\,\mathbf{k}$, $\alpha > 0$, $\alpha = |\mathbf{h}|$. But by part (a), $\alpha = |\mathbf{h}| = r^2\,(d\theta/dt)$.

(c) $A(t) = \frac{1}{2}\int_{\theta_0}^{\theta} |\mathbf{r}|^2\,d\theta = \frac{1}{2}\int_{t_0}^{t} r^2\,(d\theta/dt)\,dt$ in polar coordinates. Thus, by the Fundamental Theorem of Calculus,

$$\frac{dA}{dt} = \frac{r^2}{2}\frac{d\theta}{dt}.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $\dfrac{dA}{dt} = \dfrac{r^2}{2}\dfrac{d\theta}{dt} = \dfrac{h}{2} =$ constant since $\mathbf{h}$ is a constant vector and $h = |\mathbf{h}|$.

2. (a) Since $dA/dt = \frac{1}{2}h$, a constant, $A(t) = \frac{1}{2}ht + c_1$. But $A(0) = 0$, so $A(t) = \frac{1}{2}ht$. But $A(T)$ = area of the ellipse $= \pi ab$ and $A(T) = \frac{1}{2}hT$, so $T = 2\pi ab/h$.

(b) $h^2/(GM) = ed$ where e is the eccentricity of the ellipse. But $a = ed/(1-e^2)$ or $ed = a(1-e^2)$ and $1 - e^2 = b^2/a^2$. Hence $h^2/(GM) = ed = b^2/a$.

(c) $T^2 = \dfrac{4\pi a^2 b^2}{h^2} = 4\pi^2 a^2 b^2 \dfrac{a}{GMb^2} = \dfrac{4\pi^2}{GM}a^3$.

3. From Problem 2, $T^2 = \dfrac{4\pi^2}{GM}a^3$. $T \approx 365.25$ days $\times\, 24 \cdot 60^2 \dfrac{\text{seconds}}{\text{day}} \approx 3.1558 \times 10^7$ seconds. Therefore

$a^3 = \dfrac{GMT^2}{4\pi^2} \approx \dfrac{(6.67 \times 10^{-11})(1.99 \times 10^{30})(3.1558 \times 10^7)^2}{4\pi^2} \approx 3.348 \times 10^{33}\ \text{m}^3 \quad \Rightarrow \quad a \approx 1.496 \times 10^{11}$ m. Thus, the length of the major axis of the earth's orbit (that is, $2a$) is approximately 2.99×10^{11} m $= 2.99 \times 10^8$ km.

4. We can adapt the equation $T^2 = \dfrac{4\pi^2}{GM}a^3$ from Problem 2(c) with the earth at the center of the system, so T is the period of the satellite's orbit about the earth, M is the mass of the earth, and a is the length of the semimajor axis of the satellite's orbit (measured from the earth's center). Since we want the satellite to remain fixed above a particular point on the earth's equator, T must coincide with the period of the earth's own rotation, so $T = 24$ h $= 86{,}400$ s. The mass of the earth is $M = 5.98 \times 10^{24}$ kg, so $a = \left(\dfrac{T^2GM}{4\pi^2}\right)^{1/3} \approx \left[\dfrac{(86{,}400)^2(6.67 \times 10^{-11})(5.98 \times 10^{24})}{4\pi^2}\right]^{1/3} \approx 4.23 \times 10^7$ m. If we assume a circular orbit, the radius of the orbit is a, and since the radius of the earth is 6.37×10^6 m, the required altitude above the earth's surface for the satellite is $4.23 \times 10^7 - 6.37 \times 10^6 \approx 3.59 \times 10^7$ m, or 35,900 km.

13 Review

TRUE-FALSE QUIZ

1. True. If we reparametrize the curve by replacing $u = t^3$, we have $\mathbf{r}(u) = u\,\mathbf{i} + 2u\,\mathbf{j} + 3u\,\mathbf{k}$, which is a line through the origin with direction vector $\mathbf{i} + 2\,\mathbf{j} + 3\,\mathbf{k}$.

2. True. Parametric equations for the curve are $x = 0$, $y = t^2$, $z = 4t$, and since $t = z/4$ we have $y = t^2 = (z/4)^2$ or $y = \frac{1}{16}z^2$, $x = 0$. This is an equation of a parabola in the yz-plane.

3. False. The vector function represents a line, but the line does not pass through the origin; the x-component is 0 only for $t = 0$ which corresponds to the point $(0, 3, 0)$ not $(0, 0, 0)$.

4. True. See Theorem 13.2.2.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

5. False. By Formula 5 of Theorem 13.2.3, $\dfrac{d}{dt}\left[\mathbf{u}(t)\times\mathbf{v}(t)\right]=\mathbf{u}'(t)\times\mathbf{v}(t)+\mathbf{u}(t)\times\mathbf{v}'(t)$.

6. False. For example, let $\mathbf{r}(t)=\langle\cos t,\sin t\rangle$. Then $|\mathbf{r}(t)|=\sqrt{\cos^2 t+\sin^2 t}=1 \;\Rightarrow\; \dfrac{d}{dt}\,|\mathbf{r}(t)|=0$, but $|\mathbf{r}'(t)|=|\langle-\sin t,\cos t\rangle|=\sqrt{(-\sin t)^2+\cos^2 t}=1$.

7. False. κ is the magnitude of the rate of change of the unit tangent vector $\mathbf{T}$ with respect to arc length s, not with respect to t.

8. False. The binormal vector, by the definition given in Section 13.3, is $\mathbf{B}(t)=\mathbf{T}(t)\times\mathbf{N}(t)=-\left[\mathbf{N}(t)\times\mathbf{T}(t)\right]$.

9. True. At an inflection point where f is twice continuously differentiable we must have $f''(x)=0$, and by Equation 13.3.11, the curvature is 0 there.

10. True. From Equation 13.3.9 , $\kappa(t)=0 \;\Leftrightarrow\; |\mathbf{T}'(t)|=0 \;\Leftrightarrow\; \mathbf{T}'(t)=\mathbf{0}$ for all t. But then $\mathbf{T}(t)=\mathbf{C}$, a constant vector, which is true only for a straight line.

11. False. If $\mathbf{r}(t)$ is the position of a moving particle at time t and $|\mathbf{r}(t)|=1$ then the particle lies on the unit circle or the unit sphere, but this does not mean that the speed $|\mathbf{r}'(t)|$ must be constant. As a counterexample, let $\mathbf{r}(t)=\left\langle t,\sqrt{1-t^2}\right\rangle$, then $\mathbf{r}'(t)=\left\langle 1,-t/\sqrt{1-t^2}\right\rangle$ and $|\mathbf{r}(t)|=\sqrt{t^2+1-t^2}=1$ but $|\mathbf{r}'(t)|=\sqrt{1+t^2/(1-t^2)}=1/\sqrt{1-t^2}$ which is not constant.

12. True. See Example 4 in Section 13.2 .

13. True. See the discussion preceding Example 7 in Section 13.3.

14. False. For example, $\mathbf{r}_1(t)=\langle t,t\rangle$ and $\mathbf{r}_2(t)=\langle 2t,2t\rangle$ both represent the same plane curve (the line $y=x$), but the tangent vector $\mathbf{r}_1'(t)=\langle 1,1\rangle$ for all t, while $\mathbf{r}_2'(t)=\langle 2,2\rangle$. In fact, different parametrizations give parallel tangent vectors at a point, but their magnitudes may differ.

EXERCISES

1. (a) The corresponding parametric equations for the curve are $x=t$, $y=\cos\pi t$, $z=\sin\pi t$. Since $y^2+z^2=1$, the curve is contained in a circular cylinder with axis the x-axis. Since $x=t$, the curve is a helix.



(b) $\mathbf{r}(t)=t\,\mathbf{i}+\cos\pi t\,\mathbf{j}+\sin\pi t\,\mathbf{k} \;\Rightarrow$
$\mathbf{r}'(t)=\mathbf{i}-\pi\sin\pi t\,\mathbf{j}+\pi\cos\pi t\,\mathbf{k} \;\Rightarrow$
$\mathbf{r}''(t)=-\pi^2\cos\pi t\,\mathbf{j}-\pi^2\sin\pi t\,\mathbf{k}$

2. (a) The expressions $\sqrt{2-t}$, $(e^t-1)/t$, and $\ln(t+1)$ are all defined when $2-t\geq 0 \;\Rightarrow\; t\leq 2, t\neq 0$, and $t+1>0 \;\Rightarrow\; t>-1$. Thus the domain of $\mathbf{r}$ is $(-1,0)\cup(0,2]$.

(b) $\displaystyle\lim_{t\to 0}\mathbf{r}(t)=\left\langle\lim_{t\to 0}\sqrt{2-t},\lim_{t\to 0}\frac{e^t-1}{t},\lim_{t\to 0}\ln(t+1)\right\rangle=\left\langle\sqrt{2-0},\lim_{t\to 0}\frac{e^t}{1},\ln(0+1)\right\rangle$
$=\langle\sqrt{2},1,0\rangle$ [using l'Hospital's Rule in the y-component]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\mathbf{r}'(t) = \left\langle \frac{d}{dt}\sqrt{2-t}, \frac{d}{dt}\frac{e^t-1}{t}, \frac{d}{dt}\ln(t+1) \right\rangle = \left\langle -\frac{1}{2\sqrt{2-t}}, \frac{te^t - e^t + 1}{t^2}, \frac{1}{t+1} \right\rangle$

3. The projection of the curve C of intersection onto the xy-plane is the circle $x^2 + y^2 = 16, z = 0$. So we can write $x = 4\cos t,\ y = 4\sin t,\ 0 \le t \le 2\pi$. From the equation of the plane, we have $z = 5 - x = 5 - 4\cos t$, so parametric equations for C are $x = 4\cos t,\ y = 4\sin t,\ z = 5 - 4\cos t, 0 \le t \le 2\pi$, and the corresponding vector function is $\mathbf{r}(t) = 4\cos t\,\mathbf{i} + 4\sin t\,\mathbf{j} + (5 - 4\cos t)\,\mathbf{k}, 0 \le t \le 2\pi$.

4. The curve is given by $\mathbf{r}(t) = \langle 2\sin t, 2\sin 2t, 2\sin 3t\rangle$, so $\mathbf{r}'(t) = \langle 2\cos t, 4\cos 2t, 6\cos 3t\rangle$. The point $\left(1, \sqrt{3}, 2\right)$ corresponds to $t = \frac{\pi}{6}$ (or $\frac{\pi}{6} + 2k\pi$, k an integer), so the tangent vector there is $\mathbf{r}'(\frac{\pi}{6}) = \langle \sqrt{3}, 2, 0\rangle$. Then the tangent line has direction vector $\langle \sqrt{3}, 2, 0\rangle$ and includes the point $\left(1, \sqrt{3}, 2\right)$, so parametric equations are $x = 1 + \sqrt{3}\,t,\ y = \sqrt{3} + 2t,\ z = 2$.



5. $\int_0^1 (t^2\,\mathbf{i} + t\cos\pi t\,\mathbf{j} + \sin\pi t\,\mathbf{k})\,dt = \left(\int_0^1 t^2\,dt\right)\mathbf{i} + \left(\int_0^1 t\cos\pi t\,dt\right)\mathbf{j} + \left(\int_0^1 \sin\pi t\,dt\right)\mathbf{k}$

$= \left[\frac{1}{3}t^3\right]_0^1 \mathbf{i} + \left(\frac{t}{\pi}\sin\pi t\right]_0^1 - \int_0^1 \frac{1}{\pi}\sin\pi t\,dt\Big)\mathbf{j} + \left[-\frac{1}{\pi}\cos\pi t\right]_0^1 \mathbf{k}$

$= \frac{1}{3}\mathbf{i} + \left[\frac{1}{\pi^2}\cos\pi t\right]_0^1 \mathbf{j} + \frac{2}{\pi}\mathbf{k} = \frac{1}{3}\mathbf{i} - \frac{2}{\pi^2}\mathbf{j} + \frac{2}{\pi}\mathbf{k}$

where we integrated by parts in the y-component.

6. (a) C intersects the xz-plane where $y = 0 \ \Rightarrow\ 2t - 1 = 0 \ \Rightarrow\ t = \frac{1}{2}$, so the point is $\left(2 - \left(\frac{1}{2}\right)^3, 0, \ln\frac{1}{2}\right) = \left(\frac{15}{8}, 0, -\ln 2\right)$.

(b) The curve is given by $\mathbf{r}(t) = \langle 2 - t^3, 2t - 1, \ln t\rangle$, so $\mathbf{r}'(t) = \langle -3t^2, 2, 1/t\rangle$. The point $(1, 1, 0)$ corresponds to $t = 1$, so the tangent vector there is $\mathbf{r}'(1) = \langle -3, 2, 1\rangle$. Then the tangent line has direction vector $\langle -3, 2, 1\rangle$ and includes the point $(1, 1, 0)$, so parametric equations are $x = 1 - 3t,\ y = 1 + 2t,\ z = t$.

(c) The normal plane has normal vector $\mathbf{r}'(1) = \langle -3, 2, 1\rangle$ and equation $-3(x - 1) + 2(y - 1) + z = 0$ or $3x - 2y - z = 1$.

7. $\mathbf{r}(t) = \langle t^2, t^3, t^4\rangle \ \Rightarrow\ \mathbf{r}'(t) = \langle 2t, 3t^2, 4t^3\rangle \ \Rightarrow\ |\mathbf{r}'(t)| = \sqrt{4t^2 + 9t^4 + 16t^6}$ and $L = \int_0^3 |\mathbf{r}'(t)|\,dt = \int_0^3 \sqrt{4t^2 + 9t^4 + 16t^6}\,dt$. Using Simpson's Rule with $f(t) = \sqrt{4t^2 + 9t^4 + 16t^6}$ and $n = 6$ we have $\Delta t = \frac{3-0}{6} = \frac{1}{2}$ and

$$\begin{aligned} L &\approx \tfrac{\Delta t}{3}\left[f(0) + 4f\left(\tfrac{1}{2}\right) + 2f(1) + 4f\left(\tfrac{3}{2}\right) + 2f(2) + 4f\left(\tfrac{5}{2}\right) + f(3)\right] \\ &= \tfrac{1}{6}\Bigg[\sqrt{0+0+0} + 4\cdot\sqrt{4\left(\tfrac{1}{2}\right)^2 + 9\left(\tfrac{1}{2}\right)^4 + 16\left(\tfrac{1}{2}\right)^6} + 2\cdot\sqrt{4(1)^2 + 9(1)^4 + 16(1)^6} \\ &\qquad + 4\cdot\sqrt{4\left(\tfrac{3}{2}\right)^2 + 9\left(\tfrac{3}{2}\right)^4 + 16\left(\tfrac{3}{2}\right)^6} + 2\cdot\sqrt{4(2)^2 + 9(2)^4 + 16(2)^6} \\ &\qquad + 4\cdot\sqrt{4\left(\tfrac{5}{2}\right)^2 + 9\left(\tfrac{5}{2}\right)^4 + 16\left(\tfrac{5}{2}\right)^6} + \sqrt{4(3)^2 + 9(3)^4 + 16(3)^6}\Bigg] \\ &\approx 86.631 \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

8. $\mathbf{r}'(t) = \left\langle 3t^{1/2}, -2\sin 2t, 2\cos 2t \right\rangle$, $|\mathbf{r}'(t)| = \sqrt{9t + 4(\sin^2 2t + \cos^2 2t)} = \sqrt{9t+4}$.

Thus $L = \int_0^1 \sqrt{9t+4}\,dt = \int_4^{13} \frac{1}{9} u^{1/2}\,du = \frac{1}{9}\cdot\frac{2}{3}u^{3/2}\Big]_4^{13} = \frac{2}{27}(13^{3/2} - 8)$.

9. The angle of intersection of the two curves, θ, is the angle between their respective tangents at the point of intersection.

For both curves the point $(1, 0, 0)$ occurs when $t = 0$.

$\mathbf{r}_1'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j} + \mathbf{k} \;\Rightarrow\; \mathbf{r}_1'(0) = \mathbf{j} + \mathbf{k}$ and $\mathbf{r}_2'(t) = \mathbf{i} + 2t\,\mathbf{j} + 3t^2\,\mathbf{k} \;\Rightarrow\; \mathbf{r}_2'(0) = \mathbf{i}$.

$\mathbf{r}_1'(0)\cdot\mathbf{r}_2'(0) = (\mathbf{j}+\mathbf{k})\cdot\mathbf{i} = 0$. Therefore, the curves intersect in a right angle, that is, $\theta = 90°$.

10. The parametric value corresponding to the point $(1, 0, 1)$ is $t = 0$.

$\mathbf{r}'(t) = e^t\,\mathbf{i} + e^t(\cos t + \sin t)\,\mathbf{j} + e^t(\cos t - \sin t)\,\mathbf{k} \;\Rightarrow\; |\mathbf{r}'(t)| = e^t\sqrt{1 + (\cos t + \sin t)^2 + (\cos t - \sin t)^2} = \sqrt{3}\,e^t$

and $s(t) = \int_0^t e^u\sqrt{3}\,du = \sqrt{3}(e^t - 1) \;\Rightarrow\; t = \ln\left(1 + \frac{1}{\sqrt{3}}s\right)$.

Therefore, $\mathbf{r}(t(s)) = \left(1 + \frac{1}{\sqrt{3}}s\right)\mathbf{i} + \left(1 + \frac{1}{\sqrt{3}}s\right)\sin\ln\left(1 + \frac{1}{\sqrt{3}}s\right)\mathbf{j} + \left(1 + \frac{1}{\sqrt{3}}s\right)\cos\ln\left(1 + \frac{1}{\sqrt{3}}s\right)\mathbf{k}$.

11. (a) $\mathbf{r}(t) = \langle \sin^3 t, \cos^3 t, \sin^2 t\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 3\sin^2 t\cos t, -3\cos^2 t\sin t, 2\sin t\cos t\rangle$,

$$|\mathbf{r}'(t)| = \sqrt{9\sin^4 t\cos^2 t + 9\cos^4 t\sin^2 t + 4\sin^2 t\cos^2 t}$$
$$= \sqrt{\sin^2 t\cos^2 t\,(9\sin^2 t + 9\cos^2 t + 4)} = \sqrt{13}\,\sin t\cos t \quad [\text{since } 0 \le t \le \pi/2 \;\Rightarrow\; \sin t, \cos t \ge 0]$$

Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{13}\,\sin t\cos t}\langle 3\sin^2 t\cos t, -3\cos^2 t\sin t, 2\sin t\cos t\rangle = \frac{1}{\sqrt{13}}\langle 3\sin t, -3\cos t, 2\rangle$.

(b) $\mathbf{T}'(t) = \frac{1}{\sqrt{13}}\langle 3\cos t, 3\sin t, 0\rangle$, $|\mathbf{T}'(t)| = \frac{1}{\sqrt{13}}\sqrt{9\cos^2 t + 9\sin^2 t + 0} = \frac{3}{\sqrt{13}}$, and

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \frac{1}{3}\langle 3\cos t, 3\sin t, 0\rangle = \langle \cos t, \sin t, 0\rangle$.

(c) $\mathbf{B}(t) = \mathbf{T}(t)\times\mathbf{N}(t) = \frac{1}{\sqrt{13}}\langle 3\sin t, -3\cos t, 2\rangle \times \langle \cos t, \sin t, 0\rangle = \frac{1}{\sqrt{13}}\langle -2\sin t, 2\cos t, 3\rangle$

(d) $\kappa(t) \quad \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{3\sqrt{13}}{\sqrt{13}\,\sin t\cos t} = \dfrac{3}{13\,\sin t\cos t}$ or $\frac{3}{13}\sec t\csc t$

12. Using Exercise 13.3.42, we have $\mathbf{r}'(t) = \langle -3\sin t, 4\cos t\rangle$, $\mathbf{r}''(t) = \langle -3\cos t, -4\sin t\rangle$,

$|\mathbf{r}'(t)|^3 = \left(\sqrt{9\sin^2 t + 4\cos^2 t}\right)^3$ and then

$$\kappa(t) = \frac{|(-3\sin t)(-4\sin t) - (4\cos t)(-3\cos t)|}{(9\sin^2 t + 16\cos^2 t)^{3/2}} = \frac{12}{(9\sin^2 t + 16\cos^2 t)^{3/2}}.$$

At $(3, 0)$, $t = 0$ and $\kappa(0) = 12/(16)^{3/2} = \frac{12}{64} = \frac{3}{16}$. At $(0, 4)$, $t = \frac{\pi}{2}$ and $\kappa\left(\frac{\pi}{2}\right) = 12/9^{3/2} = \frac{12}{27} = \frac{4}{9}$.

13. $y' = 4x^3$, $y'' = 12x^2$ and $\kappa(x) = \dfrac{|y''|}{[1 + (y')^2]^{3/2}} = \dfrac{|12x^2|}{(1 + 16x^6)^{3/2}}$, so $\kappa(1) = \dfrac{12}{17^{3/2}}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $\kappa(x) = \dfrac{|12x^2 - 2|}{[1 + (4x^3 - 2x)^2]^{3/2}} \Rightarrow \kappa(0) = 2.$

So the osculating circle has radius $\frac{1}{2}$ and center $(0, -\frac{1}{2})$.

Thus its equation is $x^2 + \left(y + \frac{1}{2}\right)^2 = \frac{1}{4}$.



15. $\mathbf{r}(t) = \langle \sin 2t, t, \cos 2t \rangle \Rightarrow \mathbf{r}'(t) = \langle 2\cos 2t, 1, -2\sin 2t \rangle \Rightarrow \mathbf{T}(t) = \frac{1}{\sqrt{5}} \langle 2\cos 2t, 1, -2\sin 2t \rangle \Rightarrow$

$\mathbf{T}'(t) = \frac{1}{\sqrt{5}} \langle -4\sin 2t, 0, -4\cos 2t \rangle \Rightarrow \mathbf{N}(t) = \langle -\sin 2t, 0, -\cos 2t \rangle$. So $\mathbf{N} = \mathbf{N}(\pi) = \langle 0, 0, -1 \rangle$ and

$\mathbf{B} = \mathbf{T} \times \mathbf{N} = \frac{1}{\sqrt{5}} \langle -1, 2, 0 \rangle$. So a normal to the osculating plane is $\langle -1, 2, 0 \rangle$ and an equation is

$-1(x - 0) + 2(y - \pi) + 0(z - 1) = 0$ or $x - 2y + 2\pi = 0$.

16. (a) The average velocity over $[3, 3.2]$ is given by

$\dfrac{\mathbf{r}(3.2) - \mathbf{r}(3)}{3.2 - 3} = 5[\mathbf{r}(3.2) - \mathbf{r}(3)]$, so we draw a vector with the same direction but 5 times the length of the vector $[\mathbf{r}(3.2) - \mathbf{r}(3)]$.



(b) $\mathbf{v}(3) = \mathbf{r}'(3) = \lim\limits_{h \to 0} \dfrac{\mathbf{r}(3+h) - \mathbf{r}(3)}{h}$

(c) $\mathbf{T}(3) = \dfrac{\mathbf{r}'(3)}{|\mathbf{r}'(3)|}$, a unit vector in the same direction as $\mathbf{r}'(3)$, that is, parallel to the tangent line to the curve at $\mathbf{r}(3)$, pointing in the direction corresponding to increasing t, and with length 1.



17. $\mathbf{r}(t) = t\ln t\,\mathbf{i} + t\,\mathbf{j} + e^{-t}\,\mathbf{k}$, $\mathbf{v}(t) = \mathbf{r}'(t) = (1 + \ln t)\,\mathbf{i} + \mathbf{j} - e^{-t}\,\mathbf{k}$,

$|\mathbf{v}(t)| = \sqrt{(1 + \ln t)^2 + 1^2 + (-e^{-t})^2} = \sqrt{2 + 2\ln t + (\ln t)^2 + e^{-2t}}$, $\mathbf{a}(t) = \mathbf{v}'(t) = \frac{1}{t}\,\mathbf{i} + e^{-t}\,\mathbf{k}$

18. $\mathbf{r}(t) = (2t^2 - 3)\,\mathbf{i} + 2t\,\mathbf{j} \Rightarrow \mathbf{v}(t) = \mathbf{r}'(t) = 4t\,\mathbf{i} + 2\,\mathbf{j}$,

speed $= |\mathbf{v}(t)| = \sqrt{16t^2 + 4} = 2\sqrt{4t^2 + 1}$, and $\mathbf{a}(t) = \mathbf{v}'(t) = \mathbf{r}''(t) = 4\,\mathbf{i}$.

At $t = 1$ we have $\mathbf{r}(1) = -\mathbf{i} + 2\,\mathbf{j}$, $\mathbf{v}(1) = 4\,\mathbf{i} + 2\,\mathbf{j}$, $\mathbf{a}(1) = 4\,\mathbf{i}$.

Notice that $y/2 = t \Rightarrow x = 2(y/2)^2 - 3 = \frac{1}{2}y^2 - 3$, so the path is a parabola.



19. $\mathbf{v}(t) = \int \mathbf{a}(t)\,dt = \int (6t\,\mathbf{i} + 12t^2\,\mathbf{j} - 6t\,\mathbf{k})\,dt = 3t^2\,\mathbf{i} + 4t^3\,\mathbf{j} - 3t^2\,\mathbf{k} + \mathbf{C}$, but $\mathbf{i} - \mathbf{j} + 3\,\mathbf{k} = \mathbf{v}(0) = 0 + \mathbf{C}$,

so $\mathbf{C} = \mathbf{i} - \mathbf{j} + 3\,\mathbf{k}$ and $\mathbf{v}(t) = (3t^2 + 1)\,\mathbf{i} + (4t^3 - 1)\,\mathbf{j} + (3 - 3t^2)\,\mathbf{k}$.

$\mathbf{r}(t) = \int \mathbf{v}(t)\,dt = (t^3 + t)\,\mathbf{i} + (t^4 - t)\,\mathbf{j} + (3t - t^3)\,\mathbf{k} + \mathbf{D}$.

But $\mathbf{r}(0) = \mathbf{0}$, so $\mathbf{D} = \mathbf{0}$ and $\mathbf{r}(t) = (t^3 + t)\,\mathbf{i} + (t^4 - t)\,\mathbf{j} + (3t - t^3)\,\mathbf{k}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. We set up the axes so that the shot leaves the athlete's hand 7 ft above the origin. Then we are given $\mathbf{r}(0) = 7\mathbf{j}$, $|\mathbf{v}(0)| = 43$ ft/s, and $\mathbf{v}(0)$ has direction given by a 45° angle of elevation. Then a unit vector in the direction of $\mathbf{v}(0)$ is $\frac{1}{\sqrt{2}}(\mathbf{i}+\mathbf{j}) \Rightarrow \mathbf{v}(0) = \frac{43}{\sqrt{2}}(\mathbf{i}+\mathbf{j})$. Assuming air resistance is negligible, the only external force is due to gravity, so as in Example 13.4.5 we have $\mathbf{a} = -g\,\mathbf{j}$ where here $g \approx 32$ ft/s^2. Since $\mathbf{v}'(t) = \mathbf{a}(t)$, we integrate, giving $\mathbf{v}(t) = -gt\,\mathbf{j} + \mathbf{C}$ where $\mathbf{C} = \mathbf{v}(0) = \frac{43}{\sqrt{2}}(\mathbf{i}+\mathbf{j}) \Rightarrow \mathbf{v}(t) = \frac{43}{\sqrt{2}}\,\mathbf{i} + \left(\frac{43}{\sqrt{2}} - gt\right)\mathbf{j}$. Since $\mathbf{r}'(t) = \mathbf{v}(t)$ we integrate again, so $\mathbf{r}(t) = \frac{43}{\sqrt{2}}t\,\mathbf{i} + \left(\frac{43}{\sqrt{2}}t - \frac{1}{2}gt^2\right)\mathbf{j} + \mathbf{D}$. But $\mathbf{D} = \mathbf{r}(0) = 7\,\mathbf{j} \Rightarrow \mathbf{r}(t) = \frac{43}{\sqrt{2}}t\,\mathbf{i} + \left(\frac{43}{\sqrt{2}}t - \frac{1}{2}gt^2 + 7\right)\mathbf{j}$.

(a) At 2 seconds, the shot is at $\mathbf{r}(2) = \frac{43}{\sqrt{2}}(2)\,\mathbf{i} + \left(\frac{43}{\sqrt{2}}(2) - \frac{1}{2}g(2)^2 + 7\right)\mathbf{j} \approx 60.8\,\mathbf{i} + 3.8\,\mathbf{j}$, so the shot is about 3.8 ft above the ground, at a horizontal distance of 60.8 ft from the athlete.

(b) The shot reaches its maximum height when the vertical component of velocity is 0: $\frac{43}{\sqrt{2}} - gt = 0 \Rightarrow$ $t = \dfrac{43}{\sqrt{2}\,g} \approx 0.95$ s. Then $\mathbf{r}(0.95) \approx 28.9\,\mathbf{i} + 21.4\,\mathbf{j}$, so the maximum height is approximately 21.4 ft.

(c) The shot hits the ground when the vertical component of $\mathbf{r}(t)$ is 0, so $\frac{43}{\sqrt{2}}t - \frac{1}{2}gt^2 + 7 = 0 \Rightarrow$ $-16t^2 + \frac{43}{\sqrt{2}}t + 7 = 0 \Rightarrow t \approx 2.11$ s. $\mathbf{r}(2.11) \approx 64.2\,\mathbf{i} - 0.08\,\mathbf{j}$, thus the shot lands approximately 64.2 ft from the athlete.

21. Example 13.4.5 showed that the maximum horizontal range is achieved with an angle of elevation of 45°. In this case, however, the projectile would hit the top of the tunnel using that angle. The horizontal range will be maximized with the largest angle of elevation that keeps the projectile within a height of 30 m. From Example 13.4.5 we know that the position function of the projectile is $\mathbf{r}(t) = (v_0\cos\alpha)t\,\mathbf{i} + \left[(v_0\sin\alpha)t - \frac{1}{2}gt^2\right]\mathbf{j}$ and the velocity is $\mathbf{v}(t) = \mathbf{r}'(t) = (v_0\cos\alpha)\,\mathbf{i} + [(v_0\sin\alpha) - gt]\,\mathbf{j}$. The projectile achieves its maximum height when the vertical component of velocity is zero, so $(v_0\sin\alpha) - gt = 0 \Rightarrow t = \dfrac{v_0\sin\alpha}{g}$. We want the vertical height of the projectile at that time to be 30 m: $(v_0\sin\alpha)\left(\dfrac{v_0\sin\alpha}{g}\right) - \frac{1}{2}g\left(\dfrac{v_0\sin\alpha}{g}\right)^2 = 30 \Rightarrow$ $\left(\dfrac{v_0^2\sin^2\alpha}{g}\right) - \dfrac{1}{2}\left(\dfrac{v_0^2\sin^2\alpha}{g}\right) = 30 \Rightarrow \dfrac{v_0^2\sin^2\alpha}{2g} = 30 \Rightarrow \sin^2\alpha = \dfrac{30(2g)}{v_0^2} = \dfrac{60(9.8)}{40^2} = 0.3675 \Rightarrow$ $\sin\alpha = \sqrt{0.3675}$. Thus the desired angle of elevation is $\alpha = \sin^{-1}\sqrt{0.3675} \approx 37.3^\circ$.

From the same example, the horizontal distance traveled is $d = \dfrac{v_0^2\sin 2\alpha}{g} \approx \dfrac{40^2\sin(74.6^\circ)}{9.8} \approx 157.4$ m.

22. $\mathbf{r}'(t) = \mathbf{i} + 2\,\mathbf{j} + 2t\,\mathbf{k}$, $\mathbf{r}''(t) = 2\,\mathbf{k}$, $|\mathbf{r}'(t)| = \sqrt{1+4+4t^2} = \sqrt{4t^2+5}$.

Then $a_T = \dfrac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \dfrac{4t}{\sqrt{4t^2+5}}$ and $a_N = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{|4\,\mathbf{i} - 2\,\mathbf{j}|}{\sqrt{4t^2+5}} = \dfrac{2\sqrt{5}}{\sqrt{4t^2+5}}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. (a) Instead of proceeding directly, we use Formula 3 of Theorem 13.2.3: $\mathbf{r}(t) = t\,\mathbf{R}(t) \;\Rightarrow$

$$\mathbf{v} = \mathbf{r}'(t) = \mathbf{R}(t) + t\,\mathbf{R}'(t) = \cos\omega t\,\mathbf{i} + \sin\omega t\,\mathbf{j} + t\,\mathbf{v}_d.$$

(b) Using the same method as in part (a) and starting with $\mathbf{v} = \mathbf{R}(t) + t\,\mathbf{R}'(t)$, we have

$$\mathbf{a} = \mathbf{v}' = \mathbf{R}'(t) + \mathbf{R}'(t) + t\,\mathbf{R}''(t) = 2\,\mathbf{R}'(t) + t\,\mathbf{R}''(t) = 2\,\mathbf{v}_d + t\,\mathbf{a}_d.$$

(c) Here we have $\mathbf{r}(t) = e^{-t}\cos\omega t\,\mathbf{i} + e^{-t}\sin\omega t\,\mathbf{j} = e^{-t}\,\mathbf{R}(t)$. So, as in parts (a) and (b),

$$\mathbf{v} = \mathbf{r}'(t) = e^{-t}\,\mathbf{R}'(t) - e^{-t}\,\mathbf{R}(t) = e^{-t}[\mathbf{R}'(t) - \mathbf{R}(t)] \;\Rightarrow$$

$$\mathbf{a} = \mathbf{v}' = e^{-t}[\mathbf{R}''(t) - \mathbf{R}'(t)] - e^{-t}[\mathbf{R}'(t) - \mathbf{R}(t)] = e^{-t}[\mathbf{R}''(t) - 2\,\mathbf{R}'(t) + \mathbf{R}(t)]$$
$$= e^{-t}\,\mathbf{a}_d - 2e^{-t}\,\mathbf{v}_d + e^{-t}\,\mathbf{R}$$

Thus, the Coriolis acceleration (the sum of the "extra" terms not involving $\mathbf{a}_d$) is $-2e^{-t}\,\mathbf{v}_d + e^{-t}\,\mathbf{R}$.

24. (a) $F(x) = \begin{cases} 1 & \text{if } x \le 0 \\ \sqrt{1-x^2} & \text{if } 0 < x < \frac{1}{\sqrt{2}} \\ \sqrt{2} - x & \text{if } x \ge \frac{1}{\sqrt{2}} \end{cases} \quad \Rightarrow \quad F'(x) = \begin{cases} 0 & \text{if } x < 0 \\ -x/\sqrt{1-x^2} & \text{if } 0 < x < \frac{1}{\sqrt{2}} \\ -1 & \text{if } x > \frac{1}{\sqrt{2}} \end{cases} \quad \Rightarrow$

$$F''(x) = \begin{cases} 0 & \text{if } x < 0 \\ -1/(1-x^2)^{3/2} & \text{if } 0 < x < \frac{1}{\sqrt{2}} \\ 0 & \text{if } x > \frac{1}{\sqrt{2}} \end{cases}$$

since $\dfrac{d}{dx}[-x(1-x^2)^{-1/2}] = -(1-x^2)^{-1/2} - x^2(1-x^2)^{-3/2} = -(1-x^2)^{-3/2}$.

Now $\lim\limits_{x\to 0^+}\sqrt{1-x^2} = 1 = F(0)$ and $\lim\limits_{x\to(1/\sqrt{2})^-}\sqrt{1-x^2} = \frac{1}{\sqrt{2}} = F\left(\frac{1}{\sqrt{2}}\right)$, so F is continuous. Also, since

$\lim\limits_{x\to 0^+} F'(x) = 0 = \lim\limits_{x\to 0} F'(x)$ and $\lim\limits_{x\to(1/\sqrt{2})^-} F'(x) = -1 = \lim\limits_{x\to(1/\sqrt{2})^+} F'(x)$, F' is continuous. But

$\lim\limits_{x\to 0^+} F''(x) = -1 \ne 0 = \lim\limits_{x\to 0^-} F''(x)$, so F'' is not continuous at $x = 0$. (The same is true at $x = \frac{1}{\sqrt{2}}$.)

So F does not have continuous curvature.

(b) Set $P(x) = ax^5 + bx^4 + cx^3 + dx^2 + ex + f$. The continuity conditions on P are $P(0) = 0$, $P(1) = 1$, $P'(0) = 0$ and $P'(1) = 1$. Also the curvature must be continuous. For $x \le 0$ and $x \ge 1$, $\kappa(x) = 0$; elsewhere

$\kappa(x) = \dfrac{|P''(x)|}{(1 + [P'(x)]^2)^{3/2}}$, so we need $P''(0) = 0$ and $P''(1) = 0$.

The conditions $P(0) = P'(0) = P''(0) = 0$ imply that $d = e = f = 0$.
The other conditions imply that $a + b + c = 1$, $5a + 4b + 3c = 1$, and
$10a + 6b + 3c = 0$. From these, we find that $a = 3$, $b = -8$, and $c = 6$.
Therefore $P(x) = 3x^5 - 8x^4 + 6x^3$. Since there was no solution with
$a = 0$, this could not have been done with a polynomial of degree 4.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

PROBLEMS PLUS

1. (a) $\mathbf{r}(t) = R\cos\omega t\,\mathbf{i} + R\sin\omega t\,\mathbf{j} \;\Rightarrow\; \mathbf{v} = \mathbf{r}'(t) = -\omega R\sin\omega t\,\mathbf{i} + \omega R\cos\omega t\,\mathbf{j}$, so $\mathbf{r} = R(\cos\omega t\,\mathbf{i} + \sin\omega t\,\mathbf{j})$ and $\mathbf{v} = \omega R(-\sin\omega t\,\mathbf{i} + \cos\omega t\,\mathbf{j})$. $\mathbf{v}\cdot\mathbf{r} = \omega R^2(-\cos\omega t\sin\omega t + \sin\omega t\cos\omega t) = 0$, so $\mathbf{v}\perp\mathbf{r}$. Since $\mathbf{r}$ points along a radius of the circle, and $\mathbf{v}\perp\mathbf{r}$, $\mathbf{v}$ is tangent to the circle. Because it is a velocity vector, $\mathbf{v}$ points in the direction of motion.

(b) In (a), we wrote $\mathbf{v}$ in the form $\omega R\,\mathbf{u}$, where $\mathbf{u}$ is the unit vector $-\sin\omega t\,\mathbf{i} + \cos\omega t\,\mathbf{j}$. Clearly $|\mathbf{v}| = \omega R\,|\mathbf{u}| = \omega R$. At speed ωR, the particle completes one revolution, a distance $2\pi R$, in time $T = \dfrac{2\pi R}{\omega R} = \dfrac{2\pi}{\omega}$.

(c) $\mathbf{a} = \dfrac{d\mathbf{v}}{dt} = -\omega^2 R\cos\omega t\,\mathbf{i} - \omega^2 R\sin\omega t\,\mathbf{j} = -\omega^2 R(\cos\omega t\,\mathbf{i} + \sin\omega t\,\mathbf{j})$, so $\mathbf{a} = -\omega^2\mathbf{r}$. This shows that $\mathbf{a}$ is proportional to $\mathbf{r}$ and points in the opposite direction (toward the origin). Also, $|\mathbf{a}| = \omega^2\,|\mathbf{r}| = \omega^2 R$. $\leq$

(d) By Newton's Second Law (see Section 13.4), $\mathbf{F} = m\mathbf{a}$, so $|\mathbf{F}| = m\,|\mathbf{a}| = mR\omega^2 = \dfrac{m\,(\omega R)^2}{R} = \dfrac{m\,|\mathbf{v}|^2}{R}$.

2. (a) Dividing the equation $|\mathbf{F}|\sin\theta = \dfrac{mv_R^2}{R}$ by the equation $|\mathbf{F}|\cos\theta = mg$, we obtain $\tan\theta = \dfrac{v_R^2}{Rg}$, so $v_R^2 = Rg\tan\theta$.

(b) $R = 400$ ft and $\theta = 12°$, so $v_R = \sqrt{Rg\tan\theta} \approx \sqrt{400\cdot 32\cdot\tan 12°} \approx 52.16$ ft/s ≈ 36 mi/h.

(c) We want to choose a new radius R_1 for which the new rated speed is $\frac{3}{2}$ of the old one: $\sqrt{R_1 g\tan 12°} = \frac{3}{2}\sqrt{Rg\tan 12°}$. Squaring, we get $R_1 g\tan 12° = \frac{9}{4}Rg\tan 12°$, so $R_1 = \frac{9}{4}R = \frac{9}{4}(400) = 900$ ft.

3. (a) The projectile reaches maximum height when $0 = \dfrac{dy}{dt} = \dfrac{d}{dt}\left[(v_0\sin\alpha)t - \frac{1}{2}gt^2\right] = v_0\sin\alpha - gt$; that is, when $t = \dfrac{v_0\sin\alpha}{g}$ and $y = (v_0\sin\alpha)\left(\dfrac{v_0\sin\alpha}{g}\right) - \dfrac{1}{2}g\left(\dfrac{v_0\sin\alpha}{g}\right)^2 = \dfrac{v_0^2\sin^2\alpha}{2g}$. This is the maximum height attained when the projectile is fired with an angle of elevation α. This maximum height is largest when $\alpha = 90°$. In that case, $\sin\alpha = 1$ and the maximum height is $\dfrac{v_0^2}{2g}$.

(b) Let $R = v_0^2/g$. We are asked to consider the parabola $x^2 + 2Ry - R^2 = 0$ which can be rewritten as $y = -\dfrac{1}{2R}x^2 + \dfrac{R}{2}$. The points on or inside this parabola are those for which $-R \le x \le R$ and $0 \le y \le \dfrac{-1}{2R}x^2 + \dfrac{R}{2}$. When the projectile is fired at angle of elevation α, the points (x, y) along its path satisfy the relations $x = (v_0\cos\alpha)\,t$ and $y = (v_0\sin\alpha)t - \frac{1}{2}gt^2$, where $0 \le t \le (2v_0\sin\alpha)/g$ (as in Example 13.4.5). Thus $|x| \le \left|v_0\cos\alpha\left(\dfrac{2v_0\sin\alpha}{g}\right)\right| = \left|\dfrac{v_0^2}{g}\sin 2\alpha\right| \le \left|\dfrac{v_0^2}{g}\right| = |R|$. This shows that $-R \le x \le R$.

For t in the specified range, we also have $y = t\left(v_0\sin\alpha - \frac{1}{2}gt\right) = \frac{1}{2}gt\left(\dfrac{2v_0\sin\alpha}{g} - t\right) \ge 0$ and

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$y = (v_0 \sin\alpha)\dfrac{x}{v_0\cos\alpha} - \dfrac{g}{2}\left(\dfrac{x}{v_0\cos\alpha}\right)^2 = (\tan\alpha)\,x - \dfrac{g}{2v_0^2\cos^2\alpha}x^2 = -\dfrac{1}{2R\cos^2\alpha}\,x^2 + (\tan\alpha)\,x$. Thus

$$\begin{aligned} y - \left(\frac{-1}{2R}x^2 + \frac{R}{2}\right) &= \frac{-1}{2R\cos^2\alpha}\,x^2 + \frac{1}{2R}x^2 + (\tan\alpha)\,x - \frac{R}{2} \\ &= \frac{x^2}{2R}\left(1 - \frac{1}{\cos^2\alpha}\right) + (\tan\alpha)\,x - \frac{R}{2} = \frac{x^2(1-\sec^2\alpha) + 2R(\tan\alpha)\,x - R^2}{2R} \\ &= \frac{-(\tan^2\alpha)\,x^2 + 2R(\tan\alpha)\,x - R^2}{2R} = \frac{-\left[(\tan\alpha)\,x - R\right]^2}{2R} \le 0 \end{aligned}$$

We have shown that every target that can be hit by the projectile lies on or inside the parabola $y = -\dfrac{1}{2R}\,x^2 + \dfrac{R}{2}$.

Now let (a, b) be any point on or inside the parabola $y = -\dfrac{1}{2R}\,x^2 + \dfrac{R}{2}$. Then $-R \le a \le R$ and $0 \le b \le -\dfrac{1}{2R}\,a^2 + \dfrac{R}{2}$.

We seek an angle α such that (a, b) lies in the path of the projectile; that is, we wish to find an angle α such that $b = -\dfrac{1}{2R\cos^2\alpha}\,a^2 + (\tan\alpha)\,a$ or equivalently $b = \dfrac{-1}{2R}\,(\tan^2\alpha + 1)a^2 + (\tan\alpha)\,a$. Rearranging this equation we get $\dfrac{a^2}{2R}\tan^2\alpha - a\tan\alpha + \left(\dfrac{a^2}{2R} + b\right) = 0$ or $a^2(\tan\alpha)^2 - 2aR(\tan\alpha) + (a^2 + 2bR) = 0$ $(\star)$. This quadratic equation for $\tan\alpha$ has real solutions exactly when the discriminant is nonnegative. Now $B^2 - 4AC \ge 0 \;\Leftrightarrow$ $(-2aR)^2 - 4a^2(a^2 + 2bR) \ge 0 \;\Leftrightarrow\; 4a^2(R^2 - a^2 - 2bR) \ge 0 \;\Leftrightarrow\; -a^2 - 2bR + R^2 \ge 0 \;\Leftrightarrow$ $b \le \dfrac{1}{2R}\,(R^2 - a^2) \;\Leftrightarrow\; b \le \dfrac{-1}{2R}\,a^2 + \dfrac{R}{2}$. This condition is satisfied since (a, b) is on or inside the parabola $y = -\dfrac{1}{2R}\,x^2 + \dfrac{R}{2}$. It follows that (a, b) lies in the path of the projectile when $\tan\alpha$ satisfies $(\star)$, that is, when

$$\tan\alpha = \frac{2aR \pm \sqrt{4a^2(R^2 - a^2 - 2bR)}}{2a^2} = \frac{R \pm \sqrt{R^2 - 2bR - a^2}}{a}.$$

(c)



If the gun is pointed at a target with height h at a distance D downrange, then $\tan\alpha = h/D$. When the projectile reaches a distance D downrange (remember we are assuming that it doesn't hit the ground first), we have $D = x = (v_0\cos\alpha)t$, so $t = \dfrac{D}{v_0\cos\alpha}$ and $y = (v_0\sin\alpha)t - \frac{1}{2}gt^2 = D\tan\alpha - \dfrac{gD^2}{2v_0^2\cos^2\alpha}$.

Meanwhile, the target, whose x-coordinate is also D, has fallen from height h to height $h - \frac{1}{2}gt^2 = D\tan\alpha - \dfrac{gD^2}{2v_0^2\cos^2\alpha}$. Thus the projectile hits the target.

4. (a) As in Problem 3, $\mathbf{r}(t) = (v_0\cos\alpha)t\,\mathbf{i} + \left[(v_0\sin\alpha)t - \frac{1}{2}gt^2\right]\mathbf{j}$, so $x = (v_0\cos\alpha)t$ and $y = (v_0\sin\alpha)t - \frac{1}{2}gt^2$. The difference here is that the projectile travels until it reaches a point where $x > 0$ and $y = -(\tan\theta)x$. (Here $0 \le \theta \le \frac{\pi}{2}$.)

From the parametric equations, we obtain $t = \dfrac{x}{v_0\cos\alpha}$ and $y = \dfrac{(v_0\sin\alpha)x}{v_0\cos\alpha} - \dfrac{gx^2}{2v_0^2\cos^2\alpha} = (\tan\alpha)x - \dfrac{gx^2}{2v_0^2\cos^2\alpha}$.

Thus the projectile hits the inclined plane at the point where $(\tan\alpha)x - \dfrac{gx^2}{2v_0^2\cos^2\alpha} = -(\tan\theta)x$. Since $\dfrac{gx^2}{2v_0^2\cos^2\alpha} = (\tan\alpha + \tan\theta)x$ and $x > 0$, we must have $\dfrac{gx}{2v_0^2\cos^2\alpha} = \tan\alpha + \tan\theta$. It follows that

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x = \dfrac{2v_0^2 \cos^2\alpha}{g}(\tan\alpha + \tan\theta)$ and $t = \dfrac{x}{v_0 \cos\alpha} = \dfrac{2v_0 \cos\alpha}{g}(\tan\alpha + \tan\theta)$. This means that the parametric equations are defined for t in the interval $\left[0, \dfrac{2v_0 \cos\alpha}{g}(\tan\alpha + \tan\theta)\right]$.

(b) The downhill range (that is, the distance to the projectile's landing point as measured along the inclined plane) is $R(\alpha) = x \sec\theta$, where x is the coordinate of the landing point calculated in part (a). Thus



$$R(\alpha) = \frac{2v_0^2 \cos^2\alpha}{g}(\tan\alpha + \tan\theta)\sec\theta = \frac{2v_0^2}{g}\left(\frac{\sin\alpha\,\cos\alpha}{\cos\theta} + \frac{\cos^2\alpha\,\sin\theta}{\cos^2\theta}\right)$$

$$= \frac{2v_0^2 \cos\alpha}{g\cos^2\theta}(\sin\alpha\,\cos\theta + \cos\alpha\,\sin\theta) = \frac{2v_0^2 \cos\alpha\,\sin(\alpha+\theta)}{g\cos^2\theta}$$

$R(\alpha)$ is maximized when

$$0 = R'(\alpha) = \frac{2v_0^2}{g\cos^2\theta}[-\sin\alpha\,\sin(\alpha+\theta) + \cos\alpha\,\cos(\alpha+\theta)]$$

$$= \frac{2v_0^2}{g\cos^2\theta}\cos[(\alpha+\theta)+\alpha] = \frac{2v_0^2\cos(2\alpha+\theta)}{g\cos^2\theta}$$

This condition implies that $\cos(2\alpha+\theta) = 0 \;\Rightarrow\; 2\alpha+\theta = \frac{\pi}{2} \;\Rightarrow\; \alpha = \frac{1}{2}\left(\frac{\pi}{2} - \theta\right)$.

(c) The solution is similar to the solutions to parts (a) and (b). This time the projectile travels until it reaches a point where $x > 0$ and $y = (\tan\theta)x$. Since $\tan\theta = -\tan(-\theta)$, we obtain the solution from the previous one by replacing θ with $-\theta$. The desired angle is $\alpha = \frac{1}{2}\left(\frac{\pi}{2} + \theta\right)$.

(d) As observed in part (c), firing the projectile up an inclined plane with angle of inclination θ involves the same equations as in parts (a) and (b) but with θ replaced by $-\theta$. So if R is the distance up an inclined plane, we know from part (b) that $R = \dfrac{2v_0^2 \cos\alpha \sin(\alpha-\theta)}{g\cos^2(-\theta)} \;\Rightarrow\; v_0^2 = \dfrac{Rg\cos^2\theta}{2\cos\alpha\sin(\alpha-\theta)}$. v_0^2 is minimized (and hence v_0 is minimized) with respect to α when

$$0 = \frac{d}{d\alpha}(v_0^2) = \frac{Rg\cos^2\theta}{2}\cdot\frac{-(\cos\alpha\cos(\alpha-\theta) - \sin\alpha\sin(\alpha-\theta))}{[\cos\alpha\sin(\alpha-\theta)]^2}$$

$$= \frac{-Rg\cos^2\theta}{2}\cdot\frac{\cos[\alpha+(\alpha-\theta)]}{[\cos\alpha\sin(\alpha-\theta)]^2} = \frac{-Rg\cos^2\theta}{2}\cdot\frac{\cos(2\alpha-\theta)}{[\cos\alpha\sin(\alpha-\theta)]^2}$$

Since $\theta < \alpha < \frac{\pi}{2}$, this implies $\cos(2\alpha-\theta) = 0 \;\Leftrightarrow\; 2\alpha-\theta = \frac{\pi}{2} \;\Rightarrow\; \alpha = \frac{1}{2}\left(\frac{\pi}{2} + \theta\right)$. Thus the initial speed, and hence the energy required, is minimized for $\alpha = \frac{1}{2}\left(\frac{\pi}{2} + \theta\right)$.

5. (a) $\mathbf{a} = -g\,\mathbf{j} \;\Rightarrow\; \mathbf{v} = \mathbf{v}_0 - gt\,\mathbf{j} = 2\,\mathbf{i} - gt\,\mathbf{j} \;\Rightarrow\; \mathbf{s} = \mathbf{s}_0 + 2t\,\mathbf{i} - \frac{1}{2}gt^2\,\mathbf{j} = 3.5\,\mathbf{j} + 2t\,\mathbf{i} - \frac{1}{2}gt^2\,\mathbf{j} \;\Rightarrow$ $\mathbf{s} = 2t\,\mathbf{i} + \left(3.5 - \frac{1}{2}gt^2\right)\mathbf{j}$. Therefore $y = 0$ when $t = \sqrt{7/g}$ seconds. At that instant, the ball is $2\sqrt{7/g} \approx 0.94$ ft to the right of the table top. Its coordinates (relative to an origin on the floor directly under the table's edge) are $(0.94, 0)$. At impact, the velocity is $\mathbf{v} = 2\,\mathbf{i} - \sqrt{7g}\,\mathbf{j}$, so the speed is $|\mathbf{v}| = \sqrt{4+7g} \approx 15$ ft/s.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) The slope of the curve when $t = \sqrt{\frac{7}{g}}$ is $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{-gt}{2} = \frac{-g\sqrt{7/g}}{2} = \frac{-\sqrt{7g}}{2}$. Thus $\cot\theta = \frac{\sqrt{7g}}{2}$ and $\theta \approx 7.6°$.

(c) From (a), $|\mathbf{v}| = \sqrt{4+7g}$. So the ball rebounds with speed $0.8\sqrt{4+7g} \approx 12.08$ ft/s at angle of inclination $90° - \theta \approx 82.3886°$. By Example 13.4.5, the horizontal distance traveled between bounces is $d = \frac{v_0^2 \sin 2\alpha}{g}$, where $v_0 \approx 12.08$ ft/s and $\alpha \approx 82.3886°$. Therefore, $d \approx 1.197$ ft. So the ball strikes the floor at about $2\sqrt{7/g} + 1.197 \approx 2.13$ ft to the right of the table's edge.

6. By the Fundamental Theorem of Calculus, $\mathbf{r}'(t) = \langle \sin(\frac{1}{2}\pi t^2), \cos(\frac{1}{2}\pi t^2)\rangle$, $|\mathbf{r}'(t)| = 1$ and so $\mathbf{T}(t) = \mathbf{r}'(t)$. Thus $\mathbf{T}'(t) = \pi t\langle \cos(\frac{1}{2}\pi t^2), -\sin(\frac{1}{2}\pi t^2)\rangle$ and the curvature is $\kappa = |\mathbf{T}'(t)| = \sqrt{(\pi t)^2(1)} = \pi|t|$.

7. The trajectory of the projectile is given by $\mathbf{r}(t) = (v\cos\alpha)t\,\mathbf{i} + \left[(v\sin\alpha)t - \frac{1}{2}gt^2\right]\mathbf{j}$, so $\mathbf{v}(t) = \mathbf{r}'(t) = v\cos\alpha\,\mathbf{i} + (v\sin\alpha - gt)\,\mathbf{j}$ and

$$|\mathbf{v}(t)| = \sqrt{(v\cos\alpha)^2 + (v\sin\alpha - gt)^2} = \sqrt{v^2 - (2vg\sin\alpha)\,t + g^2t^2} = \sqrt{g^2\left(t^2 - \frac{2v}{g}(\sin\alpha)\,t + \frac{v^2}{g^2}\right)}$$

$$= g\sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \frac{v^2}{g^2} - \frac{v^2}{g^2}\sin^2\alpha} = g\sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \frac{v^2}{g^2}\cos^2\alpha}$$

The projectile hits the ground when $(v\sin\alpha)t - \frac{1}{2}gt^2 = 0 \;\Rightarrow\; t = \frac{2v}{g}\sin\alpha$, so the distance traveled by the projectile is

$$L(\alpha) = \int_0^{(2v/g)\sin\alpha} |\mathbf{v}(t)|\,dt = \int_0^{(2v/g)\sin\alpha} g\sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \frac{v^2}{g^2}\cos^2\alpha}\,dt$$

$$= g\left[\frac{t - (v/g)\sin\alpha}{2}\sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2}\right.$$

$$\left. + \frac{[(v/g)\cos\alpha]^2}{2}\ln\left(t - \frac{v}{g}\sin\alpha + \sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2}\right)\right]_0^{(2v/g)\sin\alpha}$$

[using Formula 21 in the Table of Integrals]

$$= \frac{g}{2}\left[\frac{v}{g}\sin\alpha\sqrt{\left(\frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2} + \left(\frac{v}{g}\cos\alpha\right)^2\ln\left(\frac{v}{g}\sin\alpha + \sqrt{\left(\frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2}\right)\right.$$

$$\left. + \frac{v}{g}\sin\alpha\sqrt{\left(\frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2} - \left(\frac{v}{g}\cos\alpha\right)^2\ln\left(-\frac{v}{g}\sin\alpha + \sqrt{\left(\frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2}\right)\right]$$

$$= \frac{g}{2}\left[\frac{v}{g}\sin\alpha\cdot\frac{v}{g} + \frac{v^2}{g^2}\cos^2\alpha\ln\left(\frac{v}{g}\sin\alpha + \frac{v}{g}\right) + \frac{v}{g}\sin\alpha\cdot\frac{v}{g} - \frac{v^2}{g^2}\cos^2\alpha\ln\left(-\frac{v}{g}\sin\alpha + \frac{v}{g}\right)\right]$$

$$= \frac{v^2}{g}\sin\alpha + \frac{v^2}{2g}\cos^2\alpha\ln\left(\frac{(v/g)\sin\alpha + v/g}{-(v/g)\sin\alpha + v/g}\right) = \frac{v^2}{g}\sin\alpha + \frac{v^2}{2g}\cos^2\alpha\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)$$

We want to maximize $L(\alpha)$ for $0 \le \alpha \le \pi/2$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$L'(\alpha) = \frac{v^2}{g}\cos\alpha + \frac{v^2}{2g}\left[\cos^2\alpha \cdot \frac{1-\sin\alpha}{1+\sin\alpha} \cdot \frac{2\cos\alpha}{(1-\sin\alpha)^2} - 2\cos\alpha\,\sin\alpha\,\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)\right]$$

$$= \frac{v^2}{g}\cos\alpha + \frac{v^2}{2g}\left[\cos^2\alpha \cdot \frac{2}{\cos\alpha} - 2\cos\alpha\,\sin\alpha\,\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)\right]$$

$$= \frac{v^2}{g}\cos\alpha + \frac{v^2}{g}\cos\alpha\left[1 - \sin\alpha\,\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)\right] = \frac{v^2}{g}\cos\alpha\left[2 - \sin\alpha\,\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)\right]$$

$L(\alpha)$ has critical points for $0 < \alpha < \pi/2$ when $L'(\alpha) = 0 \;\Rightarrow\; 2 - \sin\alpha \ln\left(\dfrac{1+\sin\alpha}{1-\sin\alpha}\right) = 0$ [since $\cos\alpha \neq 0$]. Solving by graphing (or using a CAS) gives $\alpha \approx 0.9855$. Compare values at the critical point and the endpoints: $L(0) = 0$, $L(\pi/2) = v^2/g$, and $L(0.9855) \approx 1.20v^2/g$. Thus the distance traveled by the projectile is maximized for $\alpha \approx 0.9855$ or $\approx 56°$.

8. As the cable is wrapped around the spool, think of the top or bottom of the cable forming a helix of radius $R + r$. Let h be the vertical distance between coils. Then, from similar triangles,



$$\frac{2r}{\sqrt{h^2-4r^2}} = \frac{2\pi(r+R)}{h} \quad\Rightarrow\quad h^2r^2 = \pi^2(r+R)^2(h^2-4r^2) \quad\Rightarrow$$

$$h = \frac{2\pi r(r+R)}{\sqrt{\pi^2(r+R)^2 - r^2}}.$$

If we parametrize the helix by $x(t) = (R+r)\cos t$, $y(t) = (R+r)\sin t$, then we must have $z(t) = [h/(2\pi)]t$. The length of one complete cycle is

$$\ell = \int_0^{2\pi} \sqrt{[x'(t)]^2 + [y'(t)]^2 + [z'(t)]^2}\,dt = \int_0^{2\pi} \sqrt{(R+r)^2 + \left(\frac{h}{2\pi}\right)^2}\,dt = 2\pi\sqrt{(R+r)^2 + \left(\frac{h}{2\pi}\right)^2}$$

$$= 2\pi\sqrt{(R+r)^2 + \frac{r^2(R+r)^2}{\pi^2(R+r)^2 - r^2}} = 2\pi(R+r)\sqrt{1 + \frac{r^2}{\pi^2(R+r)^2 - r^2}} = \frac{2\pi^2(R+r)^2}{\sqrt{\pi^2(R+r)^2 - r^2}}$$

The number of complete cycles is $[\![L/\ell]\!]$, and so the shortest length along the spool is

$$h\left[\!\left[\frac{L}{\ell}\right]\!\right] = \frac{2\pi r(R+r)}{\sqrt{\pi^2(R+r)^2 - r^2}}\left[\!\left[\frac{L\sqrt{\pi^2(R+r)^2 - r^2}}{2\pi^2(R+r)^2}\right]\!\right]$$

9. We can write the vector equation as $\mathbf{r}(t) = \mathbf{a}t^2 + \mathbf{b}t + \mathbf{c}$ where $\mathbf{a} = \langle a_1, a_2, a_3\rangle$, $\mathbf{b} = \langle b_1, b_2, b_3\rangle$, and $\mathbf{c} = \langle c_1, c_2, c_3\rangle$. Then $\mathbf{r}'(t) = 2t\,\mathbf{a} + \mathbf{b}$ which says that each tangent vector is the sum of a scalar multiple of $\mathbf{a}$ and the vector $\mathbf{b}$. Thus the tangent vectors are all parallel to the plane determined by $\mathbf{a}$ and $\mathbf{b}$ so the curve must be parallel to this plane. [Here we assume that $\mathbf{a}$ and $\mathbf{b}$ are nonparallel. Otherwise the tangent vectors are all parallel and the curve lies along a single line.] A normal vector for the plane is $\mathbf{a} \times \mathbf{b} = \langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1\rangle$. The point (c_1, c_2, c_3) lies on the plane (when $t = 0$), so an equation of the plane is

$$(a_2b_3 - a_3b_2)(x - c_1) + (a_3b_1 - a_1b_3)(y - c_2) + (a_1b_2 - a_2b_1)(z - c_3) = 0$$

or

$$(a_2b_3 - a_3b_2)x + (a_3b_1 - a_1b_3)y + (a_1b_2 - a_2b_1)z = a_2b_3c_1 - a_3b_2c_1 + a_3b_1c_2 - a_1b_3c_2 + a_1b_2c_3 - a_2b_1c_3$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

14 □ PARTIAL DERIVATIVES

14.1 Functions of Several Variables

1. (a) From Table 1, $f(-15, 40) = -27$, which means that if the temperature is $-15°\text{C}$ and the wind speed is 40 km/h, then the air would feel equivalent to approximately $-27°\text{C}$ without wind.

   (b) The question is asking: when the temperature is $-20°\text{C}$, what wind speed gives a wind-chill index of $-30°\text{C}$? From Table 1, the speed is 20 km/h.

   (c) The question is asking: when the wind speed is 20 km/h, what temperature gives a wind-chill index of $-49°\text{C}$? From Table 1, the temperature is $-35°\text{C}$.

   (d) The function $W = f(-5, v)$ means that we fix T at -5 and allow v to vary, resulting in a function of one variable. In other words, the function gives wind-chill index values for different wind speeds when the temperature is $-5°\text{C}$. From Table 1 (look at the row corresponding to $T = -5$), the function decreases and appears to approach a constant value as v increases.

   (e) The function $W = f(T, 50)$ means that we fix v at 50 and allow T to vary, again giving a function of one variable. In other words, the function gives wind-chill index values for different temperatures when the wind speed is 50 km/h . From Table 1 (look at the column corresponding to $v = 50$), the function increases almost linearly as T increases.

2. (a) From Table 3, $f(95, 70) = 124$, which means that when the actual temperature is $95°\text{F}$ and the relative humidity is 70%, the perceived air temperature is approximately $124°\text{F}$.

   (b) Looking at the row corresponding to $T = 90$, we see that $f(90, h) = 100$ when $h = 60$.

   (c) Looking at the column corresponding to $h = 50$, we see that $f(T, 50) = 88$ when $T = 85$.

   (d) $I = f(80, h)$ means that T is fixed at 80 and h is allowed to vary, resulting in a function of h that gives the humidex values for different relative humidities when the actual temperature is $80°\text{F}$. Similarly, $I = f(100, h)$ is a function of one variable that gives the humidex values for different relative humidities when the actual temperature is $100°\text{F}$. Looking at the rows of the table corresponding to $T = 80$ and $T = 100$, we see that $f(80, h)$ increases at a relatively constant rate of approximately $1°\text{F}$ per 10% relative humidity, while $f(100, h)$ increases more quickly (at first with an average rate of change of $5°\text{F}$ per 10% relative humidity) and at an increasing rate (approximately $12°\text{F}$ per 10% relative humidity for larger values of h).

3. $P(120, 20) = 1.47(120)^{0.65}(20)^{0.35} \approx 94.2$, so when the manufacturer invests \$20 million in capital and 120,000 hours of labor are completed yearly, the monetary value of the production is about \$94.2 million.

4. If the amounts of labor and capital are both doubled, we replace L, K in the function with $2L, 2K$, giving

$$P(2L, 2K) = 1.01(2L)^{0.75}(2K)^{0.25} = 1.01(2^{0.75})(2^{0.25})L^{0.75}K^{0.25} = (2^1)1.01L^{0.75}K^{0.25} = 2P(L, K)$$

   Thus, the production is doubled. It is also true for the general case $P(L, K) = bL^{\alpha}K^{1-\alpha}$:

$$P(2L, 2K) = b(2L)^{\alpha}(2K)^{1-\alpha} = b(2^{\alpha})(2^{1-\alpha})L^{\alpha}K^{1-\alpha} = (2^{\alpha+1-\alpha})bL^{\alpha}K^{1-\alpha} = 2P(L, K).$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. (a) $f(160, 70) = 0.1091(160)^{0.425}(70)^{0.725} \approx 20.5$, which means that the surface area of a person 70 inches (5 feet 10 inches) tall who weighs 160 pounds is approximately 20.5 square feet.

(b) Answers will vary depending on the height and weight of the reader.

6. We compare the values for the wind-chill index given by Table 1 with those given by the model function:

Modeled Wind-Chill Index Values $W(T, v)$

Wind Speed (km/h)

Actual temperature (°C) T \ V	5	10	15	20	25	30	40	50	60	70	80
5	4.08	2.66	1.74	1.07	0.52	0.05	−0.71	−1.33	−1.85	2.30	−2.70
0	−1.59	3.31	4.42	5.24	−5.91	−6.47	7.40	8.14	−8.77	−9.32	−9.80
−5	−7.26	−9.29	−10.58	11.55	−12.34	−13.00	−14.08	−14.96	−15.70	−16.34	16.91
−10	−12.93	−15.26	−16.75	−17.86	−18.76	−19.52	−20.77	−21.77	−22.62	−23.36	−24.01
−15	−18.61	−21.23	−22.91	−24.17	25.19	26.04	−27.45	28.59	−29.54	−30.38	−31.11
20	−24.28	−27.21	−29.08	−30.48	−31.61	−32.57	−34.13	−35.40	−36.47	−37.40	−38.22
−25	−29.95	−33.18	−35.24	36.79	−38.04	−39.09	−40.82	−42.22	−43.39	44.42	45.32
−30	35.62	−39.15	−41.41	−43.10	44.46	−45.62	−47.50	−49.03	−50.32	−51.44	−52.43
−35	−41.30	−45.13	−47.57	−49.41	−50.89	−52.14	−54.19	−55.84	−57.24	−58.46	−59.53
−40	−46.97	51.10	−53.74	−55.72	−57.31	−58.66	60.87	−62.66	−64.17	−65.48	−66.64

The values given by the function appear to be fairly close (within 0.5) to the values in Table 1.

7. (a) According to Table 4, $f(40, 15) = 25$, which means that if a 40-knot wind has been blowing in the open sea for 15 hours, it will create waves with estimated heights of 25 feet.

(b) $h = f(30, t)$ means we fix v at 30 and allow t to vary, resulting in a function of one variable. Thus here, $h = f(30, t)$ gives the wave heights produced by 30-knot winds blowing for t hours. From the table (look at the row corresponding to $v = 30$), the function increases but at a declining rate as t increases. In fact, the function values appear to be approaching a limiting value of approximately 19, which suggests that 30-knot winds cannot produce waves higher than about 19 feet.

(c) $h = f(v, 30)$ means we fix t at 30, again giving a function of one variable. So, $h = f(v, 30)$ gives the wave heights produced by winds of speed v blowing for 30 hours. From the table (look at the column corresponding to $t = 30$), the function appears to increase at an increasing rate, with no apparent limiting value. This suggests that faster winds (lasting 30 hours) always create higher waves.

8. (a) The cost of making x small boxes, y medium boxes, and z large boxes is $C = f(x, y, z) = 8000 + 2.5x + 4y + 4.5z$ dollars.

(b) $f(3000, 5000, 4000) = 8000 + 2.5(3000) + 4(5000) + 4.5(4000) = 53{,}500$ which means that it costs \$53,500 to make 3000 small boxes, 5000 medium boxes, and 4000 large boxes.

(c) Because no partial boxes will be produced, each of x, y, and z must be a positive integer or zero.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. (a) $g(2,-1) = \cos(2+2(-1)) = \cos(0) = 1$

(b) $x + 2y$ is defined for all choices of values for x and y and the cosine function is defined for all input values, so the domain of g is $\mathbb{R}^2$.

(c) The range of the cosine function is $[-1, 1]$ and $x + 2y$ generates all possible input values for the cosine function, so the range of $\cos(x + 2y)$ is $[-1, 1]$.

10. (a) $F(3,1) = 1 + \sqrt{4 - 1^2} = 1 + \sqrt{3}$

(b) $\sqrt{4 - y^2}$ is defined only when $4 - y^2 \geq 0$, or $y^2 \leq 4 \iff -2 \leq y \leq 2$. So the domain of F is $\{(x, y) \mid -2 \leq y \leq 2\}$.



(c) We know $0 \leq \sqrt{4 - y^2} \leq 2$ so $1 \leq 1 + \sqrt{4 - y^2} \leq 3$. Thus the range of F is $[1, 3]$.

11. (a) $f(1,1,1) = \sqrt{1} + \sqrt{1} + \sqrt{1} + \ln(4 - 1^2 - 1^2 - 1^2) = 3 + \ln 1 = 3$

(b) $\sqrt{x}$, $\sqrt{y}$, $\sqrt{z}$ are defined only when $x \geq 0$, $y \geq 0$, $z \geq 0$, and $\ln(4 - x^2 - y^2 - z^2)$ is defined when $4 - x^2 - y^2 - z^2 > 0 \iff x^2 + y^2 + z^2 < 4$, thus the domain is $\{(x, y, z) \mid x^2 + y^2 + z^2 < 4,\ x \geq 0,\ y \geq 0,\ z \geq 0\}$, the portion of the interior of a sphere of radius 2, centered at the origin, that is in the first octant.

12. (a) $g(1,2,3) = 1^3 \cdot 2^2 \cdot 3\sqrt{10 - 1 - 2 - 3} = 12\sqrt{4} = 24$

(b) g is defined only when $10 - x - y - z \geq 0 \iff z \leq 10 - x - y$, so the domain is $\{(x, y, z) \mid z \leq 10 - x - y\}$, the points on or below the plane $x + y + z = 10$.

13. $\sqrt{x - 2}$ is defined only when $x - 2 \geq 0$, or $x \geq 2$, and $\sqrt{y - 1}$ is defined only when $y - 1 \geq 0$, or $y \geq 1$. So the domain of f is $\{(x, y) \mid x \geq 2,\ y \geq 1\}$.



14. $\sqrt[4]{x - 3y}$ is defined only when $x - 3y \geq 0$, or $x \geq 3y$. So the domain of f is $\{(x, y) \mid x \geq 3y\}$ or equivalently $\{(x, y) \mid y \leq \frac{1}{3}x\}$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

15. $\ln(9 - x^2 - 9y^2)$ is defined only when $9 - x^2 - 9y^2 > 0$, or $\frac{1}{9}x^2 + y^2 < 1$. So the domain of f is $\{(x, y) \mid \frac{1}{9}x^2 + y^2 < 1\}$, the interior of an ellipse.



16. $\sqrt{x^2 + y^2 - 4}$ is defined only when $x^2 + y^2 - 4 \geq 0$ $\Leftrightarrow$ $x^2 + y^2 \geq 4$. So the domain of f is $\{(x, y) \mid x^2 + y^2 \geq 4\}$, the set of points on or outside a circle of radius 2 centered at the origin.



17. g is not defined if $x + y = 0$ $\Leftrightarrow$ $y = -x$ (and is defined otherwise). Thus the domain of g is $\{(x, y) \mid y \neq -x\}$, the set of all points in $\mathbb{R}^2$ that are not on the line $y = -x$.



18. $\ln(2 - x)$ is defined only when $2 - x > 0$, or $x < 2$. In addition, g is not defined if $1 - x^2 - y^2 = 0$ $\Leftrightarrow$ $x^2 + y^2 = 1$. Thus the domain of g is $\{(x, y) \mid x < 2,\ x^2 + y^2 \neq 1\}$, the set of all points to the left of the line $x = 2$ and not on the unit circle.



19. $\sqrt{y - x^2}$ is defined only when $y - x^2 \geq 0$, or $y \geq x^2$. In addition, f is not defined if $1 - x^2 = 0$ $\Leftrightarrow$ $x = \pm 1$. Thus the domain of f is $\{(x, y) \mid y \geq x^2,\ x \neq \pm 1\}$.



20. $\sin^{-1}(x + y)$ is defined only when $-1 \leq x + y \leq 1$ $\Leftrightarrow$ $-1 - x \leq y \leq 1 - x$. Thus the domain of f is $\{(x, y) \mid -1 - x \leq y \leq 1 - x\}$, consisting of those points on or between the parallel lines $y = -1 - x$ and $y = 1 - x$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. f is defined only when $4 - x^2 \geq 0 \Leftrightarrow -2 \leq x \leq 2$ and $9 - y^2 \geq 0 \Leftrightarrow -3 \leq y \leq 3$ and $1 - z^2 \geq 0 \Leftrightarrow -1 \leq z \leq 1$. Thus the domain of f is $\{(x, y, z) \mid -2 \leq x \leq 2,\ -3 \leq y \leq 3,\ -1 \leq z \leq 1\}$, a solid rectangular box with vertices $(\pm 2, \pm 3, \pm 1)$ (all combinations).



22. f is defined only when $16 - 4x^2 - 4y^2 - z^2 > 0 \Rightarrow \dfrac{x^2}{4} + \dfrac{y^2}{4} + \dfrac{z^2}{16} < 1$. Thus, $D = \left\{(x, y, z) \,\middle|\, \dfrac{x^2}{4} + \dfrac{y^2}{4} + \dfrac{z^2}{16} < 1\right\}$, that is, the points inside the ellipsoid $\dfrac{x^2}{4} + \dfrac{y^2}{4} + \dfrac{z^2}{16} = 1$.



23. The graph of f has equation $z = y$, a plane which intersects the yz-plane in the line $z = y$, $x = 0$. The portion of this plane in the first octant is shown.



24. The graph of f has equation $z = x^2$, a parabolic cylinder.



25. $z = 10 - 4x - 5y$ or $4x + 5y + z = 10$, a plane with intercepts 2.5, 2, and 10.



26. $z = \cos y$, a cylinder.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

27. $z = \sin x$, a cylinder.



28. $z = 2 - x^2 - y^2$, a circular paraboloid opening downward with vertex at $(0, 0, 2)$.



29. $z = x^2 + 4y^2 + 1$, an elliptic paraboloid opening upward with vertex at $(0, 0, 1)$.



30. $z = \sqrt{4x^2 + y^2}$ so $4x^2 + y^2 = z^2$ and $z \geq 0$, the top half of an elliptic cone.



31. $z = \sqrt{4 - 4x^2 - y^2}$ so $4x^2 + y^2 + z^2 = 4$ or $x^2 + \dfrac{y^2}{4} + \dfrac{z^2}{4} = 1$ and $z \geq 0$, the top half of an ellipsoid.



32. (a) $f(x, y) = \dfrac{1}{1 + x^2 + y^2}$. The trace in $x = 0$ is $z = \dfrac{1}{1 + y^2}$, and the trace in $y = 0$ is $z = \dfrac{1}{1 + x^2}$. The only possibility is graph III. Notice also that the level curves of f are $\dfrac{1}{1 + x^2 + y^2} = k \quad \Leftrightarrow \quad x^2 + y^2 = \dfrac{1}{k} - 1$, a family of circles for $k < 1$.

(b) $f(x, y) = \dfrac{1}{1 + x^2 y^2}$. The trace in $x = 0$ is the horizontal line $z = 1$, and the trace in $y = 0$ is also $z = 1$. Both graphs I and II have these traces; however, notice that here $z > 0$, so the graph is I.

(c) $f(x, y) = \ln(x^2 + y^2)$. The trace in $x = 0$ is $z = \ln y^2$, and the trace in $y = 0$ is $z = \ln x^2$. The level curves of f are $\ln(x^2 + y^2) = k \quad \Leftrightarrow \quad x^2 + y^2 = e^k$, a family of circles. In addition, f is large negative when $x^2 + y^2$ is small, so this is graph IV.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $f(x, y) = \cos\sqrt{x^2 + y^2}$. The trace in $x = 0$ is $z = \cos\sqrt{y^2} = \cos|y| = \cos y$, and the trace in $y = 0$ is $z = \cos\sqrt{x^2} = \cos|x| = \cos x$. Notice also that the level curve $f(x, y) = 0$ is $\cos\sqrt{x^2 + y^2} = 0 \Leftrightarrow x^2 + y^2 = \left(\frac{\pi}{2} + n\pi\right)^2$, a family of circles, so this is graph V.

(e) $f(x, y) = |xy|$. The trace in $x = 0$ is $z = 0$, and the trace in $y = 0$ is $z = 0$, so it must be graph VI.

(f) $f(x, y) = \cos(xy)$. The trace in $x = 0$ is $z = \cos 0 = 1$, and the trace in $y = 0$ is $z = 1$. As mentioned in part (b), these traces match both graphs I and II. Here z can be negative, so the graph is II. (Also notice that the trace in $x = 1$ is $z = \cos y$, and the trace in $y = 1$ is $z = \cos x$.)

33. The point $(-3, 3)$ lies between the level curves with z-values 50 and 60. Since the point is a little closer to the level curve with $z = 60$, we estimate that $f(-3, 3) \approx 56$. The point $(3, -2)$ appears to be just about halfway between the level curves with z-values 30 and 40, so we estimate $f(3, -2) \approx 35$. The graph rises as we approach the origin, gradually from above, steeply from below.

34. (a) C (Chicago) lies between level curves with pressures 1012 and 1016 mb, and since C appears to be located about one-fourth the distance from the 1012 mb isobar to the 1016 mb isobar, we estimate the pressure at Chicago to be about 1013 mb. N lies very close to a level curve with pressure 1012 mb so we estimate the pressure at Nashville to be approximately 1012 mb. S appears to be just about halfway between level curves with pressures 1008 and 1012 mb, so we estimate the pressure at San Francisco to be about 1010 mb. V lies close to a level curve with pressure 1016 mb but we can't see a level curve to its left so it is more difficult to make an accurate estimate. There are lower pressures to the right of V and V is a short distance to the left of the level curve with pressure 1016 mb, so we might estimate that the pressure at Vancouver is about 1017 mb.

(b) Winds are stronger where the isobars are closer together (see Figure 13), and the level curves are closer near S than at the other locations, so the winds were strongest at San Francisco.

35. The point $(160, 10)$, corresponding to day 160 and a depth of 10 m, lies between the isothermals with temperature values of 8 and 12°C. Since the point appears to be located about three-fourths the distance from the 8°C isothermal to the 12°C isothermal, we estimate the temperature at that point to be approximately 11°C. The point $(180, 5)$ lies between the 16 and 20°C isothermals, very close to the 20°C level curve, so we estimate the temperature there to be about 19.5°C.

36. If we start at the origin and move along the x-axis, for example, the z-values of a cone centered at the origin increase at a constant rate, so we would expect its level curves to be equally spaced. A paraboloid with vertex the origin, on the other hand, has z-values which change slowly near the origin and more quickly as we move farther away. Thus, we would expect its level curves near the origin to be spaced more widely apart than those farther from the origin. Therefore contour map I must correspond to the paraboloid, and contour map II the cone.

37. Near A, the level curves are very close together, indicating that the terrain is quite steep. At B, the level curves are much farther apart, so we would expect the terrain to be much less steep than near A, perhaps almost flat.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

38.



39. The level curves of $B(m,h) = \dfrac{m}{h^2}$ are $\dfrac{m}{h^2} = k \iff m = kh^2$ or equivalently $h = \sqrt{m/k} = \dfrac{1}{\sqrt{k}}\sqrt{m}$ since $m > 0$, $h > 0$. We draw the level curves for $k = 18.5$, 25, 30, and 40.

The shaded region corresponds to BMI values between 18.5 and 25, those considered optimal. For a mass of 62 kg and a height of 152 cm (1.52 m), the BMI is $B(62, 1.52) = \dfrac{62}{1.52^2} \approx 26.8$, which is outside the optimal range.



40. From Exercise 39, the body mass index function is $B(m,h) = m/h^2$. The BMI for a person 200 cm (2.0 m, about 6 ft 7 in) tall and with mass 80 kg (about 176 lb) is $B(80, 2.0) = 80/(2.0)^2 = 20$. The level curve $B(m,h) = 20 \iff m = 20h^2$ is shown in the graph.

A person 1.5 m tall (about 4 ft 11 in) has a BMI on the same level curve if their mass is $m = 20(1.5)^2 = 45$ kg (about 99 lb), and a person 1.8 m (about 5 ft 11 in) tall would have mass $m = 20(1.8)^2 = 64.8$ kg (about 143 lb). (See the graph.)



41.



42.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43.



44.



45. The level curves are $x^2 - y^2 = k$. When $k = 0$ the level curve is the pair of lines $y = \pm x$, and when $k \neq 0$ the level curves are a family of hyperbolas (oriented differently for $k > 0$ than for $k < 0$).



46. The level curves are $xy = k$ or $y = k/x$. When $k \neq 0$ the level curves are a family of hyperbolas. When $k = 0$ the level curve is the pair of lines $x = 0$, $y = 0$.



47. The level curves are $\sqrt{x} + y = k$ or $y = -\sqrt{x} + k$, a family of vertical translations of the graph of the root function $y = -\sqrt{x}$.



48. The level curves are $\ln(x^2 + 4y^2) = k$ or $x^2 + 4y^2 = e^k$, a family of ellipses.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. The level curves are $ye^x = k$ or $y = ke^{-x}$, a family of exponential curves.



50. The level curves are $y - \arctan x = k$ or $y = (\arctan x) + k$, a family of vertical translations of the graph of the inverse tangent function $y = \arctan x$.



51. The level curves are $\sqrt[3]{x^2 + y^2} = k$ or $x^2 + y^2 = k^3$ $(k \geq 0)$, a family of circles centered at the origin with radius $k^{3/2}$.



52. For $k \neq 0$ and $(x, y) \neq (0, 0)$, $k = \dfrac{y}{x^2 + y^2} \iff x^2 + y^2 - \dfrac{y}{k} = 0 \iff x^2 + \left(y - \frac{1}{2k}\right)^2 = \dfrac{1}{4k^2}$, a family of circles with center $\left(0, \frac{1}{2k}\right)$ and radius $\frac{1}{2k}$ (without the origin). If $k = 0$, the level curve is the x-axis.



53. The contour map consists of the level curves $k = x^2 + 9y^2$, a family of ellipses with major axis the x-axis. (Or, if $k = 0$, the origin.)

The graph of $f(x, y)$ is the surface $z = x^2 + 9y^2$, an elliptic paraboloid.



If we visualize lifting each ellipse $k = x^2 + 9y^2$ of the contour map to the plane $z = k$, we have horizontal traces that indicate the shape of the graph of f.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54.



The contour map consists of the level curves $k = \sqrt{36 - 9x^2 - 4y^2} \Rightarrow 9x^2 + 4y^2 = 36 - k^2$, $k \geq 0$, a family of ellipses with major axis the y-axis. (Or, if $k = 6$, the origin.)



The graph of $f(x, y)$ is the surface $z = \sqrt{36 - 9x^2 - 4y^2}$, or equivalently the upper half of the ellipsoid $9x^2 + 4y^2 + z^2 = 36$. If we visualize lifting each ellipse $k = \sqrt{36 - 9x^2 - 4y^2}$ of the contour map to the plane $z = k$, we have horizontal traces that indicate the shape of the graph of f.

55. The isothermals are given by $k = 100/(1 + x^2 + 2y^2)$ or $x^2 + 2y^2 = (100 - k)/k$ $[0 < k \leq 100]$, a family of ellipses.



56. The equipotential curves are $k = \dfrac{c}{\sqrt{r^2 - x^2 - y^2}}$ or $x^2 + y^2 = r^2 - \left(\dfrac{c}{k}\right)^2$, a family of circles $(k \geq c/r)$.

Note: As $k \to \infty$, the radius of the circle approaches r.



57. $f(x, y) = xy^2 - x^3$





The traces parallel to the yz-plane (such as the left-front trace in the graph above) are parabolas; those parallel to the xz-plane (such as the right-front trace) are cubic curves. The surface is called a monkey saddle because a monkey sitting on the surface near the origin has places for both legs and tail to rest.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

58. $f(x, y) = xy^3 - yx^3$

The traces parallel to either the yz-plane or the xz-plane are cubic curves.





59. $f(x, y) = e^{-(x^2+y^2)/3}\left(\sin(x^2) + \cos(y^2)\right)$





60. $f(x, y) = \cos x \cos y$

The traces parallel to either the yz- or xz-plane are cosine curves with amplitudes that vary from 0 to 1.





61. $z = \sin(xy)$ (a) C (b) II

Reasons: This function is periodic in both x and y, and the function is the same when x is interchanged with y, so its graph is symmetric about the plane $y = x$. In addition, the function is 0 along the x- and y-axes. These conditions are satisfied only by C and II.

62. $z = e^x \cos y$ (a) A (b) IV

Reasons: This function is periodic in y but not x, a condition satisfied only by A and IV. Also, note that traces in $x = k$ are cosine curves with amplitude that increases as x increases.

63. $z = \sin(x - y)$ (a) F (b) I

Reasons: This function is periodic in both x and y but is constant along the lines $y = x + k$, a condition satisfied only by F and I.

64. $z = \sin x - \sin y$ (a) E (b) III

Reasons: This function is periodic in both x and y, but unlike the function in Exercise 63, it is not constant along lines such as $y = x + \pi$, so the contour map is III. Also notice that traces in $y = k$ are vertically shifted copies of the sine wave $z = \sin x$, so the graph must be E.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

65. $z = (1 - x^2)(1 - y^2)$ (a) B (b) VI

Reasons: This function is 0 along the lines $x = \pm 1$ and $y = \pm 1$. The only contour map in which this could occur is VI. Also note that the trace in the xz-plane is the parabola $z = 1 - x^2$ and the trace in the yz-plane is the parabola $z = 1 - y^2$, so the graph is B.

66. $z = \dfrac{x - y}{1 + x^2 + y^2}$ (a) D (b) V

Reasons: This function is not periodic, ruling out the graphs in A, C, E, and F. Also, the values of z approach 0 as we use points farther from the origin. The only graph that shows this behavior is D, which corresponds to V.

67. $k = x + 3y + 5z$ is a family of parallel planes with normal vector $\langle 1, 3, 5 \rangle$.

68. $k = x^2 + 3y^2 + 5z^2$ is a family of ellipsoids for $k > 0$ and the origin for $k = 0$.

69. Equations for the level surfaces are $k = y^2 + z^2$. For $k > 0$, we have a family of circular cylinders with axis the x-axis and radius $\sqrt{k}$. When $k = 0$ the level surface is the x-axis. (There are no level surfaces for $k < 0$.)

70. Equations for the level surfaces are $x^2 - y^2 - z^2 = k$. For $k = 0$, the equation becomes $y^2 + z^2 = x^2$ and the surface is a right circular cone with vertex the origin and axis the x-axis. For $k > 0$, we have a family of hyperboloids of two sheets with axis the x-axis, and for $k < 0$, we have a family of hyperboloids of one sheet with axis the x-axis.

71. (a) The graph of g is the graph of f shifted upward 2 units.

(b) The graph of g is the graph of f stretched vertically by a factor of 2.

(c) The graph of g is the graph of f reflected about the xy-plane.

(d) The graph of $g(x, y) = -f(x, y) + 2$ is the graph of f reflected about the xy-plane and then shifted upward 2 units.

72. (a) The graph of g is the graph of f shifted 2 units in the positive x-direction.

(b) The graph of g is the graph of f shifted 2 units in the negative y-direction.

(c) The graph of g is the graph of f shifted 3 units in the negative x-direction and 4 units in the positive y-direction.

73. $f(x, y) = 3x - x^4 - 4y^2 - 10xy$



Three-dimensional view



Front view

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

It does appear that the function has a maximum value, at the higher of the two "hilltops." From the front view graph, the maximum value appears to be approximately 15. Both hilltops could be considered local maximum points, as the values of f there are larger than at the neighboring points. There does not appear to be any local minimum point; although the valley shape between the two peaks looks like a minimum of some kind, some neighboring points have lower function values.

74. $f(x, y) = xye^{-x^2-y^2}$



Three-dimensional view



Front view

The function does have a maximum value, which it appears to achieve at two different points (the two "hilltops"). From the front view graph, we can estimate the maximum value to be approximately 0.18. These same two points can also be considered local maximum points. The two "valley bottoms" visible in the graph can be considered local minimum points, as all the neighboring points give greater values of f.

75.



$f(x, y) = \dfrac{x + y}{x^2 + y^2}$. As both x and y become large, the function values appear to approach 0, regardless of which direction is considered. As (x, y) approaches the origin, the graph exhibits asymptotic behavior. From some directions, $f(x, y) \to \infty$, while in others $f(x, y) \to -\infty$. (These are the vertical spikes visible in the graph.) If the graph is examined carefully, however, one can see that $f(x, y)$ approaches 0 along the line $y = -x$.

76.



$f(x, y) = \dfrac{xy}{x^2 + y^2}$. The graph exhibits different limiting values as x and y become large or as (x, y) approaches the origin, depending on the direction being examined. For example, although f is undefined at the origin, the function values appear to be $\frac{1}{2}$ along the line $y = x$, regardless of the distance from the origin. Along the line $y = -x$, the value is always $-\frac{1}{2}$. Along the axes, $f(x, y) = 0$ for all values of (x, y) except the origin. Other directions, heading toward the origin or away from the origin, give various limiting values between $-\frac{1}{2}$ and $\frac{1}{2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

77. $f(x, y) = e^{cx^2+y^2}$. First, if $c = 0$, the graph is the cylindrical surface $z = e^{y^2}$ (whose level curves are parallel lines). When $c > 0$, the vertical trace above the y-axis remains fixed while the sides of the surface in the x-direction "curl" upward, giving the graph a shape resembling an elliptic paraboloid. The level curves of the surface are ellipses centered at the origin.



$c = 0$

For $0 < c < 1$, the ellipses have major axis the x-axis and the eccentricity increases as $c \to 0$.





$c = 0.5$ (level curves in increments of 1)

For $c = 1$ the level curves are circles centered at the origin.





$c = 1$ (level curves in increments of 1)

When $c > 1$, the level curves are ellipses with major axis the y-axis, and the eccentricity increases as c increases.





$c = 2$ (level curves in increments of 4)

For values of $c < 0$, the sides of the surface in the x-direction curl downward and approach the xy-plane (while the vertical trace $x = 0$ remains fixed), giving a saddle-shaped appearance to the graph near the point $(0, 0, 1)$. The level curves consist of

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

a family of hyperbolas. As c decreases, the surface becomes flatter in the x-direction and the surface's approach to the curve in the trace $x = 0$ becomes steeper, as the graphs demonstrate.





$c = -0.5$ (level curves in increments of 0.25)





$c = -2$ (level curves in increments of 0.25)

78. $z = (ax^2 + by^2)e^{-x^2-y^2}$. There are only three basic shapes which can be obtained (the fourth and fifth graphs are the reflections of the first and second ones in the xy-plane). Interchanging a and b rotates the graph by 90° about the z-axis.



$a = 1, b = 1$



$a = 2, b = 1$



$a = 1, b = -1$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.



$a = -1, b = -1$



$a = -2, b = -1$

If a and b are both positive ($a \neq b$), we see that the graph has two maximum points whose height increases as a and b increase. If a and b have opposite signs, the graph has two maximum points and two minimum points, and if a and b are both negative, the graph has one maximum point and two minimum points.

79. $z = x^2 + y^2 + cxy$. When $c < -2$, the surface intersects the plane $z = k \neq 0$ in a hyperbola. (See the following graph.) It intersects the plane $x = y$ in the parabola $z = (2 + c)x^2$, and the plane $x = -y$ in the parabola $z = (2 - c)x^2$. These parabolas open in opposite directions, so the surface is a hyperbolic paraboloid.

When $c = -2$ the surface is $z = x^2 + y^2 - 2xy = (x - y)^2$. So the surface is constant along each line $x - y = k$. That is, the surface is a cylinder with axis $x - y = 0$, $z = 0$. The shape of the cylinder is determined by its intersection with the plane $x + y = 0$, where $z = 4x^2$, and hence the cylinder is parabolic with minima of 0 on the line $y = x$.



$c = -5, z = 2$



$c = -10$



$c = -2$

When $-2 < c \leq 0$, $z \geq 0$ for all x and y. If x and y have the same sign, then $x^2 + y^2 + cxy \geq x^2 + y^2 - 2xy = (x - y)^2 \geq 0$. If they have opposite signs, then $cxy \geq 0$. The intersection with the surface and the plane $z = k > 0$ is an ellipse (see graph below). The intersection with the surface and the planes $x = 0$ and $y = 0$ are parabolas $z = y^2$ and $z = x^2$ respectively, so the surface is an elliptic paraboloid.

When $c > 0$ the graphs have the same shape, but are reflected in the plane $x = 0$, because $x^2 + y^2 + cxy = (-x)^2 + y^2 + (-c)(-x)y$. That is, the value of z is the same for c at (x, y) as it is for $-c$ at $(-x, y)$.

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part



$c = -1, z = 2$



$c = 0$



$c = 10$

So the surface is an elliptic paraboloid for $0 < c < 2$, a parabolic cylinder for $c = 2$, and a hyperbolic paraboloid for $c > 2$.

80. First, we graph $f(x, y) = \sqrt{x^2 + y^2}$.



$f(x, y) = \sqrt{x^2 + y^2}$

Graphs of the other four functions follow.



$f(x, y) = e^{\sqrt{x^2 + y^2}}$



$f(x, y) = \ln\sqrt{x^2 + y^2}$



$f(x, y) = \sin\left(\sqrt{x^2 + y^2}\right)$



$f(x, y) = \dfrac{1}{\sqrt{x^2 + y^2}}$

Notice that each graph $f(x, y) = g\left(\sqrt{x^2 + y^2}\right)$ exhibits radial symmetry about the z-axis and the trace in the xz-plane for

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x \geq 0$ is the graph of $z = g(x)$, $x \geq 0$. This suggests that the graph of $f(x, y) = g\left(\sqrt{x^2 + y^2}\right)$ is obtained from the graph of g by graphing $z = g(x)$ in the xz-plane and rotating the curve about the z-axis.

81. (a) $P = bL^{\alpha}K^{1-\alpha} \Rightarrow \dfrac{P}{K} = bL^{\alpha}K^{-\alpha} \Rightarrow \dfrac{P}{K} = b\left(\dfrac{L}{K}\right)^{\alpha} \Rightarrow \ln\dfrac{P}{K} = \ln\left(b\left(\dfrac{L}{K}\right)^{\alpha}\right) \Rightarrow$

$$\ln\frac{P}{K} = \ln b + \alpha\ln\left(\frac{L}{K}\right)$$

(b) We list the values for $\ln(L/K)$ and $\ln(P/K)$ for the years 1899–1922. (Historically, these values were rounded to 2 decimal places.)

Year	$x = \ln(L/K)$	$y = \ln(P/K)$
1899	0	0
1900	−0.02	−0.06
1901	−0.04	−0.02
1902	−0.04	0
1903	−0.07	−0.05
1904	−0.13	−0.12
1905	−0.18	−0.04
1906	−0.20	−0.07
1907	−0.23	−0.15
1908	−0.41	−0.38
1909	−0.33	−0.24
1910	−0.35	−0.27

Year	$x = \ln(L/K)$	$y = \ln(P/K)$
1911	−0.38	−0.34
1912	−0.38	−0.24
1913	−0.41	−0.25
1914	−0.47	−0.37
1915	−0.53	−0.34
1916	−0.49	−0.28
1917	−0.53	−0.39
1918	−0.60	−0.50
1919	−0.68	−0.57
1920	−0.74	−0.57
1921	−1.05	−0.85
1922	−0.98	−0.59

After entering the (x, y) pairs into a calculator or CAS, the resulting least squares regression line through the points is approximately $y = 0.75136x + 0.01053$, which we round to $y = 0.75x + 0.01$.

(c) Comparing the regression line from part (b) to the equation $y = \ln b + \alpha x$ with $x = \ln(L/K)$ and $y = \ln(P/K)$, we have $\alpha = 0.75$ and $\ln b = 0.01 \Rightarrow b = e^{0.01} \approx 1.01$. Thus, the Cobb-Douglas production function is $P = bL^{\alpha}K^{1-\alpha} = 1.01L^{0.75}K^{0.25}$.

14.2 Limits and Continuity

1. In general, we can't say anything about $f(3, 1)$! $\lim\limits_{(x,y)\to(3,1)} f(x, y) = 6$ means that the values of $f(x, y)$ approach 6 as (x, y) approaches, but is not equal to, $(3, 1)$. If f is continuous, we know that $\lim\limits_{(x,y)\to(a,b)} f(x, y) = f(a, b)$, so $\lim\limits_{(x,y)\to(3,1)} f(x, y) = f(3, 1) = 6$.

2. (a) The outdoor temperature as a function of longitude, latitude, and time is continuous. Small changes in longitude, latitude, or time can produce only small changes in temperature, as the temperature doesn't jump abruptly from one value to another.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) Elevation is not necessarily continuous. If we think of a cliff with a sudden drop-off, a very small change in longitude or latitude can produce a comparatively large change in elevation, without all the intermediate values being attained. Elevation *can* jump from one value to another.

(c) The cost of a taxi ride is usually discontinuous. The cost normally increases in jumps, so small changes in distance traveled or time can produce a jump in cost. A graph of the function would show breaks in the surface.

3. We make a table of values of $f(x,y) = \dfrac{x^2y^3 + x^3y^2 - 5}{2 - xy}$ for a set of (x,y) points near the origin.

x \ y	−0.2	−0.1	−0.05	0	0.05	0.1	0.2
−0.2	−2.551	−2.525	−2.513	−2.500	−2.488	−2.475	−2.451
−0.1	−2.525	−2.513	−2.506	−2.500	−2.494	−2.488	−2.475
−0.05	2.513	−2.506	2.503	−2.500	−2.497	−2.494	−2.488
0	2.500	2.500	2.500		−2.500	−2.500	−2.500
0.05	−2.488	−2.494	−2.497	−2.500	−2.503	−2.506	−2.513
0.1	−2.475	−2.488	−2.494	−2.500	−2.506	−2.513	2.525
0.2	−2.451	−2.475	−2.488	−2.500	−2.513	−2.525	2.551

As the table shows, the values of $f(x,y)$ seem to approach -2.5 as (x,y) approaches the origin from a variety of different directions. This suggests that $\lim\limits_{(x,y)\to(0,0)} f(x,y) = -2.5$. Since f is a rational function, it is continuous on its domain. f is defined at $(0,0)$, so we can use direct substitution to establish that $\lim\limits_{(x,y)\to(0,0)} f(x,y) = \dfrac{0^2 0^3 + 0^3 0^2 - 5}{2 - 0 \cdot 0} = -\dfrac{5}{2}$, verifying our guess.

4. We make a table of values of $f(x,y) = \dfrac{2xy}{x^2 + 2y^2}$ for a set of (x,y) points near the origin.

x \ y	−0.3	−0.2	−0.1	0	0.1	0.2	0.3
−0.3	0.667	0.706	0.545	0.000	−0.545	0.706	−0.667
−0.2	0.545	0.667	0.667	0.000	−0.667	0.667	−0.545
0.1	0.316	0.444	0.667	0.000	−0.667	0.444	0.316
0	0.000	0.000	0.000		0.000	0.000	0.000
0.1	0.316	−0.444	−0.667	0.000	0.667	0.444	0.316
0.2	0.545	−0.667	−0.667	0.000	0.667	0.667	0.545
0.3	0.667	0.706	0.545	0.000	0.545	0.706	0.667

It appears from the table that the values of $f(x,y)$ are not approaching a single value as (x,y) approaches the origin. For verification, if we first approach $(0,0)$ along the x-axis, we have $f(x,0) = 0$, so $f(x,y) \to 0$. But if we approach $(0,0)$ along the line $y = x$, $f(x,x) = \dfrac{2x^2}{x^2 + 2x^2} = \dfrac{2}{3}$ $(x \neq 0)$, so $f(x,y) \to \frac{2}{3}$. Since f approaches different values along different paths to the origin, this limit does not exist.

5. $f(x,y) = x^2y^3 - 4y^2$ is a polynomial, and hence continuous, so we can find the limit by direct substitution:

$$\lim_{(x,y)\to(3,2)} f(x,y) = f(3,2) = (3)^2(2)^3 - 4(2)^2 = 56.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $f(x,y) = \dfrac{x^2y + xy^2}{x^2 - y^2}$ is a rational function and hence continuous on its domain.

$(2,-1)$ is in the domain of f, so f is continuous there and $\lim\limits_{(x,y)\to(2,-1)} f(x,y) = f(2,-1) = \dfrac{(2)^2(-1)+(2)(-1)^2}{(2)^2-(-1)^2} = -\dfrac{2}{3}$.

7. $x - y$ is a polynomial and therefore continuous. Since $\sin t$ is a continuous function, the composition $\sin(x-y)$ is also continuous. The function y is a polynomial, and hence continuous, and the product of continuous functions is continuous, so $f(x,y) = y\sin(x-y)$ is a continuous function. Then $\lim\limits_{(x,y)\to(\pi,\pi/2)} f(x,y) = f\left(\pi, \frac{\pi}{2}\right) = \frac{\pi}{2}\sin\left(\pi - \frac{\pi}{2}\right) = \frac{\pi}{2}\sin\frac{\pi}{2} = \frac{\pi}{2}$.

8. $2x - y$ is a polynomial and therefore continuous. Since $\sqrt{t}$ is continuous for $t \geq 0$, the composition $\sqrt{2x-y}$ is continuous where $2x - y \geq 0$. The function e^u is continuous everywhere, so the composition $f(x,y) = e^{\sqrt{2x-y}}$ is a continuous function for $2x - y \geq 0$. If $x = 3$ and $y = 2$ then $2x - y \geq 0$, so $\lim\limits_{(x,y)\to(3,2)} f(x,y) = f(3,2) = e^{\sqrt{2(3)-2}} = e^2$.

9. $f(x,y) = (x^4 - 4y^2)/(x^2 + 2y^2)$. First approach $(0,0)$ along the x-axis. Then $f(x,0) = x^4/x^2 = x^2$ for $x \neq 0$, so $f(x,y) \to 0$. Now approach $(0,0)$ along the y-axis. For $y \neq 0$, $f(0,y) = -4y^2/2y^2 = -2$, so $f(x,y) \to -2$. Since f has two different limits along two different lines, the limit does not exist.

10. $f(x,y) = (5y^4\cos^2 x)/(x^4 + y^4)$. First approach $(0,0)$ along the x-axis. Then $f(x,0) = 0/x^4 = 0$ for $x \neq 0$, so $f(x,y) \to 0$. Next approach $(0,0)$ along the y-axis. For $y \neq 0$, $f(0,y) = 5y^4/y^4 = 5$, so $f(x,y) \to 5$. Since f has two different limits along two different lines, the limit does not exist.

11. $f(x,y) = (y^2\sin^2 x)/(x^4 + y^4)$. On the x-axis, $f(x,0) = 0$ for $x \neq 0$, so $f(x,y) \to 0$ as $(x,y) \to (0,0)$ along the x-axis. Approaching $(0,0)$ along the line $y = x$, $f(x,x) = \dfrac{x^2\sin^2 x}{x^4 + x^4} = \dfrac{\sin^2 x}{2x^2} = \dfrac{1}{2}\left(\dfrac{\sin x}{x}\right)^2$ for $x \neq 0$ and $\lim\limits_{x\to 0}\dfrac{\sin x}{x} = 1$, so $f(x,y) \to \frac{1}{2}$. Since f has two different limits along two different lines, the limit does not exist.

12. $f(x,y) = \dfrac{xy - y}{(x-1)^2 + y^2}$. On the x-axis, $f(x,0) = 0/(x-1)^2 = 0$ for $x \neq 1$, so $f(x,y) \to 0$ as $(x,y) \to (1,0)$ along the x-axis. Approaching $(1,0)$ along the line $y = x - 1$, $f(x, x-1) = \dfrac{x(x-1)-(x-1)}{(x-1)^2 + (x-1)^2} = \dfrac{(x-1)^2}{2(x-1)^2} = \dfrac{1}{2}$ for $x \neq 1$, so $f(x,y) \to \frac{1}{2}$ along this line. Thus the limit does not exist.

13. $f(x,y) = \dfrac{xy}{\sqrt{x^2+y^2}}$. We can see that the limit along any line through $(0,0)$ is 0, as well as along other paths through $(0,0)$ such as $x = y^2$ and $y = x^2$. So we suspect that the limit exists and equals 0; we use the Squeeze Theorem to prove our assertion. Since $|y| \leq \sqrt{x^2+y^2}$, we have $\dfrac{|y|}{\sqrt{x^2+y^2}} \leq 1$ and so $0 \leq \left|\dfrac{xy}{\sqrt{x^2+y^2}}\right| \leq |x|$. Now $|x| \to 0$ as $(x,y) \to (0,0)$, so $\left|\dfrac{xy}{\sqrt{x^2+y^2}}\right| \to 0$ and hence $\lim\limits_{(x,y)\to(0,0)} f(x,y) = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

14. $f(x,y) = \dfrac{x^3 - y^3}{x^2 + xy + y^2} = \dfrac{(x-y)(x^2+xy+y^2)}{x^2+xy+y^2} = x - y$ for $(x,y) \neq (0,0)$. [Note that $x^2 + xy + y^2 = 0$ only when $(x,y) = (0,0)$.] Thus $\lim\limits_{(x,y)\to(0,0)} f(x,y) = \lim\limits_{(x,y)\to(0,0)} (x-y) = 0 - 0 = 0$.

15. Let $f(x,y) = \dfrac{xy^2 \cos y}{x^2 + y^4}$. Then $f(x,0) = 0$ for $x \neq 0$, so $f(x,y) \to 0$ as $(x,y) \to (0,0)$ along the x-axis. Approaching $(0,0)$ along the y-axis or the line $y = x$ also gives a limit of 0. But $f(y^2, y) = \dfrac{y^2 y^2 \cos y}{(y^2)^2 + y^4} = \dfrac{y^4 \cos y}{2y^4} = \dfrac{\cos y}{2}$ for $y \neq 0$, so $f(x,y) \to \frac{1}{2}\cos 0 = \frac{1}{2}$ as $(x,y) \to (0,0)$ along the parabola $x = y^2$. Thus the limit doesn't exist.

16. We can use the Squeeze Theorem to show that $\lim\limits_{(x,y)\to(0,0)} \dfrac{xy^4}{x^4 + y^4} = 0$:

$0 \le \dfrac{|x|\,y^4}{x^4+y^4} \le |x|$ since $0 \le \dfrac{y^4}{x^4+y^4} \le 1$, and $|x| \to 0$ as $(x,y) \to (0,0)$, so $\dfrac{|x|\,y^4}{x^4+y^4} \to 0 \quad \Rightarrow \quad \dfrac{xy^4}{x^4+y^4} \to 0$ as $(x,y) \to (0,0)$.

17.
$$\lim_{(x,y)\to(0,0)} \frac{x^2+y^2}{\sqrt{x^2+y^2+1}-1} = \lim_{(x,y)\to(0,0)} \frac{x^2+y^2}{\sqrt{x^2+y^2+1}-1} \cdot \frac{\sqrt{x^2+y^2+1}+1}{\sqrt{x^2+y^2+1}+1}$$
$$= \lim_{(x,y)\to(0,0)} \frac{(x^2+y^2)\left(\sqrt{x^2+y^2+1}+1\right)}{x^2+y^2} = \lim_{(x,y)\to(0,0)} \left(\sqrt{x^2+y^2+1}+1\right) = 2$$

18. $f(x,y) = xy^4/(x^2+y^8)$. On the x-axis, $f(x,0) = 0$ for $x \neq 0$, so $f(x,y) \to 0$ as $(x,y) \to (0,0)$ along the x-axis. Approaching $(0,0)$ along the curve $x = y^4$ gives $f(y^4, y) = y^8/2y^8 = \frac{1}{2}$ for $y \neq 0$, so along this path $f(x,y) \to \frac{1}{2}$ as $(x,y) \to (0,0)$. Thus the limit does not exist.

19. e^{y^2} is a composition of continuous functions and hence continuous. xz is a continuous function and $\tan t$ is continuous for $t \neq \frac{\pi}{2} + n\pi$ (n an integer), so the composition $\tan(xz)$ is continuous for $xz \neq \frac{\pi}{2} + n\pi$. Thus the product $f(x,y,z) = e^{y^2}\tan(xz)$ is a continuous function for $xz \neq \frac{\pi}{2} + n\pi$. If $x = \pi$ and $z = \frac{1}{3}$ then $xz \neq \frac{\pi}{2} + n\pi$, so $\lim\limits_{(x,y,z)\to(\pi,0,1/3)} f(x,y,z) = f(\pi, 0, 1/3) = e^{0^2}\tan(\pi \cdot 1/3) = 1 \cdot \tan(\pi/3) = \sqrt{3}$.

20. $f(x,y,z) = \dfrac{xy+yz}{x^2+y^2+z^2}$. Then $f(x,0,0) = 0/x^2 = 0$ for $x \neq 0$, so as $(x,y,z) \to (0,0,0)$ along the x-axis, $f(x,y,z) \to 0$. But $f(x,x,0) = x^2/(2x^2) = \frac{1}{2}$ for $x \neq 0$, so as $(x,y,z) \to (0,0,0)$ along the line $y = x$, $z = 0$, $f(x,y,z) \to \frac{1}{2}$. Thus the limit doesn't exist.

21. $f(x,y,z) = \dfrac{xy+yz^2+xz^2}{x^2+y^2+z^4}$. Then $f(x,0,0) = 0/x^2 = 0$ for $x \neq 0$, so as $(x,y,z) \to (0,0,0)$ along the x-axis, $f(x,y,z) \to 0$. But $f(x,x,0) = x^2/(2x^2) = \frac{1}{2}$ for $x \neq 0$, so as $(x,y,z) \to (0,0,0)$ along the line $y = x$, $z = 0$, $f(x,y,z) \to \frac{1}{2}$. Thus the limit doesn't exist.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. We can use the Squeeze Theorem to show that $\lim_{(x,y,z)\to(0,0,0)} \dfrac{x^2y^2z^2}{x^2+y^2+z^2} = 0$:

$0 \le \dfrac{x^2y^2z^2}{x^2+y^2+z^2} \le x^2y^2$ since $0 \le \dfrac{z^2}{x^2+y^2+z^2} \le 1$, and $x^2y^2 \to 0$ as $(x,y,z) \to (0,0,0)$, so $\dfrac{x^2y^2z^2}{x^2+y^2+z^2} \to 0$ as $(x,y,z) \to (0,0,0)$.

23.



From the ridges on the graph, we see that as $(x,y) \to (0,0)$ along the lines under the two ridges, $f(x,y)$ approaches different values. So the limit does not exist.

24.



From the graph, it appears that as we approach the origin along the lines $x = 0$ or $y = 0$, the function is everywhere 0, whereas if we approach the origin along a certain curve it has a constant value of about $\frac{1}{2}$. [In fact, $f(y^3, y) = y^6/(2y^6) = \frac{1}{2}$ for $y \neq 0$, so $f(x,y) \to \frac{1}{2}$ as $(x,y) \to (0,0)$ along the curve $x = y^3$.] Since the function approaches different values depending on the path of approach, the limit does not exist.

25. $h(x,y) = g(f(x,y)) = (2x+3y-6)^2 + \sqrt{2x+3y-6}$. Since f is a polynomial, it is continuous on $\mathbb{R}^2$ and g is continuous on its domain $\{t \mid t \ge 0\}$. Thus h is continuous on its domain $\{(x,y) \mid 2x+3y-6 \ge 0\} = \{(x,y) \mid y \ge -\frac{2}{3}x + 2\}$, which consists of all points on or above the line $y = -\frac{2}{3}x + 2$.

26. $h(x,y) = g(f(x,y)) = \dfrac{1 \cdot xy}{1+x^2y^2} + \ln\left(\dfrac{1-xy}{1+x^2y^2}\right)$. f is a rational function, so it is continuous on its domain. Because $1 + x^2y^2 > 0$, the domain of f is $\mathbb{R}^2$, so f is continuous everywhere. g is continuous on its domain $\{t \mid t > 0\}$. Thus h is continuous on its domain $\left\{(x,y) \;\middle|\; \dfrac{1-xy}{1+x^2y^2} > 0\right\} = \{(x,y) \mid xy < 1\}$ which consists of all points between (but not on) the two branches of the hyperbola $y = 1/x$.

27.



From the graph, it appears that f is discontinuous along the line $y = x$. If we consider $f(x,y) = e^{1/(x-y)}$ as a composition of functions, $g(x,y) = 1/(x-y)$ is a rational function and therefore continuous except where $x - y = 0 \iff y = x$. Since the function $h(t) = e^t$ is continuous everywhere, the composition $h(g(x,y)) = e^{1/(x-y)} = f(x,y)$ is continuous except along the line $y = x$, as we suspected.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

28.



We can see a circular break in the graph, corresponding approximately to the unit circle, where f is discontinuous. Since $f(x,y) = \dfrac{1}{1-x^2-y^2}$ is a rational function, it is continuous except where $1 - x^2 - y^2 = 0 \;\Leftrightarrow\; x^2 + y^2 = 1$, confirming our observation that f is discontinuous on the circle $x^2 + y^2 = 1$.

29. The functions xy and $1 + e^{x-y}$ are continuous everywhere, and $1 + e^{x-y}$ is never zero, so $F(x,y) = \dfrac{xy}{1+e^{x-y}}$ is continuous on its domain $\mathbb{R}^2$.

30. $F(x,y) = \cos\sqrt{1+x-y} = g(f(x,y))$ where $f(x,y) = \sqrt{1+x-y}$, continuous on its domain $\{(x,y) \mid 1+x-y \geq 0\} = \{(x,y) \mid y \leq x+1\}$, and $g(t) = \cos t$ is continuous everywhere. Thus F is continuous on its domain $\{(x,y) \mid y \leq x+1\}$.

31. $F(x,y) = \dfrac{1+x^2+y^2}{1-x^2-y^2}$ is a rational function and thus is continuous on its domain $\{(x,y) \mid 1-x^2-y^2 \neq 0\} = \{(x,y) \mid x^2+y^2 \neq 1\}$.

32. The functions $e^x + e^y$ and $e^{xy} - 1$ are continuous everywhere, so $H(x,y) = \dfrac{e^x+e^y}{e^{xy}-1}$ is continuous except where $e^{xy} - 1 = 0 \;\Rightarrow\; xy = 0 \;\Rightarrow\; x = 0$ or $y = 0$. Thus H is continuous on its domain $\{(x,y) \mid x \neq 0,\ y \neq 0\}$.

33. $\sqrt{x}$ is continuous on its domain $\{(x,y) \mid x \geq 0\}$ and $\sqrt{1-x^2-y^2}$ is continuous on its domain $\{(x,y) \mid 1-x^2-y^2 \geq 0\} = \{(x,y) \mid x^2+y^2 \leq 1\}$, so the sum $G(x,y) = \sqrt{x} + \sqrt{1-x^2-y^2}$ is continuous for $x \geq 0$ and $x^2 + y^2 \leq 1$, that is, $\{(x,y) \mid x^2+y^2 \leq 1,\ x \geq 0\}$. This is the right half of the unit disk.

34. $G(x,y) = \ln(1+x-y) = g(f(x,y))$ where $f(x,y) = 1+x-y$, a polynomial and hence continuous on $\mathbb{R}^2$, and $g(t) = \ln t$, continuous on its domain $\{t \mid t > 0\}$. Thus G is continuous on its domain $\{(x,y) \mid 1+x-y > 0\} = \{(x,y) \mid y < x+1\}$, the region in $\mathbb{R}^2$ below the line $y = x + 1$.

35. $f(x,y,z) = h(g(x,y,z))$ where $g(x,y,z) = x^2+y^2+z^2$, a polynomial that is continuous everywhere, and $h(t) = \arcsin t$, continuous on $[-1,1]$. Thus f is continuous on its domain $\{(x,y,z) \mid -1 \leq x^2+y^2+z^2 \leq 1\} = \{(x,y,z) \mid x^2+y^2+z^2 \leq 1\}$, so f is continuous on the unit ball.

36. $\sqrt{y-x^2}$ is continuous on its domain $\{(x,y) \mid y - x^2 \geq 0\} = \{(x,y) \mid y \geq x^2\}$ and $\ln z$ is continuous on its domain $\{z \mid z > 0\}$, so the product $f(x,y,z) = \sqrt{y-x^2}\ln z$ is continuous for $y \geq x^2$ and $z > 0$, that is, $\{(x,y,z) \mid y \geq x^2,\ z > 0\}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $f(x,y) = \begin{cases} \dfrac{x^2y^3}{2x^2+y^2} & \text{if } (x,y) \neq (0,0) \\ 1 & \text{if } (x,y) = (0,0) \end{cases}$ The first piece of f is a rational function defined everywhere except at the origin, so f is continuous on $\mathbb{R}^2$ except possibly at the origin. Since $x^2 \le 2x^2 + y^2$, we have $|x^2y^3/(2x^2+y^2)| \le |y^3|$. We know that $|y^3| \to 0$ as $(x,y) \to (0,0)$. So, by the Squeeze Theorem, $\lim\limits_{(x,y)\to(0,0)} f(x,y) = \lim\limits_{(x,y)\to(0,0)} \dfrac{x^2y^3}{2x^2+y^2} = 0$. But $f(0,0) = 1$, so f is discontinuous at $(0,0)$. Therefore, f is continuous on the set $\{(x,y) \mid (x,y) \neq (0,0)\}$.

38. $f(x,y) = \begin{cases} \dfrac{xy}{x^2+xy+y^2} & \text{if } (x,y) \neq (0,0) \\ 0 & \text{if } (x,y) = (0,0) \end{cases}$ The first piece of f is a rational function defined everywhere except at the origin, so f is continuous on $\mathbb{R}^2$ except possibly at the origin. $f(x,0) = 0/x^2 = 0$ for $x \neq 0$, so $f(x,y) \to 0$ as $(x,y) \to (0,0)$ along the x-axis. But $f(x,x) = x^2/(3x^2) = \frac{1}{3}$ for $x \neq 0$, so $f(x,y) \to \frac{1}{3}$ as $(x,y) \to (0,0)$ along the line $y = x$. Thus $\lim\limits_{(x,y)\to(0,0)} f(x,y)$ doesn't exist, so f is not continuous at $(0,0)$ and the largest set on which f is continuous is $\{(x,y) \mid (x,y) \neq (0,0)\}$.

39. $\lim\limits_{(x,y)\to(0,0)} \dfrac{x^3+y^3}{x^2+y^2} = \lim\limits_{r\to0^+} \dfrac{(r\cos\theta)^3 + (r\sin\theta)^3}{r^2} = \lim\limits_{r\to0^+} (r\cos^3\theta + r\sin^3\theta) = 0$

40. $\lim\limits_{(x,y)\to(0,0)} (x^2+y^2)\ln(x^2+y^2) = \lim\limits_{r\to0^+} r^2 \ln r^2 = \lim\limits_{r\to0^+} \dfrac{\ln r^2}{1/r^2} = \lim\limits_{r\to0^+} \dfrac{(1/r^2)(2r)}{-2/r^3}$ [using l'Hospital's Rule]

$= \lim\limits_{r\to0^+} (-r^2) = 0$

41. $\lim\limits_{(x,y)\to(0,0)} \dfrac{e^{-x^2-y^2}-1}{x^2+y^2} = \lim\limits_{r\to0^+} \dfrac{e^{-r^2}-1}{r^2} = \lim\limits_{r\to0^+} \dfrac{e^{-r^2}(-2r)}{2r}$ [using l'Hospital's Rule]

$= \lim\limits_{r\to0^+} -e^{-r^2} = -e^0 = -1$

42. $\lim\limits_{(x,y)\to(0,0)} \dfrac{\sin(x^2+y^2)}{x^2+y^2} = \lim\limits_{r\to0^+} \dfrac{\sin(r^2)}{r^2}$, which is an indeterminate form of type $0/0$. Using l'Hospital's Rule, we get

$\lim\limits_{r\to0^+} \dfrac{\sin(r^2)}{r^2} \overset{\text{H}}{=} \lim\limits_{r\to0^+} \dfrac{2r\cos(r^2)}{2r} = \lim\limits_{r\to0^+} \cos(r^2) = 1.$

Or: Use the fact that $\lim\limits_{\theta\to0} \dfrac{\sin\theta}{\theta} = 1$.



43. $f(x,y) = \begin{cases} \dfrac{\sin(xy)}{xy} & \text{if } (x,y) \neq (0,0) \\ 1 & \text{if } (x,y) = (0,0) \end{cases}$

From the graph, it appears that f is continuous everywhere. We know xy is continuous on $\mathbb{R}^2$ and $\sin t$ is continuous everywhere, so $\sin(xy)$ is continuous on $\mathbb{R}^2$ and $\dfrac{\sin(xy)}{xy}$ is continuous on $\mathbb{R}^2$



[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

except possibly where $xy = 0$. To show that f is continuous at those points, consider any point (a, b) in $\mathbb{R}^2$ where $ab = 0$. Because xy is continuous, $xy \to ab = 0$ as $(x, y) \to (a, b)$. If we let $t = xy$, then $t \to 0$ as $(x, y) \to (a, b)$ and $\lim_{(x,y)\to(a,b)} \dfrac{\sin(xy)}{xy} = \lim_{t\to 0} \dfrac{\sin(t)}{t} = 1$ by Equation 2.4.2 [ET 3.3.2]. Thus $\lim_{(x,y)\to(a,b)} f(x, y) = f(a, b)$ and f is continuous on $\mathbb{R}^2$.

44. (a) $f(x, y) = \begin{cases} 0 & \text{if } y \le 0 \quad \text{or} \quad y \ge x^4 \\ 1 & \text{if } 0 < y < x^4 \end{cases}$ Consider the path $y = mx^a$, $0 < a < 4$. [The path does not pass through $(0, 0)$ if $a \le 0$ except for the trivial case where $m = 0$.] If $mx^a \le 0$ then $f(x, mx^a) = 0$. If $mx^a > 0$ then $mx^a = |mx^a| = |m|\,|x^a|$ and $mx^a \ge x^4 \iff |m|\,|x^a| \ge x^4 \iff \dfrac{x^4}{|x^a|} \le |m| \iff |x|^{4-a} \le |m|$ whenever x^a is defined. Then $mx^a \ge x^4 \iff |x| \le |m|^{1/(4-a)}$ so $f(x, mx^a) = 0$ for $|x| \le |m|^{1/(4-a)}$ and $f(x, y) \to 0$ as $(x, y) \to (0, 0)$ along this path.

(b) If we approach $(0, 0)$ along the path $y = x^5$, $x > 0$ then we have $f(x, x^5) = 1$ for $0 < x < 1$ because $0 < x^5 < x^4$ there. Thus $f(x, y) \to 1$ as $(x, y) \to (0, 0)$ along this path, but in part (a) we found a limit of 0 along other paths, so $\lim_{(x,y)\to(0,0)} f(x, y)$ doesn't exist and f is discontinuous at $(0, 0)$.

(c) First we show that f is discontinuous at any point $(a, 0)$ on the x-axis. If we approach $(a, 0)$ along the path $x = a$, $y > 0$ then $f(a, y) = 1$ for $0 < y < a^4$, so $f(x, y) \to 1$ as $(x, y) \to (a, 0)$ along this path. If we approach $(a, 0)$ along the path $x = a$, $y < 0$ then $f(a, y) = 0$ since $y < 0$ and $f(x, y) \to 0$ as $(x, y) \to (a, 0)$. Thus the limit does not exist and f is discontinuous on the line $y = 0$. f is also discontinuous on the curve $y = x^4$: For any point (a, a^4) on this curve, approaching the point along the path $x = a$, $y > a^4$ gives $f(a, y) = 0$ since $y > a^4$, so $f(x, y) \to 0$ as $(x, y) \to (a, a^4)$. But approaching the point along the path $x = a$, $y < a^4$ gives $f(a, y) = 1$ for $y > 0$, so $f(x, y) \to 1$ as $(x, y) \to (a, a^4)$ and the limit does not exist there.

45. Since $|\mathbf{x} - \mathbf{a}|^2 = |\mathbf{x}|^2 + |\mathbf{a}|^2 - 2\,|\mathbf{x}|\,|\mathbf{a}| \cos\theta \ge |\mathbf{x}|^2 + |\mathbf{a}|^2 - 2\,|\mathbf{x}|\,|\mathbf{a}| = (|\mathbf{x}| - |\mathbf{a}|)^2$, we have $\big||\mathbf{x}| - |\mathbf{a}|\big| \le |\mathbf{x} - \mathbf{a}|$. Let $\epsilon > 0$ be given and set $\delta = \epsilon$. Then if $0 < |\mathbf{x} - \mathbf{a}| < \delta$, $\big||\mathbf{x}| - |\mathbf{a}|\big| \le |\mathbf{x} - \mathbf{a}| < \delta = \epsilon$. Hence $\lim_{\mathbf{x}\to\mathbf{a}} |\mathbf{x}| = |\mathbf{a}|$ and $f(\mathbf{x}) = |\mathbf{x}|$ is continuous on $\mathbb{R}^n$.

46. Let $\epsilon > 0$ be given. We need to find $\delta > 0$ such that if $0 < |\mathbf{x} - \mathbf{a}| < \delta$ then $|f(\mathbf{x}) - f(\mathbf{a})| = |\mathbf{c} \cdot \mathbf{x} - \mathbf{c} \cdot \mathbf{a}| < \epsilon$. But $|\mathbf{c} \cdot \mathbf{x} - \mathbf{c} \cdot \mathbf{a}| = |\mathbf{c} \cdot (\mathbf{x} - \mathbf{a})|$ and $|\mathbf{c} \cdot (\mathbf{x} - \mathbf{a})| \le |\mathbf{c}|\,|\mathbf{x} - \mathbf{a}|$ by Exercise 12.3.61 (the Cauchy-Schwartz Inequality). Set $\delta = \epsilon / |\mathbf{c}|$. Then if $0 < |\mathbf{x} - \mathbf{a}| < \delta$, $|f(\mathbf{x}) - f(\mathbf{a})| = |\mathbf{c} \cdot \mathbf{x} - \mathbf{c} \cdot \mathbf{a}| \le |\mathbf{c}|\,|\mathbf{x} - \mathbf{a}| < |\mathbf{c}|\,\delta = |\mathbf{c}|\,(\epsilon / |\mathbf{c}|) = \epsilon$. So f is continuous on $\mathbb{R}^n$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.3 Partial Derivatives

1. (a) $\partial T/\partial x$ represents the rate of change of T when we fix y and t and consider T as a function of the single variable x, which describes how quickly the temperature changes when longitude changes but latitude and time are constant. $\partial T/\partial y$ represents the rate of change of T when we fix x and t and consider T as a function of y, which describes how quickly the temperature changes when latitude changes but longitude and time are constant. $\partial T/\partial t$ represents the rate of change of T when we fix x and y and consider T as a function of t, which describes how quickly the temperature changes over time for a constant longitude and latitude.

   (b) $f_x(158, 21, 9)$ represents the rate of change of temperature at longitude 158°W, latitude 21°N at 9:00 AM when only longitude varies. Since the air is warmer to the west than to the east, increasing longitude results in an increased air temperature, so we would expect $f_x(158, 21, 9)$ to be positive. $f_y(158, 21, 9)$ represents the rate of change of temperature at the same time and location when only latitude varies. Since the air is warmer to the south and cooler to the north, increasing latitude results in a decreased air temperature, so we would expect $f_y(158, 21, 9)$ to be negative. $f_t(158, 21, 9)$ represents the rate of change of temperature at the same time and location when only time varies. Since typically air temperature increases from the morning to the afternoon as the sun warms it, we would expect $f_t(158, 21, 9)$ to be positive.

2. By Definition 4, $f_T(92, 60) = \lim_{h\to 0} \dfrac{f(92+h, 60) - f(92, 60)}{h}$, which we can approximate by considering $h = 2$ and $h = -2$ and using the values given in Table 1: $f_T(92, 60) \approx \dfrac{f(94, 60) - f(92, 60)}{2} = \dfrac{111 - 105}{2} = 3$, $f_T(92, 60) \approx \dfrac{f(90, 60) - f(92, 60)}{-2} = \dfrac{100 - 105}{-2} = 2.5$. Averaging these values, we estimate $f_T(92, 60)$ to be approximately 2.75. Thus, when the actual temperature is 92°F and the relative humidity is 60%, the apparent temperature rises by about 2.75°F for every degree that the actual temperature rises.

   Similarly, $f_H(92, 60) = \lim_{h\to 0} \dfrac{f(92, 60+h) - f(92, 60)}{h}$ which we can approximate by considering $h = 5$ and $h = -5$: $f_H(92, 60) \approx \dfrac{f(92, 65) - f(92, 60)}{5} = \dfrac{108 - 105}{5} = 0.6$, $f_H(92, 60) \approx \dfrac{f(92, 55) - f(92, 60)}{-5} = \dfrac{103 - 105}{-5} = 0.4$. Averaging these values, we estimate $f_H(92, 60)$ to be approximately 0.5. Thus, when the actual temperature is 92°F and the relative humidity is 60%, the apparent temperature rises by about 0.5°F for every percent that the relative humidity increases.

3. (a) By Definition 4, $f_T(-15, 30) = \lim_{h\to 0} \dfrac{f(-15+h, 30) - f(-15, 30)}{h}$, which we can approximate by considering $h = 5$ and $h = -5$ and using the values given in the table:

   $f_T(-15, 30) \approx \dfrac{f(-10, 30) - f(-15, 30)}{5} = \dfrac{-20 - (-26)}{5} = \dfrac{6}{5} = 1.2$,

   $f_T(-15, 30) \approx \dfrac{f(-20, 30) - f(-15, 30)}{-5} = \dfrac{-33 - (-26)}{-5} = \dfrac{-7}{-5} = 1.4$. Averaging these values, we estimate

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$f_t(-15, 30)$ to be approximately 1.3. Thus, when the actual temperature is $-15°C$ and the wind speed is 30 km/h, the apparent temperature rises by about $1.3°C$ for every degree that the actual temperature rises.

Similarly, $f_v(-15, 30) = \lim_{h \to 0} \dfrac{f(-15, 30 + h) - f(-15, 30)}{h}$ which we can approximate by considering $h = 10$ and $h = -10$: $f_v(-15, 30) \approx \dfrac{f(-15, 40) - f(-15, 30)}{10} = \dfrac{-27 - (-26)}{10} = \dfrac{-1}{10} = -0.1$, $f_v(-15, 30) \approx \dfrac{f(-15, 20) - f(-15, 30)}{-10} = \dfrac{-24 - (-26)}{-10} = \dfrac{2}{-10} = -0.2$. Averaging these values, we estimate $f_v(-15, 30)$ to be approximately -0.15. Thus, when the actual temperature is $-15°C$ and the wind speed is 30 km/h, the apparent temperature decreases by about $0.15°C$ for every km/h that the wind speed increases.

(b) For a fixed wind speed v, the values of the wind-chill index W increase as temperature T increases (look at a column of the table), so $\dfrac{\partial W}{\partial T}$ is positive. For a fixed temperature T, the values of W decrease (or remain constant) as v increases (look at a row of the table), so $\dfrac{\partial W}{\partial v}$ is negative (or perhaps 0).

(c) For fixed values of T, the function values $f(T, v)$ appear to become constant (or nearly constant) as v increases, so the corresponding rate of change is 0 or near 0 as v increases. This suggests that $\lim_{v \to \infty} (\partial W/\partial v) = 0$.

4. (a) $\partial h/\partial v$ represents the rate of change of h when we fix t and consider h as a function of v, which describes how quickly the wave heights change when the wind speed changes for a fixed time duration. $\partial h/\partial t$ represents the rate of change of h when we fix v and consider h as a function of t, which describes how quickly the wave heights change when the duration of time changes, but the wind speed is constant.

(b) By Definition 4, $f_v(40, 15) = \lim_{h \to 0} \dfrac{f(40 + h, 15) - f(40, 15)}{h}$ which we can approximate by considering $h = 10$ and $h = -10$ and using the values given in the table: $f_v(40, 15) \approx \dfrac{f(50, 15) - f(40, 15)}{10} = \dfrac{36 - 25}{10} = 1.1$, $f_v(40, 15) \approx \dfrac{f(30, 15) - f(40, 15)}{-10} = \dfrac{16 - 25}{-10} = 0.9$. Averaging these values, we have $f_v(40, 15) \approx 1.0$. Thus, when a 40-knot wind has been blowing for 15 hours, the wave heights should increase by about 1 foot for every knot that the wind speed increases (with the same time duration). Similarly, $f_t(40, 15) = \lim_{h \to 0} \dfrac{f(40, 15 + h) - f(40, 15)}{h}$ which we can approximate by considering $h = 5$ and $h = -5$: $f_t(40, 15) \approx \dfrac{f(40, 20) - f(40, 15)}{5} = \dfrac{28 - 25}{5} = 0.6$, $f_t(40, 15) \approx \dfrac{f(40, 10) - f(40, 15)}{-5} = \dfrac{21 - 25}{-5} = 0.8$. Averaging these values, we have $f_t(40, 15) \approx 0.7$. Thus, when a 40-knot wind has been blowing for 15 hours, the wave heights increase by about 0.7 feet for every additional hour that the wind blows.

(c) For fixed values of v, the function values $f(v, t)$ appear to increase in smaller and smaller increments, becoming nearly constant as t increases. Thus, the corresponding rate of change is nearly 0 as t increases, suggesting that $\lim_{t \to \infty} (\partial h/\partial t) = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. (a) If we start at $(1, 2)$ and move in the positive x-direction, the graph of f increases. Thus $f_x(1, 2)$ is positive.

(b) If we start at $(1, 2)$ and move in the positive y-direction, the graph of f decreases. Thus $f_y(1, 2)$ is negative.

6. (a) The graph of f decreases if we start at $(-1, 2)$ and move in the positive x-direction, so $f_x(-1, 2)$ is negative.

(b) The graph of f decreases if we start at $(-1, 2)$ and move in the positive y-direction, so $f_y(-1, 2)$ is negative.

7. (a) $f_{xx} = \frac{\partial}{\partial x}(f_x)$, so f_{xx} is the rate of change of f_x in the x-direction. f_x is negative at $(-1, 2)$ and if we move in the positive x-direction, the surface becomes less steep. Thus the values of f_x are increasing and $f_{xx}(-1, 2)$ is positive.

(b) f_{yy} is the rate of change of f_y in the y-direction. f_y is negative at $(-1, 2)$ and if we move in the positive y-direction, the surface becomes steeper. Thus the values of f_y are decreasing, and $f_{yy}(-1, 2)$ is negative.

8. (a) $f_{xy} = \frac{\partial}{\partial y}(f_x)$, so f_{xy} is the rate of change of f_x in the y-direction. f_x is positive at $(1, 2)$ and if we move in the positive y-direction, the surface becomes steeper, looking in the positive x-direction. Thus the values of f_x are increasing and $f_{xy}(1, 2)$ is positive.

(b) f_x is negative at $(-1, 2)$ and if we move in the positive y-direction, the surface gets steeper (with negative slope), looking in the positive x-direction. This means that the values of f_x are decreasing as y increases, so $f_{xy}(-1, 2)$ is negative.

9. First of all, if we start at the point $(3, -3)$ and move in the positive y-direction, we see that both b and c decrease, while a increases. Both b and c have a low point at about $(3, -1.5)$, while a is 0 at this point. So a is definitely the graph of f_y, and one of b and c is the graph of f. To see which is which, we start at the point $(-3, -1.5)$ and move in the positive x-direction. b traces out a line with negative slope, while c traces out a parabola opening downward. This tells us that b is the x-derivative of c. So c is the graph of f, b is the graph of f_x, and a is the graph of f_y.

10. $f_x(2, 1)$ is the rate of change of f at $(2, 1)$ in the x-direction. If we start at $(2, 1)$, where $f(2, 1) = 10$, and move in the positive x-direction, we reach the next contour line [where $f(x, y) = 12$] after approximately 0.6 units. This represents an average rate of change of about $\frac{2}{0.6}$. If we approach the point $(2, 1)$ from the left (moving in the positive x-direction) the output values increase from 8 to 10 with an increase in x of approximately 0.9 units, corresponding to an average rate of change of $\frac{2}{0.9}$. A good estimate for $f_x(2, 1)$ would be the average of these two, so $f_x(2, 1) \approx 2.8$. Similarly, $f_y(2, 1)$ is the rate of change of f at $(2, 1)$ in the y-direction. If we approach $(2, 1)$ from below, the output values decrease from 12 to 10 with a change in y of approximately 1 unit, corresponding to an average rate of change of -2. If we start at $(2, 1)$ and move in the positive y-direction, the output values decrease from 10 to 8 after approximately 0.9 units, a rate of change of $\frac{-2}{0.9}$. Averaging these two results, we estimate $f_y(2, 1) \approx -2.1$.

11. $f(x, y) = 16 - 4x^2 - y^2 \Rightarrow f_x(x, y) = -8x$ and $f_y(x, y) = -2y \Rightarrow f_x(1, 2) = -8$ and $f_y(1, 2) = -4$. The graph of f is the paraboloid $z = 16 - 4x^2 - y^2$ and the vertical plane $y = 2$ intersects it in the parabola $z = 12 - 4x^2$, $y = 2$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(the curve C_1 in the first figure). The slope of the tangent line to this parabola at $(1, 2, 8)$ is $f_x(1, 2) = -8$. Similarly the plane $x = 1$ intersects the paraboloid in the parabola $z = 12 - y^2$, $x = 1$ (the curve C_2 in the second figure) and the slope of the tangent line at $(1, 2, 8)$ is $f_y(1, 2) = -4$.



12. $f(x, y) = (4 - x^2 - 4y^2)^{1/2} \Rightarrow f_x(x, y) = -x(4 - x^2 - 4y^2)^{-1/2}$ and $f_y(x, y) = -4y(4 - x^2 - 4y^2)^{-1/2} \Rightarrow$ $f_x(1, 0) = -\frac{1}{\sqrt{3}}$, $f_y(1, 0) = 0$. The graph of f is the upper half of the ellipsoid $z^2 + x^2 + 4y^2 = 4$ and the plane $y = 0$ intersects the graph in the semicircle $x^2 + z^2 = 4$, $z \geq 0$ and the slope of the tangent line T_1 to this semicircle at $\left(1, 0, \sqrt{3}\right)$ is $f_x(1, 0) = -\frac{1}{\sqrt{3}}$. Similarly the plane $x = 1$ intersects the graph in the semi-ellipse $z^2 + 4y^2 = 3$, $z \geq 0$ and the slope of the tangent line T_2 to this semi-ellipse at $\left(1, 0, \sqrt{3}\right)$ is $f_y(1, 0) = 0$.



13. $f(x, y) = x^2y^3 \Rightarrow f_x = 2xy^3, \quad f_y = 3x^2y^2$





Note that traces of f in planes parallel to the xz-plane are parabolas which open downward for $y < 0$ and upward for $y > 0$, and the traces of f_x in these planes are straight lines, which have negative slopes for $y < 0$ and positive slopes for $y > 0$. The traces of f in planes parallel to the yz-plane are cubic curves, and the traces of f_y in these planes are parabolas.

14. $f(x, y) = \dfrac{y}{1 + x^2y^2} \Rightarrow f_x = \dfrac{(1 + x^2y^2)(0) - y(2xy^2)}{(1 + x^2y^2)^2} = -\dfrac{2xy^3}{(1 + x^2y^2)^2}$,

$f_y = \dfrac{(1 + x^2y^2)(1) - y(2x^2y)}{(1 + x^2y^2)^2} = \dfrac{1 - x^2y^2}{(1 + x^2y^2)^2}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.



Note that traces of f in planes parallel to the xz-plane have only one extreme value (a minimum for $y < 0$, a maximum for $y > 0$), and the traces of f_x in these planes have only one zero (going from negative to positive if $y < 0$ and from positive to negative if $y > 0$). The traces of f in planes parallel to the yz-plane have two extreme values, and the traces of f_y in these planes have two zeros.

15. $f(x,y) = x^4 + 5xy^3 \;\Rightarrow\; f_x(x,y) = 4x^3 + 5y^3,\; f_y(x,y) = 0 + 5x \cdot 3y^2 = 15xy^2$

16. $f(x,y) = x^2y - 3y^4 \;\Rightarrow\; f_x(x,y) = 2x \cdot y - 0 = 2xy,\; f_y(x,y) = x^2 \cdot 1 - 3 \cdot 4y^3 = x^2 - 12y^3$

17. $f(x,t) = t^2e^{-x} \;\Rightarrow\; f_x(x,t) = t^2 \cdot e^{-x}(-1) = -t^2e^{-x},\; f_t(x,t) = 2te^{-x}$

18. $f(x,t) = \sqrt{3x+4t} \;\Rightarrow\; f_x(x,t) = \frac{1}{2}(3x+4t)^{-1/2}(3) = \dfrac{3}{2\sqrt{3x+4t}},\; f_t(x,t) = \frac{1}{2}(3x+4t)^{-1/2}(4) = \dfrac{2}{\sqrt{3x+4t}}$

19. $z = \ln(x+t^2) \;\Rightarrow\; \dfrac{\partial z}{\partial x} = \dfrac{1}{x+t^2},\; \dfrac{\partial z}{\partial t} = \dfrac{2t}{x+t^2}$

20. $z = x\sin(xy) \;\Rightarrow\; \dfrac{\partial z}{\partial x} = x \cdot [\cos(xy)](y) + [\sin(xy)] \cdot 1 = xy\cos(xy) + \sin(xy),\; \dfrac{\partial z}{\partial y} = x\,[\cos(xy)](x) = x^2\cos(xy)$

21. $f(x,y) = x/y = xy^{-1} \;\Rightarrow\; f_x(x,y) = y^{-1} = 1/y,\; f_y(x,y) = -xy^{-2} = -x/y^2$

22. $f(x,y) = \dfrac{x}{(x+y)^2} \;\Rightarrow\; f_x(x,y) = \dfrac{(x+y)^2(1) - (x)(2)(x+y)}{[(x+y)^2]^2} = \dfrac{x+y-2x}{(x+y)^3} = \dfrac{y-x}{(x+y)^3}$,

$f_y(x,y) = \dfrac{(x+y)^2(0) - (x)(2)(x+y)}{[(x+y)^2]^2} = -\dfrac{2x}{(x+y)^3}$

23. $f(x,y) = \dfrac{ax+by}{cx+dy} \;\Rightarrow\; f_x(x,y) = \dfrac{(cx+dy)(a) - (ax+by)(c)}{(cx+dy)^2} = \dfrac{(ad-bc)y}{(cx+dy)^2}$.

$f_y(x,y) = \dfrac{(cx+dy)(b) - (ax+by)(d)}{(cx+dy)^2} = \dfrac{(bc-ad)x}{(cx+dy)^2}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

24. $w = \dfrac{e^v}{u+v^2} \quad \Rightarrow \quad \dfrac{\partial w}{\partial u} = \dfrac{0(u+v^2)-e^v(1)}{(u+v^2)^2} = -\dfrac{e^v}{(u+v^2)^2}, \quad \dfrac{\partial w}{\partial v} = \dfrac{e^v(u+v^2)-e^v(2v)}{(u+v^2)^2} = \dfrac{e^v(u+v^2-2v)}{(u+v^2)^2}$

25. $g(u,v) = (u^2v - v^3)^5 \quad \Rightarrow \quad g_u(u,v) = 5(u^2v-v^3)^4 \cdot 2uv = 10uv(u^2v-v^3)^4,$

$g_v(u,v) = 5(u^2v-v^3)^4(u^2-3v^2) = 5(u^2-3v^2)(u^2v-v^3)^4$

26. $u(r,\theta) = \sin(r\cos\theta) \quad \Rightarrow \quad u_r(r,\theta) = \cos(r\cos\theta)\cdot\cos\theta = \cos\theta\cos(r\cos\theta),$

$u_\theta(r,\theta) = \cos(r\cos\theta)(-r\sin\theta) = -r\sin\theta\cos(r\cos\theta)$

27. $R(p,q) = \tan^{-1}(pq^2) \quad \Rightarrow \quad R_p(p,q) = \dfrac{1}{1+(pq^2)^2}\cdot q^2 = \dfrac{q^2}{1+p^2q^4}, \quad R_q(p,q) = \dfrac{1}{1+(pq^2)^2}\cdot 2pq = \dfrac{2pq}{1+p^2q^4}$

28. $f(x,y) = x^y \quad \Rightarrow \quad f_x(x,y) = yx^{y-1}, \; f_y(x,y) = x^y \ln x$

29. $F(x,y) = \displaystyle\int_y^x \cos(e^t)\,dt \quad \Rightarrow \quad F_x(x,y) = \frac{\partial}{\partial x}\int_y^x \cos(e^t)\,dt = \cos(e^x)$ by the Fundamental Theorem of Calculus, Part 1;

$F_y(x,y) = \dfrac{\partial}{\partial y}\displaystyle\int_y^x \cos(e^t)\,dt = \frac{\partial}{\partial y}\left[-\int_x^y \cos(e^t)\,dt\right] = -\frac{\partial}{\partial y}\int_x^y \cos(e^t)\,dt = -\cos(e^y).$

30. $F(\alpha,\beta) = \displaystyle\int_\alpha^\beta \sqrt{t^3+1}\,dt \quad \Rightarrow$

$F_\alpha(\alpha,\beta) = \dfrac{\partial}{\partial\alpha}\displaystyle\int_\alpha^\beta \sqrt{t^3+1}\,dt = \frac{\partial}{\partial\alpha}\left[-\int_\beta^\alpha \sqrt{t^3+1}\,dt\right] = -\frac{\partial}{\partial\alpha}\int_\beta^\alpha \sqrt{t^3+1}\,dt = -\sqrt{\alpha^3+1}$ by the Fundamental

Theorem of Calculus, Part 1; $F_\beta(\alpha,\beta) = \dfrac{\partial}{\partial\beta}\displaystyle\int_\alpha^\beta \sqrt{t^3+1}\,dt = \sqrt{\beta^3+1}.$

31. $f(x,y,z) = x^3yz^2 + 2yz \quad \Rightarrow \quad f_x(x,y,z) = 3x^2yz^2, \; f_y(x,y,z) = x^3z^2 + 2z, \; f_z(x,y,z) = 2x^3yz + 2y$

32. $f(x,y,z) = xy^2e^{-xz} \quad \Rightarrow \quad f_x(x,y,z) = y^2\left[x\cdot e^{-xz}(-z) + e^{-xz}\cdot 1\right] = (1-xz)y^2e^{-xz}, \; f_y(x,y,z) = 2xye^{-xz},$

$f_z(x,y,z) = xy^2e^{-xz}(-x) = -x^2y^2e^{-xz}$

33. $w = \ln(x+2y+3z) \quad \Rightarrow \quad \dfrac{\partial w}{\partial x} = \dfrac{1}{x+2y+3z}, \; \dfrac{\partial w}{\partial y} = \dfrac{2}{x+2y+3z}, \; \dfrac{\partial w}{\partial z} = \dfrac{3}{x+2y+3z}$

34. $w = y\tan(x+2z) \quad \Rightarrow \quad \dfrac{\partial w}{\partial x} = y\left[\sec^2(x+2z)\right](1) = y\sec^2(x+2z), \; \dfrac{\partial w}{\partial y} = \tan(x+2z),$

$\dfrac{\partial w}{\partial z} = y\left[\sec^2(x+2z)\right](2) = 2y\sec^2(x+2z)$

35. $p = \sqrt{t^4+u^2\cos v} \quad \Rightarrow \quad \dfrac{\partial p}{\partial t} = \frac{1}{2}(t^4+u^2\cos v)^{-1/2}(4t^3) = \dfrac{2t^3}{\sqrt{t^4+u^2\cos v}},$

$\dfrac{\partial p}{\partial u} = \frac{1}{2}(t^4+u^2\cos v)^{-1/2}(2u\cos v) = \dfrac{u\cos v}{\sqrt{t^4+u^2\cos v}}, \; \dfrac{\partial p}{\partial v} = \frac{1}{2}(t^4+u^2\cos v)^{-1/2}[u^2(-\sin v)] = -\dfrac{u^2\sin v}{2\sqrt{t^4+u^2\cos v}}$

36. $u = x^{y/z} \quad \Rightarrow \quad u_x = \dfrac{y}{z}\,x^{(y/z)-1}, \; u_y = x^{y/z}\ln x\cdot\dfrac{1}{z} = \dfrac{x^{y/z}}{z}\ln x, \; u_z = x^{y/z}\ln x\cdot\dfrac{-y}{z^2} = -\dfrac{yx^{y/z}}{z^2}\ln x$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $h(x,y,z,t) = x^2 y\cos(z/t) \;\Rightarrow\; h_x(x,y,z,t) = 2xy\cos(z/t),\ h_y(x,y,z,t) = x^2\cos(z/t),$

$h_z(x,y,z,t) = -x^2 y\sin(z/t)(1/t) = (-x^2y/t)\sin(z/t),\ h_t(x,y,z,t) = -x^2 y\sin(z/t)(-zt^{-2}) = (x^2yz/t^2)\sin(z/t)$

38. $\phi(x,y,z,t) = \dfrac{\alpha x+\beta y^2}{\gamma z+\delta t^2} \;\Rightarrow\; \phi_x(x,y,z,t) = \dfrac{1}{\gamma z+\delta t^2}(\alpha) = \dfrac{\alpha}{\gamma z+\delta t^2},$

$\phi_y(x,y,z,t) = \dfrac{1}{\gamma z+\delta t^2}(2\beta y) = \dfrac{2\beta y}{\gamma z+\delta t^2},\ \phi_z(x,y,z,t) = \dfrac{(\gamma z+\delta t^2)(0)-(\alpha x+\beta y^2)(\gamma)}{(\gamma z+\delta t^2)^2} = \dfrac{-\gamma(\alpha x+\beta y^2)}{(\gamma z+\delta t^2)^2},$

$\phi_t(x,y,z,t) = \dfrac{(\gamma z+\delta t^2)(0)-(\alpha x+\beta y^2)(2\delta t)}{(\gamma z+\delta t^2)^2} = -\dfrac{2\delta t(\alpha x+\beta y^2)}{(\gamma z+\delta t^2)^2}$

39. $u = \sqrt{x_1^2+x_2^2+\cdots+x_n^2}$. For each $i = 1,\ldots,n$, $u_{x_i} = \frac{1}{2}\left(x_1^2+x_2^2+\cdots+x_n^2\right)^{-1/2}(2x_i) = \dfrac{x_i}{\sqrt{x_1^2+x_2^2+\cdots+x_n^2}}$.

40. $u = \sin(x_1+2x_2+\cdots+nx_n)$. For each $i = 1,\ldots,n$, $u_{x_i} = i\cos(x_1+2x_2+\cdots+nx_n)$.

41. $R(s,t) = te^{s/t} \;\Rightarrow\; R_t(s,t) = t\cdot e^{s/t}(-s/t^2)+e^{s/t}\cdot 1 = \left(1-\dfrac{s}{t}\right)e^{s/t}$, so $R_t(0,1) = \left(1-\dfrac{0}{1}\right)e^{0/1} = 1$.

42. $f(x,y) = y\sin^{-1}(xy) \;\Rightarrow\; f_y(x,y) = y\cdot\dfrac{1}{\sqrt{1-(xy)^2}}(x)+\sin^{-1}(xy)\cdot 1 = \dfrac{xy}{\sqrt{1-x^2y^2}}+\sin^{-1}(xy),$

so $f_y\left(1,\frac{1}{2}\right) = \dfrac{1\cdot\frac{1}{2}}{\sqrt{1-1^2\left(\frac{1}{2}\right)^2}}+\sin^{-1}\left(1\cdot\frac{1}{2}\right) = \dfrac{\frac{1}{2}}{\sqrt{\frac{3}{4}}}+\sin^{-1}\frac{1}{2} = \frac{1}{\sqrt{3}}+\frac{\pi}{6}$.

43. $f(x,y,z) = \ln\dfrac{1-\sqrt{x^2+y^2+z^2}}{1+\sqrt{x^2+y^2+z^2}} \;\Rightarrow$

$$f_y(x,y,z) = \frac{1}{\dfrac{1-\sqrt{x^2+y^2+z^2}}{1+\sqrt{x^2+y^2+z^2}}}\cdot \frac{\left(1+\sqrt{x^2+y^2+z^2}\right)\left(-\frac{1}{2}(x^2+y^2+z^2)^{-1/2}\cdot 2y\right)-\left(1-\sqrt{x^2+y^2+z^2}\right)\left(\frac{1}{2}(x^2+y^2+z^2)^{-1/2}\cdot 2y\right)}{\left(1+\sqrt{x^2+y^2+z^2}\right)^2}$$

$$= \frac{1+\sqrt{x^2+y^2+z^2}}{1-\sqrt{x^2+y^2+z^2}}\cdot\frac{-y(x^2+y^2+z^2)^{-1/2}\left(1+\sqrt{x^2+y^2+z^2}+1-\sqrt{x^2+y^2+z^2}\right)}{\left(1+\sqrt{x^2+y^2+z^2}\right)^2}$$

$$= \frac{-y(x^2+y^2+z^2)^{-1/2}(2)}{\left(1-\sqrt{x^2+y^2+z^2}\right)\left(1+\sqrt{x^2+y^2+z^2}\right)} = \frac{-2y}{\sqrt{x^2+y^2+z^2}\,[1-(x^2+y^2+z^2)]}$$

so $f_y(1,2,2) = \dfrac{-2(2)}{\sqrt{1^2+2^2+2^2}\,[1-(1^2+2^2+2^2)]} = \dfrac{-4}{\sqrt{9}\,(1-9)} = \dfrac{1}{6}$.

44. $f(x,y,z) = x^{yz} \;\Rightarrow\; f_z(x,y,z) = (x^{yz}\ln x)(y) = yx^{yz}\ln x$, so $f_z(e,1,0) = 1e^{(1)(0)}\ln e = 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

45. $f(x,y) = xy^2 - x^3y \;\Rightarrow$

$$f_x(x,y) = \lim_{h\to 0}\frac{f(x+h,y)-f(x,y)}{h} = \lim_{h\to 0}\frac{(x+h)y^2-(x+h)^3y-(xy^2-x^3y)}{h}$$

$$= \lim_{h\to 0}\frac{h(y^2-3x^2y-3xyh-yh^2)}{h} = \lim_{h\to 0}(y^2-3x^2y-3xyh-yh^2) = y^2-3x^2y$$

$$f_y(x,y) = \lim_{h\to 0}\frac{f(x,y+h)-f(x,y)}{h} = \lim_{h\to 0}\frac{x(y+h)^2-x^3(y+h)-(xy^2-x^3y)}{h} = \lim_{h\to 0}\frac{h(2xy+xh-x^3)}{h}$$

$$= \lim_{h\to 0}(2xy+xh-x^3) = 2xy-x^3$$

46. $f(x,y) = \dfrac{x}{x+y^2} \;\Rightarrow$

$$f_x(x,y) = \lim_{h\to 0}\frac{f(x+h,y)-f(x,y)}{h} = \lim_{h\to 0}\frac{\frac{x+h}{x+h+y^2}-\frac{x}{x+y^2}}{h}\cdot\frac{(x+h+y^2)(x+y^2)}{(x+h+y^2)(x+y^2)}$$

$$= \lim_{h\to 0}\frac{(x+h)(x+y^2)-x(x+h+y^2)}{h(x+h+y^2)(x+y^2)} = \lim_{h\to 0}\frac{y^2h}{h(x+h+y^2)(x+y^2)}$$

$$= \lim_{h\to 0}\frac{y^2}{(x+h+y^2)(x+y^2)} = \frac{y^2}{(x+y^2)^2}$$

$$f_y(x,y) = \lim_{h\to 0}\frac{f(x,y+h)-f(x,y)}{h} = \lim_{h\to 0}\frac{\frac{x}{x+(y+h)^2}-\frac{x}{x+y^2}}{h}\cdot\frac{\left[x+(y+h)^2\right](x+y^2)}{\left[x+(y+h)^2\right](x+y^2)}$$

$$= \lim_{h\to 0}\frac{x(x+y^2)-x\left[x+(y+h)^2\right]}{h[x+(y+h)^2](x+y^2)} = \lim_{h\to 0}\frac{h(-2xy-xh)}{h[x+(y+h)^2](x+y^2)}$$

$$= \lim_{h\to 0}\frac{-2xy-xh}{[x+(y+h)^2](x+y^2)} = \frac{-2xy}{(x+y^2)^2}$$

47. $x^2+2y^2+3z^2=1 \;\Rightarrow\; \dfrac{\partial}{\partial x}(x^2+2y^2+3z^2) = \dfrac{\partial}{\partial x}(1) \;\Rightarrow\; 2x+0+6z\dfrac{\partial z}{\partial x}=0 \;\Rightarrow\; 6z\dfrac{\partial z}{\partial x} = -2x \;\Rightarrow$

$\dfrac{\partial z}{\partial x} = \dfrac{-2x}{6z} = -\dfrac{x}{3z}$, and $\dfrac{\partial}{\partial y}(x^2+2y^2+3z^2) = \dfrac{\partial}{\partial y}(1) \;\Rightarrow\; 0+4y+6z\dfrac{\partial z}{\partial y}=0 \;\Rightarrow\; 6z\dfrac{\partial z}{\partial y} = -4y \;\Rightarrow$

$\dfrac{\partial z}{\partial y} = \dfrac{-4y}{6z} = -\dfrac{2y}{3z}$.

48. $x^2-y^2+z^2-2z=4 \;\Rightarrow\; \dfrac{\partial}{\partial x}(x^2-y^2+z^2-2z) = \dfrac{\partial}{\partial x}(4) \;\Rightarrow\; 2x-0+2z\dfrac{\partial z}{\partial x}-2\dfrac{\partial z}{\partial x}=0 \;\Rightarrow$

$(2z-2)\dfrac{\partial z}{\partial x} = -2x \;\Rightarrow\; \dfrac{\partial z}{\partial x} = \dfrac{-2x}{2z-2} = \dfrac{x}{1-z}$, and $\dfrac{\partial}{\partial y}(x^2-y^2+z^2-2z) = \dfrac{\partial}{\partial y}(4) \;\Rightarrow$

$0-2y+2z\dfrac{\partial z}{\partial y}-2\dfrac{\partial z}{\partial y}=0 \;\Rightarrow\; (2z-2)\dfrac{\partial z}{\partial y} = 2y \;\Rightarrow\; \dfrac{\partial z}{\partial y} = \dfrac{2y}{2z-2} = \dfrac{y}{z-1}$.

49. $e^z = xyz \;\Rightarrow\; \dfrac{\partial}{\partial x}(e^z) = \dfrac{\partial}{\partial x}(xyz) \;\Rightarrow\; e^z\dfrac{\partial z}{\partial x} = y\left(x\dfrac{\partial z}{\partial x}+z\cdot 1\right) \;\Rightarrow\; e^z\dfrac{\partial z}{\partial x}-xy\dfrac{\partial z}{\partial x} = yz \;\Rightarrow$

$(e^z - xy)\dfrac{\partial z}{\partial x} = yz$, so $\dfrac{\partial z}{\partial x} = \dfrac{yz}{e^z - xy}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\dfrac{\partial}{\partial y}(e^z) = \dfrac{\partial}{\partial y}(xyz) \quad \Rightarrow \quad e^z \dfrac{\partial z}{\partial y} = x\left(y \dfrac{\partial z}{\partial x} + z \cdot 1\right) \quad \Rightarrow \quad e^z \dfrac{\partial z}{\partial y} - xy \dfrac{\partial z}{\partial y} = xz \quad \Rightarrow \quad (e^z - xy) \dfrac{\partial z}{\partial y} = xz$, so

$\dfrac{\partial z}{\partial y} = \dfrac{xz}{e^z - xy}$.

50. $yz + x \ln y = z^2 \quad \Rightarrow \quad \dfrac{\partial}{\partial x}(yz + x \ln y) = \dfrac{\partial}{\partial x}(z^2) \quad \Rightarrow \quad y \dfrac{\partial z}{\partial x} + \ln y = 2z \dfrac{\partial z}{\partial x} \quad \Rightarrow \quad \ln y = 2z \dfrac{\partial z}{\partial x} - y \dfrac{\partial z}{\partial x} \quad \Rightarrow$

$\ln y = (2z - y) \dfrac{\partial z}{\partial x}$, so $\dfrac{\partial z}{\partial x} = \dfrac{\ln y}{2z - y}$.

$\dfrac{\partial}{\partial y}(yz + x \ln y) = \dfrac{\partial}{\partial y}(z^2) \quad \Rightarrow \quad y \dfrac{\partial z}{\partial y} + z \cdot 1 + x \cdot \dfrac{1}{y} = 2z \dfrac{\partial z}{\partial y} \quad \Rightarrow \quad z + \dfrac{x}{y} = 2z \dfrac{\partial z}{\partial y} - y \dfrac{\partial z}{\partial y} \quad \Rightarrow$

$z + \dfrac{x}{y} = (2z - y) \dfrac{\partial z}{\partial y}$, so $\dfrac{\partial z}{\partial y} = \dfrac{z + (x/y)}{2z - y} = \dfrac{x + yz}{y(2z - y)}$.

51. (a) $z = f(x) + g(y) \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = f'(x), \quad \dfrac{\partial z}{\partial y} = g'(y)$

(b) $z = f(x + y)$. Let $u = x + y$. Then $\dfrac{\partial z}{\partial x} = \dfrac{df}{du}\dfrac{\partial u}{\partial x} = \dfrac{df}{du}(1) = f'(u) = f'(x + y)$,

$\dfrac{\partial z}{\partial y} = \dfrac{df}{du}\dfrac{\partial u}{\partial y} = \dfrac{df}{du}(1) = f'(u) = f'(x + y)$.

52. (a) $z = f(x)g(y) \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = f'(x)g(y), \quad \dfrac{\partial z}{\partial y} = f(x)g'(y)$

(b) $z = f(xy)$. Let $u = xy$. Then $\dfrac{\partial u}{\partial x} = y$ and $\dfrac{\partial u}{\partial y} = x$. Hence $\dfrac{\partial z}{\partial x} = \dfrac{df}{du}\dfrac{\partial u}{\partial x} = \dfrac{df}{du} \cdot y = yf'(u) = yf'(xy)$

and $\dfrac{\partial z}{\partial y} = \dfrac{df}{du}\dfrac{\partial u}{\partial y} = \dfrac{df}{du} \cdot x = xf'(u) = xf'(xy)$.

(c) $z = f\left(\dfrac{x}{y}\right)$. Let $u = \dfrac{x}{y}$. Then $\dfrac{\partial u}{\partial x} = \dfrac{1}{y}$ and $\dfrac{\partial u}{\partial y} = -\dfrac{x}{y^2}$. Hence $\dfrac{\partial z}{\partial x} = \dfrac{df}{du}\dfrac{\partial u}{\partial x} = f'(u)\dfrac{1}{y} = \dfrac{f'(x/y)}{y}$

and $\dfrac{\partial z}{\partial y} = \dfrac{df}{du}\dfrac{\partial u}{\partial y} = f'(u)\left(-\dfrac{x}{y^2}\right) = -\dfrac{xf'(x/y)}{y^2}$.

53. $f(x, y) = x^4y - 2x^3y^2 \quad \Rightarrow \quad f_x(x, y) = 4x^3y - 6x^2y^2, \quad f_y(x, y) = x^4 - 4x^3y$. Then $f_{xx}(x, y) = 12x^2y - 12xy^2$,

$f_{xy}(x, y) = 4x^3 - 12x^2y$, $f_{yx}(x, y) = 4x^3 - 12x^2y$, and $f_{yy}(x, y) = -4x^3$.

54. $f(x, y) = \ln(ax + by) \quad \Rightarrow \quad f_x(x, y) = \dfrac{a}{ax + by} = a(ax + by)^{-1}, \quad f_y(x, y) = \dfrac{b}{ax + by} = b(ax + by)^{-1}$. Then

$f_{xx}(x, y) = -a(ax + by)^{-2}(a) = -\dfrac{a^2}{(ax + by)^2}, \quad f_{xy}(x, y) = -a(ax + by)^{-2}(b) = -\dfrac{ab}{(ax + by)^2}$,

$f_{yx}(x, y) = -b(ax + by)^{-2}(a) = -\dfrac{ab}{(ax + by)^2}$, and $f_{yy}(x, y) = -b(ax + by)^{-2}(b) = -\dfrac{b^2}{(ax + by)^2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

55. $z = \dfrac{y}{2x+3y} = y(2x+3y)^{-1} \quad \Rightarrow \quad z_x = y(-1)(2x+3y)^{-2}(2) = -\dfrac{2y}{(2x+3y)^2}$,

$z_y = \dfrac{(2x+3y)\cdot 1 - y\cdot 3}{(2x+3y)^2} = \dfrac{2x}{(2x+3y)^2}$. Then $z_{xx} = -2y(-2)(2x+3y)^{-3}(2) = \dfrac{8y}{(2x+3y)^3}$,

$z_{xy} = -\dfrac{(2x+3y)^2\cdot 2 - 2y\cdot 2(2x+3y)(3)}{[(2x+3y)^2]^2} = -\dfrac{(2x+3y)\,(4x+6y-12y)}{(2x+3y)^4} = \dfrac{6y-4x}{(2x+3y)^3}$,

$z_{yx} = \dfrac{(2x+3y)^2\cdot 2 - 2x\cdot 2(2x+3y)(2)}{[(2x+3y)^2]^2} = \dfrac{6y-4x}{(2x+3y)^3}$, $z_{yy} = 2x(-2)(2x+3y)^{-3}(3) = -\dfrac{12x}{(2x+3y)^3}$.

56. $T = e^{-2r}\cos\theta \quad \Rightarrow \quad T_r = -2e^{-2r}\cos\theta$, $T_\theta = -e^{-2r}\sin\theta$. Then $T_{rr} = -2e^{-2r}(-2)\cos\theta = 4e^{-2r}\cos\theta$,

$T_{r\theta} = 2e^{-2r}\sin\theta$, $T_{\theta r} = -e^{-2r}(-2)\sin\theta = 2e^{-2r}\sin\theta$, $T_{\theta\theta} = -e^{-2r}\cos\theta$.

57. $v = \sin(s^2-t^2) \quad \Rightarrow \quad v_s = \cos(s^2-t^2)\cdot 2s = 2s\cos(s^2-t^2)$, $v_t = \cos(s^2-t^2)\cdot(-2t) = -2t\cos(s^2-t^2)$. Then

$v_{ss} = 2s\left[-\sin(s^2-t^2)\cdot 2s\right] + \cos(s^2-t^2)\cdot 2 = 2\cos(s^2-t^2) - 4s^2\sin(s^2-t^2)$,

$v_{st} = 2s\left[-\sin(s^2-t^2)\cdot(-2t)\right] = 4st\sin(s^2-t^2)$, $v_{ts} = -2t\left[-\sin(s^2-t^2)\cdot 2s\right] = 4st\sin(s^2-t^2)$,

$v_{tt} = -2t\cdot\left[-\sin(s^2-t^2)\cdot(-2t)\right] + \cos(s^2-t^2)\cdot(-2) = -2\cos(s^2-t^2) - 4t^2\sin(s^2-t^2)$.

58. $w = \sqrt{1+uv^2} \quad \Rightarrow \quad w_u = \frac{1}{2}(1+uv^2)^{-1/2}\cdot v^2 = \dfrac{v^2}{2\sqrt{1+uv^2}}$, $w_v = \frac{1}{2}(1+uv^2)^{-1/2}\cdot 2uv = \dfrac{uv}{\sqrt{1+uv^2}}$.

Then $w_{uu} = \frac{1}{2}v^2\left(-\frac{1}{2}\right)(1+uv^2)^{-3/2}(v^2) = -\dfrac{v^4}{4(1+uv^2)^{3/2}}$,

$$w_{uv} = \frac{2\sqrt{1+uv^2}\cdot 2v - v^2\cdot 2\left(\frac{1}{2}\right)(1+uv^2)^{-1/2}(2uv)}{\left(2\sqrt{1+uv^2}\right)^2} = \frac{4v\sqrt{1+uv^2} - 2uv^3/\sqrt{1+uv^2}}{4(1+uv^2)}$$

$$= \frac{4v(1+uv^2) - 2uv^3}{4\,(1+uv^2)^{3/2}} - \frac{2v+uv^3}{2\,(1+uv^2)^{3/2}}$$

$$w_{vv} = \frac{\sqrt{1+uv^2}\cdot v - uv\cdot\frac{1}{2}(1+uv^2)^{-1/2}(v^2)}{\left(\sqrt{1+uv^2}\right)^2} = \frac{v\sqrt{1+uv^2} - \frac{1}{2}uv^3/\sqrt{1+uv^2}}{(1+uv^2)}$$

$$= \frac{v(1+uv^2) - \frac{1}{2}uv^3}{(1+uv^2)^{3/2}} - \frac{2v+uv^3}{2\,(1+uv^2)^{3/2}}$$

$$w_{uv} = \frac{\sqrt{1+uv^2}\cdot u - uv\cdot\frac{1}{2}(1+uv^2)^{-1/2}(2uv)}{\left(\sqrt{1+uv^2}\right)^2} = \frac{u\sqrt{1+uv^2} - u^2v^2/\sqrt{1+uv^2}}{(1+uv^2)}$$

$$- \frac{u(1+uv^2) - u^2v^2}{(1+uv^2)^{3/2}} - \frac{u}{(1+uv^2)^{3/2}}$$

59. $u = x^4y^3 - y^4 \quad \Rightarrow \quad u_x = 4x^3y^3$, $u_{xy} = 12x^3y^2$ and $u_y = 3x^4y^2 - 4y^3$, $u_{yx} = 12x^3y^2$.

Thus $u_{xy} = u_{yx}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. $u = e^{xy}\sin y \;\Rightarrow\; u_x = ye^{xy}\sin y,\; u_{xy} = ye^{xy}\cos y + (\sin y)(y\cdot xe^{xy} + e^{xy}\cdot 1) = e^{xy}(y\cos y + xy\sin y + \sin y)$,

$u_y = e^{xy}\cos y + (\sin y)(xe^{xy}) = e^{xy}(\cos y + x\sin y)$,

$u_{yx} = e^{xy}\cdot\sin y + (\cos y + x\sin y)\cdot ye^{xy} = e^{xy}(\sin y + y\cos y + xy\sin y)$. Thus $u_{xy} = u_{yx}$.

61. $u = \cos(x^2y) \;\Rightarrow\; u_x = -\sin(x^2y)\cdot 2xy = -2xy\sin(x^2y)$,

$u_{xy} = -2xy\cdot\cos(x^2y)\cdot x^2 + \sin(x^2y)\cdot(-2x) = -2x^3y\cos(x^2y) - 2x\sin(x^2y)$ and

$u_y = -\sin(x^2y)\cdot x^2 = -x^2\sin(x^2y),\; u_{yx} = -x^2\cdot\cos(x^2y)\cdot 2xy + \sin(x^2y)\cdot(-2x) = -2x^3y\cos(x^2y) - 2x\sin(x^2y)$.

Thus $u_{xy} = u_{yx}$.

62. $u = \ln(x+2y) \;\Rightarrow\; u_x = \dfrac{1}{x+2y} = (x+2y)^{-1},\; u_{xy} = (-1)(x+2y)^{-2}(2) = -\dfrac{2}{(x+2y)^2}$ and

$u_y = \dfrac{1}{x+2y}\cdot 2 = 2(x+2y)^{-1},\; u_{yx} = (-2)(x+2y)^{-2} = -\dfrac{2}{(x+2y)^2}$. Thus $u_{xy} = u_{yx}$.

63. $f(x,y) = x^4y^2 - x^3y \;\Rightarrow\; f_x = 4x^3y^2 - 3x^2y,\; f_{xx} = 12x^2y^2 - 6xy,\; f_{xxx} = 24xy^2 - 6y$ and

$f_{xy} = 8x^3y - 3x^2,\; f_{xyx} = 24x^2y - 6x$.

64. $f(x,y) = \sin(2x+5y) \;\Rightarrow\; f_y = \cos(2x+5y)\cdot 5 = 5\cos(2x+5y),\; f_{yx} = -5\sin(2x+5y)\cdot 2 = -10\sin(2x+5y)$,

$f_{yxy} = -10\cos(2x+5y)\cdot 5 = -50\cos(2x+5y)$

65. $f(x,y,z) = e^{xyz^2} \;\Rightarrow\; f_x = e^{xyz^2}\cdot yz^2 = yz^2e^{xyz^2},\; f_{xy} = yz^2\cdot e^{xyz^2}(xz^2) + e^{xyz^2}\cdot z^2 = (xyz^4 + z^2)e^{xyz^2}$,

$f_{xyz} = (xyz^4 + z^2)\cdot e^{xyz^2}(2xyz) + e^{xyz^2}\cdot(4xyz^3 + 2z) = (2x^2y^2z^5 + 6xyz^3 + 2z)e^{xyz^2}$.

66. $g(r,s,t) = e^r\sin(st) \;\Rightarrow\; g_r = e^r\sin(st),\; g_{rs} = e^r\cos(st)\cdot t = te^r\cos(st)$,

$g_{rst} = te^r(-\sin(st)\cdot s) + \cos(st)\cdot e^r = e^r[\cos(st) - st\sin(st)]$.

67. $W = \sqrt{u+v^2} \;\Rightarrow\; \dfrac{\partial W}{\partial v} = \frac{1}{2}(u+v^2)^{-1/2}(2v) = v(u+v^2)^{-1/2}$,

$\dfrac{\partial^2 W}{\partial u\,\partial v} = v\left(-\frac{1}{2}\right)(u+v^2)^{-3/2}(1) = -\frac{1}{2}v(u+v^2)^{-3/2},\; \dfrac{\partial^3 W}{\partial u^2\,\partial v} = -\frac{1}{2}v\left(-\frac{3}{2}\right)(u+v^2)^{-5/2}(1) = \frac{3}{4}v(u+v^2)^{-5/2}$.

68. $V = \ln(r+s^2+t^3) \;\Rightarrow\; \dfrac{\partial V}{\partial t} = \dfrac{3t^2}{r+s^2+t^3} = 3t^2(r+s^2+t^3)^{-1}$,

$\dfrac{\partial^2 V}{\partial s\,\partial t} = 3t^2(-1)(r+s^2+t^3)^{-2}(2s) = -6st^2(r+s^2+t^3)^{-2}$,

$\dfrac{\partial^3 V}{\partial r\,\partial s\,\partial t} = -6st^2(-2)(r+s^2+t^3)^{-3}(1) = 12st^2(r+s^2+t^3)^{-3} = \dfrac{12st^2}{(r+s^2+t^3)^3}$.

69. $w = \dfrac{x}{y+2z} = x(y+2z)^{-1} \;\Rightarrow\; \dfrac{\partial w}{\partial x} = (y+2z)^{-1},\; \dfrac{\partial^2 w}{\partial y\,\partial x} = -(y+2z)^{-2}(1) = -(y+2z)^{-2}$,

$\dfrac{\partial^3 w}{\partial z\,\partial y\,\partial x} = -(-2)(y+2z)^{-3}(2) = 4(y+2z)^{-3} = \dfrac{4}{(y+2z)^3}$ and $\dfrac{\partial w}{\partial y} = x(-1)(y+2z)^{-2}(1) = -x(y+2z)^{-2}$,

$\dfrac{\partial^2 w}{\partial x\,\partial y} = -(y+2z)^{-2},\; \dfrac{\partial^3 w}{\partial x^2\,\partial y} = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

70. $u = x^a y^b z^c$. If $a = 0$, or if $b = 0$ or 1, or if $c = 0$, 1, or 2, then $\dfrac{\partial^6 u}{\partial x\,\partial y^2\,\partial z^3} = 0$. Otherwise $\dfrac{\partial u}{\partial z} = cx^a y^b z^{c-1}$,

$\dfrac{\partial^2 u}{\partial z^2} = c(c-1)x^a y^b z^{c-2}$, $\dfrac{\partial^3 u}{\partial z^3} = c(c-1)(c-2)x^a y^b z^{c-3}$, $\dfrac{\partial^4 u}{\partial y\,\partial z^3} = bc(c-1)(c-2)x^a y^{b-1} z^{c-3}$,

$\dfrac{\partial^5 u}{\partial y^2\,\partial z^3} = b(b-1)c(c-1)(c-2)x^a y^{b-2} z^{c-3}$, and $\dfrac{\partial^6 u}{\partial x\,\partial y^2\,\partial z^3} = ab(b-1)c(c-1)(c-2)x^{a-1} y^{b-2} z^{c-3}$.

71. Assuming that the third partial derivatives of f are continuous (easily verified), we can write $f_{xzy} = f_{yxz}$. Then

$f(x, y, z) = xy^2 z^3 + \arcsin\left(x\sqrt{z}\right) \;\Rightarrow\; f_y = 2xyz^3 + 0$, $f_{yx} = 2yz^3$, and $f_{yxz} = 6yz^2 = f_{xzy}$.

72. Let $f(x, y, z) = \sqrt{1 + xz}$ and $h(x, y, z) = \sqrt{1 - xy}$ so that $g = f + h$. Then $f_y = 0 = f_{yx} = f_{yxz}$ and $h_z = 0 = h_{zx} = h_{zxy}$. But (since the partial derivatives are continous on their domains) $f_{xyz} = f_{yxz}$ and $h_{xyz} = h_{zxy}$, so $g_{xyz} = f_{xyz} + h_{xyz} = 0 + 0 = 0$.

73. By Definition 4, $f_x(3, 2) = \lim\limits_{h \to 0} \dfrac{f(3 + h, 2) - f(3, 2)}{h}$ which we can approximate by considering $h = 0.5$ and $h = -0.5$:

$f_x(3, 2) \approx \dfrac{f(3.5, 2) - f(3, 2)}{0.5} = \dfrac{22.4 - 17.5}{0.5} = 9.8$, $f_x(3, 2) \approx \dfrac{f(2.5, 2) - f(3, 2)}{-0.5} = \dfrac{10.2 - 17.5}{-0.5} = 14.6$. Averaging these values, we estimate $f_x(3, 2)$ to be approximately 12.2. Similarly, $f_x(3, 2.2) = \lim\limits_{h \to 0} \dfrac{f(3 + h, 2.2) - f(3, 2.2)}{h}$ which we can approximate by considering $h = 0.5$ and $h = -0.5$: $f_x(3, 2.2) \approx \dfrac{f(3.5, 2.2) - f(3, 2.2)}{0.5} = \dfrac{26.1 - 15.9}{0.5} = 20.4$,

$f_x(3, 2.2) \approx \dfrac{f(2.5, 2.2) - f(3, 2.2)}{-0.5} = \dfrac{9.3 - 15.9}{-0.5} = 13.2$. Averaging these values, we have $f_x(3, 2.2) \approx 16.8$.

To estimate $f_{xy}(3, 2)$, we first need an estimate for $f_x(3, 1.8)$:

$f_x(3, 1.8) \approx \dfrac{f(3.5, 1.8) - f(3, 1.8)}{0.5} = \dfrac{20.0 - 18.1}{0.5} = 3.8$, $f_x(3, 1.8) \approx \dfrac{f(2.5, 1.8) - f(3, 1.8)}{-0.5} = \dfrac{12.5 - 18.1}{-0.5} = 11.2$.

Averaging these values, we get $f_x(3, 1.8) \approx 7.5$. Now $f_{xy}(x, y) = \dfrac{\partial}{\partial y}\left[f_x(x, y)\right]$ and $f_x(x, y)$ is itself a function of two variables, so Definition 4 says that $f_{xy}(x, y) = \dfrac{\partial}{\partial y}\left[f_x(x, y)\right] = \lim\limits_{h \to 0} \dfrac{f_x(x, y + h) - f_x(x, y)}{h}$ $\Rightarrow$

$f_{xy}(3, 2) = \lim\limits_{h \to 0} \dfrac{f_x(3, 2 + h) - f_x(3, 2)}{h}$. We can estimate this value using our previous work with $h = 0.2$ and $h = -0.2$:

$f_{xy}(3, 2) \approx \dfrac{f_x(3, 2.2) - f_x(3, 2)}{0.2} = \dfrac{16.8 - 12.2}{0.2} = 23$, $f_{xy}(3, 2) \approx \dfrac{f_x(3, 1.8) - f_x(3, 2)}{-0.2} = \dfrac{7.5 - 12.2}{-0.2} = 23.5$.

Averaging these values, we estimate $f_{xy}(3, 2)$ to be approximately 23.25.

74. (a) If we fix y and allow x to vary, the level curves indicate that the value of f decreases as we move through P in the positive x-direction, so f_x is negative at P.

(b) If we fix x and allow y to vary, the level curves indicate that the value of f increases as we move through P in the positive y-direction, so f_y is positive at P.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $f_{xx} = \dfrac{\partial}{\partial x}(f_x)$, so if we fix y and allow x to vary, f_{xx} is the rate of change of f_x as x increases. Note that at points to the right of P the level curves are spaced farther apart (in the x-direction) than at points to the left of P, demonstrating that f decreases less quickly with respect to x to the right of P. So as we move through P in the positive x-direction the (negative) value of f_x increases, hence $\dfrac{\partial}{\partial x}(f_x) = f_{xx}$ is positive at P.

(d) $f_{xy} = \dfrac{\partial}{\partial y}(f_x)$, so if we fix x and allow y to vary, f_{xy} is the rate of change of f_x as y increases. The level curves are closer together (in the x-direction) at points above P than at those below P, demonstrating that f decreases more quickly with respect to x for y-values above P. So as we move through P in the positive y-direction, the (negative) value of f_x decreases, hence f_{xy} is negative.

(e) $f_{yy} = \dfrac{\partial}{\partial y}(f_y)$, so if we fix x and allow y to vary, f_{yy} is the rate of change of f_y as y increases. The level curves are closer together (in the y-direction) at points above P than at those below P, demonstrating that f increases more quickly with respect to y above P. So as we move through P in the positive y-direction the (positive) value of f_y increases, hence $\dfrac{\partial}{\partial y}(f_y) = f_{yy}$ is positive at P.

75. $u = e^{-\alpha^2 k^2 t}\sin kx \;\Rightarrow\; u_x = ke^{-\alpha^2 k^2 t}\cos kx,\; u_{xx} = -k^2 e^{-\alpha^2 k^2 t}\sin kx,$ and $u_t = -\alpha^2 k^2 e^{-\alpha^2 k^2 t}\sin kx$. Thus $\alpha^2 u_{xx} = u_t$.

76. (a) $u = x^2 + y^2 \;\Rightarrow\; u_x = 2x,\; u_{xx} = 2;\quad u_y = 2y,\; u_{yy} = 2$. Thus $u_{xx} + u_{yy} \neq 0$ and $u = x^2 + y^2$ does not satisfy Laplace's Equation.

(b) $u = x^2 - y^2$ is a solution: $u_{xx} = 2,\; u_{yy} = -2$ so $u_{xx} + u_{yy} = 0$.

(c) $u = x^3 + 3xy^2$ is not a solution: $u_x = 3x^2 + 3y^2,\; u_{xx} = 6x;\quad u_y = 6xy,\; u_{yy} = 6x$.

(d) $u = \ln\sqrt{x^2+y^2}$ is a solution: $u_x = \dfrac{1}{\sqrt{x^2+y^2}}\left(\dfrac{1}{2}\right)(x^2+y^2)^{-1/2}(2x) = \dfrac{x}{x^2+y^2}$,

$u_{xx} = \dfrac{(x^2+y^2) - x(2x)}{(x^2+y^2)^2} = \dfrac{y^2 - x^2}{(x^2+y^2)^2}$. By symmetry, $u_{yy} = \dfrac{x^2 - y^2}{(x^2+y^2)^2}$, so $u_{xx} + u_{yy} = 0$.

(e) $u = \sin x\,\cosh y + \cos x\,\sinh y$ is a solution: $u_x = \cos x\,\cosh y - \sin x\,\sinh y, u_{xx} = -\sin x\,\cosh y - \cos x\,\sinh y$, and $u_y = \sin x\,\sinh y + \cos x\,\cosh y, u_{yy} = \sin x\,\cosh y + \cos x\,\sinh y$.

(f) $u = e^{-x}\cos y - e^{-y}\cos x$ is a solution: $u_x = -e^{-x}\cos y + e^{-y}\sin x,\quad u_{xx} = e^{-x}\cos y + e^{-y}\cos x$, and $u_y = -e^{-x}\sin y + e^{-y}\cos x,\quad u_{yy} = -e^{-x}\cos y - e^{-y}\cos x$.

77. $u = \dfrac{1}{\sqrt{x^2+y^2+z^2}} \;\Rightarrow\; u_x = \left(-\tfrac{1}{2}\right)(x^2+y^2+z^2)^{-3/2}(2x) = -x(x^2+y^2+z^2)^{-3/2}$ and

$u_{xx} = -(x^2+y^2+z^2)^{-3/2} - x\left(-\tfrac{3}{2}\right)(x^2+y^2+z^2)^{-5/2}(2x) = \dfrac{2x^2 - y^2 - z^2}{(x^2+y^2+z^2)^{5/2}}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

By symmetry, $u_{yy} = \dfrac{2y^2 - x^2 - z^2}{(x^2+y^2+z^2)^{5/2}}$ and $u_{zz} = \dfrac{2z^2 - x^2 - y^2}{(x^2+y^2+z^2)^{5/2}}$.

Thus $u_{xx} + u_{yy} + u_{zz} = \dfrac{2x^2 - y^2 - z^2 + 2y^2 - x^2 - z^2 + 2z^2 - x^2 - y^2}{(x^2+y^2+z^2)^{5/2}} = 0.$

78. (a) $u = \sin(kx)\,\sin(akt) \;\Rightarrow\; u_t = ak\sin(kx)\,\cos(akt)$, $u_{tt} = -a^2k^2\sin(kx)\,\sin(akt)$, $u_x = k\cos(kx)\,\sin(akt)$, $u_{xx} = -k^2\sin(kx)\,\sin(akt)$. Thus $u_{tt} = a^2u_{xx}$.

(b) $u = \dfrac{t}{a^2t^2 - x^2} \;\Rightarrow\; u_t = \dfrac{(a^2t^2 - x^2) - t(2a^2t)}{(a^2t^2 - x^2)^2} = -\dfrac{a^2t^2 + x^2}{(a^2t^2 - x^2)^2}$,

$u_{tt} = \dfrac{-2a^2t(a^2t^2 - x^2)^2 + (a^2t^2 + x^2)(2)(a^2t^2 - x^2)(2a^2t)}{(a^2t^2 - x^2)^4} = \dfrac{2a^4t^3 + 6a^2tx^2}{(a^2t^2 - x^2)^3}$,

$u_x = t(-1)(a^2t^2 - x^2)^{-2}(-2x) = \dfrac{2tx}{(a^2t^2 - x^2)^2}$,

$u_{xx} = \dfrac{2t(a^2t^2 - x^2)^2 - 2tx\,(2)(a^2t^2 - x^2)(-2x)}{(a^2t^2 - x^2)^4} = \dfrac{2a^2t^3 - 2tx^2 + 8tx^2}{(a^2t^2 - x^2)^3} = \dfrac{2a^2t^3 + 6tx^2}{(a^2t^2 - x^2)^3}$.

Thus $u_{tt} = a^2u_{xx}$.

(c) $u = (x - at)^6 + (x + at)^6 \;\Rightarrow\; u_t = -6a(x - at)^5 + 6a(x + at)^5$, $u_{tt} = 30a^2(x - at)^4 + 30a^2(x + at)^4$, $u_x = 6(x - at)^5 + 6(x + at)^5$, $u_{xx} = 30(x - at)^4 + 30(x + at)^4$. Thus $u_{tt} = a^2u_{xx}$.

(d) $u = \sin(x - at) + \ln(x + at) \;\Rightarrow\; u_t = -a\cos(x - at) + \dfrac{a}{x + at}$, $u_{tt} = -a^2\sin(x - at) - \dfrac{a^2}{(x + at)^2}$, $u_x = \cos(x - at) + \dfrac{1}{x + at}$, $u_{xx} = -\sin(x - at) - \dfrac{1}{(x + at)^2}$. Thus $u_{tt} = a^2u_{xx}$.

79. Let $v = x + at$, $w = x - at$. Then $u_t = \dfrac{\partial[f(v) + g(w)]}{\partial t} = \dfrac{df(v)}{dv}\dfrac{\partial v}{\partial t} + \dfrac{dg(w)}{dw}\dfrac{\partial w}{\partial t} = af'(v) - ag'(w)$ and $u_{tt} = \dfrac{\partial[af'(v) - ag'(w)]}{\partial t} = a[af''(v) + ag''(w)] = a^2[f''(v) + g''(w)]$. Similarly, by using the Chain Rule we have $u_x = f'(v) + g'(w)$ and $u_{xx} = f''(v) + g''(w)$. Thus $u_{tt} = a^2u_{xx}$.

80. For each i, $i = 1, \ldots, n$, $\partial u/\partial x_i = a_ie^{a_1x_1 + a_2x_2 + \cdots + a_nx_n}$ and $\partial^2u/\partial x_i^2 = a_i^2e^{a_1x_1 + a_2x_2 + \cdots + a_nx_n}$.

Then $\dfrac{\partial^2u}{\partial x_1^2} + \dfrac{\partial^2u}{\partial x_2^2} + \cdots + \dfrac{\partial^2u}{\partial x_n^2} = (a_1^2 + a_2^2 + \cdots + a_n^2)e^{a_1x_1 + a_2x_2 + \cdots + a_nx_n} = e^{a_1x_1 + a_2x_2 + \cdots + a_nx_n} = u$

since $a_1^2 + a_2^2 + \cdots + a_n^2 = 1$.

81. $c(x,t) = \dfrac{1}{\sqrt{4\pi Dt}}\,e^{-x^2/(4Dt)} \;\Rightarrow$

$$\frac{\partial c}{\partial t} = \frac{1}{\sqrt{4\pi Dt}}\cdot e^{-x^2/(4Dt)}\left[-x^2(-1)(4Dt)^{-2}(4D)\right] + e^{-x^2/(4Dt)}\cdot\left(-\tfrac{1}{2}\right)(4\pi Dt)^{-3/2}(4\pi D)$$

$$= (4\pi Dt)^{-3/2}\left(4\pi Dt\cdot\frac{x^2}{4Dt^2} - 2\pi D\right)e^{-x^2/(4Dt)} = \frac{2\pi D}{(4\pi Dt)^{3/2}}\left(\frac{x^2}{2Dt} - 1\right)e^{-x^2/(4Dt)},$$

$$\frac{\partial c}{\partial x} = \frac{1}{\sqrt{4\pi Dt}}\,e^{-x^2/(4Dt)}\cdot\frac{-2x}{4Dt} = \frac{-2\pi x}{(4\pi Dt)^{3/2}}\,e^{-x^2/(4Dt)}, \text{ and}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\frac{\partial^2 c}{\partial x^2} = \frac{-2\pi}{(4\pi Dt)^{3/2}}\left[x \cdot e^{-x^2/(4Dt)} \cdot \frac{-2x}{4Dt} + e^{-x^2/(4Dt)} \cdot 1\right]$$

$$= \frac{-2\pi}{(4\pi Dt)^{3/2}}\left(-\frac{x^2}{2Dt} + 1\right)e^{-x^2/(4Dt)} = \frac{2\pi}{(4\pi Dt)^{3/2}}\left(\frac{x^2}{2Dt} - 1\right)e^{-x^2/(4Dt)}.$$

Thus $\dfrac{\partial c}{\partial t} = \dfrac{2\pi D}{(4\pi Dt)^{3/2}}\left(\dfrac{x^2}{2Dt} - 1\right)e^{-x^2/(4Dt)} = D\left[\dfrac{2\pi}{(4\pi Dt)^{3/2}}\left(\dfrac{x^2}{2Dt} - 1\right)e^{-x^2/(4Dt)}\right] = D\dfrac{\partial^2 c}{\partial x^2}$.

82. (a) $\partial T/\partial x = -60(2x)/(1 + x^2 + y^2)^2$, so at $(2, 1)$, $T_x = -240/(1 + 4 + 1)^2 = -\frac{20}{3}$.

(b) $\partial T/\partial y = -60(2y)/(1 + x^2 + y^2)^2$, so at $(2, 1)$, $T_y = -120/36 = -\frac{10}{3}$. Thus from the point $(2, 1)$ the temperature is decreasing at a rate of $\frac{20}{3}$°C/m in the x-direction and is decreasing at a rate of $\frac{10}{3}$°C/m in the y-direction.

83. By the Chain Rule, taking the partial derivative of both sides with respect to R_1 gives

$$\frac{\partial R^{-1}}{\partial R}\frac{\partial R}{\partial R_1} = \frac{\partial\left[(1/R_1) + (1/R_2) + (1/R_3)\right]}{\partial R_1} \quad \text{or} \quad -R^{-2}\frac{\partial R}{\partial R_1} = -R_1^{-2}. \text{ Thus } \frac{\partial R}{\partial R_1} = \frac{R^2}{R_1^2}.$$

84. $P = bL^\alpha K^\beta$, so $\dfrac{\partial P}{\partial L} = \alpha b L^{\alpha-1}K^\beta$ and $\dfrac{\partial P}{\partial K} = \beta b L^\alpha K^{\beta-1}$. Then

$$L\frac{\partial P}{\partial L} + K\frac{\partial P}{\partial K} = L(\alpha b L^{\alpha-1}K^\beta) + K(\beta b L^\alpha K^{\beta-1}) = \alpha b L^{1+\alpha-1}K^\beta + \beta b L^\alpha K^{1+\beta-1} = (\alpha+\beta)bL^\alpha K^\beta = (\alpha+\beta)P$$

85. If we fix $K = K_0$, $P(L, K_0)$ is a function of a single variable L, and $\dfrac{dP}{dL} = \alpha\dfrac{P}{L}$ is a separable differential equation. Then

$$\frac{dP}{P} = \alpha\frac{dL}{L} \quad \Rightarrow \quad \int\frac{dP}{P} = \int\alpha\frac{dL}{L} \quad \Rightarrow \quad \ln|P| = \alpha\ln|L| + C\,(K_0), \text{ where } C(K_0) \text{ can depend on } K_0. \text{ Then}$$

$|P| = e^{\alpha\ln|L| + C(K_0)}$, and since $P > 0$ and $L > 0$, we have $P = e^{\alpha\ln L}e^{C(K_0)} = e^{C(K_0)}e^{\ln L^\alpha} = C_1(K_0)L^\alpha$ where $C_1(K_0) = e^{C(K_0)}$.

86. (a) $P(L, K) = 1.01L^{0.75}K^{0.25} \quad \Rightarrow \quad P_L(L, K) = 1.01(0.75L^{-0.25})K^{0.25} = 0.7575L^{-0.25}K^{0.25}$ and

$P_K(L, K) = 1.01L^{0.75}(0.25K^{-0.75}) = 0.2525L^{0.75}K^{-0.75}$.

(b) The marginal productivity of labor in 1920 is $P_L(194, 407) = 0.7575(194)^{-0.25}(407)^{0.25} \approx 0.912$. Recall that P, L, and K are expressed as percentages of the respective amounts in 1899, so this means that in 1920, if the amount of labor is increased, production increases at a rate of about 0.912 percentage points per percentage point increase in labor. The marginal productivity of capital in 1920 is $P_K(194, 407) = 0.2525(194)^{0.75}(407)^{-0.75} \approx 0.145$, so an increase in capital investment would cause production to increase at a rate of about 0.145 percentage points per percentage point increase in capital.

(c) The value of $P_L(194, 407)$ is greater than the value of $P_K(194, 407)$, suggesting that increasing labor in 1920 would have increased production more than increasing capital.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

87. $\left(P+\frac{n^2a}{V^2}\right)(V-nb)=nRT \;\Rightarrow\; T=\frac{1}{nR}\left(P+\frac{n^2a}{V^2}\right)(V-nb)$, so $\frac{\partial T}{\partial P}=\frac{1}{nR}(1)(V-nb)=\frac{V-nb}{nR}$.

We can also write $P+\frac{n^2a}{V^2}=\frac{nRT}{V-nb} \;\Rightarrow\; P=\frac{nRT}{V-nb}-\frac{n^2a}{V^2}=nRT(V-nb)^{-1}-n^2aV^{-2}$, so

$\frac{\partial P}{\partial V}=-nRT(V-nb)^{-2}(1)+2n^2aV^{-3}=\frac{2n^2a}{V^3}-\frac{nRT}{(V-nb)^2}$.

88. $P=\frac{mRT}{V}$ so $\frac{\partial P}{\partial V}=\frac{-mRT}{V^2}$; $V=\frac{mRT}{P}$, so $\frac{\partial V}{\partial T}=\frac{mR}{P}$; $T=\frac{PV}{mR}$, so $\frac{\partial T}{\partial P}=\frac{V}{mR}$.

Thus $\frac{\partial P}{\partial V}\frac{\partial V}{\partial T}\frac{\partial T}{\partial P}=\frac{-mRT}{V^2}\frac{mR}{P}\frac{V}{mR}=\frac{-mRT}{PV}=-1$, since $PV=mRT$.

89. By Exercise 88, $PV=mRT \;\Rightarrow\; P=\frac{mRT}{V}$, so $\frac{\partial P}{\partial T}=\frac{mR}{V}$. Also, $PV=mRT \;\Rightarrow\; V=\frac{mRT}{P}$ and $\frac{\partial V}{\partial T}=\frac{mR}{P}$.

Since $T=\frac{PV}{mR}$, we have $T\,\frac{\partial P}{\partial T}\,\frac{\partial V}{\partial T}=\frac{PV}{mR}\cdot\frac{mR}{V}\cdot\frac{mR}{P}=mR$.

90. $\frac{\partial W}{\partial T}=0.6215+0.3965v^{0.16}$. When $T=-15^\circ\text{C}$ and $v=30$ km/h, $\frac{\partial W}{\partial T}=0.6215+0.3965(30)^{0.16}\approx 1.3048$, so we would expect the apparent temperature to drop by approximately 1.3°C if the actual temperature decreases by 1°C.

$\frac{\partial W}{\partial v}=-11.37(0.16)v^{-0.84}+0.3965T(0.16)v^{-0.84}$ and when $T=-15^\circ\text{C}$ and $v=30$ km/h,

$\frac{\partial W}{\partial v}=-11.37(0.16)(30)^{-0.84}+0.3965(-15)(0.16)(30)^{-0.84}\approx -0.1592$, so we would expect the apparent temperature to drop by approximately 0.16°C if the wind speed increases by 1 km/h.

91. (a) $S=f(w,h)=0.1091w^{0.425}h^{0.725} \;\Rightarrow\; \frac{\partial S}{\partial w}=0.1091(0.425)w^{0.425-1}h^{0.725}=0.0463675w^{-0.575}h^{0.725}$, so

$\frac{\partial S}{\partial w}(160,70)=0.0463675(160)^{-0.575}(70)^{0.725}\approx 0.0545$. This means that for a person 70 inches tall who weighs 160 pounds, an increase in weight (while height remains constant) causes the surface area to increase at a rate of about 0.0545 square feet (about 7.85 square inches) per pound.

(b) $\frac{\partial S}{\partial h}=0.1091(0.725)w^{0.425}h^{0.725-1}=0.0790975w^{0.425}h^{-0.275}$, so

$\frac{\partial S}{\partial h}(160,70)=0.0790975(160)^{0.425}(70)^{-0.275}\approx 0.213$. This means that for a person 70 inches tall who weighs 160 pounds, an increase in height (while weight remains unchanged at 160 pounds) causes the surface area to increase at a rate of about 0.213 square feet (about 30.7 square inches) per inch of height.

92. $R=C\frac{L}{r^4} \;\Rightarrow\; \frac{\partial R}{\partial L}=\frac{C}{r^4}$ and $\frac{\partial R}{\partial r}=CL\left(-4r^{-5}\right)=-4C\frac{L}{r^5}$.

$\partial R/\partial L$ is the rate at which the resistance of the flowing blood increases with respect to the length of the artery when the radius stays constant. $\partial R/\partial r$ is the rate of change of the resistance with respect to the radius of the artery when the length remains unchanged. Because $\partial R/\partial r$ is negative, the resistance decreases if the radius increases.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

93. $P(v,x,m) = Av^3 + \dfrac{B(mg/x)^2}{v} = Av^3 + Bm^2g^2x^{-2}v^{-1}$.

$\partial P/\partial v = 3Av^2 - \dfrac{B(mg/x)^2}{v^2}$ is the rate of change of the power needed during flapping mode with respect to the bird's velocity when the mass and fraction of flapping time remain constant. $\partial P/\partial x = -2Bm^2g^2x^{-3}v^{-1} = -\dfrac{2Bm^2g^2}{x^3v}$ is the rate at which the power changes with respect to the fraction of time spent in flapping mode when the mass and velocity are held constant. $\partial P/\partial m = 2Bmg^2x^{-2}v^{-1} = \dfrac{2Bmg^2}{x^2v}$ is the rate of change of the power with respect to mass when the velocity and fraction of flapping time remain constant.

94. $E(m,v) = 2.65m^{0.66} + \dfrac{3.5m^{0.75}}{v} \quad \Rightarrow$

$E_m(m,v) = 2.65(0.66)m^{0.66-1} + \dfrac{3.5(0.75)m^{0.75-1}}{v} = 1.749m^{-0.34} + \dfrac{2.625m^{-0.25}}{v}$,

$E_v(m,v) = 3.5m^{0.75}\left(-v^{\ -2}\right) = -\dfrac{3.5m^{0.75}}{v^2}$. Then $E_m(400,8) = 1.749(400)^{-0.34} + \dfrac{2.625(400)^{-0.25}}{8} \approx 0.301$ which means that the average energy needed for a lizard to walk or run 1 km increases at a rate of about 0.301 kcal per gram of body mass increase from 400 g if the speed is 8 km/h. $E_v(400,8) = -\dfrac{3.5(400)^{0.75}}{8^2} \approx -4.89$, which means that the average energy needed by a lizard with body mass 400 g decreases at a rate of about 4.89 kcal per km/h when the speed increases from 8 km/h.

95. $\dfrac{\partial K}{\partial m} = \frac{1}{2}v^2$, $\dfrac{\partial K}{\partial v} = mv$, $\dfrac{\partial^2 K}{\partial v^2} = m$. Thus $\dfrac{\partial K}{\partial m}\cdot\dfrac{\partial^2 K}{\partial v^2} = \frac{1}{2}v^2 m = K$.

96. The Law of Cosines says that $a^2 = b^2 + c^2 - 2bc\cos A$. Thus $\dfrac{\partial(a^2)}{\partial a} = \dfrac{\partial(b^2 + c^2 - 2ab\cos A)}{\partial a}$ or

$2a = -2bc\,(-\sin A)\dfrac{\partial A}{\partial a}$, implying that $\dfrac{\partial A}{\partial a} = \dfrac{a}{bc\sin A}$. Taking the partial derivative of both sides with respect to b gives $0 = 2b - 2c(\cos A) - 2bc\,(-\sin A)\dfrac{\partial A}{\partial b}$. Thus $\dfrac{\partial A}{\partial b} = \dfrac{c\cos A - b}{bc\sin A}$. By symmetry, $\dfrac{\partial A}{\partial c} = \dfrac{b\cos A - c}{bc\sin A}$.

97. $f_x(x,y) = x + 4y \quad \Rightarrow \quad f_{xy}(x,y) = 4$ and $f_y(x,y) = 3x - y \quad \Rightarrow \quad f_{yx}(x,y) = 3$. Since f_{xy} and f_{yx} are continuous everywhere but $f_{xy}(x,y) \neq f_{yx}(x,y)$, Clairaut's Theorem implies that such a function $f(x,y)$ does not exist.

98. Setting $x = 1$, the equation of the parabola of intersection is $z = 6 - 1 - 1 - 2y^2 = 4 - 2y^2$. The slope of the tangent is $\partial z/\partial y = -4y$, so at $(1,2,-4)$ the slope is -8. Parametric equations for the line are therefore $x = 1,\ y = 2 + t$, $z = -4 - 8t$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

99. By the geometry of partial derivatives, the slope of the tangent line is $f_x(1,2)$. By implicit differentiation of $4x^2 + 2y^2 + z^2 = 16$, we get $8x + 2z\,(\partial z/\partial x) = 0 \;\Rightarrow\; \partial z/\partial x = -4x/z$, so when $x = 1$ and $z = 2$ we have $\partial z/\partial x = -2$. So the slope is $f_x(1,2) = -2$. Thus the tangent line is given by $z - 2 = -2(x-1),\ y = 2$. Taking the parameter to be $t = x - 1$, we can write parametric equations for this line: $x = 1 + t,\ y = 2,\ z = 2 - 2t$.

100. $T(x,t) = T_0 + T_1 e^{-\lambda x}\sin(\omega t - \lambda x)$

(a) $\partial T/\partial x = T_1 e^{-\lambda x}\,[\cos(\omega t - \lambda x)(-\lambda)] + T_1(-\lambda e^{-\lambda x})\sin(\omega t - \lambda x) = -\lambda T_1 e^{-\lambda x}\,[\sin(\omega t - \lambda x) + \cos(\omega t - \lambda x)]$. This quantity represents the rate of change of temperature with respect to depth below the surface, at a given time t.

(b) $\partial T/\partial t = T_1 e^{-\lambda x}\,[\cos(\omega t - \lambda x)(\omega)] = \omega T_1 e^{-\lambda x}\cos(\omega t - \lambda x)$. This quantity represents the rate of change of temperature with respect to time at a fixed depth x.

(c) $T_{xx} = \dfrac{\partial}{\partial x}\left(\dfrac{\partial T}{\partial x}\right)$

$$= -\lambda T_1\left(e^{-\lambda x}\,[\cos(\omega t - \lambda x)(-\lambda) - \sin(\omega t - \lambda x)(-\lambda)] + e^{-\lambda x}(-\lambda)\,[\sin(\omega t - \lambda x) + \cos(\omega t - \lambda x)]\right)$$

$$= 2\lambda^2 T_1 e^{-\lambda x}\cos(\omega t - \lambda x)$$

But from part (b), $T_t = \omega T_1 e^{-\lambda x}\cos(\omega t - \lambda x) = \dfrac{\omega}{2\lambda^2} T_{xx}$. So with $k = \dfrac{\omega}{2\lambda^2}$, the function T satisfies the heat equation.

(d)



Note that near the surface (that is, for small x) the temperature varies greatly as t changes, but deeper (for large x) the temperature is more stable.

(e) The term $-\lambda x$ is a phase shift: it represents the fact that since heat diffuses slowly through soil, it takes time for changes in the surface temperature to affect the temperature at deeper points. As x increases, the phase shift also increases. For example, when $\lambda = 0.2$, the highest temperature at the surface is reached when $t \approx 91$, whereas at a depth of 5 feet the peak temperature is attained at $t \approx 149$, and at a depth of 10 feet, at $t \approx 207$.

101. By Clairaut's Theorem, $f_{xyy} = (f_{xy})_y = (f_{yx})_y = f_{yxy} = (f_y)_{xy} = (f_y)_{yx} = f_{yyx}$.

102. (a) Since we are differentiating n times, with two choices of variable at each differentiation, there are 2^n nth-order partial derivatives.

(b) If these partial derivatives are all continuous, then the order in which the partials are taken doesn't affect the value of the result, that is, all nth-order partial derivatives with p partials with respect to x and $n - p$ partials with respect to y are equal. Since the number of partials taken with respect to x for an nth-order partial derivative can range from 0 to n, a function of two variables has $n + 1$ distinct partial derivatives of order n if these partial derivatives are all continuous.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) Since n differentiations are to be performed with three choices of variable at each differentiation, there are 3^n nth-order partial derivatives of a function of three variables.

103. Let $g(x) = f(x,0) = x(x^2)^{-3/2}e^0 = x\,|x|^{-3}$. But we are using the point $(1,0)$, so near $(1,0)$, $g(x) = x^{-2}$. Then $g'(x) = -2x^{-3}$ and $g'(1) = -2$, so using (1) we have $f_x(1,0) = g'(1) = -2$.

104. $f_x(0,0) = \lim\limits_{h\to 0} \dfrac{f(0+h,0) - f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{(h^3+0)^{1/3} - 0}{h} = \lim\limits_{h\to 0} \dfrac{h}{h} = 1.$

Or: Let $g(x) = f(x,0) = \sqrt[3]{x^3+0} = x$. Then $g'(x) = 1$ and $g'(0) = 1$ so, by (1), $f_x(0,0) = g'(0) = 1$.

105. (a)



(b) For $(x,y) \neq (0,0)$,

$$f_x(x,y) = \frac{(3x^2y - y^3)(x^2+y^2) - (x^3y - xy^3)(2x)}{(x^2+y^2)^2} = \frac{x^4y + 4x^2y^3 - y^5}{(x^2+y^2)^2}$$

and by symmetry $f_y(x,y) = \dfrac{x^5 - 4x^3y^2 - xy^4}{(x^2+y^2)^2}$.

(c) $f_x(0,0) = \lim\limits_{h\to 0} \dfrac{f(h,0) - f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{(0/h^2) - 0}{h} = 0$ and $f_y(0,0) = \lim\limits_{h\to 0} \dfrac{f(0,h) - f(0,0)}{h} = 0.$

(d) By (3), $f_{xy}(0,0) = \dfrac{\partial f_x}{\partial y} = \lim\limits_{h\to 0} \dfrac{f_x(0,h) - f_x(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{(-h^5 - 0)/h^4}{h} = -1$ while by (2),

$$f_{yx}(0,0) = \frac{\partial f_y}{\partial x} = \lim_{h\to 0} \frac{f_y(h,0) - f_y(0,0)}{h} = \lim_{h\to 0} \frac{h^5/h^4}{h} = 1.$$

(e) For $(x,y) \neq (0,0)$, we use a CAS to compute

$$f_{xy}(x,y) = \frac{x^6 + 9x^4y^2 - 9x^2y^4 - y^6}{(x^2+y^2)^3}$$

Now as $(x,y) \to (0,0)$ along the x-axis, $f_{xy}(x,y) \to 1$ while as $(x,y) \to (0,0)$ along the y-axis, $f_{xy}(x,y) \to -1$. Thus f_{xy} isn't continuous at $(0,0)$ and Clairaut's Theorem doesn't apply, so there is no contradiction. The graphs of f_{xy} and f_{yx} are identical except at the origin, where we observe the discontinuity.



14.4 Tangent Planes and Linear Approximations

1. $z = f(x,y) = 2x^2 + y^2 - 5y \;\Rightarrow\; f_x(x,y) = 4x,\ f_y(x,y) = 2y - 5,$ so $f_x(1,2) = 4,\ f_y(1,2) = -1.$

By Equation 2, an equation of the tangent plane is $z - (-4) = f_x(1,2)(x-1) + f_y(1,2)(y-2) \;\Rightarrow$

$z + 4 = 4(x-1) + (-1)(y-2)$ or $z = 4x - y - 6$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

2. $z = f(x,y) = (x+2)^2 - 2(y-1)^2 - 5 \Rightarrow f_x(x,y) = 2(x+2),\ f_y(x,y) = -4(y-1)$, so $f_x(2,3) = 8$ and $f_y(2,3) = -8$. By Equation 2, an equation of the tangent plane is $z - 3 = f_x(2,3)(x-2) + f_y(2,3)(y-3) \Rightarrow z - 3 = 8(x-2) + (-8)(y-3)$ or $z = 8x - 8y + 11$.

3. $z = f(x,y) = e^{x-y} \Rightarrow f_x(x,y) = e^{x-y}(1) = e^{x-y},\ f_y(x,y) = e^{x-y}(-1) = -e^{x-y}$, so $f_x(2,2) = 1$ and $f_y(2,2) = -1$. Thus an equation of the tangent plane is $z - 1 = f_x(2,2)(x-2) + f_y(2,2)(y-2) \Rightarrow z - 1 = 1(x-2) + (-1)(y-2)$ or $z = x - y + 1$.

4. $z = f(x,y) = x/y^2 = xy^{-2} \Rightarrow f_x(x,y) = 1/y^2,\ f_y(x,y) = -2xy^{-3} = -2x/y^3$, so $f_x(-4,2) = \frac{1}{4}$ and $f_y(-4,2) = 1$. Thus an equation of the tangent plane is $z - (-1) = f_x(-4,2)\,[x - (-4)] + f_y(-4,2)(y-2) \Rightarrow z + 1 = \frac{1}{4}(x+4) + 1(y-2)$ or $z = \frac{1}{4}x + y - 2$.

5. $z = f(x,y) = x\sin(x+y) \Rightarrow f_x(x,y) = x \cdot \cos(x+y) + \sin(x+y) \cdot 1 = x\cos(x+y) + \sin(x+y)$, $f_y(x,y) = x\cos(x+y)$, so $f_x(-1,1) = (-1)\cos 0 + \sin 0 = -1,\ f_y(-1,1) = (-1)\cos 0 = -1$ and an equation of the tangent plane is $z - 0 = (-1)(x+1) + (-1)(y-1)$ or $x + y + z = 0$.

6. $z = f(x,y) = \ln(x - 2y) \Rightarrow f_x(x,y) = 1/(x-2y),\ f_y(x,y) = -2/(x-2y)$, so $f_x(3,1) = 1,\ f_y(3,1) = -2$, and an equation of the tangent plane is $z - 0 = f_x(3,1)(x-3) + f_y(3,1)(y-1) \Rightarrow z = 1(x-3) + (-2)(y-1)$ or $z = x - 2y - 1$.

7. $z = f(x,y) = x^2 + xy + 3y^2$, so $f_x(x,y) = 2x + y \Rightarrow f_x(1,1) = 3,\ f_y(x,y) = x + 6y \Rightarrow f_y(1,1) = 7$ and an equation of the tangent plane is $z - 5 = 3(x-1) + 7(y-1)$ or $z = 3x + 7y - 5$. After zooming in, the surface and the tangent plane become almost indistinguishable. (Here, the tangent plane is below the surface.) If we zoom in farther, the surface and the tangent plane will appear to coincide.





8. $z = f(x,y) = \sqrt{9 + x^2y^2} \Rightarrow f_x(x,y) = \frac{1}{2}\left(9 + x^2y^2\right)^{-1/2}(2xy^2) = xy^2/\sqrt{9 + x^2y^2}$, $f_y(x,y) = \frac{1}{2}\left(9 + x^2y^2\right)^{-1/2}(2x^2y) = x^2y/\sqrt{9 + x^2y^2}$, so $f_x(2,2) = \frac{8}{5}$ and $f_y(2,2) = \frac{8}{5}$. Thus an equation of the tangent plane is $z - 5 = f_x(2,2)(x-2) + f_y(2,2)(y-2) \Rightarrow z - 5 = \frac{8}{5}(x-2) + \frac{8}{5}(y-2)$ or $z = \frac{8}{5}x + \frac{8}{5}y - \frac{7}{5}$. After zooming in, the surface and the tangent plane become almost indistinguishable. (Here the tangent plane is shown with

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

fewer traces than the surface.) If we zoom in farther, the surface and the tangent plane will appear to coincide.





9. $f(x,y) = \dfrac{1+\cos^2(x-y)}{1+\cos^2(x+y)}$. A CAS gives

$$f_x(x,y) = -\frac{2\cos(x-y)\sin(x-y)}{1+\cos^2(x+y)} + \frac{2\left[1+\cos^2(x-y)\right]\cos(x+y)\sin(x+y)}{\left[1+\cos^2(x+y)\right]^2} \quad \text{and}$$

$$f_y(x,y) = \frac{2\cos(x-y)\sin(x-y)}{1+\cos^2(x+y)} + \frac{2\left[1+\cos^2(x-y)\right]\cos(x+y)\sin(x+y)}{\left[1+\cos^2(x+y)\right]^2}.$$ We use the CAS to evaluate these at $(\pi/3, \pi/6)$, giving $f_x(\pi/3,\pi/6) = -\sqrt{3}/2$ and $f_y(\pi/3,\pi/6) = \sqrt{3}/2$. Substituting into Equation 2, an equation of the tangent plane is $z = -\frac{\sqrt{3}}{2}\left(x-\frac{\pi}{3}\right) + \frac{\sqrt{3}}{2}\left(y-\frac{\pi}{6}\right) + \frac{7}{4}$. The surface and tangent plane are shown in the first graph below. After zooming in, the surface and the tangent plane become almost indistinguishable, as shown in the second graph. (Here, the tangent plane is above the surface.) If we zoom in farther, the surface and the tangent plane will appear to coincide.





10. $f(x,y) = e^{-xy/10}\left(\sqrt{x}+\sqrt{y}+\sqrt{xy}\right)$. A CAS gives

$$f_x(x,y) = -\tfrac{1}{10}ye^{-xy/10}\left(\sqrt{x}+\sqrt{y}+\sqrt{xy}\right) + e^{-xy/10}\left(\frac{1}{2\sqrt{x}}+\frac{y}{2\sqrt{xy}}\right) \quad \text{and}$$

$$f_y(x,y) = -\tfrac{1}{10}xe^{-xy/10}\left(\sqrt{x}+\sqrt{y}+\sqrt{xy}\right) + e^{-xy/10}\left(\frac{1}{2\sqrt{y}}+\frac{x}{2\sqrt{xy}}\right).$$ We use the CAS to evaluate these at $(1,1)$, and then substitute the results into Equation 2 to get an equation of the tangent plane: $z = 0.7e^{-0.1}x + 0.7e^{-0.1}y + 1.6e^{-0.1}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

The surface and tangent plane are shown in the first graph below. After zooming in, the surface and the tangent plane become almost indistinguishable, as shown in the second graph. (Here, the tangent plane is above the surface.) If we zoom in farther, the surface and the tangent plane will appear to coincide.





11. $f(x,y) = 1 + x\ln(xy-5)$. The partial derivatives are $f_x(x,y) = x \cdot \dfrac{1}{xy-5}\,(y) + \ln(xy-5)\cdot 1 = \dfrac{xy}{xy-5} + \ln(xy-5)$ and $f_y(x,y) = x \cdot \dfrac{1}{xy-5}\,(x) = \dfrac{x^2}{xy-5}$, so $f_x(2,3) = 6$ and $f_y(2,3) = 4$. Both f_x and f_y are continuous functions for $xy > 5$, so by Theorem 8, f is differentiable at $(2,3)$. By Equation 3, the linearization of f at $(2,3)$ is given by $L(x,y) = f(2,3) + f_x(2,3)(x-2) + f_y(2,3)(y-3) = 1 + 6(x-2) + 4(y-3) = 6x + 4y - 23$.

12. $f(x,y) = \sqrt{xy} = (xy)^{1/2}$. The partial derivatives are $f_x(x,y) = \frac{1}{2}(xy)^{-1/2}(y) = y/\left(2\sqrt{xy}\right)$ and $f_y(x,y) = \frac{1}{2}(xy)^{-1/2}(x) = x/\left(2\sqrt{xy}\right)$, so $f_x(1,4) = 4/\left(2\sqrt{4}\right) = 1$ and $f_y(1,4) = 1/\left(2\sqrt{4}\right) = \frac{1}{4}$. Both f_x and f_y are continuous functions for $xy > 0$, so f is differentiable at $(1,4)$ by Theorem 8. The linearization of f at $(1,4)$ is $L(x,y) = f(1,4) + f_x(1,4)(x-1) + f_y(1,4)(y-4) = 2 + 1(x-1) + \frac{1}{4}(y-4) = x + \frac{1}{4}y$.

13. $f(x,y) = x^2e^y$. The partial derivatives are $f_x(x,y) = 2xe^y$ and $f_y(x,y) = x^2e^y$, so $f_x(1,0) = 2$ and $f_y(1,0) = 1$. Both f_x and f_y are continuous functions, so by Theorem 8, f is differentiable at $(1,0)$. By Equation 3, the linearization of f at $(1,0)$ is given by $L(x,y) = f(1,0) + f_x(1,0)(x-1) + f_y(1,0)(y-0) = 1 + 2(x-1) + 1(y-0) = 2x + y - 1$.

14. $f(x,y) = \dfrac{1+y}{1+x} = (1+y)(1+x)^{-1}$. The partial derivatives are $f_x(x,y) = (1+y)(-1)(1+x)^{-2} = -\dfrac{1+y}{(1+x)^2}$ and $f_y(x,y) = (1)(1+x)^{-1} = \dfrac{1}{1+x}$, so $f_x(1,3) = -1$ and $f_y(1,3) = \frac{1}{2}$. Both f_x and f_y are continuous functions for $x \neq -1$, so f is differentiable at $(1,3)$ by Theorem 8. The linearization of f at $(1,3)$ is $L(x,y) = f(1,3) + f_x(1,3)(x-1) + f_y(1,3)(y-3) = 2 + (-1)(x-1) + \frac{1}{2}(y-3) = -x + \frac{1}{2}y + \frac{3}{2}$.

15. $f(x,y) = 4\arctan(xy)$. The partial derivatives are $f_x(x,y) = 4 \cdot \dfrac{1}{1+(xy)^2}(y) = \dfrac{4y}{1+x^2y^2}$, and $f_y(x,y) = \dfrac{4x}{1+x^2y^2}$, so $f_x(1,1) = 2$ and $f_y(1,1) = 2$. Both f_x and f_y are continuous functions, so f is differentiable at $(1,1)$ by Theorem 8. The linearization of f at $(1,1)$ is $L(x,y) = f(1,1) + f_x(1,1)(x-1) + f_y(1,1)(y-1) = 4(\pi/4) + 2(x-1) + 2(y-1) = 2x + 2y + \pi - 4$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. $f(x,y) = y + \sin(x/y)$. The partial derivatives are $f_x(x,y) = (1/y)\cos(x/y)$ and $f_y(x,y) = 1 + (-x/y^2)\cos(x/y)$, so $f_x(0,3) = \frac{1}{3}$ and $f_y(0,3) = 1$. Both f_x and f_y are continuous functions for $y \neq 0$, so f is differentiable at $(0,3)$, and the linearization of f at $(0,3)$ is

$$L(x,y) = f(0,3) + f_x(0,3)(x-0) + f_y(0,3)(y-3) = 3 + \tfrac{1}{3}(x-0) + 1(y-3) = \tfrac{1}{3}x + y.$$

17. Let $f(x,y) = e^x \cos(xy)$. Then $f_x(x,y) = e^x[-\sin(xy)](y) + e^x \cos(xy) = e^x[\cos(xy) - y\sin(xy)]$ and $f_y(x,y) = e^x[-\sin(xy)](x) = -xe^x \sin(xy)$. Both f_x and f_y are continuous functions, so by Theorem 8, f is differentiable at $(0,0)$. We have $f_x(0,0) = e^0(\cos 0 - 0) = 1$, $f_y(0,0) = 0$ and the linear approximation of f at $(0,0)$ is $f(x,y) \approx f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) = 1 + 1x + 0y = x + 1$.

18. Let $f(x,y) = \dfrac{y-1}{x+1}$. Then $f_x(x,y) = (y-1)(-1)(x+1)^{-2} = \dfrac{1-y}{(x+1)^2}$ and $f_y(x,y) = \dfrac{1}{x+1}$. Both f_x and f_y are continuous functions for $x \neq -1$, so by Theorem 8, f is differentiable at $(0,0)$. We have $f_x(0,0) = 1$, $f_y(0,0) = 1$ and the linear approximation of f at $(0,0)$ is $f(x,y) \approx f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) = -1 + 1x + 1y = x + y - 1$.

19. We can estimate $f(2.2, 4.9)$ using a linear approximation of f at $(2,5)$, given by $f(x,y) \approx f(2,5) + f_x(2,5)(x-2) + f_y(2,5)(y-5) = 6 + 1(x-2) + (-1)(y-5) = x - y + 9$. Thus $f(2.2, 4.9) \approx 2.2 - 4.9 + 9 = 6.3$.

20. $f(x,y) = 1 - xy\cos\pi y \;\Rightarrow\; f_x(x,y) = -y\cos\pi y$ and $f_y(x,y) = -x[y(-\pi\sin\pi y) + (\cos\pi y)(1)] = \pi xy\sin\pi y - x\cos\pi y$, so $f_x(1,1) = 1$, $f_y(1,1) = 1$. Then the linear approximation of f at $(1,1)$ is given by

$$\begin{aligned} f(x,y) &\approx f(1,1) + f_x(1,1)(x-1) + f_y(1,1)(y-1) \\ &= 2 + (1)(x-1) + (1)(y-1) = x + y \end{aligned}$$

Thus $f(1.02, 0.97) \approx 1.02 + 0.97 = 1.99$. We graph f and its tangent plane near the point $(1,1,2)$ below. Notice near $y = 1$ the surfaces are almost identical.



21. $f(x,y,z) = \sqrt{x^2+y^2+z^2} \;\Rightarrow\; f_x(x,y,z) = \dfrac{x}{\sqrt{x^2+y^2+z^2}}$, $f_y(x,y,z) = \dfrac{y}{\sqrt{x^2+y^2+z^2}}$, and $f_z(x,y,z) = \dfrac{z}{\sqrt{x^2+y^2+z^2}}$, so $f_x(3,2,6) = \frac{3}{7}$, $f_y(3,2,6) = \frac{2}{7}$, $f_z(3,2,6) = \frac{6}{7}$. Then the linear approximation of f

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

at $(3, 2, 6)$ is given by

$$f(x, y, z) \approx f(3, 2, 6) + f_x(3, 2, 6)(x - 3) + f_y(3, 2, 6)(y - 2) + f_z(3, 2, 6)(z - 6)$$
$$= 7 + \tfrac{3}{7}(x - 3) + \tfrac{2}{7}(y - 2) + \tfrac{6}{7}(z - 6) = \tfrac{3}{7}x + \tfrac{2}{7}y + \tfrac{6}{7}z$$

Thus $\sqrt{(3.02)^2 + (1.97)^2 + (5.99)^2} = f(3.02, 1.97, 5.99) \approx \frac{3}{7}(3.02) + \frac{2}{7}(1.97) + \frac{6}{7}(5.99) \approx 6.9914$.

22. From the table, $f(40, 20) = 28$. To estimate $f_v(40, 20)$ and $f_t(40, 20)$ we follow the procedure used in Exercise 14.3.4. Since $f_v(40, 20) = \lim_{h \to 0} \dfrac{f(40 + h, 20) - f(40, 20)}{h}$, we approximate this quantity with $h = \pm 10$ and use the values given in the table:

$$f_v(40, 20) \approx \frac{f(50, 20) - f(40, 20)}{10} = \frac{40 - 28}{10} = 1.2, \qquad f_v(40, 20) \approx \frac{f(30, 20) - f(40, 20)}{-10} = \frac{17 - 28}{-10} = 1.1$$

Averaging these values gives $f_v(40, 20) \approx 1.15$. Similarly, $f_t(40, 20) = \lim_{h \to 0} \dfrac{f(40, 20 + h) - f(40, 20)}{h}$, so we use $h = 10$ and $h = -5$:

$$f_t(40, 20) \approx \frac{f(40, 30) - f(40, 20)}{10} = \frac{31 - 28}{10} = 0.3, \qquad f_t(40, 20) \approx \frac{f(40, 15) - f(40, 20)}{-5} = \frac{25 - 28}{-5} = 0.6$$

Averaging these values gives $f_t(40, 15) \approx 0.45$. The linear approximation, then, is

$$f(v, t) \approx f(40, 20) + f_v(40, 20)(v - 40) + f_t(40, 20)(t - 20) \approx 28 + 1.15(v - 40) + 0.45(t - 20)$$

When $v = 43$ and $t = 24$, we estimate $f(43, 24) \approx 28 + 1.15(43 - 40) + 0.45(24 - 20) = 33.25$, so we would expect the wave heights to be approximately 33.25 ft.

23. From the table, $f(94, 80) = 127$. To estimate $f_T(94, 80)$ and $f_H(94, 80)$ we follow the procedure used in Section 14.3. Since $f_T(94, 80) = \lim_{h \to 0} \dfrac{f(94 + h, 80) - f(94, 80)}{h}$, we approximate this quantity with $h = \pm 2$ and use the values given in the table:

$$f_T(94, 80) \approx \frac{f(96, 80) - f(94, 80)}{2} = \frac{135 - 127}{2} = 4, \qquad f_T(94, 80) \approx \frac{f(92, 80) - f(94, 80)}{-2} = \frac{119 - 127}{-2} = 4$$

Averaging these values gives $f_T(94, 80) \approx 4$. Similarly, $f_H(94, 80) = \lim_{h \to 0} \dfrac{f(94, 80 + h) - f(94, 80)}{h}$, so we use $h = \pm 5$:

$$f_H(94, 80) \approx \frac{f(94, 85) - f(94, 80)}{5} = \frac{132 - 127}{5} = 1, \qquad f_H(94, 80) \approx \frac{f(94, 75) - f(94, 80)}{-5} = \frac{122 - 127}{-5} = 1$$

Averaging these values gives $f_H(94, 80) \approx 1$. The linear approximation, then, is

$$f(T, H) \approx f(94, 80) + f_T(94, 80)(T - 94) + f_H(94, 80)(H - 80)$$
$$\approx 127 + 4(T - 94) + 1(H - 80) \qquad [\text{or } 4T + H - 329]$$

Thus when $T = 95$ and $H = 78$, $f(95, 78) \approx 127 + 4(95 - 94) + 1(78 - 80) = 129$, so we estimate the heat index to be approximately 129°F.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. From the table, $f(-15,50) = -29$. To estimate $f_T(-15,50)$ and $f_v(-15,50)$ we follow the procedure used in Section 14.3.

Since $f_T(-15,50) = \lim_{h\to 0} \dfrac{f(-15+h,50) - f(-15,50)}{h}$, we approximate this quantity with $h = \pm 5$ and use the values given in the table:

$$f_T(-15,50) \approx \frac{f(-10,50) - f(-15,50)}{5} = \frac{-22-(-29)}{5} = 1.4$$

$$f_T(-15,50) \approx \frac{f(-20,50) - f(-15,50)}{-5} = \frac{-35-(-29)}{-5} = 1.2$$

Averaging these values gives $f_T(-15,50) \approx 1.3$. Similarly $f_v(-15,50) = \lim_{h\to 0} \dfrac{f(-15,50+h) - f(-15,50)}{h}$, so we use $h = \pm 10$:

$$f_v(-15,50) \approx \frac{f(-15,60) - f(-15,50)}{10} = \frac{-30-(-29)}{10} = -0.1$$

$$f_v(-15,50) \approx \frac{f(-15,40) - f(-15,50)}{-10} = \frac{-27-(-29)}{-10} = -0.2$$

Averaging these values gives $f_v(-15,50) \approx -0.15$. The linear approximation to the wind-chill index function, then, is $f(T,v) \approx f(-15,50) + f_T(-15,50)(T-(-15)) + f_v(-15,50)(v-50) \approx -29 + (1.3)(T+15) - (0.15)(v-50)$. Thus when $T = -17^\circ\text{C}$ and $v = 55$ km/h, $f(-17,55) \approx -29 + (1.3)(-17+15) - (0.15)(55-50) = -32.35$, so we estimate the wind-chill index to be approximately -32.35°C.

25. $z = e^{-2x}\cos 2\pi t \;\Rightarrow$

$$dz = \frac{\partial z}{\partial x}dx + \frac{\partial z}{\partial t}dt = e^{-2x}(-2)\cos 2\pi t\,dx + e^{-2x}(-\sin 2\pi t)(2\pi)\,dt = -2e^{-2x}\cos 2\pi t\,dx - 2\pi e^{-2x}\sin 2\pi t\,dt$$

26. $u = \sqrt{x^2+3y^2} = (x^2+3y^2)^{1/2} \;\Rightarrow$

$$du = \frac{\partial u}{\partial x}dx + \frac{\partial u}{\partial y}dy = \tfrac{1}{2}(x^2+3y^2)^{-1/2}(2x)\,dx + \tfrac{1}{2}(x^2+3y^2)^{-1/2}(6y)\,dy = \frac{x}{\sqrt{x^2+3y^2}}\,dx + \frac{3y}{\sqrt{x^2+3y^2}}\,dy$$

27. $m = p^5q^3 \;\Rightarrow\; dm = \dfrac{\partial m}{\partial p}\,dp + \dfrac{\partial m}{\partial q}\,dq = 5p^4q^3\,dp + 3p^5q^2\,dq$

28. $T = \dfrac{v}{1+uvw} \;\Rightarrow$

$$\begin{aligned} dT &= \frac{\partial T}{\partial u}\,du + \frac{\partial T}{\partial v}\,dv + \frac{\partial T}{\partial w}\,dw \\ &= v(-1)(1+uvw)^{-2}(vw)\,du + \frac{1(1+uvw)-v(uw)}{(1+uvw)^2}\,dv + v(-1)(1+uvw)^{-2}(uv)\,dw \\ &= -\frac{v^2w}{(1+uvw)^2}\,du + \frac{1}{(1+uvw)^2}\,dv - \frac{uv^2}{(1+uvw)^2}\,dw \end{aligned}$$

29. $R = \alpha\beta^2\cos\gamma \;\Rightarrow\; dR = \dfrac{\partial R}{\partial \alpha}\,d\alpha + \dfrac{\partial R}{\partial \beta}\,d\beta + \dfrac{\partial R}{\partial \gamma}\,d\gamma = \beta^2\cos\gamma\,d\alpha + 2\alpha\beta\cos\gamma\,d\beta - \alpha\beta^2\sin\gamma\,d\gamma$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

30. $L = xze^{y^2-z^2} \Rightarrow$

$$dL = \frac{\partial L}{\partial x}\,dx + \frac{\partial L}{\partial y}\,dy + \frac{\partial L}{\partial z}\,dz = ze^{-y^2-z^2}\,dx + xze^{-y^2-z^2}(-2y)\,dy + x[z\cdot e^{-y^2-z^2}(-2z) + e^{-y^2-z^2}\cdot 1]\,dz$$

$$= ze^{-y^2-z^2}\,dx - 2xyze^{-y^2-z^2}\,dy + x(1-2z^2)e^{-y^2-z^2}\,dz$$

31. $dx = \Delta x = 0.05$, $dy = \Delta y = 0.1$, $z = 5x^2 + y^2$, $z_x = 10x$, $z_y = 2y$. Thus when $x = 1$ and $y = 2$,

$dz = z_x(1,2)\,dx + z_y(1,2)\,dy = (10)(0.05) + (4)(0.1) = 0.9$ while

$\Delta z = f(1.05, 2.1) - f(1,2) = 5(1.05)^2 + (2.1)^2 - 5 - 4 = 0.9225$.

32. $dx = \Delta x = -0.04$, $dy = \Delta y = 0.05$, $z = x^2 - xy + 3y^2$, $z_x = 2x - y$, $z_y = 6y - x$. Thus when $x = 3$ and $y = -1$,

$dz = (7)(-0.04) + (-9)(0.05) = -0.73$ while $\Delta z = (2.96)^2 - (2.96)(-0.95) + 3(-0.95)^2 - (9 + 3 + 3) = -0.7189$.

33. $dA = \dfrac{\partial A}{\partial x}\,dx + \dfrac{\partial A}{\partial y}\,dy = y\,dx + x\,dy$ and $|\Delta x| \le 0.1$, $|\Delta y| \le 0.1$. We use $dx = 0.1$, $dy = 0.1$ with $x = 30$, $y = 24$; then the maximum error in the area is about $dA = 24(0.1) + 30(0.1) = 5.4$ cm^2.

34. Let V be the volume. Then $V = \pi r^2 h$ and $\Delta V \approx dV = 2\pi rh\,dr + \pi r^2\,dh$ is an estimate of the amount of metal. With $dr = 0.05$ and $dh = 0.2$ we get $dV = 2\pi(2)(10)(0.05) + \pi(2)^2(0.2) = 2.80\pi \approx 8.8$ cm^3.

35. The volume of a can is $V = \pi r^2 h$ and $\Delta V \approx dV$ is an estimate of the amount of tin. Here $dV = 2\pi rh\,dr + \pi r^2\,dh$, so put $dr = 0.04$, $dh = 0.08$ (0.04 on top, 0.04 on bottom) and then $\Delta V \approx dV = 2\pi(48)(0.04) + \pi(16)(0.08) \approx 16.08$ cm^3. Thus the amount of tin is about 16 cm^3.

36. $W = 13.12 + 0.6215T - 11.37v^{0.16} + 0.3965Tv^{0.16}$, so the differential of W is

$$dW = \frac{\partial W}{\partial T}\,dT + \frac{\partial W}{\partial v}\,dv = (0.6215 + 0.3965v^{0.16})\,dT + \left[-11.37(0.16)v^{-0.84} + 0.3965T(0.16)v^{-0.84}\right]dv$$

$$= (0.6215 + 0.3965v^{0.16})\,dT + (-1.8192 + 0.06344T)v^{-0.84}\,dv$$

Here we have $|\Delta T| \le 1$, $|\Delta v| \le 2$, so we take $dT = 1$, $dv = 2$ with $T = -11$, $v = 26$. The maximum error in the calculated value of W is about $dW = (0.6215 + 0.3965(26)^{0.16})(1) + (-1.8192 + 0.06344(-11))(26)^{-0.84}(2) \approx 0.96$.

37. $T = \dfrac{mgR}{2r^2 + R^2}$, so the differential of T is

$$dT = \frac{\partial T}{\partial R}\,dR + \frac{\partial T}{\partial r}\,dr = \frac{(2r^2 + R^2)(mg) - mgR(2R)}{(2r^2 + R^2)^2}\,dR + \frac{(2r^2 + R^2)(0) - mgR(4r)}{(2r^2 + R^2)^2}\,dr$$

$$= \frac{mg(2r^2 - R^2)}{(2r^2 + R^2)^2}\,dR - \frac{4mgRr}{(2r^2 + R^2)^2}\,dr$$

Here we have $\Delta R = 0.1$ and $\Delta r = 0.1$, so we take $dR = 0.1$, $dr = 0.1$ with $R = 3$, $r = 0.7$. Then the change in the tension T is approximately

$$dT = \frac{mg[2(0.7)^2 - (3)^2]}{[2(0.7)^2 + (3)^2]^2}(0.1) - \frac{4mg(3)(0.7)}{[2(0.7)^2 + (3)^2]^2}(0.1)$$

$$= -\frac{0.802mg}{(9.98)^2} - \frac{0.84mg}{(9.98)^2} = -\frac{1.642}{99.6004}mg \approx -0.0165mg$$

Because the change is negative, tension decreases.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. Here $dV = \Delta V = 0.3$, $dT = \Delta T = -5$, $P = 8.31\dfrac{T}{V}$, so

$dP = \left(\dfrac{8.31}{V}\right)dT - \dfrac{8.31 \cdot T}{V^2}\,dV = 8.31\left[-\dfrac{5}{12} - \dfrac{310}{144}\cdot\dfrac{3}{10}\right] \approx -8.83$. Thus the pressure will drop by about 8.83 kPa.

39. First we find $\dfrac{\partial R}{\partial R_1}$ implicitly by taking partial derivatives of both sides with respect to R_1:

$\dfrac{\partial}{\partial R_1}\left(\dfrac{1}{R}\right) = \dfrac{\partial\left[(1/R_1)+(1/R_2)+(1/R_3)\right]}{\partial R_1} \;\Rightarrow\; -R^{-2}\dfrac{\partial R}{\partial R_1} = -R_1^{-2} \;\Rightarrow\; \dfrac{\partial R}{\partial R_1} = \dfrac{R^2}{R_1^2}$. Then by symmetry,

$\dfrac{\partial R}{\partial R_2} = \dfrac{R^2}{R_2^2}$, $\dfrac{\partial R}{\partial R_3} = \dfrac{R^2}{R_3^2}$. When $R_1 = 25$, $R_2 = 40$ and $R_3 = 50$, $\dfrac{1}{R} = \dfrac{17}{200} \;\Leftrightarrow\; R = \frac{200}{17}\ \Omega$. Since the possible error for each R_i is 0.5%, the maximum error of R is attained by setting $\Delta R_i = 0.005R_i$. So

$\Delta R \approx dR = \dfrac{\partial R}{\partial R_1}\Delta R_1 + \dfrac{\partial R}{\partial R_2}\Delta R_2 + \dfrac{\partial R}{\partial R_3}\Delta R_3 = (0.005)R^2\left(\dfrac{1}{R_1}+\dfrac{1}{R_2}+\dfrac{1}{R_3}\right) = (0.005)R = \frac{1}{17} \approx 0.059\ \Omega.$

40. The errors in measurement are at most 2%, so $\left|\dfrac{\Delta w}{w}\right| \le 0.02$ and $\left|\dfrac{\Delta h}{h}\right| \le 0.02$. The relative error in the calculated surface area is

$$\frac{\Delta S}{S} \approx \frac{dS}{S} = \frac{0.1091(0.425w^{0.425-1})h^{0.725}\,dw + 0.1091w^{0.425}(0.725h^{0.725-1})\,dh}{0.1091w^{0.425}h^{0.725}} = 0.425\frac{dw}{w} + 0.725\frac{dh}{h}$$

To estimate the maximum relative error, we use $\dfrac{dw}{w} = \left|\dfrac{\Delta w}{w}\right| = 0.02$ and $\dfrac{dh}{h} = \left|\dfrac{\Delta h}{h}\right| = 0.02 \;\Rightarrow$

$\dfrac{dS}{S} = 0.425\,(0.02) + 0.725\,(0.02) = 0.023$. Thus the maximum percentage error is approximately 2.3%.

41. (a) $B(m,h) = m/h^2 \;\Rightarrow\; B_m(m,h) = 1/h^2$ and $B_h(m,h) = -2m/h^3$. Since $h > 0$, both B_m and B_h are continuous functions, so B is differentiable at $(23, 1.10)$. We have $B(23,1.10) = 23/(1.10)^2 \approx 19.01$, $B_m(23,1.10) = 1/(1.10)^2 \approx 0.8264$, and $B_h(23,1.10) = -2(23)/(1.10)^3 \approx -34.56$, so the linear approximation of B at $(23,1.10)$ is

$B(m,h) \approx B(23,1.10) + B_m(23,1.10)(m-23) + B_h(23,1.10)(h-1.10) \approx 19.01 + 0.8264(m-23) - 34.56(h-1.10)$

or $B(m,h) \approx 0.8264m - 34.56h + 38.02$.

(b) From part (a), for values near $m = 23$ and $h = 1.10$, $B(m,h) \approx 0.8264m - 34.56h + 38.02$. If m increases by 1 kg to 24 kg and h increases by 0.03 m to 1.13 m, we estimate the BMI to be $B(24,1.13) \approx 0.8264(24) - 34.56(1.13) + 38.02 \approx 18.801$. This is very close to the actual computed BMI: $B(24,1.13) = 24/(1.13)^2 \approx 18.796$.

42. $\mathbf{r}_1(t) = \langle 2+3t, 1-t^2, 3-4t+t^2\rangle \;\Rightarrow\; \mathbf{r}_1'(t) = \langle 3, -2t, -4+2t\rangle$, $\mathbf{r}_2(u) = \langle 1+u^2, 2u^3-1, 2u+1\rangle \;\Rightarrow$ $\mathbf{r}_2'(u) = \langle 2u, 6u^2, 2\rangle$. Both curves pass through P since $\mathbf{r}_1(0) = \mathbf{r}_2(1) = \langle 2,1,3\rangle$, so the tangent vectors $\mathbf{r}_1'(0) = \langle 3,0,-4\rangle$ and $\mathbf{r}_2'(1) = \langle 2,6,2\rangle$ are both parallel to the tangent plane to S at P. A normal vector for the tangent plane is

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$\mathbf{r}_1'(0) \times \mathbf{r}_2'(1) = \langle 3, 0, -4\rangle \times \langle 2, 6, 2\rangle = \langle 24, -14, 18\rangle$, so an equation of the tangent plane is $24(x-2) - 14(y-1) + 18(z-3) = 0$ or $12x - 7y + 9z = 44$.

43. $\Delta z = f(a+\Delta x, b+\Delta y) - f(a,b) = (a+\Delta x)^2 + (b+\Delta y)^2 - (a^2+b^2)$

$= a^2 + 2a\,\Delta x + (\Delta x)^2 + b^2 + 2b\,\Delta y + (\Delta y)^2 - a^2 - b^2 = 2a\,\Delta x + (\Delta x)^2 + 2b\,\Delta y + (\Delta y)^2$

But $f_x(a,b) = 2a$ and $f_y(a,b) = 2b$ and so $\Delta z = f_x(a,b)\,\Delta x + f_y(a,b)\,\Delta y + \Delta x\,\Delta x + \Delta y\,\Delta y$, which is Definition 7 with $\varepsilon_1 = \Delta x$ and $\varepsilon_2 = \Delta y$. Hence f is differentiable.

44. $\Delta z = f(a+\Delta x, b+\Delta y) - f(a,b) = (a+\Delta x)(b+\Delta y) - 5(b+\Delta y)^2 - (ab - 5b^2)$

$= ab + a\,\Delta y + b\,\Delta x + \Delta x\,\Delta y - 5b^2 - 10b\,\Delta y - 5(\Delta y)^2 - ab + 5b^2$

$= (a - 10b)\,\Delta y + b\,\Delta x + \Delta x\,\Delta y - 5\,\Delta y\,\Delta y,$

but $f_x(a,b) = b$ and $f_y(a,b) = a - 10b$ and so $\Delta z = f_x(a,b)\,\Delta x + f_y(a,b)\,\Delta y + \Delta x\,\Delta y - 5\Delta y\,\Delta y$, which is Definition 7 with $\varepsilon_1 = \Delta y$ and $\varepsilon_2 = -5\,\Delta y$. Hence f is differentiable.

45. To show that f is continuous at (a,b) we need to show that $\lim\limits_{(x,y)\to(a,b)} f(x,y) = f(a,b)$ or equivalently $\lim\limits_{(\Delta x,\Delta y)\to(0,0)} f(a+\Delta x, b+\Delta y) = f(a,b)$. Since f is differentiable at (a,b), $f(a+\Delta x, b+\Delta y) - f(a,b) = \Delta z = f_x(a,b)\,\Delta x + f_y(a,b)\,\Delta y + \varepsilon_1\,\Delta x + \varepsilon_2\,\Delta y$, where ϵ_1 and $\epsilon_2 \to 0$ as $(\Delta x, \Delta y) \to (0,0)$. Thus $f(a+\Delta x, b+\Delta y) = f(a,b) + f_x(a,b)\,\Delta x + f_y(a,b)\,\Delta y + \varepsilon_1\,\Delta x + \varepsilon_2\,\Delta y$. Taking the limit of both sides as $(\Delta x, \Delta y) \to (0,0)$ gives $\lim\limits_{(\Delta x,\Delta y)\to(0,0)} f(a+\Delta x, b+\Delta y) = f(a,b)$. Thus f is continuous at (a,b).

46. (a) $\lim\limits_{h\to 0} \dfrac{f(h,0) - f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{0-0}{h} = 0$ and $\lim\limits_{h\to 0} \dfrac{f(0,h) - f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{0-0}{h} = 0$. Thus $f_x(0,0) = f_y(0,0) = 0$.

To show that f isn't differentiable at $(0,0)$ we need only show that f is not continuous at $(0,0)$ and apply Exercise 45. As $(x,y) \to (0,0)$ along the x-axis $f(x,y) = 0/x^2 = 0$ for $x \neq 0$ so $f(x,y) \to 0$ as $(x,y) \to (0,0)$ along the x-axis. But as $(x,y) \to (0,0)$ along the line $y = x$, $f(x,x) = x^2/(2x^2) = \frac{1}{2}$ for $x \neq 0$ so $f(x,y) \to \frac{1}{2}$ as $(x,y) \to (0,0)$ along this line. Thus $\lim\limits_{(x,y)\to(0,0)} f(x,y)$ doesn't exist, so f is discontinuous at $(0,0)$ and thus not differentiable there.

(b) For $(x,y) \neq (0,0)$, $f_x(x,y) = \dfrac{(x^2+y^2)y - xy(2x)}{(x^2+y^2)^2} = \dfrac{y(y^2 - x^2)}{(x^2+y^2)^2}$. If we approach $(0,0)$ along the y-axis, then $f_x(x,y) = f_x(0,y) = \dfrac{y^3}{y^4} = \dfrac{1}{y}$, so $f_x(x,y) \to \pm\infty$ as $(x,y) \to (0,0)$. Thus $\lim\limits_{(x,y)\to(0,0)} f_x(x,y)$ does not exist and $f_x(x,y)$ is not continuous at $(0,0)$. Similarly, $f_y(x,y) = \dfrac{(x^2+y^2)x - xy(2y)}{(x^2+y^2)^2} = \dfrac{x(x^2-y^2)}{(x^2+y^2)^2}$ for $(x,y) \neq (0,0)$, and if we approach $(0,0)$ along the x-axis, then $f_y(x,y) = f_x(x,0) = \dfrac{x^3}{x^4} = \dfrac{1}{x}$. Thus $\lim\limits_{(x,y)\to(0,0)} f_y(x,y)$ does not exist and $f_y(x,y)$ is not continuous at $(0,0)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT The Speedo LZR Racer

1. $v(P,C) = \left(\dfrac{2P}{kC}\right)^{1/3} \Rightarrow$

$$f(x,y) = \frac{v(P+xP, C+yC) - v(P,C)}{v(P,C)} = \frac{v(P+xP, C+yC)}{v(P,C)} - \frac{v(P,C)}{v(P,C)} = \frac{\left(\dfrac{2(P+xP)}{k(C+yC)}\right)^{1/3}}{\left(\dfrac{2P}{kC}\right)^{1/3}} - 1$$

$$= \left(\frac{2P(1+x)}{kC(1+y)} \cdot \frac{kC}{2P}\right)^{1/3} - 1 = \left(\frac{1+x}{1+y}\right)^{1/3} - 1$$

Both power and drag cannot be reduced by more than 100%, but both could be increased by any percentage, so $x \geq -1$ and $y \geq -1$. But f is undefined when $y = -1$, so the domain is $\{(x,y) \mid x \geq -1,\ y > -1\}$.

2. If x and y are small, then we can say they are near zero and we can use a linear approximation to f at $(0,0)$.

We have $f(x,y) = (1+x)^{1/3}(1+y)^{-1/3} - 1$ so the partial derivatives are

$f_x(x,y) = \frac{1}{3}(1+x)^{-2/3}(1+y)^{-1/3} = \dfrac{1}{3(1+x)^{2/3}(1+y)^{1/3}}$ and

$f_y(x,y) = -\frac{1}{3}(1+x)^{1/3}(1+y)^{-4/3} = -\dfrac{(1+x)^{1/3}}{3(1+y)^{4/3}}$. Note that f_x and f_y are continuous functions for $x > -1$, $y > -1$ so f is differentiable at $(0,0)$. Then $f_x(0,0) = \frac{1}{3}$ and $f_y(0,0) = -\frac{1}{3}$, and the linear approximation is $f(x,y) \approx f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) = 0 + \frac{1}{3}(x-0) - \frac{1}{3}(y-0) = \frac{1}{3}x - \frac{1}{3}y$. According to the linear approximation, a small fractional increase in power results in 1/3 that fractional increase in speed, and a small decrease in drag has the same effect.

3. $f_{xx}(x,y) = \dfrac{1}{3(1+y)^{1/3}} \cdot \left(-\frac{2}{3}\right)(1+x)^{-5/3} = -\dfrac{2}{9(1+x)^{5/3}(1+y)^{1/3}}$,

$f_{yy}(x,y) = -\frac{1}{3}(1+x)^{1/3} \cdot \left(-\frac{4}{3}\right)(1+y)^{-7/3} = \dfrac{4(1+x)^{1/3}}{9(1+y)^{7/3}}$. Because $f_x(x,y)$ is positive in the domain of f, an increase in power results in an increase in speed, but $f_{xx}(x,y)$ is negative, so as the fractional power increases, the fractional speed increases at a declining rate. (We can say that in the positive x-direction, f is increasing and concave downward.) Thus the linear approximation gives an overestimate for an increase in power. Since $f_y(x,y)$ is negative, a *decrease* in drag *increases* speed. But $f_{yy}(x,y)$ is positive, so f_y increases as y increases and f_y decreases (f_y becomes larger and larger negative) as y decreases. (In the positive y-direction, f is decreasing and concave upward.) Thus as the fractional drag decreases, the fractional speed increases at an accelerating pace and the linear approximation gives an underestimate of the increase in power. This explains why a decrease in drag is more effective than an increase in power: Reducing drag improves speed at an increasing rate while adding power improves speed at a declining rate.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. The level curves of $f(x,y) = \left(\dfrac{1+x}{1+y}\right)^{1/3} - 1$ are

$$\left(\frac{1+x}{1+y}\right)^{1/3} - 1 = c \quad \Rightarrow \quad \frac{1+x}{1+y} = (1+c)^3 \quad \Rightarrow$$

$$y = \frac{1+x}{(1+c)^3} - 1.$$

From the level curves, we see that increasing x (from 0) by a small amount has a similar effect on the value of f as decreasing y by a small amount. However, for larger changes, a decrease in y gives greater values of f than a similar increase in x.



14.5 The Chain Rule

1. $z = xy^3 - x^2y, \quad x = t^2 + 1, \quad y = t^2 - 1 \quad \Rightarrow$

$$\frac{dz}{dt} = \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt} = (y^3 - 2xy)(2t) + (3xy^2 - x^2)(2t) = 2t(y^3 - 2xy + 3xy^2 - x^2)$$

2. $z = \dfrac{x-y}{x+2y}, \quad x = e^{\pi t}, \quad y = e^{-\pi t} \quad \Rightarrow$

$$\frac{dz}{dt} = \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt} = \frac{(x+2y)(1) - (x-y)(1)}{(x+2y)^2}(\pi e^{\pi t}) + \frac{(x+2y)(-1) - (x-y)(2)}{(x+2y)^2}(-\pi e^{-\pi t})$$

$$= \frac{3y}{(x+2y)^2}(\pi e^{\pi t}) + \frac{-3x}{(x+2y)^2}(-\pi e^{-\pi t}) = \frac{3\pi}{(x+2y)^2}\left(ye^{\pi t} + xe^{-\pi t}\right)$$

3. $z = \sin x \cos y, \quad x = \sqrt{t}, \quad y = 1/t \quad \Rightarrow$

$$\frac{dz}{dt} = \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt} = (\cos x \cos y)\left(\tfrac{1}{2}t^{-1/2}\right) + (-\sin x \sin y)\left(-t^{-2}\right) = \frac{1}{2\sqrt{t}}\cos x \cos y + \frac{1}{t^2}\sin x \sin y$$

4. $z = \sqrt{1+xy}, \quad x = \tan t, \quad y = \arctan t \quad \Rightarrow$

$$\frac{dz}{dt} = \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt} = \tfrac{1}{2}(1+xy)^{-1/2}(y)\cdot \sec^2 t + \tfrac{1}{2}(1+xy)^{-1/2}(x)\cdot \frac{1}{1+t^2}$$

$$= \frac{1}{2\sqrt{1+xy}}\left(y\sec^2 t + \frac{x}{1+t^2}\right)$$

5. $w = xe^{y/z}, \; x = t^2, \; y = 1 - t, \; z = 1 + 2t \quad \Rightarrow$

$$\frac{dw}{dt} = \frac{\partial w}{\partial x}\frac{dx}{dt} + \frac{\partial w}{\partial y}\frac{dy}{dt} + \frac{\partial w}{\partial z}\frac{dz}{dt} = e^{y/z}\cdot 2t + xe^{y/z}\left(\frac{1}{z}\right)\cdot(-1) + xe^{y/z}\left(-\frac{y}{z^2}\right)\cdot 2 = e^{y/z}\left(2t - \frac{x}{z} - \frac{2xy}{z^2}\right)$$

6. $w = \ln\sqrt{x^2+y^2+z^2} = \tfrac{1}{2}\ln(x^2+y^2+z^2), \quad x = \sin t, \quad y = \cos t, \quad z = \tan t \quad \Rightarrow$

$$\frac{dw}{dt} = \frac{\partial w}{\partial x}\frac{dx}{dt} + \frac{\partial w}{\partial y}\frac{dy}{dt} + \frac{\partial w}{\partial z}\frac{dz}{dt} = \frac{1}{2}\cdot\frac{2x}{x^2+y^2+z^2}\cdot\cos t + \frac{1}{2}\cdot\frac{2y}{x^2+y^2+z^2}\cdot(-\sin t) + \frac{1}{2}\cdot\frac{2z}{x^2+y^2+z^2}\cdot\sec^2 t$$

$$= \frac{x\cos t - y\sin t + z\sec^2 t}{x^2+y^2+z^2}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $z = (x-y)^5, \quad x = s^2t, \quad y = st^2 \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = 5(x-y)^4(1)\cdot 2st + 5(x-y)^4(-1)\cdot t^2 = 5(x-y)^4\left(2st - t^2\right)$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = 5(x-y)^4(1)\cdot s^2 + 5(x-y)^4(-1)\cdot 2st = 5(x-y)^4\left(s^2 - 2st\right)$$

8. $z = \tan^{-1}(x^2+y^2), \quad x = s\ln t, \quad y = te^s \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = \frac{2x}{1+(x^2+y^2)^2}\cdot \ln t + \frac{2y}{1+(x^2+y^2)^2}\cdot te^s$$

$$= \frac{2}{1+(x^2+y^2)^2}\left(x\ln t + yte^s\right)$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = \frac{2x}{1+(x^2+y^2)^2}\cdot \frac{s}{t} + \frac{2y}{1+(x^2+y^2)^2}\cdot e^s$$

$$= \frac{2}{1+(x^2+y^2)^2}\left(\frac{xs}{t} + ye^s\right)$$

9. $z = \ln(3x+2y), \quad x = s\sin t, \quad y = t\cos s \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = \frac{3}{3x+2y}(\sin t) + \frac{2}{3x+2y}(-t\sin s) = \frac{3\sin t - 2t\sin s}{3x+2y}$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = \frac{3}{3x+2y}(s\cos t) + \frac{2}{3x+2y}(\cos s) = \frac{3s\cos t + 2\cos s}{3x+2y}$$

10. $z = \sqrt{x}\,e^{xy}, \quad x = 1+st, \quad y = s^2 - t^2 \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = \left(\sqrt{x}\cdot e^{xy}(y) + e^{xy}\cdot \tfrac{1}{2}x^{-1/2}\right)(t) + \sqrt{x}\,e^{xy}(x)\,(2s) = \left(yt\sqrt{x} + \frac{t}{2\sqrt{x}} + 2x^{3/2}s\right)e^{xy}$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = \left(\sqrt{x}\cdot e^{xy}(y) + e^{xy}\cdot \tfrac{1}{2}x^{-1/2}\right)(s) + \sqrt{x}\,e^{xy}(x)\,(-2t) = \left(ys\sqrt{x} + \frac{s}{2\sqrt{x}} - 2x^{3/2}t\right)e^{xy}$$

11. $z = e^r\cos\theta, \ r = st, \ \theta = \sqrt{s^2+t^2} \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial r}\frac{\partial r}{\partial s} + \frac{\partial z}{\partial \theta}\frac{\partial \theta}{\partial s} = e^r\cos\theta\cdot t + e^r(-\sin\theta)\cdot \tfrac{1}{2}(s^2+t^2)^{-1/2}(2s) = te^r\cos\theta - e^r\sin\theta\cdot\frac{s}{\sqrt{s^2+t^2}}$$

$$= e^r\left(t\cos\theta - \frac{s}{\sqrt{s^2+t^2}}\sin\theta\right)$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial r}\frac{\partial r}{\partial t} + \frac{\partial z}{\partial \theta}\frac{\partial \theta}{\partial t} = e^r\cos\theta\cdot s + e^r(-\sin\theta)\cdot \tfrac{1}{2}(s^2+t^2)^{-1/2}(2t) = se^r\cos\theta - e^r\sin\theta\cdot\frac{t}{\sqrt{s^2+t^2}}$$

$$= e^r\left(s\cos\theta - \frac{t}{\sqrt{s^2+t^2}}\sin\theta\right)$$

12. $z = \tan(u/v), \ u = 2s+3t, \ v = 3s-2t \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial u}\frac{\partial u}{\partial s} + \frac{\partial z}{\partial v}\frac{\partial v}{\partial s} = \sec^2(u/v)(1/v)\cdot 2 + \sec^2(u/v)(-uv^{-2})\cdot 3$$

$$= \frac{2}{v}\sec^2\left(\frac{u}{v}\right) - \frac{3u}{v^2}\sec^2\left(\frac{u}{v}\right) = \frac{2v-3u}{v^2}\sec^2\left(\frac{u}{v}\right)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial u}\frac{\partial u}{\partial t} + \frac{\partial z}{\partial v}\frac{\partial v}{\partial t} = \sec^2(u/v)(1/v)\cdot 3 + \sec^2(u/v)(-uv^{-2})\cdot(-2)$$
$$= \frac{3}{v}\sec^2\left(\frac{u}{v}\right) + \frac{2u}{v^2}\sec^2\left(\frac{u}{v}\right) = \frac{2u+3v}{v^2}\sec^2\left(\frac{u}{v}\right)$$

13. Let $x = g(t)$ and $y = h(t)$. Then $p(t) = f(x, y)$ and the Chain Rule (2) gives $\dfrac{dp}{dt} = \dfrac{\partial f}{\partial x}\dfrac{dx}{dt} + \dfrac{\partial f}{\partial y}\dfrac{dy}{dt}$. When $t = 2$, $x = g(2) = 4$ and $y = h(2) = 5$, so $p'(2) = f_x(4,5)\,g'(2) + f_y(4,5)\,h'(2) = (2)(-3) + (8)(6) = 42$.

14. $R(s,t) = G\left(u(s,t), v(s,t)\right) \quad \Rightarrow \quad \dfrac{\partial R}{\partial s} = \dfrac{\partial R}{\partial u}\dfrac{\partial u}{\partial s} + \dfrac{\partial R}{\partial v}\dfrac{\partial v}{\partial s}$ and $\dfrac{\partial R}{\partial t} = \dfrac{\partial R}{\partial u}\dfrac{\partial u}{\partial t} + \dfrac{\partial R}{\partial v}\dfrac{\partial v}{\partial t}$ by the Chain Rule (3). When $s = 1$ and $t = 2$, $u(1,2) = 5$ and $v(1,2) = 7$.

Thus $R_s(1,2) = G_u(5,7)\,u_s(1,2) + G_v(5,7)\,v_s(1,2) = (9)(4) + (-2)(2) = 32$ and

$R_t(1,2) = G_u(5,7)\,u_t(1,2) + G_v(5,7)\,v_t(1,2) = (9)(-3) + (-2)(6) = -39$.

15. $g(u,v) = f(x(u,v), y(u,v))$ where $x = e^u + \sin v$, $y = e^u + \cos v \quad \Rightarrow$

$\dfrac{\partial x}{\partial u} = e^u$, $\dfrac{\partial x}{\partial v} = \cos v$, $\dfrac{\partial y}{\partial u} = e^u$, $\dfrac{\partial y}{\partial v} = -\sin v$. By the Chain Rule (3), $\dfrac{\partial g}{\partial u} = \dfrac{\partial f}{\partial x}\dfrac{\partial x}{\partial u} + \dfrac{\partial f}{\partial y}\dfrac{\partial y}{\partial u}$. Then

$g_u(0,0) = f_x(x(0,0), y(0,0))\,x_u(0,0) + f_y(x(0,0), y(0,0))\,y_u(0,0) = f_x(1,2)(e^0) + f_y(1,2)(e^0) = 2(1) + 5(1) = 7$.

Similarly, $\dfrac{\partial g}{\partial v} = \dfrac{\partial f}{\partial x}\dfrac{\partial x}{\partial v} + \dfrac{\partial f}{\partial y}\dfrac{\partial y}{\partial v}$. Then

$g_v(0,0) = f_x(x(0,0), y(0,0))\,x_v(0,0) + f_y(x(0,0), y(0,0))\,y_v(0,0) = f_x(1,2)(\cos 0) + f_y(1,2)(-\sin 0)$
$= 2(1) + 5(0) = 2$

16. $g(r,s) = f(x(r,s), y(r,s))$ where $x = 2r - s$, $y = s^2 - 4r \quad \Rightarrow \quad \dfrac{\partial x}{\partial r} = 2$, $\dfrac{\partial x}{\partial s} = -1$, $\dfrac{\partial y}{\partial r} = -4$, $\dfrac{\partial y}{\partial s} = 2s$.

By the Chain Rule (3) $\dfrac{\partial g}{\partial r} = \dfrac{\partial f}{\partial x}\dfrac{\partial x}{\partial r} + \dfrac{\partial f}{\partial y}\dfrac{\partial y}{\partial r}$. Then

$g_r(1,2) = f_x(x(1,2), y(1,2))\,x_r(1,2) + f_y(x(1,2), y(1,2))\,y_r(1,2) = f_x(0,0)(2) + f_y(0,0)(-4)$
$= 4(2) + 8(-4) = -24$

Similarly, $\dfrac{\partial g}{\partial s} = \dfrac{\partial f}{\partial x}\dfrac{\partial x}{\partial s} + \dfrac{\partial f}{\partial y}\dfrac{\partial y}{\partial s}$. Then

$g_s(1,2) = f_x(x(1,2), y(1,2))\,x_s(1,2) + f_y(x(1,2), y(1,2))\,y_s(1,2) = f_x(0,0)(-1) + f_y(0,0)(4)$
$= 4(-1) + 8(4) = 28$

17.

$u = f(x,y)$, $x = x(r,s,t)$, $y = y(r,s,t) \quad \Rightarrow$

$$\frac{\partial u}{\partial r} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial r} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial r}, \quad \frac{\partial u}{\partial s} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial s},$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial t}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

18.



$$w = f(x,y,z),\ x = x(u,v),\ y = y(u,v),\ z = z(u,v) \ \Rightarrow$$

$$\frac{\partial w}{\partial u} = \frac{\partial w}{\partial x}\frac{\partial x}{\partial u} + \frac{\partial w}{\partial y}\frac{\partial y}{\partial u} + \frac{\partial w}{\partial z}\frac{\partial z}{\partial u},$$

$$\frac{\partial w}{\partial v} = \frac{\partial w}{\partial x}\frac{\partial x}{\partial v} + \frac{\partial w}{\partial y}\frac{\partial y}{\partial v} + \frac{\partial w}{\partial z}\frac{\partial z}{\partial v}$$

19.



$$T = F(p,q,r),\ p = p(x,y,z),\ q = q(x,y,z),\ r = r(x,y,z) \ \Rightarrow$$

$$\frac{\partial T}{\partial x} = \frac{\partial T}{\partial p}\frac{\partial p}{\partial x} + \frac{\partial T}{\partial q}\frac{\partial q}{\partial x} + \frac{\partial T}{\partial r}\frac{\partial r}{\partial x},$$

$$\frac{\partial T}{\partial y} = \frac{\partial T}{\partial p}\frac{\partial p}{\partial y} + \frac{\partial T}{\partial q}\frac{\partial q}{\partial y} + \frac{\partial T}{\partial r}\frac{\partial r}{\partial y},$$

$$\frac{\partial T}{\partial z} = \frac{\partial T}{\partial p}\frac{\partial p}{\partial z} + \frac{\partial T}{\partial q}\frac{\partial q}{\partial z} + \frac{\partial T}{\partial r}\frac{\partial r}{\partial z}$$

20.



$$R = F(t,u),\ t = t(w,x,y,z),\ u = u(w,x,y,z) \ \Rightarrow$$

$$\frac{\partial R}{\partial w} = \frac{\partial R}{\partial t}\frac{\partial t}{\partial w} + \frac{\partial R}{\partial u}\frac{\partial u}{\partial w},\ \frac{\partial R}{\partial x} = \frac{\partial R}{\partial t}\frac{\partial t}{\partial x} + \frac{\partial R}{\partial u}\frac{\partial u}{\partial x},$$

$$\frac{\partial R}{\partial y} = \frac{\partial R}{\partial t}\frac{\partial t}{\partial y} + \frac{\partial R}{\partial u}\frac{\partial u}{\partial y},\ \frac{\partial R}{\partial z} = \frac{\partial R}{\partial t}\frac{\partial t}{\partial z} + \frac{\partial R}{\partial u}\frac{\partial u}{\partial z}$$

21. $z = x^4 + x^2y,\ x = s + 2t - u,\ y = stu^2 \ \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = (4x^3 + 2xy)(1) + (x^2)(tu^2),$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = (4x^3 + 2xy)(2) + (x^2)(su^2),$$

$$\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial u} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial u} = (4x^3 + 2xy)(-1) + (x^2)(2stu).$$

When $s = 4$, $t = 2$, and $u = 1$ we have $x = 7$ and $y = 8$,

so $\dfrac{\partial z}{\partial s} = (1484)(1) + (49)(2) = 1582,\ \dfrac{\partial z}{\partial t} = (1484)(2) + (49)(4) = 3164,\ \dfrac{\partial z}{\partial u} = (1484)(-1) + (49)(16) = -700.$

22. $T = v/(2u+v) = v(2u+v)^{-1},\ u = pq\sqrt{r},\ v = p\sqrt{q}\,r \ \Rightarrow$

$$\frac{\partial T}{\partial p} = \frac{\partial T}{\partial u}\frac{\partial u}{\partial p} + \frac{\partial T}{\partial v}\frac{\partial v}{\partial p} = [-v(2u+v)^{-2}(2)](q\sqrt{r}) + \frac{(2u+v)(1) - v(1)}{(2u+v)^2}(\sqrt{q}\,r)$$

$$= \frac{-2v}{(2u+v)^2}(q\sqrt{r}) + \frac{2u}{(2u+v)^2}(\sqrt{q}\,r),$$

$$\frac{\partial T}{\partial q} = \frac{\partial T}{\partial u}\frac{\partial u}{\partial q} + \frac{\partial T}{\partial v}\frac{\partial v}{\partial q} = \frac{-2v}{(2u+v)^2}(p\sqrt{r}) + \frac{2u}{(2u+v)^2}\frac{pr}{2\sqrt{q}},$$

$$\frac{\partial T}{\partial r} = \frac{\partial T}{\partial u}\frac{\partial u}{\partial r} + \frac{\partial T}{\partial v}\frac{\partial v}{\partial r} = \frac{-2v}{(2u+v)^2}\frac{pq}{2\sqrt{r}} + \frac{2u}{(2u+v)^2}(p\sqrt{q}).$$

When $p = 2$, $q = 1$, and $r = 4$ we have $u = 4$ and $v = 8$,

so $\dfrac{\partial T}{\partial p} = \left(-\frac{1}{16}\right)(2) + \left(\frac{1}{32}\right)(4) = 0,\ \dfrac{\partial T}{\partial q} = \left(-\frac{1}{16}\right)(4) + \left(\frac{1}{32}\right)(4) = -\frac{1}{8},\ \dfrac{\partial T}{\partial r} = \left(-\frac{1}{16}\right)\left(\frac{1}{2}\right) + \left(\frac{1}{32}\right)(2) = \frac{1}{32}.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $w = xy + yz + zx,\ x = r\cos\theta,\ y = r\sin\theta,\ z = r\theta \ \Rightarrow$

$$\frac{\partial w}{\partial r} = \frac{\partial w}{\partial x}\frac{\partial x}{\partial r} + \frac{\partial w}{\partial y}\frac{\partial y}{\partial r} + \frac{\partial w}{\partial z}\frac{\partial z}{\partial r} = (y+z)(\cos\theta) + (x+z)(\sin\theta) + (y+x)(\theta),$$

$$\frac{\partial w}{\partial \theta} = \frac{\partial w}{\partial x}\frac{\partial x}{\partial \theta} + \frac{\partial w}{\partial y}\frac{\partial y}{\partial \theta} + \frac{\partial w}{\partial z}\frac{\partial z}{\partial \theta} = (y+z)(-r\sin\theta) + (x+z)(r\cos\theta) + (y+x)(r).$$

When $r = 2$ and $\theta = \pi/2$ we have $x = 0$, $y = 2$, and $z = \pi$, so $\dfrac{\partial w}{\partial r} = (2+\pi)(0) + (0+\pi)(1) + (2+0)(\pi/2) = 2\pi$ and

$$\frac{\partial w}{\partial \theta} = (2+\pi)(-2) + (0+\pi)(0) + (2+0)(2) = -2\pi.$$

24. $P = \sqrt{u^2+v^2+w^2} = (u^2+v^2+w^2)^{1/2},\ u = xe^y,\ v = ye^x,\ w = e^{xy} \ \Rightarrow$

$$\begin{aligned}\frac{\partial P}{\partial x} &= \frac{\partial P}{\partial u}\frac{\partial u}{\partial x} + \frac{\partial P}{\partial v}\frac{\partial v}{\partial x} + \frac{\partial P}{\partial w}\frac{\partial w}{\partial x}\\ &= \tfrac{1}{2}(u^2+v^2+w^2)^{-1/2}(2u)(e^y) + \tfrac{1}{2}(u^2+v^2+w^2)^{-1/2}(2v)(ye^x) + \tfrac{1}{2}(u^2+v^2+w^2)^{-1/2}(2w)(ye^{xy})\\ &= \frac{ue^y + vye^x + wye^{xy}}{\sqrt{u^2+v^2+w^2}}.\end{aligned}$$

$$\begin{aligned}\frac{\partial P}{\partial y} &= \frac{\partial P}{\partial u}\frac{\partial u}{\partial y} + \frac{\partial P}{\partial v}\frac{\partial v}{\partial y} + \frac{\partial P}{\partial w}\frac{\partial w}{\partial y} = \frac{u}{\sqrt{u^2+v^2+w^2}}(xe^y) + \frac{v}{\sqrt{u^2+v^2+w^2}}(e^x) + \frac{w}{\sqrt{u^2+v^2+w^2}}(xe^{xy})\\ &= \frac{uxe^y + ve^x + wxe^{xy}}{\sqrt{u^2+v^2+w^2}}.\end{aligned}$$

When $x = 0$ and $y = 2$ we have $u = 0$, $v = 2$, and $w = 1$, so $\dfrac{\partial P}{\partial x} = \dfrac{0+4+2}{\sqrt{5}} = \dfrac{6}{\sqrt{5}}$ and $\dfrac{\partial P}{\partial y} = \dfrac{0+2+0}{\sqrt{5}} = \dfrac{2}{\sqrt{5}}$.

25. $N = \dfrac{p+q}{p+r},\ p = u+vw,\ q = v+uw,\ r = w+uv \ \Rightarrow$

$$\begin{aligned}\frac{\partial N}{\partial u} &= \frac{\partial N}{\partial p}\frac{\partial p}{\partial u} + \frac{\partial N}{\partial q}\frac{\partial q}{\partial u} + \frac{\partial N}{\partial r}\frac{\partial r}{\partial u}\\ &= \frac{(p+r)(1)-(p+q)(1)}{(p+r)^2}(1) + \frac{(p+r)(1)-(p+q)(0)}{(p+r)^2}(w) + \frac{(p+r)(0)-(p+q)(1)}{(p+r)^2}(v)\\ &= \frac{(r-q)+(p+r)w-(p+q)v}{(p+r)^2},\end{aligned}$$

$$\frac{\partial N}{\partial v} = \frac{\partial N}{\partial p}\frac{\partial p}{\partial v} + \frac{\partial N}{\partial q}\frac{\partial q}{\partial v} + \frac{\partial N}{\partial r}\frac{\partial r}{\partial v} = \frac{r-q}{(p+r)^2}(w) + \frac{p+r}{(p+r)^2}(1) + \frac{-(p+q)}{(p+r)^2}(u) = \frac{(r-q)w+(p+r)-(p+q)u}{(p+r)^2},$$

$$\frac{\partial N}{\partial w} = \frac{\partial N}{\partial p}\frac{\partial p}{\partial w} + \frac{\partial N}{\partial q}\frac{\partial q}{\partial w} + \frac{\partial N}{\partial r}\frac{\partial r}{\partial w} = \frac{r-q}{(p+r)^2}(v) + \frac{p+r}{(p+r)^2}(u) + \frac{-(p+q)}{(p+r)^2}(1) = \frac{(r-q)v+(p+r)u-(p+q)}{(p+r)^2}.$$

When $u = 2$, $v = 3$, and $w = 4$ we have $p = 14$, $q = 11$, and $r = 10$, so $\dfrac{\partial N}{\partial u} = \dfrac{-1+(24)(4)-(25)(3)}{(24)^2} = \dfrac{20}{576} = \dfrac{5}{144}$,

$\dfrac{\partial N}{\partial v} = \dfrac{(-1)(4)+24-(25)(2)}{(24)^2} = \dfrac{-30}{576} = -\dfrac{5}{96}$, and $\dfrac{\partial N}{\partial w} = \dfrac{(-1)(3)+(24)(2)-25}{(24)^2} = \dfrac{20}{576} = \dfrac{5}{144}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

26. $u = xe^{ty},\ x = \alpha^2\beta,\ y = \beta^2\gamma,\ t = \gamma^2\alpha \ \Rightarrow$

$$\frac{\partial u}{\partial \alpha} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial \alpha} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial \alpha} + \frac{\partial u}{\partial t}\frac{\partial t}{\partial \alpha} = e^{ty}(2\alpha\beta) + xte^{ty}(0) + xye^{ty}(\gamma^2) = e^{ty}(2\alpha\beta + xy\gamma^2),$$

$$\frac{\partial u}{\partial \beta} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial \beta} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial \beta} + \frac{\partial u}{\partial t}\frac{\partial t}{\partial \beta} = e^{ty}(\alpha^2) + xte^{ty}(2\beta\gamma) + xye^{ty}(0) = e^{ty}(\alpha^2 + 2xt\beta\gamma),$$

$$\frac{\partial u}{\partial \gamma} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial \gamma} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial \gamma} + \frac{\partial u}{\partial t}\frac{\partial t}{\partial \gamma} = e^{ty}(0) + xte^{ty}(\beta^2) + xye^{ty}(2\gamma\alpha) = e^{ty}(xt\beta^2 + 2xy\alpha\gamma).$$

When $\alpha = -1$, $\beta = 2$, and $\gamma = 1$ we have $x = 2$, $y = 4$, and $t = -1$, so $\dfrac{\partial u}{\partial \alpha} = e^{-4}(-4 + 8) = 4e^{-4}$,

$\dfrac{\partial u}{\partial \beta} = e^{-4}(1 - 8) = -7e^{-4}$, and $\dfrac{\partial u}{\partial \gamma} = e^{-4}(-8 - 16) = -24e^{-4}$.

27. $y\cos x = x^2 + y^2$, so let $F(x, y) = y\cos x - x^2 - y^2 = 0$. Then by Equation 6

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{-y\sin x - 2x}{\cos x - 2y} = \frac{2x + y\sin x}{\cos x - 2y}.$$

28. $\cos(xy) = 1 + \sin y$, so let $F(x, y) = \cos(xy) - 1 - \sin y = 0$. Then by Equation 6

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{-\sin(xy)(y)}{-\sin(xy)(x) - \cos y} = -\frac{y\sin(xy)}{\cos y + x\sin(xy)}.$$

29. $\tan^{-1}(x^2y) = x + xy^2$, so let $F(x, y) = \tan^{-1}(x^2y) - x - xy^2 = 0$. Then

$$F_x(x, y) = \frac{1}{1 + (x^2y)^2}(2xy) - 1 - y^2 = \frac{2xy}{1 + x^4y^2} - 1 - y^2 = \frac{2xy - (1 + y^2)(1 + x^4y^2)}{1 + x^4y^2},$$

$$F_y(x, y) = \frac{1}{1 + (x^2y)^2}(x^2) - 2xy = \frac{x^2}{1 + x^4y^2} - 2xy = \frac{x^2 - 2xy(1 + x^4y^2)}{1 + x^4y^2}$$

and
$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{[2xy - (1 + y^2)(1 + x^4y^2)]/(1 + x^4y^2)}{[x^2 - 2xy(1 + x^4y^2)]/(1 + x^4y^2)} = \frac{(1 + y^2)(1 + x^4y^2) - 2xy}{x^2 - 2xy(1 + x^4y^2)}$$

$$= \frac{1 + x^4y^2 + y^2 + x^4y^4 - 2xy}{x^2 - 2xy - 2x^5y^3}$$

30. $e^y\sin x = x + xy$, so let $F(x, y) = e^y\sin x - x - xy = 0$. Then $\dfrac{dy}{dx} = -\dfrac{F_x}{F_y} = -\dfrac{e^y\cos x - 1 - y}{e^y\sin x - x} = \dfrac{1 + y - e^y\cos x}{e^y\sin x - x}$.

31. $x^2 + 2y^2 + 3z^2 = 1$, so let $F(x, y, z) = x^2 + 2y^2 + 3z^2 - 1 = 0$. Then by Equations 7

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{2x}{6z} = -\frac{x}{3z} \quad \text{and} \quad \frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{4y}{6z} = -\frac{2y}{3z}.$$

32. $x^2 - y^2 + z^2 - 2z = 4$, so let $F(x, y, z) = x^2 - y^2 + z^2 - 2z - 4 = 0$. Then by Equations 7

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{2x}{2z - 2} = \frac{x}{1 - z} \quad \text{and} \quad \frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{-2y}{2z - 2} = \frac{y}{z - 1}.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

33. $e^z = xyz$, so let $F(x,y,z) = e^z - xyz = 0$. Then $\dfrac{\partial z}{\partial x} = -\dfrac{F_x}{F_z} = -\dfrac{-yz}{e^z - xy} = \dfrac{yz}{e^z - xy}$ and

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{-xz}{e^z - xy} = \frac{xz}{e^z - xy}.$$

34. $yz + x\ln y = z^2$, so let $F(x,y,z) = yz + x\ln y - z^2 = 0$. Then $\dfrac{\partial z}{\partial x} = -\dfrac{F_x}{F_z} = -\dfrac{\ln y}{y - 2z} = \dfrac{\ln y}{2z - y}$ and

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{z + (x/y)}{y - 2z} = \frac{x + yz}{2yz - y^2}.$$

35. Since x and y are each functions of t, $T(x,y)$ is a function of t, so by the Chain Rule, $\dfrac{dT}{dt} = \dfrac{\partial T}{\partial x}\dfrac{dx}{dt} + \dfrac{\partial T}{\partial y}\dfrac{dy}{dt}$. After 3 seconds, $x = \sqrt{1+t} = \sqrt{1+3} = 2$, $y = 2 + \frac{1}{3}t = 2 + \frac{1}{3}(3) = 3$, $\dfrac{dx}{dt} = \dfrac{1}{2\sqrt{1+t}} = \dfrac{1}{2\sqrt{1+3}} = \dfrac{1}{4}$, and $\dfrac{dy}{dt} = \dfrac{1}{3}$. Then $\dfrac{dT}{dt} = T_x(2,3)\dfrac{dx}{dt} + T_y(2,3)\dfrac{dy}{dt} = 4\left(\frac{1}{4}\right) + 3\left(\frac{1}{3}\right) = 2$. Thus the temperature is rising at a rate of $2°\text{C/s}$.

36. (a) Since $\partial W/\partial T$ is negative, a rise in average temperature (while annual rainfall remains constant) causes a decrease in wheat production at the current production levels. Since $\partial W/\partial R$ is positive, an increase in annual rainfall (while the average temperature remains constant) causes an increase in wheat production.

(b) Since the average temperature is rising at a rate of $0.15°\text{C/year}$, we know that $dT/dt = 0.15$. Since rainfall is decreasing at a rate of 0.1 cm/year, we know $dR/dt = -0.1$. Then, by the Chain Rule, $\dfrac{dW}{dt} = \dfrac{\partial W}{\partial T}\dfrac{dT}{dt} + \dfrac{\partial W}{\partial R}\dfrac{dR}{dt} = (-2)(0.15) + (8)(-0.1) = -1.1$. Thus we estimate that wheat production will decrease at a rate of 1.1 units/year.

37. $C = 1449.2 + 4.6T - 0.055T^2 + 0.00029T^3 + 0.016D$, so $\dfrac{\partial C}{\partial T} = 4.6 - 0.11T + 0.00087T^2$ and $\dfrac{\partial C}{\partial D} = 0.016$.

According to the graph, the diver is experiencing a temperature of approximately $12.5°\text{C}$ at $t = 20$ minutes, so $\dfrac{\partial C}{\partial T} = 4.6 - 0.11(12.5) + 0.00087(12.5)^2 \approx 3.36$. By sketching tangent lines at $t = 20$ to the graphs given, we estimate $\dfrac{dD}{dt} \approx \dfrac{1}{2}$ and $\dfrac{dT}{dt} \approx -\dfrac{1}{10}$. Then, by the Chain Rule, $\dfrac{dC}{dt} = \dfrac{\partial C}{\partial T}\dfrac{dT}{dt} + \dfrac{\partial C}{\partial D}\dfrac{dD}{dt} \approx (3.36)\left(-\frac{1}{10}\right) + (0.016)\left(\frac{1}{2}\right) \approx -0.33$. Thus the speed of sound experienced by the diver is decreasing at a rate of approximately 0.33 m/s per minute.

38. $V = \pi r^2 h/3$, so $\dfrac{dV}{dt} = \dfrac{\partial V}{\partial r}\dfrac{dr}{dt} + \dfrac{\partial V}{\partial h}\dfrac{dh}{dt} = \dfrac{2\pi rh}{3}\,1.8 + \dfrac{\pi r^2}{3}(-2.5) = 20{,}160\pi - 12{,}000\pi = 8160\pi \text{ in}^3\text{/s}$.

39. (a) $V = \ell wh$, so by the Chain Rule,

$$\frac{dV}{dt} = \frac{\partial V}{\partial \ell}\frac{d\ell}{dt} + \frac{\partial V}{\partial w}\frac{dw}{dt} + \frac{\partial V}{\partial h}\frac{dh}{dt} = wh\frac{d\ell}{dt} + \ell h\frac{dw}{dt} + \ell w\frac{dh}{dt} = 2\cdot 2\cdot 2 + 1\cdot 2\cdot 2 + 1\cdot 2\cdot(-3) = 6 \text{ m}^3\text{/s}.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $S = 2(\ell w + \ell h + wh)$, so by the Chain Rule,

$$\begin{aligned}\frac{dS}{dt} &= \frac{\partial S}{\partial \ell}\frac{d\ell}{dt} + \frac{\partial S}{\partial w}\frac{dw}{dt} + \frac{\partial S}{\partial h}\frac{dh}{dt} = 2(w+h)\frac{d\ell}{dt} + 2(\ell+h)\frac{dw}{dt} + 2(\ell+w)\frac{dh}{dt} \\ &= 2(2+2)2 + 2(1+2)2 + 2(1+2)(-3) = 10 \text{ m}^2/\text{s}\end{aligned}$$

(c) $L^2 = \ell^2 + w^2 + h^2 \;\Rightarrow\; 2L\dfrac{dL}{dt} = 2\ell\dfrac{d\ell}{dt} + 2w\dfrac{dw}{dt} + 2h\dfrac{dh}{dt} = 2(1)(2) + 2(2)(2) + 2(2)(-3) = 0 \;\Rightarrow$

$dL/dt = 0$ m/s.

40. $I = \dfrac{V}{R} \;\Rightarrow$

$$\frac{dI}{dt} = \frac{\partial I}{\partial V}\frac{dV}{dt} + \frac{\partial I}{\partial R}\frac{dR}{dt} = \frac{1}{R}\frac{dV}{dt} - \frac{V}{R^2}\frac{dR}{dt} = \frac{1}{R}\frac{dV}{dt} - \frac{I}{R}\frac{dR}{dt} = \frac{1}{400}(-0.01) - \frac{0.08}{400}(0.03) = -0.000031 \text{ A/s}$$

41. $\dfrac{dP}{dt} = 0.05$, $\dfrac{dT}{dt} = 0.15$, $V = 8.31\dfrac{T}{P}$ and $\dfrac{dV}{dt} = \dfrac{8.31}{P}\dfrac{dT}{dt} - 8.31\dfrac{T}{P^2}\dfrac{dP}{dt}$. Thus when $P = 20$ and $T = 320$,

$$\frac{dV}{dt} = 8.31\left[\frac{0.15}{20} - \frac{(0.05)(320)}{400}\right] \approx -0.27 \text{ L/s}.$$

42. $P = 1.47L^{0.65}K^{0.35}$ and considering P, L, and K as functions of time t we have

$\dfrac{dP}{dt} = \dfrac{\partial P}{\partial L}\dfrac{dL}{dt} + \dfrac{\partial P}{\partial K}\dfrac{dK}{dt} = 1.47(0.65)L^{-0.35}K^{0.35}\dfrac{dL}{dt} + 1.47(0.35)L^{0.65}K^{-0.65}\dfrac{dK}{dt}$. We are given that $\dfrac{dL}{dt} = -2$ and $\dfrac{dK}{dt} = 0.5$, so when $L = 30$ and $K = 8$, the rate of change of production $\dfrac{dP}{dt}$ is $1.47(0.65)(30)^{-0.35}(8)^{0.35}(-2) + 1.47(0.35)(30)^{0.65}(8)^{-0.65}(0.5) \approx -0.596$. Thus production at that time is decreasing at a rate of about \$596,000 per year.

43. Let x be the length of the first side of the triangle and y the length of the second side. The area A of the triangle is given by $A = \frac{1}{2}xy\sin\theta$ where θ is the angle between the two sides. Thus A is a function of x, y, and θ, and x, y, and θ are each in turn functions of time t. We are given that $\dfrac{dx}{dt} = 3$, $\dfrac{dy}{dt} = -2$, and because A is constant, $\dfrac{dA}{dt} = 0$. By the Chain Rule, $\dfrac{dA}{dt} = \dfrac{\partial A}{\partial x}\dfrac{dx}{dt} + \dfrac{\partial A}{\partial y}\dfrac{dy}{dt} + \dfrac{\partial A}{\partial \theta}\dfrac{d\theta}{dt} \;\Rightarrow\; \dfrac{dA}{dt} = \frac{1}{2}y\sin\theta\cdot\dfrac{dx}{dt} + \frac{1}{2}x\sin\theta\cdot\dfrac{dy}{dt} + \frac{1}{2}xy\cos\theta\cdot\dfrac{d\theta}{dt}$. When $x = 20$, $y = 30$, and $\theta = \pi/6$ we have

$$\begin{aligned}0 &= \tfrac{1}{2}(30)\left(\sin\tfrac{\pi}{6}\right)(3) + \tfrac{1}{2}(20)\left(\sin\tfrac{\pi}{6}\right)(-2) + \tfrac{1}{2}(20)(30)\left(\cos\tfrac{\pi}{6}\right)\frac{d\theta}{dt} \\ &= 45\cdot\tfrac{1}{2} - 20\cdot\tfrac{1}{2} + 300\cdot\frac{\sqrt{3}}{2}\cdot\frac{d\theta}{dt} = \tfrac{25}{2} + 150\sqrt{3}\,\frac{d\theta}{dt}\end{aligned}$$

Solving for $\dfrac{d\theta}{dt}$ gives $\dfrac{d\theta}{dt} = \dfrac{-25/2}{150\sqrt{3}} = -\dfrac{1}{12\sqrt{3}}$, so the angle between the sides is decreasing at a rate of $1/(12\sqrt{3}) \approx 0.048$ rad/s.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

44. $f_o = \left(\dfrac{c+v_o}{c-v_s}\right) f_s = \left(\frac{332+34}{332-40}\right) 460 \approx 576.6$ Hz. v_o and v_s are functions of time t, so

$$\frac{df_o}{dt} = \frac{\partial f_o}{\partial v_o}\frac{dv_o}{dt} + \frac{\partial f_o}{\partial v_s}\frac{dv_s}{dt} = \left(\frac{1}{c-v_s}\right) f_s \cdot \frac{dv_o}{dt} + \frac{c+v_o}{(c-v_s)^2} f_s \cdot \frac{dv_s}{dt}$$
$$= \left(\tfrac{1}{332-40}\right)(460)\,(1.2) + \tfrac{332+34}{(332-40)^2}\,(460)\,(1.4) \approx 4.65 \text{ Hz/s}$$

45. (a) By the Chain Rule, $\dfrac{\partial z}{\partial r} = \dfrac{\partial z}{\partial x}\cos\theta + \dfrac{\partial z}{\partial y}\sin\theta$, $\dfrac{\partial z}{\partial \theta} = \dfrac{\partial z}{\partial x}(-r\sin\theta) + \dfrac{\partial z}{\partial y} r\cos\theta$.

(b) $\left(\dfrac{\partial z}{\partial r}\right)^2 = \left(\dfrac{\partial z}{\partial x}\right)^2 \cos^2\theta + 2\dfrac{\partial z}{\partial x}\dfrac{\partial z}{\partial y}\cos\theta\sin\theta + \left(\dfrac{\partial z}{\partial y}\right)^2 \sin^2\theta$,

$\left(\dfrac{\partial z}{\partial \theta}\right)^2 = \left(\dfrac{\partial z}{\partial x}\right)^2 r^2\sin^2\theta - 2\dfrac{\partial z}{\partial x}\dfrac{\partial z}{\partial y} r^2\cos\theta\sin\theta + \left(\dfrac{\partial z}{\partial y}\right)^2 r^2\cos^2\theta$. Thus

$$\left(\frac{\partial z}{\partial r}\right)^2 + \frac{1}{r^2}\left(\frac{\partial z}{\partial \theta}\right)^2 = \left[\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2\right](\cos^2\theta + \sin^2\theta) = \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2.$$

46. By the Chain Rule, $\dfrac{\partial u}{\partial s} = \dfrac{\partial u}{\partial x}e^s\cos t + \dfrac{\partial u}{\partial y}e^s\sin t$, $\dfrac{\partial u}{\partial t} = \dfrac{\partial u}{\partial x}(-e^s\sin t) + \dfrac{\partial u}{\partial y}e^s\cos t$. Then

$\left(\dfrac{\partial u}{\partial s}\right)^2 = \left(\dfrac{\partial u}{\partial x}\right)^2 e^{2s}\cos^2 t + 2\dfrac{\partial u}{\partial x}\dfrac{\partial u}{\partial y}e^{2s}\cos t\sin t + \left(\dfrac{\partial u}{\partial y}\right)^2 e^{2s}\sin^2 t$ and

$\left(\dfrac{\partial u}{\partial t}\right)^2 = \left(\dfrac{\partial u}{\partial x}\right)^2 e^{2s}\sin^2 t - 2\dfrac{\partial u}{\partial x}\dfrac{\partial u}{\partial y}e^{2s}\cos t\sin t + \left(\dfrac{\partial u}{\partial y}\right)^2 e^{2s}\sin^2 t$. Thus

$$\left[\left(\frac{\partial u}{\partial s}\right)^2 + \left(\frac{\partial u}{\partial t}\right)^2\right] e^{-2s} = \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2.$$

47. Let $u = x - y$ and $v = x + y$. Then $z = \dfrac{1}{x}[f(u) + g(v)]$ and

$$\frac{\partial z}{\partial x} = \frac{1}{x}\left[\frac{df}{du}\frac{\partial u}{\partial x} + \frac{dg}{dv}\frac{\partial v}{\partial x}\right] + [f(u) + g(v)]\left(-\frac{1}{x^2}\right)$$
$$= \frac{1}{x}[f'(u)(1) + g'(v)(1)] - \frac{1}{x^2}[f(u) + g(v)] = \frac{1}{x}[f'(u) + g'(v)] - \frac{1}{x^2}[f(u) + g(v)]$$

$$\frac{\partial z}{\partial y} = \frac{1}{x}\left[\frac{df}{du}\frac{\partial u}{\partial y} + \frac{dg}{dv}\frac{\partial v}{\partial y}\right] = \frac{1}{x}[f'(u)(-1) + g'(v)(1)] = \frac{1}{x}[-f'(u) + g'(v)]$$

$$\frac{\partial^2 z}{\partial y^2} = \frac{1}{x}\left[\frac{d}{du}[-f'(u)]\frac{\partial u}{\partial y} + \frac{d}{dv}[g'(v)]\frac{\partial v}{\partial y}\right] = \frac{1}{x}[-f''(u)(-1) + g''(v)(1)] = \frac{1}{x}[f''(u) + g''(v)]$$

Thus

$$\frac{\partial}{\partial x}\left(x^2\frac{\partial z}{\partial x}\right) = \frac{\partial}{\partial x}(x[f'(u) + g'(v)] - [f(u) + g(v)])$$
$$= x[f''(u)(1) + g''(v)(1)] + [f'(u) + g'(v)](1) - [f'(u)(1) + g'(v)(1)]$$
$$= x[f''(u) + g''(v)] + f'(u) + g'(v) - f'(u) - g'(v) = x[f''(u) + g''(v)]$$
$$\therefore x^2 \cdot \frac{1}{x}[f''(u) + g''(v)] = x^2\frac{\partial^2 z}{\partial y^2}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. Let $u = ax + y$ and $v = ax - y$. Then $z = \dfrac{1}{y}\,[f(u) + g(v)]$ and

$$\frac{\partial z}{\partial x} = \frac{1}{y}\left[\frac{df}{du}\frac{\partial u}{\partial x} + \frac{dg}{dv}\frac{\partial v}{\partial x}\right] = \frac{1}{y}\,[f'(u)(a) + g'(v)(a)] = \frac{a}{y}\,[f'(u) + g'(v)]$$

$$\frac{\partial^2 z}{\partial x^2} = \frac{a}{y}\left[\frac{d}{du}\,[f'(u)]\,\frac{\partial u}{\partial x} + \frac{d}{dv}\,[g'(v)]\,\frac{\partial v}{\partial x}\right] = \frac{a}{y}\,[f''(u)(a) + g''(v)(a)] = \frac{a^2}{y}\,[f''(u) + g''(v)]$$

$$\frac{\partial z}{\partial y} = \frac{1}{y}\left[\frac{df}{du}\frac{\partial u}{\partial y} + \frac{dg}{dv}\frac{\partial v}{\partial y}\right] + [f(u) + g(v)]\left(-\frac{1}{y^2}\right)$$

$$= \frac{1}{y}\,[f'(u)(1) + g'(v)(-1)] - \frac{1}{y^2}\,[f(u) + g(v)] = \frac{1}{y}\,[f'(u) - g'(v)] - \frac{1}{y^2}\,[f(u) + g(v)]$$

$$\frac{\partial}{\partial y}\left(y^2\frac{\partial z}{\partial y}\right) = \frac{\partial}{\partial y}\left(y\,[f'(u) - g'(v)] - [f(u) + g(v)]\right)$$

$$= y\,[f''(u)(1) - g''(v)(-1)] + [f'(u) - g'(v)]\,(1) - [f'(u)(1) + g'(v)(-1)]$$

$$= y\,[f''(u) + g''(v)] + f'(u) - g'(v) - f'(u) + g'(v) = y\,[f''(u) + g''(v)]$$

Thus $\dfrac{\partial^2 z}{\partial x^2} = \dfrac{a^2}{y}\,[f''(u) + g''(v)] = \dfrac{a^2}{y^2}\cdot y\,[f''(u) + g''(v)] = \dfrac{a^2}{y^2}\dfrac{\partial}{\partial y}\left(y^2\dfrac{\partial z}{\partial y}\right)$.

49. Let $u = x + at,\ v = x - at$. Then $z = f(u) + g(v)$, so $\partial z/\partial u = f'(u)$ and $\partial z/\partial v = g'(v)$.

Thus $\dfrac{\partial z}{\partial t} = \dfrac{\partial z}{\partial u}\dfrac{\partial u}{\partial t} + \dfrac{\partial z}{\partial v}\dfrac{\partial v}{\partial t} = af'(u) - ag'(v)$ and

$$\frac{\partial^2 z}{\partial t^2} = a\,\frac{\partial}{\partial t}\,[f'(u) - g'(v)] = a\left(\frac{df'(u)}{du}\frac{\partial u}{\partial t} - \frac{dg'(v)}{dv}\frac{\partial v}{\partial t}\right) = a^2 f''(u) + a^2 g''(v).$$

Similarly, $\dfrac{\partial z}{\partial x} = f'(u) + g'(v)$ and $\dfrac{\partial^2 z}{\partial x^2} = f''(u) + g''(v)$. Thus $\dfrac{\partial^2 z}{\partial t^2} = a^2\,\dfrac{\partial^2 z}{\partial x^2}$.

50. By the Chain Rule, $\dfrac{\partial u}{\partial s} = e^s\cos t\,\dfrac{\partial u}{\partial x} + e^s\sin t\,\dfrac{\partial u}{\partial y}$ and $\dfrac{\partial u}{\partial t} = -e^s\sin t\,\dfrac{\partial u}{\partial x} + e^s\cos t\,\dfrac{\partial u}{\partial y}$.

Then $\dfrac{\partial^2 u}{\partial s^2} = e^s\cos t\,\dfrac{\partial u}{\partial x} + e^s\cos t\,\dfrac{\partial}{\partial s}\left(\dfrac{\partial u}{\partial x}\right) + e^s\sin t\,\dfrac{\partial u}{\partial y} + e^s\sin t\,\dfrac{\partial}{\partial s}\left(\dfrac{\partial u}{\partial y}\right)$. But

$$\frac{\partial}{\partial s}\left(\frac{\partial u}{\partial x}\right) = \frac{\partial^2 u}{\partial x^2}\frac{\partial x}{\partial s} + \frac{\partial^2 u}{\partial y\,\partial x}\frac{\partial y}{\partial s} = e^s\cos t\,\frac{\partial^2 u}{\partial x^2} + e^s\sin t\,\frac{\partial^2 u}{\partial y\,\partial x} \text{ and}$$

$$\frac{\partial}{\partial s}\left(\frac{\partial u}{\partial y}\right) = \frac{\partial^2 u}{\partial y^2}\frac{\partial y}{\partial s} + \frac{\partial^2 u}{\partial x\,\partial y}\frac{\partial x}{\partial s} = e^s\sin t\,\frac{\partial^2 u}{\partial y^2} + e^s\cos t\,\frac{\partial^2 u}{\partial x\,\partial y}.$$

Also, by continuity of the partials, $\dfrac{\partial^2 u}{\partial x\,\partial y} = \dfrac{\partial^2 u}{\partial y\,\partial x}$. Thus

$$\frac{\partial^2 u}{\partial s^2} = e^s\cos t\,\frac{\partial u}{\partial x} + e^s\cos t\left(e^s\cos t\,\frac{\partial^2 u}{\partial x^2} + e^s\sin t\,\frac{\partial^2 u}{\partial x\,\partial y}\right) + e^s\sin t\,\frac{\partial u}{\partial y} + e^s\sin t\left(e^s\sin t\,\frac{\partial^2 u}{\partial y^2} + e^s\cos t\,\frac{\partial^2 u}{\partial x\,\partial y}\right)$$

$$= e^s\cos t\,\frac{\partial u}{\partial x} + e^s\sin t\,\frac{\partial u}{\partial y} + e^{2s}\cos^2 t\,\frac{\partial^2 u}{\partial x^2} + 2e^{2s}\cos t\sin t\,\frac{\partial^2 u}{\partial x\,\partial y} + e^{2s}\sin^2 t\,\frac{\partial^2 u}{\partial y^2}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Similarly

$$\begin{aligned}\frac{\partial^2 u}{\partial t^2} &= -e^s\cos t\,\frac{\partial u}{\partial x} - e^s\sin t\,\frac{\partial}{\partial t}\left(\frac{\partial u}{\partial x}\right) - e^s\sin t\,\frac{\partial u}{\partial y} + e^s\cos t\,\frac{\partial}{\partial t}\left(\frac{\partial u}{\partial y}\right)\\ &= -e^s\cos t\,\frac{\partial u}{\partial x} - e^s\sin t\left(-e^s\sin t\,\frac{\partial^2 u}{\partial x^2} + e^s\cos t\,\frac{\partial^2 u}{\partial x\,\partial y}\right)\\ &\qquad -e^s\sin t\,\frac{\partial u}{\partial y} + e^s\cos t\left(e^s\cos t\,\frac{\partial^2 u}{\partial y^2} - e^s\sin t\,\frac{\partial^2 u}{\partial x\,\partial y}\right)\\ &= -e^s\cos t\,\frac{\partial u}{\partial x} - e^s\sin t\,\frac{\partial u}{\partial y} + e^{2s}\sin^2 t\,\frac{\partial^2 u}{\partial x^2} - 2e^{2s}\cos t\sin t\,\frac{\partial^2 u}{\partial x\,\partial y} + e^{2s}\cos^2 t\,\frac{\partial^2 u}{\partial y^2}\end{aligned}$$

Thus $e^{-2s}\left(\dfrac{\partial^2 u}{\partial s^2} + \dfrac{\partial^2 u}{\partial t^2}\right) = (\cos^2 t + \sin^2 t)\left(\dfrac{\partial^2 u}{\partial x^2} + \dfrac{\partial^2 u}{\partial y^2}\right) = \dfrac{\partial^2 u}{\partial x^2} + \dfrac{\partial^2 u}{\partial y^2}$, as desired.

51. $\dfrac{\partial z}{\partial s} = \dfrac{\partial z}{\partial x}\,2s + \dfrac{\partial z}{\partial y}\,2r$. Then

$$\begin{aligned}\frac{\partial^2 z}{\partial r\,\partial s} &= \frac{\partial}{\partial r}\left(\frac{\partial z}{\partial x}2s\right) + \frac{\partial}{\partial r}\left(\frac{\partial z}{\partial y}2r\right)\\ &= \frac{\partial^2 z}{\partial x^2}\frac{\partial x}{\partial r}\,2s + \frac{\partial}{\partial y}\left(\frac{\partial z}{\partial x}\right)\frac{\partial y}{\partial r}\,2s + \frac{\partial z}{\partial x}\frac{\partial}{\partial r}\,2s + \frac{\partial^2 z}{\partial y^2}\frac{\partial y}{\partial r}\,2r + \frac{\partial}{\partial x}\left(\frac{\partial z}{\partial y}\right)\frac{\partial x}{\partial r}\,2r + \frac{\partial z}{\partial y}\,2\\ &= 4rs\,\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y\,\partial x}\,4s^2 + 0 + 4rs\,\frac{\partial^2 z}{\partial y^2} + \frac{\partial^2 z}{\partial x\,\partial y}\,4r^2 + 2\,\frac{\partial z}{\partial y}\end{aligned}$$

By the continuity of the partials, $\dfrac{\partial^2 z}{\partial r\partial s} = 4rs\,\dfrac{\partial^2 z}{\partial x^2} + 4rs\,\dfrac{\partial^2 z}{\partial y^2} + (4r^2 + 4s^2)\,\dfrac{\partial^2 z}{\partial x\,\partial y} + 2\dfrac{\partial z}{\partial y}$.

52. By the Chain Rule,

(a) $\dfrac{\partial z}{\partial r} = \dfrac{\partial z}{\partial x}\cos\theta + \dfrac{\partial z}{\partial y}\sin\theta$ (b) $\dfrac{\partial z}{\partial \theta} = -\dfrac{\partial z}{\partial x}\,r\sin\theta + \dfrac{\partial z}{\partial y}\,r\cos\theta$

(c) $$\begin{aligned}\frac{\partial^2 z}{\partial r\,\partial\theta} &= \frac{\partial^2 z}{\partial\theta\,\partial r} = \frac{\partial}{\partial\theta}\left(\frac{\partial z}{\partial x}\cos\theta + \frac{\partial z}{\partial y}\sin\theta\right) = -\sin\theta\,\frac{\partial z}{\partial x} + \cos\theta\,\frac{\partial}{\partial\theta}\left(\frac{\partial z}{\partial x}\right) + \cos\theta\,\frac{\partial z}{\partial y} + \sin\theta\,\frac{\partial}{\partial\theta}\left(\frac{\partial z}{\partial y}\right)\\ &= -\sin\theta\,\frac{\partial z}{\partial x} + \cos\theta\left(\frac{\partial^2 z}{\partial x^2}\frac{\partial x}{\partial\theta} + \frac{\partial^2 z}{\partial y\,\partial x}\frac{\partial y}{\partial\theta}\right) + \cos\theta\,\frac{\partial z}{\partial y} + \sin\theta\,\frac{\partial^2 z}{\partial y^2}\frac{\partial y}{\partial\theta} + \frac{\partial^2 z}{\partial x\,\partial y}\frac{\partial x}{\partial\theta}\\ &= -\sin\theta\,\frac{\partial z}{\partial x} + \cos\theta\left(-r\sin\theta\,\frac{\partial^2 z}{\partial x^2} + r\cos\theta\,\frac{\partial^2 z}{\partial y\,\partial x}\right) + \cos\theta\,\frac{\partial z}{\partial y} + \sin\theta\left(r\cos\theta\,\frac{\partial^2 z}{\partial y^2} - r\sin\theta\,\frac{\partial^2 z}{\partial x\,\partial y}\right)\\ &= -\sin\theta\,\frac{\partial z}{\partial x} - r\cos\theta\sin\theta\,\frac{\partial^2 z}{\partial x^2} + r\cos^2\theta\,\frac{\partial^2 z}{\partial y\,\partial x} + \cos\theta\,\frac{\partial z}{\partial y} + r\cos\theta\sin\theta\,\frac{\partial^2 z}{\partial y^2} - r\sin^2\theta\,\frac{\partial^2 z}{\partial y\,\partial x}\\ &= \cos\theta\,\frac{\partial z}{\partial y} - \sin\theta\,\frac{\partial z}{\partial x} + r\cos\theta\sin\theta\left(\frac{\partial^2 z}{\partial y^2} - \frac{\partial^2 z}{\partial x^2}\right) + r(\cos^2\theta - \sin^2\theta)\,\frac{\partial^2 z}{\partial y\,\partial x}\end{aligned}$$

53. $\dfrac{\partial z}{\partial r} = \dfrac{\partial z}{\partial x}\cos\theta + \dfrac{\partial z}{\partial y}\sin\theta$ and $\dfrac{\partial z}{\partial\theta} = -\dfrac{\partial z}{\partial x}\,r\sin\theta + \dfrac{\partial z}{\partial y}\,r\cos\theta$. Then

$$\begin{aligned}\frac{\partial^2 z}{\partial r^2} &= \cos\theta\left(\frac{\partial^2 z}{\partial x^2}\cos\theta + \frac{\partial^2 z}{\partial y\,\partial x}\sin\theta\right) + \sin\theta\left(\frac{\partial^2 z}{\partial y^2}\sin\theta + \frac{\partial^2 z}{\partial x\,\partial y}\cos\theta\right)\\ &= \cos^2\theta\,\frac{\partial^2 z}{\partial x^2} + 2\cos\theta\sin\theta\,\frac{\partial^2 z}{\partial x\,\partial y} + \sin^2\theta\,\frac{\partial^2 z}{\partial y^2}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and

$$\begin{aligned}\frac{\partial^2 z}{\partial \theta^2} &= -r\cos\theta\,\frac{\partial z}{\partial x} + (-r\sin\theta)\left(\frac{\partial^2 z}{\partial x^2}(-r\sin\theta) + \frac{\partial^2 z}{\partial y\,\partial x}\,r\cos\theta\right)\\ &\qquad -r\sin\theta\,\frac{\partial z}{\partial y} + r\cos\theta\left(\frac{\partial^2 z}{\partial y^2}\,r\cos\theta + \frac{\partial^2 z}{\partial x\,\partial y}(-r\sin\theta)\right)\\ &= -r\cos\theta\,\frac{\partial z}{\partial x} - r\sin\theta\,\frac{\partial z}{\partial y} + r^2\sin^2\theta\,\frac{\partial^2 z}{\partial x^2} - 2r^2\cos\theta\,\sin\theta\,\frac{\partial^2 z}{\partial x\,\partial y} + r^2\cos^2\theta\,\frac{\partial^2 z}{\partial y^2}\end{aligned}$$

Thus

$$\begin{aligned}\frac{\partial^2 z}{\partial r^2} + \frac{1}{r^2}\frac{\partial^2 z}{\partial \theta^2} + \frac{1}{r}\frac{\partial z}{\partial r} &= (\cos^2\theta + \sin^2\theta)\,\frac{\partial^2 z}{\partial x^2} + (\sin^2\theta + \cos^2\theta)\,\frac{\partial^2 z}{\partial y^2}\\ &\qquad -\frac{1}{r}\cos\theta\,\frac{\partial z}{\partial x} - \frac{1}{r}\sin\theta\,\frac{\partial z}{\partial y} + \frac{1}{r}\left(\cos\theta\,\frac{\partial z}{\partial x} + \sin\theta\,\frac{\partial z}{\partial y}\right)\\ &= \frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} \text{ as desired.}\end{aligned}$$

54. (a) $\dfrac{\partial z}{\partial t} = \dfrac{\partial z}{\partial x}\dfrac{\partial x}{\partial t} + \dfrac{\partial z}{\partial y}\dfrac{\partial y}{\partial t}$. Then

$$\begin{aligned}\frac{\partial^2 z}{\partial t^2} &= \frac{\partial}{\partial t}\left(\frac{\partial z}{\partial x}\frac{\partial x}{\partial t}\right) + \frac{\partial}{\partial t}\left(\frac{\partial z}{\partial y}\frac{\partial y}{\partial t}\right) = \frac{\partial}{\partial t}\left(\frac{\partial z}{\partial x}\right)\frac{\partial x}{\partial t} + \frac{\partial^2 x}{\partial t^2}\frac{\partial z}{\partial x} + \frac{\partial}{\partial t}\left(\frac{\partial z}{\partial y}\right)\frac{\partial y}{\partial t} + \frac{\partial^2 y}{\partial t^2}\frac{\partial z}{\partial y}\\ &= \frac{\partial^2 z}{\partial x^2}\left(\frac{\partial x}{\partial t}\right)^2 + \frac{\partial^2 z}{\partial y\,\partial x}\frac{\partial x}{\partial t}\frac{\partial y}{\partial t} + \frac{\partial^2 x}{\partial t^2}\frac{\partial z}{\partial x} + \frac{\partial^2 z}{\partial y^2}\left(\frac{\partial y}{\partial t}\right)^2 + \frac{\partial^2 z}{\partial x\,\partial y}\frac{\partial y}{\partial t}\frac{\partial x}{\partial t} + \frac{\partial^2 y}{\partial t^2}\frac{\partial z}{\partial y}\\ &= \frac{\partial^2 z}{\partial x^2}\left(\frac{\partial x}{\partial t}\right)^2 + 2\,\frac{\partial^2 z}{\partial x\,\partial y}\frac{\partial x}{\partial t}\frac{\partial y}{\partial t} + \frac{\partial^2 z}{\partial y^2}\left(\frac{\partial y}{\partial t}\right)^2 + \frac{\partial^2 x}{\partial t^2}\frac{\partial z}{\partial x} + \frac{\partial^2 y}{\partial t^2}\frac{\partial z}{\partial y}\end{aligned}$$

(b) $\dfrac{\partial^2 z}{\partial s\,\partial t} = \dfrac{\partial}{\partial s}\left(\dfrac{\partial z}{\partial x}\dfrac{\partial x}{\partial t} + \dfrac{\partial z}{\partial y}\dfrac{\partial y}{\partial t}\right)$

$$\begin{aligned}&= \left(\frac{\partial^2 z}{\partial x^2}\frac{\partial x}{\partial s} + \frac{\partial^2 z}{\partial y\,\partial x}\frac{\partial y}{\partial s}\right)\frac{\partial x}{\partial t} + \frac{\partial z}{\partial x}\frac{\partial^2 x}{\partial s\,\partial t} + \left(\frac{\partial^2 z}{\partial y^2}\frac{\partial y}{\partial s} + \frac{\partial^2 z}{\partial x\,\partial y}\frac{\partial x}{\partial s}\right)\frac{\partial y}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial^2 y}{\partial s\,\partial t}\\ &= \frac{\partial^2 z}{\partial x^2}\frac{\partial x}{\partial s}\frac{\partial x}{\partial t} + \frac{\partial^2 z}{\partial x\,\partial y}\left(\frac{\partial y}{\partial s}\frac{\partial x}{\partial t} + \frac{\partial y}{\partial t}\frac{\partial x}{\partial s}\right) + \frac{\partial z}{\partial x}\frac{\partial^2 x}{\partial s\,\partial t} + \frac{\partial z}{\partial y}\frac{\partial^2 y}{\partial s\,\partial t} + \frac{\partial^2 z}{\partial y^2}\frac{\partial y}{\partial s}\frac{\partial y}{\partial t}\end{aligned}$$

55. (a) Since f is a polynomial, it has continuous second-order partial derivatives, and

$$f(tx, ty) = (tx)^2(ty) + 2(tx)(ty)^2 + 5(ty)^3 = t^3x^2y + 2t^3xy^2 + 5t^3y^3 = t^3(x^2y + 2xy^2 + 5y^3) = t^3 f(x, y).$$

Thus, f is homogeneous of degree 3.

(b) Differentiating both sides of $f(tx, ty) = t^n f(x, y)$ with respect to t using the Chain Rule, we get

$$\frac{\partial}{\partial t} f(tx, ty) = \frac{\partial}{\partial t}\left[t^n f(x, y)\right] \quad \Leftrightarrow$$

$$\frac{\partial}{\partial (tx)} f(tx, ty)\cdot\frac{\partial (tx)}{\partial t} + \frac{\partial}{\partial (ty)} f(tx, ty)\cdot\frac{\partial (ty)}{\partial t} = x\,\frac{\partial}{\partial (tx)} f(tx, ty) + y\,\frac{\partial}{\partial (ty)} f(tx, ty) = nt^{n-1} f(x, y).$$

Setting $t = 1$: $x\,\dfrac{\partial}{\partial x} f(x, y) + y\,\dfrac{\partial}{\partial y} f(x, y) = nf(x, y)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

56. Differentiating both sides of $f(tx,ty) = t^n f(x,y)$ with respect to t using the Chain Rule, we get

$$\frac{\partial}{\partial(tx)} f(tx,ty)\cdot\frac{\partial(tx)}{\partial t} + \frac{\partial}{\partial(ty)} f(tx,ty)\cdot\frac{\partial(ty)}{\partial t} = x\frac{\partial}{\partial(tx)} f(tx,ty) + y\frac{\partial}{\partial(ty)} f(tx,ty) = nt^{n-1} f(x,y) \text{ and}$$

differentiating again with respect to t gives

$$x\left[\frac{\partial^2}{\partial(tx)^2} f(tx,ty)\cdot\frac{\partial(tx)}{\partial t} + \frac{\partial^2}{\partial(ty)\,\partial(tx)} f(tx,ty)\cdot\frac{\partial(ty)}{\partial t}\right]$$

$$+ y\left[\frac{\partial^2}{\partial(tx)\,\partial(ty)} f(tx,ty)\cdot\frac{\partial(tx)}{\partial t} + \frac{\partial^2}{\partial(ty)^2} f(tx,ty)\cdot\frac{\partial(ty)}{\partial t}\right] = n(n-1)t^{n-1} f(x,y).$$

Setting $t = 1$ and using the fact that $f_{yx} = f_{xy}$, we have $x^2 f_{xx} + 2xy f_{xy} + y^2 f_{yy} = n(n-1) f(x,y)$.

57. Differentiating both sides of $f(tx,ty) = t^n f(x,y)$ with respect to x using the Chain Rule, we get

$$\frac{\partial}{\partial x} f(tx,ty) = \frac{\partial}{\partial x}\left[t^n f(x,y)\right] \quad \Leftrightarrow$$

$$\frac{\partial}{\partial(tx)} f(tx,ty)\cdot\frac{\partial(tx)}{\partial x} + \frac{\partial}{\partial(ty)} f(tx,ty)\cdot\frac{\partial(ty)}{\partial x} = t^n\frac{\partial}{\partial x} f(x,y) \quad \Leftrightarrow \quad t f_x(tx,ty) = t^n f_x(x,y).$$

Thus $f_x(tx,ty) = t^{n-1} f_x(x,y)$.

58. $F(x,y,z) = 0$ is assumed to define z as a function of x and y, that is, $z = f(x,y)$. So by (7), $\dfrac{\partial z}{\partial x} = -\dfrac{F_x}{F_z}$ since $F_z \neq 0$.

Similarly, it is assumed that $F(x,y,z) = 0$ defines x as a function of y and z, that is $x = h(x,z)$. Then $F(h(y,z),y,z) = 0$ and by the Chain Rule, $F_x\dfrac{\partial x}{\partial y} + F_y\dfrac{\partial y}{\partial y} + F_z\dfrac{\partial z}{\partial y} = 0$. But $\dfrac{\partial z}{\partial y} = 0$ and $\dfrac{\partial y}{\partial y} = 1$, so $F_x\dfrac{\partial x}{\partial y} + F_y = 0 \;\Rightarrow\; \dfrac{\partial x}{\partial y} = -\dfrac{F_y}{F_x}$.

A similar calculation shows that $\dfrac{\partial y}{\partial z} = -\dfrac{F_z}{F_y}$. Thus $\dfrac{\partial z}{\partial x}\dfrac{\partial x}{\partial y}\dfrac{\partial y}{\partial z} = \left(-\dfrac{F_x}{F_z}\right)\left(-\dfrac{F_y}{F_x}\right)\left(-\dfrac{F_z}{F_y}\right) = -1$.

59. Given a function defined implicitly by $F(x,y) = 0$, where F is differentiable and $F_y \neq 0$, we know that $\dfrac{dy}{dx} = -\dfrac{F_x}{F_y}$. Let

$G(x,y) = -\dfrac{F_x}{F_y}$ so $\dfrac{dy}{dx} = G(x,y)$. Differentiating both sides with respect to x and using the Chain Rule gives

$$\frac{d^2y}{dx^2} = \frac{\partial G}{\partial x}\frac{dx}{dx} + \frac{\partial G}{\partial y}\frac{dy}{dx} \text{ where } \frac{\partial G}{\partial x} = \frac{\partial}{\partial x}\left(-\frac{F_x}{F_y}\right) = -\frac{F_y F_{xx} - F_x F_{yx}}{F_y^2},\ \frac{\partial G}{\partial y} = \frac{\partial}{\partial y}\left(-\frac{F_x}{F_y}\right) = -\frac{F_y F_{xy} - F_x F_{yy}}{F_y^2}.$$

Thus

$$\frac{d^2y}{dx^2} = \left(-\frac{F_y F_{xx} - F_x F_{yx}}{F_y^2}\right)(1) + \left(-\frac{F_y F_{xy} - F_x F_{yy}}{F_y^2}\right)\left(-\frac{F_x}{F_y}\right)$$

$$= -\frac{F_{xx}F_y^2 - F_{yx}F_xF_y - F_{xy}F_yF_x + F_{yy}F_x^2}{F_y^3}$$

But F has continuous second derivatives, so by Clauraut's Theorem, $F_{yx} = F_{xy}$ and we have

$$\frac{d^2y}{dx^2} = -\frac{F_{xx}F_y^2 - 2F_{xy}F_xF_y + F_{yy}F_x^2}{F_y^3} \text{ as desired.}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.6 Directional Derivatives and the Gradient Vector

1. We can approximate the directional derivative of the pressure function at K in the direction of S by the average rate of change of pressure between the points where the red line intersects the contour lines closest to K (extend the red line slightly at the left). In the direction of S, the pressure changes from 1000 millibars to 996 millibars and we estimate the distance between these two points to be approximately 50 km (using the fact that the distance from K to S is 300 km). Then the rate of change of pressure in the direction given is approximately $\frac{996-1000}{50} = -0.08$ millibar/km.

2. First we draw a line passing through Dubbo and Sydney. We approximate the directional derivative at Dubbo in the direction of Sydney by the average rate of change of temperature between the points where the line intersects the contour lines closest to Dubbo. In the direction of Sydney, the temperature changes from 30°C to 27°C. We estimate the distance between these two points to be approximately 120 km, so the rate of change of maximum temperature in the direction given is approximately $\frac{27-30}{120} = -0.025$°C/km.

3. $D_{\mathbf{u}} f(-20,30) = \nabla f(-20,30) \cdot \mathbf{u} = f_T(-20,30)\left(\frac{1}{\sqrt{2}}\right) + f_v(-20,30)\left(\frac{1}{\sqrt{2}}\right)$.

   $f_T(-20,30) = \lim_{h\to 0} \dfrac{f(-20+h,30) - f(-20,30)}{h}$, so we can approximate $f_T(-20,30)$ by considering $h = \pm 5$ and using the values given in the table: $f_T(-20,30) \approx \dfrac{f(-15,30) - f(-20,30)}{5} = \dfrac{-26-(-33)}{5} = 1.4$,

   $f_T(-20,30) \approx \dfrac{f(-25,30) - f(-20,30)}{-5} = \dfrac{-39-(-33)}{-5} = 1.2$. Averaging these values gives $f_T(-20,30) \approx 1.3$.

   Similarly, $f_v(-20,30) = \lim_{h\to 0} \dfrac{f(-20,30+h) - f(-20,30)}{h}$, so we can approximate $f_v(-20,30)$ with $h = \pm 10$:

   $f_v(-20,30) \approx \dfrac{f(-20,40) - f(-20,30)}{10} = \dfrac{-34-(-33)}{10} = -0.1$,

   $f_v(-20,30) \approx \dfrac{f(-20,20) - f(-20,30)}{-10} = \dfrac{-30-(-33)}{-10} = -0.3$. Averaging these values gives $f_v(-20,30) \approx -0.2$.

   Then $D_{\mathbf{u}} f(-20,30) \approx 1.3\left(\frac{1}{\sqrt{2}}\right) + (-0.2)\left(\frac{1}{\sqrt{2}}\right) \approx 0.778$.

4. $f(x,y) = xy^3 - x^2 \;\Rightarrow\; f_x(x,y) = y^3 - 2x$ and $f_y(x,y) = 3xy^2$. If $\mathbf{u}$ is a unit vector in the direction of $\theta = \pi/3$, then from Equation 6, $D_{\mathbf{u}} f(1,2) = f_x(1,2)\cos\left(\frac{\pi}{3}\right) + f_y(1,2)\sin\left(\frac{\pi}{3}\right) = 6 \cdot \frac{1}{2} + 12 \cdot \frac{\sqrt{3}}{2} = 3 + 6\sqrt{3}$.

5. $f(x,y) = y\cos(xy) \;\Rightarrow\; f_x(x,y) = y[-\sin(xy)](y) = -y^2\sin(xy)$ and $f_y(x,y) = y[-\sin(xy)](x) + [\cos(xy)](1) = \cos(xy) - xy\sin(xy)$. If $\mathbf{u}$ is a unit vector in the direction of $\theta = \pi/4$, then from Equation 6, $D_{\mathbf{u}} f(0,1) = f_x(0,1)\cos\left(\frac{\pi}{4}\right) + f_y(0,1)\sin\left(\frac{\pi}{4}\right) = 0 \cdot \frac{\sqrt{2}}{2} + 1 \cdot \frac{\sqrt{2}}{2} = \frac{\sqrt{2}}{2}$.

6. $f(x,y) = \sqrt{2x+3y} \;\Rightarrow\; f_x(x,y) = \frac{1}{2}(2x+3y)^{-1/2}(2) = 1/\sqrt{2x+3y}$ and $f_y(x,y) = \frac{1}{2}(2x+3y)^{-1/2}(3) = 3/\left(2\sqrt{2x+3y}\right)$. If $\mathbf{u}$ is a unit vector in the direction of $\theta = -\pi/6$, then from Equation 6, $D_{\mathbf{u}} f(3,1) = f_x(3,1)\cos\left(-\frac{\pi}{6}\right) + f_y(3,1)\sin\left(-\frac{\pi}{6}\right) = \frac{1}{3} \cdot \frac{\sqrt{3}}{2} + \frac{1}{2} \cdot \left(-\frac{1}{2}\right) = \frac{\sqrt{3}}{6} - \frac{1}{4}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

7. $f(x,y) = x/y = xy^{-1}$

(a) $\nabla f(x,y) = \dfrac{\partial f}{\partial x}\,\mathbf{i} + \dfrac{\partial f}{\partial y}\,\mathbf{j} = y^{-1}\,\mathbf{i} + (-xy^{-2})\,\mathbf{j} = \dfrac{1}{y}\,\mathbf{i} - \dfrac{x}{y^2}\,\mathbf{j}$

(b) $\nabla f(2,1) = \dfrac{1}{1}\,\mathbf{i} - \dfrac{2}{1^2}\,\mathbf{j} = \mathbf{i} - 2\,\mathbf{j}$

(c) By Equation 9, $D_{\mathbf{u}} f(2,1) = \nabla f(2,1)\cdot\mathbf{u} = (\mathbf{i} - 2\,\mathbf{j})\cdot\left(\frac{3}{5}\,\mathbf{i} + \frac{4}{5}\,\mathbf{j}\right) = \frac{3}{5} - \frac{8}{5} = -1.$

8. $f(x,y) = x^2 \ln y$

(a) $\nabla f(x,y) = \dfrac{\partial f}{\partial x}\,\mathbf{i} + \dfrac{\partial f}{\partial y}\,\mathbf{j} = 2x\ln y\,\mathbf{i} + (x^2/y)\,\mathbf{j}$

(b) $\nabla f(3,1) = 0\,\mathbf{i} + (9/1)\,\mathbf{j} = 9\,\mathbf{j}$

(c) By Equation 9, $D_{\mathbf{u}} f(3,1) = \nabla f(3,1)\cdot\mathbf{u} = 9\,\mathbf{j}\cdot\left(-\frac{5}{13}\,\mathbf{i} + \frac{12}{13}\,\mathbf{j}\right) = 0 + \frac{108}{13} = \frac{108}{13}.$

9. $f(x,y,z) = x^2yz - xyz^3$

(a) $\nabla f(x,y,z) = \langle f_x(x,y,z), f_y(x,y,z), f_z(x,y,z)\rangle = \left\langle 2xyz - yz^3, x^2z - xz^3, x^2y - 3xyz^2\right\rangle$

(b) $\nabla f(2,-1,1) = \langle -4+1, 4-2, -4+6\rangle = \langle -3,2,2\rangle$

(c) By Equation 14, $D_{\mathbf{u}} f(2,-1,1) = \nabla f(2,-1,1)\cdot\mathbf{u} = \langle -3,2,2\rangle\cdot\left\langle 0, \frac{4}{5}, -\frac{3}{5}\right\rangle = 0 + \frac{8}{5} - \frac{6}{5} = \frac{2}{5}.$

10. $f(x,y,z) = y^2 e^{xyz}$

(a) $\nabla f(x,y,z) = \langle f_x(x,y,z), f_y(x,y,z), f_z(x,y,z)\rangle = \left\langle y^2e^{xyz}(yz), y^2\cdot e^{xyz}(xz) + e^{xyz}\cdot 2y, y^2e^{xyz}(xy)\right\rangle$

$= \left\langle y^3ze^{xyz}, (xy^2z + 2y)e^{xyz}, xy^3e^{xyz}\right\rangle$

(b) $\nabla f(0,1,-1) = \langle -1,2,0\rangle$

(c) $D_{\mathbf{u}} f(0,1,-1) = \nabla f(0,1,-1)\cdot\mathbf{u} = \langle -1,2,0\rangle\cdot\left\langle \frac{3}{13}, \frac{4}{13}, \frac{12}{13}\right\rangle = -\frac{3}{13} + \frac{8}{13} + 0 = \frac{5}{13}$

11. $f(x,y) = e^x \sin y \;\Rightarrow\; \nabla f(x,y) = \langle e^x\sin y, e^x\cos y\rangle$, $\nabla f(0,\pi/3) = \left\langle \frac{\sqrt{3}}{2}, \frac{1}{2}\right\rangle$, and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{(-6)^2+8^2}}\langle -6,8\rangle = \frac{1}{10}\langle -6,8\rangle = \left\langle -\frac{3}{5}, \frac{4}{5}\right\rangle$, so

$D_{\mathbf{u}} f(0,\pi/3) = \nabla f(0,\pi/3)\cdot\mathbf{u} = \left\langle \frac{\sqrt{3}}{2}, \frac{1}{2}\right\rangle\cdot\left\langle -\frac{3}{5}, \frac{4}{5}\right\rangle = -\frac{3\sqrt{3}}{10} + \frac{4}{10} = \frac{4-3\sqrt{3}}{10}.$

12. $f(x,y) = \dfrac{x}{x^2+y^2} \;\Rightarrow\; \nabla f(x,y) = \left\langle \dfrac{(x^2+y^2)(1) - x(2x)}{(x^2+y^2)^2}, \dfrac{0 - x(2y)}{(x^2+y^2)^2}\right\rangle = \left\langle \dfrac{y^2-x^2}{(x^2+y^2)^2}, -\dfrac{2xy}{(x^2+y^2)^2}\right\rangle,$

$\nabla f(1,2) = \left\langle \frac{3}{25}, -\frac{4}{25}\right\rangle$, and a unit vector in the direction of $\mathbf{v} = \langle 3,5\rangle$ is $\mathbf{u} = \frac{1}{\sqrt{9+25}}\langle 3,5\rangle = \left\langle \frac{3}{\sqrt{34}}, \frac{5}{\sqrt{34}}\right\rangle$, so

$D_{\mathbf{u}} f(1,2) = \nabla f(1,2)\cdot\mathbf{u} = \left\langle \frac{3}{25}, -\frac{4}{25}\right\rangle\cdot\left\langle \frac{3}{\sqrt{34}}, \frac{5}{\sqrt{34}}\right\rangle = \frac{9}{25\sqrt{34}} - \frac{20}{25\sqrt{34}} = -\frac{11}{25\sqrt{34}}.$

13. $g(s,t) = s\sqrt{t} \;\Rightarrow\; \nabla g(s,t) = \left(\sqrt{t}\right)\mathbf{i} + \left(s/(2\sqrt{t})\right)\mathbf{j}$, $\nabla g(2,4) = 2\,\mathbf{i} + \frac{1}{2}\,\mathbf{j}$, and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{2^2+(-1)^2}}(2\,\mathbf{i} - \mathbf{j}) = \frac{1}{\sqrt{5}}(2\,\mathbf{i} - \mathbf{j})$, so $D_{\mathbf{u}}\, g(2,4) = \nabla g(2,4)\cdot\mathbf{u} = (2\,\mathbf{i} + \frac{1}{2}\,\mathbf{j})\cdot\frac{1}{\sqrt{5}}(2\,\mathbf{i} - \mathbf{j}) = \frac{1}{\sqrt{5}}\left(4 - \frac{1}{2}\right) = \frac{7}{2\sqrt{5}}$ or $\frac{7\sqrt{5}}{10}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $g(u,v) = u^2e^{-v} \Rightarrow \nabla g(u,v) = \left(2ue^{-v}\right)\mathbf{i} + \left(-u^2e^{-v}\right)\mathbf{j}$, $\nabla g(3,0) = 6\,\mathbf{i} - 9\,\mathbf{j}$, and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{3^2+4^2}}(3\,\mathbf{i}+4\,\mathbf{j}) = \frac{1}{5}(3\,\mathbf{i}+4\,\mathbf{j})$, so $D_{\mathbf{u}}\,g(3,0) = \nabla g(3,0)\cdot\mathbf{u} = (6\,\mathbf{i}-9\,\mathbf{j})\cdot\frac{1}{5}(3\,\mathbf{i}+4\,\mathbf{j}) = \frac{1}{5}(18-36) = -\frac{18}{5}$.

15. $f(x,y,z) = x^2y + y^2z \Rightarrow \nabla f(x,y,z) = \langle 2xy, x^2+2yz, y^2\rangle$, $\nabla f(1,2,3) = \langle 4,13,4\rangle$, and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{4+1+4}}\langle 2,-1,2\rangle = \frac{1}{3}\langle 2,-1,2\rangle$, so

$D_{\mathbf{u}}\,f(1,2,3) = \nabla f(1,2,3)\cdot\mathbf{u} = \langle 4,13,4\rangle\cdot\frac{1}{3}\langle 2,-1,2\rangle = \frac{1}{3}(8-13+8) = \frac{3}{3} = 1$.

16. $f(x,y,z) = xy^2\tan^{-1}z \Rightarrow \nabla f(x,y,z) = \left\langle y^2\tan^{-1}z, 2xy\tan^{-1}z, \frac{xy^2}{1+z^2}\right\rangle$,

$\nabla f(2,1,1) = \left\langle 1\cdot\frac{\pi}{4}, 4\cdot\frac{\pi}{4}, \frac{2}{1+1}\right\rangle = \left\langle\frac{\pi}{4},\pi,1\right\rangle$, and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{1+1+1}}\langle 1,1,1\rangle = \frac{1}{\sqrt{3}}\langle 1,1,1\rangle$,

so $D_{\mathbf{u}}\,f(2,1,1) = \nabla f(2,1,1)\cdot\mathbf{u} = \left\langle\frac{\pi}{4},\pi,1\right\rangle\cdot\frac{1}{\sqrt{3}}\langle 1,1,1\rangle = \frac{1}{\sqrt{3}}\left(\frac{\pi}{4}+\pi+1\right) = \frac{1}{\sqrt{3}}\left(\frac{5\pi}{4}+1\right)$.

17. $h(r,s,t) = \ln(3r+6s+9t) \Rightarrow \nabla h(r,s,t) = \langle 3/(3r+6s+9t), 6/(3r+6s+9t), 9/(3r+6s+9t)\rangle$,

$\nabla h(1,1,1) = \left\langle\frac{1}{6},\frac{1}{3},\frac{1}{2}\right\rangle$, and a unit vector in the direction of $\mathbf{v} = 4\,\mathbf{i}+12\,\mathbf{j}+6\,\mathbf{k}$

is $\mathbf{u} = \frac{1}{\sqrt{16+144+36}}(4\,\mathbf{i}+12\,\mathbf{j}+6\,\mathbf{k}) = \frac{2}{7}\,\mathbf{i}+\frac{6}{7}\,\mathbf{j}+\frac{3}{7}\,\mathbf{k}$, so

$D_{\mathbf{u}}\,h(1,1,1) = \nabla h(1,1,1)\cdot\mathbf{u} = \left\langle\frac{1}{6},\frac{1}{3},\frac{1}{2}\right\rangle\cdot\left\langle\frac{2}{7},\frac{6}{7},\frac{3}{7}\right\rangle = \frac{1}{21}+\frac{2}{7}+\frac{3}{14} = \frac{23}{42}$.

18. $D_{\mathbf{u}}f(2,2) = \nabla f(2,2)\cdot\mathbf{u}$, the scalar projection of $\nabla f(2,2)$ onto $\mathbf{u}$, so we draw a perpendicular from the tip of $\nabla f(2,2)$ to the line containing $\mathbf{u}$. We can use the point $(2,2)$ to determine the scale of the axes, and we estimate the length of the projection to be approximately 3.0 units. Since the angle between $\nabla f(2,2)$ and $\mathbf{u}$ is greater than 90°, the scalar projection is negative. Thus $D_{\mathbf{u}}\,f(2,2) \approx -3$.



19. $f(x,y) = \sqrt{xy} \Rightarrow \nabla f(x,y) = \left\langle\frac{1}{2}(xy)^{-1/2}(y), \frac{1}{2}(xy)^{-1/2}(x)\right\rangle = \left\langle\frac{y}{2\sqrt{xy}}, \frac{x}{2\sqrt{xy}}\right\rangle$, so $\nabla f(2,8) = \left\langle 1,\frac{1}{4}\right\rangle$.

The unit vector in the direction of $\overrightarrow{PQ} = \langle 5-2, 4-8\rangle = \langle 3,-4\rangle$ is $\mathbf{u} = \left\langle\frac{3}{5},-\frac{4}{5}\right\rangle$, so

$D_{\mathbf{u}}\,f(2,8) = \nabla f(2,8)\cdot\mathbf{u} = \left\langle 1,\frac{1}{4}\right\rangle\cdot\left\langle\frac{3}{5},-\frac{4}{5}\right\rangle = \frac{2}{5}$.

20. $f(x,y,z) = xy^2z^3 \Rightarrow \nabla f(x,y,z) = \langle y^2z^3, 2xyz^3, 3xy^2z^2\rangle$, so $\nabla f(2,1,1) = \langle 1,4,6\rangle$. The unit vector in the direction of $\overrightarrow{PQ} = \langle -2,-4,4\rangle$ is $\mathbf{u} = \frac{1}{\sqrt{4+16+16}}\langle -2,-4,4\rangle = \frac{1}{6}\langle -2,-4,4\rangle$, so

$D_{\mathbf{u}}\,f(2,1,1) = \nabla f(2,1,1)\cdot\mathbf{u} = \langle 1,4,6\rangle\cdot\frac{1}{6}\langle -2,-4,4\rangle = \frac{1}{6}(-2-16+24) = 1$.

21. $f(x,y) = 4y\sqrt{x} \Rightarrow \nabla f(x,y) = \left\langle 4y\cdot\frac{1}{2}x^{-1/2}, 4\sqrt{x}\right\rangle = \langle 2y/\sqrt{x}, 4\sqrt{x}\,\rangle$.

$\nabla f(4,1) = \langle 1,8\rangle$ is the direction of maximum rate of change, and the maximum rate is $|\nabla f(4,1)| = \sqrt{1+64} = \sqrt{65}$.

22. $f(s,t) = te^{st} \Rightarrow \nabla f(s,t) = \left\langle te^{st}(t), te^{st}(s)+e^{st}(1)\right\rangle = \left\langle t^2e^{st}, (st+1)e^{st}\right\rangle$.

$\nabla f(0,2) = \langle 4,1\rangle$ is the direction of maximum rate of change, and the maximum rate is $|\nabla f(0,2)| = \sqrt{16+1} = \sqrt{17}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

23. $f(x,y)=\sin(xy) \Rightarrow \nabla f(x,y)=\langle y\cos(xy), x\cos(xy)\rangle$, $\nabla f(1,0)=\langle 0,1\rangle$. Thus the maximum rate of change is $|\nabla f(1,0)|=1$ in the direction $\langle 0,1\rangle$.

24. $f(x,y,z)=x\ln(yz) \Rightarrow \nabla f(x,y,z)=\left\langle \ln(yz), x\cdot\frac{z}{yz}, x\cdot\frac{y}{yz}\right\rangle=\left\langle \ln(yz), \frac{x}{y}, \frac{x}{z}\right\rangle$, $\nabla f(1,2,\frac{1}{2})=\langle 0,\frac{1}{2},2\rangle$. Thus the maximum rate of change is $\left|\nabla f(1,2,\frac{1}{2})\right|=\sqrt{0+\frac{1}{4}+4}=\sqrt{\frac{17}{4}}=\frac{\sqrt{17}}{2}$ in the direction $\langle 0,\frac{1}{2},2\rangle$ or equivalently $\langle 0,1,4\rangle$.

25. $f(x,y,z)=x/(y+z)=x(y+z)^{-1} \Rightarrow$

$\nabla f(x,y,z)=\langle 1/(y+z), -x(y+z)^{-2}(1), -x(y+z)^{-2}(1)\rangle=\left\langle \frac{1}{y+z}, -\frac{x}{(y+z)^2}, -\frac{x}{(y+z)^2}\right\rangle$,

$\nabla f(8,1,3)=\langle \frac{1}{4}, -\frac{8}{4^2}, -\frac{8}{4^2}\rangle=\langle \frac{1}{4}, -\frac{1}{2}, -\frac{1}{2}\rangle$. Thus the maximum rate of change is

$|\nabla f(8,1,3)|=\sqrt{\frac{1}{16}+\frac{1}{4}+\frac{1}{4}}=\sqrt{\frac{9}{16}}=\frac{3}{4}$ in the direction $\langle \frac{1}{4}, -\frac{1}{2}, -\frac{1}{2}\rangle$ or equivalently $\langle 1,-2,-2\rangle$.

26. $f(p,q,r)=\arctan(pqr) \Rightarrow \nabla f(p,q,r)=\left\langle \frac{qr}{1+(pqr)^2}, \frac{pr}{1+(pqr)^2}, \frac{pq}{1+(pqr)^2}\right\rangle$, $\nabla f(1,2,1)=\langle \frac{2}{5}, \frac{1}{5}, \frac{2}{5}\rangle$. Thus the maximum rate of change is $|\nabla f(1,2,1)|=\sqrt{\frac{4}{25}+\frac{1}{25}+\frac{4}{25}}=\sqrt{\frac{9}{25}}=\frac{3}{5}$ in the direction $\langle \frac{2}{5}, \frac{1}{5}, \frac{2}{5}\rangle$ or equivalently $\langle 2,1,2\rangle$.

27. (a) As in the proof of Theorem 15, $D_{\mathbf{u}}f=|\nabla f|\cos\theta$. Since the minimum value of $\cos\theta$ is -1 occurring when $\theta=\pi$, the minimum value of $D_{\mathbf{u}}f$ is $-|\nabla f|$ occurring when $\theta=\pi$, that is when $\mathbf{u}$ is in the opposite direction of ∇f (assuming $\nabla f\neq\mathbf{0}$).

(b) $f(x,y)=x^4y-x^2y^3 \Rightarrow \nabla f(x,y)=\langle 4x^3y-2xy^3, x^4-3x^2y^2\rangle$, so f decreases fastest at the point $(2,-3)$ in the direction $-\nabla f(2,-3)=-\langle 12,-92\rangle=\langle -12,92\rangle$.

28. $f(x,y)=x^2+xy^3 \Rightarrow \nabla f(x,y)=\langle 2x+y^3, 3xy^2\rangle$ so $\nabla f(2,1)=\langle 5,6\rangle$. If $\mathbf{u}=\langle a,b\rangle$ is a unit vector in the desired direction then $D_{\mathbf{u}}f(2,1)=2 \Leftrightarrow \langle 5,6\rangle\cdot\langle a,b\rangle=2 \Leftrightarrow 5a+6b=2 \Leftrightarrow b=\frac{1}{3}-\frac{5}{6}a$. But $a^2+b^2=1 \Leftrightarrow a^2+\left(\frac{1}{3}-\frac{5}{6}a\right)^2=1 \Leftrightarrow \frac{61}{36}a^2-\frac{5}{9}a+\frac{1}{9}=1 \Leftrightarrow 61a^2-20a-32=0$. By the quadratic formula, the solutions are $a=\dfrac{-(-20)\pm\sqrt{(-20)^2-4(61)(-32)}}{2(61)}=\dfrac{20\pm\sqrt{8208}}{122}=\dfrac{10\pm 6\sqrt{57}}{61}$. If $a=\dfrac{10+6\sqrt{57}}{61}\approx 0.9065$ then $b=\dfrac{1}{3}-\dfrac{5}{6}\left(\dfrac{10+6\sqrt{57}}{61}\right)\approx -0.4221$, and if $a=\dfrac{10-6\sqrt{57}}{61}\approx -0.5787$ then $b=\dfrac{1}{3}-\dfrac{5}{6}\left(\dfrac{10-6\sqrt{57}}{61}\right)\approx 0.8156$. Thus the two directions giving a directional derivative of 2 are approximately $\langle 0.9065, -0.4221\rangle$ and $\langle -0.5787, 0.8156\rangle$.

29. The direction of fastest change is $\nabla f(x,y)=(2x-2)\,\mathbf{i}+(2y-4)\,\mathbf{j}$, so we need to find all points (x,y) where $\nabla f(x,y)$ is parallel to $\mathbf{i}+\mathbf{j} \Leftrightarrow (2x-2)\,\mathbf{i}+(2y-4)\,\mathbf{j}=k\,(\mathbf{i}+\mathbf{j}) \Leftrightarrow k=2x-2$ and $k=2y-4$. Then $2x-2=2y-4 \Rightarrow y=x+1$, so the direction of fastest change is $\mathbf{i}+\mathbf{j}$ at all points on the line $y=x+1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. The fisherman is traveling in the direction $\langle -80, -60\rangle$. A unit vector in this direction is $\mathbf{u} = \frac{1}{100}\langle -80, -60\rangle = \left\langle -\frac{4}{5}, -\frac{3}{5}\right\rangle$, and if the depth of the lake is given by $f(x, y) = 200 + 0.02x^2 - 0.001y^3$, then $\nabla f(x, y) = \left\langle 0.04x, -0.003y^2\right\rangle$.

$D_{\mathbf{u}} f(80, 60) = \nabla f(80, 60) \cdot \mathbf{u} = \langle 3.2, -10.8\rangle \cdot \left\langle -\frac{4}{5}, -\frac{3}{5}\right\rangle = 3.92$. Since $D_{\mathbf{u}} f(80, 60)$ is positive, the depth of the lake is increasing near $(80, 60)$ in the direction toward the buoy.

31. $T = \dfrac{k}{\sqrt{x^2 + y^2 + z^2}}$ and $120 = T(1, 2, 2) = \dfrac{k}{3}$ so $k = 360$.

(a) $\mathbf{u} = \dfrac{\langle 1, -1, 1\rangle}{\sqrt{3}}$,

$D_{\mathbf{u}} T(1, 2, 2) = \nabla T(1, 2, 2) \cdot \mathbf{u} = \left[-360\left(x^2 + y^2 + z^2\right)^{-3/2}\langle x, y, z\rangle\right]_{(1,2,2)} \cdot \mathbf{u} = -\frac{40}{3}\langle 1, 2, 2\rangle \cdot \frac{1}{\sqrt{3}}\langle 1, -1, 1\rangle = -\frac{40}{3\sqrt{3}}$

(b) From (a), $\nabla T = -360\left(x^2 + y^2 + z^2\right)^{-3/2}\langle x, y, z\rangle$, and since $\langle x, y, z\rangle$ is the position vector of the point (x, y, z), the vector $-\langle x, y, z\rangle$, and thus ∇T, always points toward the origin.

32. $\nabla T = -400e^{-x^2 - 3y^2 - 9z^2}\langle x, 3y, 9z\rangle$

(a) $\mathbf{u} = \frac{1}{\sqrt{6}}\langle 1, -2, 1\rangle$, $\nabla T(2, -1, 2) = -400e^{-43}\langle 2, -3, 18\rangle$ and

$$D_{\mathbf{u}} T(2, -1, 2) = \left(-\frac{400e^{-43}}{\sqrt{6}}\right)(26) = -\frac{5200\sqrt{6}}{3e^{43}} \ ^\circ\text{C/m}.$$

(b) $\nabla T(2, -1, 2) = 400e^{-43}\langle -2, 3, -18\rangle$ or equivalently $\langle -2, 3, -18\rangle$.

(c) $|\nabla T| = 400e^{-x^2 - 3y^2 - 9z^2}\sqrt{x^2 + 9y^2 + 81z^2}$ °C/m is the maximum rate of increase. At $(2, -1, 2)$ the maximum rate of increase is $400e^{-43}\sqrt{337}$ °C/m.

33. $\nabla V(x, y, z) = \langle 10x - 3y + yz, xz - 3x, xy\rangle$, $\nabla V(3, 4, 5) = \langle 38, 6, 12\rangle$

(a) $D_{\mathbf{u}} V(3, 4, 5) = \langle 38, 6, 12\rangle \cdot \frac{1}{\sqrt{3}}\langle 1, 1, -1\rangle = \frac{32}{\sqrt{3}}$

(b) $\nabla V(3, 4, 5) = \langle 38, 6, 12\rangle$, or equivalently, $\langle 19, 3, 6\rangle$.

(c) $|\nabla V(3, 4, 5)| = \sqrt{38^2 + 6^2 + 12^2} = \sqrt{1624} = 2\sqrt{406}$

34. $z = f(x, y) = 1000 - 0.005x^2 - 0.01y^2 \Rightarrow \nabla f(x, y) = \langle -0.01x, -0.02y\rangle$ and $\nabla f(60, 40) = \langle -0.6, -0.8\rangle$.

(a) Due south is in the direction of the unit vector $\mathbf{u} = -\mathbf{j}$ and $D_{\mathbf{u}} f(60, 40) = \nabla f\,(60, 40) \cdot \langle 0, -1\rangle = \langle -0.6, -0.8\rangle \cdot \langle 0, -1\rangle = 0.8$. Thus, if you walk due south from $(60, 40, 966)$ you will ascend at a rate of 0.8 vertical meters per horizontal meter.

(b) Northwest is in the direction of the unit vector $\mathbf{u} = \frac{1}{\sqrt{2}}\langle -1, 1\rangle$ and $D_{\mathbf{u}} f(60, 40) = \nabla f\,(60, 40) \cdot \frac{1}{\sqrt{2}}\langle -1, 1\rangle = \langle -0.6, -0.8\rangle \cdot \frac{1}{\sqrt{2}}\langle -1, 1\rangle = -\frac{0.2}{\sqrt{2}} \approx -0.14$. Thus, if you walk northwest from $(60, 40, 966)$ you will descend at a rate of approximately 0.14 vertical meters per horizontal meter.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c) $\nabla f(60, 40) = \langle -0.6, -0.8 \rangle$ is the direction of largest slope with a rate of ascent given by $|\nabla f(60, 40)| = \sqrt{(-0.6)^2 + (-0.8)^2} = 1$. The angle above the horizontal in which the path begins is given by $\tan\theta = 1 \quad \Rightarrow \quad \theta = 45°$.

35. A unit vector in the direction of $\overrightarrow{AB}$ is $\mathbf{i}$ and a unit vector in the direction of $\overrightarrow{AC}$ is $\mathbf{j}$. Thus $D_{\overrightarrow{AB}} f(1, 3) = f_x(1, 3) = 3$ and $D_{\overrightarrow{AC}} f(1, 3) = f_y(1, 3) = 26$. Therefore $\nabla f(1, 3) = \langle f_x(1, 3), f_y(1, 3) \rangle = \langle 3, 26 \rangle$, and by definition, $D_{\overrightarrow{AD}} f(1, 3) = \nabla f \cdot \mathbf{u}$ where $\mathbf{u}$ is a unit vector in the direction of $\overrightarrow{AD}$, which is $\left\langle \frac{5}{13}, \frac{12}{13} \right\rangle$. Therefore, $D_{\overrightarrow{AD}} f(1, 3) = \langle 3, 26 \rangle \cdot \left\langle \frac{5}{13}, \frac{12}{13} \right\rangle = 3 \cdot \frac{5}{13} + 26 \cdot \frac{12}{13} = \frac{327}{13}$.

36. The curves of steepest ascent or descent are perpendicular to all of the contour lines (see Figure 12) so we sketch curves beginning at A and B that head toward lower elevations, crossing each contour line at a right angle.



37. (a) $\nabla(au + bv) = \left\langle \frac{\partial(au + bv)}{\partial x}, \frac{\partial(au + bv)}{\partial y} \right\rangle = \left\langle a\frac{\partial u}{\partial x} + b\frac{\partial v}{\partial x}, a\frac{\partial u}{\partial y} + b\frac{\partial v}{\partial y} \right\rangle = a\left\langle \frac{\partial u}{\partial x}, \frac{\partial u}{\partial y} \right\rangle + b\left\langle \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y} \right\rangle$

$= a\,\nabla u + b\,\nabla v$

(b) $\nabla(uv) = \left\langle v\frac{\partial u}{\partial x} + u\frac{\partial v}{\partial x}, v\frac{\partial u}{\partial y} + u\frac{\partial v}{\partial y} \right\rangle = v\left\langle \frac{\partial u}{\partial x}, \frac{\partial u}{\partial y} \right\rangle + u\left\langle \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y} \right\rangle = v\,\nabla u + u\,\nabla v$

(c) $\nabla\left(\frac{u}{v}\right) = \left\langle \frac{v\frac{\partial u}{\partial x} - u\frac{\partial v}{\partial x}}{v^2}, \frac{v\frac{\partial u}{\partial y} - u\frac{\partial v}{\partial y}}{v^2} \right\rangle = \frac{v\left\langle \frac{\partial u}{\partial x}, \frac{\partial u}{\partial y} \right\rangle - u\left\langle \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y} \right\rangle}{v^2} = \frac{v\,\nabla u - u\,\nabla v}{v^2}$

(d) $\nabla u^n = \left\langle \frac{\partial(u^n)}{\partial x}, \frac{\partial(u^n)}{\partial y} \right\rangle = \left\langle nu^{n-1}\frac{\partial u}{\partial x}, nu^{n-1}\frac{\partial u}{\partial y} \right\rangle = nu^{n-1}\nabla u$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. If we place the initial point of the gradient vector $\nabla f(4,6)$ at $(4,6)$, the vector is perpendicular to the level curve of f that includes $(4,6)$, so we sketch a portion of the level curve through $(4,6)$ (using the nearby level curves as a guideline) and draw a line perpendicular to the curve at $(4,6)$. The gradient vector is parallel to this line, pointing in the direction of increasing function values, and with length equal to the maximum value of the directional derivative of f at $(4,6)$. We can estimate this length by finding the average rate of change in the direction of the gradient. The line intersects the contour lines corresponding to -2 and -3 with an estimated distance of 0.5 units. Thus the rate of change is approximately $\dfrac{-2-(-3)}{0.5}=2$, and we sketch the gradient vector with length 2.



39. $f(x,y)=x^3+5x^2y+y^3 \quad \Rightarrow$

$D_{\mathbf{u}}f(x,y)=\nabla f(x,y)\cdot\mathbf{u}=\langle 3x^2+10xy,\,5x^2+3y^2\rangle\cdot\left\langle\frac{3}{5},\frac{4}{5}\right\rangle=\frac{9}{5}x^2+6xy+4x^2+\frac{12}{5}y^2=\frac{29}{5}x^2+6xy+\frac{12}{5}y^2$. Then

$$D_{\mathbf{u}}^2f(x,y)=D_{\mathbf{u}}\left[D_{\mathbf{u}}f(x,y)\right]=\nabla\left[D_{\mathbf{u}}f(x,y)\right]\cdot\mathbf{u}=\left\langle\tfrac{58}{5}x+6y,\,6x+\tfrac{24}{5}y\right\rangle\cdot\left\langle\tfrac{3}{5},\tfrac{4}{5}\right\rangle$$
$$=\tfrac{174}{25}x+\tfrac{18}{5}y+\tfrac{24}{5}x+\tfrac{96}{25}y=\tfrac{294}{25}x+\tfrac{186}{25}y$$

and $D_{\mathbf{u}}^2f(2,1)=\frac{294}{25}(2)+\frac{186}{25}(1)=\frac{774}{25}$.

40. (a) From Equation 9 we have $D_{\mathbf{u}}f=\nabla f\cdot\mathbf{u}=\langle f_x,f_y\rangle\cdot\langle a,b\rangle=f_xa+f_yb$ and from Exercise 39 we have

$D_{\mathbf{u}}^2f=D_{\mathbf{u}}\left[D_{\mathbf{u}}f\right]=\nabla\left[D_{\mathbf{u}}f\right]\cdot\mathbf{u}=\langle f_{xx}a+f_{yx}b,\,f_{xy}a+f_{yy}b\rangle\cdot\langle a,b\rangle=f_{xx}a^2+f_{yx}ab+f_{xy}ab+f_{yy}b^2$.

But $f_{yx}=f_{xy}$ by Clairaut's Theorem, so $D_{\mathbf{u}}^2f=f_{xx}a^2+2f_{xy}ab+f_{yy}b^2$.

(b) $f(x,y)=xe^{2y} \quad \Rightarrow \quad f_x=e^{2y},\ f_y=2xe^{2y},\ f_{xx}=0,\ f_{xy}=2e^{2y},\ f_{yy}=4xe^{2y}$ and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u}=\frac{1}{\sqrt{4^2+6^2}}\langle 4,6\rangle=\left\langle\frac{2}{\sqrt{13}},\frac{3}{\sqrt{13}}\right\rangle=\langle a,b\rangle$. Then

$$D_{\mathbf{u}}^2f=f_{xx}a^2+2f_{xy}ab+f_{yy}b^2=0\cdot\left(\tfrac{2}{\sqrt{13}}\right)^2+2\cdot 2e^{2y}\left(\tfrac{2}{\sqrt{13}}\right)\left(\tfrac{3}{\sqrt{13}}\right)+4xe^{2y}\left(\tfrac{3}{\sqrt{13}}\right)^2=\tfrac{24}{13}e^{2y}+\tfrac{36}{13}xe^{2y}.$$

41. Let $F(x,y,z)=2(x-2)^2+(y-1)^2+(z-3)^2$. Then $2(x-2)^2+(y-1)^2+(z-3)^2=10$ is a level surface of F.

$F_x(x,y,z)=4(x-2) \quad \Rightarrow \quad F_x(3,3,5)=4$, $F_y(x,y,z)=2(y-1) \quad \Rightarrow \quad F_y(3,3,5)=4$, and

$F_z(x,y,z)=2(z-3) \quad \Rightarrow \quad F_z(3,3,5)=4$.

(a) Equation 19 gives an equation of the tangent plane at $(3,3,5)$ as $4(x-3)+4(y-3)+4(z-5)=0 \quad \Leftrightarrow$

$4x+4y+4z=44$ or equivalently $x+y+z=11$.

(b) By Equation 20, the normal line has symmetric equations $\dfrac{x-3}{4}=\dfrac{y-3}{4}=\dfrac{z-5}{4}$ or equivalently

$x-3=y-3=z-5$. Corresponding parametric equations are $x=3+t,\ y=3+t,\ z=5+t$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

42. Let $F(x,y,z) = y^2 + z^2 - x$. Then $x = y^2 + z^2 + 1 \Leftrightarrow y^2 + z^2 - x = -1$ is a level surface of F.

$F_x(x,y,z) = -1 \Rightarrow F_x(3,1,-1) = -1$, $F_y(x,y,z) = 2y \Rightarrow F_y(3,1,-1) = 2$, and $F_z(x,y,z) = 2z \Rightarrow F_z(3,1,-1) = -2$.

(a) By Equation 19, an equation of the tangent plane at $(3,1,-1)$ is $(-1)(x-3) + 2(y-1) + (-2)[z-(-1)] = 0$ or $-x + 2y - 2z = 1$ or $x - 2y + 2z = -1$.

(b) By Equation 20, the normal line has symmetric equations $\dfrac{x-3}{-1} = \dfrac{y-1}{2} = \dfrac{z-(-1)}{-2}$ or equivalently $x - 3 = \dfrac{y-1}{-2} = \dfrac{z+1}{2}$ and parametric equations $x = 3 - t$, $y = 1 + 2t$, $z = -1 - 2t$.

43. Let $F(x,y,z) = xy^2z^3$. Then $xy^2z^3 = 8$ is a level surface of F and $\nabla F(x,y,z) = \langle y^2z^3, 2xyz^3, 3xy^2z^2 \rangle$.

(a) $\nabla F(2,2,1) = \langle 4, 8, 24 \rangle$ is a normal vector for the tangent plane at $(2,2,1)$, so an equation of the tangent plane is $4(x-2) + 8(y-2) + 24(z-1) = 0$ or $4x + 8y + 24z = 48$ or equivalently $x + 2y + 6z = 12$.

(b) The normal line has direction $\nabla F(2,2,1) = \langle 4,8,24 \rangle$ or equivalently $\langle 1,2,6 \rangle$, so parametric equations are $x = 2 + t$, $y = 2 + 2t$, $z = 1 + 6t$, and symmetric equations are $x - 2 = \dfrac{y-2}{2} = \dfrac{z-1}{6}$.

44. Let $F(x,y,z) = xy + yz + zx$. Then $xy + yz + zx = 5$ is a level surface of F and $\nabla F(x,y,z) = \langle y+z, x+z, x+y \rangle$.

(a) $\nabla F(1,2,1) = \langle 3,2,3 \rangle$ is a normal vector for the tangent plane at $(1,2,1)$, so an equation of the tangent plane is $3(x-1) + 2(y-2) + 3(z-1) = 0$ or $3x + 2y + 3z = 10$.

(b) The normal line has direction $\langle 3,2,3 \rangle$, so parametric equations are $x = 1 + 3t$, $y = 2 + 2t$, $z = 1 + 3t$, and symmetric equations are $\dfrac{x-1}{2} = \dfrac{y-2}{1} = \dfrac{z-1}{3}$.

45. Let $F(x,y,z) = x + y + z - e^{xyz}$. Then $x + y + z = e^{xyz}$ is the level surface $F(x,y,z) = 0$, and $\nabla F(x,y,z) = \langle 1 - yze^{xyz}, 1 - xze^{xyz}, 1 - xye^{xyz} \rangle$.

(a) $\nabla F(0,0,1) = \langle 1,1,1 \rangle$ is a normal vector for the tangent plane at $(0,0,1)$, so an equation of the tangent plane is $1(x-0) + 1(y-0) + 1(z-1) = 0$ or $x + y + z = 1$.

(b) The normal line has direction $\langle 1,1,1 \rangle$, so parametric equations are $x = t$, $y = t$, $z = 1 + t$, and symmetric equations are $x = y = z - 1$.

46. Let $F(x,y,z) = x^4 + y^4 + z^4 - 3x^2y^2z^2$. Then $x^4 + y^4 + z^4 = 3x^2y^2z^2$ is the level surface $F(x,y,z) = 0$, and $\nabla F(x,y,z) = \langle 4x^3 - 6xy^2z^2, 4y^3 - 6x^2yz^2, 4z^3 - 6x^2y^2z \rangle$.

(a) $\nabla F(1,1,1) = \langle -2,-2,-2 \rangle$ or equivalently $\langle 1,1,1 \rangle$ is a normal vector for the tangent plane at $(1,1,1)$, so an equation of the tangent plane is $1(x-1) + 1(y-1) + 1(z-1) = 0$ or $x + y + z = 3$.

(b) The normal line has direction $\langle 1,1,1 \rangle$, so parametric equations are $x = 1 + t$, $y = 1 + t$, $z = 1 + t$, and symmetric equations are $x - 1 = y - 1 = z - 1$ or equivalently $x = y = z$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $F(x,y,z) = xy + yz + zx$,

$\nabla F(x,y,z) = \langle y+z, x+z, y+x \rangle$,

$\nabla F(1,1,1) = \langle 2,2,2 \rangle$, so an equation of the tangent plane is $2x + 2y + 2z = 6$ or $x + y + z = 3$, and the normal line is given by $x - 1 = y - 1 = z - 1$ or $x = y = z$. To graph the surface we solve for z:

$z = \dfrac{3 - xy}{x + y}$.



48. $F(x,y,z) = xyz$, $\nabla F(x,y,z) = \langle yz, xz, yx \rangle$,

$\nabla F(1,2,3) = \langle 6,3,2 \rangle$, so an equation of the tangent plane is $6x + 3y + 2z = 18$, and the normal line is given by $\dfrac{x-1}{6} = \dfrac{y-2}{3} = \dfrac{z-3}{2}$ or $x = 1 + 6t$, $y = 2 + 3t$, $z = 3 + 2t$. To graph the surface we solve for z: $z = \dfrac{6}{xy}$.



49. $f(x,y) = xy \;\Rightarrow\; \nabla f(x,y) = \langle y, x \rangle$, $\nabla f(3,2) = \langle 2,3 \rangle$. $\nabla f(3,2)$ is perpendicular to the tangent line, so the tangent line has equation $\nabla f(3,2) \cdot \langle x-3, y-2 \rangle = 0 \;\Rightarrow\; \langle 2,3 \rangle \cdot \langle x-3, x-2 \rangle = 0 \;\Rightarrow\; 2(x-3) + 3(y-2) = 0$ or $2x + 3y = 12$.



50. $g(x,y) = x^2 + y^2 - 4x \;\Rightarrow\; \nabla g(x,y) = \langle 2x-4, 2y \rangle$,

$\nabla g(1,2) = \langle -2,4 \rangle$. $\nabla g(1,2)$ is perpendicular to the tangent line, so the tangent line has equation $\nabla g(1,2) \cdot \langle x-1, y-2 \rangle = 0 \;\Rightarrow\; \langle -2,4 \rangle \cdot \langle x-1, y-2 \rangle = 0 \;\Rightarrow\; -2(x-1) + 4(y-2) = 0 \;\Leftrightarrow\; -2x + 4y = 6$ or equivalently $-x + 2y = 3$.



51. $\nabla F(x_0,y_0,z_0) = \left\langle \dfrac{2x_0}{a^2}, \dfrac{2y_0}{b^2}, \dfrac{2z_0}{c^2} \right\rangle$. Thus an equation of the tangent plane at (x_0,y_0,z_0) is

$\dfrac{2x_0}{a^2}x + \dfrac{2y_0}{b^2}y + \dfrac{2z_0}{c^2}z = 2\left(\dfrac{x_0^2}{a^2} + \dfrac{y_0^2}{b^2} + \dfrac{z_0^2}{c^2}\right) = 2(1) = 2$ since (x_0,y_0,z_0) is a point on the ellipsoid. Hence

$\dfrac{x_0}{a^2}x + \dfrac{y_0}{b^2}y + \dfrac{z_0}{c^2}z = 1$ is an equation of the tangent plane.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

52. $\nabla F(x_0, y_0, z_0) = \left\langle \frac{2x_0}{a^2}, \frac{2y_0}{b^2}, \frac{-2z_0}{c^2} \right\rangle$, so an equation of the tangent plane at (x_0, y_0, z_0) is

$\frac{2x_0}{a^2}\,x + \frac{2y_0}{b^2}\,y - \frac{2z_0}{c^2}\,z = 2\left(\frac{x_0^2}{a^2} + \frac{y_0^2}{b^2} - \frac{z_0^2}{c^2}\right) = 2$ or $\frac{x_0}{a^2}\,x + \frac{y_0}{b^2}\,y - \frac{z_0}{c^2}\,z = 1$.

53. $\nabla F(x_0, y_0, z_0) = \left\langle \frac{2x_0}{a^2}, \frac{2y_0}{b^2}, \frac{-1}{c} \right\rangle$, so an equation of the tangent plane is $\frac{2x_0}{a^2}\,x + \frac{2y_0}{b^2}\,y - \frac{1}{c}\,z = \frac{2x_0^2}{a^2} + \frac{2y_0^2}{b^2} - \frac{z_0}{c}$

or $\frac{2x_0}{a^2}\,x + \frac{2y_0}{b^2}\,y = \frac{z}{c} + 2\left(\frac{x_0^2}{a^2} + \frac{y_0^2}{b^2}\right) - \frac{z_0}{c}$. But $\frac{z_0}{c} = \frac{x_0^2}{a^2} + \frac{y_0^2}{b^2}$, so the equation can be written as

$\frac{2x_0}{a^2}\,x + \frac{2y_0}{b^2}\,y = \frac{z + z_0}{c}$.

54. Let $F(x, y, z) = x^2 + y^2 + 2z^2$; then the ellipsoid $x^2 + y^2 + 2z^2 = 1$ is a level surface of F. $\nabla F(x, y, z) = \langle 2x, 2y, 4z \rangle$ is a normal vector to the surface at (x, y, z) and so it is a normal vector for the tangent plane there. The tangent plane is parallel to the plane $x + 2y + z = 1$ when the normal vectors of the planes are parallel, so we need a point (x_0, y_0, z_0) on the ellipsoid where $\langle 2x_0, 2y_0, 4z_0 \rangle = k\,\langle 1, 2, 1 \rangle$ for some $k \neq 0$. Comparing components we have $2x_0 = k \;\Rightarrow\; x_0 = k/2$, $2y_0 = 2k \;\Rightarrow\; y_0 = k$, $4z_0 = k \;\Rightarrow\; z_0 = k/4$. $(x_0, y_0, z_0) = (k/2, k, k/4)$ lies on the ellipsoid, so $(k/2)^2 + k^2 + 2(k/4)^2 = 1 \;\Rightarrow\; \frac{11}{8}k^2 = 1 \;\Rightarrow\; k^2 = \frac{8}{11} \;\Rightarrow\; k = \pm 2\sqrt{\frac{2}{11}}$. Thus the tangent planes at the points $\left(\sqrt{\frac{2}{11}}, 2\sqrt{\frac{2}{11}}, \frac{1}{2}\sqrt{\frac{2}{11}}\right)$ and $\left(-\sqrt{\frac{2}{11}}, -2\sqrt{\frac{2}{11}}, -\frac{1}{2}\sqrt{\frac{2}{11}}\right)$ are parallel to the given plane.

55. The hyperboloid $x^2 - y^2 - z^2 = 1$ is a level surface of $F(x, y, z) = x^2 - y^2 - z^2$ and $\nabla F\,(x, y, z) = \langle 2x, -2y, -2z \rangle$ is a normal vector to the surface and hence a normal vector for the tangent plane at (x, y, z). The tangent plane is parallel to the plane $z = x + y$ or $x + y - z = 0$ if and only if the corresponding normal vectors are parallel, so we need a point (x_0, y_0, z_0) on the hyperboloid where $\langle 2x_0, -2y_0, -2z_0 \rangle = c\,\langle 1, 1, -1 \rangle$ or equivalently $\langle x_0, -y_0, -z_0 \rangle = k\,\langle 1, 1, -1 \rangle$ for some $k \neq 0$. Then we must have $x_0 = k$, $y_0 = -k$, $z_0 = k$ and substituting into the equation of the hyperboloid gives $k^2 - (-k)^2 - k^2 = 1 \;\Leftrightarrow\; -k^2 = 1$, an impossibility. Thus there is no such point on the hyperboloid.

56. First note that the point $(1, 1, 2)$ is on both surfaces. The ellipsoid is a level surface of $F(x, y, z) = 3x^2 + 2y^2 + z^2$ and $\nabla F(x, y, z) = \langle 6x, 4y, 2z \rangle$. A normal vector to the surface at $(1, 1, 2)$ is $\nabla F(1, 1, 2) = \langle 6, 4, 4 \rangle$ and an equation of the tangent plane there is $6(x - 1) + 4(y - 1) + 4(z - 2) = 0$ or $6x + 4y + 4z = 18$ or $3x + 2y + 2z = 9$. The sphere is a level surface of $G(x, y, z) = x^2 + y^2 + z^2 - 8x - 6y - 8z + 24$ and $\nabla G(x, y, z) = \langle 2x - 8, 2y - 6, 2z - 8 \rangle$. A normal vector to the sphere at $(1, 1, 2)$ is $\nabla G(1, 1, 2) = \langle -6, -4, -4 \rangle$ and the tangent plane there is $-6(x - 1) - 4(y - 1) - 4(z - 2) = 0$ or $3x + 2y + 2z = 9$. Since these tangent planes are identical, the surfaces are tangent to each other at the point $(1, 1, 2)$.

57. Let (x_0, y_0, z_0) be a point on the cone [other than $(0, 0, 0)$]. The cone is a level surface of $F(x, y, z) = x^2 + y^2 - z^2$ and $\nabla F(x, y, z) = \langle 2x, 2y, -2z \rangle$, so $\nabla F(x_0, y_0, z_0) = \langle 2x_0, 2y_0, -2z_0 \rangle$ is a normal vector to the cone at this point and an

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

equation of the tangent plane there is $2x_0(x-x_0)+2y_0(y-y_0)-2z_0(z-z_0)=0$ or $x_0x+y_0y-z_0z=x_0^2+y_0^2-z_0^2$. But $x_0^2+y_0^2=z_0^2$ so the tangent plane is given by $x_0x+y_0y-z_0z=0$, a plane which always contains the origin.

58. Let (x_0,y_0,z_0) be a point on the sphere. Then the normal line is given by $\dfrac{x-x_0}{2x_0}=\dfrac{y-y_0}{2y_0}=\dfrac{z-z_0}{2z_0}$. For the center $(0,0,0)$ to be on the line, we need $-\dfrac{x_0}{2x_0}=-\dfrac{y_0}{2y_0}=-\dfrac{z_0}{2z_0}$ or equivalently $1=1=1$, which is true.

59. Let $F(x,y,z)=x^2+y^2-z$. Then the paraboloid is the level surface $F(x,y,z)=0$ and $\nabla F(x,y,z)=\langle 2x,2y,-1\rangle$, so $\nabla F(1,1,2)=\langle 2,2,-1\rangle$ is a normal vector to the surface. Thus the normal line at $(1,1,2)$ is given by $x=1+2t$, $y=1+2t$, $z=2-t$. Substitution into the equation of the paraboloid $z=x^2+y^2$ gives $2-t=(1+2t)^2+(1+2t)^2 \;\Leftrightarrow\; 2-t=2+8t+8t^2 \;\Leftrightarrow\; 8t^2+9t=0 \;\Leftrightarrow\; t(8t+9)=0$. Thus the line intersects the paraboloid when $t=0$, corresponding to the given point $(1,1,2)$, or when $t=-\frac{9}{8}$, corresponding to the point $\left(-\frac{5}{4},-\frac{5}{4},\frac{25}{8}\right)$.

60. The ellipsoid is a level surface of $F(x,y,z)=4x^2+y^2+4z^2$ and $\nabla F(x,y,z)=\langle 8x,2y,8z\rangle$, so $\nabla F(1,2,1)=\langle 8,4,8\rangle$ or equivalently $\langle 2,1,2\rangle$ is a normal vector to the surface. Thus the normal line to the ellipsoid at $(1,2,1)$ is given by $x=1+2t$, $y=2+t$, $z=1+2t$. Substitution into the equation of the sphere gives $(1+2t)^2+(2+t)^2+(1+2t)^2=102 \;\Leftrightarrow\; 6+12t+9t^2=102 \;\Leftrightarrow\; 9t^2+12t-96=0 \;\Leftrightarrow\; 3(t+4)(3t-8)=0$. Thus the line intersects the sphere when $t=-4$, corresponding to the point $(-7,-2,-7)$, and when $t=\frac{8}{3}$, corresponding to the point $\left(\frac{19}{3},\frac{14}{3},\frac{19}{3}\right)$.

61. Let (x_0,y_0,z_0) be a point on the surface. Then an equation of the tangent plane at the point is $\dfrac{x}{2\sqrt{x_0}}+\dfrac{y}{2\sqrt{y_0}}+\dfrac{z}{2\sqrt{z_0}}=\dfrac{\sqrt{x_0}+\sqrt{y_0}+\sqrt{z_0}}{2}$. But $\sqrt{x_0}+\sqrt{y_0}+\sqrt{z_0}=\sqrt{c}$, so the equation is $\dfrac{x}{\sqrt{x_0}}+\dfrac{y}{\sqrt{y_0}}+\dfrac{z}{\sqrt{z_0}}=\sqrt{c}$. The x-, y-, and z-intercepts are $\sqrt{cx_0}$, $\sqrt{cy_0}$ and $\sqrt{cz_0}$ respectively. (The x-intercept is found by setting $y=z=0$ and solving the resulting equation for x, and the y- and z-intercepts are found similarly.) So the sum of the intercepts is $\sqrt{c}\left(\sqrt{x_0}+\sqrt{y_0}+\sqrt{z_0}\right)=c$, a constant.

62. The surface $xyz=1$ is a level surface of $F(x,y,z)=xyz$ and $\nabla F(x,y,z)=\langle yz,xz,xy\rangle$ is normal to the surface, so a normal vector for the tangent plane to the surface at (x_0,y_0,z_0) is $\langle y_0z_0,x_0z_0,x_0y_0\rangle$. An equation for the tangent plane there is $y_0z_0(x-x_0)+x_0z_0(y-y_0)+x_0y_0(z-z_0)=0 \;\Rightarrow\; y_0z_0x+x_0z_0y+x_0y_0z=3x_0y_0z_0$ or $\dfrac{x}{x_0}+\dfrac{y}{y_0}+\dfrac{z}{z_0}=3$. If (x_0,y_0,z_0) is in the first octant, then the tangent plane cuts off a pyramid in the first octant with vertices $(0,0,0)$, $(3x_0,0,0)$, $(0,3y_0,0)$, $(0,0,3z_0)$. The base in the xy-plane is a triangle with area $\frac{1}{2}(3x_0)(3y_0)$ and the height (along the z-axis) of the pyramid is $3z_0$. The volume of the pyramid for any point (x_0,y_0,z_0) on the surface $xyz=1$ in the first octant is $\frac{1}{3}$ (base) (height) $=\frac{1}{3}\cdot\frac{1}{2}(3x_0)(3y_0)\cdot 3z_0=\frac{9}{2}x_0y_0z_0=\frac{9}{2}$ since $x_0y_0z_0=1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

63. If $f(x,y,z) = z - x^2 - y^2$ and $g(x,y,z) = 4x^2 + y^2 + z^2$, then the tangent line is perpendicular to both ∇f and ∇g at $(-1,1,2)$. The vector $\mathbf{v} = \nabla f \times \nabla g$ will therefore be parallel to the tangent line.

We have $\nabla f(x,y,z) = \langle -2x, -2y, 1\rangle \Rightarrow \nabla f(-1,1,2) = \langle 2,-2,1\rangle$, and $\nabla g(x,y,z) = \langle 8x, 2y, 2z\rangle \Rightarrow$

$\nabla g(-1,1,2) = \langle -8,2,4\rangle$. Hence $\mathbf{v} = \nabla f \times \nabla g = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & 1 \\ -8 & 2 & 4 \end{vmatrix} = -10\,\mathbf{i} - 16\,\mathbf{j} - 12\,\mathbf{k}$.

Parametric equations are: $x = -1 - 10t,\ y = 1 - 16t,\ z = 2 - 12t$.

64. (a) Let $f(x,y,z) = y + z$ and $g(x,y,z) = x^2 + y^2$. Then the required tangent line is perpendicular to both ∇f and ∇g at $(1,2,1)$ and the vector $\mathbf{v} = \nabla f \times \nabla g$ is parallel to the tangent line. We have

$\nabla f(x,y,z) = \langle 0,1,1\rangle \Rightarrow \nabla f(1,2,1) = \langle 0,1,1\rangle$, and

$\nabla g(x,y,z) = \langle 2x, 2y, 0\rangle \Rightarrow \nabla g(1,2,1) = \langle 2,4,0\rangle$. Hence

$\mathbf{v} = \nabla f \times \nabla g = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & 1 \\ 2 & 4 & 0 \end{vmatrix} = -4\,\mathbf{i} + 2\,\mathbf{j} - 2\,\mathbf{k}$. So parametric equations of the desired tangent line are $x = 1 - 4t,\ y = 2 + 2t,\ z = 1 - 2t$.

(b)



65. Parametric equations for the helix are $x = \cos \pi t,\ y = \sin \pi t,\ z = t$, and substituting into the equation of the paraboloid gives $t = \cos^2 \pi t + \sin^2 \pi t \Rightarrow t = 1$. Thus the helix intersects the surface at the point $(\cos \pi, \sin \pi, 1) = (-1,0,1)$. Here $\mathbf{r}'(t) = \langle -\pi \sin \pi t, \pi \cos \pi t, 1\rangle$, so the tangent vector to the helix at that point is $\mathbf{r}'(1) = \langle -\pi \sin \pi, \pi \cos \pi, 1\rangle = \langle 0, -\pi, 1\rangle$. The paraboloid $z = x^2 + y^2 \Leftrightarrow x^2 + y^2 - z = 0$ is a level surface of $F(x,y,z) = x^2 + y^2 - z$ and $\nabla F(x,y,z) = \langle 2x, 2y, -1\rangle$, so a normal vector to the tangent plane at $(-1,0,1)$ is $\nabla F(-1,0,1) = \langle -2,0,-1\rangle$. The angle θ between $\mathbf{r}'(1)$ and $\nabla F(-1,0,1)$ is given by

$$\cos\theta = \frac{\langle 0,-\pi,1\rangle \cdot \langle -2,0,-1\rangle}{|\langle 0,-\pi,1\rangle|\ |\langle -2,0,-1\rangle|} = \frac{0+0-1}{\sqrt{0+\pi^2+1}\sqrt{4+0+1}} = \frac{-1}{\sqrt{5(\pi^2+1)}} \Rightarrow$$

$\theta = \cos^{-1} \dfrac{-1}{\sqrt{5(\pi^2+1)}} \approx 97.8°$. Because $\nabla F(-1,0,1)$ is perpendicular to the tangent plane, the angle of intersection between the helix and the paraboloid is approximately $97.8° - 90° = 7.8°$.

66. Parametric equations for the helix are $x = \cos(\pi t/2),\ y = \sin(\pi t/2),\ z = t$, and substituting into the equation of the sphere gives $\cos^2(\pi t/2) + \sin^2(\pi t/2) + t^2 = 2 \Rightarrow 1 + t^2 = 2 \Rightarrow t = \pm 1$. Thus the helix intersects the sphere at two points: $(\cos(\pi/2), \sin(\pi/2), 1) = (0,1,1)$, when $t = 1$, and $(\cos(-\pi/2), \sin(-\pi/2), -1) = (0,-1,-1)$, when $t = -1$. Here $\mathbf{r}'(t) = \langle -\frac{\pi}{2}\sin(\pi t/2), \frac{\pi}{2}\cos(\pi t/2), 1\rangle$, so the tangent vector to the helix at $(0,1,1)$ is $\mathbf{r}'(1) = \langle -\pi/2, 0, 1\rangle$. The sphere $x^2 + y^2 + z^2 = 2$ is a level surface of $F(x,y,z) = x^2 + y^2 + z^2$ and $\nabla F(x,y,z) = \langle 2x, 2y, 2z\rangle$, so a normal

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

vector to the tangent plane at $(0, 1, 1)$ is $\nabla F(0, 1, 1) = \langle 0, 2, 2\rangle$. As in Exercise 65, the angle of intersection between the helix and the sphere is the angle between the tangent vector to the helix and the tangent plane to the sphere. The angle θ between $\mathbf{r}'(1)$ and $\nabla F(0, 1, 1)$ is given by

$$\cos\theta = \frac{\langle -\pi/2, 0, 1\rangle \cdot \langle 0, 2, 2\rangle}{|\langle -\pi/2, 0, 1\rangle|\ |\langle 0, 2, 2\rangle|} = \frac{2}{\sqrt{(\pi^2/4)+1}\sqrt{8}} = \frac{2}{\sqrt{2\pi^2+8}} \quad \Rightarrow \quad \theta = \cos^{-1}\frac{2}{\sqrt{2\pi^2+8}} \approx 67.7^\circ$$

Because $\nabla F(0, 1, 1)$ is perpendicular to the tangent plane, the angle between $\mathbf{r}'(1)$ and the tangent plane is approximately $90^\circ - 67.7^\circ = 22.3^\circ$.

At $(0, -1, -1)$, $\mathbf{r}'(-1) = \langle \pi/2, 0, 1\rangle$ and $\nabla F(0, -1, -1) = \langle 0, -2, -2\rangle$, and the angle ϕ between these vectors is given by $\cos\phi = \dfrac{\langle \pi/2, 0, 1\rangle \cdot \langle 0, -2, -2\rangle}{|\langle \pi/2, 0, 1\rangle|\ |\langle 0, -2, -2\rangle|} = \dfrac{-2}{\sqrt{2\pi^2+8}} \quad \Rightarrow \quad \phi = \cos^{-1}\dfrac{-2}{\sqrt{2\pi^2+8}} \approx 112.3^\circ$. Thus the angle between the helix and the sphere at $(0, -1, -1)$ is approximately $112.3^\circ - 90^\circ = 22.3^\circ$. (By symmetry, we would expect the angles to be identical.)

67. (a) The direction of the normal line of F is given by ∇F, and that of G by ∇G. Assuming that $\nabla F \neq 0 \neq \nabla G$, the two normal lines are perpendicular at P if $\nabla F \cdot \nabla G = 0$ at $P \quad \Leftrightarrow$

$\langle \partial F/\partial x, \partial F/\partial y, \partial F/\partial z\rangle \cdot \langle \partial G/\partial x, \partial G/\partial y, \partial G/\partial z\rangle = 0$ at $P \quad \Leftrightarrow \quad F_x G_x + F_y G_y + F_z G_z = 0$ at P.

(b) Here $F = x^2 + y^2 - z^2$ and $G = x^2 + y^2 + z^2 - r^2$, so

$\nabla F \cdot \nabla G = \langle 2x, 2y, -2z\rangle \cdot \langle 2x, 2y, 2z\rangle = 4x^2 + 4y^2 - 4z^2 = 4F = 0$, since the point (x, y, z) lies on the graph of $F = 0$. To see that this is true without using calculus, note that $G = 0$ is the equation of a sphere centered at the origin and $F = 0$ is the equation of a right circular cone with vertex at the origin (which is generated by lines through the origin). At any point of intersection, the sphere's normal line (which passes through the origin) lies on the cone, and thus is perpendicular to the cone's normal line. So the surfaces with equations $F = 0$ and $G = 0$ are everywhere orthogonal.

68. (a) The function $f(x, y) = (xy)^{1/3}$ is continuous on $\mathbb{R}^2$ since it is a composition of a polynomial and the cube root function, both of which are continuous. (See the text just after Example 14.2.8.)

$$f_x(0,0) = \lim_{h\to 0}\frac{f(0+h,0)-f(0,0)}{h} = \lim_{h\to 0}\frac{(h\cdot 0)^{1/3}-0}{h} = 0,$$

$$f_y(0,0) = \lim_{h\to 0}\frac{f(0,0+h)-f(0,0)}{h} = \lim_{h\to 0}\frac{(0\cdot h)^{1/3}-0}{h} = 0.$$

Therefore, $f_x(0, 0)$ and $f_y(0, 0)$ do exist and are equal to 0. Now let $\mathbf{u}$ be any unit vector other than $\mathbf{i}$ and $\mathbf{j}$ (these correspond to f_x and f_y respectively.) Then $\mathbf{u} = a\,\mathbf{i} + b\,\mathbf{j}$ where $a \neq 0$ and $b \neq 0$. Thus

$D_{\mathbf{u}} f(0, 0) = \lim_{h\to 0}\dfrac{f(0+ha, 0+hb)-f(0,0)}{h} = \lim_{h\to 0}\dfrac{\sqrt[3]{(ha)(hb)}}{h} = \lim_{h\to 0}\dfrac{\sqrt[3]{ab}}{h^{1/3}}$ and this limit does not exist, so $D_{\mathbf{u}} f(0, 0)$ does not exist.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b)



Notice that if we start at the origin and proceed in the direction of the x- or y-axis, then the graph is flat. But if we proceed in any other direction, then the graph is extremely steep.

69. Let $\mathbf{u} = \langle a, b\rangle$ and $\mathbf{v} = \langle c, d\rangle$. Then we know that at the given point, $D_{\mathbf{u}} f = \nabla f \cdot \mathbf{u} = af_x + bf_y$ and $D_{\mathbf{v}} f = \nabla f \cdot \mathbf{v} = cf_x + df_y$. But these are just two linear equations in the two unknowns f_x and f_y, and since $\mathbf{u}$ and $\mathbf{v}$ are not parallel, we can solve the equations to find $\nabla f = \langle f_x, f_y\rangle$ at the given point. In fact,

$$\nabla f = \left\langle \frac{d\,D_{\mathbf{u}} f - b\,D_{\mathbf{v}} f}{ad - bc}, \frac{a\,D_{\mathbf{v}} f - c\,D_{\mathbf{u}} f}{ad - bc} \right\rangle.$$

70. Since $z = f(x, y)$ is differentiable at $\mathbf{x}_0 = (x_0, y_0)$, by Definition 14.4.7 we have $\Delta z = f_x(x_0, y_0)\,\Delta x + f_y(x_0, y_0)\,\Delta y + \varepsilon_1\,\Delta x + \varepsilon_2\,\Delta y$ where $\varepsilon_1, \varepsilon_2 \to 0$ as $(\Delta x, \Delta y) \to (0, 0)$. Now $\Delta z = f(\mathbf{x}) - f(\mathbf{x}_0)$, $\langle \Delta x, \Delta y\rangle = \mathbf{x} - \mathbf{x}_0$ so $(\Delta x, \Delta y) \to (0, 0)$ is equivalent to $\mathbf{x} \to \mathbf{x}_0$ and $\langle f_x(x_0, y_0), f_y(x_0, y_0)\rangle = \nabla f(\mathbf{x}_0)$. Substituting into 14.4.7 gives $f(\mathbf{x}) - f(\mathbf{x}_0) = \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) + \langle \varepsilon_1, \varepsilon_2\rangle \cdot \langle \Delta x, \Delta y\rangle$ or $\langle \varepsilon_1, \varepsilon_2\rangle \cdot (\mathbf{x} - \mathbf{x}_0) = f(\mathbf{x}) - f(\mathbf{x}_0) - \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0)$,

and so $\dfrac{f(\mathbf{x}) - f(\mathbf{x}_0) - \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0)}{|\mathbf{x} - \mathbf{x}_0|} = \dfrac{\langle \varepsilon_1, \varepsilon_2\rangle \cdot (\mathbf{x} - \mathbf{x}_0)}{|\mathbf{x} - \mathbf{x}_0|}$. But $\dfrac{\mathbf{x} - \mathbf{x}_0}{|\mathbf{x} - \mathbf{x}_0|}$ is a unit vector so

$\lim\limits_{\mathbf{x} \to \mathbf{x}_0} \dfrac{\langle \varepsilon_1, \varepsilon_2\rangle \cdot (\mathbf{x} - \mathbf{x}_0)}{|\mathbf{x} - \mathbf{x}_0|} = 0$ since $\varepsilon_1, \varepsilon_2 \to 0$ as $\mathbf{x} \to \mathbf{x}_0$. Hence $\lim\limits_{\mathbf{x} \to \mathbf{x}_0} \dfrac{f(\mathbf{x}) - f(\mathbf{x}_0) - \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0)}{|\mathbf{x} - \mathbf{x}_0|} = 0.$

14.7 Maximum and Minimum Values

1. (a) First we compute $D(1, 1) = f_{xx}(1, 1)\,f_{yy}(1, 1) - [f_{xy}(1, 1)]^2 = (4)(2) - (1)^2 = 7$. Since $D(1, 1) > 0$ and $f_{xx}(1, 1) > 0$, f has a local minimum at $(1, 1)$ by the Second Derivatives Test.

(b) $D(1, 1) = f_{xx}(1, 1)\,f_{yy}(1, 1) - [f_{xy}(1, 1)]^2 = (4)(2) - (3)^2 = -1$. Since $D(1, 1) < 0$, f has a saddle point at $(1, 1)$ by the Second Derivatives Test.

2. (a) $D = g_{xx}(0, 2)\,g_{yy}(0, 2) - [g_{xy}(0, 2)]^2 = (-1)(1) - (6)^2 = -37$. Since $D < 0$, g has a saddle point at $(0, 2)$ by the Second Derivatives Test.

(b) $D = g_{xx}(0, 2)\,g_{yy}(0, 2) - [g_{xy}(0, 2)]^2 = (-1)(-8) - (2)^2 = 4$. Since $D > 0$ and $g_{xx}(0, 2) < 0$, g has a local maximum at $(0, 2)$ by the Second Derivatives Test.

(c) $D = g_{xx}(0, 2)\,g_{yy}(0, 2) - [g_{xy}(0, 2)]^2 = (4)(9) - (6)^2 = 0$. In this case the Second Derivatives Test gives no information about g at the point $(0, 2)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. In the figure, a point at approximately $(1, 1)$ is enclosed by level curves which are oval in shape and indicate that as we move away from the point in any direction the values of f are increasing. Hence we would expect a local minimum at or near $(1, 1)$. The level curves near $(0, 0)$ resemble hyperbolas, and as we move away from the origin, the values of f increase in some directions and decrease in others, so we would expect to find a saddle point there.

To verify our predictions, we have $f(x, y) = 4 + x^3 + y^3 - 3xy \Rightarrow f_x(x, y) = 3x^2 - 3y$, $f_y(x, y) = 3y^2 - 3x$. We have critical points where these partial derivatives are equal to 0: $3x^2 - 3y = 0$, $3y^2 - 3x = 0$. Substituting $y = x^2$ from the first equation into the second equation gives $3(x^2)^2 - 3x = 0 \Rightarrow 3x(x^3 - 1) = 0 \Rightarrow x = 0$ or $x = 1$. Then we have two critical points, $(0, 0)$ and $(1, 1)$. The second partial derivatives are $f_{xx}(x, y) = 6x$, $f_{xy}(x, y) = -3$, and $f_{yy}(x, y) = 6y$, so $D(x, y) = f_{xx}(x, y)\, f_{yy}(x, y) - [f_{xy}(x, y)]^2 = (6x)(6y) - (-3)^2 = 36xy - 9$. Then $D(0, 0) = 36(0)(0) - 9 = -9$, and $D(1, 1) = 36(1)(1) - 9 = 27$. Since $D(0, 0) < 0$, f has a saddle point at $(0, 0)$ by the Second Derivatives Test. Since $D(1, 1) > 0$ and $f_{xx}(1, 1) > 0$, f has a local minimum at $(1, 1)$.

4. In the figure, points at approximately $(-1, 1)$ and $(-1, -1)$ are enclosed by oval-shaped level curves which indicate that as we move away from either point in any direction, the values of f are increasing. Hence we would expect local minima at or near $(-1, \pm 1)$. Similarly, the point $(1, 0)$ appears to be enclosed by oval-shaped level curves which indicate that as we move away from the point in any direction the values of f are decreasing, so we should have a local maximum there. We also show hyperbola-shaped level curves near the points $(-1, 0)$, $(1, 1)$, and $(1, -1)$. The values of f increase along some paths leaving these points and decrease in others, so we should have a saddle point at each of these points.

To confirm our predictions, we have $f(x, y) = 3x - x^3 - 2y^2 + y^4 \Rightarrow f_x(x, y) = 3 - 3x^2$, $f_y(x, y) = -4y + 4y^3$. Setting these partial derivatives equal to 0, we have $3 - 3x^2 = 0 \Rightarrow x = \pm 1$ and $-4y + 4y^3 = 0 \Rightarrow$ $y(y^2 - 1) = 0 \Rightarrow y = 0, \pm 1$. So our critical points are $(\pm 1, 0)$, $(\pm 1, 1)$, $(\pm 1, -1)$.

The second partial derivatives are $f_{xx}(x, y) = -6x$, $f_{xy}(x, y) = 0$, and $f_{yy}(x, y) = 12y^2 - 4$, so

$D(x, y) = f_{xx}(x, y)\, f_{yy}(x, y) - [f_{xy}(x, y)]^2 = (-6x)(12y^2 - 4) - (0)^2 = -72xy^2 + 24x$.

We use the Second Derivatives Test to classify the 6 critical points:

Critical Point	D	f_{xx}	Conclusion
$(1, 0)$	24	-6	$D > 0$, $f_{xx} < 0 \Rightarrow$ f has a local maximum at $(1, 0)$
$(1, 1)$	-48		$D < 0 \Rightarrow$ f has a saddle point at $(1, 1)$
$(1, -1)$	-48		$D < 0 \Rightarrow$ f has a saddle point at $(1, -1)$
$(-1, 0)$	-24		$D < 0 \Rightarrow$ f has a saddle point at $(-1, 0)$
$(-1, 1)$	48	6	$D > 0$, $f_{xx} > 0 \Rightarrow$ f has a local minimum at $(-1, 1)$
$(-1, -1)$	48	6	$D > 0$, $f_{xx} > 0 \Rightarrow$ f has a local minimum at $(-1, -1)$

5. $f(x, y) = x^2 + xy + y^2 + y \Rightarrow f_x = 2x + y$, $f_y = x + 2y + 1$, $f_{xx} = 2$, $f_{xy} = 1$, $f_{yy} = 2$. Then $f_x = 0$ implies $y = -2x$, and substitution into $f_y = x + 2y + 1 = 0$ gives $x + 2(-2x) + 1 = 0 \Rightarrow -3x = -1 \Rightarrow x = \frac{1}{3}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Then $y = -\frac{2}{3}$ and the only critical point is $\left(\frac{1}{3}, -\frac{2}{3}\right)$.

$D(x,y) = f_{xx}f_{yy} - (f_{xy})^2 = (2)(2) - (1)^2 = 3$, and since

$D\left(\frac{1}{3}, -\frac{2}{3}\right) = 3 > 0$ and $f_{xx}\left(\frac{1}{3}, -\frac{2}{3}\right) = 2 > 0$, $f\left(\frac{1}{3}, -\frac{2}{3}\right) = -\frac{1}{3}$ is a local minimum by the Second Derivatives Test.



6. $f(x,y) = xy - 2x - 2y - x^2 - y^2 \quad \Rightarrow \quad f_x = y - 2 - 2x$,

$f_y = x - 2 - 2y$, $f_{xx} = -2$, $f_{xy} = 1$, $f_{yy} = -2$. Then $f_x = 0$ implies $y = 2x + 2$, and substitution into $f_y = 0$ gives $x - 2 - 2(2x + 2) = 0 \quad \Rightarrow$ $-3x = 6 \quad \Rightarrow \quad x = -2$. Then $y = -2$ and the only critical point is $(-2, -2)$. $D(x,y) = f_{xx}f_{yy} - (f_{xy})^2 = (-2)(-2) - 1^2 = 3$, and since $D(-2,-2) = 3 > 0$ and $f_{xx}(-2,-2) = -2 < 0$, $f(-2,-2) = 4$ is a local maximum by the Second Derivatives Test.



7. $f(x,y) = (x-y)(1-xy) = x - y - x^2y + xy^2 \quad \Rightarrow \quad f_x = 1 - 2xy + y^2$, $f_y = -1 - x^2 + 2xy$, $f_{xx} = -2y$, $f_{xy} = -2x + 2y$, $f_{yy} = 2x$. Then $f_x = 0$ implies $1 - 2xy + y^2 = 0$ and $f_y = 0$ implies $-1 - x^2 + 2xy = 0$. Adding the two equations gives $1 + y^2 - 1 - x^2 = 0 \quad \Rightarrow \quad y^2 = x^2 \quad \Rightarrow \quad y = \pm x$, but if $y = -x$ then $f_x = 0$ implies $1 + 2x^2 + x^2 = 0 \quad \Rightarrow \quad 3x^2 = -1$ which has no real solution. If $y = x$ then substitution into $f_x = 0$ gives $1 - 2x^2 + x^2 = 0 \quad \Rightarrow \quad x^2 = 1 \quad \Rightarrow$ $x = \pm 1$, so the critical points are $(1,1)$ and $(-1,-1)$. Now $D(1,1) = (-2)(2) - 0^2 = -4 < 0$ and $D(-1,-1) = (2)(-2) - 0^2 = -4 < 0$, so $(1,1)$ and $(-1,-1)$ are saddle points.



8. $f(x,y) = y(e^x - 1) \quad \Rightarrow \quad f_x = ye^x$, $f_y = e^x - 1$, $f_{xx} = ye^x$, $f_{xy} = e^x$, $f_{yy} = 0$. Because e^x is never zero, $f_x = 0$ only when $y = 0$, and $f_y = 0$ when $e^x = 1 \quad \Rightarrow \quad x = 0$, so the only critical point is $(0,0)$. $D(x,y) = f_{xx}f_{yy} - (f_{xy})^2 = (ye^x)(0) - (e^x)^2 = -e^{2x}$, and since $D(0,0) = -1 < 0$, $(0,0)$ is a saddle point.



9. $f(x,y) = x^2 + y^4 + 2xy \quad \Rightarrow \quad f_x = 2x + 2y$, $f_y = 4y^3 + 2x$, $f_{xx} = 2$, $f_{xy} = 2$, $f_{yy} = 12y^2$. Then $f_x = 0$ implies $y = -x$, and substitution into $f_y = 4y^3 + 2x = 0$ gives $-4x^3 + 2x = 0 \quad \Rightarrow \quad 2x\left(1 - 2x^2\right) = 0 \quad \Rightarrow \quad x = 0$ or $x = \pm\frac{1}{\sqrt{2}}$. Thus the critical points are $(0,0)$, $\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$, and $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$. Now

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$D(x,y) = f_{xx}f_{yy} - (f_{xy})^2 = (2)(12y^2) - (2)^2 = 24y^2 - 4$,
so $D(0,0) = -4 < 0$ and $(0,0)$ is a saddle point.

$D\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right) = D\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) = 24\left(\frac{1}{2}\right) - 4 = 8 > 0$ and

$f_{xx}\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right) = f_{xx}\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) = 2 > 0$, so $f\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) = -\frac{1}{4}$

and $f\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) = -\frac{1}{4}$ are local minima.



10. $f(x,y) = 2 - x^4 + 2x^2 - y^2 \Rightarrow f_x = -4x^3 + 4x,\ f_y = -2y,\ f_{xx} = -12x^2 + 4,\ f_{xy} = 0,\ f_{yy} = -2$. Then $f_x = 0$ implies $-4x(x^2 - 1) = 0$, so $x = 0$ or $x = \pm 1$, and $f_y = 0$ implies $y = 0$. Thus the critical points are $(0,0)$, $(\pm 1, 0)$.

$D(0,0) = (4)(-2) - 0^2 = -8 < 0$, so $(0,0)$ is a saddle point.

$D(1,0) = D(-1,0) = (-8)(-2) - (0)^2 = 16 > 0$, and

$f_{xx}(1,0) = f_{xx}(-1,0) = -8 < 0$, so $f(1,0) = 3$ and $f(-1,0) = 3$ are local maxima.



11. $f(x,y) = x^3 - 3x + 3xy^2 \Rightarrow f_x = 3x^2 - 3 + 3y^2,\ f_y = 6xy,\ f_{xx} = 6x,\ f_{xy} = 6y,\ f_{yy} = 6x$. Then $f_y = 0$ implies $x = 0$ or $y = 0$. If $x = 0$, substitution into $f_x = 0$ gives $3y^2 = 3 \Rightarrow y = \pm 1$, and if $y = 0$, substitution into $f_x = 0$ gives $x = \pm 1$. Thus the critical points are $(0, \pm 1)$ and $(\pm 1, 0)$.

$D(0, \pm 1) = 0 - 36 < 0$, so $(0, \pm 1)$ are saddle points.

$D(\pm 1, 0) = 36 - 0 > 0$, $f_{xx}(1,0) = 6 > 0$, and $f_{xx}(-1,0) = -6 < 0$, so $f(1,0) = -2$ is a local minimum and $f(-1,0) = 2$ is a local maximum.



12. $f(x,y) = x^3 + y^3 - 3x^2 - 3y^2 - 9x \Rightarrow f_x = 3x^2 - 6x - 9,\ f_y = 3y^2 - 6y,\ f_{xx} = 6x - 6,\ f_{xy} = 0,\ f_{yy} = 6y - 6$. Then $f_x = 0$ implies $3(x+1)(x-3) = 0 \Rightarrow x = -1$ or $x = 3$, and $f_y = 0$ implies $3y(y-2) = 0 \Rightarrow y = 0$ or $y = 2$. Thus the critical points are $(-1,0)$, $(-1,2)$, $(3,0)$, and $(3,2)$. $D(-1,2) = (-12)(6) - (0)^2 = -72 < 0$ and $D(3,0) = (12)(-6) - (0)^2 = -72 < 0$, so $(-1,2)$ and $(3,0)$ are saddle points. $D(-1,0) = (-12)(-6) - (0)^2 = 72 > 0$ and $f_{xx}(-1,0) = -12 < 0$, so $f(-1,0) = 5$ is a local maximum.

$D(3,2) = (12)(6) - (0)^2 = 72 > 0$ and $f_{xx}(3,2) = 12 > 0$, so $f(3,2) = -31$ is a local minimum.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

13. $f(x,y) = x^4 - 2x^2 + y^3 - 3y \;\Rightarrow\; f_x = 4x^3 - 4x,\ f_y = 3y^2 - 3,\ f_{xx} = 12x^2 - 4,\ f_{xy} = 0,\ f_{yy} = 6y.$

Then $f_x = 0$ implies $4x(x^2-1) = 0 \;\Rightarrow\; x = 0$ or $x = \pm 1$, and $f_y = 0$ implies $3(y^2-1) = 0 \;\Rightarrow\; y = \pm 1$.

Thus there are six critical points: $(0,\pm 1)$, $(\pm 1, 1)$, and $(\pm 1, -1)$.

$D(0,1) = (-4)(6) - (0)^2 = -24 < 0$ and

$D(\pm 1,-1) = (8)(-6) = -48 < 0$, so $(0,1)$ and $(\pm 1,-1)$ are saddle points. $D(0,-1) = (-4)(-6) - 24 > 0$ and $f_{xx}(0,-1) = -4 < 0$, so $f(0,-1) = 2$ is a local maximum. $D(\pm 1, 1) = (8)(6) = 48 > 0$ and $f_{xx}(\pm 1, 1) = 8 > 0$, so $f(\pm 1, 1) = -3$ are local minima.



14. $f(x,y) = y\cos x \;\Rightarrow\; f_x = -y\sin x,\ f_y = \cos x,\ f_{xx} = -y\cos x,$ $f_{xy} = -\sin x,\ f_{yy} = 0$. Then $f_y = 0$ if and only if $x = \frac{\pi}{2} + n\pi$ for n an integer. But $\sin\left(\frac{\pi}{2} + n\pi\right) \neq 0$, so $f_x = 0 \;\Rightarrow\; y = 0$ and the critical points are $\left(\frac{\pi}{2} + n\pi, 0\right)$, n an integer.

$D\left(\frac{\pi}{2} + n\pi, 0\right) = (0)(0) - (\pm 1)^2 = -1 < 0$, so each critical point is a saddle point.



15. $f(x,y) = e^x \cos y \;\Rightarrow\; f_x = e^x\cos y,\ f_y = -e^x \sin y.$

Now $f_x = 0$ implies $\cos y = 0$ or $y = \frac{\pi}{2} + n\pi$ for n an integer.

But $\sin\left(\frac{\pi}{2} + n\pi\right) \neq 0$, so there are no critical points.



16. $f(x,y) = xye^{-(x^2+y^2)/2} \;\Rightarrow\; f_x = xy\cdot e^{-(x^2+y^2)/2}(-x) + e^{-(x^2+y^2)/2}\cdot y = y(1-x^2)e^{-(x^2+y^2)/2},$

$f_y = xy\cdot e^{-(x^2+y^2)/2}(-y) + e^{-(x^2+y^2)/2}\cdot x = x(1-y^2)e^{-(x^2+y^2)/2},$

$f_{xx} = y\left[(1-x^2)\cdot e^{-(x^2+y^2)/2}(-x) + e^{-(x^2+y^2)/2}(-2x)\right] = xy(x^2-3)e^{-(x^2+y^2)/2},$

$f_{xy} = (1-x^2)\left[y\cdot e^{-(x^2+y^2)/2}(-y) + e^{-(x^2+y^2)/2}(1)\right] = (1-x^2)(1-y^2)e^{-(x^2+y^2)/2},$

$f_{yy} = x\left[(1-y^2)\cdot e^{-(x^2+y^2)/2}(-y) + e^{-(x^2+y^2)/2}(-2y)\right] = xy(y^2-3)e^{-(x^2+y^2)/2}.$

Then $f_x = 0$ implies $y(1-x^2) = 0 \;\Rightarrow\; y = 0$ or $x = \pm 1$. Substituting $y = 0$ into $f_y = 0$ gives $xe^{-x^2/2} = 0 \;\Rightarrow\; x = 0$, and substituting $x = \pm 1$ into $f_y = 0$ gives $\pm(1-y^2)e^{-(1+y^2)/2} = 0 \;\Rightarrow\; y = \pm 1$, so the critical points are $(0,0)$, $(1,\pm 1)$, and $(-1,\pm 1)$. $D(0,0) = (0)(0) - (1)^2 = -1 < 0$, so $(0,0)$ is a saddle point.

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$D(1,1) = D(-1,-1) = (-2e^{-1})(-2e^{-1}) - (0)^2 = 4e^{-2} > 0$ and $f_{xx}(1,1) = f_{xx}(-1,-1) = -2e^{-1} < 0$, so $f(1,1) = f(-1,-1) = e^{-1}$ are local maxima.

$D(1,-1) = D(-1,1) = (2e^{-1})(2e^{-1}) - (0)^2 = 4e^{-2} > 0$ and $f_{xx}(1,-1) = f_{xx}(-1,1) = 2e^{-1} > 0$, so $f(1,-1) = f(-1,1) = -e^{-1}$ are local minima.



17. $f(x,y) = xy + e^{-xy} \Rightarrow f_x = y - ye^{-xy}$, $f_y = x - xe^{-xy}$, $f_{xx} = y^2e^{-xy}$, $f_{xy} = 1 - \left[y(-xe^{-xy}) + e^{-xy}(1)\right] = 1 + (xy-1)e^{-xy}$, $f_{yy} = x^2e^{-xy}$. Then $f_x = 0$ implies $y(1 - e^{-xy}) = 0 \Rightarrow$ $y = 0$ or $e^{-xy} = 1 \Rightarrow x = 0$ or $y = 0$. If $x = 0$ then $f_y = 0$ for any y-value, so all points of the form $(0, y_0)$ are critical points. If $y = 0$, then $f_y = x - xe^0 = 0$ for any x-value, so all points of the form $(x_0, 0)$ are critical points. We have $D(x_0, 0) = (0)(x_0^2) - (0)^2 = 0$ and $D(0, y_0) = (y_0^2)(0) - (0)^2 = 0$, so the Second Derivatives Test gives no information.

Notice that if we let $t = xy$, then $f(x,y) = g(t) = t + e^{-t} \Rightarrow$ $g'(t) = 1 - e^{-t}$. Now $g'(t) = 0$ only for $t = 0$, and $g'(t) < 0$ for $t < 0$, $g'(t) > 0$ for $t > 0$. Thus $g(0) = 1$ is a local and absolute minimum, so $f(x,y) = xy + e^{-xy} \geq 1$ for all (x,y) with equality if and only if $x = 0$ or $y = 0$. Hence all points on the x- and y-axes are local (and absolute) minima, where $f(x,y) = 1$.



18. $f(x,y) = (x^2 + y^2)e^{-x} \Rightarrow f_x = (x^2 + y^2)(-e^{-x}) + e^{-x}(2x) = (2x - x^2 - y^2)e^{-x}$, $f_y = 2ye^{-x}$, $f_{xx} = (2x - x^2 - y^2)(-e^{-x}) + e^{-x}(2 - 2x) = (x^2 + y^2 - 4x + 2)e^{-x}$, $f_{xy} = -2ye^{-x}$, $f_{yy} = 2e^{-x}$. Then $f_y = 0$ implies $y = 0$ and substituting into $f_x = 0$ gives $(2x - x^2)e^{-x} = 0 \Rightarrow$ $x(2 - x) = 0 \Rightarrow x = 0$ or $x = 2$, so the critical points are $(0,0)$ and $(2,0)$. $D(0,0) = (2)(2) - (0)^2 = 4 > 0$ and $f_{xx}(0,0) = 2 > 0$, so $f(0,0) = 0$ is a local minimum.

$D(2,0) = (-2e^{-2})(2e^{-2}) - (0)^2 = -4e^{-4} < 0$ so $(2,0)$ is a saddle point.



19. $f(x,y) = y^2 - 2y\cos x \Rightarrow f_x = 2y\sin x$, $f_y = 2y - 2\cos x$, $f_{xx} = 2y\cos x$, $f_{xy} = 2\sin x$, $f_{yy} = 2$. Then $f_x = 0$ implies $y = 0$ or $\sin x = 0 \Rightarrow x = 0, \pi$, or 2π for $-1 \leq x \leq 7$. Substituting $y = 0$ into $f_y = 0$ gives $\cos x = 0 \Rightarrow x = \frac{\pi}{2}$ or $\frac{3\pi}{2}$, substituting $x = 0$ or $x = 2\pi$ into $f_y = 0$ gives $y = 1$, and substituting $x = \pi$ into $f_y = 0$ gives $y = -1$. Thus the critical points are $(0,1)$, $\left(\frac{\pi}{2}, 0\right)$, $(\pi, -1)$, $\left(\frac{3\pi}{2}, 0\right)$, and $(2\pi, 1)$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$D\left(\frac{\pi}{2},0\right) = D\left(\frac{3\pi}{2},0\right) = -4 < 0$ so $\left(\frac{\pi}{2},0\right)$ and $\left(\frac{3\pi}{2},0\right)$ are saddle points. $D(0,1) = D(\pi,-1) = D(2\pi,1) = 4 > 0$ and $f_{xx}(0,1) = f_{xx}(\pi,-1) = f_{xx}(2\pi,1) = 2 > 0$, so $f(0,1) = f(\pi,-1) = f(2\pi,1) = -1$ are local minima.

20. $f(x,y) = \sin x \sin y \quad \Rightarrow \quad f_x = \cos x \sin y$, $f_y = \sin x \cos y$, $f_{xx} = -\sin x \sin y$, $f_{xy} = \cos x \cos y$, $f_{yy} = -\sin x \sin y$. Here we have $-\pi < x < \pi$ and $-\pi < y < \pi$, so $f_x = 0$ implies $\cos x = 0$ or $\sin y = 0$. If $\cos x = 0$ then $x = -\frac{\pi}{2}$ or $\frac{\pi}{2}$, and if $\sin y = 0$ then $y = 0$. Substituting $x = \pm\frac{\pi}{2}$ into $f_y = 0$ gives $\cos y = 0 \quad \Rightarrow \quad y = -\frac{\pi}{2}$ or $\frac{\pi}{2}$, and substituting $y = 0$ into $f_y = 0$ gives $\sin x = 0 \quad \Rightarrow \quad x = 0$. Thus the critical points are $\left(-\frac{\pi}{2}, \pm\frac{\pi}{2}\right)$, $\left(\frac{\pi}{2}, \pm\frac{\pi}{2}\right)$, and $(0,0)$.

$D(0,0) = -1 < 0$ so $(0,0)$ is a saddle point.

$D\left(-\frac{\pi}{2}, \pm\frac{\pi}{2}\right) = D\left(\frac{\pi}{2}, \pm\frac{\pi}{2}\right) = 1 > 0$ and

$f_{xx}\left(-\frac{\pi}{2}, -\frac{\pi}{2}\right) = f_{xx}\left(\frac{\pi}{2}, \frac{\pi}{2}\right) = -1 < 0$ while

$f_{xx}\left(-\frac{\pi}{2}, \frac{\pi}{2}\right) = f_{xx}\left(\frac{\pi}{2}, -\frac{\pi}{2}\right) = 1 > 0$, so $f\left(-\frac{\pi}{2}, -\frac{\pi}{2}\right) = f\left(\frac{\pi}{2}, \frac{\pi}{2}\right) = 1$ are local maxima and $f\left(-\frac{\pi}{2}, \frac{\pi}{2}\right) = f\left(\frac{\pi}{2}, -\frac{\pi}{2}\right) = 1$ are local minima.



21. $f(x,y) = x^2 + 4y^2 - 4xy + 2 \quad \Rightarrow \quad f_x = 2x - 4y$, $f_y = 8y - 4x$, $f_{xx} = 2$, $f_{xy} = -4$, $f_{yy} = 8$. Then $f_x = 0$ and $f_y = 0$ each implies $y = \frac{1}{2}x$, so all points of the form $\left(x_0, \frac{1}{2}x_0\right)$ are critical points and for each of these we have $D\left(x_0, \frac{1}{2}x_0\right) = (2)(8) - (-4)^2 = 0$. The Second Derivatives Test gives no information, but $f(x,y) = x^2 + 4y^2 - 4xy + 2 = (x - 2y)^2 + 2 \geq 2$ with equality if and only if $y = \frac{1}{2}x$. Thus $f\left(x_0, \frac{1}{2}x_0\right) = 2$ are all local (and absolute) minima.

22. $f(x,y) = x^2 y e^{-x^2-y^2} \quad \Rightarrow$

$f_x = x^2 y e^{-x^2-y^2}(-2x) + 2xy e^{-x^2-y^2} = 2xy(1 - x^2)e^{-x^2-y^2}$,

$f_y = x^2 y e^{-x^2-y^2}(-2y) + x^2 e^{-x^2-y^2} = x^2(1 - 2y^2)e^{-x^2-y^2}$,

$f_{xx} = 2y(2x^4 - 5x^2 + 1)e^{-x^2-y^2}$,

$f_{xy} = 2x(1 - x^2)(1 - 2y^2)e^{-x^2-y^2}$, $f_{yy} = 2x^2 y(2y^2 - 3)e^{-x^2-y^2}$.



$f_x = 0$ implies $x = 0$, $y = 0$, or $x = \pm 1$. If $x = 0$ then $f_y = 0$ for any y-value, so all points of the form $(0,y)$ are critical points. If $y = 0$ then $f_y = 0 \quad \Rightarrow \quad x^2 e^{-x^2} = 0 \quad \Rightarrow \quad x = 0$, so $(0,0)$ (already included above) is a critical point. If $x = \pm 1$ then $(1 - 2y^2)e^{-1-y^2} = 0 \quad \Rightarrow \quad y = \pm\frac{1}{\sqrt{2}}$, so $\left(\pm 1, \frac{1}{\sqrt{2}}\right)$ and $\left(\pm 1, -\frac{1}{\sqrt{2}}\right)$ are critical points. Now

$D\left(\pm 1, \frac{1}{\sqrt{2}}\right) = 8e^{-3} > 0$, $f_{xx}\left(\pm 1, \frac{1}{\sqrt{2}}\right) = -2\sqrt{2}\,e^{-3/2} < 0$ and $D\left(\pm 1, -\frac{1}{\sqrt{2}}\right) = 8e^{-3} > 0$,

$f_{xx}\left(\pm 1, -\frac{1}{\sqrt{2}}\right) = 2\sqrt{2}\,e^{-3/2} > 0$, so $f\left(\pm 1, \frac{1}{\sqrt{2}}\right) = \frac{1}{\sqrt{2}}e^{-3/2}$ are local maximum points while

$f\left(\pm 1, -\frac{1}{\sqrt{2}}\right) = -\frac{1}{\sqrt{2}}e^{-3/2}$ are local minimum points. At all critical points $(0,y)$ we have $D(0,y) = 0$, so the Second Derivatives Test gives no information. However, if $y > 0$ then $x^2 y e^{-x^2-y^2} \geq 0$ with equality only when $x = 0$, so we have local minimum values $f(0,y) = 0$, $y > 0$. Similarly, if $y < 0$ then $x^2 y e^{-x^2-y^2} \leq 0$ with equality when $x = 0$ so $f(0,y) = 0$, $y < 0$ are local maximum values, and $(0,0)$ is a saddle point.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $f(x, y) = x^2 + y^2 + x^{-2}y^{-2}$







From the graphs, there appear to be local minima of about $f(1, \pm 1) = f(-1, \pm 1) \approx 3$ (and no local maxima or saddle points). $f_x = 2x - 2x^{-3}y^{-2}$, $f_y = 2y - 2x^{-2}y^{-3}$, $f_{xx} = 2 + 6x^{-4}y^{-2}$, $f_{xy} = 4x^{-3}y^{-3}$, $f_{yy} = 2 + 6x^{-2}y^{-4}$. Then $f_x = 0$ implies $2x^4y^2 - 2 = 0$ or $x^4y^2 = 1$ or $y^2 = x^{-4}$. Note that neither x nor y can be zero. Now $f_y = 0$ implies $2x^2y^4 - 2 = 0$, and with $y^2 = x^{-4}$ this implies $2x^{-6} - 2 = 0$ or $x^6 = 1$. Thus $x = \pm 1$ and if $x = 1$, $y = \pm 1$; if $x = -1$, $y = \pm 1$. So the critical points are $(1, 1)$, $(1, -1)$,$(-1, 1)$ and $(-1, -1)$. Now $D(1, \pm 1) = D(-1, \pm 1) = 64 - 16 > 0$ and $f_{xx} > 0$ always, so $f(1, \pm 1) = f(-1, \pm 1) = 3$ are local minima.

24. $f(x, y) = (x - y)e^{-x^2-y^2}$







From the graphs, there appears to be a local maximum of about $f(0.5, -0.5) \approx 0.6$ and a local minimum of about $f(-0.5, 0.5) \approx -0.6$.

$f_x = (x - y)e^{-x^2-y^2}(-2x) + e^{-x^2-y^2}(1) = e^{-x^2-y^2}(1 - 2x^2 + 2xy)$,

$f_y = (x - y)e^{-x^2-y^2}(-2y) + e^{-x^2-y^2}(-1) = -e^{-x^2-y^2}(1 - 2y^2 + 2xy)$, $f_{xx} = 2e^{-x^2-y^2}(2x^3 - 3x + y - 2x^2y)$, $f_{xy} = 2e^{-x^2-y^2}(x - y + 2x^2y - 2xy^2)$, $f_{yy} = -2e^{-x^2-y^2}(2y^3 - 3y + x - 2xy^2)$. Then $f_x = 0$ implies $1 - 2x^2 + 2xy = 0$ and $f_y = 0$ implies $1 - 2y^2 + 2xy = 0$. Subtracting these two equations gives $-2x^2 + 2y^2 = 0 \quad \Rightarrow \quad y = \pm x$. If $y = x$ then substituting into $f_x = 0$ gives $1 - 2x^2 + 2x^2 = 0$, an impossibility. Substituting $y = -x$ gives $1 - 2x^2 - 2x^2 = 0 \quad \Rightarrow \quad x^2 = \frac{1}{4} \quad \Rightarrow \quad x = \pm\frac{1}{2}$. Thus the critical points are $\left(\frac{1}{2}, -\frac{1}{2}\right)$ and $\left(-\frac{1}{2}, \frac{1}{2}\right)$. Now $D\left(\frac{1}{2}, -\frac{1}{2}\right) = (-3e^{-1/2})(-3e^{-1/2}) - (e^{-1/2})^2 = 8e^{-1} > 0$ with $f_{xx}\left(\frac{1}{2}, -\frac{1}{2}\right) = -3e^{-1/2} < 0$, so $f\left(\frac{1}{2}, -\frac{1}{2}\right) = e^{-1/2} \approx 0.607$ is a local maximum, and $D\left(-\frac{1}{2}, \frac{1}{2}\right) = (3e^{-1/2})(3e^{-1/2}) - (-e^{-1/2})^2 = 8e^{-1} > 0$ with $f_{xx}\left(-\frac{1}{2}, \frac{1}{2}\right) = 3e^{-1/2} > 0$, so $f\left(-\frac{1}{2}, \frac{1}{2}\right) = -e^{-1/2} \approx -0.607$ is a local minimum.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

25. $f(x, y) = \sin x + \sin y + \sin(x + y),\ 0 \le x \le 2\pi,\ 0 \le y \le 2\pi$



From the graphs it appears that f has a local maximum at about $(1, 1)$ with value approximately 2.6, a local minimum at about $(5, 5)$ with value approximately -2.6, and a saddle point at about $(3, 3)$.

$f_x = \cos x + \cos(x + y),\ f_y = \cos y + \cos(x + y),\ f_{xx} = -\sin x - \sin(x + y),\ f_{yy} = -\sin y - \sin(x + y),$ $f_{xy} = -\sin(x + y)$. Setting $f_x = 0$ and $f_y = 0$ and subtracting gives $\cos x - \cos y = 0$ or $\cos x = \cos y$. Thus $x = y$ or $x = 2\pi - y$. If $x = y$, $f_x = 0$ becomes $\cos x + \cos 2x = 0$ or $2\cos^2 x + \cos x - 1 = 0$, a quadratic in $\cos x$. Thus $\cos x = -1$ or $\frac{1}{2}$ and $x = \pi$, $\frac{\pi}{3}$, or $\frac{5\pi}{3}$, giving the critical points (π, π), $\left(\frac{\pi}{3}, \frac{\pi}{3}\right)$ and $\left(\frac{5\pi}{3}, \frac{5\pi}{3}\right)$. Similarly if $x = 2\pi - y$, $f_x = 0$ becomes $(\cos x) + 1 = 0$ and the resulting critical point is (π, π). Now $D(x, y) = \sin x \sin y + \sin x \sin(x + y) + \sin y \sin(x + y)$. So $D(\pi, \pi) = 0$ and the Second Derivatives Test doesn't apply. However, along the line $y = x$ we have $f(x, x) = 2\sin x + \sin 2x = 2\sin x + 2\sin x \cos x = 2\sin x(1 + \cos x)$, and $f(x, x) > 0$ for $0 < x < \pi$ while $f(x, x) < 0$ for $\pi < x < 2\pi$. Thus every disk with center (π, π) contains points where f is positive as well as points where f is negative, so the graph crosses its tangent plane ($z = 0$) there and (π, π) is a saddle point.

$D\left(\frac{\pi}{3}, \frac{\pi}{3}\right) = \frac{9}{4} > 0$ and $f_{xx}\left(\frac{\pi}{3}, \frac{\pi}{3}\right) < 0$ so $f\left(\frac{\pi}{3}, \frac{\pi}{3}\right) = \frac{3\sqrt{3}}{2}$ is a local maximum while $D\left(\frac{5\pi}{3}, \frac{5\pi}{3}\right) = \frac{9}{4} > 0$ and $f_{xx}\left(\frac{5\pi}{3}, \frac{5\pi}{3}\right) > 0$, so $f\left(\frac{5\pi}{3}, \frac{5\pi}{3}\right) = -\frac{3\sqrt{3}}{2}$ is a local minimum.

26. $f(x, y) = \sin x + \sin y + \cos(x + y),\ 0 < x \le \frac{\pi}{4},\ 0 < y \le \frac{\pi}{4}$



[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

From the graphs, it seems that f has a local maximum at about $(0.5, 0.5)$.

$f_x = \cos x - \sin(x+y)$, $f_y = \cos y - \sin(x+y)$, $f_{xx} = -\sin x - \cos(x+y)$, $f_{yy} = -\sin y - \cos(x+y)$, $f_{xy} = -\cos(x+y)$. Setting $f_x = 0$ and $f_y = 0$ and subtracting gives $\cos x = \cos y$. Thus $x = y$. Substituting $x = y$ into $f_x = 0$ gives $\cos x - \sin 2x = 0$ or $\cos x(1 - 2\sin x) = 0$. But $\cos x \neq 0$ for $0 \le x \le \frac{\pi}{4}$ and $1 - 2\sin x = 0$ implies $x = \frac{\pi}{6}$, so the only critical point is $\left(\frac{\pi}{6}, \frac{\pi}{6}\right)$. Here $f_{xx}\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = -1 < 0$ and $D\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = (-1)^2 - \frac{1}{4} > 0$. Thus $f\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = \frac{3}{2}$ is a local maximum.

27. $f(x,y) = x^4 + y^4 - 4x^2y + 2y \Rightarrow f_x(x,y) = 4x^3 - 8xy$ and $f_y(x,y) = 4y^3 - 4x^2 + 2$. $f_x = 0 \Rightarrow 4x(x^2 - 2y) = 0$, so $x = 0$ or $x^2 = 2y$. If $x = 0$ then substitution into $f_y = 0$ gives $4y^3 = -2 \Rightarrow y = -\frac{1}{\sqrt[3]{2}}$, so $\left(0, -\frac{1}{\sqrt[3]{2}}\right)$ is a critical point. Substituting $x^2 = 2y$ into $f_y = 0$ gives $4y^3 - 8y + 2 = 0$. Using a graph, solutions are approximately $y = -1.526$, 0.259, and 1.267. (Alternatively, we could have used a calculator or a CAS to find these roots.) We have $x^2 = 2y \Rightarrow x = \pm\sqrt{2y}$, so $y = -1.526$ gives no real-valued solution for x, but $y = 0.259 \Rightarrow x \approx \pm 0.720$ and $y = 1.267 \Rightarrow x \approx \pm 1.592$. Thus to three decimal places, the critical points are $\left(0, -\frac{1}{\sqrt[3]{2}}\right) \approx (0, -0.794)$, $(\pm 0.720, 0.259)$, and $(\pm 1.592, 1.267)$. Now since $f_{xx} = 12x^2 - 8y$, $f_{xy} = -8x$, $f_{yy} = 12y^2$, and $D = (12x^2 - 8y)(12y^2) - 64x^2$, we have $D(0, -0.794) > 0$, $f_{xx}(0, -0.794) > 0$, $D(\pm 0.720, 0.259) < 0$, $D(\pm 1.592, 1.267) > 0$, and $f_{xx}(\pm 1.592, 1.267) > 0$. Therefore $f(0, -0.794) \approx -1.191$ and $f(\pm 1.592, 1.267) \approx -1.310$ are local minima, and $(\pm 0.720, 0.259)$ are saddle points. There is no highest point on the graph, but the lowest points are approximately $(\pm 1.592, 1.267, -1.310)$.





28. $f(x,y) = y^6 - 2y^4 + x^2 - y^2 + y \Rightarrow f_x(x,y) = 2x$ and $f_y(x,y) = 6y^5 - 8y^3 - 2y + 1$. $f_x = 0$ implies $x = 0$, and the graph of f_y shows that the roots of $f_y = 0$ are approximately $y = -1.273$, 0.347, and 1.211. (Alternatively, we could have found the roots of $f_y = 0$ directly, using a calculator or CAS.) So to three decimal places, the critical points are $(0, -1.273)$, $(0, 0.347)$, and $(0, 1.211)$. Now since $f_{xx} = 2$, $f_{xy} = 0$, $f_{yy} = 30y^4 - 24y^2 - 2$, and $D = 60y^4 - 48y^2 - 4$, we have $D(0, -1.273) > 0$, $f_{xx}(0, -1.273) > 0$, $D(0, 0.347) < 0$, $D(0, 1.211) > 0$, and $f_{xx}(0, 1.211) > 0$, so $f(0, -1.273) \approx -3.890$ and $f(0, 1.211) \approx -1.403$ are local minima, and $(0, 0.347)$ is a saddle point. The lowest point on

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

the graph is approximately $(0, -1.273, -3.890)$.





29. $f(x, y) = x^4 + y^3 - 3x^2 + y^2 + x - 2y + 1 \Rightarrow f_x(x, y) = 4x^3 - 6x + 1$ and $f_y(x, y) = 3y^2 + 2y - 2$. From the graphs, we see that to three decimal places, $f_x = 0$ when $x \approx -1.301$, 0.170, or 1.131, and $f_y = 0$ when $y \approx -1.215$ or 0.549. (Alternatively, we could have used a calculator or a CAS to find these roots. We could also use the quadratic formula to find the solutions of $f_y = 0$.) So, to three decimal places, f has critical points at $(-1.301, -1.215)$, $(-1.301, 0.549)$, $(0.170, -1.215)$, $(0.170, 0.549)$, $(1.131, -1.215)$, and $(1.131, 0.549)$. Now since $f_{xx} = 12x^2 - 6$, $f_{xy} = 0$, $f_{yy} = 6y + 2$, and $D = (12x^2 - 6)(6y + 2)$, we have $D(-1.301, -1.215) < 0$, $D(-1.301, 0.549) > 0$, $f_{xx}(-1.301, 0.549) > 0$, $D(0.170, -1.215) > 0$, $f_{xx}(0.170, -1.215) < 0$, $D(0.170, 0.549) < 0$, $D(1.131, -1.215) < 0$, $D(1.131, 0.549) > 0$, and $f_{xx}(1.131, 0.549) > 0$. Therefore, to three decimal places, $f(-1.301, 0.549) \approx -3.145$ and $f(1.131, 0.549) \approx -0.701$ are local minima, $f(0.170, -1.215) \approx 3.197$ is a local maximum, and $(-1.301, -1.215)$, $(0.170, 0.549)$, and $(1.131, -1.215)$ are saddle points. There is no highest or lowest point on the graph.







30. $f(x, y) = 20e^{-x^2-y^2} \sin 3x \cos 3y \Rightarrow$

$$f_x(x, y) = 20 \cos 3y \left[e^{-x^2-y^2}(3 \cos 3x) + (\sin 3x)e^{-x^2-y^2}(-2x)\right]$$
$$= 20e^{-x^2-y^2} \cos 3y\,(3 \cos 3x - 2x \sin 3x)$$

$$f_y(x, y) = 20 \sin 3x \left[e^{-x^2-y^2}(-3 \sin 3y) + (\cos 3y)e^{-x^2-y^2}(-2y)\right]$$
$$= -20e^{-x^2-y^2} \sin 3x\,(3 \sin 3y + 2y \cos 3y)$$

Now $f_x = 0$ implies $\cos 3y = 0$ or $3 \cos 3x - 2x \sin 3x = 0$. For $|y| \leq 1$, the solutions to $\cos 3y = 0$ are $y = \pm\frac{\pi}{6} \approx \pm 0.524$. Using a graph (or a calculator or CAS), we estimate the roots of $3 \cos 3x - 2x \sin 3x$ for $|x| \leq 1$ to be

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x \approx \pm 0.430$. $f_y = 0$ implies $\sin 3x = 0$, so $x = 0$, or $3\sin 3y + 2y\cos 3y = 0$. From a graph (or calculator or CAS), the roots of $3\sin 3y + 2y\cos 3y$ between -1 and 1 are approximately 0 and ± 0.872. So to three decimal places, f has critical points at $(\pm 0.430, 0)$, $(0.430, \pm 0.872)$, $(-0.430, \pm 0.872)$, and $(0, \pm 0.524)$. Now

$$f_{xx} = 20e^{-x^2-y^2}\cos 3y[(4x^2 - 11)\sin 3x - 12x\cos 3x]$$

$$f_{xy} = -20e^{-x^2-y^2}(3\cos 3x - 2x\sin 3x)(3\sin 3y + 2y\cos 3y)$$

$$f_{yy} = 20e^{-x^2-y^2}\sin 3x[(4y^2 - 11)\cos 3y - 12y\sin 3y]$$

and $D = f_{xx}f_{yy} - f_{xy}^2$. Then $D(\pm 0.430, 0) > 0$, $f_{xx}(0.430, 0) < 0$, $f_{xx}(-0.430, 0) > 0$, $D(0.430, \pm 0.872) > 0$, $f_{xx}(0.430, \pm 0.872) > 0$, $D(-0.430, \pm 0.872) > 0$, $f_{xx}(-0.430, \pm 0.872) < 0$, and $D(0, \pm 0.524) < 0$, so $f(0.430, 0) \approx 15.973$ and $f(-0.430, \pm 0.872) \approx 6.459$ are local maxima, $f(-0.430, 0) \approx -15.973$ and $f(0.430, \pm 0.872) \approx -6.459$ are local minima, and $(0, \pm 0.524)$ are saddle points. The highest point on the graph is approximately $(0.430, 0, 15.973)$ and the lowest point is approximately $(-0.430, 0, -15.973)$.



31. Since f is a polynomial it is continuous on D, so an absolute maximum and minimum exist. Here $f_x = 2x - 2$, $f_y = 2y$, and setting $f_x = f_y = 0$ gives $(1, 0)$ as the only critical point (which is inside D), where $f(1, 0) = -1$. Along L_1: $x = 0$ and $f(0, y) = y^2$ for $-2 \le y \le 2$, a quadratic function which attains its minimum at $y = 0$, where $f(0, 0) = 0$, and its maximum at $y = \pm 2$, where $f(0, \pm 2) = 4$. Along L_2: $y = x - 2$ for $0 \le x \le 2$, and $f(x, x - 2) = 2x^2 - 6x + 4 = 2\left(x - \frac{3}{2}\right)^2 - \frac{1}{2}$, a quadratic which attains its minimum at $x = \frac{3}{2}$, where $f\left(\frac{3}{2}, -\frac{1}{2}\right) = -\frac{1}{2}$, and its maximum at $x = 0$, where $f(0, -2) = 4$. Along L_3: $y = 2 - x$ for $0 \le x \le 2$, and $f(x, 2 - x) = 2x^2 - 6x + 4 = 2\left(x - \frac{3}{2}\right)^2 - \frac{1}{2}$, a quadratic which attains its minimum at $x = \frac{3}{2}$, where $f\left(\frac{3}{2}, \frac{1}{2}\right) = -\frac{1}{2}$, and its maximum at $x = 0$, where $f(0, 2) = 4$. Thus the absolute maximum of f on D is $f(0, \pm 2) = 4$ and the absolute minimum is $f(1, 0) = -1$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

32. Since f is a polynomial it is continuous on D, so an absolute maximum and minimum exist. $f_x = 1 - y$, $f_y = 1 - x$, and setting $f_x = f_y = 0$ gives $(1, 1)$ as the only critical point (which is inside D), where $f(1, 1) = 1$. Along L_1: $y = 0$ and $f(x, 0) = x$ for $0 \le x \le 4$, an increasing function in x, so the maximum value is $f(4, 0) = 4$ and the minimum value is $f(0, 0) = 0$. Along L_2: $y = 2 - \frac{1}{2}x$ and $f\left(x, 2 - \frac{1}{2}x\right) = \frac{1}{2}x^2 - \frac{3}{2}x + 2 = \frac{1}{2}\left(x - \frac{3}{2}\right)^2 + \frac{7}{8}$ for $0 \le x \le 4$, a quadratic function which has a minimum at $x = \frac{3}{2}$, where $f\left(\frac{3}{2}, \frac{5}{4}\right) = \frac{7}{8}$, and a maximum at $x = 4$, where $f(4, 0) = 4$.

Along L_3: $x = 0$ and $f(0, y) = y$ for $0 \le y \le 2$, an increasing function in y, so the maximum value is $f(0, 2) = 2$ and the minimum value is $f(0, 0) = 0$. Thus the absolute maximum of f on D is $f(4, 0) = 4$ and the absolute minimum is $f(0, 0) = 0$.



33. $f_x(x, y) = 2x + 2xy$, $f_y(x, y) = 2y + x^2$, and setting $f_x = f_y = 0$ gives $(0, 0)$ as the only critical point in D, with $f(0, 0) = 4$.

On L_1: $y = -1$, $f(x, -1) = 5$, a constant.

On L_2: $x = 1$, $f(1, y) = y^2 + y + 5$, a quadratic in y which attains its maximum at $(1, 1)$, $f(1, 1) = 7$ and its minimum at $\left(1, -\frac{1}{2}\right)$, $f\left(1, -\frac{1}{2}\right) = \frac{19}{4}$.

On L_3: $f(x, 1) = 2x^2 + 5$ which attains its maximum at $(-1, 1)$ and $(1, 1)$ with $f(\pm 1, 1) = 7$ and its minimum at $(0, 1)$, $f(0, 1) = 5$.



On L_4: $f(-1, y) = y^2 + y + 5$ with maximum at $(-1, 1)$, $f(-1, 1) = 7$ and minimum at $\left(-1, -\frac{1}{2}\right)$, $f\left(-1, -\frac{1}{2}\right) = \frac{19}{4}$. Thus the absolute maximum is attained at both $(\pm 1, 1)$ with $f(\pm 1, 1) = 7$ and the absolute minimum on D is attained at $(0, 0)$ with $f(0, 0) = 4$.

34. $f(x, y) = x^2 + xy + y^2 - 6y \quad \Rightarrow \quad f_x = 2x + y$, $f_y = x + 2y - 6$. Then $f_x = 0$ implies $y = -2x$, and substituting into $f_y = 0$ gives $x - 4x - 6 = 0 \quad \Rightarrow \quad x = -2$, so the only critical point is $(-2, 4)$ (which is in D) where $f(-2, 4) = -12$.

Along L_1: $y = 0$, so $f(x, 0) = x^2$, $-3 \le x \le 3$, which has a maximum value at $x = \pm 3$ where $f(\pm 3, 0) = 9$ and a minimum value at $x = 0$, where $f(0, 0) = 0$. Along L_2: $x = 3$, so $f(3, y) = 9 - 3y + y^2 = \left(y - \frac{3}{2}\right)^2 + \frac{27}{4}$, $0 \le y \le 5$, which has a maximum value at $y = 5$ where $f(3, 5) = 19$ and a minimum value at $y = \frac{3}{2}$ where $f\left(3, \frac{3}{2}\right) = \frac{27}{4}$.

Along L_3: $y = 5$, so $f(x, 5) = x^2 + 5x - 5 = \left(x + \frac{5}{2}\right)^2 - \frac{45}{4}$, $-3 \le x \le 3$, which has a maximum value at $x = 3$ where $f(3, 5) = 19$ and a minimum value at $x = -\frac{5}{2}$, where $f\left(-\frac{5}{2}, 5\right) = -\frac{45}{4}$. Along L_4: $x = -3$, so $f(-3, y) = 9 - 9y + y^2 = \left(y - \frac{9}{2}\right)^2 - \frac{45}{4}$, $0 \le y \le 5$, which has a maximum value at $y = 0$ where $f(-3, 0) = 9$ and a minimum value at $y = \frac{9}{2}$ where $f\left(-3, \frac{9}{2}\right) = -\frac{45}{4}$. Thus the absolute maximum of f on D is $f(3, 5) = 19$ and the absolute minimum is $f(-2, 4) = -12$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $f(x,y) = x^2 + 2y^2 - 2x - 4y + 1 \quad \Rightarrow \quad f_x = 2x - 2,\ f_y = 4y - 4$. Setting $f_x = 0$ and $f_y = 0$ gives $(1,1)$ as the only critical point (which is inside D), where $f(1,1) = -2$. Along L_1: $y = 0$, so $f(x,0) = x^2 - 2x + 1 = (x-1)^2$, $0 \le x \le 2$, which has a maximum value both at $x = 0$ and $x = 2$ where $f(0,0) = f(2,0) = 1$ and a minimum value at $x = 1$, where $f(1,0) = 0$. Along L_2: $x = 2$, so $f(2,y) = 2y^2 - 4y + 1 = 2(y-1)^2 - 1$, $0 \le y \le 3$, which has a maximum value at $y = 3$ where $f(2,3) = 7$ and a minimum value at $y = 1$ where $f(2,1) = -1$. Along L_3: $y = 3$, so $f(x,3) = x^2 - 2x + 7 = (x-1)^2 + 6$, $0 \le x \le 2$, which has a maximum value both at $x = 0$ and $x = 2$ where $f(0,3) = f(2,3) = 7$ and a minimum value at $x = 1$, where $f(1,3) = 6$. Along L_4: $x = 0$, so $f(0,y) = 2y^2 - 4y + 1 = 2(y-1)^2 - 1$, $0 \le y \le 3$, which has a maximum value at $y = 3$ where $f(0,3) = 7$ and a minimum value at $y = 1$ where $f(0,1) = -1$. Thus the absolute maximum is attained at both $(0,3)$ and $(2,3)$, where $f(0,3) = f(2,3) = 7$, and the absolute minimum is $f(1,1) = -2$.



36. $f_x = y^2$ and $f_y = 2xy$, and since $f_x = 0 \quad \Leftrightarrow \quad y = 0$, there are no critical points in the interior of D. Along L_1: $y = 0$ and $f(x,0) = 0$. Along L_2: $x = 0$ and $f(0,y) = 0$. Along L_3: $y = \sqrt{3 - x^2}$, so let $g(x) = f\left(x, \sqrt{3-x^2}\right) = 3x - x^3$ for $0 \le x \le \sqrt{3}$. Then $g'(x) = 3 - 3x^2 = 0 \quad \Leftrightarrow \quad x = 1$. The maximum value is $f\left(1, \sqrt{2}\right) = 2$ and the minimum occurs both at $x = 0$ and $x = \sqrt{3}$ where $f\left(0, \sqrt{3}\right) = f\left(\sqrt{3}, 0\right) = 0$. Thus the absolute maximum of f on D is $f\left(1, \sqrt{2}\right) = 2$, and the absolute minimum is 0 which occurs at all points along L_1 and L_2.



37. $f_x(x,y) = 6x^2$ and $f_y(x,y) = 4y^3$. And so $f_x = 0$ and $f_y = 0$ only occur when $x = y = 0$. Hence, the only critical point inside the disk is at $x = y = 0$ where $f(0,0) = 0$. Now on the circle $x^2 + y^2 = 1$, $y^2 = 1 - x^2$ so let $g(x) = f(x,y) = 2x^3 + (1 - x^2)^2 = x^4 + 2x^3 - 2x^2 + 1$, $-1 \le x \le 1$. Then $g'(x) = 4x^3 + 6x^2 - 4x = 0 \quad \Rightarrow \quad x = 0$, -2, or $\frac{1}{2}$. $f(0, \pm 1) = g(0) = 1$, $f\left(\frac{1}{2}, \pm\frac{\sqrt{3}}{2}\right) = g\left(\frac{1}{2}\right) = \frac{13}{16}$, and $(-2,-3)$ is not in D. Checking the endpoints, we get $f(-1,0) = g(-1) = -2$ and $f(1,0) = g(1) = 2$. Thus the absolute maximum and minimum of f on D are $f(1,0) = 2$ and $f(-1,0) = -2$.

Another method: On the boundary $x^2 + y^2 = 1$ we can write $x = \cos\theta$, $y = \sin\theta$, so $f(\cos\theta, \sin\theta) = 2\cos^3\theta + \sin^4\theta$, $0 \le \theta \le 2\pi$.

38. $f_x(x,y) = 3x^2 - 3$ and $f_y(x,y) = -3y^2 + 12$ and the critical points are $(1,2)$, $(1,-2)$, $(-1,2)$, and $(-1,-2)$. But only $(1,2)$ and $(-1,2)$ are in D and $f(1,2) = 14$, $f(-1,2) = 18$. Along L_1: $x = -2$ and $f(-2,y) = -2 - y^3 + 12y$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$-2 \le y \le 3$, which has a maximum at $y = 2$ where $f(-2, 2) = 14$ and a minimum at $y = -2$ where $f(-2, -2) = -18$. Along L_2: $x = 2$ and $f(2, y) = 2 - y^3 + 12y$, $2 \le y \le 3$, which has a maximum at $y = 2$ where $f(2, 2) = 18$ and a minimum at $y = 3$ where $f(2, 3) = 11$. Along L_3: $y = 3$ and $f(x, 3) = x^3 - 3x + 9$, $-2 \le x \le 2$, which has a maximum at $x = -1$ and $x = 2$ where $f(-1, 3) = f(2, 3) = 11$ and a minimum at $x = 1$ and $x = -2$ where $f(1, 3) = f(-2, 3) = 7$. Along L_4: $y = x$ and $f(x, x) = 9x$, $-2 \le x \le 2$, which has a maximum at $x = 2$ where $f(2, 2) = 18$ and a minimum at $x = -2$ where $f(-2, -2) = -18$. So the absolute maximum value of f on D is $f(2, 2) = 18$ and the minimum is $f(-2, -2) = -18$.



39. $f(x, y) = -(x^2 - 1)^2 - (x^2 y - x - 1)^2 \Rightarrow f_x(x, y) = -2(x^2 - 1)(2x) - 2(x^2 y - x - 1)(2xy - 1)$ and $f_y(x, y) = -2(x^2 y - x - 1)x^2$. Setting $f_y(x, y) = 0$ gives either $x = 0$ or $x^2 y - x - 1 = 0$.

There are no critical points for $x = 0$, since $f_x(0, y) = -2$, so we set $x^2 y - x - 1 = 0 \Leftrightarrow y = \dfrac{x + 1}{x^2}$ $[x \ne 0]$, so $f_x\left(x, \dfrac{x+1}{x^2}\right) = -2(x^2 - 1)(2x) - 2\left(x^2 \dfrac{x+1}{x^2} - x - 1\right)\left(2x \dfrac{x+1}{x^2} - 1\right) = -4x(x^2 - 1)$. Therefore $f_x(x, y) = f_y(x, y) = 0$ at the points $(1, 2)$ and $(-1, 0)$. To classify these critical points, we calculate $f_{xx}(x, y) = -12x^2 - 12x^2 y^2 + 12xy + 4y + 2$, $f_{yy}(x, y) = -2x^4$, and $f_{xy}(x, y) = -8x^3 y + 6x^2 + 4x$. In order to use the Second Derivatives Test we calculate $D(-1, 0) = f_{xx}(-1, 0)\, f_{yy}(-1, 0) - [f_{xy}(-1, 0)]^2 = 16 > 0$, $f_{xx}(-1, 0) = -10 < 0$, $D(1, 2) = 16 > 0$, and $f_{xx}(1, 2) = -26 < 0$, so both $(-1, 0)$ and $(1, 2)$ give local maxima.



40. $f(x, y) = 3xe^y - x^3 - e^{3y}$ is differentiable everywhere, so the requirement for critical points is that $f_x = 3e^y - 3x^2 = 0$ **(1)** and $f_y = 3xe^y - 3e^{3y} = 0$ **(2)**. From **(1)** we obtain $e^y = x^2$, and then **(2)** gives $3x^3 - 3x^6 = 0 \Rightarrow x = 1$ or 0, but only $x = 1$ is valid, since $x = 0$ makes **(1)** impossible. So substituting $x = 1$ into **(1)** gives $y = 0$, and the only critical point is $(1, 0)$.



The Second Derivatives Test shows that this gives a local maximum, since $D(1, 0) = \left[-6x(3xe^y - 9e^{3y}) - (3e^y)^2\right]_{(1,0)} = 27 > 0$ and $f_{xx}(1, 0) = [-6x]_{(1,0)} = -6 < 0$. But $f(1, 0) = 1$ is not an absolute maximum because, for instance, $f(-3, 0) = 17$. This can also be seen from the graph.

41. Let d be the distance from $(2, 0, -3)$ to any point (x, y, z) on the plane $x + y + z = 1$, so $d = \sqrt{(x-2)^2 + y^2 + (z+3)^2}$ where $z = 1 - x - y$, and we minimize $d^2 = f(x, y) = (x - 2)^2 + y^2 + (4 - x - y)^2$. Then

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f_x(x,y) = 2(x-2) + 2(4-x-y)(-1) = 4x + 2y - 12$, $f_y(x,y) = 2y + 2(4-x-y)(-1) = 2x + 4y - 8$. Solving $4x + 2y - 12 = 0$ and $2x + 4y - 8 = 0$ simultaneously gives $x = \frac{8}{3}$, $y = \frac{2}{3}$, so the only critical point is $\left(\frac{8}{3}, \frac{2}{3}\right)$. An absolute minimum exists (since there is a minimum distance from the point to the plane) and it must occur at a critical point, so the shortest distance occurs for $x = \frac{8}{3}$, $y = \frac{2}{3}$ for which $d = \sqrt{\left(\frac{8}{3} - 2\right)^2 + \left(\frac{2}{3}\right)^2 + \left(4 - \frac{8}{3} - \frac{2}{3}\right)^2} = \sqrt{\frac{4}{3}} = \frac{2}{\sqrt{3}}$.

42. Here the distance d from a point on the plane to the point $(0,1,1)$ is $d = \sqrt{x^2 + (y-1)^2 + (z-1)^2}$, where $z = 2 - \frac{1}{3}x + \frac{2}{3}y$. We can minimize $d^2 = f(x,y) = x^2 + (y-1)^2 + \left(1 - \frac{1}{3}x + \frac{2}{3}y\right)^2$, so $f_x(x,y) = 2x + 2\left(1 - \frac{1}{3}x + \frac{2}{3}y\right)\left(-\frac{1}{3}\right) = \frac{20}{9}x - \frac{4}{9}y - \frac{2}{3}$ and $f_y(x,y) = 2(y-1) + 2\left(1 - \frac{1}{3}x + \frac{2}{3}y\right)\left(\frac{2}{3}\right) = -\frac{4}{9}x + \frac{26}{9}y - \frac{2}{3}$. Solving $\frac{20}{9}x - \frac{4}{9}y - \frac{2}{3} = 0$ and $-\frac{4}{9}x + \frac{26}{9}y - \frac{2}{3} = 0$ simultaneously gives $x = \frac{5}{14}$ and $y = \frac{2}{7}$, so the only critical point is $\left(\frac{5}{14}, \frac{2}{7}\right)$.

This point must correspond to the minimum distance, so the point on the plane closest to $(0,1,1)$ is $\left(\frac{5}{14}, \frac{2}{7}, \frac{29}{14}\right)$.

43. Let d be the distance from the point $(4,2,0)$ to any point (x,y,z) on the cone, so $d = \sqrt{(x-4)^2 + (y-2)^2 + z^2}$ where $z^2 = x^2 + y^2$, and we minimize $d^2 = (x-4)^2 + (y-2)^2 + x^2 + y^2 = f(x,y)$. Then $f_x(x,y) = 2(x-4) + 2x = 4x - 8$, $f_y(x,y) = 2(y-2) + 2y = 4y - 4$, and the critical points occur when $f_x = 0 \Rightarrow x = 2$, $f_y = 0 \Rightarrow y = 1$. Thus the only critical point is $(2,1)$. An absolute minimum exists (since there is a minimum distance from the cone to the point) which must occur at a critical point, so the points on the cone closest to $(4,2,0)$ are $\left(2, 1, \pm\sqrt{5}\right)$.

44. The distance from the origin to a point (x,y,z) on the surface is $d = \sqrt{x^2 + y^2 + z^2}$ where $y^2 = 9 + xz$, so we minimize $d^2 = x^2 + 9 + xz + z^2 = f(x,z)$. Then $f_x = 2x + z$, $f_z = x + 2z$, and $f_x = 0$, $f_z = 0 \Rightarrow x = 0, z = 0$, so the only critical point is $(0,0)$. $D(0,0) = (2)(2) - 1 = 3 > 0$ with $f_{xx}(0,0) = 2 > 0$, so this is a minimum. Thus $y^2 = 9 + 0 \Rightarrow y = \pm 3$ and the points on the surface closest to the origin are $(0, \pm 3, 0)$.

45. Let x, y, z be the positive numbers. Then $x + y + z = 100 \Rightarrow z = 100 - x - y$, and we want to maximize $xyz = xy(100 - x - y) = 100xy - x^2y - xy^2 = f(x,y)$ for $0 < x, y, z < 100$. $f_x = 100y - 2xy - y^2$, $f_y = 100x - x^2 - 2xy$, $f_{xx} = -2y$, $f_{yy} = -2x$, $f_{xy} = 100 - 2x - 2y$. Then $f_x = 0$ implies $y(100 - 2x - y) = 0 \Rightarrow y = 100 - 2x$ (since $y > 0$). Substituting into $f_y = 0$ gives $x[100 - x - 2(100 - 2x)] = 0 \Rightarrow 3x - 100 = 0$ (since $x > 0$) $\Rightarrow x = \frac{100}{3}$. Then $y = 100 - 2\left(\frac{100}{3}\right) = \frac{100}{3}$, and the only critical point is $\left(\frac{100}{3}, \frac{100}{3}\right)$.

$D\left(\frac{100}{3}, \frac{100}{3}\right) = \left(-\frac{200}{3}\right)\left(-\frac{200}{3}\right) - \left(-\frac{100}{3}\right)^2 = \frac{10{,}000}{3} > 0$ and $f_{xx}\left(\frac{100}{3}, \frac{100}{3}\right) = -\frac{200}{3} < 0$. Thus $f\left(\frac{100}{3}, \frac{100}{3}\right)$ is a local maximum. It is also the absolute maximum (compare to the values of f as x, y, or $z \to 0$ or 100), so the numbers are $x = y = z = \frac{100}{3}$.

46. Let x, y, z, be the positive numbers. Then $x + y + z = 12$ and we want to minimize $x^2 + y^2 + z^2 = x^2 + y^2 + (12 - x - y)^2 = f(x,y)$ for $0 < x, y < 12$. $f_x = 2x + 2(12 - x - y)(-1) = 4x + 2y - 24$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$f_y = 2y + 2(12 - x - y)(-1) = 2x + 4y - 24$, $f_{xx} = 4$, $f_{xy} = 2$, $f_{yy} = 4$. Then $f_x = 0$ implies $4x + 2y = 24$ or $y = 12 - 2x$ and substituting into $f_y = 0$ gives $2x + 4(12 - 2x) = 24 \quad \Rightarrow \quad 6x = 24 \quad \Rightarrow \quad x = 4$ and then $y = 4$, so the only critical point is $(4, 4)$. $D(4, 4) = 16 - 4 > 0$ and $f_{xx}(4, 4) = 4 > 0$, so $f(4, 4)$ is a local minimum. $f(4, 4)$ is also the absolute minimum [compare to the values of f as $x, y \to 0$ or 12] so the numbers are $x = y = z = 4$.

47. Center the sphere at the origin so that its equation is $x^2 + y^2 + z^2 = r^2$, and orient the inscribed rectangular box so that its edges are parallel to the coordinate axes. Any vertex of the box satisfies $x^2 + y^2 + z^2 = r^2$, so take (x, y, z) to be the vertex in the first octant. Then the box has length $2x$, width $2y$, and height $2z = 2\sqrt{r^2 - x^2 - y^2}$ with volume given by $V(x, y) = (2x)(2y)\left(2\sqrt{r^2 - x^2 - y^2}\right) = 8xy\sqrt{r^2 - x^2 - y^2}$ for $0 < x < r$, $0 < y < r$. Then

$V_x = (8xy) \cdot \frac{1}{2}(r^2 - x^2 - y^2)^{\ 1/2}(-2x) + \sqrt{r^2 - x^2 - y^2} \cdot 8y = \dfrac{8y(r^2 - 2x^2 - y^2)}{\sqrt{r^2 - x^2 - y^2}}$ and $V_y = \dfrac{8x(r^2 - x^2 - 2y^2)}{\sqrt{r^2 - x^2 - y^2}}$.

Setting $V_x = 0$ gives $y = 0$ or $2x^2 + y^2 = r^2$, but $y > 0$ so only the latter solution applies. Similarly, $V_y = 0$ with $x > 0$ implies $x^2 + 2y^2 = r^2$. Substituting, we have $2x^2 + y^2 = x^2 + 2y^2 \quad \Rightarrow \quad x^2 = y^2 \quad \Rightarrow \quad y = x$. Then $x^2 + 2y^2 = r^2 \quad \Rightarrow \quad 3x^2 = r^2 \quad \Rightarrow \quad x = \sqrt{r^2/3} = r/\sqrt{3} = y$. Thus the only critical point is $\left(r/\sqrt{3}, r/\sqrt{3}\right)$. There must be a maximum volume and here it must occur at a critical point, so the maximum volume occurs when $x = y = r/\sqrt{3}$ and the maximum volume is $V\left(\frac{r}{\sqrt{3}}, \frac{r}{\sqrt{3}}\right) = 8\left(\frac{r}{\sqrt{3}}\right)\left(\frac{r}{\sqrt{3}}\right)\sqrt{r^2 - \left(\frac{r}{\sqrt{3}}\right)^2 - \left(\frac{r}{\sqrt{3}}\right)^2} = \dfrac{8}{3\sqrt{3}}r^3$.

48. Let x, y, and z be the dimensions of the box. We wish to minimize surface area $= 2xy + 2xz + 2yz$, but we have volume $= xyz = 1000 \quad \Rightarrow \quad z = \dfrac{1000}{xy}$ so we minimize

$f(x, y) = 2xy + 2x\left(\dfrac{1000}{xy}\right) + 2y\left(\dfrac{1000}{xy}\right) = 2xy + \dfrac{2000}{y} + \dfrac{2000}{x}$. Then $f_x = 2y - \dfrac{2000}{x^2}$ and $f_y = 2x - \dfrac{2000}{y^2}$. Setting $f_x = 0$ implies $y = \dfrac{1000}{x^2}$ and substituting into $f_y = 0$ gives $x - \dfrac{x^4}{1000} = 0 \quad \Rightarrow \quad x^3 = 1000$ [since $x \neq 0$] $\quad \Rightarrow \quad x = 10$. The surface area has a minimum but no maximum and it must occur at a critical point, so the minimal surface area occurs for a box with dimensions $x = 10$ cm, $y = 1000/10^2 = 10$ cm, $z = 1000/10^2 = 10$ cm.

49. Maximize $f(x, y) = \dfrac{xy}{3}(6 - x - 2y)$, then the maximum volume is $V = xyz$.

$f_x = \frac{1}{3}(6y - 2xy - y^2) = \frac{1}{3}y(6 - 2x - 2y)$ and $f_y = \frac{1}{3}x(6 - x - 4y)$. Setting $f_x = 0$ and $f_y = 0$ gives the critical point $(2, 1)$ which geometrically must give a maximum. Thus the volume of the largest such box is $V = (2)(1)\left(\frac{2}{3}\right) = \frac{4}{3}$.

50. Surface area $= 2(xy + xz + yz) = 64$ cm^2, so $xy + xz + yz = 32$ or $z = \dfrac{32 - xy}{x + y}$. Maximize the volume $f(x, y) = xy\dfrac{32 - xy}{x + y}$. Then $f_x = \dfrac{32y^2 - 2xy^3 - x^2y^2}{(x + y)^2} = y^2\dfrac{32 - 2xy - x^2}{(x + y)^2}$ and $f_y = x^2\dfrac{32 - 2xy - y^2}{(x + y)^2}$. Setting

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f_x = 0$ implies $y = \dfrac{32 - x^2}{2x}$ and substituting into $f_y = 0$ gives $32(4x^2) - (32 - x^2)(4x^2) - (32 - x^2)^2 = 0$ or $3x^4 + 64x^2 - (32)^2 = 0$. Thus $x^2 = \frac{64}{6}$ or $x = \frac{8}{\sqrt{6}}$, $y = \frac{64/3}{16/\sqrt{6}} = \frac{8}{\sqrt{6}}$ and $z = \frac{8}{\sqrt{6}}$. Thus the box is a cube with edge length $\frac{8}{\sqrt{6}}$ cm.

51. Let the dimensions be x, y, and z; then $4x + 4y + 4z = c$ and the volume is $V = xyz = xy\left(\frac{1}{4}c - x - y\right) = \frac{1}{4}cxy - x^2y - xy^2$, $x > 0$, $y > 0$. Then $V_x = \frac{1}{4}cy - 2xy - y^2$ and $V_y = \frac{1}{4}cx - x^2 - 2xy$, so $V_x = 0 = V_y$ when $2x + y = \frac{1}{4}c$ and $x + 2y = \frac{1}{4}c$. Solving, we get $x = \frac{1}{12}c$, $y = \frac{1}{12}c$ and $z = \frac{1}{4}c - x - y = \frac{1}{12}c$. From the geometrical nature of the problem, this critical point must give an absolute maximum. Thus the box is a cube with edge length $\frac{1}{12}c$.

52. The cost equals $5xy + 2(xz + yz)$ and $xyz = V$, so $C(x, y) = 5xy + 2V(x + y)/(xy) = 5xy + 2V(x^{-1} + y^{-1})$. Then $C_x = 5y - 2Vx^{-2}$, $C_y = 5x - 2\,Vy^{-2}$, $C_x = 0$ implies $y = 2\,V/(5x^2)$, $C_y = 0$ implies $x = \sqrt[3]{\frac{2}{5}V} = y$. Thus the dimensions of the aquarium which minimize the cost are $x = y = \sqrt[3]{\frac{2}{5}V}$ units, $z = V^{1/3}\left(\frac{5}{2}\right)^{2/3}$.

53. Let the dimensions be x, y and z, then minimize $xy + 2(xz + yz)$ if $xyz = 32{,}000$ cm^3. Then $f(x, y) = xy + [64{,}000(x + y)/xy] = xy + 64{,}000(x^{-1} + y^{-1})$, $f_x = y - 64{,}000x^{-2}$, $f_y = x - 64{,}000y^{-2}$. And $f_x = 0$ implies $y = 64{,}000/x^2$; substituting into $f_y = 0$ implies $x^3 = 64{,}000$ or $x = 40$ and then $y = 40$. Now $D(x, y) = [(2)(64{,}000)]^2x^{-3}y^{-3} - 1 > 0$ for $(40, 40)$ and $f_{xx}(40, 40) > 0$ so this is indeed a minimum. Thus the dimensions of the box are $x = y = 40$ cm, $z = 20$ cm.

54. Let x be the length of the north and south walls, y the length of the east and west walls, and z the height of the building. The heat loss is given by $h = 10(2yz) + 8(2xz) + 1(xy) + 5(xy) = 6xy + 16xz + 20yz$. The volume is 4000 m^3, so $xyz = 4000$, and we substitute $z = \frac{4000}{xy}$ to obtain the heat loss function $h(x, y) = 6xy + 80{,}000/x + 64{,}000/y$.

(a) Since $z = \frac{4000}{xy} \geq 4$, $xy \leq 1000 \;\Rightarrow\; y \leq 1000/x$. Also $x \geq 30$ and $y \geq 30$, so the domain of h is $D = \{(x, y) \mid x \geq 30, 30 \leq y \leq 1000/x\}$.



(b) $h(x, y) = 6xy + 80{,}000x^{-1} + 64{,}000y^{-1} \;\Rightarrow$

$h_x = 6y - 80{,}000x^{-2}$, $h_y = 6x - 64{,}000y^{-2}$.

$h_x = 0$ implies $6x^2y = 80{,}000 \;\Rightarrow\; y = \dfrac{80{,}000}{6x^2}$ and substituting into

$h_y = 0$ gives $6x = 64{,}000\left(\dfrac{6x^2}{80{,}000}\right)^2 \;\Rightarrow\; x^3 = \dfrac{80{,}000^2}{6 \cdot 64{,}000} = \dfrac{50{,}000}{3}$, so

$x = \sqrt[3]{\dfrac{50{,}000}{3}} = 10\sqrt[3]{\dfrac{50}{3}} \;\Rightarrow\; y = \dfrac{80}{\sqrt[3]{60}}$, and the only critical point of h is $\left(10\sqrt[3]{\dfrac{50}{3}}, \dfrac{80}{\sqrt[3]{60}}\right) \approx (25.54, 20.43)$

which is not in D. Next we check the boundary of D.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

On L_1: $y = 30$, $h(x, 30) = 180x + 80{,}000/x + 6400/3$, $30 \le x \le \frac{100}{3}$. Since $h'(x, 30) = 180 - 80{,}000/x^2 > 0$ for $30 \le x \le \frac{100}{3}$, $h(x, 30)$ is an increasing function with minimum $h(30, 30) = 10{,}200$ and maximum $h\left(\frac{100}{3}, 30\right) \approx 10{,}533$.

On L_2: $y = 1000/x$, $h(x, 1000/x) = 6000 + 64x + 80{,}000/x$, $30 \le x \le \frac{100}{3}$.

Since $h'(x, 1000/x) = 64 - 80{,}000/x^2 < 0$ for $30 \le x \le \frac{100}{3}$, $h(x, 1000/x)$ is a decreasing function with minimum $h\left(\frac{100}{3}, 30\right) \approx 10{,}533$ and maximum $h\left(30, \frac{100}{3}\right) \approx 10{,}587$.

On L_3: $x = 30$, $h(30, y) = 180y + 64{,}000/y + 8000/3$, $30 \le y \le \frac{100}{3}$. $h'(30, y) = 180 - 64{,}000/y^2 > 0$ for $30 \le y \le \frac{100}{3}$, so $h(30, y)$ is an increasing function of y with minimum $h(30, 30) = 10{,}200$ and maximum $h\left(30, \frac{100}{3}\right) \approx 10{,}587$.

Thus the absolute minimum of h is $h(30, 30) = 10{,}200$, and the dimensions of the building that minimize heat loss are walls 30 m in length and height $\frac{4000}{30^2} = \frac{40}{9} \approx 4.44$ m.

(c) From part (b), the only critical point of h, which gives a local (and absolute) minimum, is approximately $h(25.54, 20.43) \approx 9396$. So a building of volume 4000 m^2 with dimensions $x \approx 25.54$ m, $y \approx 20.43$ m, $z \approx \frac{4000}{(25.54)(20.43)} \approx 7.67$ m has the least amount of heat loss.

55. Let x, y, z be the dimensions of the rectangular box. Then the volume of the box is xyz and

$L = \sqrt{x^2 + y^2 + z^2} \;\Rightarrow\; L^2 = x^2 + y^2 + z^2 \;\Rightarrow\; z = \sqrt{L^2 - x^2 - y^2}$.

Substituting, we have volume $V(x, y) = xy\sqrt{L^2 - x^2 - y^2}$ $(x, y > 0)$.

$V_x = xy \cdot \frac{1}{2}(L^2 - x^2 - y^2)^{-1/2}(-2x) + y\sqrt{L^2 - x^2 - y^2} = y\sqrt{L^2 - x^2 - y^2} - \dfrac{x^2 y}{\sqrt{L^2 - x^2 - y^2}}$.

$V_y = x\sqrt{L^2 - x^2 - y^2} - \dfrac{xy^2}{\sqrt{L^2 - x^2 - y^2}}$. $V_x = 0$ implies $y(L^2 - x^2 - y^2) = x^2 y \;\Rightarrow\; y(L^2 - 2x^2 - y^2) = 0 \;\Rightarrow$

$2x^2 + y^2 = L^2$ (since $y > 0$), and $V_y = 0$ implies $x(L^2 - x^2 - y^2) = xy^2 \;\Rightarrow\; x(L^2 - x^2 - 2y^2) = 0 \;\Rightarrow$

$x^2 + 2y^2 = L^2$ (since $x > 0$). Substituting $y^2 = L^2 - 2x^2$ into $x^2 + 2y^2 = L^2$ gives $x^2 + 2L^2 - 4x^2 = L^2 \;\Rightarrow$

$3x^2 = L^2 \;\Rightarrow\; x = L/\sqrt{3}$ (since $x > 0$) and then $y = \sqrt{L^2 - 2\left(L/\sqrt{3}\right)^2} = L/\sqrt{3}$.

So the only critical point is $\left(L/\sqrt{3}, L/\sqrt{3}\right)$ which, from the geometrical nature of the problem, must give an absolute maximum. Thus the maximum volume is $V\left(L/\sqrt{3}, L/\sqrt{3}\right) = \left(L/\sqrt{3}\right)^2\sqrt{L^2 - \left(L/\sqrt{3}\right)^2 - \left(L/\sqrt{3}\right)^2} = L^3/\left(3\sqrt{3}\right)$ cubic units.

56. $Y(N, P) = kNPe^{-N-P} \;\Rightarrow\; Y_N = kP\left[N\left(-e^{-N-P}\right) + e^{-N-P}(1)\right] = kP(1 - N)e^{-N-P}$,

$Y_P = kN\left[P\left(-e^{-N-P}\right) + e^{-N-P}(1)\right] = kN(1 - P)e^{-N-P}$. Here $N \ge 0$ and $P \ge 0$, but if either $N = 0$ or $P = 0$ then

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

the yield is zero. Assuming that $N > 0$ and $P > 0$, $Y_N = 0$ implies $N = 1$ and $Y_P = 0$ implies $P = 1$, so the only critical point in $\{(N, P) \mid N > 0, P > 0\}$ is $(1, 1)$ where $Y(1, 1) = ke^{-2}$.

$D(N, P) = Y_{NN}Y_{PP} - (Y_{NP})^2 = \left[kP(N-2)e^{-N-P}\right]\left[kN(P-2)e^{-N-P}\right] - \left[k(1-N)(1-P)e^{-N-P}\right]^2 \quad \Rightarrow$

$D(1, 1) = \left(-ke^{-2}\right)\left(-ke^{-2}\right) - (0)^2 = k^2e^{-4} > 0$ and $Y_{NN}(1, 1) = -ke^{-2} < 0$, so $Y(1, 1) = ke^{-2}$ is a local maximum. $Y(1, 1)$ is also the absolute maximum (we have only one critical point, and $Y \to 0$ as $N \to 0$ or $P \to 0$ and $Y \to 0$ as N or P grow large) so the best yield is achieved when both the nitrogen and phosphorus levels are 1 (measured in appropriate units).

57. (a) We are given that $p_1 + p_2 + p_3 = 1 \quad \Rightarrow \quad p_3 = 1 - p_1 - p_2$, so

$H = -p_1 \ln p_1 - p_2 \ln p_2 - p_3 \ln p_3 = -p_1 \ln p_1 - p_2 \ln p_2 - (1 - p_1 - p_2) \ln (1 - p_1 - p_2)$.

(b) Because p_i is a proportion we have $0 \le p_i \le 1$, but H is undefined unless $p_1 > 0$, $p_2 > 0$, and $1 - p_1 - p_2 > 0 \quad \Leftrightarrow \quad p_1 + p_2 < 1$. This last restriction forces $p_1 < 1$ and $p_2 < 1$, so the domain of H is $\{(p_1, p_2) \mid 0 < p_1 < 1,\ p_2 < 1 - p_1\}$. It is the interior of the triangle drawn in the figure.



(c) $H_{p_1} = -[p_1 \cdot (1/p_1) + (\ln p_1) \cdot 1] - [(1 - p_1 - p_2) \cdot (-1)/(1 - p_1 - p_2) + \ln (1 - p_1 - p_2) \cdot (-1)]$

$= -1 - \ln p_1 + 1 + \ln (1 - p_1 - p_2) = \ln (1 - p_1 - p_2) - \ln p_1$

Similarly $H_{p_2} = \ln (1 - p_1 - p_2) - \ln p_2$. Then $H_{p_1} = 0$ implies

$\ln (1 - p_1 - p_2) = \ln p_1 \quad \Rightarrow \quad 1 - p_1 - p_2 = p_1 \quad \Rightarrow \quad p_2 = 1 - 2p_1$, and $H_{p_2} = 0$ implies

$\ln (1 - p_1 - p_2) = \ln p_2 \quad \Rightarrow \quad p_1 = 1 - 2p_2$. Substituting, we have $p_1 = 1 - 2(1 - 2p_1) \quad \Rightarrow$

$3p_1 = 1 \quad \Rightarrow \quad p_1 = \frac{1}{3}$, and then $p_2 = 1 - 2\left(\frac{1}{3}\right) = \frac{1}{3}$. Thus the only critical point is $\left(\frac{1}{3}, \frac{1}{3}\right)$.

$D(p_1, p_2) = H_{p_1p_1}H_{p_2p_2} - (H_{p_1p_2})^2 = \left(\dfrac{-1}{1 - p_1 - p_2} - \dfrac{1}{p_1}\right)\left(\dfrac{-1}{1 - p_1 - p_2} - \dfrac{1}{p_2}\right) - \left(\dfrac{-1}{1 - p_1 - p_2}\right)^2$, so

$D\left(\frac{1}{3}, \frac{1}{3}\right) = (-6)(-6) - (-3)^2 = 27 > 0$ and $H_{p_1p_1}\left(\frac{1}{3}, \frac{1}{3}\right) = -6 < 0$. Thus

$H\left(\frac{1}{3}, \frac{1}{3}\right) = -\frac{1}{3}\ln\frac{1}{3} - \frac{1}{3}\ln\frac{1}{3} - \frac{1}{3}\ln\frac{1}{3} = -\ln\frac{1}{3} = \ln 3$ is a local maximum. Here it is also the absolute maximum, so the maximum value of H is $\ln 3$, which occurs for $p_1 = p_2 = p_3 = \frac{1}{3}$ (all three species have equal proportion in the ecosystem).

58. Since $p + q + r = 1$ we can substitute $p = 1 - r - q$ into P giving

$P = P(q, r) = 2(1 - r - q)q + 2(1 - r - q)r + 2rq = 2q - 2q^2 + 2r - 2r^2 - 2rq$. Since p, q and r represent proportions and $p + q + r = 1$, we know $q \ge 0$, $r \ge 0$, and $q + r \le 1$. Thus, we want to find the absolute maximum of the continuous function $P(q, r)$ on the closed set D enclosed by the lines $q = 0$, $r = 0$, and $q + r = 1$. To find any critical points, we set the

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

partial derivatives equal to zero: $P_q(q,r) = 2 - 4q - 2r = 0$ and $P_r(q,r) = 2 - 4r - 2q = 0$. The first equation gives $r = 1 - 2q$, and substituting into the second equation we have $2 - 4(1-2q) - 2q = 0 \;\Rightarrow\; q = \frac{1}{3}$. Then we have one critical point, $\left(\frac{1}{3}, \frac{1}{3}\right)$, where $P\left(\frac{1}{3}, \frac{1}{3}\right) = \frac{2}{3}$. Next we find the maximum values of P on the boundary of D which consists of three line segments. For the segment given by $r = 0$, $0 \le q \le 1$, $P(q,r) = P(q,0) = 2q - 2q^2$, $0 \le q \le 1$. This represents a parabola with maximum value $P\left(\frac{1}{2}, 0\right) = \frac{1}{2}$. On the segment $q = 0$, $0 \le r \le 1$ we have $P(0,r) = 2r - 2r^2$, $0 \le r \le 1$. This represents a parabola with maximum value $P\left(0, \frac{1}{2}\right) = \frac{1}{2}$. Finally, on the segment $q + r = 1$, $0 \le q \le 1$, $P(q,r) = P(q, 1-q) = 2q - 2q^2$, $0 \le q \le 1$ which has a maximum value of $P\left(\frac{1}{2}, \frac{1}{2}\right) = \frac{1}{2}$. Comparing these values with the value of P at the critical point, we see that the absolute maximum value of $P(q,r)$ on D is $\frac{2}{3}$.

59. Note that here the variables are m and b, and $f(m,b) = \sum_{i=1}^{n} [y_i - (mx_i + b)]^2$. Then $f_m = \sum_{i=1}^{n} -2x_i[y_i - (mx_i + b)] = 0$ implies $\sum_{i=1}^{n} \left(x_i y_i - m x_i^2 - b x_i\right) = 0$ or $\sum_{i=1}^{n} x_i y_i = m \sum_{i=1}^{n} x_i^2 + b \sum_{i=1}^{n} x_i$ and $f_b = \sum_{i=1}^{n} -2[y_i - (mx_i + b)] = 0$ implies $\sum_{i=1}^{n} y_i = m \sum_{i=1}^{n} x_i + \sum_{i=1}^{n} b = m\left(\sum_{i=1}^{n} x_i\right) + nb$. Thus we have the two desired equations.

Now $f_{mm} = \sum_{i=1}^{n} 2x_i^2$, $f_{bb} = \sum_{i=1}^{n} 2 = 2n$ and $f_{mb} = \sum_{i=1}^{n} 2x_i$. And $f_{mm}(m,b) > 0$ always and $D(m,b) = 4n\left(\sum_{i=1}^{n} x_i^2\right) - 4\left(\sum_{i=1}^{n} x_i\right)^2 = 4\left[n\left(\sum_{i=1}^{n} x_i^2\right) - \left(\sum_{i=1}^{n} x_i\right)^2\right] > 0$ always so the solutions of these two equations do indeed minimize $\sum_{i=1}^{n} d_i^2$.

60. Any such plane must cut out a tetrahedron in the first octant. We need to minimize the volume of the tetrahedron that passes through the point $(1,2,3)$. Writing the equation of the plane as $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$, the volume of the tetrahedron is given by $V = \frac{abc}{6}$. But $(1,2,3)$ must lie on the plane, so we need $\frac{1}{a} + \frac{2}{b} + \frac{3}{c} = 1$ $(\star)$ and thus can think of c as a function of a and b. Then $V_a = \frac{b}{6}\left(c + a\frac{\partial c}{\partial a}\right)$ and $V_b = \frac{a}{6}\left(c + b\frac{\partial c}{\partial b}\right)$. Differentiating $(\star)$ with respect to a we get $-a^{-2} - 3c^{-2}\frac{\partial c}{\partial a} = 0 \;\Rightarrow\; \frac{\partial c}{\partial a} = \frac{-c^2}{3a^2}$, and differentiating $(\star)$ with respect to b gives $-2b^{-2} - 3c^{-2}\frac{\partial c}{\partial b} = 0 \;\Rightarrow\; \frac{\partial c}{\partial b} = \frac{-2c^2}{3b^2}$. Then $V_a = \frac{b}{6}\left(c + a\frac{-c^2}{3a^2}\right) = 0 \;\Rightarrow\; c = 3a$, and $V_b = \frac{a}{6}\left(c + b\frac{-2c^2}{3b^2}\right) = 0 \;\Rightarrow\; c = \frac{3}{2}b$. Thus $3a = \frac{3}{2}b$ or $b = 2a$. Putting these into $(\star)$ gives $\frac{3}{a} = 1$ or $a = 3$ and then $b = 6$, $c = 9$. Thus the equation of the required plane is $\frac{x}{3} + \frac{y}{6} + \frac{z}{9} = 1$ or $6x + 3y + 2z = 18$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT Designing a Dumpster

Note: The difficulty and results of this project vary widely with the type of container studied. In addition to the variation of basic shapes of containers, dumpsters may include additional constructed parts such as supports, lift pockets, wheels, etc. Also, a CAS or graphing utility may be needed to solve the resulting equations.

Here we present a typical solution for one particular trash Dumpster.

1. The basic shape and dimensions (in inches) of an actual trash Dumpster are as shown in the figure.



The front and back, as well as both sides, have an extra one-inch-wide flap that is folded under and welded to the base. In addition, the side panels each fold over one inch onto the front and back pieces where they are welded. Each side has a rectangular lift pocket, with cross-section 5 by 8 inches, made of the same material. These are attached with an extra one-inch width of steel on both top and bottom where each pocket is welded to the side sheet. All four sides have a "lip" at the top; the front and back panels have an extra 5 inches of steel at the top which is folded outward in three creases to form a rectangular tube. The edge is then welded back to the main sheet. The two sides form a top lip with separate sheets of steel 5 inches wide, similarly bent into three sides and welded to the main sheets (requiring two welds each). These extend beyond the main side sheets by 1.5 inches at each end in order to join with the lips on the front and back panels. The container has a hinged lid, extra steel supports on the base at each corner, metal "fins" serving as extra support for the side lift pockets, and wheels underneath. The volume of the container is $V = \frac{1}{2}(40 + 49) \times 42 \times 72 = 134{,}568$ in^3 or 77.875 ft^3.

2. First, we assume that some aspects of the construction do not change with different dimensions, so they may be considered fixed costs. This includes the lid (with hinges), wheels, and extra steel supports. Also, the upper "lip" we previously described extends beyond the side width to connect to the other pieces. We can safely assume that this extra portion, including any associated welds, costs the same regardless of the container's dimensions, so we will consider just the portion matching the measurement of the side panels in our calculations. We will further assume that the angle of the top of the container should be preserved. Then to compute the variable costs, let x be the width, y the length, and z the height of the front of the container. The back of the container is 9 inches, or $\frac{3}{4}$ ft, taller than the front, so using similar triangles we can say the back panel has height $z + \frac{3}{14}x$. Measuring in feet, we want the volume to remain constant, so $V = \frac{1}{2}\left(z + z + \frac{3}{14}x\right)(x)(y) = xyz + \frac{3}{28}x^2y = 77.875$. To determine a function for the variable cost, we first find the area of each sheet of metal needed. The base has area xy ft^2. The front panel has visible area yz plus $\frac{1}{12}y$ for the portion folded onto the base and $\frac{5}{12}y$ for the steel at the top used to form the lip, so $\left(yz + \frac{1}{2}y\right)$ ft^2 in total. Similarly, the back sheet has area $y\left(z + \frac{3}{14}x\right) + \frac{1}{12}y + \frac{5}{12}y = yz + \frac{3}{14}xy + \frac{1}{2}y$. Each side has visible area $\frac{1}{2}\left[z + \left(z + \frac{3}{14}x\right)\right](x)$, and the sheet includes

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

one-inch flaps folding onto the front and back panels, so with area $\frac{1}{12}z$ and $\frac{1}{12}\left(z+\frac{3}{14}x\right)$, and a one-inch flap to fold onto the base with area $\frac{1}{12}x$. The lift pocket is constructed of a piece of steel 20 inches by x ft (including the 2 extra inches used by the welds). The additional metal used to make the lip at the top of the panel has width 5 inches and length that we can determine using the Pythagorean Theorem: $x^2+\left(\frac{3}{14}x\right)^2=\text{length}^2$, so length $=\frac{\sqrt{205}}{14}x\approx 1.0227x$. Thus the area of steel needed for each side panel is approximately

$$\tfrac{1}{2}\left[z+\left(z+\tfrac{3}{14}x\right)\right](x)+\tfrac{1}{12}z+\tfrac{1}{12}\left(z+\tfrac{3}{14}x\right)+\tfrac{1}{12}x+\tfrac{5}{3}x+\tfrac{5}{12}(1.0227x)\approx xz+\tfrac{3}{28}x^2+\tfrac{1}{6}z+2.194x$$

We also have the following welds:

Weld	Length
Front, back welded to base	$2y$
Sides welded to base	$2x$
Sides welded to front	$2z$
Sides welded to back	$2\left(z+\frac{3}{14}x\right)$
Weld on front and back lip	$2y$
Two welds on each side lip	$4(1.0227x)$
Two welds for each lift pocket	$4x$

Thus the total length of welds needed is

$$2y+2x+2z+2\left(z+\tfrac{3}{14}x\right)+2y+4(1.0227x)+4x\approx 10.519x+4y+4z$$

Finally, the total variable cost is approximately

$$\begin{aligned}0.90(xy)+0.70\left[\left(yz+\tfrac{1}{2}y\right)+\left(yz+\tfrac{3}{14}xy+\tfrac{1}{2}y\right)+2\left(xz+\tfrac{3}{28}x^2+\tfrac{1}{6}z+2.194x\right)\right]+0.18(10.519x+4y+4z)\\ \approx 1.05xy+1.4yz+1.42y+1.4xz+0.15x^2+0.953z+4.965x\end{aligned}$$

We would like to minimize this function while keeping volume constant, so since $xyz+\frac{3}{28}x^2y=77.875$ we can substitute $z=\dfrac{77.875}{xy}-\dfrac{3}{28}x$ giving variable cost as a function of x and y:

$$C(x,y)\approx 0.9xy+\frac{109.0}{x}+1.42y+\frac{109.0}{y}+\frac{74.2}{xy}+4.86x$$

Using a CAS, we solve the system of equations $C'_x(x,y)=0$ and $C'_y(x,y)=0$; the only critical point within an appropriate domain is approximately $(3.58, 5.29)$. From the nature of the function C (or from a graph) we can determine that C has an absolute minimum at $(3.58, 5.29)$, and so the minimum cost is attained for $x\approx 3.58$ ft (or 43.0 in), $y\approx 5.29$ ft (or 63.5 in), and $z\approx\frac{77.875}{3.58(5.29)}-\frac{3}{28}(3.58)\approx 3.73$ ft (or 44.8 in).

3. The fixed cost aspects of the container which we did not include in our calculations, such as the wheels and lid, don't affect the validity of our results. Some of our other assumptions, however, may influence the accuracy of our findings. We simplified the price of the steel sheets to include cuts and bends, and we simplified the price of welding to include the labor and materials. This may not be accurate for areas of the container, such as the lip and lift pockets, that require several cuts, bends, and welds

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

in a relatively small surface area. Consequently, increasing some dimensions of the container may not increase the cost in the same manner as our computations predict. If we do not assume that the angle of the sloped top of the container must be preserved, it is likely that we could further improve our cost. Finally, our results show that the length of the container should be changed to minimize cost; this may not be possible if the two lift pockets must remain a fixed distance apart for handling by machinery.

4. The minimum variable cost using our values found in Problem 2 is $C(3.58, 5.29) \approx \$96.95$, while the current dimensions give an estimated variable cost of $C(3.5, 6.0) \approx \$97.30$. If we determine that our assumptions and simplifications are acceptable, our work shows that a slight savings can be gained by adjusting the dimensions of the container. However, the difference in cost is modest, and may not justify changes in the manufacturing process.

DISCOVERY PROJECT Quadratic Approximations and Critical Points

1. $Q(x,y) = f(a,b) + f_x(a,b)(x-a) + f_y(a,b)(y-b) + \frac{1}{2}f_{xx}(a,b)(x-a)^2$
$+ f_{xy}(a,b)(x-a)(y-b) + \frac{1}{2}f_{yy}(a,b)(y-b)^2,$

so

$$Q_x(x,y) = f_x(a,b) + \tfrac{1}{2}f_{xx}(a,b)(2)(x-a) + f_{xy}(a,b)(y-b) = f_x(a,b) + f_{xx}(a,b)(x-a) + f_{xy}(a,b)(y-b)$$

At (a,b) we have $Q_x(a,b) = f_x(a,b) + f_{xx}(a,b)(a-a) + f_{xy}(a,b)(b-b) = f_x(a,b)$.

Similarly, $Q_y(x,y) = f_y(a,b) + f_{xy}(a,b)(x-a) + f_{yy}(a,b)(y-b) \Rightarrow$

$Q_y(a,b) = f_y(a,b) + f_{xy}(a,b)(a-a) + f_{yy}(a,b)(b-b) = f_y(a,b)$.

For the second-order partial derivatives we have

$$Q_{xx}(x,y) = \frac{\partial}{\partial x}\left[f_x(a,b) + f_{xx}(a,b)(x-a) + f_{xy}(a,b)(y-b)\right] = f_{xx}(a,b) \quad \Rightarrow \quad Q_{xx}(a,b) = f_{xx}(a,b)$$

$$Q_{xy}(x,y) = \frac{\partial}{\partial y}\left[f_x(a,b) + f_{xx}(a,b)(x-a) + f_{xy}(a,b)(y-b)\right] = f_{xy}(a,b) \quad \Rightarrow \quad Q_{xy}(a,b) = f_{xy}(a,b)$$

$$Q_{yy}(x,y) = \frac{\partial}{\partial y}\left[f_y(a,b) + f_{xy}(a,b)(x-a) + f_{yy}(a,b)(y-b)\right] = f_{yy}(a,b) \quad \Rightarrow \quad Q_{yy}(a,b) = f_{yy}(a,b)$$

2. (a) First we find the partial derivatives and values that will be needed:

$$f(x,y) = e^{-x^2-y^2} \qquad f(0,0) = 1$$

$$f_x(x,y) = -2xe^{-x^2-y^2} \qquad f_x(0,0) = 0$$

$$f_y(x,y) = -2ye^{-x^2-y^2} \qquad f_y(0,0) = 0$$

$$f_{xx}(x,y) = (4x^2-2)e^{-x^2-y^2} \qquad f_{xx}(0,0) = -2$$

$$f_{xy}(x,y) = 4xye^{-x^2-y^2} \qquad f_{xy}(0,0) = 0$$

$$f_{yy}(x,y) = (4y^2-2)e^{-x^2-y^2} \qquad f_{yy}(0,0) = -2$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Then the first-degree Taylor polynomial of f at $(0, 0)$ is

$$L(x, y) = f(0, 0) + f_x(0, 0)(x - 0) + f_y(0, 0)(y - 0) = 1 + (0)(x - 0) + (0)(y - 0) = 1$$

The second-degree Taylor polynomial is given by

$$\begin{aligned} Q(x, y) &= f(0, 0) + f_x(0, 0)(x - 0) + f_y(0, 0)(y - 0) + \tfrac{1}{2} f_{xx}(0, 0)(x - 0)^2 \\ &\quad + f_{xy}(0, 0)(x - 0)(y - 0) + \tfrac{1}{2} f_{yy}(0, 0)(y - 0)^2 \\ &= 1 - x^2 - y^2 \end{aligned}$$

(b)



As we see from the graph, L approximates f well only for points (x, y) extremely close to the origin. Q is a much better approximation; the shape of its graph looks similar to that of the graph of f near the origin, and the values of Q appear to be good estimates for the values of f within a significant radius of the origin.

3. (a) First we find the partial derivatives and values that will be needed:

$$\begin{aligned} &f(x, y) = xe^y \qquad f(1, 0) = 1 \qquad f_{xx}(x, y) = 0 \qquad f_{xx}(1, 0) = 0 \\ &f_x(x, y) = e^y \qquad f_x(1, 0) = 1 \qquad f_{xy}(x, y) = e^y \qquad f_{xy}(1, 0) = 1 \\ &f_y(x, y) = xe^y \qquad f_y(1, 0) = 1 \qquad f_{yy}(x, y) = xe^y \qquad f_{yy}(1, 0) = 1 \end{aligned}$$

Then the first-degree Taylor polynomial of f at $(1, 0)$ is

$$L(x, y) = f(1, 0) + f_x(1, 0)(x - 1) + f_y(1, 0)(y - 0) = 1 + (1)(x - 1) + (1)(y - 0) = x + y$$

The second-degree Taylor polynomial is given by

$$\begin{aligned} Q(x, y) &= f(1, 0) + f_x(1, 0)(x - 1) + f_y(1, 0)(y - 0) + \tfrac{1}{2} f_{xx}(1, 0)(x - 1)^2 \\ &\quad + f_{xy}(1, 0)(x - 1)(y - 0) + \tfrac{1}{2} f_{yy}(1, 0)(y - 0)^2 \\ &= \tfrac{1}{2} y^2 + x + xy \end{aligned}$$

(b) $L(0.9, 0.1) = 0.9 + 0.1 = 1.0$

$Q(0.9, 0.1) = \frac{1}{2}(0.1)^2 + 0.9 + (0.9)(0.1) = 0.995$

$f(0.9, 0.1) = 0.9e^{0.1} \approx 0.9947$

(c)



As we see from the graph, L and Q both approximate f reasonably well near the point $(1, 0)$. As we venture farther from the point, the graph of Q follows the shape of the graph of f more closely than L.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a) $f(x,y) = ax^2 + bxy + cy^2 = a\left[x^2 + \frac{b}{a}xy + \frac{c}{a}y^2\right] = a\left[x^2 + \frac{b}{a}xy + \left(\frac{b}{2a}y\right)^2 - \left(\frac{b}{2a}y\right)^2 + \frac{c}{a}y^2\right]$

$= a\left[\left(x + \frac{b}{2a}y\right)^2 - \frac{b^2}{4a^2}y^2 + \frac{c}{a}y^2\right] = a\left[\left(x + \frac{b}{2a}y\right)^2 + \left(\frac{4ac - b^2}{4a^2}\right)y^2\right]$

(b) For $D = 4ac - b^2$, from part (a) we have $f(x,y) = a\left[\left(x + \frac{b}{2a}y\right)^2 + \left(\frac{D}{4a^2}\right)y^2\right]$. If $D > 0$, $\left(\frac{D}{4a^2}\right)y^2 \geq 0$ and $\left(x + \frac{b}{2a}y\right)^2 \geq 0$, so $\left[\left(x + \frac{b}{2a}y\right)^2 + \left(\frac{D}{4a^2}\right)y^2\right] \geq 0$. Here $a > 0$, thus $f(x,y) = a\left[\left(x + \frac{b}{2a}y\right)^2 + \left(\frac{D}{4a^2}\right)y^2\right] \geq 0$. We know $f(0,0) = 0$, so $f(0,0) \leq f(x,y)$ for all (x,y), and by definition f has a local minimum at $(0,0)$.

(c) As in part (b), $\left[\left(x + \frac{b}{2a}y\right)^2 + \left(\frac{D}{4a^2}\right)y^2\right] \geq 0$, and since $a < 0$ we have $f(x,y) = a\left[\left(x + \frac{b}{2a}y\right)^2 + \left(\frac{D}{4a^2}\right)y^2\right] \leq 0$. Since $f(0,0) = 0$, we must have $f(0,0) \geq f(x,y)$ for all (x,y), so by definition f has a local maximum at $(0,0)$.

(d) $f(x,y) = ax^2 + bxy + cy^2$, so $f_x(x,y) = 2ax + by \;\Rightarrow\; f_x(0,0) = 0$ and $f_y(x,y) = bx + 2cy \;\Rightarrow\; f_y(0,0) = 0$. Since $f(0,0) = 0$ and f and its partial derivatives are continuous, we know from Equation 14.4.2 that the tangent plane to the graph of f at $(0,0)$ is the plane $z = 0$. Then f has a saddle point at $(0,0)$ if the graph of f crosses the tangent plane at $(0,0)$, or equivalently, if some paths to the origin have positive function values while other paths have negative function values. Suppose we approach the origin along the x-axis; then we have $y = 0 \;\Rightarrow\; f(x,0) = ax^2$ which has the same sign as a. We must now find at least one path to the origin where $f(x,y)$ gives values with sign opposite that of a. Since $f(x,y) = a\left[\left(x + \frac{b}{2a}y\right)^2 + \left(\frac{D}{4a^2}\right)y^2\right]$, if we approach the origin along the line $x = -\frac{b}{2a}y$, we have $f\left(-\frac{b}{2a}y, y\right) = a\left[\left(-\frac{b}{2a}y + \frac{b}{2a}y\right)^2 + \left(\frac{D}{4a^2}\right)y^2\right] = \frac{D}{4a}y^2$. Since $D < 0$, these values have signs opposite that of a. Thus, f has a saddle point at $(0,0)$.

5. (a) Since the partial derivatives of f exist at $(0,0)$ and $(0,0)$ is a critical point, we know $f_x(0,0) = 0$ and $f_y(0,0) = 0$. Then the second-degree Taylor polynomial of f at $(0,0)$ can be expressed as

$$\begin{aligned} Q(x,y) &= f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) + \tfrac{1}{2}f_{xx}(0,0)(x-0)^2 \\ &\qquad + f_{xy}(0,0)(x-0)(y-0) + \tfrac{1}{2}f_{yy}(0,0)(y-0)^2 \\ &= \tfrac{1}{2}f_{xx}(0,0)x^2 + f_{xy}(0,0)xy + \tfrac{1}{2}f_{yy}(0,0)y^2 \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) $Q(x,y) = \frac{1}{2} f_{xx}(0,0)x^2 + f_{xy}(0,0)xy + \frac{1}{2} f_{yy}(0,0)y^2$ fits the form of the polynomial function in Problem 4 with $a = \frac{1}{2} f_{xx}(0,0)$, $b = f_{xy}(0,0)$, and $c = \frac{1}{2} f_{yy}(0,0)$. Then we know Q is a paraboloid, and that Q has a local maximum, local minimum, or saddle point at $(0,0)$. Here, $D = 4ac - b^2 = 4\left(\frac{1}{2}\right) f_{xx}(0,0)\left(\frac{1}{2}\right) f_{yy}(0,0) - [f_{xy}(0,0)]^2 = f_{xx}(0,0)f_{yy}(0,0) - [f_{xy}(0,0)]^2$, and if $D > 0$ with $a = \frac{1}{2} f_{xx}(0,0) > 0 \Rightarrow f_{xx}(0,0) > 0$, we know from Problem 4 that Q has a local minimum at $(0,0)$. Similarly, if $D > 0$ and $a < 0 \Rightarrow f_{xx}(0,0) < 0$, Q has a local maximum at $(0,0)$, and if $D < 0$, Q has a saddle point at $(0,0)$.

(c) Since $f(x,y) \approx Q(x,y)$ near $(0,0)$, part (b) suggests that for $D = f_{xx}(0,0)f_{yy}(0,0) - [f_{xy}(0,0)]^2$, if $D > 0$ and $f_{xx}(0,0) > 0$, f has a local minimum at $(0,0)$. If $D > 0$ and $f_{xx}(0,0) < 0$, f has a local maximum at $(0,0)$, and if $D < 0$, f has a saddle point at $(0,0)$. Together with the conditions given in part (a), this is precisely the Second Derivatives Test from Section 14.7.

14.8 Lagrange Multipliers

1. At the extreme values of f, the level curves of f just touch the curve $g(x,y) = 8$ with a common tangent line. (See Figure 1 and the accompanying discussion.) We can observe several such occurrences on the contour map, but the level curve $f(x,y) = c$ with the largest value of c which still intersects the curve $g(x,y) = 8$ is approximately $c = 59$, and the smallest value of c corresponding to a level curve which intersects $g(x,y) = 8$ appears to be $c = 30$. Thus we estimate the maximum value of f subject to the constraint $g(x,y) = 8$ to be about 59 and the minimum to be 30.

2. (a) The values $c = \pm 1$ and $c = 1.25$ seem to give curves which are tangent to the circle. These values represent possible extreme values of the function $x^2 + y$ subject to the constraint $x^2 + y^2 = 1$.



(b) $\nabla f = \langle 2x, 1\rangle$, $\lambda \nabla g = \langle 2\lambda x, 2\lambda y\rangle$. So $2x = 2\lambda x \Rightarrow$ either $\lambda = 1$ or $x = 0$. If $\lambda = 1$, then $y = \frac{1}{2}$ and so $x = \pm\frac{\sqrt{3}}{2}$ (from the constraint). If $x = 0$, then $y = \pm 1$. Therefore f has possible extreme values at the points $(0, \pm 1)$ and $\left(\pm\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$. We calculate $f\left(\pm\frac{\sqrt{3}}{2}, \frac{1}{2}\right) = \frac{5}{4}$ (the maximum value), $f(0,1) = 1$, and $f(0,-1) = -1$ (the minimum value). These are our answers from part (a).

3. We want to find the extreme values of $f(x,y) = x^2 - y^2$ subject to the constraint $g(x,y) = x^2 + y^2 = 1$. Then $\nabla f = \lambda \nabla g \Rightarrow \langle 2x, -2y\rangle = \lambda \langle 2x, 2y\rangle$, so we solve the equations $2x = 2\lambda x$, $-2y = 2\lambda y$, and $x^2 + y^2 = 1$. From the first equation we have $2x(\lambda - 1) = 0 \Rightarrow x = 0$ or $\lambda = 1$. If $x = 0$ then substitution into the constraint gives

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$y^2 = 1 \Rightarrow y = \pm 1$. If $\lambda = 1$ then substitution into the second equation gives $-2y = 2y \Rightarrow y = 0$, and from the constraint we must have $x = \pm 1$. Thus the possible points for the extreme values of f are $(0, \pm 1)$ and $(\pm 1, 0)$. Evaluating f at these points, we see that the maximum value of f is $f(\pm 1, 0) = 1$ and the minimum is $f(0, \pm 1) = -1$.

4. $f(x, y) = 3x + y$, $g(x, y) = x^2 + y^2 = 10$, and $\nabla f = \lambda \nabla g \Rightarrow \langle 3, 1 \rangle = \langle 2\lambda x, 2\lambda y \rangle$, so $3 = 2\lambda x$, $1 = 2\lambda y$, and $x^2 + y^2 = 10$. From the first two equations we have $\dfrac{3}{2x} = \lambda = \dfrac{1}{2y} \Rightarrow x = 3y$ (note that the first two equations imply $x \neq 0$ and $y \neq 0$) and substitution into the third equation gives $9y^2 + y^2 = 10 \Rightarrow y^2 = 1 \Rightarrow y = \pm 1$. Then f has possible extreme values at the points $(3, 1)$ and $(-3, -1)$. We compute $f(3, 1) = 10$ and $f(-3, -1) = -10$, so the maximum value of f on $x^2 + y^2 = 10$ is $f(3, 1) = 10$ and the minimum value is $f(-3, -1) = -10$.

5. $f(x, y) = xy$, $g(x, y) = 4x^2 + y^2 = 8$, and $\nabla f = \lambda \nabla g \Rightarrow \langle y, x \rangle = \langle 8\lambda x, 2\lambda y \rangle$, so $y = 8\lambda x$, $x = 2\lambda y$, and $4x^2 + y^2 = 8$. First note that if $x = 0$ then $y = 0$ by the first equation, and if $y = 0$ then $x = 0$ by the second equation. But this contradicts the third equation, so $x \neq 0$ and $y \neq 0$. Then from the first two equations we have $\dfrac{y}{8x} = \lambda = \dfrac{x}{2y} \Rightarrow 2y^2 = 8x^2 \Rightarrow y^2 = 4x^2$, and substitution into the third equation gives $4x^2 + 4x^2 = 8 \Rightarrow x = \pm 1$. If $x = \pm 1$ then $y^2 = 4 \Rightarrow y = \pm 2$, so f has possible extreme values at $(1, \pm 2)$ and $(-1, \pm 2)$. Evaluating f at these points, we see that the maximum value is $f(1, 2) = f(-1, -2) = 2$ and the minimum is $f(1, -2) = f(-1, 2) = -2$.

6. $f(x, y) = xe^y$, $g(x, y) = x^2 + y^2 = 2$, and $\nabla f = \lambda \nabla g \Rightarrow \langle e^y, xe^y \rangle = \langle 2\lambda x, 2\lambda y \rangle$, so $e^y = 2\lambda x$, $xe^y = 2\lambda y$, and $x^2 + y^2 = 2$. First note that from the first equation $x \neq 0$. If $y = 0$, the second equation implies $x = 0$, so $y \neq 0$. Then from the first two equations we have $\dfrac{e^y}{2x} = \lambda = \dfrac{xe^y}{2y} \Rightarrow 2ye^y = 2x^2e^y \Rightarrow y = x^2$, and substituting into the third equation gives $x^2 + (x^2)^2 = 2 \Rightarrow x^4 + x^2 - 2 = 0 \Rightarrow (x^2 + 2)(x^2 - 1) = 0 \Rightarrow x = \pm 1$. From $y = x^2$ we have $y = 1$, so f has possible extreme values at $(\pm 1, 1)$. Evaluating f at these points, we see that the maximum value is $f(1, 1) = e$ and the minimum is $f(-1, 1) = -e$.

7. $f(x, y, z) = 2x + 2y + z$, $g(x, y, z) = x^2 + y^2 + z^2 = 9$, and $\nabla f = \lambda \nabla g \Rightarrow \langle 2, 2, 1 \rangle = \langle 2\lambda x, 2\lambda y, 2\lambda z \rangle$, so $2\lambda x = 2$, $2\lambda y = 2$, $2\lambda z = 1$, and $x^2 + y^2 + z^2 = 9$. The first three equations imply $x = \dfrac{1}{\lambda}$, $y = \dfrac{1}{\lambda}$, and $z = \dfrac{1}{2\lambda}$. But substitution into the fourth equation gives $\left(\dfrac{1}{\lambda}\right)^2 + \left(\dfrac{1}{\lambda}\right)^2 + \left(\dfrac{1}{2\lambda}\right)^2 = 9 \Rightarrow \dfrac{9}{4\lambda^2} = 9 \Rightarrow \lambda = \pm\frac{1}{2}$, so f has possible extreme values at the points $(2, 2, 1)$ and $(-2, -2, -1)$. The maximum value of f on $x^2 + y^2 + z^2 = 9$ is $f(2, 2, 1) = 9$, and the minimum is $f(-2, -2, -1) = -9$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

8. $f(x,y,z) = e^{xyz}$, $g(x,y,z) = 2x^2 + y^2 + z^2 = 24$, and $\nabla f = \lambda \nabla g \;\Rightarrow\; \langle yze^{xyz}, xze^{xyz}, xye^{xyz}\rangle = \langle 4\lambda x, 2\lambda y, 2\lambda z\rangle$. Then $yze^{xyz} = 4\lambda x$, $xze^{xyz} = 2\lambda y$, $xye^{xyz} = 2\lambda z$, and $2x^2 + y^2 + z^2 = 24$. If any of x, y, z, or λ is zero, then the first three equations imply that two of the variables x, y, z must be zero. If $x = y = z = 0$ it contradicts the fourth equation, so exactly two are zero, and from the fourth equation the possibilities are $\left(\pm 2\sqrt{3}, 0, 0\right)$, $\left(0, \pm 2\sqrt{6}, 0\right)$, $\left(0, 0, \pm 2\sqrt{6}\right)$, all with an f-value of $e^0 = 1$. If none of x, y, z, λ is zero then from the first three equations we have $\dfrac{4\lambda x}{yz} = e^{xyz} = \dfrac{2\lambda y}{xz} = \dfrac{2\lambda z}{xy} \;\Rightarrow\; \dfrac{2x}{yz} = \dfrac{y}{xz} = \dfrac{z}{xy}$. This gives $2x^2 z = y^2 z \;\Rightarrow\; 2x^2 = y^2$ and $xy^2 = xz^2 \;\Rightarrow\; y^2 = z^2$. Substituting into the fourth equation, we have $y^2 + y^2 + y^2 = 24 \;\Rightarrow\; y^2 = 8 \;\Rightarrow\; y = \pm 2\sqrt{2}$, so $x^2 = 4 \;\Rightarrow\; x = \pm 2$ and $z^2 = y^2 \;\Rightarrow\; z = \pm 2\sqrt{2}$, giving possible points $\left(\pm 2, \pm 2\sqrt{2}, \pm 2\sqrt{2}\right)$ (all combinations). The value of f is e^{16} when all coordinates are positive or exactly two are negative, and the value is e^{-16} when all are negative or exactly one of the coordinates is negative. Thus the maximum of f subject to the constraint is e^{16} and the minimum is e^{-16}.

9. $f(x,y,z) = xy^2 z$, $g(x,y,z) = x^2 + y^2 + z^2 = 4$, and $\nabla f = \lambda \nabla g \;\Rightarrow\; \langle y^2 z, 2xyz, xy^2\rangle = \lambda \langle 2x, 2y, 2z\rangle$. Then $y^2 z = 2\lambda x$, $2xyz = 2\lambda y$, $xy^2 = 2\lambda z$, and $x^2 + y^2 + z^2 = 4$.

Case 1: If $\lambda = 0$, then the first equation implies that $y = 0$ or $z = 0$. If $y = 0$, then any values of x and z satisfy the first three equations, so from the fourth equation all points $(x, 0, z)$ such that $x^2 + z^2 = 4$ are possible points. If $z = 0$ then from the third equation $x = 0$ or $y = 0$, and from the fourth equation, the possible points are $(0, \pm 2, 0)$, $(\pm 2, 0, 0)$. The f-value in all these cases is 0.

Case 2: If $\lambda \neq 0$ but any one of x, y, z is zero, the first three equations imply that all three coordinates must be zero, contradicting the fourth equation. Thus if $\lambda \neq 0$, none of x, y, z is zero and from the first three equations we have $\dfrac{y^2 z}{2x} = \lambda = xz = \dfrac{xy^2}{2z}$. This gives $y^2 z = 2x^2 z \;\Rightarrow\; y^2 = 2x^2$ and $2y^2 z^2 = 2x^2 y^2 \;\Rightarrow\; z^2 = x^2$. Substituting into the fourth equation, we have $x^2 + 2x^2 + x^2 = 4 \;\Rightarrow\; x^2 = 1 \;\Rightarrow\; x = \pm 1$, so $y = \pm\sqrt{2}$ and $z = \pm 1$, giving possible points $\left(\pm 1, \pm\sqrt{2}, \pm 1\right)$ (all combinations). The value of f is 2 when x and z are the same sign and -2 when they are opposite. Thus the maximum of f subject to the constraint is $f(1, \pm\sqrt{2}, 1) = f(-1, \pm\sqrt{2}, -1) = 2$ and the minimum is $f(1, \pm\sqrt{2}, -1) = f(-1, \pm\sqrt{2}, 1) = -2$.

10. $f(x,y,z) = \ln(x^2+1) + \ln(y^2+1) + \ln(z^2+1)$, $g(x,y,z) = x^2 + y^2 + z^2 = 12$. Then $\nabla f = \lambda \nabla g \;\Rightarrow$ $\left\langle \dfrac{2x}{x^2+1}, \dfrac{2y}{y^2+1}, \dfrac{2z}{z^2+1} \right\rangle = \lambda \langle 2x, 2y, 2z\rangle$, so $\dfrac{2x}{x^2+1} = 2\lambda x$, $\dfrac{2y}{y^2+1} = 2\lambda y$, $\dfrac{2z}{z^2+1} = 2\lambda z$, and $x^2 + y^2 + z^2 = 12$. First, if $\lambda = 0$ then $x = y = z = 0$ which contradicts the last equation, so we may assume that $\lambda \neq 0$.

Case 1: If $x \neq 0$, $y \neq 0$, and $z \neq 0$, then from the first three equations we have $\dfrac{1}{x^2+1} = \lambda = \dfrac{1}{y^2+1} = \dfrac{1}{z^2+1} \;\Rightarrow$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x^2 = y^2 = z^2$, and substitution into the last equation gives $3x^2 = 12 \Rightarrow x = \pm 2$. Thus possible points are $(\pm 2, \pm 2, \pm 2)$ (all combinations), all of which have an f-value of $3 \ln 5$.

Case 2: If exactly one of x, y, z is zero, say $x = 0$, then from the second and third equations we have $y^2 = z^2$. Substitution into the last equation gives $2y^2 = 12 \Rightarrow y = \pm\sqrt{6}$. The situation is similar for $y = 0$ or $z = 0$, giving possible points $\left(0, \pm\sqrt{6}, \pm\sqrt{6}\right)$, $\left(\pm\sqrt{6}, 0, \pm\sqrt{6}\right)$, $\left(\pm\sqrt{6}, \pm\sqrt{6}, 0\right)$ (all combinations), all with an f-value of $2 \ln 7$.

Case 3: If exactly two of x, y, z are zero, then the square of the nonzero variable is 12, giving possible points $\left(0, 0, \pm 2\sqrt{3}\right)$, $\left(0, \pm 2\sqrt{3}, 0\right)$, $\left(\pm 2\sqrt{3}, 0, 0\right)$, all with an f-value of $\ln 13$.

Thus the maximum of f subject to the constraint is $3 \ln 5 \approx 4.83$ and the minimum is $\ln 13 \approx 2.56$.

11. $f(x, y, z) = x^2 + y^2 + z^2$, $g(x, y, z) = x^4 + y^4 + z^4 = 1 \Rightarrow \nabla f = \langle 2x, 2y, 2z \rangle$, $\lambda \nabla g = \langle 4\lambda x^3, 4\lambda y^3, 4\lambda z^3 \rangle$.

Case 1: If $x \neq 0$, $y \neq 0$, and $z \neq 0$, then $\nabla f = \lambda \nabla g$ implies $\lambda = 1/(2x^2) = 1/(2y^2) = 1/(2z^2)$ or $x^2 = y^2 = z^2$ and $3x^4 = 1$ or $x = \pm \frac{1}{\sqrt[4]{3}}$ giving the points $\left(\pm \frac{1}{\sqrt[4]{3}}, \frac{1}{\sqrt[4]{3}}, \frac{1}{\sqrt[4]{3}}\right)$, $\left(\pm \frac{1}{\sqrt[4]{3}}, -\frac{1}{\sqrt[4]{3}}, \frac{1}{\sqrt[4]{3}}\right)$, $\left(\pm \frac{1}{\sqrt[4]{3}}, \frac{1}{\sqrt[4]{3}}, -\frac{1}{\sqrt[4]{3}}\right)$, $\left(\pm \frac{1}{\sqrt[4]{3}}, -\frac{1}{\sqrt[4]{3}}, -\frac{1}{\sqrt[4]{3}}\right)$ all with an f-value of $\sqrt{3}$.

Case 2: If one of the variables equals zero and the other two are not zero, then the squares of the two nonzero coordinates are equal with common value $\frac{1}{\sqrt{2}}$ and corresponding f-value of $\sqrt{2}$.

Case 3: If exactly two of the variables are zero, then the third variable has value ± 1 with the corresponding f-value of 1.

Thus on $x^4 + y^4 + z^4 = 1$, the maximum value of f is $\sqrt{3}$ and the minimum value is 1.

12. $f(x, y, z) = x^4 + y^4 + z^4$, $g(x, y, z) = x^2 + y^2 + z^2 = 1 \Rightarrow \nabla f = \langle 4x^3, 4y^3, 4z^3 \rangle$, $\lambda \nabla g = \langle 2\lambda x, 2\lambda y, 2\lambda z \rangle$.

Case 1: If $x \neq 0$, $y \neq 0$, and $z \neq 0$ then $\nabla f = \lambda \nabla g$ implies $\lambda = 2x^2 = 2y^2 = 2z^2$ or $x^2 = y^2 = z^2 = \frac{1}{3}$ giving 8 points each with an f-value of $\frac{1}{3}$.

Case 2: If one of the variables is 0 and the other two are not, then the squares of the two nonzero coordinates are equal with common value $\frac{1}{2}$ and the corresponding f-value is $\frac{1}{2}$.

Case 3: If exactly two of the variables are 0, then the third variable has value ± 1 with corresponding f-value of 1.

Thus on $x^2 + y^2 + z^2 = 1$, the maximum value of f is 1 and the minimum value is $\frac{1}{3}$.

13. $f(x, y, z, t) = x + y + z + t$, $g(x, y, z, t) = x^2 + y^2 + z^2 + t^2 = 1 \Rightarrow \langle 1, 1, 1, 1 \rangle = \langle 2\lambda x, 2\lambda y, 2\lambda z, 2\lambda t \rangle$, so $\lambda = 1/(2x) = 1/(2y) = 1/(2z) = 1/(2t)$ and $x = y = z = t$. But $x^2 + y^2 + z^2 + t^2 = 1$, so the possible points are $\left(\pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}\right)$. Thus the maximum value of f is $f\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) = 2$ and the minimum value is $f\left(-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right) = -2$.

14. $f(x_1, x_2, \ldots, x_n) = x_1 + x_2 + \cdots + x_n$, $g(x_1, x_2, \ldots, x_n) = x_1^2 + x_2^2 + \cdots + x_n^2 = 1 \Rightarrow$ $\langle 1, 1, \ldots, 1 \rangle = \langle 2\lambda x_1, 2\lambda x_2, \ldots, 2\lambda x_n \rangle$, so $\lambda = 1/(2x_1) = 1/(2x_2) = \cdots = 1/(2x_n)$ and $x_1 = x_2 = \cdots = x_n$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

But $x_1^2 + x_2^2 + \cdots + x_n^2 = 1$, so $x_i = \pm 1/\sqrt{n}$ for $i = 1, \ldots, n$. Thus the maximum value of f is $f(1/\sqrt{n}, 1/\sqrt{n}, \ldots, 1/\sqrt{n}) = \sqrt{n}$ and the minimum value is $f(-1/\sqrt{n}, -1/\sqrt{n}, \ldots, -1/\sqrt{n}) = -\sqrt{n}$.

15. $f(x, y) = x^2 + y^2$, $g(x, y) = xy = 1$, and $\nabla f = \lambda \nabla g \;\Rightarrow\; \langle 2x, 2y\rangle = \langle \lambda y, \lambda x\rangle$, so $2x = \lambda y$, $2y = \lambda x$, and $xy = 1$. From the last equation, $x \neq 0$ and $y \neq 0$, so $2x = \lambda y \;\Rightarrow\; \lambda = 2x/y$. Substituting, we have $2y = (2x/y)\,x \;\Rightarrow\; y^2 = x^2 \;\Rightarrow\; y = \pm x$. But $xy = 1$, so $x = y = \pm 1$ and the possible points for the extreme values of f are $(1, 1)$ and $(-1, -1)$. Here there is no maximum value, since the constraint $xy = 1 \;\Leftrightarrow\; y = 1/x$ allows x or y to become arbitrarily large, and hence $f(x, y) = x^2 + y^2$ can be made arbitrarily large. The minimum value is $f(1, 1) = f(-1, -1) = 2$.

16. $f(x, y, z) = x^2 + 2y^2 + 3z^2$, $g(x, y) = x + 2y + 3z = 10$, and $\nabla f = \lambda \nabla g \;\Rightarrow\; \langle 2x, 4y, 6z\rangle = \langle \lambda, 2\lambda, 3\lambda\rangle$, so $2x = \lambda$, $4y = 2\lambda$, $6z = 3\lambda$, and $x + 2y + 3z = 10$. From the first three equations we have $2x = \lambda = 2y = 2z \;\Rightarrow\; x = y = z$, and substituting into the fourth equation gives $x + 2x + 3x = 10 \;\Rightarrow\; x = \frac{5}{3} = y = z$. Thus the only possible point for an extreme value of f is $\left(\frac{5}{3}, \frac{5}{3}, \frac{5}{3}\right)$. Notice here that the constraint $x + 2y + 3z = 10$ allows any of $|x|$, $|y|$, or $|z|$ to be arbitrarily large, and hence $f(x, y, z) = x^2 + 2y^2 + 3z^2$ can be made arbitrarily large. So f has no maximum value subject to the constraint. The minimum value is $f\left(\frac{5}{3}, \frac{5}{3}, \frac{5}{3}\right) = 6\left(\frac{5}{3}\right)^2 = \frac{50}{3}$.

17. $f(x, y, z) = x + y + z$, $g(x, y, z) = x^2 + z^2 = 2$, $h(x, y, z) = x + y = 1$, and $\nabla f = \lambda \nabla g + \mu \nabla h \;\Rightarrow\; \langle 1, 1, 1\rangle = \langle 2\lambda x, 0, 2\lambda z\rangle + \langle \mu, \mu, 0\rangle$. Then $1 = 2\lambda x + \mu$, $1 = \mu$, $1 = 2\lambda z$, $x^2 + z^2 = 2$, and $x + y = 1$. Substituting $\mu = 1$ into the first equation gives $\lambda = 0$ or $x = 0$. But $\lambda = 0$ contradicts $1 = 2\lambda z$, so $x = 0$. Then $x + y = 1 \;\Rightarrow\; y = 1$ and $x^2 + z^2 = 2 \;\Rightarrow\; z = \pm\sqrt{2}$, so the possible points are $\left(0, 1, \pm\sqrt{2}\right)$. The maximum value of f subject to the constraints is $f(0, 1, \sqrt{2}) = 1 + \sqrt{2} \approx 2.41$ and the minimum is $f(0, 1, -\sqrt{2}) = 1 - \sqrt{2} \approx -0.41$.

Note: Since $x + y = 1$ is one of the constraints, we could have solved the problem by solving $f(x, z) = 1 + z$ subject to $x^2 + z^2 = 2$.

18. $f(x, y, z) = z$, $g(x, y, z) = x^2 + y^2 - z^2 = 0$, $h(x, y, z) = x + y + z = 24$, and $\nabla f = \lambda \nabla g + \mu \nabla h \;\Rightarrow\; \langle 0, 0, 1\rangle = \langle 2\lambda x, 2\lambda y, -2\lambda z\rangle + \langle \mu, \mu, \mu\rangle$. Then $0 = 2\lambda x + \mu$, $0 = 2\lambda y + \mu$, $1 = -2\lambda z + \mu$, $x^2 + y^2 - z^2 = 0$, and $x + y + z = 24$. From the first two equations we have $-2\lambda x = \mu = -2\lambda y \;\Rightarrow\; \lambda = 0$ or $x = y$. But $\lambda = 0 \;\Rightarrow\; \mu = 0$ which contradicts the third equation, so $x = y$ and substitution into the last equation gives $z = 24 - 2x$. From the fourth equation we have $x^2 + x^2 - (24 - 2x)^2 = 0 \;\Rightarrow\; -2x^2 + 96x - 576 = 0 \;\Rightarrow\; x^2 - 48x + 288 = 0 \;\Rightarrow\; x = \dfrac{48 \pm \sqrt{1152}}{2} = 24 \pm 12\sqrt{2} = y$. Now $z = 24 - 2x$, so the possible points are $\left(24 + 12\sqrt{2}, 24 + 12\sqrt{2}, -24 - 24\sqrt{2}\right)$ and $\left(24 - 12\sqrt{2}, 24 - 12\sqrt{2}, -24 + 24\sqrt{2}\right)$. The maximum of f subject to the constraints is $f\left(24 - 12\sqrt{2}, 24 - 12\sqrt{2}, -24 + 24\sqrt{2}\right) = -24 + 24\sqrt{2} \approx 9.94$ and the minimum is $f\left(24 + 12\sqrt{2}, 24 + 12\sqrt{2}, -24 - 24\sqrt{2}\right) = -24 - 24\sqrt{2} \approx -57.94$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $f(x,y,z) = yz + xy,\ g(x,y,z) = xy = 1,\ h(x,y,z) = y^2 + z^2 = 1 \ \Rightarrow\ \nabla f = \langle y, x+z, y\rangle, \lambda\nabla g = \langle \lambda y, \lambda x, 0\rangle,$ $\mu\nabla h = \langle 0, 2\mu y, 2\mu z\rangle$. Then $y = \lambda y$ implies $\lambda = 1$ [$y \neq 0$ since $g(x,y,z) = 1$], $x + z = \lambda x + 2\mu y$ and $y = 2\mu z$. Thus $\mu = z/(2y) = y/(2y)$ or $y^2 = z^2$, and so $y^2 + z^2 = 1$ implies $y = \pm\frac{1}{\sqrt{2}}$, $z = \pm\frac{1}{\sqrt{2}}$. Then $xy = 1$ implies $x = \pm\sqrt{2}$ and the possible points are $\left(\pm\sqrt{2}, \pm\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$, $\left(\pm\sqrt{2}, \pm\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$. Hence the maximum of f subject to the constraints is $f\left(\pm\sqrt{2}, \pm\frac{1}{\sqrt{2}}, \pm\frac{1}{\sqrt{2}}\right) = \frac{3}{2}$ and the minimum is $f\left(\pm\sqrt{2}, \pm\frac{1}{\sqrt{2}}, \mp\frac{1}{\sqrt{2}}\right) = \frac{1}{2}$.

Note: Since $xy = 1$ is one of the constraints we could have solved the problem by solving $f(y,z) = yz + 1$ subject to $y^2 + z^2 = 1$.

20. $f(x,y,z) = x^2 + y^2 + z^2,\ g(x,y,z) = x - y = 1,\ h(x,y,z) = y^2 - z^2 = 1 \ \Rightarrow\ \nabla f = \langle 2x, 2y, 2z\rangle,$ $\lambda\nabla g = \langle \lambda, -\lambda, 0\rangle$, and $\mu\nabla h = \langle 0, 2\mu y, -2\mu z\rangle$. Then $2x = \lambda$, $2y = -\lambda + 2\mu y$, and $2z = -2\mu z \ \Rightarrow\ z = 0$ or $\mu = -1$. If $z = 0$ then $y^2 - z^2 = 1$ implies $y^2 = 1 \ \Rightarrow\ y = \pm 1$. If $y = 1$, $x - y = 1$ implies $x = 2$, and if $y = -1$ we have $x = 0$, so possible points are $(2,1,0)$ and $(0,-1,0)$. If $\mu = -1$ then $2y = -\lambda + 2\mu y$ implies $4y = -\lambda$, but $\lambda = 2x$ so $4y = -2x \ \Rightarrow\ x = -2y$ and $x - y = 1$ implies $-3y = 1 \ \Rightarrow\ y = -\frac{1}{3}$. But then $y^2 - z^2 = 1$ implies $z^2 = -\frac{8}{9}$, an impossibility. Thus the maximum value of f subject to the constraints is $f(2,1,0) = 5$ and the minimum is $f(0,-1,0) = 1$.

Note: Since $x - y = 1 \ \Rightarrow\ x = y + 1$ is one of the constraints we could have solved the problem by solving $f(y,z) = (y+1)^2 + y^2 + z^2$ subject to $y^2 - z^2 = 1$.

21. $f(x,y) = x^2 + y^2 + 4x - 4y$. For the interior of the region, we find the critical points: $f_x = 2x + 4$, $f_y = 2y - 4$, so the only critical point is $(-2,2)$ (which is inside the region) and $f(-2,2) = -8$. For the boundary, we use Lagrange multipliers. $g(x,y) = x^2 + y^2 = 9$, so $\nabla f = \lambda\nabla g \ \Rightarrow\ \langle 2x+4, 2y-4\rangle = \langle 2\lambda x, 2\lambda y\rangle$. Thus $2x + 4 = 2\lambda x$ and $2y - 4 = 2\lambda y$. Adding the two equations gives $2x + 2y = 2\lambda x + 2\lambda y \ \Rightarrow\ x + y = \lambda(x + y) \ \Rightarrow\ (x+y)(\lambda - 1) = 0$, so $x + y = 0 \ \Rightarrow\ y = -x$ or $\lambda - 1 = 0 \ \Rightarrow\ \lambda = 1$. But $\lambda = 1$ leads to a contradition in $2x + 4 = 2\lambda x$, so $y = -x$ and $x^2 + y^2 = 9$ implies $2y^2 = 9 \ \Rightarrow\ y = \pm\frac{3}{\sqrt{2}}$. We have $f\left(\frac{3}{\sqrt{2}}, -\frac{3}{\sqrt{2}}\right) = 9 + 12\sqrt{2} \approx 25.97$ and $f\left(-\frac{3}{\sqrt{2}}, \frac{3}{\sqrt{2}}\right) = 9 - 12\sqrt{2} \approx -7.97$, so the maximum value of f on the disk $x^2 + y^2 \leq 9$ is $f\left(\frac{3}{\sqrt{2}}, -\frac{3}{\sqrt{2}}\right) = 9 + 12\sqrt{2}$ and the minimum is $f(-2,2) = -8$.

22. $f(x,y) = 2x^2 + 3y^2 - 4x - 5 \ \Rightarrow\ \nabla f = \langle 4x - 4, 6y\rangle = \langle 0, 0\rangle \ \Rightarrow\ x = 1, y = 0$. Thus $(1,0)$ is the only critical point of f, and it lies in the region $x^2 + y^2 < 16$. On the boundary, $g(x,y) = x^2 + y^2 = 16 \ \Rightarrow\ \lambda\nabla g = \langle 2\lambda x, 2\lambda y\rangle$, so $6y = 2\lambda y \ \Rightarrow$ either $y = 0$ or $\lambda = 3$. If $y = 0$, then $x = \pm 4$; if $\lambda = 3$, then $4x - 4 = 2\lambda x \ \Rightarrow\ x = -2$ and $y = \pm 2\sqrt{3}$. Now $f(1,0) = -7$, $f(4,0) = 11$, $f(-4,0) = 43$, and $f\left(-2, \pm 2\sqrt{3}\right) = 47$. Thus the maximum value of $f(x,y)$ on the disk $x^2 + y^2 \leq 16$ is $f\left(-2, \pm 2\sqrt{3}\right) = 47$, and the minimum value is $f(1,0) = -7$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

23. $f(x,y) = e^{-xy}$. For the interior of the region, we find the critical points: $f_x = -ye^{-xy}$, $f_y = -xe^{-xy}$, so the only critical point is $(0,0)$, and $f(0,0) = 1$. For the boundary, we use Lagrange multipliers. $g(x,y) = x^2 + 4y^2 = 1 \Rightarrow \lambda\nabla g = \langle 2\lambda x, 8\lambda y\rangle$, so setting $\nabla f = \lambda\nabla g$ we get $-ye^{-xy} = 2\lambda x$ and $-xe^{-xy} = 8\lambda y$. The first of these gives $e^{-xy} = -2\lambda x/y$, and then the second gives $-x(-2\lambda x/y) = 8\lambda y \Rightarrow x^2 = 4y^2$. Solving this last equation with the constraint $x^2 + 4y^2 = 1$ gives $x = \pm\frac{1}{\sqrt{2}}$ and $y = \pm\frac{1}{2\sqrt{2}}$. Now $f\left(\pm\frac{1}{\sqrt{2}}, \mp\frac{1}{2\sqrt{2}}\right) = e^{1/4} \approx 1.284$ and $f\left(\pm\frac{1}{\sqrt{2}}, \pm\frac{1}{2\sqrt{2}}\right) = e^{-1/4} \approx 0.779$. The former are the maxima on the region and the latter are the minima.

24. (a) $f(x,y) = 2x + 3y$, $g(x,y) = \sqrt{x} + \sqrt{y} = 5 \Rightarrow \nabla f = \langle 2, 3\rangle = \lambda\nabla g = \lambda\left\langle \dfrac{1}{2\sqrt{x}}, \dfrac{1}{2\sqrt{y}}\right\rangle$. Then

$2 = \dfrac{\lambda}{2\sqrt{x}}$ and $3 = \dfrac{\lambda}{2\sqrt{y}}$ so $4\sqrt{x} = \lambda = 6\sqrt{y} \Rightarrow \sqrt{y} = \frac{2}{3}\sqrt{x}$. With $\sqrt{x} + \sqrt{y} = 5$ we have $\sqrt{x} + \frac{2}{3}\sqrt{x} = 5 \Rightarrow \sqrt{x} = 3 \Rightarrow x = 9$. Substituting into $\sqrt{y} = \frac{2}{3}\sqrt{x}$ gives $\sqrt{y} = 2$ or $y = 4$. Thus the only possible extreme value subject to the constraint is $f(9,4) = 30$. (The question remains whether this is indeed the maximum of f.)

(b) $f(25,0) = 50$ which is larger than the result of part (a).

(c)



We can see from the level curves of f that the maximum occurs at the left endpoint $(0,25)$ of the constraint curve g. The maximum value is $f(0,25) = 75$.

(d) Here ∇g does not exist if $x = 0$ or $y = 0$, so the method will not locate any associated points. Also, the method of Lagrange multipliers identifies points where the level curves of f share a common tangent line with the constraint curve g. This normally does not occur at an endpoint, although an absolute maximum or minimum may occur there.

(e) Here $f(9,4)$ is the absolute *minimum* of f subject to g.

25. (a) $f(x,y) = x$, $g(x,y) = y^2 + x^4 - x^3 = 0 \Rightarrow \nabla f = \langle 1, 0\rangle = \lambda\nabla g = \lambda\langle 4x^3 - 3x^2, 2y\rangle$. Then

$1 = \lambda(4x^3 - 3x^2)$ **(1)** and $0 = 2\lambda y$ **(2)**. We have $\lambda \neq 0$ from **(1)**, so **(2)** gives $y = 0$. Then, from the constraint equation, $x^4 - x^3 = 0 \Rightarrow x^3(x-1) = 0 \Rightarrow x = 0$ or $x = 1$. But $x = 0$ contradicts **(1)**, so the only possible extreme value subject to the constraint is $f(1,0) = 1$. (The question remains whether this is indeed the minimum of f.)

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The constraint is $y^2 + x^4 - x^3 = 0 \quad \Leftrightarrow \quad y^2 = x^3 - x^4$. The left side is non-negative, so we must have $x^3 - x^4 \geq 0$ which is true only for $0 \leq x \leq 1$. Therefore the minimum possible value for $f(x, y) = x$ is 0 which occurs for $x = y = 0$. However, $\lambda \nabla g(0, 0) = \lambda \langle 0 - 0, 0 \rangle = \langle 0, 0 \rangle$ and $\nabla f(0, 0) = \langle 1, 0 \rangle$, so $\nabla f(0, 0) \neq \lambda \nabla g(0, 0)$ for all values of λ.

(c) Here $\nabla g(0, 0) = \mathbf{0}$ but the method of Lagrange multipliers requires that $\nabla g \neq \mathbf{0}$ everywhere on the constraint curve.

26. (a) The graphs of $f(x, y) = 3.7$ and $f(x, y) = 350$ seem to be tangent to the circle, and so 3.7 and 350 are the approximate minimum and maximum values of the function $f(x, y)$ subject to the constraint $(x - 3)^2 + (y - 3)^2 = 9$.



(b) Let $g(x, y) = (x - 3)^2 + (y - 3)^2$. We calculate $f_x(x, y) = 3x^2 + 3y$, $f_y(x, y) = 3y^2 + 3x$, $g_x(x, y) = 2x - 6$, and $g_y(x, y) = 2y - 6$, and use a CAS to search for solutions to the equations $g(x, y) = (x - 3)^2 + (y - 3)^2 = 9$, $f_x = \lambda g_x$, and $f_y = \lambda g_y$. The solutions are $(x, y) = \left(3 - \frac{3}{2}\sqrt{2}, 3 - \frac{3}{2}\sqrt{2}\right) \approx (0.879, 0.879)$ and $(x, y) = \left(3 + \frac{3}{2}\sqrt{2}, 3 + \frac{3}{2}\sqrt{2}\right) \approx (5.121, 5.121)$. These give $f\left(3 - \frac{3}{2}\sqrt{2}, 3 - \frac{3}{2}\sqrt{2}\right) = \frac{351}{2} - \frac{243}{2}\sqrt{2} \approx 3.673$ and $f\left(3 + \frac{3}{2}\sqrt{2}, 3 + \frac{3}{2}\sqrt{2}\right) = \frac{351}{2} + \frac{243}{2}\sqrt{2} \approx 347.33$, in accordance with part (a).

27. $P(L, K) = bL^{\alpha}K^{1-\alpha}$, $g(L, K) = mL + nK = p \quad \Rightarrow \quad \nabla P = \langle \alpha b L^{\alpha-1}K^{1-\alpha}, (1-\alpha)bL^{\alpha}K^{-\alpha} \rangle$, $\lambda \nabla g = \langle \lambda m, \lambda n \rangle$. Then $\alpha b(K/L)^{1-\alpha} = \lambda m$ and $(1 - \alpha)b(L/K)^{\alpha} = \lambda n$ and $mL + nK = p$, so $\alpha b(K/L)^{1-\alpha}/m = (1 - \alpha)b(L/K)^{\alpha}/n$ or $n\alpha/[m(1 - \alpha)] = (L/K)^{\alpha}(L/K)^{1-\alpha}$ or $L = Kn\alpha/[m(1 - \alpha)]$. Substituting into $mL + nK = p$ gives $K = (1 - \alpha)p/n$ and $L = \alpha p/m$ for the maximum production.

28. $C(L, K) = mL + nK$, $g(L, K) = bL^{\alpha}K^{1-\alpha} = Q \quad \Rightarrow \quad \nabla C = \langle m, n \rangle$, $\lambda \nabla g = \langle \lambda \alpha b L^{\alpha-1}K^{1-\alpha}, \lambda(1-\alpha)bL^{\alpha}K^{-\alpha} \rangle$.

Then $\frac{m}{\alpha b}\left(\frac{L}{K}\right)^{1-\alpha} = \frac{n}{(1-\alpha)b}\left(\frac{K}{L}\right)^{\alpha}$ and $bL^{\alpha}K^{1-\alpha} = Q \quad \Rightarrow \quad \frac{n\alpha}{m(1-\alpha)} = \left(\frac{L}{K}\right)^{1-\alpha}\left(\frac{L}{K}\right)^{\alpha} \quad \rightarrow$

$L = \dfrac{Kn\alpha}{m(1-\alpha)}$ and so $b\left[\dfrac{Kn\alpha}{m(1-\alpha)}\right]^{\alpha} K^{1-\alpha} = Q$. Hence $K = \dfrac{Q}{b\left(n\alpha/[m(1-\alpha)]\right)^{\alpha}} = \dfrac{Qm^{\alpha}(1-\alpha)^{\alpha}}{bn^{\alpha}\alpha^{\alpha}}$

and $L = \dfrac{Qm^{\alpha-1}(1-\alpha)^{\alpha-1}}{bn^{\alpha-1}\alpha^{\alpha-1}} = \dfrac{Qn^{1-\alpha}\alpha^{1-\alpha}}{bm^{1-\alpha}(1-\alpha)^{1-\alpha}}$ minimizes cost.

29. Let the sides of the rectangle be x and y. Then $f(x, y) = xy$, $g(x, y) = 2x + 2y = p \quad \Rightarrow \quad \nabla f(x, y) = \langle y, x \rangle$, $\lambda \nabla g = \langle 2\lambda, 2\lambda \rangle$. Then $\lambda = \frac{1}{2}y = \frac{1}{2}x$ implies $x = y$ and the rectangle with maximum area is a square with side length $\frac{1}{4}p$.

30. Let $f(x, y, z) = s(s - x)(s - y)(s - z)$, $g(x, y, z) = x + y + z$. Then $\nabla f = \langle -s(s-y)(s-z), -s(s-x)(s-z), -s(s-x)(s-y) \rangle$, $\lambda \nabla g = \langle \lambda, \lambda, \lambda \rangle$. Thus $(s - y)(s - z) = (s - x)(s - z)$ **(1)**, and $(s - x)(s - z) = (s - x)(s - y)$ **(2)**. **(1)** implies $x = y$ while **(2)** implies $y = z$, so $x = y = z = p/3$ and the triangle with maximum area is equilateral.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

31. The distance from $(2, 0, -3)$ to a point (x, y, z) on the plane is $d = \sqrt{(x-2)^2 + y^2 + (z+3)^2}$, so we seek to minimize $d^2 = f(x, y, z) = (x-2)^2 + y^2 + (z+3)^2$ subject to the constraint that (x, y, z) lies on the plane $x + y + z = 1$, that is, that $g(x, y, z) = x + y + z = 1$. Then $\nabla f = \lambda \nabla g \Rightarrow \langle 2(x-2), 2y, 2(z+3)\rangle = \langle \lambda, \lambda, \lambda\rangle$, so $x = (\lambda + 4)/2$, $y = \lambda/2$, $z = (\lambda - 6)/2$. Substituting into the constraint equation gives $\dfrac{\lambda+4}{2} + \dfrac{\lambda}{2} + \dfrac{\lambda-6}{2} = 1 \Rightarrow 3\lambda - 2 = 2 \Rightarrow \lambda = \frac{4}{3}$, so $x = \frac{8}{3}$, $y = \frac{2}{3}$, and $z = -\frac{7}{3}$. This must correspond to a minimum, so the shortest distance is $d = \sqrt{\left(\frac{8}{3} - 2\right)^2 + \left(\frac{2}{3}\right)^2 + \left(-\frac{7}{3} + 3\right)^2} = \sqrt{\frac{4}{3}} = \frac{2}{\sqrt{3}}$.

32. The distance from $(0, 1, 1)$ to a point (x, y, z) on the plane is $d = \sqrt{x^2 + (y-1)^2 + (z-1)^2}$, so we minimize $d^2 = f(x, y, z) = x^2 + (y-1)^2 + (z-1)^2$ subject to the constraint that (x, y, z) lies on the plane $x - 2y + 3z = 6$, that is, $g(x, y, z) = x - 2y + 3z = 6$. Then $\nabla f = \lambda \nabla g \Rightarrow \langle 2x, 2(y-1), 2(z-1)\rangle = \langle \lambda, -2\lambda, 3\lambda\rangle$, so $x = \lambda/2$, $y = 1 - \lambda$, $z = (3\lambda + 2)/2$. Substituting into the constraint equation gives $\dfrac{\lambda}{2} - 2(1-\lambda) + 3 \cdot \dfrac{3\lambda+2}{2} = 6 \Rightarrow \lambda = \frac{5}{7}$, so $x = \frac{5}{14}$, $y = \frac{2}{7}$, and $z = \frac{29}{14}$. This must correspond to a minimum, so the point on the plane closest to the point $(0, 1, 1)$ is $\left(\frac{5}{14}, \frac{2}{7}, \frac{29}{14}\right)$.

33. Let $f(x, y, z) = d^2 = (x-4)^2 + (y-2)^2 + z^2$. Then we want to minimize f subject to the constraint $g(x, y, z) = x^2 + y^2 - z^2 = 0$. $\nabla f = \lambda \nabla g \Rightarrow \langle 2(x-4), 2(y-2), 2z\rangle = \langle 2\lambda x, 2\lambda y, -2\lambda z\rangle$, so $x - 4 = \lambda x$, $y - 2 = \lambda y$, and $z = -\lambda z$. From the last equation we have $z + \lambda z = 0 \Rightarrow z(1 + \lambda) = 0$, so either $z = 0$ or $\lambda = -1$. But from the constraint equation we have $z = 0 \Rightarrow x^2 + y^2 = 0 \Rightarrow x = y = 0$ which is not possible from the first two equations. So $\lambda = -1$ and $x - 4 = \lambda x \Rightarrow x = 2$, $y - 2 = \lambda y \Rightarrow y = 1$, and $x^2 + y^2 - z^2 = 0 \Rightarrow 4 + 1 - z^2 = 0 \Rightarrow z = \pm\sqrt{5}$. This must correspond to a minimum, so the points on the cone closest to $(4, 2, 0)$ are $\left(2, 1, \pm\sqrt{5}\right)$.

34. Let $f(x, y, z) = d^2 = x^2 + y^2 + z^2$. Then we want to minimize f subject to the constraint $g(x, y, z) = y^2 - xz = 9$. $\nabla f = \lambda \nabla g \Rightarrow \langle 2x, 2y, 2z\rangle = \langle -\lambda z, 2\lambda y, -\lambda x\rangle$, so $2x = -\lambda z$, $y = \lambda y$, and $2z = -\lambda x$. If $x = 0$ then the last equation implies $z = 0$, and from the constraint $y^2 - xz = 9$ we have $y = \pm 3$. If $x \neq 0$, then the first and third equations give $\lambda = -2x/z = -2z/x \Rightarrow x^2 = z^2$. From the second equation we have $y = 0$ or $\lambda = 1$. If $y = 0$ then $y^2 - xz = 9 \Rightarrow z = -9/x$ and $x^2 = z^2 \Rightarrow x^2 = 81/x^2 \Rightarrow x = \pm 3$. Since $z = -9/x$, $x = 3 \Rightarrow z = -3$ and $x = -3 \Rightarrow z = 3$. If $\lambda = 1$, then $2x = -z$ and $2z = -x$ which implies $x = z = 0$, contradicting the assumption that $x \neq 0$. Thus the possible points are $(0, \pm 3, 0)$, $(3, 0, -3)$, $(-3, 0, 3)$. We have $f(0, \pm 3, 0) = 9$ and $f(3, 0, -3) = f(-3, 0, 3) = 18$, so the points on the surface that are closest to the origin are $(0, \pm 3, 0)$.

35. $f(x, y, z) = xyz$, $g(x, y, z) = x + y + z = 100 \Rightarrow \nabla f = \langle yz, xz, xy\rangle = \lambda \nabla g = \langle \lambda, \lambda, \lambda\rangle$. Then $\lambda = yz = xz = xy$ implies $x = y = z = \frac{100}{3}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. Minimize $f(x,y,z) = x^2 + y^2 + z^2$ subject to $g(x,y,z) = x + y + z = 12$ with $x > 0$, $y > 0$, $z > 0$. Then $\nabla f = \lambda \nabla g \;\Rightarrow\; \langle 2x, 2y, 2z\rangle = \lambda\langle 1,1,1\rangle \;\Rightarrow\; 2x = \lambda,\ 2y = \lambda,\ 2z = \lambda \;\Rightarrow\; x = y = z$, so $x + y + z = 12 \;\Rightarrow\; 3x = 12 \;\Rightarrow\; x = 4 = y = z$. By comparing nearby values we can confirm that this gives a minimum and not a maximum. Thus the three numbers are 4, 4, and 4.

37. If the dimensions are $2x$, $2y$, and $2z$, then maximize $f(x,y,z) = (2x)(2y)(2z) = 8xyz$ subject to $g(x,y,z) = x^2 + y^2 + z^2 = r^2$ $(x > 0, y > 0, z > 0)$. Then $\nabla f = \lambda \nabla g \;\Rightarrow\; \langle 8yz, 8xz, 8xy\rangle = \lambda\langle 2x, 2y, 2z\rangle \;\Rightarrow\; 8yz = 2\lambda x$, $8xz = 2\lambda y$, and $8xy = 2\lambda z$, so $\lambda = \dfrac{4yz}{x} = \dfrac{4xz}{y} = \dfrac{4xy}{z}$. This gives $x^2 z = y^2 z \;\Rightarrow\; x^2 = y^2$ (since $z \neq 0$) and $xy^2 = xz^2 \;\Rightarrow\; z^2 = y^2$, so $x^2 = y^2 = z^2 \;\Rightarrow\; x = y = z$, and substituting into the constraint equation gives $3x^2 = r^2 \;\Rightarrow\; x = r/\sqrt{3} = y = z$. Thus the largest volume of such a box is $f\left(\frac{r}{\sqrt{3}}, \frac{r}{\sqrt{3}}, \frac{r}{\sqrt{3}}\right) = 8\left(\frac{r}{\sqrt{3}}\right)\left(\frac{r}{\sqrt{3}}\right)\left(\frac{r}{\sqrt{3}}\right) = \dfrac{8}{3\sqrt{3}} r^3$.

38. If the dimensions of the box are x, y, and z then minimize $f(x,y,z) = 2xy + 2xz + 2yz$ subject to $g(x,y,z) = xyz = 1000$ $(x > 0, y > 0, z > 0)$. Then $\nabla f = \lambda \nabla g \;\Rightarrow\; \langle 2y + 2z, 2x + 2z, 2x + 2y\rangle = \lambda\langle yz, xz, xy\rangle \;\Rightarrow\; 2y + 2z = \lambda yz$, $2x + 2z = \lambda xz$, $2x + 2y = \lambda xy$. Solving for λ in each equation gives $\lambda = \dfrac{2}{z} + \dfrac{2}{y} = \dfrac{2}{z} + \dfrac{2}{x} = \dfrac{2}{y} + \dfrac{2}{x} \;\Rightarrow\; x = y = z$. From $xyz = 1000$ we have $x^3 = 1000 \;\Rightarrow\; x = 10$ and the dimensions of the box are $x = y = z = 10$ cm.

39. $f(x,y,z) = xyz$, $g(x,y,z) = x + 2y + 3z = 6 \;\Rightarrow\; \nabla f = \langle yz, xz, xy\rangle = \lambda \nabla g = \langle \lambda, 2\lambda, 3\lambda\rangle$. Then $\lambda = yz = \frac{1}{2}xz = \frac{1}{3}xy$ implies $x = 2y$, $z = \frac{2}{3}y$. But $2y + 2y + 2y = 6$ so $y = 1$, $x = 2$, $z = \frac{2}{3}$ and the volume is $V = \frac{4}{3}$.

40. $f(x,y,z) = xyz$, $g(x,y,z) = xy + yz + xz = 32 \;\Rightarrow\; \nabla f = \langle yz, xz, xy\rangle = \lambda \nabla g = \langle \lambda(y+z), \lambda(x+z), \lambda(x+y)\rangle$. Then $\lambda(y+z) = yz$ **(1)**, $\lambda(x+z) = xz$ **(2)**, and $\lambda(x+y) = xy$ **(3)**. And **(1)** minus **(2)** implies $\lambda(y - x) = z(y - x)$ so $x = y$ or $\lambda = z$. If $\lambda = z$, then **(1)** implies $z(y+z) = yz$ or $z = 0$ which is false. Thus $x = y$. Similarly **(2)** minus **(3)** implies $\lambda(z - y) = x(z - y)$ so $y = z$ or $\lambda = x$. As above, $\lambda \neq x$, so $x = y = z$ and $3x^2 = 32$ or $x = y = z = \frac{8}{\sqrt{6}}$ cm.

41. $f(x,y,z) = xyz$, $g(x,y,z) = 4(x + y + z) = c \;\Rightarrow\; \nabla f = \langle yz, xz, xy\rangle$, $\lambda \nabla g = \langle 4\lambda, 4\lambda, 4\lambda\rangle$. Thus $4\lambda = yz = xz = xy$ or $x = y = z = \frac{1}{12}c$ are the dimensions giving the maximum volume.

42. $C(x,y,z) = 5xy + 2xz + 2yz$, $g(x,y,z) = xyz = V \;\Rightarrow$ $\nabla C = \langle 5y + 2z, 5x + 2z, 2x + 2y\rangle = \lambda \nabla g = \langle \lambda yz, \lambda xz, \lambda xy\rangle$. Then $\lambda yz = 5y + 2z$ **(1)**, $\lambda xz = 5x + 2z$ **(2)**, $\lambda xy = 2(x + y)$ **(3)**, and $xyz = V$ **(4)**. Now **(1)** − **(2)** implies $\lambda z(y - x) = 5(y - x)$, so $x = y$ or $\lambda = 5/z$, but z can't be 0, so $x = y$. Then twice **(2)** minus five times **(3)** together with $x = y$ implies $\lambda y(2x - 5y) = 2(2z - 5y)$ which gives

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$z = \frac{5}{2}y$ [again $\lambda \neq 2/y$ or else **(3)** implies $y = 0$]. Hence $\frac{5}{2}y^3 = V$ and the dimensions which minimize cost are $x = y = \sqrt[3]{\frac{2}{5}V}$ units, $z = V^{1/3}\left(\frac{5}{2}\right)^{2/3}$ units.

43. If the dimensions of the box are given by x, y, and z, then we need to find the maximum value of $f(x, y, z) = xyz$ $[x, y, z > 0]$ subject to the constraint $L = \sqrt{x^2 + y^2 + z^2}$ or $g(x, y, z) = x^2 + y^2 + z^2 = L^2$. $\nabla f = \lambda \nabla g \Rightarrow \langle yz, xz, xy \rangle = \lambda\langle 2x, 2y, 2z \rangle$, so $yz = 2\lambda x \Rightarrow \lambda = \dfrac{yz}{2x}$, $xz = 2\lambda y \Rightarrow \lambda = \dfrac{xz}{2y}$, and $xy = 2\lambda z \Rightarrow \lambda = \dfrac{xy}{2z}$.

Thus $\lambda = \dfrac{yz}{2x} = \dfrac{xz}{2y} \Rightarrow x^2 = y^2$ [since $z \neq 0$] $\Rightarrow x = y$ and $\lambda = \dfrac{yz}{2x} = \dfrac{xy}{2z} \Rightarrow x = z$ [since $y \neq 0$].

Substituting into the constraint equation gives $x^2 + x^2 + x^2 = L^2 \Rightarrow x^2 = L^2/3 \Rightarrow x = L/\sqrt{3} = y = z$ and the maximum volume is $\left(L/\sqrt{3}\right)^3 = L^3/\left(3\sqrt{3}\right)$.

44. Let the dimensions of the box be x, y, and z, so its volume is $f(x, y, z) = xyz$, its surface area is $2xy + 2yz + 2xz = 1500$ and its total edge length is $4x + 4y + 4z = 200$. We find the extreme values of $f(x, y, z)$ subject to the constraints $g(x, y, z) = xy + yz + xz = 750$ and $h(x, y, z) = x + y + z = 50$. Then $\nabla f = \langle yz, xz, xy \rangle = \lambda \nabla g + \mu \nabla h = \langle \lambda(y + z), \lambda(x + z), \lambda(x + y) \rangle + \langle \mu, \mu, \mu \rangle$. So $yz = \lambda(y + z) + \mu$ **(1)**, $xz = \lambda(x + z) + \mu$ **(2)**, and $xy = \lambda(x + y) + \mu$ **(3)**. Notice that the box can't be a cube or else $x = y = z = \frac{50}{3}$ but then $xy + yz + xz = \frac{2500}{3} \neq 750$. Assume x is the distinct side, that is, $x \neq y$, $x \neq z$. Then **(1)** minus **(2)** implies $z(y - x) = \lambda(y - x)$ or $\lambda = z$, and **(1)** minus **(3)** implies $y(z - x) = \lambda(z - x)$ or $\lambda = y$. So $y = z = \lambda$ and $x + y + z = 50$ implies $x = 50 - 2\lambda$; also $xy + yz + xz = 750$ implies $x(2\lambda) + \lambda^2 = 750$. Hence $50 - 2\lambda = \dfrac{750 - \lambda^2}{2\lambda}$ or $3\lambda^2 - 100\lambda + 750 = 0$ and $\lambda = \dfrac{50 \pm 5\sqrt{10}}{3}$, giving the points $\left(\frac{1}{3}\left(50 \mp 10\sqrt{10}\right), \frac{1}{3}\left(50 \pm 5\sqrt{10}\right), \frac{1}{3}\left(50 \pm 5\sqrt{10}\right)\right)$. Thus the minimum of f is $f\left(\frac{1}{3}\left(50 - 10\sqrt{3}\right), \frac{1}{3}\left(50 + 5\sqrt{10}\right), \frac{1}{3}\left(50 + 5\sqrt{10}\right)\right) = \frac{1}{27}\left(87{,}500 - 2500\sqrt{10}\right)$, and its maximum is $f\left(\frac{1}{3}\left(50 + 10\sqrt{10}\right), \frac{1}{3}\left(50 - 5\sqrt{10}\right), \frac{1}{3}\left(50 - 5\sqrt{10}\right)\right) = \frac{1}{27}\left(87{,}500 + 2500\sqrt{10}\right)$.

Note: If either y or z is the distinct side, then symmetry gives the same result.

45. We need to find the extreme values of $f(x, y, z) = x^2 + y^2 + z^2$ subject to the two constraints $g(x, y, z) = x + y + 2z = 2$ and $h(x, y, z) = x^2 + y^2 - z = 0$. $\nabla f = \langle 2x, 2y, 2z \rangle$, $\lambda \nabla g = \langle \lambda, \lambda, 2\lambda \rangle$ and $\mu \nabla h = \langle 2\mu x, 2\mu y, -\mu \rangle$. Thus we need $2x = \lambda + 2\mu x$ **(1)**, $2y = \lambda + 2\mu y$ **(2)**, $2z = 2\lambda - \mu$ **(3)**, $x + y + 2z = 2$ **(4)**, and $x^2 + y^2 - z = 0$ **(5)**. From **(1)** and **(2)**, $2(x - y) = 2\mu(x - y)$, so if $x \neq y$, $\mu = 1$. Putting this in **(3)** gives $2z = 2\lambda - 1$ or $\lambda = z + \frac{1}{2}$, but putting $\mu = 1$ into **(1)** says $\lambda = 0$. Hence $z + \frac{1}{2} = 0$ or $z = -\frac{1}{2}$. Then **(4)** and **(5)** become $x + y - 3 = 0$ and $x^2 + y^2 + \frac{1}{2} = 0$. The last equation cannot be true, so this case gives no solution. So we must have $x = y$. Then **(4)** and **(5)** become $2x + 2z = 2$ and $2x^2 - z = 0$ which imply $z = 1 - x$ and $z = 2x^2$. Thus $2x^2 = 1 - x$ or $2x^2 + x - 1 = (2x - 1)(x + 1) = 0$ so $x = \frac{1}{2}$ or

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x = -1$. The two points to check are $\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$ and $(-1, -1, 2)$: $f\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) = \frac{3}{4}$ and $f(-1, -1, 2) = 6$. Thus $\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$ is the point on the ellipse nearest the origin and $(-1, -1, 2)$ is the one farthest from the origin.

46. (a) After plotting $z = \sqrt{x^2 + y^2}$, the top half of the cone, and the plane $z = (5 - 4x + 3y)/8$ we see the ellipse formed by the intersection of the surfaces. The ellipse can be plotted explicitly using cylindrical coordinates (see Section 15.7): The cone is given by $z = r$, and the plane is $4r\cos\theta - 3r\sin\theta + 8z = 5$. Substituting $z = r$ into the plane equation gives $4r\cos\theta - 3r\sin\theta + 8r = 5 \quad \Rightarrow \quad r = \dfrac{5}{4\cos\theta - 3\sin\theta + 8}$. Since $z = r$ on the ellipse, parametric equations (in cylindrical coordinates) are $\theta = t$, $r = z = \dfrac{5}{4\cos t - 3\sin t + 8}$, $0 \le t \le 2\pi$.



(b) We need to find the extreme values of $f(x, y, z) = z$ subject to the two constraints $g(x, y, z) = 4x - 3y + 8z = 5$ and $h(x, y, z) = x^2 + y^2 - z^2 = 0$.

$\nabla f = \lambda\nabla g + \mu\nabla h \quad \Rightarrow \quad \langle 0, 0, 1\rangle = \lambda\langle 4, -3, 8\rangle + \mu\langle 2x, 2y, -2z\rangle$, so we need $4\lambda + 2\mu x = 0 \quad \Rightarrow \quad x = -\frac{2\lambda}{\mu}$ **(1)**, $-3\lambda + 2\mu y = 0 \quad \Rightarrow \quad y = \frac{3\lambda}{2\mu}$ **(2)**, $8\lambda - 2\mu z = 1 \quad \Rightarrow \quad z = \frac{8\lambda - 1}{2\mu}$ **(3)**, $4x - 3y + 8z = 5$ **(4)**, and $x^2 + y^2 = z^2$ **(5)**. [Note that $\mu \neq 0$, else $\lambda = 0$ from **(1)**, but substitution into **(3)** gives a contradiction.]

Substituting **(1)**, **(2)**, and **(3)** into **(4)** gives $4\left(-\frac{2\lambda}{\mu}\right) - 3\left(\frac{3\lambda}{2\mu}\right) + 8\left(\frac{8\lambda - 1}{2\mu}\right) = 5 \quad \Rightarrow \quad \mu = \frac{39\lambda - 8}{10}$ and into **(5)** gives $\left(-\frac{2\lambda}{\mu}\right)^2 + \left(\frac{3\lambda}{2\mu}\right)^2 = \left(\frac{8\lambda - 1}{2\mu}\right)^2 \quad \Rightarrow \quad 16\lambda^2 + 9\lambda^2 = (8\lambda - 1)^2 \quad \Rightarrow \quad 39\lambda^2 - 16\lambda + 1 = 0 \quad \Rightarrow \quad \lambda = \frac{1}{13}$ or $\lambda = \frac{1}{3}$.

If $\lambda = \frac{1}{13}$ then $\mu = -\frac{1}{2}$ and $x = \frac{4}{13}$, $y = -\frac{3}{13}$, $z = \frac{5}{13}$. If $\lambda = \frac{1}{3}$ then $\mu = \frac{1}{2}$ and $x = -\frac{4}{3}$, $y = 1$, $z = \frac{5}{3}$. Thus the highest point on the ellipse is $\left(-\frac{4}{3}, 1, \frac{5}{3}\right)$ and the lowest point is $\left(\frac{4}{13}, -\frac{3}{13}, \frac{5}{13}\right)$.

47. $f(x, y, z) = ye^{x-z}$, $g(x, y, z) = 9x^2 + 4y^2 + 36z^2 = 36$, $h(x, y, z) = xy + yz = 1$. $\nabla f = \lambda\nabla g + \mu\nabla h \quad \Rightarrow$ $\langle ye^{x-z}, e^{x-z}, -ye^{x-z}\rangle = \lambda\langle 18x, 8y, 72z\rangle + \mu\langle y, x + z, y\rangle$, so $ye^{x-z} = 18\lambda x + \mu y$, $e^{x-z} = 8\lambda y + \mu(x + z)$, $-ye^{x-z} = 72\lambda z + \mu y$, $9x^2 + 4y^2 + 36z^2 = 36$, $xy + yz = 1$. Using a CAS to solve these 5 equations simultaneously for x, y, z, λ, and μ (in Maple, use the `allvalues` command), we get 4 real-valued solutions:

$$\begin{array}{lllll}
x \approx 0.222444, & y \approx -2.157012, & z \approx -0.686049, & \lambda \approx -0.200401, & \mu \approx 2.108584 \\
x \approx -1.951921, & y \approx -0.545867, & z \approx 0.119973, & \lambda \approx 0.003141, & \mu \approx -0.076238 \\
x \approx 0.155142, & y \approx 0.904622, & z \approx 0.950293, & \lambda \approx -0.012447, & \mu \approx 0.489938 \\
x \approx 1.138731, & y \approx 1.768057, & z \approx -0.573138, & \lambda \approx 0.317141, & \mu \approx 1.862675
\end{array}$$

Substituting these values into f gives $f(0.222444, -2.157012, -0.686049) \approx -5.3506$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$f(-1.951921, -0.545867, 0.119973) \approx -0.0688$, $f(0.155142, 0.904622, 0.950293) \approx 0.4084$, $f(1.138731, 1.768057, -0.573138) \approx 9.7938$. Thus the maximum is approximately 9.7938, and the minimum is approximately -5.3506.

48. $f(x, y, z) = x + y + z$, $g(x, y, z) = x^2 - y^2 - z = 0$, $h(x, y, z) = x^2 + z^2 = 4$.

$\nabla f = \lambda \nabla g + \mu \nabla h \quad \Rightarrow \quad \langle 1, 1, 1 \rangle = \lambda \langle 2x, -2y, -1 \rangle + \mu \langle 2x, 0, 2z \rangle$, so $1 = 2\lambda x + 2\mu x$, $1 = -2\lambda y$, $1 = -\lambda + 2\mu z$, $x^2 - y^2 = z$, $x^2 + z^2 = 4$. Using a CAS to solve these 5 equations simultaneously for x, y, z, λ, and μ, we get 4 real-valued solutions:

$$\begin{array}{lllll} x \approx -1.652878, & y \approx -1.964194, & z \approx -1.126052, & \lambda \approx 0.254557, & \mu \approx -0.557060 \\ x \approx -1.502800, & y \approx 0.968872, & z \approx 1.319694, & \lambda \approx -0.516064, & \mu \approx 0.183352 \\ x \approx -0.992513, & y \approx 1.649677, & z \approx -1.736352, & \lambda \approx -0.303090, & \mu \approx -0.200682 \\ x \approx 1.895178, & y \approx 1.718347, & z \approx 0.638984, & \lambda \approx -0.290977, & \mu \approx 0.554805 \end{array}$$

Substituting these values into f gives $f(-1.652878, -1.964194, -1.126052) \approx -4.7431$, $f(-1.502800, 0.968872, 1.319694) \approx 0.7858$, $f(-0.992513, 1.649677, -1.736352) \approx -1.0792$, $f(1.895178, 1.718347, 0.638984) \approx 4.2525$. Thus the maximum is approximately 4.2525, and the minimum is approximately -4.7431.

49. (a) We wish to maximize $f(x_1, x_2, \ldots, x_n) = \sqrt[n]{x_1 x_2 \cdots x_n}$ subject to

$g(x_1, x_2, \ldots, x_n) = x_1 + x_2 + \cdots + x_n = c$ and $x_i > 0$.

$\nabla f = \left\langle \frac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_2 \cdots x_n), \frac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_1 x_3 \cdots x_n), \ldots, \frac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_1 \cdots x_{n-1}) \right\rangle$

and $\lambda \nabla g = \langle \lambda, \lambda, \ldots, \lambda \rangle$, so we need to solve the system of equations

$$\begin{aligned} \tfrac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_2 \cdots x_n) = \lambda \quad &\Rightarrow \quad x_1^{1/n} x_2^{1/n} \cdots x_n^{1/n} = n\lambda x_1 \\ \tfrac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_1 x_3 \cdots x_n) = \lambda \quad &\Rightarrow \quad x_1^{1/n} x_2^{1/n} \cdots x_n^{1/n} = n\lambda x_2 \\ &\vdots \\ \tfrac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_1 \cdots x_{n-1}) = \lambda \quad &\Rightarrow \quad x_1^{1/n} x_2^{1/n} \cdots x_n^{1/n} = n\lambda x_n \end{aligned}$$

This implies $n\lambda x_1 = n\lambda x_2 = \cdots = n\lambda x_n$. Note $\lambda \neq 0$, otherwise we can't have all $x_i > 0$. Thus $x_1 = x_2 = \cdots = x_n$. But $x_1 + x_2 + \cdots + x_n = c \quad \Rightarrow \quad nx_1 = c \quad \Rightarrow \quad x_1 = \dfrac{c}{n} = x_2 = x_3 = \cdots = x_n$. Then the only point where f can have an extreme value is $\left(\frac{c}{n}, \frac{c}{n}, \ldots, \frac{c}{n}\right)$. Since we can choose values for $(x_1, x_2, \ldots, x_n)$ that make f as close to zero (but not equal) as we like, f has no minimum value. Thus the maximum value is

$$f\left(\frac{c}{n}, \frac{c}{n}, \ldots, \frac{c}{n}\right) = \sqrt[n]{\frac{c}{n} \cdot \frac{c}{n} \cdots \frac{c}{n}} = \frac{c}{n}.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) From part (a), $\dfrac{c}{n}$ is the maximum value of f. Thus $f(x_1, x_2, \ldots, x_n) = \sqrt[n]{x_1 x_2 \cdots x_n} \le \dfrac{c}{n}$. But $x_1 + x_2 + \cdots + x_n = c$, so $\sqrt[n]{x_1 x_2 \cdots x_n} \le \dfrac{x_1 + x_2 + \cdots + x_n}{n}$. These two means are equal when f attains its maximum value $\dfrac{c}{n}$, but this can occur only at the point $\left(\dfrac{c}{n}, \dfrac{c}{n}, \ldots, \dfrac{c}{n}\right)$ we found in part (a). So the means are equal only when $x_1 = x_2 = x_3 = \cdots = x_n = \dfrac{c}{n}$.

50. (a) Let $f(x_1, \ldots, x_n, y_1, \ldots, y_n) = \sum_{i=1}^{n} x_i y_i$, $g(x_1, \ldots, x_n) = \sum_{i=1}^{n} x_i^2$, and $h(x_1, \ldots, x_n) = \sum_{i=1}^{n} y_i^2$. Then $\nabla f = \nabla \sum_{i=1}^{n} x_i y_i = \langle y_1, y_2, \ldots, y_n, x_1, x_2, \ldots, x_n \rangle$, $\nabla g = \nabla \sum_{i=1}^{n} x_i^2 = \langle 2x_1, 2x_2, \ldots, 2x_n, 0, 0, \ldots, 0 \rangle$ and $\nabla h = \nabla \sum_{i=1}^{n} y_i^2 = \langle 0, 0, \ldots, 0, 2y_1, 2y_2, \ldots, 2y_n \rangle$. So $\nabla f = \lambda \nabla g + \mu \nabla h \;\Leftrightarrow\; y_i = 2\lambda x_i$ and $x_i = 2\mu y_i$, $1 \le i \le n$. Then $1 = \sum_{i=1}^{n} y_i^2 = \sum_{i=1}^{n} 4\lambda^2 x_i^2 = 4\lambda^2 \sum_{i=1}^{n} x_i^2 = 4\lambda^2 \;\Rightarrow\; \lambda = \pm\frac{1}{2}$. If $\lambda = \frac{1}{2}$ then $y_i = 2\left(\frac{1}{2}\right)x_i = x_i$, $1 \le i \le n$. Thus $\sum_{i=1}^{n} x_i y_i = \sum_{i=1}^{n} x_i^2 = 1$. Similarly if $\lambda = -\frac{1}{2}$ we get $y_i = -x_i$ and $\sum_{i=1}^{n} x_i y_i = -1$. Similarly we get $\mu = \pm\frac{1}{2}$ giving $y_i = \pm x_i$, $1 \le i \le n$, and $\sum_{i=1}^{n} x_i y_i = \pm 1$. Thus the maximum value of $\sum_{i=1}^{n} x_i y_i$ is 1.

(b) Here we assume $\sum_{i=1}^{n} a_i^2 \ne 0$ and $\sum_{i=1}^{n} b_i^2 \ne 0$. (If $\sum_{i=1}^{n} a_i^2 = 0$, then each $a_i = 0$ and so the inequality is trivially true.)

$x_i = \dfrac{a_i}{\sqrt{\sum a_j^2}} \;\Rightarrow\; \sum x_i^2 = \dfrac{\sum a_i^2}{\sum a_j^2} = 1$, and $y_i = \dfrac{b_i}{\sqrt{\sum b_j^2}} \;\Rightarrow\; \sum y_i^2 = \dfrac{\sum b_i^2}{\sum b_j^2} = 1$. Therefore, from part (a),

$$\sum x_i y_i = \sum \frac{a_i b_i}{\sqrt{\sum a_j^2}\sqrt{\sum b_j^2}} \le 1 \;\Leftrightarrow\; \sum a_i b_i \le \sqrt{\sum a_j^2}\sqrt{\sum b_j^2}.$$

APPLIED PROJECT Rocket Science

1. Initially the rocket engine has mass $M_r = M_1$ and payload mass $P = M_2 + M_3 + A$. Then the change in velocity resulting from the first stage is $\Delta V_1 = -c \ln\left(1 - \dfrac{(1 - S)M_1}{M_2 + M_3 + A + M_1}\right)$. After the first stage is jettisoned we can consider the rocket engine to have mass $M_r = M_2$ and the payload to have mass $P = M_3 + A$. The resulting change in velocity from the second stage is $\Delta V_2 = -c \ln\left(1 - \dfrac{(1 - S)M_2}{M_3 + A + M_2}\right)$. When only the third stage remains, we have $M_r = M_3$ and $P = A$, so the resulting change in velocity is $\Delta V_3 = -c \ln\left(1 - \dfrac{(1 - S)M_3}{A + M_3}\right)$. Since the rocket started from rest, the final velocity

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

attained is

$$\begin{aligned} v_f &= \Delta V_1 + \Delta V_2 + \Delta V_3 \\ &= -c\ln\left(1 - \frac{(1-S)M_1}{M_2+M_3+A+M_1}\right) + (-c)\ln\left(1 - \frac{(1-S)M_2}{M_3+A+M_2}\right) + (-c)\ln\left(1 - \frac{(1-S)M_3}{A+M_3}\right) \\ &= -c\left[\ln\left(\frac{M_1+M_2+M_3+A-(1-S)M_1}{M_1+M_2+M_3+A}\right) + \ln\left(\frac{M_2+M_3+A-(1-S)M_2}{M_2+M_3+A}\right)\right. \\ &\qquad \left. + \ln\left(\frac{M_3+A-(1-S)M_3}{M_3+A}\right)\right] \\ &= c\left[\ln\left(\frac{M_1+M_2+M_3+A}{SM_1+M_2+M_3+A}\right) + \ln\left(\frac{M_2+M_3+A}{SM_2+M_3+A}\right) + \ln\left(\frac{M_3+A}{SM_3+A}\right)\right] \end{aligned}$$

2. Define $N_1 = \dfrac{M_1+M_2+M_3+A}{SM_1+M_2+M_3+A}$, $N_2 = \dfrac{M_2+M_3+A}{SM_2+M_3+A}$, and $N_3 = \dfrac{M_3+A}{SM_3+A}$. Then

$$\begin{aligned} \frac{(1-S)N_1}{1-SN_1} &= \frac{(1-S)\dfrac{M_1+M_2+M_3+A}{SM_1+M_2+M_3+A}}{1-S\dfrac{M_1+M_2+M_3+A}{SM_1+M_2+M_3+A}} = \frac{(1-S)(M_1+M_2+M_3+A)}{SM_1+M_2+M_3+A-S(M_1+M_2+M_3+A)} \\ &= \frac{(1-S)(M_1+M_2+M_3+A)}{(1-S)(M_2+M_3+A)} = \frac{M_1+M_2+M_3+A}{M_2+M_3+A} \end{aligned}$$

as desired.

Similarly,

$$\frac{(1-S)N_2}{1-SN_2} = \frac{(1-S)(M_2+M_3+A)}{SM_2+M_3+A-S(M_2+M_3+A)} = \frac{(1-S)(M_2+M_3+A)}{(1-S)(M_3+A)} = \frac{M_2+M_3+A}{M_3+A}$$

and

$$\frac{(1-S)N_3}{1-SN_3} = \frac{(1-S)(M_3+A)}{SM_3+A-S(M_3+A)} = \frac{(1-S)(M_3+A)}{(1-S)(A)} = \frac{M_3+A}{A}$$

Then

$$\begin{aligned} \frac{M+A}{A} &= \frac{M_1+M_2+M_3+A}{A} = \frac{M_1+M_2+M_3+A}{M_2+M_3+A}\cdot\frac{M_2+M_3+A}{M_3+A}\cdot\frac{M_3+A}{A} \\ &= \frac{(1-S)N_1}{1-SN_1}\cdot\frac{(1-S)N_2}{1-SN_2}\cdot\frac{(1-S)N_3}{1-SN_3} = \frac{(1-S)^3N_1N_2N_3}{(1-SN_1)(1-SN_2)(1-SN_3)} \end{aligned}$$

3. Since $A > 0$, $M + A$ and consequently $\dfrac{M+A}{A}$ is minimized for the same values as M. $\ln x$ is a strictly increasing function, so $\ln\left(\dfrac{M+A}{A}\right)$ must give a minimum for the same values as $\dfrac{M+A}{A}$ and hence M. We then wish to minimize $\ln\left(\dfrac{M+A}{A}\right)$ subject to the constraint $c(\ln N_1 + \ln N_2 + \ln N_3) = v_f$. From Problem 2,

$$\begin{aligned} \ln\left(\frac{M+A}{A}\right) &= \ln\left(\frac{(1-S)^3 N_1N_2N_3}{(1-SN_1)(1-SN_2)(1-SN_3)}\right) \\ &= 3\ln(1-S) + \ln N_1 + \ln N_2 + \ln N_3 - \ln(1-SN_1) - \ln(1-SN_2) - \ln(1-SN_3) \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Using the method of Lagrange multipliers, we need to solve $\nabla\left[\ln\left(\frac{M+A}{A}\right)\right] = \lambda\nabla[c(\ln N_1 + \ln N_2 + \ln N_3)]$ with $c(\ln N_1 + \ln N_2 + \ln N_3) = v_f$ in terms of N_1, N_2, and N_3. The resulting system is

$$\frac{1}{N_1} + \frac{S}{1-SN_1} = \lambda\frac{c}{N_1} \qquad \frac{1}{N_2} + \frac{S}{1-SN_2} = \lambda\frac{c}{N_2} \qquad \frac{1}{N_3} + \frac{S}{1-SN_3} = \lambda\frac{c}{N_3}$$

$$c(\ln N_1 + \ln N_2 + \ln N_3) = v_f$$

One approach to solving the system is isolating $c\lambda$ in the first three equations which gives

$1 + \frac{SN_1}{1-SN_1} = c\lambda = 1 + \frac{SN_2}{1-SN_2} = 1 + \frac{SN_3}{1-SN_3} \Rightarrow \frac{N_1}{1-SN_1} = \frac{N_2}{1-SN_2} = \frac{N_3}{1-SN_3} \Rightarrow$

$N_1 = N_2 = N_3$ (Verify!). This says the fourth equation can be expressed as $c(\ln N_1 + \ln N_1 + \ln N_1) = v_f \Rightarrow$

$3c\ln N_1 = v_f \Rightarrow \ln N_1 = \frac{v_f}{3c}$. Thus the minimum mass M of the rocket engine is attained for

$N_1 = N_2 = N_3 = e^{v_f/(3c)}$.

4. Using the previous results, $\frac{M+A}{A} = \frac{(1-S)^3 N_1N_2N_3}{(1-SN_1)(1-SN_2)(1-SN_3)} = \frac{(1-S)^3\left[e^{v_f/(3c)}\right]^3}{\left[1-Se^{v_f/(3c)}\right]^3} = \frac{(1-S)^3 e^{v_f/c}}{\left[1-Se^{v_f/(3c)}\right]^3}$.

Then $M = \frac{A(1-S)^3 e^{v_f/c}}{\left[1-Se^{v_f/(3c)}\right]^3} - A$.

5. (a) From Problem 4, $M = \frac{A(1-0.2)^3 e^{(17{,}500/6000)}}{(1-0.2e^{[17{,}500/(3\cdot 6000)]})^3} - A \approx 90.4A - A = 89.4A$.

(b) First, $N_3 = \frac{M_3+A}{SM_3+A} \Rightarrow e^{[17{,}500/(3\cdot 6000)]} = \frac{M_3+A}{0.2M_3+A} \Rightarrow M_3 = \frac{A(1-e^{35/36})}{0.2e^{35/36}-1} \approx 3.49A$.

Then $N_2 = \frac{M_2+M_3+A}{SM_2+M_3+A} = \frac{M_2+3.49A+A}{0.2M_2+3.49A+A} \Rightarrow M_2 = \frac{4.49A(1-e^{35/36})}{0.2e^{35/36}-1} \approx 15.67A$ and

$N_3 = \frac{M_1+M_2+M_3+A}{SM_1+M_2+M_3+A} = \frac{M_1+15.67A+3.49A+A}{0.2M_1+15.67A+3.49A+A} \Rightarrow M_1 = \frac{20.16A(1-e^{35/36})}{0.2e^{35/36}-1} \approx 70.36A$.

6. As in Problem 5, $N_3 = \frac{M_3+A}{SM_3+A} \Rightarrow e^{24{,}700/(3\cdot 6000)} = \frac{M_3+A}{0.2M_3+A} \Rightarrow M_3 = \frac{A(1-e^{247/180})}{0.2e^{247/180}-1} \approx 13.9A$,

$N_2 = \frac{M_2+M_3+A}{SM_2+M_3+A} = \frac{M_2+13.9A+A}{0.2M_2+13.9A+A} \Rightarrow M_2 = \frac{14.9A(1-e^{247/180})}{0.2e^{247/180}-1} \approx 208A$, and

$N_3 = \frac{M_1+M_2+M_3+A}{SM_1+M_2+M_3+A} = \frac{M_1+208A+13.9A+A}{0.2M_1+208A+13.9A+A} \Rightarrow M_1 = \frac{222.9A(1-e^{247/180})}{0.2e^{247/180}-1} \approx 3110A$.

Here $A = 500$, so the mass of each stage of the rocket engine is approximately $M_1 = 3110(500) = 1{,}550{,}000$ lb, $M_2 = 208(500) = 104{,}000$ lb, and $M_3 = 13.9(500) = 6950$ lb.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

APPLIED PROJECT Hydro-Turbine Optimization

1. We wish to maximize the total energy production for a given total flow, so we can say Q_T is fixed and we want to maximize $KW_1 + KW_2 + KW_3$. Notice each KW_i has a constant factor $\left(170 - 1.6 \cdot 10^{-6} Q_T^2\right)$, so to simplify the computations we can equivalently maximize

$$\begin{aligned} f(Q_1, Q_2, Q_3) &= \frac{KW_1 + KW_2 + KW_3}{170 - 1.6 \cdot 10^{-6} Q_T^2} \\ &= \left(-18.89 + 0.1277Q_1 - 4.08 \cdot 10^{-5} Q_1^2\right) + \left(-24.51 + 0.1358Q_2 - 4.69 \cdot 10^{-5} Q_2^2\right) \\ &\qquad + \left(-27.02 + 0.1380Q_3 - 3.84 \cdot 10^{-5} Q_3^2\right) \end{aligned}$$

subject to the constraint $g(Q_1, Q_2, Q_3) = Q_1 + Q_2 + Q_3 = Q_T$. So first we find the values of Q_1, Q_2, Q_3 where $\nabla f(Q_1, Q_2, Q_3) = \lambda \nabla g(Q_1, Q_2, Q_3)$ and $Q_1 + Q_2 + Q_3 = Q_T$ which is equivalent to solving the system

$$\begin{aligned} 0.1277 - 2(4.08 \cdot 10^{-5}) Q_1 &= \lambda \\ 0.1358 - 2(4.69 \cdot 10^{-5}) Q_2 &= \lambda \\ 0.1380 - 2(3.84 \cdot 10^{-5}) Q_3 &= \lambda \\ Q_1 + Q_2 + Q_3 &= Q_T \end{aligned}$$

Comparing the first and third equations, we have $0.1277 - 2(4.08 \cdot 10^{-5}) Q_1 = 0.1380 - 2(3.84 \cdot 10^{-5}) Q_3 \Rightarrow Q_1 = -126.2255 + 0.9412Q_3$. From the second and third equations, $0.1358 - 2(4.69 \cdot 10^{-5}) Q_2 = 0.1380 - 2(3.84 \cdot 10^{-5}) Q_3 \Rightarrow Q_2 = -23.4542 + 0.8188Q_3$. Substituting into $Q_1 + Q_2 + Q_3 = Q_T$ gives $(-126.2255 + 0.9412Q_3) + (-23.4542 + 0.8188Q_3) + Q_3 = Q_T \Rightarrow 2.76Q_3 = Q_T + 149.6797 \Rightarrow Q_3 = 0.3623Q_T + 54.23$. Then $Q_1 = -126.2255 + 0.9412Q_3 = -126.2255 + 0.9412(0.3623Q_T + 54.23) = 0.3410Q_T - 75.18$ and $Q_2 = -23.4542 + 0.8188(0.3623Q_T + 54.23) = 0.2967Q_T + 20.95$. As long as we maintain $250 \le Q_1 \le 1110$, $250 \le Q_2 \le 1110$, and $250 \le Q_3 \le 1225$, we can reason from the nature of the functions KW_i that these values give a maximum of f, and hence a maximum energy production, and not a minimum.

2. From Problem 1, the value of Q_1 that maximizes energy production is $0.3410Q_T - 75.18$, but since $250 \le Q_1 \le 1110$, we must have $250 \le 0.3410Q_T - 75.18 \le 1110 \Rightarrow 325.18 \le 0.3410Q_T \le 1185.18 \Rightarrow 953.6 \le Q_T \le 3475.6$. Similarly, $250 \le Q_2 \le 1110 \Rightarrow 250 \le 0.2967Q_T + 20.95 \le 1110 \Rightarrow 772.0 \le Q_T \le 3670.5$, and $250 \le Q_3 \le 1225 \Rightarrow 250 \le 0.3623Q_T + 54.23 \le 1225 \Rightarrow 540.4 \le Q_T \le 3231.5$. Consolidating these results, we see that the values from Problem 1 are applicable only for $953.6 \le Q_T \le 3231.5$.

3. If $Q_T = 2500$, the results from Problem 1 show that the maximum energy production occurs for

$$\begin{aligned} Q_1 &= 0.3410Q_T - 75.18 = 0.3410(2500) - 75.18 = 777.3 \\ Q_2 &= 0.2967Q_T + 20.95 = 0.2967(2500) + 20.95 = 762.7 \\ Q_3 &= 0.3623Q_T + 54.23 = 0.3623(2500) + 54.23 = 960.0 \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The energy produced for these values is $KW_1 + KW_2 + KW_3 \approx 8915.2 + 8285.1 + 11{,}211.3 \approx 28{,}411.6$.

We compute the energy production for a nearby distribution, $Q_1 = 770$, $Q_2 = 760$, and $Q_3 = 970$: $KW_1 + KW_2 + KW_3 \approx 8839.8 + 8257.4 + 11{,}313.5 = 28{,}410.7$. For another example, we take $Q_1 = 780$, $Q_2 = 765$, and $Q_3 = 955$: $KW_1 + KW_2 + KW_3 \approx 8942.9 + 8308.8 + 11{,}159.7 = 28{,}411.4$. These distributions are both close to the distribution from Problem 1 and both give slightly lower energy productions, suggesting that $Q_1 = 777.3$, $Q_2 = 762.7$, and $Q_3 = 960.0$ is indeed the optimal distribution.

4. First we graph each power function in its domain if all of the flow is directed to that turbine (so $Q_i = Q_T$). If we use only one turbine, the graph indicates that for a water flow of 1000 ft^3/s, Turbine 3 produces the most power, approximately 12,200 kW. In comparison, if we use all three turbines, the results of Problem 1 with $Q_T = 1000$ give $Q_1 = 265.8$, $Q_2 = 317.7$, and $Q_3 = 416.5$, resulting in a total energy production of $KW_1 + KW_2 + KW_3 \approx 8397.4$ kW. Here, using only one turbine produces significantly more energy! If the flow is only 600 ft^3/s, we do not have the option of using all three turbines, as the domain restrictions require a minimum of 250 ft^3/s in each turbine. We can use just one turbine, then, and from the graph Turbine 1 produces the most energy for a water flow of 600 ft^3.



5. If we examine the graph from Problem 4, we see that for water flows above approximately 450 ft^3/s, Turbine 2 produces the least amount of power. Therefore it seems reasonable to assume that we should distribute the incoming flow of 1500 ft^3/s between Turbines 1 and 3. (This can be verified by computing the power produced with the other pairs of turbines for comparison.) So now we wish to maximize $KW_1 + KW_3$ subject to the constraint $Q_1 + Q_3 = Q_T$ where $Q_T = 1500$.

As in Problem 1, we can equivalently maximize

$$f(Q_1, Q_3) = \frac{KW_1 + KW_3}{170 - 1.6 \cdot 10^{-6} Q_T^2}$$
$$= \left(-18.89 + 0.1277Q_1 - 4.08 \cdot 10^{-5} Q_1^2\right) + \left(-27.02 + 0.1380Q_3 - 3.84 \cdot 10^{-5} Q_3^2\right)$$

subject to the constraint $g(Q_1, Q_3) = Q_1 + Q_3 = Q_T$.

Then we solve $\nabla f(Q_1, Q_3) = \lambda \nabla g(Q_1, Q_3) \quad \Rightarrow \quad 0.1277 - 2\left(4.08 \cdot 10^{-5}\right) Q_1 = \lambda$ and $0.1380 - 2(3.84 \cdot 10^{-5})Q_3 = \lambda$, thus $0.1277 - 2(4.08 \cdot 10^{-5})Q_1 = 0.1380 - 2(3.84 \cdot 10^{-5})Q_3 \quad \Rightarrow$ $Q_1 = -126.2255 + 0.9412Q_3$. Substituting into $Q_1 + Q_3 = Q_T$ gives $-126.2255 + 0.9412Q_3 + Q_3 = 1500 \quad \Rightarrow$ $Q_3 \approx 837.7$, and then $Q_1 = Q_T - Q_3 \approx 1500 - 837.7 = 662.3$. So we should apportion approximately 662.3 ft^3/s to Turbine 1 and the remaining 837.7 ft^3/s to Turbine 3. The resulting energy production is $KW_1 + KW_3 \approx 7952.1 + 10{,}256.2 = 18{,}208.3$ kW. (We can verify that this is indeed a maximum energy production by

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

checking nearby distributions.) In comparison, if we use all three turbines with $Q_T = 1500$ we get $Q_1 = 436.3$, $Q_2 = 466.0$, and $Q_3 = 597.7$, resulting in a total energy production of $KW_1 + KW_2 + KW_3 \approx 16{,}538.7$ kW. Clearly, for this flow level it is beneficial to use only two turbines.

6. Note that an incoming flow of 3400 ft^3/s is not within the domain we established in Problem 2, so we cannot simply use our previous work to give the optimal distribution. We will need to use all three turbines, due to the capacity limitations of each individual turbine, but 3400 is less than the maximum combined capacity of 3445 ft^3/s, so we still must decide how to distribute the flows. From the graph in Problem 4, Turbine 3 produces the most power for the higher flows, so it seems reasonable to use Turbine 3 at its maximum capacity of 1225 and distribute the remaining 2175 ft^3/s flow between Turbines 1 and 2. We can again use the technique of Lagrange multipliers to determine the optimal distribution. Following the procedure we used in Problem 5, we wish to maximize $KW_1 + KW_2$ subject to the constraint $Q_1 + Q_2 = Q_T$ where $Q_T = 2175$. We can equivalently maximize

$$\begin{aligned} f(Q_1, Q_2) &= \frac{KW_1 + KW_2}{170 - 1.6 \cdot 10^{-6} Q_T^2} \\ &= \left(-18.89 + 0.1277Q_1 - 4.08 \cdot 10^{-5} Q_1^2\right) + \left(-24.51 + 0.1358Q_2 - 4.69 \cdot 10^{-5} Q_2^2\right) \end{aligned}$$

subject to the constraint $g(Q_1, Q_2) = Q_1 + Q_2 = Q_T$. Then we solve $\nabla f(Q_1, Q_2) = \lambda \nabla g(Q_1, Q_2) \quad \Rightarrow$ $0.1277 - 2(4.08 \cdot 10^{-5})Q_1 = \lambda$ and $0.1358 - 2(4.69 \cdot 10^{-5})Q_2 = \lambda$, thus $0.1277 - 2(4.08 \cdot 10^{-5})Q_1 = 0.1358 - 2(4.69 \cdot 10^{-5})Q_2 \quad \Rightarrow \quad Q_1 = -99.2647 + 1.1495Q_2$. Substituting into $Q_1 + Q_2 = Q_T$ gives $-99.2647 + 1.1495Q_2 + Q_2 = 2175 \quad \Rightarrow \quad Q_2 \approx 1058.0$, and then $Q_1 \approx 1117.0$. This value for Q_1 is larger than the allowable maximum flow to Turbine 1, but the result indicates that the flow to Turbine 1 should be maximized. Thus we should recommend that the company apportion the maximum allowable flows to Turbines 1 and 3, 1110 and 1225 ft^3/s, and the remaining 1065 ft^3/s to Turbine 2. Checking nearby distributions within the domain verifies that we have indeed found the optimal distribution.

14 Review

TRUE-FALSE QUIZ

1. True. $f_y(a,b) = \lim_{h \to 0} \dfrac{f(a, b+h) \quad f(a,b)}{h}$ from Equation 14.3.3. Let $h = y - b$. As $h \to 0$, $y \to b$. Then by substituting, we get $f_y(a,b) = \lim_{y \to b} \dfrac{f(a,y) - f(a,b)}{y - b}$.

2. False. If there were such a function, then $f_{xy} = 2y$ and $f_{yx} = 1$. So $f_{xy} \neq f_{yx}$, which contradicts Clairaut's Theorem.

3. False. $f_{xy} = \dfrac{\partial^2 f}{\partial y \, \partial x}$.

4. True. From Equation 14.6.14 we get $D_{\mathbf{k}} f(x,y,z) = \nabla f(x,y,z) \cdot \langle 0, 0, 1 \rangle = f_z(x,y,z)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. False. See Example 14.2.3.

6. False. See Exercise 14.4.46(a).

7. True. If f has a local minimum and f is differentiable at (a, b) then by Theorem 14.7.2, $f_x(a, b) = 0$ and $f_y(a, b) = 0$, so $\nabla f(a, b) = \langle f_x(a, b), f_y(a, b) \rangle = \langle 0, 0 \rangle = \mathbf{0}$.

8. False. If f is not continuous at $(2, 5)$, then we can have $\lim_{(x,y)\to(2,5)} f(x, y) \neq f(2, 5)$. (See Example 14.2.7)

9. False. $\nabla f(x, y) = \langle 0, 1/y \rangle$.

10. True. This is part (c) of the Second Derivatives Test (14.7.3).

11. True. $\nabla f = \langle \cos x, \cos y \rangle$, so $|\nabla f| = \sqrt{\cos^2 x + \cos^2 y}$. But $|\cos \theta| \leq 1$, so $|\nabla f| \leq \sqrt{2}$. Now $D_{\mathbf{u}} f(x, y) = \nabla f \cdot \mathbf{u} = |\nabla f|\,|\mathbf{u}| \cos \theta$, but $\mathbf{u}$ is a unit vector, so $|D_{\mathbf{u}} f(x, y)| \leq \sqrt{2} \cdot 1 \cdot 1 = \sqrt{2}$.

12. False. See Exercise 14.7.39.

EXERCISES

1. $\ln(x + y + 1)$ is defined only when $x + y + 1 > 0 \quad \Leftrightarrow \quad y > -x - 1$, so the domain of f is $\{(x, y) \mid y > -x - 1\}$, all those points above the line $y = -x - 1$.



2. $\sqrt{4 - x^2 - y^2}$ is defined only when $4 - x^2 - y^2 \geq 0 \quad \Leftrightarrow \quad x^2 + y^2 \leq 4$, and $\sqrt{1 - x^2}$ is defined only when $1 - x^2 \geq 0 \quad \Leftrightarrow \quad -1 \leq x \leq 1$, so the domain of f is $\left\{(x, y) \mid -1 \leq x \leq 1, -\sqrt{4 - x^2} \leq y \leq \sqrt{4 - x^2}\right\}$, which consists of those points on or inside the circle $x^2 + y^2 = 4$ for $-1 \leq x \leq 1$.



3. $z = f(x, y) = 1 - y^2$, a parabolic cylinder



4. $z = f(x, y) = x^2 + (y - 2)^2$, a circular paraboloid with vertex $(0, 2, 0)$ and axis parallel to the z-axis



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. The level curves are $\sqrt{4x^2 + y^2} = k$ or $4x^2 + y^2 = k^2$, $k \geq 0$, a family of ellipses.



6. The level curves are $e^x + y = k$ or $y = -e^x + k$, a family of exponential curves.



7.



8. (a) The point $(3, 2)$ lies partway between the level curves with z-values 50 and 60, and it appears that $(3, 2)$ is about the same distance from either level curve. So we estimate that $f(3, 2) \approx 55$.

(b) At the point $(3, 2)$, if we fix y at $y = 2$ and allow x to vary, the level curves indicate that the z-values decrease as x increases, so $f_x(3, 2)$ is negative. In other words, if we start at $(3, 2)$ and move right (in the positive x-direction), the contours show that our path along the surface $z = f(x, y)$ is descending.

(c) Both $f_y(2, 1)$ and $f_y(2, 2)$ are positive, because if we start from either point and move in the positive y-direction, the contour map indicates that the path is ascending. But the level curves are closer together in the y-direction at $(2, 1)$ than at $(2, 2)$, so the path is steeper (the z-values increase more rapidly) at $(2, 1)$ and hence $f_y(2, 1) > f_y(2, 2)$.

9. f is a rational function, so it is continuous on its domain. Since f is defined at $(1, 1)$, we use direct substitution to evaluate the limit: $\lim\limits_{(x,y)\to(1,1)} \dfrac{2xy}{x^2 + 2y^2} = \dfrac{2(1)(1)}{1^2 + 2(1)^2} = \dfrac{2}{3}$.

10. As $(x, y) \to (0, 0)$ along the x-axis, $f(x, 0) = 0/x^2 = 0$ for $x \neq 0$, so $f(x, y) \to 0$ along this line. But $f(x, x) = 2x^2/(3x^2) = \frac{2}{3}$, so as $(x, y) \to (0, 0)$ along the line $x = y$, $f(x, y) \to \frac{2}{3}$. Thus the limit doesn't exist.

11. (a) $T_x(6, 4) = \lim\limits_{h\to 0} \dfrac{T(6 + h, 4) - T(6, 4)}{h}$, so we can approximate $T_x(6, 4)$ by considering $h = \pm 2$ and using the values given in the table: $T_x(6, 4) \approx \dfrac{T(8, 4) - T(6, 4)}{2} = \dfrac{86 - 80}{2} = 3$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$T_x(6,4) \approx \dfrac{T(4,4)-T(6,4)}{-2} = \dfrac{72-80}{-2} = 4$. Averaging these values, we estimate $T_x(6,4)$ to be approximately $3.5°\text{C/m}$. Similarly, $T_y(6,4) = \lim\limits_{h\to 0} \dfrac{T(6,4+h)-T(6,4)}{h}$, which we can approximate with $h = \pm 2$:

$T_y(6,4) \approx \dfrac{T(6,6)-T(6,4)}{2} = \dfrac{75-80}{2} = -2.5$, $T_y(6,4) \approx \dfrac{T(6,2)-T(6,4)}{-2} = \dfrac{87-80}{-2} = -3.5$. Averaging these values, we estimate $T_y(6,4)$ to be approximately $-3.0°\text{C/m}$.

(b) Here $\mathbf{u} = \left\langle \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right\rangle$, so by Equation 14.6.9, $D_{\mathbf{u}}T(6,4) = \nabla T(6,4)\cdot\mathbf{u} = T_x(6,4)\frac{1}{\sqrt{2}} + T_y(6,4)\frac{1}{\sqrt{2}}$. Using our estimates from part (a), we have $D_{\mathbf{u}}T(6,4) \approx (3.5)\frac{1}{\sqrt{2}} + (-3.0)\frac{1}{\sqrt{2}} = \frac{1}{2\sqrt{2}} \approx 0.35$. This means that as we move through the point $(6,4)$ in the direction of $\mathbf{u}$, the temperature increases at a rate of approximately $0.35°\text{C/m}$.

Alternatively, we can use Definition 14.6.2: $D_{\mathbf{u}}T(6,4) = \lim\limits_{h\to 0} \dfrac{T\left(6+h\frac{1}{\sqrt{2}}, 4+h\frac{1}{\sqrt{2}}\right) - T(6,4)}{h}$,

which we can estimate with $h = \pm 2\sqrt{2}$. Then $D_{\mathbf{u}}T(6,4) \approx \dfrac{T(8,6)-T(6,4)}{2\sqrt{2}} = \dfrac{80-80}{2\sqrt{2}} = 0$,

$D_{\mathbf{u}}T(6,4) \approx \dfrac{T(4,2)-T(6,4)}{-2\sqrt{2}} = \dfrac{74-80}{-2\sqrt{2}} = \dfrac{3}{\sqrt{2}}$. Averaging these values, we have $D_{\mathbf{u}}T(6,4) \approx \frac{3}{2\sqrt{2}} \approx 1.1°\text{C/m}$.

(c) $T_{xy}(x,y) = \dfrac{\partial}{\partial y}[T_x(x,y)] = \lim\limits_{h\to 0} \dfrac{T_x(x,y+h)-T_x(x,y)}{h}$, so $T_{xy}(6,4) = \lim\limits_{h\to 0} \dfrac{T_x(6,4+h)-T_x(6,4)}{h}$ which we can estimate with $h = \pm 2$. We have $T_x(6,4) \approx 3.5$ from part (a), but we will also need values for $T_x(6,6)$ and $T_x(6,2)$. If we use $h = \pm 2$ and the values given in the table, we have

$T_x(6,6) \approx \dfrac{T(8,6)-T(6,6)}{2} = \dfrac{80-75}{2} = 2.5$, $T_x(6,6) \approx \dfrac{T(4,6)-T(6,6)}{-2} = \dfrac{68-75}{-2} = 3.5$.

Averaging these values, we estimate $T_x(6,6) \approx 3.0$. Similarly,

$T_x(6,2) \approx \dfrac{T(8,2)-T_x(6,2)}{2} = \dfrac{90-87}{2} = 1.5$, $T_x(6,2) \approx \dfrac{T(4,2)-T(6,2)}{-2} = \dfrac{74-87}{-2} = 6.5$.

Averaging these values, we estimate $T_x(6,2) \approx 4.0$. Finally, we estimate $T_{xy}(6,4)$:

$T_{xy}(6,4) \approx \dfrac{T_x(6,6)-T_x(6,4)}{2} = \dfrac{3.0-3.5}{2} = -0.25$, $T_{xy}(6,4) \approx \dfrac{T_x(6,2)-T_x(6,4)}{-2} = \dfrac{4.0-3.5}{-2} = -0.25$.

Averaging these values, we have $T_{xy}(6,4) \approx -0.25$.

12. From the table, $T(6,4) = 80$, and from Exercise 11 we estimated $T_x(6,4) \approx 3.5$ and $T_y(6,4) \approx -3.0$. The linear approximation then is

$$T(x,y) \approx T(6,4) + T_x(6,4)(x-6) + T_y(6,4)(y-4) \approx 80 + 3.5(x-6) - 3(y-4) = 3.5x - 3y + 71$$

Thus at the point $(5, 3.8)$, we can use the linear approximation to estimate $T(5,3.8) \approx 3.5(5) - 3(3.8) + 71 \approx 77.1°\text{C}$.

13. $f(x,y) = (5y^3+2x^2y)^8 \quad \Rightarrow \quad f_x = 8(5y^3+2x^2y)^7(4xy) = 32xy(5y^3+2x^2y)^7$,

$f_y = 8(5y^3+2x^2y)^7(15y^2+2x^2) = (16x^2+120y^2)(5y^3+2x^2y)^7$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

14. $g(u,v)=\dfrac{u+2v}{u^2+v^2} \quad \Rightarrow \quad g_u=\dfrac{(u^2+v^2)(1)-(u+2v)(2u)}{(u^2+v^2)^2}=\dfrac{v^2-u^2-4uv}{(u^2+v^2)^2}$,

$g_v=\dfrac{(u^2+v^2)(2)-(u+2v)(2v)}{(u^2+v^2)^2}=\dfrac{2u^2-2v^2-2uv}{(u^2+v^2)^2}$

15. $F(\alpha,\beta)=\alpha^2\ln(\alpha^2+\beta^2) \quad \Rightarrow \quad F_\alpha=\alpha^2\cdot\dfrac{1}{\alpha^2+\beta^2}\,(2\alpha)+\ln(\alpha^2+\beta^2)\cdot 2\alpha=\dfrac{2\alpha^3}{\alpha^2+\beta^2}+2\alpha\ln(\alpha^2+\beta^2)$,

$F_\beta=\alpha^2\cdot\dfrac{1}{\alpha^2+\beta^2}\,(2\beta)=\dfrac{2\alpha^2\beta}{\alpha^2+\beta^2}$

16. $G(x,y,z)=e^{xz}\sin(y/z) \quad \Rightarrow \quad G_x=ze^{xz}\sin(y/z),\ G_y=e^{xz}\cos(y/z)(1/z)=(e^{xz}/z)\cos(y/z)$,

$G_z=e^{xz}\cdot\cos(y/z)(-y/z^2)+\sin(y/z)\cdot xe^{xz}=e^{xz}\left[x\sin(y/z)-(y/z^2)\cos(y/z)\right]$

17. $S(u,v,w)=u\arctan(v\sqrt{w}) \quad \Rightarrow \quad S_u=\arctan(v\sqrt{w}),\ S_v=u\cdot\dfrac{1}{1+(v\sqrt{w})^2}\,(\sqrt{w})=\dfrac{u\sqrt{w}}{1+v^2w}$,

$S_w=u\cdot\dfrac{1}{1+(v\sqrt{w})^2}\left(v\cdot\tfrac{1}{2}w^{-1/2}\right)=\dfrac{uv}{2\sqrt{w}\,(1+v^2w)}$

18. $C=1449.2+4.6T-0.055T^2+0.00029T^3+(1.34-0.01T)(S-35)+0.016D \quad \Rightarrow$

$\partial C/\partial T=4.6-0.11T+0.00087T^2-0.01(S-35)$, $\partial C/\partial S=1.34-0.01T$, and $\partial C/\partial D=0.016$. When $T=10$, $S=35$, and $D=100$ we have $\partial C/\partial T=4.6-0.11(10)+0.00087(10)^2-0.01(35-35)\approx 3.587$, thus in 10°C water with salinity 35 parts per thousand and a depth of 100 m, the speed of sound increases by about 3.59 m/s for every degree Celsius that the water temperature rises. Similarly, $\partial C/\partial S=1.34-0.01(10)=1.24$, so the speed of sound increases by about 1.24 m/s for every part per thousand the salinity of the water increases. $\partial C/\partial D=0.016$, so the speed of sound increases by about 0.016 m/s for every meter that the depth is increased.

19. $f(x,y)=4x^3-xy^2 \quad \Rightarrow \quad f_x=12x^2-y^2,\ f_y=-2xy,\ f_{xx}=24x,\ f_{yy}=-2x,\ f_{xy}=f_{yx}=-2y$

20. $z=xe^{-2y} \quad \Rightarrow \quad z_x=e^{-2y},\ z_y=-2xe^{-2y},\ z_{xx}=0,\ z_{yy}=4xe^{-2y},\ z_{xy}=z_{yx}=-2e^{-2y}$

21. $f(x,y,z)=x^ky^lz^m \quad \Rightarrow \quad f_x=kx^{k-1}y^lz^m,\ f_y=lx^ky^{l-1}z^m,\ f_z=mx^ky^lz^{m-1},\ f_{xx}=k(k-1)x^{k-2}y^lz^m$,

$f_{yy}=l(l-1)x^ky^{l-2}z^m,\ f_{zz}=m(m-1)x^ky^lz^{m-2},\ f_{xy}=f_{yx}=klx^{k-1}y^{l-1}z^m,\ f_{xz}=f_{zx}=kmx^{k-1}y^lz^{m-1}$.

$f_{yz}=f_{zy}=lmx^ky^{l-1}z^{m-1}$

22. $v=r\cos(s+2t) \quad \Rightarrow \quad v_r=\cos(s+2t),\ v_s=-r\sin(s+2t),\ v_t=-2r\sin(s+2t),\ v_{rr}=0,\ v_{ss}=-r\cos(s+2t)$,

$v_{tt}=-4r\cos(s+2t),\ v_{rs}=v_{sr}=-\sin(s+2t),\ v_{rt}=v_{tr}=-2\sin(s+2t),\ v_{st}=v_{ts}=-2r\cos(s+2t)$

23. $z=xy+xe^{y/x} \quad \Rightarrow \quad \dfrac{\partial z}{\partial x}=y-\dfrac{y}{x}e^{y/x}+e^{y/x},\ \dfrac{\partial z}{\partial y}=x+e^{y/x}$ and

$$x\,\frac{\partial z}{\partial x}+y\,\frac{\partial z}{\partial y}=x\Big(y-\frac{y}{x}e^{y/x}+e^{y/x}\Big)+y\Big(x+e^{y/x}\Big)=xy-ye^{y/x}+xe^{y/x}+xy+ye^{y/x}=xy+xy+xe^{y/x}=xy+z.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $z = \sin(x + \sin t) \Rightarrow \dfrac{\partial z}{\partial x} = \cos(x + \sin t),\ \dfrac{\partial z}{\partial t} = \cos(x + \sin t)\cos t,$

$\dfrac{\partial^2 z}{\partial x \partial t} = -\sin(x + \sin t)\cos t,\ \dfrac{\partial^2 z}{\partial x^2} = -\sin(x + \sin t)$ and

$\dfrac{\partial z}{\partial x}\dfrac{\partial^2 z}{\partial x \partial t} = \cos(x + \sin t)\,[-\sin(x + \sin t)\cos t] = \cos(x + \sin t)(\cos t)\,[-\sin(x + \sin t)] = \dfrac{\partial z}{\partial t}\dfrac{\partial^2 z}{\partial x^2}.$

25. (a) $z_x = 6x + 2 \Rightarrow z_x(1, -2) = 8$ and $z_y = -2y \Rightarrow z_y(1, -2) = 4$, so an equation of the tangent plane is $z - 1 = 8(x - 1) + 4(y + 2)$ or $z = 8x + 4y + 1$.

(b) A normal vector to the tangent plane (and the surface) at $(1, -2, 1)$ is $\langle 8, 4, -1\rangle$. Then parametric equations for the normal line there are $x = 1 + 8t$, $y = -2 + 4t$, $z = 1 - t$, and symmetric equations are $\dfrac{x - 1}{8} = \dfrac{y + 2}{4} = \dfrac{z - 1}{-1}$.

26. (a) $z_x = e^x \cos y \Rightarrow z_x(0, 0) = 1$ and $z_y = -e^x \sin y \Rightarrow z_y(0, 0) = 0$, so an equation of the tangent plane is $z - 1 = 1(x - 0) + 0(y - 0)$ or $z = x + 1$.

(b) A normal vector to the tangent plane (and the surface) at $(0, 0, 1)$ is $\langle 1, 0, -1\rangle$. Then parametric equations for the normal line there are $x = t$, $y = 0$, $z = 1 - t$, and symmetric equations are $x = 1 - z$, $y = 0$.

27. (a) Let $F(x, y, z) = x^2 + 2y^2 - 3z^2$. Then $F_x = 2x$, $F_y = 4y$, $F_z = -6z$, so $F_x(2, -1, 1) = 4$, $F_y(2, -1, 1) = -4$, $F_z(2, -1, 1) = -6$. From Equation 14.6.19, an equation of the tangent plane is $4(x - 2) - 4(y + 1) - 6(z - 1) = 0$ or, equivalently, $2x - 2y - 3z = 3$.

(b) From Equations 14.6.20, symmetric equations for the normal line are $\dfrac{x - 2}{4} = \dfrac{y + 1}{-4} = \dfrac{z - 1}{-6}$.

28. (a) Let $F(x, y, z) = xy + yz + zx$. Then $F_x = y + z$, $F_y = x + z$, $F_z = x + y$, so $F_x(1, 1, 1) = F_y(1, 1, 1) = F_z(1, 1, 1) = 2$. From Equation 14.6.19, an equation of the tangent plane is $2(x - 1) + 2(y - 1) + 2(z - 1) = 0$ or, equivalently, $x + y + z = 3$.

(b) From Equations 14.6.20, symmetric equations for the normal line are $\dfrac{x - 1}{2} = \dfrac{y - 1}{2} = \dfrac{z - 1}{2}$ or, equivalently, $x = y = z$.

29. (a) Let $F(x, y, z) = x + 2y + 3z - \sin(xyz)$. Then $F_x = 1 - yz\cos(xyz)$, $F_y = 2 - xz\cos(xyz)$, $F_z = 3 - xy\cos(xyz)$, so $F_x(2, -1, 0) = 1$, $F_y(2, -1, 0) = 2$, $F_z(2, -1, 0) = 5$. From Equation 14.6.19, an equation of the tangent plane is $1(x - 2) + 2(y + 1) + 5(z - 0) = 0$ or $x + 2y + 5z = 0$.

(b) From Equations 14.6.20, symmetric equations for the normal line are $\dfrac{x - 2}{1} = \dfrac{y + 1}{2} = \dfrac{z}{5}$ or $x - 2 = \dfrac{y + 1}{2} = \dfrac{z}{5}$.

Parametric equations are $x = 2 + t$, $y = -1 + 2t$, $z = 5t$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

30. Let $f(x,y) = x^2 + y^4$. Then $f_x(x,y) = 2x$ and $f_y(x,y) = 4y^3$, so $f_x(1,1) = 2$, $f_y(1,1) = 4$ and an equation of the tangent plane is $z - 2 = 2(x-1) + 4(y-1)$ or $2x + 4y - z = 4$. A normal vector to the tangent plane is $\langle 2, 4, -1\rangle$ so the normal line is given by $\dfrac{x-1}{2} = \dfrac{y-1}{4} = \dfrac{z-2}{-1}$ or $x = 1 + 2t,\ y = 1 + 4t,$ $z = 2 - t$.



31. The hyperboloid is a level surface of the function $F(x,y,z) = x^2 + 4y^2 - z^2$, so a normal vector to the surface at (x_0, y_0, z_0) is $\nabla F(x_0, y_0, z_0) = \langle 2x_0, 8y_0, -2z_0\rangle$. A normal vector for the plane $2x + 2y + z = 5$ is $\langle 2, 2, 1\rangle$. For the planes to be parallel, we need the normal vectors to be parallel, so $\langle 2x_0, 8y_0, -2z_0\rangle = k\,\langle 2, 2, 1\rangle$, or $x_0 = k$, $y_0 = \frac{1}{4}k$, and $z_0 = -\frac{1}{2}k$. But $x_0^2 + 4y_0^2 - z_0^2 = 4 \quad\Rightarrow\quad k^2 + \frac{1}{4}k^2 - \frac{1}{4}k^2 = 4 \quad\Rightarrow\quad k^2 = 4 \quad\Rightarrow\quad k = \pm 2$. So there are two such points: $\left(2, \frac{1}{2}, -1\right)$ and $\left(-2, -\frac{1}{2}, 1\right)$.

32. $u = \ln(1 + se^{2t}) \quad\Rightarrow\quad du = \dfrac{\partial u}{\partial s}\,ds + \dfrac{\partial u}{\partial t}\,dt = \dfrac{e^{2t}}{1 + se^{2t}}\,ds + \dfrac{2se^{2t}}{1 + se^{2t}}\,dt$

33. $f(x,y,z) = x^3\sqrt{y^2 + z^2} \quad\Rightarrow\quad f_x(x,y,z) = 3x^2\sqrt{y^2 + z^2},\ f_y(x,y,z) = \dfrac{yx^3}{\sqrt{y^2 + z^2}},\ f_z(x,y,z) = \dfrac{zx^3}{\sqrt{y^2 + z^2}},$

so $f(2,3,4) = 8(5) = 40$, $f_x(2,3,4) = 3(4)\sqrt{25} = 60$, $f_y(2,3,4) = \frac{3(8)}{\sqrt{25}} = \frac{24}{5}$, and $f_z(2,3,4) = \frac{4(8)}{\sqrt{25}} = \frac{32}{5}$. Then the linear approximation of f at $(2,3,4)$ is

$$\begin{aligned} f(x,y,z) &\approx f(2,3,4) + f_x(2,3,4)(x-2) + f_y(2,3,4)(y-3) + f_z(2,3,4)(z-4) \\ &= 40 + 60(x-2) + \tfrac{24}{5}(y-3) + \tfrac{32}{5}(z-4) = 60x + \tfrac{24}{5}y + \tfrac{32}{5}z - 120 \end{aligned}$$

Then $(1.98)^3\sqrt{(3.01)^2 + (3.97)^2} = f(1.98, 3.01, 3.97) \approx 60(1.98) + \frac{24}{5}(3.01) + \frac{32}{5}(3.97) - 120 = 38.656$.

34. (a) $dA = \dfrac{\partial A}{\partial x}\,dx + \dfrac{\partial A}{\partial y}\,dy = \frac{1}{2}y\,dx + \frac{1}{2}x\,dy$ and $|\Delta x| \le 0.002$, $|\Delta y| \le 0.002$. Thus the maximum error in the calculated area is about $dA = 6(0.002) + \frac{5}{2}(0.002) = 0.017$ m^2 or 170 cm^2.

(b) $z = \sqrt{x^2 + y^2}$, $dz = \dfrac{x}{\sqrt{x^2 + y^2}}\,dx + \dfrac{y}{\sqrt{x^2 + y^2}}\,dy$ and $|\Delta x| \le 0.002$, $|\Delta y| \le 0.002$. Thus the maximum error in the calculated hypotenuse length is about $dz = \frac{5}{13}(0.002) + \frac{12}{13}(0.002) = \frac{0.17}{65} \approx 0.0026$ m or 0.26 cm.

35. $\dfrac{du}{dp} = \dfrac{\partial u}{\partial x}\dfrac{dx}{dp} + \dfrac{\partial u}{\partial y}\dfrac{dy}{dp} + \dfrac{\partial u}{\partial z}\dfrac{dz}{dp} = 2xy^3(1 + 6p) + 3x^2y^2(pe^p + e^p) + 4z^3(p\cos p + \sin p)$

36. $\dfrac{\partial v}{\partial s} = \dfrac{\partial v}{\partial x}\dfrac{\partial x}{\partial s} + \dfrac{\partial v}{\partial y}\dfrac{\partial y}{\partial s} = \left(2x\sin y + y^2e^{xy}\right)(1) + \left(x^2\cos y + xye^{xy} + e^{xy}\right)(t).$

$s = 0, t = 1 \quad\Rightarrow\quad x = 2, y = 0$, so $\dfrac{\partial v}{\partial s} = 0 + (4 + 1)(1) = 5.$

$\dfrac{\partial v}{\partial t} = \dfrac{\partial v}{\partial x}\dfrac{\partial x}{\partial t} + \dfrac{\partial v}{\partial y}\dfrac{\partial y}{\partial t} = \left(2x\sin y + y^2e^{xy}\right)(2) + \left(x^2\cos y + xye^{xy} + e^{xy}\right)(s) = 0 + 0 = 0.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. By the Chain Rule, $\dfrac{\partial z}{\partial s} = \dfrac{\partial z}{\partial x}\dfrac{\partial x}{\partial s} + \dfrac{\partial z}{\partial y}\dfrac{\partial y}{\partial s}$. When $s = 1$ and $t = 2$, $x = g(1,2) = 3$ and $y = h(1,2) = 6$, so

$\dfrac{\partial z}{\partial s} = f_x(3,6)g_s(1,2) + f_y(3,6)\,h_s(1,2) = (7)(-1) + (8)(-5) = -47$. Similarly, $\dfrac{\partial z}{\partial t} = \dfrac{\partial z}{\partial x}\dfrac{\partial x}{\partial t} + \dfrac{\partial z}{\partial y}\dfrac{\partial y}{\partial t}$, so

$\dfrac{\partial z}{\partial t} = f_x(3,6)g_t(1,2) + f_y(3,6)\,h_t(1,2) = (7)(4) + (8)(10) = 108$.

38.

Using the tree diagram as a guide, we have

$$\frac{\partial w}{\partial p} = \frac{\partial w}{\partial t}\frac{\partial t}{\partial p} + \frac{\partial w}{\partial u}\frac{\partial u}{\partial p} + \frac{\partial w}{\partial v}\frac{\partial v}{\partial p} \qquad \frac{\partial w}{\partial q} = \frac{\partial w}{\partial t}\frac{\partial t}{\partial q} + \frac{\partial w}{\partial u}\frac{\partial u}{\partial q} + \frac{\partial w}{\partial v}\frac{\partial v}{\partial q}$$

$$\frac{\partial w}{\partial r} = \frac{\partial w}{\partial t}\frac{\partial t}{\partial r} + \frac{\partial w}{\partial u}\frac{\partial u}{\partial r} + \frac{\partial w}{\partial v}\frac{\partial v}{\partial r} \qquad \frac{\partial w}{\partial s} = \frac{\partial w}{\partial t}\frac{\partial t}{\partial s} + \frac{\partial w}{\partial u}\frac{\partial u}{\partial s} + \frac{\partial w}{\partial v}\frac{\partial v}{\partial s}$$

39. $\dfrac{\partial z}{\partial x} = 2xf'(x^2 - y^2)$, $\dfrac{\partial z}{\partial y} = 1 - 2yf'(x^2 - y^2)$ $\left[\text{where } f' = \dfrac{df}{d(x^2 - y^2)}\right]$. Then

$y\dfrac{\partial z}{\partial x} + x\dfrac{\partial z}{\partial y} = 2xyf'(x^2 - y^2) + x - 2xyf'(x^2 - y^2) = x$.

40. $A = \frac{1}{2}xy\sin\theta$, $dx/dt = 3$, $dy/dt = -2$, $d\theta/dt = 0.05$, and $\dfrac{dA}{dt} = \dfrac{1}{2}\left[(y\sin\theta)\dfrac{dx}{dt} + (x\sin\theta)\dfrac{dy}{dt} + (xy\cos\theta)\dfrac{d\theta}{dt}\right]$.

So when $x = 40$, $y = 50$ and $\theta = \frac{\pi}{6}$, $\dfrac{dA}{dt} = \dfrac{1}{2}\left[(25)(3) + (20)(-2) + \left(1000\sqrt{3}\right)(0.05)\right] = \dfrac{35 + 50\sqrt{3}}{2} \approx 60.8$ in^2/s.

41. $\dfrac{\partial z}{\partial x} = \dfrac{\partial z}{\partial u}\,y + \dfrac{\partial z}{\partial v}\dfrac{-y}{x^2}$ and

$$\frac{\partial^2 z}{\partial x^2} = y\frac{\partial}{\partial x}\left(\frac{\partial z}{\partial u}\right) + \frac{2y}{x^3}\frac{\partial z}{\partial v} + \frac{-y}{x^2}\frac{\partial}{\partial x}\left(\frac{\partial z}{\partial v}\right) = \frac{2y}{x^3}\frac{\partial z}{\partial v} + y\left(\frac{\partial^2 z}{\partial u^2}\,y + \frac{\partial^2 z}{\partial v\,\partial u}\frac{-y}{x^2}\right) + \frac{-y}{x^2}\left(\frac{\partial^2 z}{\partial v^2}\frac{-y}{x^2} + \frac{\partial^2 z}{\partial u\,\partial v}\,y\right)$$

$$= \frac{2y}{x^3}\frac{\partial z}{\partial v} + y^2\frac{\partial^2 z}{\partial u^2} - \frac{2y^2}{x^2}\frac{\partial^2 z}{\partial u\,\partial v} + \frac{y^2}{x^4}\frac{\partial^2 z}{\partial v^2}$$

Also $\dfrac{\partial z}{\partial y} = x\dfrac{\partial z}{\partial u} + \dfrac{1}{x}\dfrac{\partial z}{\partial v}$ and

$$\frac{\partial^2 z}{\partial y^2} = x\frac{\partial}{\partial y}\left(\frac{\partial z}{\partial u}\right) + \frac{1}{x}\frac{\partial}{\partial y}\left(\frac{\partial z}{\partial v}\right) = x\left(\frac{\partial^2 z}{\partial u^2}x + \frac{\partial^2 z}{\partial v\,\partial u}\frac{1}{x}\right) + \frac{1}{x}\left(\frac{\partial^2 z}{\partial v^2}\frac{1}{x} + \frac{\partial^2 z}{\partial u\,\partial v}\,x\right) = x^2\frac{\partial^2 z}{\partial u^2} + 2\frac{\partial^2 z}{\partial u\,\partial v} + \frac{1}{x^2}\frac{\partial^2 z}{\partial v^2}$$

Thus

$$x^2\frac{\partial^2 z}{\partial x^2} - y^2\frac{\partial^2 z}{\partial y^2} = \frac{2y}{x}\frac{\partial z}{\partial v} + x^2y^2\frac{\partial^2 z}{\partial u^2} - 2y^2\frac{\partial^2 z}{\partial u\,\partial v} + \frac{y^2}{x^2}\frac{\partial^2 z}{\partial v^2} - x^2y^2\frac{\partial^2 z}{\partial u^2} - 2y^2\frac{\partial^2 z}{\partial u\,\partial v} - \frac{y^2}{x^2}\frac{\partial^2 z}{\partial v^2}$$

$$= \frac{2y}{x}\frac{\partial z}{\partial v} - 4y^2\frac{\partial^2 z}{\partial u\,\partial v} = 2v\frac{\partial z}{\partial v} - 4uv\frac{\partial^2 z}{\partial u\,\partial v}$$

since $y = xv = \dfrac{uv}{y}$ or $y^2 = uv$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

42. $\cos(xyz) = 1 + x^2y^2 + z^2$, so let $F(x, y, z) = 1 + x^2y^2 + z^2 - \cos(xyz) = 0$. Then by Equations 14.5.7 we have $\dfrac{\partial z}{\partial x} = -\dfrac{F_x}{F_z} = -\dfrac{2xy^2 + \sin(xyz) \cdot yz}{2z + \sin(xyz) \cdot xy} = -\dfrac{2xy^2 + yz\sin(xyz)}{2z + xy\sin(xyz)}$,

$\dfrac{\partial z}{\partial y} = -\dfrac{F_y}{F_z} = -\dfrac{2x^2y + \sin(xyz) \cdot xz}{2z + \sin(xyz) \cdot xy} = -\dfrac{2x^2y + xz\sin(xyz)}{2z + xy\sin(xyz)}$.

43. $f(x, y, z) = x^2e^{yz^2} \quad \Rightarrow \quad \nabla f = \langle f_x, f_y, f_z \rangle = \left\langle 2xe^{yz^2}, x^2e^{yz^2} \cdot z^2, x^2e^{yz^2} \cdot 2yz \right\rangle = \left\langle 2xe^{yz^2}, x^2z^2e^{yz^2}, 2x^2yze^{yz^2} \right\rangle$

44. (a) By Theorem 14.6.15, the maximum value of the directional derivative occurs when $\mathbf{u}$ has the same direction as the gradient vector.

(b) It is a minimum when $\mathbf{u}$ is in the direction opposite to that of the gradient vector (that is, $\mathbf{u}$ is in the direction of $-\nabla f$), since $D_{\mathbf{u}} f = |\nabla f| \cos\theta$ (see the proof of Theorem 14.6.15) has a minimum when $\theta = \pi$.

(c) The directional derivative is 0 when $\mathbf{u}$ is perpendicular to the gradient vector, since then $D_{\mathbf{u}} f = \nabla f \cdot \mathbf{u} = 0$.

(d) The directional derivative is half of its maximum value when $D_{\mathbf{u}} f = |\nabla f| \cos\theta = \frac{1}{2} |\nabla f| \quad \Leftrightarrow \quad \cos\theta = \frac{1}{2} \quad \Leftrightarrow \quad \theta = \frac{\pi}{3}$.

45. $f(x, y) = x^2e^{-y} \quad \Rightarrow \quad \nabla f = \langle 2xe^{-y}, -x^2e^{-y} \rangle$, $\nabla f(-2, 0) = \langle -4, -4 \rangle$. The direction is given by $\langle 4, -3 \rangle$, so $\mathbf{u} = \frac{1}{\sqrt{4^2 + (-3)^2}} \langle 4, -3 \rangle = \frac{1}{5} \langle 4, -3 \rangle$ and $D_{\mathbf{u}} f(-2, 0) = \nabla f(-2, 0) \cdot \mathbf{u} = \langle -4, -4 \rangle \cdot \frac{1}{5} \langle 4, -3 \rangle = \frac{1}{5}(-16 + 12) = -\frac{4}{5}$.

46. $\nabla f = \left\langle 2xy + \sqrt{1 + z}, x^2, x/\left(2\sqrt{1 + z}\right) \right\rangle$, $\nabla f(1, 2, 3) = \left\langle 6, 1, \frac{1}{4} \right\rangle$, $\mathbf{u} = \left\langle \frac{2}{3}, \frac{1}{3}, -\frac{2}{3} \right\rangle$. Then $D_{\mathbf{u}} f(1, 2, 3) = \frac{25}{6}$.

47. $\nabla f = \left\langle 2xy, x^2 + 1/\left(2\sqrt{y}\right) \right\rangle$, $|\nabla f(2, 1)| = \left|\left\langle 4, \frac{9}{2} \right\rangle\right|$. Thus the maximum rate of change of f at $(2, 1)$ is $\frac{\sqrt{145}}{2}$ in the direction $\left\langle 4, \frac{9}{2} \right\rangle$.

48. $\nabla f = \langle zye^{xy}, zxe^{xy}, e^{xy} \rangle$, $\nabla f(0, 1, 2) = \langle 2, 0, 1 \rangle$ is the direction of most rapid increase while the rate is $|\langle 2, 0, 1 \rangle| = \sqrt{5}$.

49. First we draw a line passing through Homestead and the eye of the hurricane. We can approximate the directional derivative at Homestead in the direction of the eye of the hurricane by the average rate of change of wind speed between the points where this line intersects the contour lines closest to Homestead. In the direction of the eye of the hurricane, the wind speed changes from 45 to 50 knots. We estimate the distance between these two points to be approximately 8 miles, so the rate of change of wind speed in the direction given is approximately $\frac{50 - 45}{8} = \frac{5}{8} = 0.625$ knot/mi.

50. The surfaces are $f(x, y, z) = z - 2x^2 + y^2 = 0$ and $g(x, y, z) = z - 4 = 0$. The tangent line is perpendicular to both ∇f and ∇g at $(-2, 2, 4)$. The vector $\mathbf{v} = \nabla f \times \nabla g$ is therefore parallel to the line. $\nabla f(x, y, z) = \langle -4x, 2y, 1 \rangle \quad \Rightarrow$ $\nabla f(-2, 2, 4) = \langle 8, 4, 1 \rangle$, $\nabla g(x, y, z) = \langle 0, 0, 1 \rangle \quad \Rightarrow \quad \nabla g\langle -2, 2, 4 \rangle = \langle 0, 0, 1 \rangle$. Hence

$\mathbf{v} = \nabla f \times \nabla g = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 8 & 4 & 1 \\ 0 & 0 & 1 \end{vmatrix} = 4\,\mathbf{i} - 8\,\mathbf{j}$. Thus, parametric equations are: $x = -2 + 4t$, $y = 2 - 8t$, $z = 4$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. $f(x,y) = x^2 - xy + y^2 + 9x - 6y + 10 \Rightarrow f_x = 2x - y + 9$, $f_y = -x + 2y - 6$, $f_{xx} = 2 = f_{yy}$, $f_{xy} = -1$. Then $f_x = 0$ and $f_y = 0$ imply $y = 1$, $x = -4$. Thus the only critical point is $(-4, 1)$ and $f_{xx}(-4,1) > 0$, $D(-4,1) = 3 > 0$, so $f(-4,1) = -11$ is a local minimum.



52. $f(x,y) = x^3 - 6xy + 8y^3 \Rightarrow f_x = 3x^2 - 6y$, $f_y = -6x + 24y^2$, $f_{xx} = 6x$, $f_{yy} = 48y$, $f_{xy} = -6$. Then $f_x = 0$ implies $y = x^2/2$, substituting into $f_y = 0$ implies $6x(x^3 - 1) = 0$, so the critical points are $(0,0)$, $(1, \frac{1}{2})$. $D(0,0) = -36 < 0$ so $(0,0)$ is a saddle point while $f_{xx}(1, \frac{1}{2}) = 6 > 0$ and $D(1, \frac{1}{2}) = 108 > 0$ so $f(1, \frac{1}{2}) = -1$ is a local minimum.



53. $f(x,y) = 3xy - x^2y - xy^2 \Rightarrow f_x = 3y - 2xy - y^2$, $f_y = 3x - x^2 - 2xy$, $f_{xx} = -2y$, $f_{yy} = -2x$, $f_{xy} = 3 - 2x - 2y$. Then $f_x = 0$ implies $y(3 - 2x - y) = 0$ so $y = 0$ or $y = 3 - 2x$. Substituting into $f_y = 0$ implies $x(3 - x) = 0$ or $3x(-1 + x) = 0$. Hence the critical points are $(0,0)$, $(3,0)$, $(0,3)$ and $(1,1)$. $D(0,0) = D(3,0) = D(0,3) = -9 < 0$ so $(0,0)$, $(3,0)$, and $(0,3)$ are saddle points. $D(1,1) = 3 > 0$ and $f_{xx}(1,1) = -2 < 0$, so $f(1,1) = 1$ is a local maximum.



54. $f(x,y) = (x^2 + y)e^{y/2} \Rightarrow f_x = 2xe^{y/2}$, $f_y = e^{y/2}(2 + x^2 + y)/2$, $f_{xx} = 2e^{y/2}$, $f_{yy} = e^{y/2}(4 + x^2 + y)/4$, $f_{xy} = xe^{y/2}$. Then $f_x = 0$ implies $x = 0$, so $f_y = 0$ implies $y = -2$. But $f_{xx}(0,-2) > 0$, $D(0,-2) = e^{-2} - 0 > 0$ so $f(0,-2) = -2/e$ is a local minimum.



55. First solve inside D. Here $f_x = 4y^2 - 2xy^2 - y^3$, $f_y = 8xy - 2x^2y - 3xy^2$. Then $f_x = 0$ implies $y = 0$ or $y = 4 - 2x$, but $y = 0$ isn't inside D. Substituting $y = 4 - 2x$ into $f_y = 0$ implies $x = 0$, $x = 2$ or $x = 1$, but $x = 0$ isn't inside D, and when $x = 2$, $y = 0$ but $(2,0)$ isn't inside D. Thus the only critical point inside D is $(1,2)$ and $f(1,2) = 4$. Secondly we consider the boundary of D. On L_1: $f(x,0) = 0$ and so $f = 0$ on L_1. On L_2: $x = -y + 6$ and $f(-y + 6, y) = y^2(6 - y)(-2) = -2(6y^2 - y^3)$ which has critical points at $y = 0$ and $y = 4$. Then $f(6,0) = 0$ while $f(2,4) = -64$. On L_3: $f(0,y) = 0$, so $f = 0$ on L_3. Thus on D the absolute maximum of f is $f(1,2) = 4$ while the absolute minimum is $f(2,4) = -64$.



56. Inside D: $f_x = 2xe^{-x^2-y^2}(1 - x^2 - 2y^2) = 0$ implies $x = 0$ or $x^2 + 2y^2 = 1$. Then if $x = 0$, $f_y = 2ye^{-x^2-y^2}(2 - x^2 - 2y^2) = 0$ implies $y = 0$ or $2 - 2y^2 = 0$ giving the critical points $(0,0)$, $(0, \pm 1)$. If

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$x^2 + 2y^2 = 1$, then $f_y = 0$ implies $y = 0$ giving the critical points $(\pm 1, 0)$. Now $f(0,0) = 0$, $f(\pm 1, 0) = e^{-1}$ and $f(0, \pm 1) = 2e^{-1}$. On the boundary of D: $x^2 + y^2 = 4$, so $f(x,y) = e^{-4}(4 + y^2)$ and f is smallest when $y = 0$ and largest when $y^2 = 4$. But $f(\pm 2, 0) = 4e^{-4}$, $f(0, \pm 2) = 8e^{-4}$. Thus on D the absolute maximum of f is $f(0, \pm 1) = 2e^{-1}$ and the absolute minimum is $f(0,0) = 0$.

57. $f(x,y) = x^3 - 3x + y^4 - 2y^2$







From the graphs, it appears that f has a local maximum $f(-1,0) \approx 2$, local minima $f(1, \pm 1) \approx -3$, and saddle points at $(-1, \pm 1)$ and $(1, 0)$.

To find the exact quantities, we calculate $f_x = 3x^2 - 3 = 0 \Leftrightarrow x = \pm 1$ and $f_y = 4y^3 - 4y = 0 \Leftrightarrow y = 0, \pm 1$, giving the critical points estimated above. Also $f_{xx} = 6x$, $f_{xy} = 0$, $f_{yy} = 12y^2 - 4$, so using the Second Derivatives Test, $D(-1,0) = 24 > 0$ and $f_{xx}(-1,0) = -6 < 0$ indicating a local maximum $f(-1,0) = 2$; $D(1, \pm 1) = 48 > 0$ and $f_{xx}(1, \pm 1) = 6 > 0$ indicating local minima $f(1, \pm 1) = -3$; and $D(-1, \pm 1) = -48$ and $D(1,0) = -24$, indicating saddle points.

58. $f(x,y) = 12 + 10y - 2x^2 - 8xy - y^4 \Rightarrow f_x(x,y) = -4x - 8y$, $f_y(x,y) = 10 - 8x - 4y^3$. Now $f_x(x,y) = 0 \Rightarrow x = -2x$, and substituting this into $f_y(x,y) = 0$ gives $10 + 16y - 4y^3 = 0 \Leftrightarrow 5 + 8y - 2y^3 = 0$.





From the first graph, we see that this is true when $y \approx -1.542$, 0.717, or 2.260. (Alternatively, we could have found the solutions to $f_x = f_y = 0$ using a CAS.) So to three decimal places, the critical points are $(3.085, -1.542)$, $(1.434, -0.717)$, and $(-4.519, 2.260)$. Now in order to use the Second Derivatives Test, we calculate $f_{xx} = -4$, $f_{xy} = -8$, $f_{yy} = -12y^2$, and $D = 48y^2 - 64$. So since $D(3.085, -1.542) > 0$, $D(1.434, -0.717) < 0$, and $D(-4.519, 2.260) > 0$, and f_{xx} is always negative, $f(x,y)$ has local maxima $f(-4.519, 2.260) \approx 49.373$ and $f(3.085, -1.542) \approx 9.948$, and a saddle point at approximately $(1.434, -0.717)$. The highest point on the graph is approximately $(-4.519, 2.260, 49.373)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $f(x,y) = x^2y,\ g(x,y) = x^2 + y^2 = 1 \ \Rightarrow\ \nabla f = \langle 2xy, x^2\rangle = \lambda\nabla g = \langle 2\lambda x, 2\lambda y\rangle$. Then $2xy = 2\lambda x$ implies $x = 0$ or $y = \lambda$. If $x = 0$ then $x^2 + y^2 = 1$ gives $y = \pm 1$ and we have possible points $(0, \pm 1)$ where $f(0, \pm 1) = 0$. If $y = \lambda$ then $x^2 = 2\lambda y$ implies $x^2 = 2y^2$ and substitution into $x^2 + y^2 = 1$ gives $3y^2 = 1 \ \Rightarrow\ y = \pm\frac{1}{\sqrt{3}}$ and $x = \pm\sqrt{\frac{2}{3}}$. The corresponding possible points are $\left(\pm\sqrt{\frac{2}{3}}, \pm\frac{1}{\sqrt{3}}\right)$. The absolute maximum is $f\left(\pm\sqrt{\frac{2}{3}}, \frac{1}{\sqrt{3}}\right) = \frac{2}{3\sqrt{3}}$ while the absolute minimum is $f\left(\pm\sqrt{\frac{2}{3}}, -\frac{1}{\sqrt{3}}\right) = -\frac{2}{3\sqrt{3}}$.

60. $f(x,y) = 1/x + 1/y,\ g(x,y) = 1/x^2 + 1/y^2 = 1 \ \Rightarrow\ \nabla f = \langle -x^{-2}, -y^{-2}\rangle = \lambda\nabla g = \langle -2\lambda x^{-3}, -2\lambda y^{-3}\rangle$. Then $-x^{-2} = -2\lambda x^3$ or $x = 2\lambda$ and $-y^{-2} = -2\lambda y^{-3}$ or $y = 2\lambda$. Thus $x = y$, so $1/x^2 + 1/y^2 = 2/x^2 = 1$ implies $x = \pm\sqrt{2}$ and the possible points are $(\pm\sqrt{2}, \pm\sqrt{2})$. The absolute maximum of f subject to $x^{-2} + y^{-2} = 1$ is then $f(\sqrt{2}, \sqrt{2}) = \sqrt{2}$ and the absolute minimum is $f(-\sqrt{2}, -\sqrt{2}) = -\sqrt{2}$.

61. $f(x,y,z) = xyz,\ g(x,y,z) = x^2 + y^2 + z^2 = 3.\ \ \nabla f = \lambda\nabla g \ \Rightarrow\ \langle yz, xz, xy\rangle = \lambda\langle 2x, 2y, 2z\rangle$. If any of x, y, or z is zero, then $x = y = z = 0$ which contradicts $x^2 + y^2 + z^2 = 3$. Then $\lambda = \dfrac{yz}{2x} = \dfrac{xz}{2y} = \dfrac{xy}{2z} \ \Rightarrow\ 2y^2z = 2x^2z \ \Rightarrow$ $y^2 = x^2$, and similarly $2yz^2 = 2x^2y \ \Rightarrow\ z^2 = x^2$. Substituting into the constraint equation gives $x^2 + x^2 + x^2 = 3 \ \Rightarrow$ $x^2 = 1 = y^2 = z^2$. Thus the possible points are $(1, 1, \pm 1)$, $(1, -1, \pm 1)$, $(-1, 1, \pm 1)$, $(-1, -1, \pm 1)$. The absolute maximum is $f(1,1,1) = f(1,-1,-1) = f(-1,1,-1) = f(-1,-1,1) = 1$ and the absolute minimum is $f(1,1,-1) = f(1,-1,1) = f(-1,1,1) = f(-1,-1,-1) = -1$.

62. $f(x,y,z) = x^2 + 2y^2 + 3z^2,\ g(x,y,z) = x + y + z = 1,\ h(x,y,z) = x - y + 2z = 2 \ \Rightarrow$ $\nabla f = \langle 2x, 4y, 6z\rangle = \lambda\nabla g + \mu\nabla h = \langle \lambda + \mu, \lambda - \mu, \lambda + 2\mu\rangle$ and $2x = \lambda + \mu$ **(1)**, $4y = \lambda - \mu$ **(2)**, $6z = \lambda + 2\mu$ **(3)**, $x + y + z = 1$ **(4)**, $x - y + 2z = 2$ **(5)**. Then six times **(1)** plus three times **(2)** plus two times **(3)** implies $12(x + y + z) = 11\lambda + 7\mu$, so **(4)** gives $11\lambda + 7\mu = 12$. Also six times **(1)** minus three times **(2)** plus four times **(3)** implies $12(x - y + 2z) = 7\lambda + 17\mu$, so **(5)** gives $7\lambda + 17\mu = 24$. Solving $11\lambda + 7\mu = 12$, $7\lambda + 17\mu = 24$ simultaneously gives $\lambda = \frac{6}{23}$, $\mu = \frac{30}{23}$. Substituting into **(1)**, **(2)**, and **(3)** implies $x = \frac{18}{23}$, $y = -\frac{6}{23}$, $z = \frac{11}{23}$ giving only one point. Then $f\left(\frac{18}{23}, -\frac{6}{23}, \frac{11}{23}\right) = \frac{33}{23}$. Now since $(0,0,1)$ satisfies both constraints and $f(0,0,1) = 3 > \frac{33}{23}$, $f\left(\frac{18}{23}, -\frac{6}{23}, \frac{11}{23}\right) = \frac{33}{23}$ is an absolute minimum, and there is no absolute maximum.

63. $f(x,y,z) = x^2 + y^2 + z^2,\ g(x,y,z) = xy^2z^3 = 2 \ \Rightarrow\ \nabla f = \langle 2x, 2y, 2z\rangle = \lambda\nabla g = \langle \lambda y^2z^3, 2\lambda xyz^3, 3\lambda xy^2z^2\rangle$. Since $xy^2z^3 = 2$, $x \neq 0$, $y \neq 0$ and $z \neq 0$, so $2x = \lambda y^2z^3$ **(1)**, $1 = \lambda xz^3$ **(2)**, $2 = 3\lambda xy^2z$ **(3)**. Then **(2)** and **(3)** imply $\dfrac{1}{xz^3} = \dfrac{2}{3xy^2z}$ or $y^2 = \frac{2}{3}z^2$ so $y = \pm z\sqrt{\frac{2}{3}}$. Similarly **(1)** and **(3)** imply $\dfrac{2x}{y^2z^3} = \dfrac{2}{3xy^2z}$ or $3x^2 = z^2$ so $x = \pm\frac{1}{\sqrt{3}}z$. But $xy^2z^3 = 2$ so x and z must have the same sign, that is, $x = \frac{1}{\sqrt{3}}z$. Thus $g(x,y,z) = 2$ implies $\frac{1}{\sqrt{3}}z\left(\frac{2}{3}z^2\right)z^3 = 2$ or

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$z = \pm 3^{1/4}$ and the possible points are $(\pm 3^{-1/4}, 3^{-1/4}\sqrt{2}, \pm 3^{1/4})$, $(\pm 3^{-1/4}, -3^{-1/4}\sqrt{2}, \pm 3^{1/4})$. However at each of these points f takes on the same value, $2\sqrt{3}$. But $(2, 1, 1)$ also satisfies $g(x, y, z) = 2$ and $f(2, 1, 1) = 6 > 2\sqrt{3}$. Thus f has an absolute minimum value of $2\sqrt{3}$ and no absolute maximum subject to the constraint $xy^2z^3 = 2$.

Alternate solution: $g(x, y, z) = xy^2z^3 = 2$ implies $y^2 = \dfrac{2}{xz^3}$, so minimize $f(x, z) = x^2 + \dfrac{2}{xz^3} + z^2$. Then $f_x = 2x - \dfrac{2}{x^2z^3}$, $f_z = -\dfrac{6}{xz^4} + 2z$, $f_{xx} = 2 + \dfrac{4}{x^3z^3}$, $f_{zz} = \dfrac{24}{xz^5} + 2$ and $f_{xz} = \dfrac{6}{x^2z^4}$. Now $f_x = 0$ implies $2x^3z^3 - 2 = 0$ or $z = 1/x$. Substituting into $f_y = 0$ implies $-6x^3 + 2x^{-1} = 0$ or $x = \frac{1}{\sqrt[4]{3}}$, so the two critical points are $\left(\pm\frac{1}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right)$. Then $D\left(\pm\frac{1}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right) = (2+4)\left(2+\frac{24}{3}\right) - \left(\frac{6}{\sqrt{3}}\right)^2 > 0$ and $f_{xx}\left(\pm\frac{1}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right) = 6 > 0$, so each point is a minimum. Finally, $y^2 = \dfrac{2}{xz^3}$, so the four points closest to the origin are $\left(\pm\frac{1}{\sqrt[4]{3}}, \frac{\sqrt{2}}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right)$, $\left(\pm\frac{1}{\sqrt[4]{3}}, -\frac{\sqrt{2}}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right)$.

64. $V = xyz$, say x is the length and $x + 2y + 2z \le 108$, $x > 0$, $y > 0$, $z > 0$. First maximize V subject to $x + 2y + 2z = 108$ with x, y, z all positive. Then $\langle yz, xz, xy\rangle = \langle \lambda, 2\lambda, 2\lambda\rangle$ implies $2yz = xz$ or $x = 2y$ and $xz = xy$ or $z = y$. Thus $g(x, y, z) = 108$ implies $6y = 108$ or $y = 18 = z$, $x = 36$, so the volume is $V = 11{,}664$ cubic units. Since $(104, 1, 1)$ also satisfies $g(x, y, z) = 108$ and $V(104, 1, 1) = 104$ cubic units, $(36, 18, 18)$ gives an absolute maximum of V subject to $g(x, y, z) = 108$. But if $x + 2y + 2z < 108$, there exists $\alpha > 0$ such that $x + 2y + 2z = 108 - \alpha$ and as above $6y = 108 - \alpha$ implies $y = (108 - \alpha)/6 = z$, $x = (108 - \alpha)/3$ with $V = (108 - \alpha)^3/(6^2 \cdot 3) < (108)^3/(6^2 \cdot 3) = 11{,}664$. Hence we have shown that the maximum of V subject to $g(x, y, z) \le 108$ is the maximum of V subject to $g(x, y, z) = 108$ (an intuitively obvious fact).

65.



The area of the triangle is $\frac{1}{2}ca\sin\theta$ and the area of the rectangle is bc. Thus, the area of the whole object is $f(a, b, c) = \frac{1}{2}ca\sin\theta + bc$. The perimeter of the object is $g(a, b, c) = 2a + 2b + c = P$. To simplify $\sin\theta$ in terms of a, b, and c notice that $a^2\sin^2\theta + \left(\frac{1}{2}c\right)^2 = a^2 \;\Rightarrow\; \sin\theta = \dfrac{1}{2a}\sqrt{4a^2 - c^2}$. Thus $f(a, b, c) = \dfrac{c}{4}\sqrt{4a^2 - c^2} + bc$. (Instead of using θ, we could just have used the Pythagorean Theorem.) As a result, by Lagrange's method, we must find a, b, c, and λ by solving $\nabla f = \lambda\nabla g$ which gives the following equations: $ca(4a^2 - c^2)^{-1/2} = 2\lambda$ **(1)**, $c = 2\lambda$ **(2)**, $\frac{1}{4}(4a^2 - c^2)^{1/2} - \frac{1}{4}c^2(4a^2 - c^2)^{-1/2} + b = \lambda$ **(3)**, and $2a + 2b + c = P$ **(4)**. From **(2)**, $\lambda = \frac{1}{2}c$ and so **(1)** produces $ca(4a^2 - c^2)^{-1/2} = c \;\Rightarrow\; (4a^2 - c^2)^{1/2} = a \;\Rightarrow\; 4a^2 - c^2 = a^2 \;\Rightarrow\; c = \sqrt{3}\,a$ **(5)**. Similarly, since $(4a^2 - c^2)^{1/2} = a$ and $\lambda = \frac{1}{2}c$, **(3)** gives $\dfrac{a}{4} - \dfrac{c^2}{4a} + b = \dfrac{c}{2}$, so from **(5)**, $\dfrac{a}{4} - \dfrac{3a}{4} + b = \dfrac{\sqrt{3}\,a}{2} \;\Rightarrow\; -\dfrac{a}{2} - \dfrac{\sqrt{3}\,a}{2} = -b \;\Rightarrow\; b = \dfrac{a}{2}\left(1 + \sqrt{3}\right)$ **(6)**. Substituting **(5)** and **(6)** into **(4)** we get:

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$2a + a\left(1+\sqrt{3}\right) + \sqrt{3}\,a = P \;\Rightarrow\; 3a + 2\sqrt{3}\,a = P \;\Rightarrow\; a = \dfrac{P}{3+2\sqrt{3}} = \dfrac{2\sqrt{3}-3}{3}P$ and thus

$b = \dfrac{\left(2\sqrt{3}-3\right)\left(1+\sqrt{3}\right)}{6}P = \dfrac{3-\sqrt{3}}{6}P$ and $c = \left(2-\sqrt{3}\right)P$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

PROBLEMS PLUS

1. The areas of the smaller rectangles are $A_1 = xy$, $A_2 = (L-x)y$, $A_3 = (L-x)(W-y)$, $A_4 = x(W-y)$. For $0 \le x \le L$, $0 \le y \le W$, let



$$\begin{aligned} f(x,y) &= A_1^2 + A_2^2 + A_3^2 + A_4^2 \\ &= x^2y^2 + (L-x)^2y^2 + (L-x)^2(W-y)^2 + x^2(W-y)^2 \\ &= [x^2 + (L-x)^2][y^2 + (W-y)^2] \end{aligned}$$

Then we need to find the maximum and minimum values of $f(x,y)$. Here

$f_x(x,y) = [2x - 2(L-x)][y^2 + (W-y)^2] = 0 \quad \Rightarrow \quad 4x - 2L = 0$ or $x = \frac{1}{2}L$, and

$f_y(x,y) = [x^2 + (L-x)^2][2y - 2(W-y)] = 0 \quad \Rightarrow \quad 4y - 2W = 0$ or $y = W/2$. Also

$f_{xx} = 4[y^2 + (W-y)^2]$, $f_{yy} = 4[x^2 + (L-x)^2]$, and $f_{xy} = (4x - 2L)(4y - 2W)$. Then

$D = 16[y^2 + (W-y)^2][x^2 + (L-x)^2] - (4x - 2L)^2(4y - 2W)^2$. Thus when $x = \frac{1}{2}L$ and $y = \frac{1}{2}W$, $D > 0$ and $f_{xx} = 2W^2 > 0$. Thus a minimum of f occurs at $\left(\frac{1}{2}L, \frac{1}{2}W\right)$ and this minimum value is $f\left(\frac{1}{2}L, \frac{1}{2}W\right) = \frac{1}{4}L^2W^2$.

There are no other critical points, so the maximum must occur on the boundary. Now along the width of the rectangle let $g(y) = f(0,y) = f(L,y) = L^2[y^2 + (W-y)^2]$, $0 \le y \le W$. Then $g'(y) = L^2[2y - 2(W-y)] = 0 \quad \Leftrightarrow \quad y = \frac{1}{2}W$. And $g\left(\frac{1}{2}\right) = \frac{1}{2}L^2W^2$. Checking the endpoints, we get $g(0) = g(W) = L^2W^2$. Along the length of the rectangle let $h(x) = f(x,0) = f(x,W) = W^2[x^2 + (L-x)^2]$, $0 \le x \le L$. By symmetry $h'(x) = 0 \quad \Leftrightarrow \quad x = \frac{1}{2}L$ and $h\left(\frac{1}{2}L\right) = \frac{1}{2}L^2W^2$. At the endpoints we have $h(0) = h(L) = L^2W^2$. Therefore L^2W^2 is the maximum value of f. This maximum value of f occurs when the "cutting" lines correspond to sides of the rectangle.

2. (a) The level curves of the function $C(x,y) = e^{-(x^2+2y^2)/10^4}$ are the curves $e^{-(x^2+2y^2)/10^4} = k$ (k is a positive constant). This equation is equivalent to $x^2 + 2y^2 = K \quad \Rightarrow \quad \dfrac{x^2}{\left(\sqrt{K}\right)^2} + \dfrac{y^2}{\left(\sqrt{K/2}\right)^2} = 1$, where $K = -10^4 \ln k$, a family of ellipses. We sketch level curves for $K = 1$, 2, 3, and 4. If the shark always swims in the direction of maximum increase of blood concentration, its direction at any point would coincide with the gradient vector. Then we know the shark's path is perpendicular to the level curves it intersects. We sketch one example of such a path.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\nabla C = \frac{2}{10^4} e^{-(x^2+2y^2)/10^4}(x\,\mathbf{i} + 2y\,\mathbf{j})$. And ∇C points in the direction of most rapid increase in concentration; that is, ∇C is tangent to the most rapid increase curve. If $r(t) = x(t)\,\mathbf{i} + y(t)\,\mathbf{j}$ is a parametrization of the most rapid increase curve, then $\dfrac{d\mathbf{r}}{dt} = \dfrac{dx}{dt}\,\mathbf{i} + \dfrac{dy}{dt}\,\mathbf{j}$ is tangent to the curve, so $\dfrac{d\mathbf{r}}{dt} = \lambda \nabla C \quad \Rightarrow \quad \dfrac{dx}{dt} = \lambda\left[-\dfrac{2}{10^4}\, e^{-(x^2+2y^2)/10^4}\right] x$ and $\dfrac{dy}{dt} = \lambda\left[-\dfrac{2}{10^4} e^{-(x^2+2y^2)/10^4}\right](2y)$. Therefore $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = 2\,\dfrac{y}{x} \quad \Rightarrow \quad \dfrac{dy}{y} = 2\,\dfrac{dx}{x} \quad \Rightarrow \quad \ln|y| = 2\ln|x|$ so that $y = kx^2$ for some constant k. But $y(x_0) = y_0 \quad \Rightarrow \quad y_0 = kx_0^2 \quad \Rightarrow \quad k = y_0/x_0^2 \quad (x_0 = 0 \quad \Rightarrow \quad y_0 = 0 \quad \Rightarrow$ the shark is already at the origin, so we can assume $x_0 \neq 0$.) Therefore the path the shark will follow is along the parabola $y = y_0(x/x_0)^2$.

3. (a) The area of a trapezoid is $\frac{1}{2}h(b_1 + b_2)$, where h is the height (the distance between the two parallel sides) and b_1, b_2 are the lengths of the bases (the parallel sides). From the figure in the text, we see that $h = x\sin\theta$, $b_1 = w - 2x$, and $b_2 = w - 2x + 2x\cos\theta$. Therefore the cross-sectional area of the rain gutter is

$$\begin{aligned} A(x,\theta) &= \tfrac{1}{2}x\sin\theta\,[(w-2x) + (w-2x+2x\cos\theta)] = (x\sin\theta)(w - 2x + x\cos\theta) \\ &= wx\sin\theta - 2x^2\sin\theta + x^2\sin\theta\cos\theta,\ 0 < x \le \tfrac{1}{2}w,\ 0 < \theta \le \tfrac{\pi}{2} \end{aligned}$$

We look for the critical points of A: $\partial A/\partial x = w\sin\theta - 4x\sin\theta + 2x\sin\theta\cos\theta$ and $\partial A/\partial\theta = wx\cos\theta - 2x^2\cos\theta + x^2(\cos^2\theta - \sin^2\theta)$, so $\partial A/\partial x = 0 \quad \Leftrightarrow \quad \sin\theta\,(w - 4x + 2x\cos\theta) = 0 \quad \Leftrightarrow$ $\cos\theta = \dfrac{4x - w}{2x} = 2 - \dfrac{w}{2x} \quad (0 < \theta \le \frac{\pi}{2} \quad \Rightarrow \quad \sin\theta > 0)$. If, in addition, $\partial A/\partial\theta = 0$, then

$$\begin{aligned} 0 &= wx\cos\theta - 2x^2\cos\theta + x^2(2\cos^2\theta - 1) \\ &= wx\left(2 - \frac{w}{2x}\right) - 2x^2\left(2 - \frac{w}{2x}\right) + x^2\left[2\left(2 - \frac{w}{2x}\right)^2 - 1\right] \\ &= 2wx - \tfrac{1}{2}w^2 - 4x^2 + wx + x^2\left[8 - \frac{4w}{x} + \frac{w^2}{2x^2} - 1\right] = -wx + 3x^2 = x(3x - w) \end{aligned}$$

Since $x > 0$, we must have $x = \frac{1}{3}w$, in which case $\cos\theta = \frac{1}{2}$, so $\theta = \frac{\pi}{3}$, $\sin\theta = \frac{\sqrt{3}}{2}$, $k = \frac{\sqrt{3}}{6}w$, $b_1 = \frac{1}{3}w$, $b_2 = \frac{2}{3}w$, and $A = \frac{\sqrt{3}}{12}w^2$. As in Example 14.7.6, we can argue from the physical nature of this problem that we have found a local maximum of A. Now checking the boundary of A, let $g(\theta) = A(w/2,\theta) = \frac{1}{2}w^2\sin\theta - \frac{1}{2}w^2\sin\theta + \frac{1}{4}w^2\sin\theta\cos\theta = \frac{1}{8}w^2\sin 2\theta$, $0 < \theta \le \frac{\pi}{2}$. Clearly g is maximized when $\sin 2\theta = 1$ in which case $A = \frac{1}{8}w^2$. Also along the line $\theta = \frac{\pi}{2}$, let $h(x) = A\left(x, \frac{\pi}{2}\right) = wx - 2x^2,\ 0 < x < \frac{1}{2}w \quad \Rightarrow$ $h'(x) = w - 4x = 0 \quad \Leftrightarrow \quad x = \frac{1}{4}w$, and $h\left(\frac{1}{4}w\right) = w\left(\frac{1}{4}w\right) - 2\left(\frac{1}{4}w\right)^2 = \frac{1}{8}w^2$. Since $\frac{1}{8}w^2 < \frac{\sqrt{3}}{12}w^2$, we conclude that the local maximum found earlier was an absolute maximum.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) If the metal were bent into a semi-circular gutter of radius r, we would have $w = \pi r$ and $A = \frac{1}{2}\pi r^2 = \frac{1}{2}\pi\left(\frac{w}{\pi}\right)^2 = \frac{w^2}{2\pi}$.

Since $\frac{w^2}{2\pi} > \frac{\sqrt{3}\,w^2}{12}$, it *would* be better to bend the metal into a gutter with a semicircular cross-section.

4. Since $(x+y+z)^r/(x^2+y^2+z^2)$ is a rational function with domain $\{(x,y,z) \mid (x,y,z) \neq (0,0,0)\}$, f is continuous on $\mathbb{R}^3$ if and only if $\lim_{(x,y,z)\to(0,0,0)} f(x,y,z) = f(0,0,0) = 0$. Recall that $(a+b)^2 \le 2a^2+2b^2$ and a double application of this inequality to $(x+y+z)^2$ gives $(x+y+z)^2 \le 4x^2+4y^2+2z^2 \le 4(x^2+y^2+z^2)$. Now for each r,

$$|(x+y+z)^r| = \left(|x+y+z|^2\right)^{r/2} = \left[(x+y+z)^2\right]^{r/2} \le \left[4(x^2+y^2+z^2)\right]^{r/2} = 2^r(x^2+y^2+z^2)^{r/2}$$

for $(x,y,z) \neq (0,0,0)$. Thus

$$|f(x,y,z)-0| = \left|\frac{(x+y+z)^r}{x^2+y^2+z^2}\right| = \frac{|(x+y+z)^r|}{x^2+y^2+z^2} \le 2^r\frac{(x^2+y^2+z^2)^{r/2}}{x^2+y^2+z^2} = 2^r(x^2+y^2+z^2)^{(r/2)-1}$$

for $(x,y,z) \neq (0,0,0)$. Thus if $(r/2)-1 > 0$, that is $r > 2$, then $2^r(x^2+y^2+z^2)^{(r/2)-1} \to 0$ as $(x,y,z) \to (0,0,0)$ and so $\lim_{(x,y,z)\to(0,0,0)} (x+y+z)^r/(x^2+y^2+z^2) = 0$. Hence for $r > 2$, f is continuous on $\mathbb{R}^3$. Now if $r \le 2$, then as $(x,y,z) \to (0,0,0)$ along the x-axis, $f(x,0,0) = x^r/x^2 = x^{r-2}$ for $x \neq 0$. So when $r = 2$, $f(x,y,z) \to 1 \neq 0$ as $(x,y,z) \to (0,0,0)$ along the x-axis and when $r < 2$ the limit of $f(x,y,z)$ as $(x,y,z) \to (0,0,0)$ along the x-axis doesn't exist and thus can't be zero. Hence for $r \le 2$ f isn't continuous at $(0,0,0)$ and thus is not continuous on $\mathbb{R}^3$.

5. Let $g(x,y) = xf\left(\frac{y}{x}\right)$. Then $g_x(x,y) = f\left(\frac{y}{x}\right) + xf'\left(\frac{y}{x}\right)\left(-\frac{y}{x^2}\right) = f\left(\frac{y}{x}\right) - \frac{y}{x}f'\left(\frac{y}{x}\right)$ and $g_y(x,y) = xf'\left(\frac{y}{x}\right)\left(\frac{1}{x}\right) = f'\left(\frac{y}{x}\right)$. Thus the tangent plane at (x_0,y_0,z_0) on the surface has equation

$$z - x_0 f\left(\frac{y_0}{x_0}\right) = \left[f\left(\frac{y_0}{x_0}\right) - y_0 x_0^{-1} f'\left(\frac{y_0}{x_0}\right)\right](x-x_0) + f'\left(\frac{y_0}{x_0}\right)(y-y_0) \quad \Rightarrow$$

$$\left[f\left(\frac{y_0}{x_0}\right) - y_0 x_0^{-1} f'\left(\frac{y_0}{x_0}\right)\right]x + \left[f'\left(\frac{y_0}{x_0}\right)\right]y - z = 0.$$ But any plane whose equation is of the form $ax+by+cz=0$ passes through the origin. Thus the origin is the common point of intersection.

6. (a) At $(x_1,y_1,0)$ the equations of the tangent planes to $z = f(x,y)$ and $z = g(x,y)$ are

$$P_1\colon\ z - f(x_1,y_1) = f_x(x_1,y_1)(x-x_1) + f_y(x_1,y_1)(y-y_1)$$

and

$$P_2\colon\ z - g(x_1,y_1) = g_x(x_1,y_1)(x-x_1) + g_y(x_1,y_1)(y-y_1)$$

respectively. P_1 intersects the xy-plane in the line given by $f_x(x_1,y_1)(x-x_1) + f_y(x_1,y_1)(y-y_1) = -f(x_1,y_1)$, $z = 0$; and P_2 intersects the xy-plane in the line given by $g_x(x_1,y_1)(x-x_1) + g_y(x_1,y_1)(y-y_1) = -g(x_1,y_1)$, $z = 0$. The point $(x_2,y_2,0)$ is the point of intersection of these two lines, since $(x_2,y_2,0)$ is the point where the line of

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

intersection of the two tangent planes intersects the xy-plane. Thus (x_2, y_2) is the solution of the simultaneous equations

$$f_x(x_1, y_1)(x_2 - x_1) + f_y(x_1, y_1)(y_2 - y_1) = -f(x_1, y_1)$$

and

$$g_x(x_1, y_1)(x_2 - x_1) + g_y(x_1, y_1)(y_2 - y_1) = -g(x_1, y_1)$$

For simplicity, rewrite $f_x(x_1, y_1)$ as f_x and similarly for f_y, g_x, g_y, f and g and solve the equations $(f_x)(x_2 - x_1) + (f_y)(y_2 - y_1) = -f$ and $(g_x)(x_2 - x_1) + (g_y)(y_2 - y_1) = -g$ simultaneously for $(x_2 - x_1)$ and $(y_2 - y_1)$. Then $y_2 - y_1 = \dfrac{gf_x - fg_x}{g_x f_y - f_x g_y}$ or $y_2 = y_1 - \dfrac{gf_x - fg_x}{f_x g_y - g_x f_y}$ and $(f_x)(x_2 - x_1) + \dfrac{(f_y)(gf_x - fg_x)}{g_x f_y - f_x g_y} = -f$ so $x_2 - x_1 = \dfrac{-f - [(f_y)(gf_x - fg_x)/(g_x f_y - f_x g_y)]}{f_x} = \dfrac{fg_y - f_y g}{g_x f_y - f_x g_y}$. Hence $x_2 = x_1 - \dfrac{fg_y - f_y g}{f_x g_y - g_x f_y}$.

(b) Let $f(x, y) = x^x + y^y - 1000$ and $g(x, y) = x^y + y^x - 100$. Then we wish to solve the system of equations $f(x, y) = 0$, $g(x, y) = 0$. Recall $\dfrac{d}{dx}[x^x] = x^x(1 + \ln x)$ (differentiate logarithmically), so $f_x(x, y) = x^x(1 + \ln x)$, $f_y(x, y) = y^y(1 + \ln y)$, $g_x(x, y) = yx^{y-1} + y^x \ln y$, and $g_y(x, y) = x^y \ln x + xy^{x-1}$. Looking at the graph, we estimate the first point of intersection of the curves, and thus the solution to the system, to be approximately $(2.5, 4.5)$. Then following the method of part (a), $x_1 = 2.5$, $y_1 = 4.5$ and

$$x_2 = 2.5 - \frac{f(2.5, 4.5)\, g_y(2.5, 4.5) - f_y(2.5, 4.5)\, g(2.5, 4.5)}{f_x(2.5, 4.5)\, g_y(2.5, 4.5) - f_y(2.5, 4.5)\, g_x(2.5, 4.5} \approx 2.447674117$$

$$y_2 = 4.5 - \frac{f_x(2.5, 4.5)\, g(2.5, 4.5) - f(2.5, 4.5)\, g_x(2.5, 4.5)}{f_x(2.5, 4.5)\, g_y(2.5, 4.5) - f_y(2.5, 4.5)\, g_x(2.5, 4.5)} \approx 4.555657467$$

Continuing this procedure, we arrive at the following values. (If you use a CAS, you may need to increase its computational precision.)

$x_1 = 2.5$	$y_1 = 4.5$
$x_2 = 2.447674117$	$y_2 = 4.555657467$
$x_3 = 2.449614877$	$y_3 = 4.551969333$
$x_4 = 2.449624628$	$y_4 = 4.551951420$
$x_5 = 2.449624628$	$y_5 = 4.551951420$

Thus, to six decimal places, the point of intersection is $(2.449625, 4.551951)$. The second point of intersection can be found similarly, or, by symmetry it is approximately $(4.551951, 2.449625)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. Since we are minimizing the area of the ellipse, and the circle lies above the x-axis, the ellipse will intersect the circle for only one value of y. This y-value must satisfy both the equation of the circle and the equation of the ellipse. Now $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1 \;\Rightarrow\; x^2=\dfrac{a^2}{b^2}\left(b^2-y^2\right)$. Substituting into the equation of the circle gives $\dfrac{a^2}{b^2}\left(b^2-y^2\right)+y^2-2y=0 \;\Rightarrow\; \left(\dfrac{b^2-a^2}{b^2}\right)y^2-2y+a^2=0.$



In order for there to be only one solution to this quadratic equation, the discriminant must be 0, so $4-4a^2\,\dfrac{b^2-a^2}{b^2}=0 \;\Rightarrow$ $b^2-a^2b^2+a^4=0$. The area of the ellipse is $A(a,b)=\pi ab$, and we minimize this function subject to the constraint $g(a,b)=b^2-a^2b^2+a^4=0$.

Now $\nabla A=\lambda\nabla g \;\Leftrightarrow\; \pi b=\lambda(4a^3-2ab^2),\ \pi a=\lambda(2b-2ba^2) \;\Rightarrow\; \lambda=\dfrac{\pi b}{2a(2a^2-b^2)}$ **(1)**,

$\lambda=\dfrac{\pi a}{2b(1-a^2)}$ **(2)**, $b^2-a^2b^2+a^4=0$ **(3)**. Comparing **(1)** and **(2)** gives $\dfrac{\pi b}{2a(2a^2-b^2)}=\dfrac{\pi a}{2b(1-a^2)} \;\Rightarrow$

$2\pi b^2=4\pi a^4 \;\Leftrightarrow\; a^2=\frac{1}{\sqrt{2}}\,b$. Substitute this into **(3)** to get $b=\frac{3}{\sqrt{2}} \;\Rightarrow\; a=\sqrt{\frac{3}{2}}$.

8. Let $\mathbf{u}=\langle a,b,c\rangle$ and $\mathbf{v}=\langle x,y,1\rangle$, so $|\mathbf{u}|=\sqrt{a^2+b^2+c^2}$, $|\mathbf{v}|=\sqrt{x^2+y^2+1}$, and $\mathbf{u}\cdot\mathbf{v}=ax+by+c$. Then by the Cauchy-Schwarz Inequality, $|\mathbf{u}\cdot\mathbf{v}|\le|\mathbf{u}|\,|\mathbf{v}| \;\Rightarrow\; |ax+by+c|\le\sqrt{a^2+b^2+c^2}\,\sqrt{x^2+y^2+1}$. Squaring both sides, we have $(ax+by+c)^2\le\left(a^2+b^2+c^2\right)\left(x^2+y^2+1\right) \;\Rightarrow\; \dfrac{(ax+by+c)^2}{x^2+y^2+1}\le a^2+b^2+c^2$

(since $x^2+y^2+1>0$). Thus $f(x,y)=\dfrac{(ax+by+c)^2}{x^2+y^2+1}\le a^2+b^2+c^2$. We have equality if $(ax+by+c)^2=\left(a^2+b^2+c^2\right)\left(x^2+y^2+1\right)$ or equivalently $c^2\left[(a/c)x+(b/c)y+1\right]^2=c^2\left[(a/c)^2+(b/c)^2+1\right]\left(x^2+y^2+1\right)$ which is true when $x=a/c$ and $y=b/c$. Thus the maximum value of f is $f(a/c,b/c)=a^2+b^2+c^2$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

15 □ MULTIPLE INTEGRALS

15.1 Double Integrals over Rectangles

1. (a) The subrectangles are shown in the figure.



The surface is the graph of $f(x, y) = xy$ and $\Delta A = 4$, so we estimate

$$V \approx \sum_{i=1}^{3} \sum_{j=1}^{2} f(x_i, y_j)\,\Delta A$$

$$= f(2,2)\,\Delta A + f(2,4)\,\Delta A + f(4,2)\,\Delta A + f(4,4)\,\Delta A + f(6,2)\,\Delta A + f(6,4)\,\Delta A$$

$$= 4(4) + 8(4) + 8(4) + 16(4) + 12(4) + 24(4) = 288$$

(b) $V \approx \sum_{i=1}^{3} \sum_{j=1}^{2} f(\overline{x}_i, \overline{y}_j)\,\Delta A = f(1,1)\,\Delta A + f(1,3)\,\Delta A + f(3,1)\,\Delta A + f(3,3)\,\Delta A + f(5,1)\,\Delta A + f(5,3)\,\Delta A$

$$= 1(4) + 3(4) + 3(4) + 9(4) + 5(4) + 15(4) = 144$$

2. (a) The subrectangles are shown in the figure.



Here $\Delta A = 2$ and we estimate

$$\iint_R (1 - xy^2)\,dA \approx \sum_{i=1}^{2} \sum_{j=1}^{3} f(x_{ij}^*, y_{ij}^*)\,\Delta A$$

$$= f(2,-1)\,\Delta A + f(2,0)\,\Delta A + f(2,1)\,\Delta A + f(4,-1)\,\Delta A + f(4,0)\,\Delta A + f(4,1)\,\Delta A$$

$$= (-1)(2) + 1(2) + (-1)(2) + (-3)(2) + 1(2) + (-3)(2) = -12$$

(b) $\iint_R (1 - xy^2)\,dA \approx \sum_{i=1}^{2} \sum_{j=1}^{3} f(x_{ij}^*, y_{ij}^*)\,\Delta A$

$$= f(0,0)\,\Delta A + f(0,1)\,\Delta A + f(0,2)\,\Delta A + f(2,0)\,\Delta A + f(2,1)\,\Delta A + f(2,2)\,\Delta A$$

$$= 1(2) + 1(2) + 1(2) + 1(2) + (-1)(2) + (-7)(2) = -8$$

3. (a) The subrectangles are shown in the figure. Since $\Delta A = 1 \cdot \frac{1}{2} = \frac{1}{2}$, we estimate



$$\iint_R xe^{-xy}\,dA \approx \sum_{i=1}^{2} \sum_{j=1}^{2} f(x_{ij}^*, y_{ij}^*)\,\Delta A$$

$$= f\left(1, \tfrac{1}{2}\right)\,\Delta A + f(1,1)\,\Delta A + f\left(2, \tfrac{1}{2}\right)\,\Delta A + f(2,1)\,\Delta A$$

$$= e^{-1/2}\left(\tfrac{1}{2}\right) + e^{-1}\left(\tfrac{1}{2}\right) + 2e^{-1}\left(\tfrac{1}{2}\right) + 2e^{-2}\left(\tfrac{1}{2}\right) \approx 0.990$$

(b) $\iint_R xe^{-xy}\,dA \approx \sum_{i=1}^{2} \sum_{j=1}^{2} f(\overline{x}_i, \overline{y}_j)\,\Delta A$



$$= f\left(\tfrac{1}{2}, \tfrac{1}{4}\right)\,\Delta A + f\left(\tfrac{1}{2}, \tfrac{3}{4}\right)\,\Delta A + f\left(\tfrac{3}{2}, \tfrac{1}{4}\right)\,\Delta A + f\left(\tfrac{3}{2}, \tfrac{3}{4}\right)\,\Delta A$$

$$= \tfrac{1}{2}e^{-1/8}\left(\tfrac{1}{2}\right) + \tfrac{1}{2}e^{-3/8}\left(\tfrac{1}{2}\right) + \tfrac{3}{2}e^{-3/8}\left(\tfrac{1}{2}\right) + \tfrac{3}{2}e^{-9/8}\left(\tfrac{1}{2}\right) \approx 1.151$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

4. (a) The subrectangles are shown in the figure.



The surface is the graph of $f(x, y) = 1 + x^2 + 3y$ and $\Delta A = \frac{1}{2} \cdot \frac{3}{2} = \frac{3}{4}$, so we estimate

$$V = \iint_R (1 + x^2 + 3y)\, dA \approx \sum_{i=1}^{2} \sum_{j=1}^{2} f(x_{ij}^*, y_{ij}^*)\, \Delta A$$

$$= f(1,0)\, \Delta A + f\left(1, \tfrac{3}{2}\right) \Delta A + f\left(\tfrac{3}{2}, 0\right) \Delta A + f\left(\tfrac{3}{2}, \tfrac{3}{2}\right) \Delta A$$

$$= 2\left(\tfrac{3}{4}\right) + \tfrac{13}{2}\left(\tfrac{3}{4}\right) + \tfrac{13}{4}\left(\tfrac{3}{4}\right) + \tfrac{31}{4}\left(\tfrac{3}{4}\right) = \tfrac{39}{2}\left(\tfrac{3}{4}\right) = \tfrac{117}{8} = 14.625$$

(b) $V = \iint_R (1 + x^2 + 3y)\, dA \approx \sum_{i=1}^{2} \sum_{j=1}^{2} f(\overline{x}_i, \overline{y}_j)\, \Delta A$



$$= f\left(\tfrac{5}{4}, \tfrac{3}{4}\right) \Delta A + f\left(\tfrac{5}{4}, \tfrac{9}{4}\right) \Delta A + f\left(\tfrac{7}{4}, \tfrac{3}{4}\right) \Delta A + f\left(\tfrac{7}{4}, \tfrac{9}{4}\right) \Delta A$$

$$= \tfrac{77}{16}\left(\tfrac{3}{4}\right) + \tfrac{149}{16}\left(\tfrac{3}{4}\right) + \tfrac{101}{16}\left(\tfrac{3}{4}\right) + \tfrac{173}{16}\left(\tfrac{3}{4}\right) = \tfrac{375}{16} = 23.4375$$

5. The values of $f(x, y) = \sqrt{52 - x^2 - y^2}$ get smaller as we move farther from the origin, so on any of the subrectangles in the problem, the function will have its largest value at the lower left corner of the subrectangle and its smallest value at the upper right corner, and any other value will lie between these two. So using these subrectangles we have $L < V < U$. (Note that this is true no matter how R is divided into subrectangles.)

6. To approximate the volume, let R be the planar region corresponding to the surface of the water in the pool, and place R on coordinate axes so that x and y correspond to the dimensions given. Then we define $f(x, y)$ to be the depth of the water at (x, y), so the volume of water in the pool is the volume of the solid that lies above the rectangle $R = [0, 20] \times [0, 30]$ and below the graph of $f(x, y)$. We can estimate this volume using the Midpoint Rule with $m = 2$ and $n = 3$, so $\Delta A = 100$. Each subrectangle with its midpoint is shown in the figure. Then



$$V \approx \sum_{i=1}^{2} \sum_{j=1}^{3} f(\overline{x}_i, \overline{y}_j)\, \Delta A = \Delta A[f(5,5) + f(5,15) + f(5,25) + f(15,5) + f(15,15) + f(15,25)]$$

$$= 100(3 + 7 + 10 + 3 + 5 + 8) = 3600$$

Thus, we estimate that the pool contains 3600 cubic feet of water.

Alternatively, we can approximate the volume with a Riemann sum where $m = 4$, $n = 6$ and the sample points are taken to be, for example, the upper right corner of each subrectangle. Then $\Delta A = 25$ and

$$V \approx \sum_{i=1}^{4} \sum_{j=1}^{6} f(x_i, y_j)\, \Delta A$$

$$= 25[3 + 4 + 7 + 8 + 10 + 8 + 4 + 6 + 8 + 10 + 12 + 10 + 3 + 4 + 5 + 6 + 8 + 7 + 2 + 2 + 2 + 3 + 4 + 4]$$

$$= 25(140) = 3500$$

So we estimate that the pool contains 3500 ft^3 of water.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. (a) With $m = n = 2$, we have $\Delta A = 4$. Using the contour map to estimate the value of f at the center of each subrectangle, we have

$$\iint_R f(x,y)\,dA \approx \sum_{i=1}^{2}\sum_{j=1}^{2} f(\overline{x}_i, \overline{y}_j)\,\Delta A = \Delta A[f(1,1) + f(1,3) + f(3,1) + f(3,3)] \approx 4(27 + 4 + 14 + 17) = 248$$

(b) $f_{\text{ave}} = \frac{1}{A(R)} \iint_R f(x,y)\,dA \approx \frac{1}{16}(248) = 15.5$

8. As in Example 9, we place the origin at the southwest corner of the state. Then $R = [0, 388] \times [0, 276]$ (in miles) is the rectangle corresponding to Colorado and we define $f(x, y)$ to be the temperature at the location (x, y). The average temperature is given by

$$f_{\text{ave}} = \frac{1}{A(R)} \iint_R f(x,y)\,dA = \frac{1}{388 \cdot 276} \iint_R f(x,y)\,dA$$

To use the Midpoint Rule with $m = n = 4$, we divide R into 16 regions of equal size, as shown in the figure, with the center of each subrectangle indicated.



The area of each subrectangle is $\Delta A = \frac{388}{4} \cdot \frac{276}{4} = 6693$, so using the contour map to estimate the function values at each midpoint, we have

$$\begin{aligned}\iint_R f(x,y)\,dA &\approx \sum_{i=1}^{4}\sum_{j=1}^{4} f(\overline{x}_i, \overline{y}_j)\,\Delta A \\ &\approx \Delta A\,[31 + 28 + 52 + 43 + 43 + 25 + 57 + 46 + 36 + 20 + 42 + 45 + 30 + 23 + 43 + 41] \\ &= 6693(605)\end{aligned}$$

Therefore, $f_{\text{ave}} \approx \dfrac{6693 \cdot 605}{388 \cdot 276} \approx 37.8$, so the average temperature in Colorado at 4:00 PM on February 26, 2007, was approximately $37.8°\text{F}$.

9. $z = \sqrt{2} > 0$, so we can interpret the double integral as the volume of the solid S that lies below the plane $z = \sqrt{2}$ and above the rectangle $[2, 6] \times [-1, 5]$. S is a rectangular solid, so $\iint_R \sqrt{2}\,dA = 4 \cdot 6 \cdot \sqrt{2} = 24\sqrt{2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

10. $z = 2x + 1 \geq 0$ for $0 \leq x \leq 2$, so we can interpret the integral as the volume of the solid S that lies below the plane $z = 2x + 1$ and above the rectangle $[0, 2] \times [0, 4]$. We can picture S as a rectangular solid (with height 1) surmounted by a triangular cylinder; thus

$$\iint_R (2x + 1)\, dA = (2)(4)(1) + \tfrac{1}{2}(2)(4)(4) = 24$$



11. $z = 4 - 2y \geq 0$ for $0 \leq y \leq 1$, so we can interpret the integral as the volume of the solid S that lies below the plane $z = 4 - 2y$ and above the square $[0, 1] \times [0, 1]$. We can picture S as a rectangular solid (with height 2) surmounted by a triangular cylinder; thus

$$\iint_R (4 - 2y)\, dA = (1)(1)(2) + \tfrac{1}{2}(1)(1)(2) = 3$$



12. Here $z = \sqrt{9 - y^2}$, so $z^2 + y^2 = 9$, $z \geq 0$. Thus the integral represents the volume of the top half of the part of the circular cylinder $z^2 + y^2 = 9$ that lies above the rectangle $[0, 4] \times [0, 2]$.



13. $\int_0^2 (x + 3x^2y^2)\, dx = \left[\dfrac{x^2}{2} + 3\,\dfrac{x^3}{3}\,y^2\right]_{x=0}^{x=2} = \left[\tfrac{1}{2}x^2 + x^3y^2\right]_{x=0}^{x=2} = \left[\tfrac{1}{2}(2)^2 + (2)^3y^2\right] - \left[\tfrac{1}{2}(0)^2 + (0)^3y^2\right] = 2 + 8y^2.$

$\int_0^3 (x + 3x^2y^2)\, dy = \left[xy + 3x^2\,\dfrac{y^3}{3}\right]_{y=0}^{y=3} = \left[xy + x^2y^3\right]_{y=0}^{y=3} = \left[x(3) + x^2(3)^3\right] - \left[x(0) + x^2(0)^3\right] = 3x + 27x^2$

14. $\int_0^2 y\sqrt{x+2}\, dx = \left[y \cdot \tfrac{2}{3}(x+2)^{3/2}\right]_{x=0}^{x=2} = \tfrac{2}{3}y(4)^{3/2} - \tfrac{2}{3}y(2)^{3/2} = \tfrac{16}{3}y - \tfrac{4}{3}\sqrt{2}\,y = \tfrac{4}{3}(4 - \sqrt{2})\,y,$

$\int_0^3 y\sqrt{x+2}\, dy = \left[\dfrac{y^2}{2}\sqrt{x+2}\right]_{y=0}^{y=3} = \tfrac{1}{2}(3)^2\sqrt{x+2} - \tfrac{1}{2}(0)^2\sqrt{x+2} = \tfrac{9}{2}\sqrt{x+2}$

15. $\int_1^4 \int_0^2 (6x^2y - 2x)\, dy\, dx = \int_1^4 \left[3x^2y^2 - 2xy\right]_{y=0}^{y=2} dx = \int_1^4 \left[(12x^2 - 4x) - (0 - 0)\right] dx$

$= \int_1^4 (12x^2 - 4x)\, dx = \left[4x^3 - 2x^2\right]_1^4 = (256 - 32) - (4 - 2) = 222$

16. $\int_0^1 \int_0^1 (x + y)^2\, dx\, dy = \int_0^1 \int_0^1 (x^2 + 2xy + y^2)\, dx\, dy = \int_0^1 \left[\tfrac{1}{3}x^3 + x^2y + xy^2\right]_{x=0}^{x=1} dy$

$= \int_0^1 (\tfrac{1}{3} + y + y^2)\, dy = \left[\tfrac{1}{3}y + \tfrac{1}{2}y^2 + \tfrac{1}{3}y^3\right]_0^1 = \tfrac{1}{3} + \tfrac{1}{2} + \tfrac{1}{3} - 0 = \tfrac{7}{6}$

17. $\int_0^1 \int_1^2 (x + e^{-y})\, dx\, dy = \int_0^1 \left[\tfrac{1}{2}x^2 + xe^{-y}\right]_{x=1}^{x=2} dy = \int_0^1 \left[(2 + 2e^{-y}) - (\tfrac{1}{2} + e^{-y})\right] dy$

$= \int_0^1 (\tfrac{3}{2} + e^{-y})\, dy = \left[\tfrac{3}{2}y - e^{-y}\right]_0^1 = \left(\tfrac{3}{2} - e^{-1}\right) - (0 - 1) = \tfrac{5}{2} - e^{-1}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $\int_0^{\pi/6}\int_0^{\pi/2}(\sin x+\sin y)\,dy\,dx=\int_0^{\pi/6}\left[y\sin x-\cos y\right]_{y=0}^{y=\pi/2}dx=\int_0^{\pi/6}\left[\left(\frac{\pi}{2}\sin x-0\right)-(0-1)\right]dx$

$=\int_0^{\pi/6}\left(\frac{\pi}{2}\sin x+1\right)dx=\left[-\frac{\pi}{2}\cos x+x\right]_0^{\pi/6}$

$=\left[\left(-\frac{\pi}{2}\cdot\frac{\sqrt{3}}{2}+\frac{\pi}{6}\right)-\left(-\frac{\pi}{2}+0\right)\right]=\left(\frac{2}{3}-\frac{\sqrt{3}}{4}\right)\pi$

19. $\int_{-3}^{3}\int_0^{\pi/2}(y+y^2\cos x)\,dx\,dy=\int_{-3}^{3}\left[xy+y^2\sin x\right]_{x=0}^{x=\pi/2}dy=\int_{-3}^{3}\left(\frac{\pi}{2}y+y^2\right)dy$

$=\left[\frac{\pi}{4}y^2+\frac{1}{3}y^3\right]_{-3}^{3}=\left[\left(\frac{9\pi}{4}+9\right)-\left(\frac{9\pi}{4}-9\right)\right]=18$

20. $\int_1^3\int_1^5\frac{\ln y}{xy}\,dy\,dx=\int_1^3\frac{1}{x}\,dx\int_1^5\frac{\ln y}{y}\,dy$ [by Equation 11]

$=\left[\ln|x|\right]_1^3\left[\frac{1}{2}(\ln y)^2\right]_1^5$ [substitute $u=\ln y\ \Rightarrow\ du=(1/y)\,dy$]

$=(\ln 3-0)\cdot\frac{1}{2}\left[(\ln 5)^2-0\right]=\frac{1}{2}(\ln 3)(\ln 5)^2$

21. $\int_1^4\int_1^2\left(\frac{x}{y}+\frac{y}{x}\right)dy\,dx=\int_1^4\left[x\ln|y|+\frac{1}{x}\cdot\frac{1}{2}y^2\right]_{y=1}^{y=2}dx=\int_1^4\left(x\ln 2+\frac{3}{2x}\right)dx=\left[\frac{1}{2}x^2\ln 2+\frac{3}{2}\ln|x|\right]_1^4$

$=\left(8\ln 2+\frac{3}{2}\ln 4\right)-\left(\frac{1}{2}\ln 2+0\right)=\frac{15}{2}\ln 2+\frac{3}{2}\ln 4$ or $\frac{15}{2}\ln 2+3\ln(4^{1/2})=\frac{21}{2}\ln 2$

22. $\int_0^1\int_0^2 ye^{x-y}\,dx\,dy=\int_0^1\int_0^2 ye^xe^{-y}\,dx\,dy=\int_0^2 e^x\,dx\int_0^1 ye^{-y}\,dy$ [by Equation 11]

$=\left[e^x\right]_0^2\left[(-y-1)e^{-y}\right]_0^1$ [by integrating by parts]

$=\left(e^2-e^0\right)\left[-2e^{-1}-\left(-e^0\right)\right]=(e^2-1)(1-2e^{-1})$ or $e^2-2e+2e^{-1}-1$

23. $\int_0^3\int_0^{\pi/2}t^2\sin^3\phi\,d\phi\,dt=\int_0^{\pi/2}\sin^3\phi\,d\phi\int_0^3 t^2\,dt$ [by Equation 11] $=\int_0^{\pi/2}(1-\cos^2\phi)\sin\phi\,d\phi\int_0^3 t^2\,dt$

$=\left[\frac{1}{3}\cos^3\phi-\cos\phi\right]_0^{\pi/2}\left[\frac{1}{3}t^3\right]_0^3=\left[(0-0)-\left(\frac{1}{3}-1\right)\right]\cdot\frac{1}{3}(27-0)=\frac{2}{3}(9)=6$

24. $\int_0^1\int_0^1 xy\sqrt{x^2+y^2}\,dy\,dx=\int_0^1 x\left[\frac{1}{3}(x^2+y^2)^{3/2}\right]_{y=0}^{y=1}dx=\frac{1}{3}\int_0^1 x[(x^2+1)^{3/2}-x^3]\,dx=\frac{1}{3}\int_0^1[x(x^2+1)^{3/2}-x^4]dx$

$=\frac{1}{3}\left[\frac{1}{5}(x^2+1)^{5/2}-\frac{1}{5}x^5\right]_0^1=\frac{1}{15}\left[\left(2^{5/2}-1\right)-(1-0)\right]=\frac{2}{15}\left(2\sqrt{2}-1\right)$

25. $\int_0^1\int_0^1 v(u+v^2)^4\,du\,dv=\int_0^1\left[\frac{1}{5}v(u+v^2)^5\right]_{u=0}^{u=1}dv=\frac{1}{5}\int_0^1 v\left[(1+v^2)^5-(0+v^2)^5\right]dv$

$=\frac{1}{5}\int_0^1\left[v(1+v^2)^5-v^{11}\right]dv=\frac{1}{5}\left[\frac{1}{2}\cdot\frac{1}{6}(1+v^2)^6-\frac{1}{12}v^{12}\right]_0^1$

[substitute $t=1+v^2\ \Rightarrow\ dt=2v\,dv$ in the first term]

$=\frac{1}{60}\left[\left(2^6-1\right)-(1-0)\right]=\frac{1}{60}(63-1)=\frac{31}{30}$

26. $\int_0^1\int_0^1\sqrt{s+t}\,ds\,dt=\int_0^1\left[\frac{2}{3}(s+t)^{3/2}\right]_{s=0}^{s=1}dt=\frac{2}{3}\int_0^1[(1+t)^{3/2}-t^{3/2}]\,dt=\frac{2}{3}\left[\frac{2}{5}(1+t)^{5/2}-\frac{2}{5}t^{5/2}\right]_0^1$

$=\frac{4}{15}[(2^{5/2}-1)-(1-0)]=\frac{4}{15}\left(2^{5/2}-2\right)$ or $\frac{8}{15}\left(2\sqrt{2}-1\right)$

27. $\iint_R x\sec^2 y\,dA=\int_0^2\int_0^{\pi/4}x\sec^2 y\,dy\,dx=\int_0^2 x\,dx\int_0^{\pi/4}\sec^2 y\,dy=\left[\frac{1}{2}x^2\right]_0^2\left[\tan y\right]_0^{\pi/4}$

$=(2-0)\left(\tan\frac{\pi}{4}-\tan 0\right)=2(1-0)=2$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

28. $\iint_R (y+xy^{-2})\,dA = \int_1^2 \int_0^2 (y+xy^{-2})\,dx\,dy = \int_1^2 \left[xy+\frac{1}{2}x^2y^{-2}\right]_{x=0}^{x=2} dy = \int_1^2 \left(2y+2y^{-2}\right) dy$

$= \left[y^2-2y^{-1}\right]_1^2 = (4-1)-(1-2) = 4$

29. $\iint_R \frac{xy^2}{x^2+1}\,dA = \int_0^1 \int_3^3 \frac{xy^2}{x^2+1}\,dy\,dx = \int_0^1 \frac{x}{x^2+1}\,dx \int_{-3}^3 y^2\,dy = \left[\frac{1}{2}\ln(x^2+1)\right]_0^1 \left[\frac{1}{3}y^3\right]_{-3}^3$

$= \frac{1}{2}(\ln 2 - \ln 1)\cdot\frac{1}{3}(27+27) = 9\ln 2$

30. $\iint_R \frac{\tan\theta}{\sqrt{1-t^2}}\,dA = \int_0^{1/2}\int_0^{\pi/3} \frac{\tan\theta}{\sqrt{1-t^2}}\,d\theta\,dt = \int_0^{1/2} \frac{1}{\sqrt{1-t^2}}\,dt \int_0^{\pi/3} \tan\theta\,d\theta = \left[\sin^{-1}t\right]_0^{1/2} \left[\ln|\sec\theta|\right]_0^{\pi/3}$

$= \left(\sin^{-1}\tfrac{1}{2} - \sin^{-1}0\right)\left(\ln\left|\sec\tfrac{\pi}{3}\right| - \ln|\sec 0|\right) = \left(\tfrac{\pi}{6}-0\right)(\ln 2 - \ln 1) = \tfrac{\pi}{6}\ln 2$

31. $\int_0^{\pi/6}\int_0^{\pi/3} x\sin(x+y)\,dy\,dx$

$= \int_0^{\pi/6} \left[-x\cos(x+y)\right]_{y=0}^{y=\pi/3} dx = \int_0^{\pi/6} \left[x\cos x - x\cos\left(x+\tfrac{\pi}{3}\right)\right] dx$

$= x\left[\sin x - \sin\left(x+\tfrac{\pi}{3}\right)\right]_0^{\pi/6} - \int_0^{\pi/6} \left[\sin x - \sin\left(x+\tfrac{\pi}{3}\right)\right] dx$ [by integrating by parts separately for each term]

$= \tfrac{\pi}{6}\left[\tfrac{1}{2}-1\right] - \left[-\cos x + \cos\left(x+\tfrac{\pi}{3}\right)\right]_0^{\pi/6} = -\tfrac{\pi}{12} - \left[-\tfrac{\sqrt{3}}{2}+0-\left(-1+\tfrac{1}{2}\right)\right] = \tfrac{\sqrt{3}-1}{2} - \tfrac{\pi}{12}$

32. $\iint_R \frac{x}{1+xy}\,dA = \int_0^1\int_0^1 \frac{x}{1+xy}\,dy\,dx = \int_0^1 \left[\ln(1+xy)\right]_{y=0}^{y=1} dx = \int_0^1 \left[\ln(1+x)-\ln 1\right] dx$

$= \int_0^1 \ln(1+x)\,dx = \left[(1+x)\ln(1+x)-x\right]_0^1$ [by integrating by parts]

$= (2\ln 2 - 1) - (\ln 1 - 0) = 2\ln 2 - 1$

33. $\iint_R ye^{-xy}\,dA = \int_0^3\int_0^2 ye^{-xy}\,dx\,dy = \int_0^3 \left[-e^{-xy}\right]_{x=0}^{x=2} dy = \int_0^3 (-e^{-2y}+1)\,dy = \left[\tfrac{1}{2}e^{-2y}+y\right]_0^3$

$= \frac{1}{2}e^{-6} + 3 - \left(\frac{1}{2}+0\right) = \frac{1}{2}e^{-6} + \frac{5}{2}$

34. $\iint_R \frac{1}{1+x+y}\,dA = \int_1^3\int_1^2 \frac{1}{1+x+y}\,dy\,dx = \int_1^3 \left[\ln(1+x+y)\right]_{y=1}^{y=2} dx = \int_1^3 \left[\ln(x+3)-\ln(x+2)\right] dx$

$= \left[\left((x+3)\ln(x+3)-(x+3)\right) - \left((x+2)\ln(x+2)-(x+2)\right)\right]_1^3$

[by integrating by parts separately for each term]

$= (6\ln 6 - 6 - 5\ln 5 + 5) - (4\ln 4 - 4 - 3\ln 3 + 3) = 6\ln 6 - 5\ln 5 - 4\ln 4 + 3\ln 3$

35. $z = f(x,y) = 4 - x - 2y \geq 0$ for $0 \leq x \leq 1$ and $0 \leq y \leq 1$. So the solid is the region in the first octant which lies below the plane $z = 4 - x - 2y$ and above $[0,1]\times[0,1]$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $z = 2 - x^2 - y^2 \geq 0$ for $0 \leq x \leq 1$ and $0 \leq y \leq 1$. So the solid is the region in the first octant which lies below the circular paraboloid $z = 2 - x^2 - y^2$ and above $[0, 1] \times [0, 1]$.



37. The solid lies under the plane $4x + 6y - 2z + 15 = 0$ or $z = 2x + 3y + \frac{15}{2}$ so

$$V = \iint_R (2x + 3y + \tfrac{15}{2})\, dA = \int_{-1}^{1} \int_{-1}^{2} (2x + 3y + \tfrac{15}{2})\, dx\, dy = \int_{-1}^{1} \left[x^2 + 3xy + \tfrac{15}{2}x\right]_{x=-1}^{x=2} dy$$

$$= \int_{-1}^{1} \left[(19 + 6y) - (-\tfrac{13}{2} - 3y)\right] dy = \int_{-1}^{1} (\tfrac{51}{2} + 9y)\, dy = \left[\tfrac{51}{2}y + \tfrac{9}{2}y^2\right]_{-1}^{1} = 30 - (-21) = 51$$

38. $V = \iint_R (3y^2 - x^2 + 2)\, dA = \int_{-1}^{1} \int_{1}^{2} (3y^2 - x^2 + 2)\, dy\, dx = \int_{-1}^{1} \left[y^3 - x^2 y + 2y\right]_{y=1}^{y=2} dx$

$$= \int_{-1}^{1} \left[(12 - 2x^2) - (3 - x^2)\right] dx = \int_{-1}^{1} (9 - x^2 dx) = \left[9x - \tfrac{1}{3}x^3\right]_{-1}^{1} = \tfrac{26}{3} + \tfrac{26}{3} = \tfrac{52}{3}$$

39. $V = \int_{-2}^{2} \int_{-1}^{1} (1 - \frac{1}{4}x^2 - \frac{1}{9}y^2)\, dx\, dy = 4 \int_{0}^{2} \int_{0}^{1} (1 - \frac{1}{4}x^2 - \frac{1}{9}y^2)\, dx\, dy$

$$= 4 \int_{0}^{2} \left[x - \tfrac{1}{12}x^3 - \tfrac{1}{9}y^2 x\right]_{x=0}^{x=1} dy = 4 \int_{0}^{2} (\tfrac{11}{12} - \tfrac{1}{9}y^2)\, dy = 4\left[\tfrac{11}{12}y - \tfrac{1}{27}y^3\right]_{0}^{2} = 4 \cdot \tfrac{83}{54} = \tfrac{166}{27}$$

40. The solid lies under the surface $z = x^2 + xy^2$ and above the rectangle $R = [0, 5] \times [-2, 2]$, so its volume is

$$V = \iint_R (x^2 + xy^2)\, dA = \int_{0}^{5} \int_{-2}^{2} (x^2 + xy^2)\, dy\, dx = \int_{0}^{5} \left[x^2 y + \tfrac{1}{3}xy^3\right]_{y=-2}^{y=2} dx$$

$$= \int_{0}^{5} \left[(2x^2 + \tfrac{8}{3}x) - (-2x^2 - \tfrac{8}{3}x)\right] dx = \int_{0}^{5} (4x^2 + \tfrac{16}{3}x)\, dx$$

$$= \left[\tfrac{4}{3}x^3 + \tfrac{8}{3}x^2\right]_{0}^{5} = \tfrac{500}{3} + \tfrac{200}{3} - 0 = \tfrac{700}{3}$$

41. The solid lies under the surface $z = 1 + x^2 y e^y$ and above the rectangle $R = [-1, 1] \times [0, 1]$, so its volume is

$$V = \iint_R (1 + x^2 y e^y)\, dA = \int_{0}^{1} \int_{-1}^{1} (1 + x^2 y e^y)\, dx\, dy = \int_{0}^{1} \left[x + \tfrac{1}{3}x^3 y e^y\right]_{x=-1}^{x=1} dy$$

$$= \int_{0}^{1} (2 + \tfrac{2}{3}y e^y)\, dy = \left[2y + \tfrac{2}{3}(y - 1)\, e^y\right]_{0}^{1} \qquad \text{[by integrating by parts in the second term]}$$

$$= (2 + 0) - (0 - \tfrac{2}{3}e^0) = 2 + \tfrac{2}{3} = \tfrac{8}{3}$$

42. The cylinder intersects the xy-plane along the line $x = 4$, so in the first octant, the solid lies below the surface $z = 16 - x^2$ and above the rectangle $R = [0, 4] \times [0, 5]$ in the xy-plane.

$$V = \int_{0}^{5} \int_{0}^{4} (16 - x^2)\, dx\, dy = \int_{0}^{4} (16 - x^2)\, dx \int_{0}^{5} dy$$

$$= \left[16x - \tfrac{1}{3}x^3\right]_{0}^{4} \left[y\right]_{0}^{5} = (64 - \tfrac{64}{3} - 0)(5 - 0) = \tfrac{640}{3}$$

43. The solid lies below the surface $z = 2 + x^2 + (y - 2)^2$ and above the plane $z = 1$ for $-1 \leq x \leq 1, 0 \leq y \leq 4$. The volume of the solid is the difference in volumes between the solid that lies under $z = 2 + x^2 + (y - 2)^2$ over the rectangle

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$R = [-1, 1] \times [0, 4]$ and the solid that lies under $z = 1$ over R.

$$\begin{aligned} V &= \int_0^4 \int_{-1}^1 [2 + x^2 + (y-2)^2]\, dx\, dy - \int_0^4 \int_{-1}^1 (1)\, dx\, dy \\ &= \int_0^4 \left[2x + \tfrac{1}{3}x^3 + x(y-2)^2\right]_{x=-1}^{x=1} dy - \int_{-1}^1 dx \int_0^4 dy \\ &= \int_0^4 \left[(2 + \tfrac{1}{3} + (y-2)^2) - (-2 - \tfrac{1}{3} - (y-2)^2)\right] dy - [x]_{-1}^1\, [y]_0^4 \\ &= \int_0^4 \left[\tfrac{14}{3} + 2(y-2)^2\right] dy \quad [1-(-1)][4-0] = \left[\tfrac{14}{3}y + \tfrac{2}{3}(y-2)^3\right]_0^4 - (2)(4) \\ &= \left[\left(\tfrac{56}{3} + \tfrac{16}{3}\right) - \left(0 - \tfrac{16}{3}\right)\right] - 8 = \tfrac{88}{3} - 8 = \tfrac{64}{3} \end{aligned}$$

44.



The solid lies below the plane $z = x + 2y$ and above the surface $z = \dfrac{2xy}{x^2+1}$ for $0 \le x \le 2, 0 \le y \le 4$. The volume of the solid is the difference in volumes between the solid that lies under $z = x + 2y$ over the rectangle $R = [0, 2] \times [0, 4]$ and the solid that lies under $z = \dfrac{2xy}{x^2+1}$ over R.

$$\begin{aligned} V &= \int_0^2 \int_0^4 (x + 2y)dy\, dx - \int_0^2 \int_0^4 \frac{2xy}{x^2+1}\, dy\, dx = \int_0^2 \left[xy + y^2\right]_{y=0}^{y=4} dx - \int_0^2 \frac{2x}{x^2+1}\, dx \int_0^4 y\, dy \\ &= \int_0^2 [(4x + 16) - (0 + 0)]\, dx - \left[\ln\left|x^2+1\right|\right]_0^2 \left[\tfrac{1}{2}y^2\right]_0^4 = \left[2x^2 + 16x\right]_0^2 - (\ln 5 - \ln 1)\,(8 - 0) \\ &= (8 + 32 - 0) - 8\ln 5 = 40 - 8\ln 5 \end{aligned}$$

45. In Maple, we can calculate the integral by defining the integrand as `f` and then using the command `int(int(f,x=0..1),y=0..1);`.
In Mathematica, we can use the command

```
Integrate[f,{x,0,1},{y,0,1}]
```

We find that $\iint_R x^5 y^3 e^{xy}\, dA = 21e - 57 \approx 0.0839$. We can use `plot3d` (in Maple) or `Plot3D` (in Mathematica) to graph the function.



46. In Maple, we can calculate the integral by defining `f:=exp(-x^2)*cos(x^2+y^2);` and `g:=2-x^2-y^2;` and then [since $2 - x^2 - y^2 > e^{-x^2}\cos(x^2 + y^2)$ for $-1 \le x \le 1, -1 \le y \le 1$] using the command `evalf(Int(Int(g-f,x=-1..1),y=-1..1));`.
Using `Int` rather than `int` forces Maple to use purely numerical techniques in evaluating the integral.



In Mathematica, we can use the command `NIntegrate[g-f,{x,-1,1},{y,-1,1}]`. We find that $\iint_R \left[(2 - x^2 - y^2) - \left(e^{-x^2}\cos(x^2 + y^2)\right)\right] dA \approx 3.0271$. We can use the `plot3d` command (in Maple) or `Plot3D` (in Mathematica) to graph both functions on the same screen.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. R is the rectangle $[-1,1]\times[0,5]$. Thus, $A(R)=2\cdot 5=10$ and

$$f_{\text{ave}}=\frac{1}{A(R)}\iint_R f(x,y)\,dA=\tfrac{1}{10}\int_0^5\int_{-1}^1 x^2y\,dx\,dy=\tfrac{1}{10}\int_0^5\left[\tfrac{1}{3}x^3y\right]_{x=-1}^{x=1}dy=\tfrac{1}{10}\int_0^5\tfrac{2}{3}y\,dy=\tfrac{1}{10}\left[\tfrac{1}{3}y^2\right]_0^5=\tfrac{5}{6}.$$

48. $A(R)=4\cdot 1=4$, so

$$\begin{aligned}f_{\text{ave}}&=\frac{1}{A(R)}\iint_R f(x,y)\,dA=\tfrac{1}{4}\int_0^4\int_0^1 e^y\sqrt{x+e^y}\,dy\,dx=\tfrac{1}{4}\int_0^4\left[\tfrac{2}{3}(x+e^y)^{3/2}\right]_{y=0}^{y=1}dx\\&=\tfrac{1}{4}\cdot\tfrac{2}{3}\int_0^4[(x+e)^{3/2}-(x+1)^{3/2}]\,dx=\tfrac{1}{6}\left[\tfrac{2}{5}(x+e)^{5/2}-\tfrac{2}{5}(x+1)^{5/2}\right]_0^4\\&=\tfrac{1}{6}\cdot\tfrac{2}{5}[(4+e)^{5/2}-5^{5/2}-e^{5/2}+1]=\tfrac{1}{15}[(4+e)^{5/2}-e^{5/2}-5^{5/2}+1]\approx 3.327\end{aligned}$$

49. $\displaystyle\iint_R\frac{xy}{1+x^4}\,dA=\int_{-1}^1\int_0^1\frac{xy}{1+x^4}\,dy\,dx=\int_{-1}^1\frac{x}{1+x^4}\,dx\int_0^1 y\,dy$ [by Equation 11] but $\displaystyle f(x)=\frac{x}{1+x^4}$ is an odd function so $\displaystyle\int_{-1}^1 f(x)\,dx=0$ (by Theorem 4.5.6 [ET 5.5.7]). Thus $\displaystyle\iint_R\frac{xy}{1+x^4}\,dA=0\cdot\int_0^1 y\,dy=0.$

50. $$\begin{aligned}\iint_R(1+x^2\sin y+y^2\sin x)\,dA&=\iint_R 1\,dA+\iint_R x^2\sin y\,dA+\iint_R y^2\sin x\,dA\\&=A(R)+\int_{-\pi}^{\pi}\int_{-\pi}^{\pi}x^2\sin y\,dy\,dx+\int_{-\pi}^{\pi}\int_{-\pi}^{\pi}y^2\sin x\,dy\,dx\\&=(2\pi)(2\pi)+\int_{-\pi}^{\pi}x^2dx\int_{-\pi}^{\pi}\sin y\,dy+\int_{-\pi}^{\pi}\sin x\,dx\int_{-\pi}^{\pi}y^2\,dy\end{aligned}$$

But $\sin x$ is an odd function, so $\int_{-\pi}^{\pi}\sin x\,dx=\int_{-\pi}^{\pi}\sin y\,dy=0$ (by Theorem 4.5.6 [ET 5.5.7]) and

$\iint_R(1+x^2\sin y+y^2\sin x)\,dA=4\pi^2+0+0=4\pi^2$.

51. Let $\displaystyle f(x,y)=\frac{x-y}{(x+y)^3}$. Then a CAS gives $\int_0^1\int_0^1 f(x,y)\,dy\,dx=\frac{1}{2}$ and $\int_0^1\int_0^1 f(x,y)\,dx\,dy=-\frac{1}{2}$.

To explain the seeming violation of Fubini's Theorem, note that f has an infinite discontinuity at $(0,0)$ and thus does not satisfy the conditions of Fubini's Theorem. In fact, both iterated integrals involve improper integrals which diverge at their lower limits of integration.

52. (a) Loosely speaking, Fubini's Theorem says that the order of integration of a function of two variables does not affect the value of the double integral, while Clairaut's Theorem says that the order of differentiation of such a function does not affect the value of the second-order derivative. Also, both theorems require continuity (though Fubini's allows a finite number of smooth curves to contain discontinuities).

(b) To find g_{xy}, we first hold y constant and use the single-variable Fundamental Theorem of Calculus, Part 1:

$g_x=\dfrac{d}{dx}\,g(x,y)=\dfrac{d}{dx}\displaystyle\int_a^x\left(\int_c^y f(s,t)\,dt\right)ds=\int_c^y f(x,t)\,dt$. Now we use the Fundamental Theorem again:

$g_{xy}=\dfrac{d}{dy}\displaystyle\int_c^y f(x,t)\,dt=f(x,y)$.

To find g_{yx}, we first use Fubini's Theorem to find that $\int_a^x\int_c^y f(s,t)\,dt\,ds=\int_c^y\int_a^x f(s,t)\,dt\,ds$, and then use the Fundamental Theorem twice, as above, to get $g_{yx}=f(x,y)$. So $g_{xy}=g_{yx}=f(x,y)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

15.2 Double Integrals over General Regions

1. $\int_1^5 \int_0^x (8x - 2y)\,dy\,dx = \int_1^5 \left[8xy - y^2\right]_{y=0}^{y=x} dx = \int_1^5 [8x(x) - (x)^2 - 8x(0) + (0)^2]dx$

$= \int_1^5 7x^2\,dx = \frac{7}{3}x^3\Big]_1^5 = \frac{7}{3}(125 - 1) = \frac{868}{3}$

2. $\int_0^2 \int_0^{y^2} x^2 y\,dx\,dy = \int_0^2 \left[\frac{1}{3}x^3 y\right]_{x=0}^{x=y^2} dy = \int_0^2 \frac{1}{3}y\left[(y^2)^3 - (0)^3\right] dy$

$= \int_0^2 \frac{1}{3}y^7\,dy = \frac{1}{3}\left[\frac{1}{8}y^8\right]_0^2 = \frac{1}{3}(32 - 0) = \frac{32}{3}$

3. $\int_0^1 \int_0^y xe^{y^3}\,dx\,dy = \int_0^1 \left[\frac{1}{2}x^2 e^{y^3}\right]_{x=0}^{x=y} dy = \int_0^1 \frac{1}{2}e^{y^3}\left[(y)^2 - (0)^2\right] dy$

$= \frac{1}{2}\int_0^1 y^2 e^{y^3}\,dy = \frac{1}{2}\left[\frac{1}{3}e^{y^3}\right]_0^1 = \frac{1}{2}\cdot\frac{1}{3}\left(e^1 - e^0\right) = \frac{1}{6}(e - 1)$

4. $\int_0^{\pi/2} \int_0^x x\sin y\,dy\,dx = \int_0^{\pi/2} [x(-\cos y)]_{y=0}^{y=x} dx = \int_0^{\pi/2} (-x\cos x + x)\,dx = \int_0^{\pi/2} (x - x\cos x)\,dx$

$= \left[\frac{1}{2}x^2 - (x\sin x + \cos x)\right]_0^{\pi/2}$ (by integrating by parts in the second term)

$= \left(\frac{1}{2}\cdot\frac{\pi^2}{4} - \frac{\pi}{2} - 0\right) - (0 - 0 - 1) = \frac{\pi^2}{8} - \frac{\pi}{2} + 1$

5. $\int_0^1 \int_0^{s^2} \cos(s^3)\,dt\,ds = \int_0^1 \left[t\cos(s^3)\right]_{t=0}^{t=s^2} ds = \int_0^1 s^2\cos(s^3)\,ds = \frac{1}{3}\sin(s^3)\Big]_0^1 = \frac{1}{3}(\sin 1 - \sin 0) = \frac{1}{3}\sin 1$

6. $\int_0^1 \int_0^{e^v} \sqrt{1 + e^v}\,dw\,dv = \int_0^1 \left[w\sqrt{1 + e^v}\,\right]_{w=0}^{w=e^v} dv = \int_0^1 e^v\sqrt{1 + e^v}\,dv = \frac{2}{3}(1 + e^v)^{3/2}\Big]_0^1$

$= \frac{2}{3}(1 + e)^{3/2} - \frac{2}{3}(1 + 1)^{3/2} = \frac{2}{3}(1 + e)^{3/2} - \frac{4}{3}\sqrt{2}$

7. $\iint_D \frac{y}{x^2 + 1}\,dA = \int_0^4 \int_0^{\sqrt{x}} \frac{y}{x^2 + 1}\,dy\,dx = \int_0^4 \left[\frac{1}{x^2 + 1}\cdot\frac{y^2}{2}\right]_{y=0}^{y=\sqrt{x}} dx = \frac{1}{2}\int_0^4 \frac{x}{x^2 + 1}\,dx$

$= \frac{1}{2}\left[\frac{1}{2}\ln\left|x^2 + 1\right|\right]_0^4 = \frac{1}{4}\left[\ln\left(x^2 + 1\right)\right]_0^4 = \frac{1}{4}(\ln 17 - \ln 1) = \frac{1}{4}\ln 17$

8. $\iint_D (2x + y)\,dA = \int_1^2 \int_{y-1}^1 (2x + y)\,dx\,dy = \int_1^2 \left[x^2 + xy\right]_{x=y-1}^{x=1} dy = \int_1^2 \left[1 + y - (y - 1)^2 - y(y - 1)\right] dy$

$= \int_1^2 (-2y^2 + 4y)\,dy = \left[-\frac{2}{3}y^3 + 2y^2\right]_1^2 = \left(-\frac{16}{3} + 8\right) - \left(-\frac{2}{3} + 2\right) = \frac{4}{3}$

9. $\iint_D e^{-y^2}\,dA = \int_0^3 \int_0^y e^{-y^2}\,dx\,dy = \int_0^3 \left[xe^{-y^2}\right]_{x=0}^{x=y} dy = \int_0^3 \left(ye^{-y^2} - 0\right) dy = \int_0^3 ye^{-y^2}\,dy$

$= -\frac{1}{2}e^{-y^2}\Big]_0^3 = -\frac{1}{2}\left(e^{-9} - e^0\right) = \frac{1}{2}\left(1 - e^{-9}\right)$

10. $\iint_D y\sqrt{x^2 - y^2}\,dA = \int_0^2 \int_0^x y\sqrt{x^2 - y^2}\,dy\,dx = \int_0^2 \left[-\frac{1}{3}(x^2 - y^2)^{3/2}\right]_{y=0}^{y=x} dx = \int_0^2 \left[0 + \frac{1}{3}(x^2)^{3/2}\right] dx$

$= \int_0^2 \frac{1}{3}x^3\,dx = \frac{1}{3}\cdot\frac{1}{4}x^4\Big]_0^2 = \frac{1}{12}(16 - 0) = \frac{4}{3}$

11. (a) At the right we sketch an example of a region D that can be described as lying between the graphs of two continuous functions of x (a type I region) but not as lying between graphs of two continuous functions of y (a type II region). The regions shown in Figures 6 and 8 in the text are additional examples.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Now we sketch an example of a region D that can be described as lying between the graphs of two continuous functions of y but not as lying between graphs of two continuous functions of x. The first region shown in Figure 7 is another example.



12. (a) At the right we sketch an example of a region D that can be described as lying between the graphs of two continuous functions of x (a type I region) and also as lying between graphs of two continuous functions of y (a type II region). For additional examples see Figures 9, 11, 12, and 14–16 in the text.



(b) Now we sketch an example of a region D that can't be described as lying between the graphs of two continuous functions of x or between graphs of two continuous functions of y. The region shown in Figure 18 is another example.



13.



As a type I region, D lies between the lower boundary $y = 0$ and the upper boundary $y = x$ for $0 \le x \le 1$, so $D = \{(x, y) \mid 0 \le x \le 1, 0 \le y \le x\}$. If we describe D as a type II region, D lies between the left boundary $x = y$ and the right boundary $x = 1$ for $0 \le y \le 1$, so $D = \{(x, y) \mid 0 \le y \le 1, y \le x \le 1\}$.

Thus $\iint_D x\,dA = \int_0^1 \int_0^x x\,dy\,dx = \int_0^1 \left[xy\right]_{y=0}^{y=x} dx = \int_0^1 x^2\,dx = \frac{1}{3}x^3\big]_0^1 = \frac{1}{3}(1-0) = \frac{1}{3}$ or

$\iint_D x\,dA = \int_0^1 \int_y^1 x\,dx\,dy = \int_0^1 \left[\frac{1}{2}x^2\right]_{x=y}^{x=1} dy = \frac{1}{2}\int_0^1 (1-y^2)\,dy = \frac{1}{2}\left[y - \frac{1}{3}y^3\right]_0^1 = \frac{1}{2}\left[\left(1-\frac{1}{3}\right) - 0\right] = \frac{1}{3}$.

14.



The curves $y = x^2$ and $y = 3x$ intersect at points $(0, 0)$, $(3, 9)$. As a type I region, D is enclosed by the lower boundary $y = x^2$ and the upper boundary $y = 3x$ for $0 \le x \le 3$, so $D = \{(x, y) \mid 0 \le x \le 3, x^2 \le y \le 3x\}$. If we describe D as a type II region, D is enclosed by the left boundary $x = y/3$ and the right boundary $x = \sqrt{y}$ for $0 \le y \le 9$, so $D = \{(x, y) \mid 0 \le y \le 9, y/3 \le x \le \sqrt{y}\}$. Thus

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$$\iint_D xy\,dA = \int_0^3 \int_{x^2}^{3x} xy\,dy\,dx = \int_0^3 \left[x \cdot \tfrac{1}{2}y^2\right]_{y=x^2}^{y=3x} dx = \tfrac{1}{2}\int_0^3 x(9x^2 - x^4)\,dx = \tfrac{1}{2}\int_0^3 (9x^3 - x^5)\,dx$$

$$= \tfrac{1}{2}\left[9 \cdot \tfrac{1}{4}x^4 - \tfrac{1}{6}x^6\right]_0^3 = \tfrac{1}{2}\left[\left(\tfrac{9}{4} \cdot 81 - \tfrac{1}{6} \cdot 729\right) - 0\right] = \tfrac{243}{8}$$

or

$$\iint_D xy\,dA = \int_0^9 \int_{y/3}^{\sqrt{y}} xy\,dx\,dy = \int_0^9 \left[\tfrac{1}{2}x^2 y\right]_{x=y/3}^{x=\sqrt{y}} dy = \tfrac{1}{2}\int_0^9 \left(y - \tfrac{1}{9}y^2\right)y\,dy = \tfrac{1}{2}\int_0^9 \left(y^2 - \tfrac{1}{9}y^3\right) dy$$

$$= \tfrac{1}{2}\left[\tfrac{1}{3}y^3 - \tfrac{1}{9} \cdot \tfrac{1}{4}y^4\right]_0^9 = \tfrac{1}{2}\left[\left(\tfrac{1}{3} \cdot 729 - \tfrac{1}{36} \cdot 6561\right) - 0\right] = \tfrac{243}{8}$$

15.



The curves $y = x - 2$ or $x = y + 2$ and $x = y^2$ intersect when $y + 2 = y^2 \Leftrightarrow y^2 - y - 2 = 0 \Leftrightarrow (y-2)(y+1) = 0 \Leftrightarrow y = -1, y = 2$, so the points of intersection are $(1, -1)$ and $(4, 2)$. If we describe D as a type I region, the upper boundary curve is $y = \sqrt{x}$ but the lower boundary curve consists of two parts, $y = -\sqrt{x}$ for $0 \le x \le 1$ and $y = x - 2$ for $1 \le x \le 4$.

Thus $D = \{(x, y) \mid 0 \le x \le 1,\ -\sqrt{x} \le y \le \sqrt{x}\,\} \cup \{(x, y) \mid 1 \le x \le 4,\ x - 2 \le y \le \sqrt{x}\,\}$ and $\iint_D y\,dA = \int_0^1 \int_{-\sqrt{x}}^{\sqrt{x}} y\,dy\,dx + \int_1^4 \int_{x-2}^{\sqrt{x}} y\,dy\,dx$. If we describe D as a type II region, D is enclosed by the left boundary $x = y^2$ and the right boundary $x = y + 2$ for $-1 \le y \le 2$, so $D = \{(x, y) \mid -1 \le y \le 2,\ y^2 \le x \le y + 2\}$ and $\iint_D y\,dA = \int_{-1}^{2} \int_{y^2}^{y+2} y\,dx\,dy$. In either case, the resulting iterated integrals are not difficult to evaluate but the region D is more simply described as a type II region, giving one iterated integral rather than a sum of two, so we evaluate the latter integral:

$$\iint_D y\,dA = \int_{-1}^{2} \int_{y^2}^{y+2} y\,dx\,dy = \int_{-1}^{2} \left[xy\right]_{x=y^2}^{x=y+2} dy = \int_{-1}^{2} (y + 2 - y^2)y\,dy = \int_{-1}^{2} (y^2 + 2y - y^3)\,dy$$

$$= \left[\tfrac{1}{3}y^3 + y^2 - \tfrac{1}{4}y^4\right]_{-1}^{2} = \left(\tfrac{8}{3} + 4 - 4\right) - \left(-\tfrac{1}{3} + 1 - \tfrac{1}{4}\right) = \tfrac{9}{4}$$

16.



As a type I region, $D = \{(x, y) \mid 0 \le x \le 4,\ x \le y \le 4\}$ and $\iint_D y^2 e^{xy}\,dA = \int_0^4 \int_x^4 y^2 e^{xy}\,dy\,dx$. As a type II region, $D = \{(x, y) \mid 0 \le y \le 4,\ 0 \le x \le y\}$ and $\iint_D y^2 e^{xy}\,dA = \int_0^4 \int_0^y y^2 e^{xy}\,dx\,dy$. Evaluating $\int y^2 e^{xy}\,dy$ requires integration by parts whereas $\int y^2 e^{xy}\,dx$ does not, so the iterated integral corresponding to D as a type II region appears easier to evaluate.

$$\iint_D y^2 e^{xy}\,dA = \int_0^4 \int_0^y y^2 e^{xy}\,dx\,dy = \int_0^4 \left[ye^{xy}\right]_{x=0}^{x=y} dy = \int_0^4 \left(ye^{y^2} - y\right) dy$$

$$= \left[\tfrac{1}{2}e^{y^2} - \tfrac{1}{2}y^2\right]_0^4 = \left(\tfrac{1}{2}e^{16} - 8\right) - \left(\tfrac{1}{2} - 0\right) = \tfrac{1}{2}e^{16} - \tfrac{17}{2}$$

17.



$$\int_0^1 \int_0^{x^2} x\cos y\,dy\,dx = \int_0^1 \left[x\sin y\right]_{y=0}^{y=x^2} dx = \int_0^1 x\sin x^2\,dx$$

$$= -\tfrac{1}{2}\cos x^2\Big]_0^1 = -\tfrac{1}{2}(\cos 1 - \cos 0) = \tfrac{1}{2}(1 - \cos 1)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18.



$$\iint_D (x^2+2y)\,dA = \int_0^1 \int_{x^3}^{x} (x^2+2y)\,dy\,dx = \int_0^1 \left[x^2y+y^2\right]_{y=x^3}^{y=x}\,dx$$

$$= \int_0^1 (x^3+x^2-x^5-x^6)\,dx = \left[\tfrac{1}{4}x^4+\tfrac{1}{3}x^3-\tfrac{1}{6}x^6-\tfrac{1}{7}x^7\right]_0^1$$

$$= \tfrac{1}{4}+\tfrac{1}{3}-\tfrac{1}{6}-\tfrac{1}{7} = \tfrac{23}{84}$$

19.



$$\iint_D y^2\,dA = \int_1^2 \int_{y-1}^{7-3y} y^2\,dx\,dy = \int_1^2 \left[xy^2\right]_{x=y-1}^{x=7-3y}\,dy$$

$$= \int_1^2 [(7-3y)-(y-1)]\,y^2\,dy = \int_1^2 (8y^2-4y^3)\,dy$$

$$= \left[\tfrac{8}{3}y^3-y^4\right]_1^2 = \tfrac{64}{3}-16-\tfrac{8}{3}+1 = \tfrac{11}{3}$$

20.



$$\iint_D xy\,dA = \int_0^1 \int_0^{\sqrt{1-x^2}} xy\,dy\,dx$$

$$= \int_0^1 \left[\tfrac{1}{2}xy^2\right]_{y=0}^{y=\sqrt{1-x^2}}\,dx = \int_0^1 \tfrac{1}{2}x(1-x^2)\,dx$$

$$= \tfrac{1}{2}\int_0^1 (x-x^3)\,dx = \tfrac{1}{2}\left[\tfrac{1}{2}x^2-\tfrac{1}{4}x^4\right]_0^1$$

$$= \tfrac{1}{2}\left(\tfrac{1}{2}-\tfrac{1}{4}-0\right) = \tfrac{1}{8}$$

21.



$$\int_{-2}^{2} \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} (2x-y)\,dy\,dx$$

$$= \int_{-2}^{2} \left[2xy-\tfrac{1}{2}y^2\right]_{y=-\sqrt{4-x^2}}^{y=\sqrt{4-x^2}}\,dx$$

$$= \int_{-2}^{2} \left[2x\sqrt{4-x^2}-\tfrac{1}{2}(4-x^2)+2x\sqrt{4-x^2}+\tfrac{1}{2}(4-x^2)\right]\,dx$$

$$= \int_{-2}^{2} 4x\sqrt{4-x^2}\,dx = -\tfrac{4}{3}(4-x^2)^{3/2}\Big]_{-2}^{2} = 0$$

[Or, note that $4x\sqrt{4-x^2}$ is an odd function, so $\int_{-2}^{2} 4x\sqrt{4-x^2}\,dx = 0$.]

22.



$$\iint_D y\,dA = \int_0^1 \int_y^{4-3y} y\,dx\,dy$$

$$= \int_0^1 [xy]_{x=y}^{x=4-3y}\,dy = \int_0^1 (4y-3y^2-y^2)\,dy$$

$$= \int_0^1 (4y-4y^2)\,dy = \left[2y^2-\tfrac{4}{3}y^3\right]_0^1 = 2-\tfrac{4}{3}-0 = \tfrac{2}{3}$$

23.



$$V = \int_0^1 \int_{x^2}^{\sqrt{x}} (3x+2y)\,dy\,dx = \int_0^1 \left[3xy+y^2\right]_{y=x^2}^{y=\sqrt{x}}\,dx$$

$$= \int_0^1 \left[(3x\sqrt{x}+x)-(3x^3+x^4)\right]\,dx = \int_0^1 (3x^{3/2}+x-3x^3-x^4)\,dx$$

$$= \left[3\cdot\tfrac{2}{5}x^{5/2}+\tfrac{1}{2}x^2-\tfrac{3}{4}x^4-\tfrac{1}{5}x^5\right]_0^1 = \tfrac{6}{5}+\tfrac{1}{2}-\tfrac{3}{4}-\tfrac{1}{5}-0 = \tfrac{3}{4}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

24.



$$V = \int_{-2}^{2}\int_{y^2}^{4}(1+x^2y^2)\,dx\,dy$$

$$= \int_{-2}^{2}\left[x+\tfrac{1}{3}x^3y^2\right]_{x=y^2}^{x=4}\,dy = \int_{-2}^{2}(4+\tfrac{61}{3}y^2-\tfrac{1}{3}y^8)\,dy$$

$$= \left[4y+\tfrac{61}{9}y^3-\tfrac{1}{27}y^9\right]_{-2}^{2} = 8+\tfrac{488}{9}-\tfrac{512}{27}+8+\tfrac{488}{9}-\tfrac{512}{27} = \tfrac{2336}{27}$$

25.



$$V = \int_1^2\int_1^{7-3y} xy\,dx\,dy = \int_1^2\left[\tfrac{1}{2}x^2y\right]_{x=1}^{x=7-3y}\,dy$$

$$= \tfrac{1}{2}\int_1^2 y\left[(7-3y)^2-1\right]dy = \tfrac{1}{2}\int_1^2(48y-42y^2+9y^3)\,dy$$

$$= \tfrac{1}{2}\left[24y^2-14y^3+\tfrac{9}{4}y^4\right]_1^2 = \tfrac{31}{8}$$

26.



$$V = \int_0^2\int_0^{2-x}(x^2+y^2+1)\,dy\,dx = \int_0^2\left[x^2y+\tfrac{1}{3}y^3+y\right]_{y=0}^{y=2-x}\,dx$$

$$= \int_0^2\left[x^2(2-x)+\tfrac{1}{3}(2-x)^3+(2-x)-0\right]dx$$

$$= \int_0^2\left(-\tfrac{4}{3}x^3+4x^2-5x+\tfrac{14}{3}\right)dx = \left[-\tfrac{1}{3}x^4+\tfrac{4}{3}x^3-\tfrac{5}{2}x^2+\tfrac{14}{3}x\right]_0^2$$

$$= -\tfrac{16}{3}+\tfrac{32}{3}-10+\tfrac{28}{3}-0 = \tfrac{14}{3}$$

27.



$$V = \int_0^2\int_0^{4-2x}(4-2x-y)\,dy\,dx = \int_0^2\left[4y-2xy-\tfrac{1}{2}y^2\right]_{y=0}^{y=4-2x}\,dx$$

$$= \int_0^2\left[4(4-2x)-2x(4-2x)-\tfrac{1}{2}(4-2x)^2-0\right]dx$$

$$= \int_0^2\left(2x^2-8x+8\right)dx = \left[\tfrac{2}{3}x^3-4x^2+8x\right]_0^2 = \tfrac{16}{3}-16+16-0 = \tfrac{16}{3}$$

28.



$$V = \int_0^1\int_x^{2-x} x\,dy\,dx$$

$$= \int_0^1\left[xy\right]_{y=x}^{y=2-x}dx = \int_0^1(2x-2x^2)\,dx$$

$$= \left[x^2-\tfrac{2}{3}x^3\right]_0^1 = \tfrac{1}{3}$$

29.



$$V = \int_{-2}^{2}\int_{x^2}^{4} x^2\,dy\,dx$$

$$= \int_{-2}^{2}\left[x^2y\right]_{y=x^2}^{y=4}dx = \int_{-2}^{2}(4x^2-x^4)\,dx$$

$$= \left[\tfrac{4}{3}x^3-\tfrac{1}{5}x^5\right]_{-2}^{2} = \tfrac{32}{3}-\tfrac{32}{5}+\tfrac{32}{3}-\tfrac{32}{5} = \tfrac{128}{15}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30.



$$V = \int_0^2 \int_0^{2y} \sqrt{4 - y^2}\, dx\, dy = \int_0^2 \left[x\sqrt{4 - y^2} \right]_{x=0}^{x=2y} dy$$

$$= \int_0^2 2y\sqrt{4 - y^2}\, dy = \left[-\tfrac{2}{3}(4 - y^2)^{3/2} \right]_0^2 = 0 + \tfrac{16}{3} = \tfrac{16}{3}$$

31.



$$V = \int_0^1 \int_0^{\sqrt{1-x^2}} y\, dy\, dx = \int_0^1 \left[\frac{y^2}{2} \right]_{y=0}^{y=\sqrt{1-x^2}} dx$$

$$= \int_0^1 \frac{1 - x^2}{2}\, dx = \tfrac{1}{2}\left[x - \tfrac{1}{3}x^3 \right]_0^1 = \tfrac{1}{3}$$

32.



By symmetry, the desired volume V is 8 times the volume V_1 in the first octant. Now

$$V_1 = \int_0^r \int_0^{\sqrt{r^2-y^2}} \sqrt{r^2 - y^2}\, dx\, dy = \int_0^r \left[x\sqrt{r^2 - y^2} \right]_{x=0}^{x=\sqrt{r^2-y^2}} dy$$

$$= \int_0^r (r^2 - y^2)\, dy = \left[r^2 y - \tfrac{1}{3}y^3 \right]_0^r = \tfrac{2}{3}r^3$$

Thus $V = \frac{16}{3}r^3$.

33.



From the graph, it appears that the two curves intersect at $x = 0$ and at $x \approx 1.213$. Thus the desired integral is

$$\iint_D x\, dA \approx \int_0^{1.213} \int_{x^4}^{3x - x^2} x\, dy\, dx = \int_0^{1.213} \left[xy \right]_{y=x^4}^{y=3x-x^2} dx$$

$$= \int_0^{1.213} (3x^2 - x^3 - x^5)\, dx = \left[x^3 - \tfrac{1}{4}x^4 - \tfrac{1}{6}x^6 \right]_0^{1.213}$$

$$\approx 0.713$$

34.





The desired solid is shown in the first graph. From the second graph, we estimate that $y = \cos x$ intersects $y = x$ at $x \approx 0.7391$. Therefore the volume of the solid is

$$V \approx \int_0^{0.7391} \int_x^{\cos x} x\, dy\, dx = \int_0^{0.7391} \left[xy \right]_{y=x}^{y=\cos x} dx$$

$$= \int_0^{0.7391} (x\cos x - x^2)\, dx = \left[\cos x + x\sin x - \tfrac{1}{3}x^3 \right]_0^{0.7391} \approx 0.1024$$

Note: There is a different solid which can also be construed to satisfy the conditions stated in the exercise. This is the solid bounded by all of the given surfaces, as well as the plane $y = 0$. In case you calculated the volume of this solid and want to check your work, its volume is $V \approx \int_0^{0.7391} \int_0^x x\, dy\, dx + \int_{0.7391}^{\pi/2} \int_0^{\cos x} x\, dy\, dx \approx 0.4684$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

35. The region of integration is bounded by the curves $y = 1 - x^2$ and $y = x^2 - 1$ which intersect at $(\pm 1, 0)$ with $1 - x^2 \geq x^2 - 1$ on $[-1, 1]$. Within this region, the plane $z = 2x + 2y + 10$ is above the plane $z = 2 - x - y$, so

$$\begin{aligned} V &= \int_{-1}^{1} \int_{x^2-1}^{1-x^2} (2x + 2y + 10)\, dy\, dx - \int_{-1}^{1} \int_{x^2-1}^{1-x^2} (2 - x - y)\, dy\, dx \\ &= \int_{-1}^{1} \int_{x^2-1}^{1-x^2} (2x + 2y + 10 - (2 - x - y))\, dy\, dx \\ &= \int_{-1}^{1} \int_{x^2-1}^{1-x^2} (3x + 3y + 8)\, dy\, dx = \int_{-1}^{1} \left[3xy + \tfrac{3}{2}y^2 + 8y\right]_{y=x^2-1}^{y=1-x^2} dx \\ &= \int_{-1}^{1} \left[3x(1 - x^2) + \tfrac{3}{2}(1 - x^2)^2 + 8(1 - x^2) - 3x(x^2 - 1) - \tfrac{3}{2}(x^2 - 1)^2 - 8(x^2 - 1)\right] dx \\ &= \int_{-1}^{1} (-6x^3 - 16x^2 + 6x + 16)\, dx = \left[-\tfrac{3}{2}x^4 - \tfrac{16}{3}x^3 + 3x^2 + 16x\right]_{-1}^{1} \\ &= -\tfrac{3}{2} - \tfrac{16}{3} + 3 + 16 + \tfrac{3}{2} - \tfrac{16}{3} - 3 + 16 = \tfrac{64}{3} \end{aligned}$$

36. The two planes intersect in the line $y = 1$, $z = 3$, so the region of integration is the plane region enclosed by the parabola $y = x^2$ and the line $y = 1$. We have $2 + y \geq 3y$ for $0 \leq y \leq 1$, so the solid region is bounded above by $z = 2 + y$ and bounded below by $z = 3y$.



$$\begin{aligned} V &= \int_{-1}^{1} \int_{x^2}^{1} (2 + y)\, dy\, dx - \int_{-1}^{1} \int_{x^2}^{1} (3y)\, dy\, dx = \int_{-1}^{1} \int_{x^2}^{1} (2 + y - 3y)\, dy\, dx \\ &= \int_{-1}^{1} \int_{x^2}^{1} (2 - 2y)\, dy\, dx = \int_{-1}^{1} \left[2y - y^2\right]_{y=x^2}^{y=1} dx \\ &= \int_{-1}^{1} (1 - 2x^2 + x^4)\, dx = x - \tfrac{2}{3}x^3 + \tfrac{1}{5}x^5\Big]_{-1}^{1} = \tfrac{16}{15} \end{aligned}$$

37. The region of integration is bounded by the curves $y = x^2$ and $y = 1 - x^2$ which intersect at $\left(\pm\frac{1}{\sqrt{2}}, \frac{1}{2}\right)$.

The solid lies under the graph of $z = 3$ and above the graph of $z = y$, so its volume is



$$\begin{aligned} V &= \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \int_{x^2}^{1-x^2} 3\, dy\, dx - \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \int_{x^2}^{1-x^2} y\, dy\, dx = \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \int_{x^2}^{1-x^2} (3 - y)\, dy\, dx \\ &= \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \left[3y - \tfrac{1}{2}y^2\right]_{y=x^2}^{y=1-x^2} dx = \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \left[\left(3(1 - x^2) - \tfrac{1}{2}(1 - x^2)^2\right) - \left(3x^2 - \tfrac{1}{2}(x^2)^2\right)\right] dx \\ &= \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \left(\tfrac{5}{2} - 5x^2\right) dx = \left[\tfrac{5}{2}x - \tfrac{5}{3}x^3\right]_{-1/\sqrt{2}}^{1/\sqrt{2}} \quad \left(\tfrac{5}{2\sqrt{2}} - \tfrac{5}{6\sqrt{2}}\right) - \left(-\tfrac{5}{2\sqrt{2}} + \tfrac{5}{6\sqrt{2}}\right) \\ &= \tfrac{10}{3\sqrt{2}} \text{ or } \tfrac{5\sqrt{2}}{3} \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. The region of integration is the portion of the first quadrant bounded by the axes and the curve $y = \sqrt{4-x^2}$. The solid lies under the graph of $z = x + y$ and above the graph of $z = xy$, so its volume is

$$V = \int_0^2 \int_0^{\sqrt{4-x^2}} (x+y)\,dy\,dx - \int_0^2 \int_0^{\sqrt{4-x^2}} xy\,dy\,dx = \int_0^2 \int_0^{\sqrt{4-x^2}} (x+y-xy)\,dy\,dx$$

$$= \int_0^2 \left[xy + \tfrac{1}{2}y^2 - \tfrac{1}{2}xy^2\right]_{y=0}^{y=\sqrt{4-x^2}} dx = \int_0^2 \left[x\sqrt{4-x^2} + \tfrac{1}{2}(4-x^2) - \tfrac{1}{2}x(4-x^2) - 0\right] dx$$

$$= \int_0^2 \left(x\sqrt{4-x^2} + 2 - \tfrac{1}{2}x^2 - 2x + \tfrac{1}{2}x^3\right) dx = \left[-\tfrac{1}{3}(4-x^2)^{3/2} + 2x - \tfrac{1}{6}x^3 - x^2 + \tfrac{1}{8}x^4\right]_0^2$$

$$= \left(4 - \tfrac{4}{3} - 4 + 2\right) - \left(-\tfrac{1}{3}\cdot 4^{3/2}\right) = \tfrac{2}{3} + \tfrac{8}{3} = \tfrac{10}{3}$$

39. The solid lies below the plane $z = 1 - x - y$ or $x + y + z = 1$ and above the region $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le 1 - x\}$ in the xy-plane. The solid is a tetrahedron.



40. The solid lies below the plane $z = 1 - x$ and above the region $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le 1 - x^2\}$ in the xy-plane.



41. The two bounding curves $y = x^3 - x$ and $y = x^2 + x$ intersect at the origin and at $x = 2$, with $x^2 + x > x^3 - x$ on $(0, 2)$. Using a CAS, we find that the volume of the solid is

$$V = \int_0^2 \int_{x^3-x}^{x^2+x} (x^3y^4 + xy^2)\,dy\,dx = \frac{13{,}984{,}735{,}616}{14{,}549{,}535}$$

42. For $|x| \le 1$ and $|y| \le 1$, $2x^2 + y^2 < 8 - x^2 - 2y^2$. Also, the cylinder is described by the inequalities $-1 \le x \le 1$, $-\sqrt{1-x^2} \le y \le \sqrt{1-x^2}$. So the volume is given by

$$V = \int_{-1}^{1} \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \left[(8 - x^2 - 2y^2) - (2x^2 + y^2)\right] dy\,dx = \frac{13\pi}{2} \qquad \text{[using a CAS]}$$

43. The two surfaces intersect in the circle $x^2 + y^2 = 1$, $z = 0$ and the region of integration is the disk D: $x^2 + y^2 \le 1$. Using a CAS, the volume is $\iint_D (1 - x^2 - y^2)\,dA = \int_{-1}^{1} \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} (1 - x^2 - y^2)\,dy\,dx = \frac{\pi}{2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

44. The projection onto the xy-plane of the intersection of the two surfaces is the circle $x^2 + y^2 = 2y \Rightarrow$ $x^2 + y^2 - 2y = 0 \Rightarrow x^2 + (y-1)^2 = 1$, so the region of integration is given by $-1 \le x \le 1$, $1 - \sqrt{1-x^2} \le y \le 1 + \sqrt{1-x^2}$. In this region, $2y \ge x^2 + y^2$ so, using a CAS, the volume is

$$V = \int_{-1}^{1} \int_{1-\sqrt{1-x^2}}^{1+\sqrt{1-x^2}} [2y - (x^2 + y^2)]\, dy\, dx = \frac{\pi}{2}$$

45.



Because the region of integration is

$$D = \{(x,y) \mid 0 \le x \le y, 0 \le y \le 1\} = \{(x,y) \mid x \le y \le 1, 0 \le x \le 1\}$$

we have $\int_0^1 \int_0^y f(x,y)\, dx\, dy = \iint_D f(x,y)\, dA = \int_0^1 \int_x^1 f(x,y)\, dy\, dx$.

46.



Because the region of integration is

$$\begin{aligned} D &= \{(x,y) \mid x^2 \le y \le 4, 0 \le x \le 2\} \\ &= \{(x,y) \mid 0 \le x \le \sqrt{y}, 0 \le y \le 4\} \end{aligned}$$

we have $\int_0^2 \int_{x^2}^4 f(x,y)\, dy\, dx = \iint_D f(x,y)\, dA = \int_0^4 \int_0^{\sqrt{y}} f(x,y)\, dx\, dy$.

47.



Because the region of integration is

$$\begin{aligned} D &= \{(x,y) \mid 0 \le y \le \cos x, 0 \le x \le \pi/2\} \\ &= \{(x,y) \mid 0 \le x \le \cos^{-1} y, 0 \le y \le 1\} \end{aligned}$$

we have

$\int_0^{\pi/2} \int_0^{\cos x} f(x,y)\, dy\, dx = \iint_D f(x,y)\, dA = \int_0^1 \int_0^{\cos^{-1} y} f(x,y)\, dx\, dy$.

48.



Because the region of integration is

$$\begin{aligned} D &= \left\{(x,y) \mid 0 \le x \le \sqrt{4-y^2}, -2 \le y \le 2\right\} \\ &= \left\{(x,y) \mid -\sqrt{4-x^2} \le y \le \sqrt{4-x^2}, 0 \le x \le 2\right\} \end{aligned}$$

we have

$\int_{-2}^2 \int_0^{\sqrt{4-y^2}} f(x,y)\, dx\, dy = \iint_D f(x,y)\, dA = \int_0^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} f(x,y)\, dy\, dx$.

49.



Because the region of integration is

$$D = \{(x,y) \mid 0 \le y \le \ln x, 1 \le x \le 2\} = \{(x,y) \mid e^y \le x \le 2, 0 \le y \le \ln 2\}$$

we have

$$\int_1^2 \int_0^{\ln x} f(x,y)\, dy\, dx = \iint_D f(x,y)\, dA = \int_0^{\ln 2} \int_{e^y}^2 f(x,y)\, dx\, dy$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50.



Because the region of integration is

$$D = \{(x, y) \mid \arctan x \le y \le \tfrac{\pi}{4}, 0 \le x \le 1\}$$
$$= \{(x, y) \mid 0 \le x \le \tan y, 0 \le y \le \tfrac{\pi}{4}\}$$

we have

$$\int_0^1 \int_{\arctan x}^{\pi/4} f(x, y)\, dy\, dx = \iint_D f(x, y)\, dA = \int_0^{\pi/4} \int_0^{\tan y} f(x, y)\, dx\, dy$$

51.



$$\int_0^1 \int_{3y}^3 e^{x^2}\, dx\, dy = \int_0^3 \int_0^{x/3} e^{x^2}\, dy\, dx = \int_0^3 \left[e^{x^2} y\right]_{y=0}^{y=x/3} dx$$
$$= \int_0^3 \left(\frac{x}{3}\right) e^{x^2}\, dx = \tfrac{1}{6}\, e^{x^2}\Big]_0^3 = \frac{e^9 - 1}{6}$$

52.



$$\int_0^1 \int_{x^2}^1 \sqrt{y} \sin y\, dy\, dx = \int_0^1 \int_0^{\sqrt{y}} \sqrt{y} \sin y\, dx\, dy = \int_0^1 \sqrt{y} \sin y\, [x]_{x=0}^{x=\sqrt{y}}\, dy$$
$$= \int_0^1 (\sqrt{y} \sin y)(\sqrt{y} - 0)\, dy = \int_0^1 y \sin y\, dy$$
$$= -y \cos y]_0^1 + \int_0^1 \cos y\, dy$$

[by integrating by parts with $u = y$, $dv = \sin y\, dy$]

$$= [-y \cos y + \sin y]_0^1 = -\cos 1 + \sin 1 - 0 = \sin 1 - \cos 1$$

53.



$$\int_0^1 \int_{\sqrt{x}}^1 \sqrt{y^3 + 1}\, dy\, dx = \int_0^1 \int_0^{y^2} \sqrt{y^3 + 1}\, dx\, dy = \int_0^1 \sqrt{y^3 + 1}\, [x]_{x=0}^{x=y^2}\, dy$$
$$= \int_0^1 y^2 \sqrt{y^3 + 1}\, dy = \tfrac{2}{9} \left(y^3 + 1\right)^{3/2}\Big]_0^1$$
$$= \tfrac{2}{9} \left(2^{3/2} - 1^{3/2}\right) = \tfrac{2}{9} \left(2\sqrt{2} - 1\right)$$

54.



$$\int_0^2 \int_{y/2}^1 y \cos(x^3 - 1)\, dx\, dy = \int_0^1 \int_0^{2x} y \cos(x^3 - 1)\, dy\, dx$$
$$= \int_0^1 \cos(x^3 - 1) \left[\tfrac{1}{2} y^2\right]_{y=0}^{y=2x} dx$$
$$= \int_0^1 2x^2 \cos(x^3 - 1)\, dx = \tfrac{2}{3} \sin(x^3 - 1)\Big]_0^1$$
$$= \tfrac{2}{3} [0 - \sin(-1)] = -\tfrac{2}{3} \sin(-1) = \tfrac{2}{3} \sin 1$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

55.



$$\int_0^1 \int_{\arcsin y}^{\pi/2} \cos x \sqrt{1+\cos^2 x}\, dx\, dy$$
$$= \int_0^{\pi/2} \int_0^{\sin x} \cos x \sqrt{1+\cos^2 x}\, dy\, dx$$
$$= \int_0^{\pi/2} \cos x \sqrt{1+\cos^2 x}\, \big[y\big]_{y=0}^{y=\sin x}\, dx$$
$$= \int_0^{\pi/2} \cos x \sqrt{1+\cos^2 x}\, \sin x\, dx \qquad \left[\begin{array}{c}\text{Let } u=\cos x,\ du=-\sin x\, dx,\\ dx=du/(-\sin x)\end{array}\right]$$
$$= \int_1^0 -u\sqrt{1+u^2}\, du = -\tfrac{1}{3}\left(1+u^2\right)^{3/2}\Big]_1^0$$
$$= \tfrac{1}{3}\left(\sqrt{8}-1\right) = \tfrac{1}{3}\left(2\sqrt{2}-1\right)$$

56.



$$\int_0^8 \int_{\sqrt[3]{y}}^2 e^{x^4}\, dx\, dy = \int_0^2 \int_0^{x^3} e^{x^4}\, dy\, dx$$
$$= \int_0^2 e^{x^4}\Big[y\Big]_{y=0}^{y=x^3}\, dx = \int_0^2 x^3 e^{x^4}\, dx$$
$$= \tfrac{1}{4}e^{x^4}\Big]_0^2 = \tfrac{1}{4}(e^{16}-1)$$

57. $D = \{(x,y) \mid 0 \le x \le 1,\ -x+1 \le y \le 1\} \cup \{(x,y) \mid -1 \le x \le 0,\ x+1 \le y \le 1\}$
$\cup \{(x,y) \mid 0 \le x \le 1,\ -1 \le y \le x-1\} \cup \{(x,y) \mid -1 \le x \le 0,\ -1 \le y \le -x-1\}$, all type I.

$$\iint_D x^2\, dA = \int_0^1 \int_{1-x}^1 x^2\, dy\, dx + \int_{-1}^0 \int_{x+1}^1 x^2\, dy\, dx + \int_0^1 \int_{-1}^{x-1} x^2\, dy\, dx + \int_{-1}^0 \int_{-1}^{-x-1} x^2\, dy\, dx$$
$$= 4\int_0^1 \int_{1-x}^1 x^2\, dy\, dx \qquad \text{[by symmetry of the regions and because } f(x,y)=x^2 \ge 0\text{]}$$
$$= 4\int_0^1 x^3\, dx = 4\left[\tfrac{1}{4}x^4\right]_0^1 = 1$$

58. $D = \left\{(x,y) \mid -1 \le y \le 0,\ -1 \le x \le y-y^3\right\} \cup \left\{(x,y) \mid 0 \le y \le 1,\ \sqrt{y}-1 \le x \le y-y^3\right\}$, both type II.

$$\iint_D y\, dA = \int_{-1}^0 \int_{-1}^{y-y^3} y\, dx\, dy + \int_0^1 \int_{\sqrt{y}-1}^{y-y^3} y\, dx\, dy = \int_{-1}^0 \big[xy\big]_{x=-1}^{x=y-y^3}\, dy + \int_0^1 \big[xy\big]_{x=\sqrt{y}-1}^{x=y-y^3}\, dy$$
$$= \int_{-1}^0 (y^2 - y^4 + y)\, dy + \int_0^1 (y^2 - y^4 - y^{3/2} + y)\, dy$$
$$= \left[\tfrac{1}{3}y^3 - \tfrac{1}{5}y^5 + \tfrac{1}{2}y^2\right]_{-1}^0 + \left[\tfrac{1}{3}y^3 - \tfrac{1}{5}y^5 - \tfrac{2}{5}y^{5/2} + \tfrac{1}{2}y^2\right]_0^1$$
$$= \left(0 - \tfrac{11}{30}\right) + \left(\tfrac{7}{30} - 0\right) = \tfrac{2}{15}$$

59. Since $x^2+y^2 \le 1$ on S, we must have $0 \le x^2 \le 1$ and $0 \le y^2 \le 1$, so $0 \le x^2y^2 \le 1 \quad \Rightarrow \quad 3 \le 4-x^2y^2 \le 4 \quad \Rightarrow$
$\sqrt{3} \le \sqrt{4-x^2y^2} \le 2$. Here we have $A(S) = \frac{1}{2}\pi(1)^2 = \frac{\pi}{2}$, so by Property 11,
$\sqrt{3}A(S) \le \iint_S \sqrt{4-x^2y^2}\, dA \le 2A(S) \quad \Rightarrow \quad \frac{\sqrt{3}}{2}\pi \le \iint_S \sqrt{4-x^2y^2}\, dA \le \pi$ or we can say
$2.720 < \iint_S \sqrt{4-x^2y^2}\, dA < 3.142$. (We have rounded the lower bound down and the upper bound up to preserve the inequalities.)

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. T is the triangle with vertices $(0,0)$, $(1,0)$, and $(1,2)$ so $A(T) = \frac{1}{2}(1)(2) = 1$. We have $0 \le \sin^4(x+y) \le 1$ for all x, y, and Property 11 gives $0 \cdot A(T) \le \iint_T \sin^4(x+y)\,dA \le 1 \cdot A(T) \quad \Rightarrow \quad 0 \le \iint_T \sin^4(x+y)\,dA \le 1$.

61. The average value of a function f of two variables defined on a rectangle R was defined in Section 15.1 as $f_{\text{ave}} = \frac{1}{A(R)} \iint_R f(x,y)\,dA$. Extending this definition to general regions D, we have $f_{\text{ave}} = \frac{1}{A(D)} \iint_D f(x,y)\,dA$.



Here $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le 3x\}$, so $A(D) = \frac{1}{2}(1)(3) = \frac{3}{2}$ and

$$f_{\text{ave}} = \frac{1}{A(D)} \iint_D f(x,y)\,dA = \frac{1}{3/2} \int_0^1 \int_0^{3x} xy\,dy\,dx$$
$$= \frac{2}{3} \int_0^1 \left[\tfrac{1}{2}xy^2\right]_{y=0}^{y=3x} dx = \frac{1}{3} \int_0^1 9x^3\,dx = \tfrac{3}{4}x^4\Big]_0^1 = \frac{3}{4}$$

62. Here $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le x^2\}$, so $A(D) = \int_0^1 x^2\,dx = \frac{1}{3}x^3\Big]_0^1 = \frac{1}{3}$ and



$$f_{\text{ave}} = \frac{1}{A(D)} \iint_D f(x,y)\,dA = \frac{1}{1/3} \int_0^1 \int_0^{x^2} x \sin y\,dy\,dx$$
$$= 3 \int_0^1 \left[-x\cos y\right]_{y=0}^{y=x^2} dx$$
$$= 3 \int_0^1 \left[x - x\cos(x^2)\right] dx = 3\left[\tfrac{1}{2}x^2 - \tfrac{1}{2}\sin(x^2)\right]_0^1$$
$$= 3\left(\tfrac{1}{2} - \tfrac{1}{2}\sin 1 - 0\right) = \tfrac{3}{2}(1 - \sin 1)$$

63. Since $m \le f(x,y) \le M$, $\iint_D m\,dA \le \iint_D f(x,y)\,dA \le \iint_D M\,dA$ by (8) $\Rightarrow$ $m \iint_D 1\,dA \le \iint_D f(x,y)\,dA \le M \iint_D 1\,dA$ by (7) $\Rightarrow$ $mA(D) \le \iint_D f(x,y)\,dA \le MA(D)$ by (10).

64.



$$\iint_D f(x,y)\,dA = \int_0^1 \int_0^{2y} f(x,y)\,dx\,dy + \int_1^3 \int_0^{3-y} f(x,y)\,dx\,dy$$
$$= \int_0^2 \int_{x/2}^{3-x} f(x,y)\,dy\,dx$$

65.



First we can write $\iint_D (x+2)\,dA = \iint_D x\,dA + \iint_D 2\,dA$. But $f(x,y) = x$ is an odd function with respect to x [that is, $f(-x,y) = -f(x,y)$] and D is symmetric with respect to x. Consequently, the volume above D and below the graph of f is the same as the volume below D and above the graph of f, so $\iint_D x\,dA = 0$. Also, $\iint_D 2\,dA = 2 \cdot A(D) = 2 \cdot \frac{1}{2}\pi(3)^2 = 9\pi$ since D is a half disk of radius 3. Thus $\iint_D (x+2)\,dA = 0 + 9\pi = 9\pi$.

66. The graph of $f(x,y) = \sqrt{R^2 - x^2 - y^2}$ is the top half of the sphere $x^2 + y^2 + z^2 = R^2$, centered at the origin with radius R, and D is the disk in the xy-plane also centered at the origin with radius R. Thus $\iint_D \sqrt{R^2 - x^2 - y^2}\,dA$ represents the volume of a half ball of radius R which is $\frac{1}{2} \cdot \frac{4}{3}\pi R^3 = \frac{2}{3}\pi R^3$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

67. We can write $\iint_D (2x + 3y)\,dA = \iint_D 2x\,dA + \iint_D 3y\,dA$. $\iint_D 2x\,dA$ represents the volume of the solid lying under the plane $z = 2x$ and above the rectangle D. This solid region is a triangular cylinder with length b and whose cross-section is a triangle with width a and height $2a$. (See the first figure.)





Thus its volume is $\frac{1}{2} \cdot a \cdot 2a \cdot b = a^2 b$. Similarly, $\iint_D 3y\,dA$ represents the volume of a triangular cylinder with length a, triangular cross-section with width b and height $3b$, and volume $\frac{1}{2} \cdot b \cdot 3b \cdot a = \frac{3}{2}ab^2$. (See the second figure.) Thus

$$\iint_D (2x + 3y)\,dA = a^2 b + \tfrac{3}{2}ab^2$$

68.



In the first quadrant, x and y are positive and the boundary of D is $x + y = 1$. But D is symmetric with respect to both axes because of the absolute values, so the region of integration is the square shown at the left. To evaluate the double integral, we first write $\iint_D (2 + x^2 y^3 - y^2 \sin x)\,dA = \iint_D 2\,dA + \iint_D x^2 y^3\,dA - \iint_D y^2 \sin x\,dA$. Now $f(x, y) = x^2 y^3$ is odd with respect to y [that is, $f(x, -y) = -f(x, y)$] and D is symmetric with respect to y, so $\iint_D x^2 y^3\,dA = 0$.

Similarly, $g(x, y) = y^2 \sin x$ is odd with respect to x [since $g(-x, y) = -g(x, y)$] and D is symmetric with respect to x, so $\iint_D y^2 \sin x\,dA = 0$. D is a square with side length $\sqrt{2}$, so $\iint_D 2\,dA = 2 \cdot A(D) = 2\left(\sqrt{2}\right)^2 = 4$, and $\iint_D (2 + x^2 y^3 - y^2 \sin x)\,dA = 4 + 0 + 0 = 4$.

69. $\iint_D \left(ax^3 + by^3 + \sqrt{a^2 - x^2}\right) dA = \iint_D ax^3\,dA + \iint_D by^3\,dA + \iint_D \sqrt{a^2 - x^2}\,dA$. Now ax^3 is odd with respect to x and by^3 is odd with respect to y, and the region of integration is symmetric with respect to both x and y, so $\iint_D ax^3\,dA = \iint_D by^3\,dA = 0$.

$\iint_D \sqrt{a^2 - x^2}\,dA$ represents the volume of the solid region under the graph of $z = \sqrt{a^2 - x^2}$ and above the rectangle D, namely a half circular cylinder with radius a and length $2b$ (see the figure) whose volume is $\frac{1}{2} \cdot \pi r^2 h = \frac{1}{2}\pi a^2 (2b) = \pi a^2 b$. Thus



$\iint_D \left(ax^3 + by^3 + \sqrt{a^2 - x^2}\right) dA = 0 + 0 + \pi a^2 b = \pi a^2 b$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

70. To find the equations of the boundary curves, we require that the z-values of the two surfaces be the same. In Maple, we use the command `solve(4-x^2-y^2=1-x-y,y);` and in Mathematica, we use `Solve[4-x^2-y^2==1-x-y,y]`. We find that the curves have equations $y = \dfrac{1 \pm \sqrt{13 + 4x - 4x^2}}{2}$. To find the two points of intersection of these curves, we use the CAS to solve $13 + 4x - 4x^2 = 0$, finding that $x = \frac{1 \pm \sqrt{14}}{2}$. So, using the CAS to evaluate the integral, the volume of intersection is



$$V = \int_{(1-\sqrt{14})/2}^{(1+\sqrt{14})/2} \int_{(1-\sqrt{13+4x-4x^2})/2}^{(1+\sqrt{13+4x-4x^2})/2} [(4 - x^2 - y^2) - (1 - x - y)]\, dy\, dx = \frac{49\pi}{8}$$

15.3 Double Integrals in Polar Coordinates

1. The region R is more easily described by polar coordinates: $R = \{(r, \theta) \mid 2 \le r \le 5, 0 \le \theta \le 2\pi\}$.

Thus $\iint_R f(x, y)\, dA = \int_0^{2\pi} \int_2^5 f(r\cos\theta, r\sin\theta)\, r\, dr\, d\theta$.

2. The region R is more easily described by rectangular coordinates: $R = \{(x, y) \mid -1 \le x \le 1,\ -x \le y \le 1\}$.

Thus $\iint_R f(x, y)\, dA = \int_{-1}^{1} \int_{-x}^{1} f(x, y)\, dy\, dx$.

3. The region R is more easily described by polar coordinates: $R = \{(r, \theta) \mid 0 \le r \le 1, \pi \le \theta \le 2\pi\}$.

Thus $\iint_R f(x, y)\, dA = \int_{\pi}^{2\pi} \int_0^1 f(r\cos\theta, r\sin\theta)\, r\, dr\, d\theta$.

4. The region R is more easily described by polar coordinates: $R = \left\{(r, \theta) \mid 0 \le r \le 3,\ -\frac{\pi}{4} \le \theta \le \frac{3\pi}{4}\right\}$.

Thus $\iint_R f(x, y)\, dA = \int_{-\pi/4}^{3\pi/4} \int_0^3 f(r\cos\theta, r\sin\theta)\, r\, dr\, d\theta$.

5. The integral $\int_{\pi/4}^{3\pi/4} \int_1^2 r\, dr\, d\theta$ represents the area of the region $R = \{(r, \theta) \mid 1 \le r \le 2, \pi/4 \le \theta \le 3\pi/4\}$, the top quarter portion of a ring (annulus).



$$\int_{\pi/4}^{3\pi/4} \int_1^2 r\, dr\, d\theta = \left(\int_{\pi/4}^{3\pi/4} d\theta\right)\left(\int_1^2 r\, dr\right)$$
$$= \left[\theta\right]_{\pi/4}^{3\pi/4} \left[\tfrac{1}{2}r^2\right]_1^2 = \left(\tfrac{3\pi}{4} - \tfrac{\pi}{4}\right) \cdot \tfrac{1}{2}(4 - 1) = \tfrac{\pi}{2} \cdot \tfrac{3}{2} = \tfrac{3\pi}{4}$$

6. The integral $\int_{\pi/2}^{\pi} \int_0^{2\sin\theta} r\, dr\, d\theta$ represents the area of the region $R = \{(r, \theta) \mid 0 \le r \le 2\sin\theta, \pi/2 \le \theta \le \pi\}$. Since $r = 2\sin\theta \ \Rightarrow\ r^2 = 2r\sin\theta \ \Leftrightarrow\ x^2 + y^2 = 2y \ \Leftrightarrow\ x^2 + (y - 1)^2 = 1$, R is the portion in the second quadrant of a disk of radius 1 with center $(0, 1)$.



$$\int_{\pi/2}^{\pi} \int_0^{2\sin\theta} r\, dr\, d\theta = \int_{\pi/2}^{\pi} \left[\tfrac{1}{2}r^2\right]_{r=0}^{r=2\sin\theta} d\theta = \int_{\pi/2}^{\pi} 2\sin^2\theta\, d\theta$$
$$= \int_{\pi/2}^{\pi} 2 \cdot \tfrac{1}{2}(1 - \cos 2\theta)\, d\theta = \left[\theta - \tfrac{1}{2}\sin 2\theta\right]_{\pi/2}^{\pi}$$
$$= \pi - 0 - \tfrac{\pi}{2} + 0 = \tfrac{\pi}{2}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

7. The half disk D can be described in polar coordinates as $D = \{(r,\theta) \mid 0 \le r \le 5, 0 \le \theta \le \pi\}$. Then

$$\iint_D x^2 y\,dA = \int_0^{\pi}\int_0^5 (r\cos\theta)^2 (r\sin\theta)\,r\,dr\,d\theta = \left(\int_0^{\pi}\cos^2\theta\sin\theta\,d\theta\right)\left(\int_0^5 r^4\,dr\right)$$

$$= \left[-\tfrac{1}{3}\cos^3\theta\right]_0^{\pi}\left[\tfrac{1}{5}r^5\right]_0^5 = -\tfrac{1}{3}(-1-1)\cdot 625 = \tfrac{1250}{3}$$

8. The region R is $\frac{1}{8}$ of a disk, as shown in the figure, and can be described by $R = \{(r,\theta) \mid 0 \le r \le 2, \pi/4 \le \theta \le \pi/2\}$. Thus

$$\iint_R (2x-y)\,dA = \int_{\pi/4}^{\pi/2}\int_0^2 (2r\cos\theta - r\sin\theta)\,r\,dr\,d\theta$$

$$= \int_{\pi/4}^{\pi/2}(2\cos\theta - \sin\theta)\,d\theta \int_0^2 r^2\,dr$$

$$= \left[2\sin\theta + \cos\theta\right]_{\pi/4}^{\pi/2}\left[\tfrac{1}{3}r^3\right]_0^2$$

$$= (2 + 0 - \sqrt{2} - \tfrac{\sqrt{2}}{2})(\tfrac{8}{3}) = \tfrac{16}{3} - 4\sqrt{2}$$



9. $\iint_R \sin(x^2+y^2)\,dA = \int_0^{\pi/2}\int_1^3 \sin(r^2)\,r\,dr\,d\theta = \int_0^{\pi/2} d\theta \int_1^3 r\sin(r^2)\,dr = \left[\theta\right]_0^{\pi/2}\left[-\tfrac{1}{2}\cos(r^2)\right]_1^3$

$$= \left(\tfrac{\pi}{2}\right)\left[-\tfrac{1}{2}(\cos 9 - \cos 1)\right] = \tfrac{\pi}{4}(\cos 1 - \cos 9)$$

10. $$\iint_R \frac{y^2}{x^2+y^2}\,dA = \int_0^{2\pi}\int_a^b \frac{(r\sin\theta)^2}{r^2}\,r\,dr\,d\theta = \int_0^{2\pi}\sin^2\theta\,d\theta\int_a^b r\,dr = \int_0^{2\pi}\frac{1}{2}(1-\cos 2\theta)\,d\theta\int_a^b r\,dr$$

$$= \tfrac{1}{2}\left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{2}r^2\right]_a^b = \tfrac{1}{2}(2\pi - 0 - 0)\cdot\tfrac{1}{2}(b^2-a^2) = \tfrac{\pi}{2}(b^2-a^2)$$

11. $\iint_D e^{-x^2-y^2}\,dA = \int_{-\pi/2}^{\pi/2}\int_0^2 e^{-r^2}\,r\,dr\,d\theta = \int_{-\pi/2}^{\pi/2} d\theta \int_0^2 re^{-r^2}\,dr$

$$= \left[\theta\right]_{-\pi/2}^{\pi/2}\left[-\tfrac{1}{2}e^{-r^2}\right]_0^2 = \pi\left(-\tfrac{1}{2}\right)(e^{-4} - e^0) = \tfrac{\pi}{2}(1-e^{-4})$$

12. $\iint_D \cos\sqrt{x^2+y^2}\,dA = \int_0^{2\pi}\int_0^2 \cos\sqrt{r^2}\;r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^2 r\cos r\,dr$. For the second integral, integrate by parts with $u = r$, $dv = \cos r\,dr$. Then $\iint_D \cos\sqrt{x^2+y^2}\,dA = \left[\theta\right]_0^{2\pi}\left[r\sin r + \cos r\right]_0^2 = 2\pi(2\sin 2 + \cos 2 - 1)$.

13. R is the region shown in the figure, and can be described by $R = \{(r,\theta) \mid 0 \le \theta \le \pi/4, 1 \le r \le 2\}$. Thus

$\iint_R \arctan(y/x)\,dA = \int_0^{\pi/4}\int_1^2 \arctan(\tan\theta)\,r\,dr\,d\theta$ since $y/x = \tan\theta$.

Also, $\arctan(\tan\theta) = \theta$ for $0 \le \theta \le \pi/4$, so the integral becomes

$\int_0^{\pi/4}\int_1^2 \theta\,r\,dr\,d\theta = \int_0^{\pi/4}\theta\,d\theta\int_1^2 r\,dr = \left[\tfrac{1}{2}\theta^2\right]_0^{\pi/4}\left[\tfrac{1}{2}r^2\right]_1^2 = \tfrac{\pi^2}{32}\cdot\tfrac{3}{2} = \tfrac{3}{64}\pi^2$.



14.



$$\iint_D x\,dA = \iint_{\substack{x^2+y^2\le 4\\ x\ge 0,\,y\ge 0}} x\,dA - \iint_{\substack{(x-1)^2+y^2\le 1\\ y\ge 0}} x\,dA$$

$$= \int_0^{\pi/2}\int_0^2 r^2\cos\theta\,dr\,d\theta - \int_0^{\pi/2}\int_0^{2\cos\theta} r^2\cos\theta\,dr\,d\theta$$

$$= \int_0^{\pi/2}\tfrac{1}{3}(8\cos\theta)\,d\theta - \int_0^{\pi/2}\tfrac{1}{3}(8\cos^4\theta)\,d\theta$$

$$= \tfrac{8}{3} - \tfrac{8}{12}\left[\cos^3\theta\sin\theta + \tfrac{3}{2}(\theta + \sin\theta\cos\theta)\right]_0^{\pi/2}$$

$$= \tfrac{8}{3} - \tfrac{2}{3}\left[0 + \tfrac{3}{2}\left(\tfrac{\pi}{2}\right)\right] = \tfrac{16-3\pi}{6}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. One loop is given by the region $D = \{(r,\theta) \mid -\pi/6 \le \theta \le \pi/6, 0 \le r \le \cos 3\theta\}$, so the area is



$$\iint_D dA = \int_{-\pi/6}^{\pi/6}\int_0^{\cos 3\theta} r\,dr\,d\theta = \int_{-\pi/6}^{\pi/6}\left[\frac{1}{2}r^2\right]_{r=0}^{r=\cos 3\theta} d\theta$$

$$= \int_{-\pi/6}^{\pi/6} \frac{1}{2}\cos^2 3\theta\,d\theta = 2\int_0^{\pi/6} \frac{1}{2}\left(\frac{1+\cos 6\theta}{2}\right) d\theta$$

$$= \frac{1}{2}\left[\theta + \frac{1}{6}\sin 6\theta\right]_0^{\pi/6} = \frac{\pi}{12}$$

16. By symmetry, the area of the region is 4 times the area of the region D in the first quadrant enclosed by the cardiod $r = 1 - \cos\theta$ (see the figure). Here $D = \{(r,\theta) \mid 0 \le r \le 1 - \cos\theta, 0 \le \theta \le \pi/2\}$, so the total area is



$$4A(D) = 4\iint_D dA = 4\int_0^{\pi/2}\int_0^{1-\cos\theta} r\,dr\,d\theta = 4\int_0^{\pi/2}\left[\tfrac{1}{2}r^2\right]_{r=0}^{r=1-\cos\theta} d\theta$$

$$= 2\int_0^{\pi/2}(1-\cos\theta)^2 d\theta = 2\int_0^{\pi/2}(1 - 2\cos\theta + \cos^2\theta)\,d\theta$$

$$= 2\int_0^{\pi/2}\left[1 - 2\cos\theta + \tfrac{1}{2}(1+\cos 2\theta)\right] d\theta$$

$$= 2\left[\theta - 2\sin\theta + \tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta\right]_0^{\pi/2}$$

$$= 2\left(\tfrac{\pi}{2} - 2 + \tfrac{\pi}{4}\right) = \tfrac{3\pi}{2} - 4$$

17. In polar coordinates the circle $(x-1)^2 + y^2 = 1 \;\Leftrightarrow\; x^2 + y^2 = 2x$ is $r^2 = 2r\cos\theta \;\Rightarrow\; r = 2\cos\theta$, and the circle $x^2 + y^2 = 1$ is $r = 1$. The curves intersect in the first quadrant when $2\cos\theta = 1 \;\Rightarrow\; \cos\theta = \tfrac{1}{2} \;\Rightarrow\; \theta = \pi/3$, so the portion of the region in the first quadrant is given by $D = \{(r,\theta) \mid 1 \le r \le 2\cos\theta, 0 \le \theta \le \pi/3\}$. By symmetry, the total area is twice the area of D:



$$2A(D) = 2\iint_D dA = 2\int_0^{\pi/3}\int_1^{2\cos\theta} r\,dr\,d\theta = 2\int_0^{\pi/3}\left[\tfrac{1}{2}r^2\right]_{r=1}^{r=2\cos\theta} d\theta$$

$$= \int_0^{\pi/3}\left(4\cos^2\theta - 1\right) d\theta = \int_0^{\pi/3}\left[4\cdot\tfrac{1}{2}(1+\cos 2\theta) - 1\right] d\theta$$

$$= \int_0^{\pi/3}(1 + 2\cos 2\theta)\,d\theta = [\theta + \sin 2\theta]_0^{\pi/3} = \tfrac{\pi}{3} + \tfrac{\sqrt{3}}{2}$$

18. The region lies between the two polar curves in quadrants I and IV, but in quadrants II and III the region is enclosed by the cardioid. In the first quadrant, $1 + \cos\theta = 3\cos\theta$ when $\cos\theta = \tfrac{1}{2} \;\Rightarrow\; \theta = \tfrac{\pi}{3}$, so the area of the region inside the cardioid and outside the circle is



$$A_1 = \int_{\pi/3}^{\pi/2}\int_{3\cos\theta}^{1+\cos\theta} r\,dr\,d\theta = \int_{\pi/3}^{\pi/2}\left[\tfrac{1}{2}r^2\right]_{r=3\cos\theta}^{r=1+\cos\theta} d\theta$$

$$= \tfrac{1}{2}\int_{\pi/3}^{\pi/2}(1 + 2\cos\theta - 8\cos^2\theta)d\theta = \tfrac{1}{2}\left[\theta + 2\sin\theta - 8\left(\tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta\right)\right]_{\pi/3}^{\pi/2}$$

$$= \left[-\tfrac{3}{2}\theta + \sin\theta - \sin 2\theta\right]_{\pi/3}^{\pi/2} = \left(-\tfrac{3\pi}{4} + 1 - 0\right) - \left(-\tfrac{\pi}{2} + \tfrac{\sqrt{3}}{2} - \tfrac{\sqrt{3}}{2}\right) = 1 - \tfrac{\pi}{4}.$$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

The area of the region in the second quadrant is

$$A_2 = \int_{\pi/2}^{\pi}\int_0^{1+\cos\theta} r\,dr\,d\theta = \int_{\pi/2}^{\pi}\left[\tfrac{1}{2}r^2\right]_{r=0}^{r=1+\cos\theta} d\theta = \tfrac{1}{2}\int_{\pi/2}^{\pi}(1+2\cos\theta+\cos^2\theta)\,d\theta$$
$$= \tfrac{1}{2}\left[\theta + 2\sin\theta + \tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta\right]_{\pi/2}^{\pi} = \tfrac{1}{2}\left(\tfrac{3\pi}{4} - 2\right) = \tfrac{3\pi}{8} - 1.$$

By symmetry, the total area is $A = 2(A_1 + A_2) = 2\left(1 - \tfrac{\pi}{4} + \tfrac{3\pi}{8} - 1\right) = \tfrac{\pi}{4}$.

19. $V = \iint_{x^2+y^2\le 25}\left(x^2+y^2\right)dA = \int_0^{2\pi}\int_0^5 r^2\cdot r\,dr\,d\theta = \int_0^{2\pi} d\theta\int_0^5 r^3\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^5 = 2\pi\left(\tfrac{625}{4}\right) = \tfrac{625}{2}\pi$

20. $V = \iint_{1\le x^2+y^2\le 4}\sqrt{x^2+y^2}\,dA = \int_0^{2\pi}\int_1^2\sqrt{r^2}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta\int_1^2 r^2\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{1}{3}r^3\right]_1^2 = 2\pi\left(\tfrac{8}{3} - \tfrac{1}{3}\right) = \tfrac{14}{3}\pi$

21. $2x + y + z = 4 \iff z = 4 - 2x - y$, so the volume of the solid is

$$V = \iint_{x^2+y^2\le 1}(4 - 2x - y)\,dA = \int_0^{2\pi}\int_0^1 (4 - 2r\cos\theta - r\sin\theta)\,r\,dr\,d\theta$$
$$= \int_0^{2\pi}\int_0^1 \left[4r - r^2(2\cos\theta + \sin\theta)\right]dr\,d\theta = \int_0^{2\pi}\left[2r^2 - \tfrac{1}{3}r^3(2\cos\theta+\sin\theta)\right]_{r=0}^{r=1} d\theta$$
$$= \int_0^{2\pi}\left[2 - \tfrac{1}{3}(2\cos\theta+\sin\theta)\right]d\theta = \left[2\theta - \tfrac{1}{3}(2\sin\theta - \cos\theta)\right]_0^{2\pi} = 4\pi + \tfrac{1}{3} - 0 - \tfrac{1}{3} = 4\pi$$

22. The sphere $x^2+y^2+z^2 = 16$ intersects the xy-plane in the circle $x^2+y^2 = 16$, so

$$V = 2\iint_{4\le x^2+y^2\le 16}\sqrt{16 - x^2 - y^2}\,dA \quad \text{[by symmetry]} \quad = 2\int_0^{2\pi}\int_2^4\sqrt{16-r^2}\,r\,dr\,d\theta = 2\int_0^{2\pi} d\theta\int_2^4 r(16-r^2)^{1/2}dr$$
$$= 2\left[\theta\right]_0^{2\pi}\left[-\tfrac{1}{3}(16-r^2)^{3/2}\right]_2^4 = -\tfrac{2}{3}(2\pi)(0 - 12^{3/2}) = \tfrac{4\pi}{3}\left(12\sqrt{12}\right) = 32\sqrt{3}\,\pi$$

23. By symmetry,

$$V = 2\iint_{x^2+y^2\le a^2}\sqrt{a^2 - x^2 - y^2}\,dA = 2\int_0^{2\pi}\int_0^a\sqrt{a^2-r^2}\,r\,dr\,d\theta = 2\int_0^{2\pi} d\theta\int_0^a r\sqrt{a^2-r^2}\,dr$$
$$= 2\left[\theta\right]_0^{2\pi}\left[-\tfrac{1}{3}(a^2-r^2)^{3/2}\right]_0^a = 2(2\pi)\left(0 + \tfrac{1}{3}a^3\right) = \tfrac{4}{3}\pi a^3$$

24. The paraboloid $z = 1 + 2x^2 + 2y^2$ intersects the plane $z = 7$ when $7 = 1 + 2x^2 + 2y^2$ or $x^2 + y^2 = 3$ and we are restricted to the first octant, so

$$V = \iint_{\substack{x^2+y^2\le 3,\\ x\ge 0, y\ge 0}}\left[7 - \left(1 + 2x^2 + 2y^2\right)\right]dA = \int_0^{\pi/2}\int_0^{\sqrt{3}}\left[7 - (1 + 2r^2)\right]r\,dr\,d\theta$$
$$= \int_0^{\pi/2} d\theta\int_0^{\sqrt{3}}(6r - 2r^3)\,dr = \left[\theta\right]_0^{\pi/2}\left[3r^2 - \tfrac{1}{2}r^4\right]_0^{\sqrt{3}} = \tfrac{\pi}{2}\cdot\tfrac{9}{2} = \tfrac{9}{4}\pi$$

25. The cone $z = \sqrt{x^2+y^2}$ intersects the sphere $x^2+y^2+z^2 = 1$ when $x^2 + y^2 + \left(\sqrt{x^2+y^2}\right)^2 = 1$ or $x^2+y^2 = \tfrac{1}{2}$. So

$$V = \iint_{x^2+y^2\le 1/2}\left(\sqrt{1 - x^2 - y^2} - \sqrt{x^2+y^2}\right)dA = \int_0^{2\pi}\int_0^{1/\sqrt{2}}\left(\sqrt{1-r^2} - r\right)r\,dr\,d\theta$$
$$= \int_0^{2\pi} d\theta\int_0^{1/\sqrt{2}}\left(r\sqrt{1-r^2} - r^2\right)dr = \left[\theta\right]_0^{2\pi}\left[-\tfrac{1}{3}(1-r^2)^{3/2} - \tfrac{1}{3}r^3\right]_0^{1/\sqrt{2}} = 2\pi\left(-\tfrac{1}{3}\right)\left(\tfrac{1}{\sqrt{2}} - 1\right) = \tfrac{\pi}{3}\left(2 - \sqrt{2}\right)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. The two paraboloids intersect when $6 - x^2 - y^2 = 2x^2 + 2y^2$ or $x^2 + y^2 = 2$. For $x^2 + y^2 \le 2$, the paraboloid $z = 6 - x^2 - y^2$ is above $z = 2x^2 + 2y^2$ so

$$V = \iint_{x^2+y^2\le 2} [(6 - x^2 - y^2) - (2x^2 + 2y^2)]\, dA = \iint_{x^2+y^2\le 2} [6 - 3(x^2 + y^2)]\, dA = \int_0^{2\pi}\int_0^{\sqrt{2}} (6 - 3r^2)\, r\, dr\, d\theta$$

$$= \int_0^{2\pi} d\theta \int_0^{\sqrt{2}} (6r - 3r^3)\, dr = \left[\theta\right]_0^{2\pi} \left[3r^2 - \tfrac{3}{4}r^4\right]_0^{\sqrt{2}} = 2\pi\,(6 - 3) = 6\pi$$

27. The given solid is the region inside the cylinder $x^2 + y^2 = 4$ between the surfaces $z = \sqrt{64 - 4x^2 - 4y^2}$ and $z = -\sqrt{64 - 4x^2 - 4y^2}$. So

$$V = \iint_{x^2+y^2\le 4} \left[\sqrt{64 - 4x^2 - 4y^2} - \left(-\sqrt{64 - 4x^2 - 4y^2}\right)\right] dA = \iint_{x^2+y^2\le 4} 2 \cdot 2\sqrt{16 - x^2 - y^2}\, dA$$

$$= 4\int_0^{2\pi}\int_0^2 \sqrt{16 - r^2}\; r\, dr\, d\theta = 4\int_0^{2\pi} d\theta \int_0^2 r\sqrt{16 - r^2}\, dr = 4\left[\theta\right]_0^{2\pi} \left[-\tfrac{1}{3}(16 - r^2)^{3/2}\right]_0^2$$

$$= 8\pi\left(-\tfrac{1}{3}\right)(12^{3/2} - 16^{2/3}) = \tfrac{8\pi}{3}\left(64 - 24\sqrt{3}\right)$$

28. (a) Here the region in the xy-plane is the annular region $r_1^2 \le x^2 + y^2 \le r_2^2$ and the desired volume is twice that above the xy-plane. Hence

$$V = 2 \iint_{r_1^2\le x^2+y^2\le r_2^2} \sqrt{r_2^2 - x^2 - y^2}\, dA = 2\int_0^{2\pi}\int_{r_1}^{r_2} \sqrt{r_2^2 - r^2}\; r\, dr\, d\theta = 2\int_0^{2\pi} d\theta \int_{r_1}^{r_2} \sqrt{r_2^2 - r^2}\; r\, dr$$

$$= 2\,(2\pi)\left[-\tfrac{1}{3}(r_2^2 - r^2)^{3/2}\right]_{r_1}^{r_2} = \tfrac{4\pi}{3}(r_2^2 - r_1^2)^{3/2}$$

(b) A cross-sectional cut is shown in the figure. So $r_2^2 = \left(\frac{1}{2}h\right)^2 + r_1^2$ or $\frac{1}{4}h^2 = r_2^2 - r_1^2$.

Thus the volume in terms of h is $V = \frac{4\pi}{3}\left(\frac{1}{4}h^2\right)^{3/2} = \frac{\pi}{6}h^3$.



29.



$$\int_0^2 \int_0^{\sqrt{4-x^2}} e^{-x^2-y^2}\, dy\, dx = \int_0^{\pi/2}\int_0^2 e^{-r^2}\, r\, dr\, d\theta$$

$$= \int_0^{\pi/2} d\theta \int_0^2 re^{-r^2}\, dr = \left[\theta\right]_0^{\pi/2} \left[-\tfrac{1}{2}e^{-r^2}\right]_0^2$$

$$= \tfrac{\pi}{2}\left[-\tfrac{1}{2}\left(e^{-4} - 1\right)\right] = \tfrac{\pi}{4}\left(1 - e^{-4}\right)$$

30.



$$\int_0^a \int_{-\sqrt{a^2-y^2}}^{\sqrt{a^2-y^2}} (2x + y)\, dx\, dy = \int_0^{\pi}\int_0^a (2r\cos\theta + r\sin\theta)\, r\, dr\, d\theta$$

$$= \int_0^{\pi} (2\cos\theta + \sin\theta)\, d\theta \int_0^a r^2\, dr$$

$$= \left[2\sin\theta - \cos\theta\right]_0^{\pi} \left[\tfrac{1}{3}r^3\right]_0^a$$

$$= [(0 + 1) - (0 - 1)] \cdot \tfrac{1}{3}(a^3 - 0) = \tfrac{2}{3}a^3$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

31. The region D of integration is shown in the figure. In polar coordinates the line $x = \sqrt{3}\,y$ is $\theta = \pi/6$, so



$$\begin{aligned}\int_0^{1/2}\int_{\sqrt{3}\,y}^{\sqrt{1-y^2}} xy^2\,dx\,dy &= \int_0^{\pi/6}\int_0^1 (r\cos\theta)(r\sin\theta)^2\, r\,dr\,d\theta \\ &= \int_0^{\pi/6} \sin^2\theta\cos\theta\,d\theta \int_0^1 r^4\,dr \\ &= \left[\tfrac{1}{3}\sin^3\theta\right]_0^{\pi/6}\left[\tfrac{1}{5}r^5\right]_0^1 \\ &= \left[\tfrac{1}{3}\left(\tfrac{1}{2}\right)^3 - 0\right]\left[\tfrac{1}{5} - 0\right] = \tfrac{1}{120}\end{aligned}$$

32.



$$\begin{aligned}\int_0^{\pi/2}\int_0^{2\cos\theta} r^2\,dr\,d\theta &= \int_0^{\pi/2}\left[\tfrac{1}{3}r^3\right]_{r=0}^{r=2\cos\theta} d\theta = \int_0^{\pi/2}\left(\tfrac{8}{3}\cos^3\theta\right)d\theta \\ &= \tfrac{8}{3}\int_0^{\pi/2}(1-\sin^2\theta)\cos\theta\,d\theta \\ &= \tfrac{8}{3}\left[\sin\theta - \tfrac{1}{3}\sin^3\theta\right]_0^{\pi/2} = \tfrac{16}{9}\end{aligned}$$

33. $D = \{(r,\theta) \mid 0 \le r \le 1, 0 \le \theta \le 2\pi\}$, so

$\iint_D e^{(x^2+y^2)^2}\,dA = \int_0^{2\pi}\int_0^1 e^{(r^2)^2}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^1 re^{r^4}\,dr = 2\pi\int_0^1 re^{r^4}\,dr$. Using a calculator, we estimate $2\pi\int_0^1 re^{r^4}\,dr \approx 4.5951$.

34. $D = \{(r,\theta) \mid 0 \le r \le 1, 0 \le \theta \le \pi/2\}$, so

$$\begin{aligned}\iint_D xy\sqrt{1+x^2+y^2}\,dA &= \int_0^{\pi/2}\int_0^1 (r\cos\theta)(r\sin\theta)\sqrt{1+r^2}\;r\,dr\,d\theta \\ &= \int_0^{\pi/2}\sin\theta\cos\theta\,d\theta\int_0^1 r^3\sqrt{1+r^2}\,dr = \left[\tfrac{1}{2}\sin^2\theta\right]_0^{\pi/2}\int_0^1 r^3\sqrt{1+r^2}\,dr \\ &= \tfrac{1}{2}\int_0^1 r^3\sqrt{1+r^2}\,dr \approx 0.1609\end{aligned}$$

35. The surface of the water in the pool is a circular disk D with radius 20 ft. If we place D on coordinate axes with the origin at the center of D and define $f(x,y)$ to be the depth of the water at (x,y), then the volume of water in the pool is the volume of the solid that lies above $D = \{(x,y) \mid x^2 + y^2 \le 400\}$ and below the graph of $f(x,y)$. We can associate north with the positive y-direction, so we are given that the depth is constant in the x-direction and the depth increases linearly in the y-direction from $f(0,-20) = 2$ to $f(0,20) = 7$. The trace in the yz-plane is a line segment from $(0,-20,2)$ to $(0,20,7)$. The slope of this line is $\frac{7-2}{20-(-20)} = \frac{1}{8}$, so an equation of the line is $z - 7 = \frac{1}{8}(y-20) \;\Rightarrow\; z = \frac{1}{8}y + \frac{9}{2}$. Since $f(x,y)$ is independent of x, $f(x,y) = \frac{1}{8}y + \frac{9}{2}$. Thus the volume is given by $\iint_D f(x,y)\,dA$, which is most conveniently evaluated using polar coordinates. Then $D = \{(r,\theta) \mid 0 \le r \le 20, 0 \le \theta \le 2\pi\}$ and substituting $x = r\cos\theta$, $y = r\sin\theta$ the integral becomes

$$\begin{aligned}\int_0^{2\pi}\int_0^{20}\left(\tfrac{1}{8}r\sin\theta + \tfrac{9}{2}\right)r\,dr\,d\theta &= \int_0^{2\pi}\left[\tfrac{1}{24}r^3\sin\theta + \tfrac{9}{4}r^2\right]_{r=0}^{r=20} d\theta = \int_0^{2\pi}\left(\tfrac{1000}{3}\sin\theta + 900\right)d\theta \\ &= \left[-\tfrac{1000}{3}\cos\theta + 900\theta\right]_0^{2\pi} = 1800\pi\end{aligned}$$

Thus the pool contains $1800\pi \approx 5655$ ft^3 of water.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. (a) If $R \le 100$, the total amount of water supplied each hour to the region within R feet of the sprinkler is

$$V = \int_0^{2\pi}\int_0^R e^{-r} r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^R re^{-r}\,dr = \left[\theta\right]_0^{2\pi}\left[-re^{-r} - e^{-r}\right]_0^R$$
$$= 2\pi[-Re^{-R} - e^{-R} + 0 + 1] = 2\pi(1 - Re^{-R} - e^{-R}) \text{ ft}^3$$

(b) The average amount of water per hour per square foot supplied to the region within R feet of the sprinkler is

$$\frac{V}{\text{area of region}} = \frac{V}{\pi R^2} = \frac{2\left(1 - Re^{-R} - e^{-R}\right)}{R^2} \text{ ft}^3 \text{ (per hour per square foot). See the definition of the average value of a function on page 1037 [ET 997].}$$

37. As in Exercise 15.2.61, $f_{\text{ave}} = \frac{1}{A(D)}\iint_D f(x,y)dA$. Here $D = \{(r,\theta) \mid a \le r \le b, 0 \le \theta \le 2\pi\}$, so $A(D) = \pi b^2 - \pi a^2 = \pi(b^2 - a^2)$ and

$$f_{\text{ave}} = \frac{1}{A(D)}\iint_D \frac{1}{\sqrt{x^2+y^2}}\,dA = \frac{1}{\pi(b^2-a^2)}\int_0^{2\pi}\int_a^b \frac{1}{\sqrt{r^2}}\,r\,dr\,d\theta = \frac{1}{\pi(b^2-a^2)}\int_0^{2\pi} d\theta \int_a^b dr$$
$$= \frac{1}{\pi(b^2-a^2)}\left[\theta\right]_0^{2\pi}\left[r\right]_a^b = \frac{1}{\pi(b^2-a^2)}(2\pi)(b-a) = \frac{2(b-a)}{(b+a)(b-a)} = \frac{2}{a+b}$$

38. The distance from a point (x,y) to the origin is $f(x,y) = \sqrt{x^2+y^2}$, so the average distance from points in D to the origin is

$$f_{\text{ave}} = \frac{1}{A(D)}\iint_D \sqrt{x^2+y^2}\,dA = \frac{1}{\pi a^2}\int_0^{2\pi}\int_0^a \sqrt{r^2}\,r\,dr\,d\theta$$
$$= \frac{1}{\pi a^2}\int_0^{2\pi} d\theta \int_0^a r^2\,dr = \frac{1}{\pi a^2}\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{3}r^3\right]_0^a = \frac{1}{\pi a^2}\cdot 2\pi\cdot\tfrac{1}{3}a^3 = \tfrac{2}{3}a$$

39.
$$\int_{1/\sqrt{2}}^1\int_{\sqrt{1-x^2}}^x xy\,dy\,dx + \int_1^{\sqrt{2}}\int_0^x xy\,dy\,dx + \int_{\sqrt{2}}^2\int_0^{\sqrt{4-x^2}} xy\,dy\,dx$$
$$= \int_0^{\pi/4}\int_1^2 r^3\cos\theta\sin\theta\,dr\,d\theta = \int_0^{\pi/4}\left[\frac{r^4}{4}\cos\theta\sin\theta\right]_{r=1}^{r=2} d\theta$$
$$= \frac{15}{4}\int_0^{\pi/4}\sin\theta\cos\theta\,d\theta = \frac{15}{4}\left[\frac{\sin^2\theta}{2}\right]_0^{\pi/4} = \frac{15}{16}$$



40. (a) $\iint_{D_a} e^{-(x^2+y^2)}dA = \int_0^{2\pi}\int_0^a re^{-r^2}\,dr\,d\theta = 2\pi\left[-\frac{1}{2}e^{-r^2}\right]_0^a = \pi\left(1 - e^{-a^2}\right)$ for each a. Then $\lim_{a\to\infty}\pi\left(1-e^{-a^2}\right) = \pi$ since $e^{-a^2} \to 0$ as $a \to \infty$. Hence $\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} e^{-(x^2+y^2)}\,dA = \pi$.

(b) $\iint_{S_a} e^{-(x^2+y^2)}\,dA = \int_{-a}^a\int_{-a}^a e^{-x^2}e^{-y^2}\,dx\,dy = \left(\int_{-a}^a e^{-x^2}\,dx\right)\left(\int_{-a}^a e^{-y^2}\,dy\right)$ for each a.

Then, from (a), $\pi = \iint_{\mathbb{R}^2} -(x^2+y^2)\,dA$, so

$$\pi = \lim_{a\to\infty}\iint_{S_a} e^{-(x^2+y^2)}\,dA = \lim_{a\to\infty}\left(\int_{-a}^a e^{-x^2}\,dx\right)\left(\int_{-a}^a e^{-y^2}\,dy\right) = \left(\int_{-\infty}^{\infty} e^{-x^2}\,dx\right)\left(\int_{-\infty}^{\infty} e^{-y^2}\,dy\right).$$

To evaluate $\lim_{a\to\infty}\left(\int_{-a}^a e^{-x^2}\,dx\right)\left(\int_{-a}^a e^{-y^2}\,dy\right)$, we are using the fact that these integrals are bounded. This is true since on $[-1,1]$, $0 < e^{-x^2} \le 1$ while on $(-\infty,-1)$, $0 < e^{-x^2} \le e^x$ and on $(1,\infty)$, $0 < e^{-x^2} < e^{-x}$. Hence $0 \le \int_{-\infty}^{\infty} e^{-x^2}\,dx \le \int_{-\infty}^{-1} e^x\,dx + \int_{-1}^1 dx + \int_1^{\infty} e^{-x}\,dx = 2(e^{-1}+1)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c) Since $\left(\int_{-\infty}^{\infty} e^{-x^2}\,dx\right)\left(\int_{-\infty}^{\infty} e^{-y^2}\,dy\right) = \pi$ and y can be replaced by x, $\left(\int_{-\infty}^{\infty} e^{-x^2}\,dx\right)^2 = \pi$ implies that $\int_{-\infty}^{\infty} e^{-x^2}\,dx = \pm\sqrt{\pi}$. But $e^{-x^2} \geq 0$ for all x, so $\int_{-\infty}^{\infty} e^{-x^2}\,dx = \sqrt{\pi}$.

(d) Letting $t = \sqrt{2}\,x$, $\int_{-\infty}^{\infty} e^{-x^2}\,dx = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2}}\left(e^{-t^2/2}\right)dt$, so that $\sqrt{\pi} = \frac{1}{\sqrt{2}}\int_{-\infty}^{\infty} e^{-t^2/2}\,dt$ or $\int_{-\infty}^{\infty} e^{-t^2/2}\,dt = \sqrt{2\pi}$.

41. (a) We integrate by parts with $u = x$ and $dv = xe^{-x^2}\,dx$. Then $du = dx$ and $v = -\frac{1}{2}e^{-x^2}$, so

$$\begin{aligned}\int_0^{\infty} x^2e^{-x^2}\,dx &= \lim_{t\to\infty}\int_0^t x^2e^{-x^2}\,dx = \lim_{t\to\infty}\left(-\tfrac{1}{2}xe^{-x^2}\Big]_0^t + \int_0^t \tfrac{1}{2}e^{-x^2}\,dx\right)\\ &= \lim_{t\to\infty}\left(-\tfrac{1}{2}te^{-t^2}\right) + \tfrac{1}{2}\int_0^{\infty} e^{-x^2}\,dx = 0 + \tfrac{1}{2}\int_0^{\infty} e^{-x^2}\,dx \qquad \text{[by l'Hospital's Rule]}\\ &= \tfrac{1}{4}\int_{-\infty}^{\infty} e^{-x^2}\,dx \qquad \text{[since } e^{-x^2}\text{ is an even function]}\\ &= \tfrac{1}{4}\sqrt{\pi} \qquad \text{[by Exercise 40(c)]}\end{aligned}$$

(b) Let $u = \sqrt{x}$. Then $u^2 = x \;\Rightarrow\; dx = 2u\,du \;\Rightarrow$

$$\int_0^{\infty}\sqrt{x}\,e^{-x}\,dx = \lim_{t\to\infty}\int_0^t \sqrt{x}\,e^{-x}\,dx = \lim_{t\to\infty}\int_0^{\sqrt{t}} ue^{-u^2}2u\,du = 2\int_0^{\infty} u^2e^{-u^2}\,du = 2\left(\tfrac{1}{4}\sqrt{\pi}\right) \quad \text{[by part(a)]} \;= \tfrac{1}{2}\sqrt{\pi}.$$

15.4 Applications of Double Integrals

1. $Q = \iint_D \sigma(x,y)\,dA = \int_0^5\int_2^5 (2x+4y)\,dy\,dx = \int_0^5 \left[2xy+2y^2\right]_{y=2}^{y=5} dx$

$= \int_0^5 (10x+50-4x-8)\,dx = \int_0^5 (6x+42)\,dx = \left[3x^2+42x\right]_0^5 = 75+210 = 285$ C

2. $Q = \iint_D \sigma(x,y)\,dA = \iint_D \sqrt{x^2+y^2}\,dA = \int_0^{2\pi}\int_0^1 \sqrt{r^2}\; r\,dr\,d\theta$

$= \int_0^{2\pi} d\theta \int_0^1 r^2\,dr = [\theta]_0^{2\pi}\left[\tfrac{1}{3}r^3\right]_0^1 = 2\pi\cdot\tfrac{1}{3} = \tfrac{2\pi}{3}$ C

3. $m = \iint_D \rho(x,y)\,dA = \int_1^3\int_1^4 ky^2\,dy\,dx = k\int_1^3 dx\int_1^4 y^2\,dy = k\,[x]_1^3\left[\tfrac{1}{3}y^3\right]_1^4 = k(2)(21) = 42k,$

$\overline{x} = \frac{1}{m}\iint_D x\rho(x,y)\,dA = \frac{1}{42k}\int_1^3\int_1^4 kxy^2\,dy\,dx = \frac{1}{42}\int_1^3 x\,dx\int_1^4 y^2\,dy = \frac{1}{42}\left[\tfrac{1}{2}x^2\right]_1^3\left[\tfrac{1}{3}y^3\right]_1^4 = \frac{1}{42}(4)(21) = 2,$

$\overline{y} = \frac{1}{m}\iint_D y\rho(x,y)\,dA = \frac{1}{42k}\int_1^3\int_1^4 ky^3\,dy\,dx = \frac{1}{42}\int_1^3 dx\int_1^4 y^3\,dy = \frac{1}{42}[x]_1^3\left[\tfrac{1}{4}y^4\right]_1^4 = \frac{1}{42}(2)\left(\tfrac{255}{4}\right) = \frac{85}{28}$

Hence $m = 42k$, $(\overline{x},\overline{y}) = \left(2, \tfrac{85}{28}\right)$.

4. $m = \iint_D \rho(x,y)\,dA = \int_0^a\int_0^b (1+x^2+y^2)\,dy\,dx = \int_0^a \left[y+x^2y+\tfrac{1}{3}y^3\right]_{y=0}^{y=b} dx = \int_0^a \left(b+bx^2+\tfrac{1}{3}b^3\right)dx$

$= \left[bx+\tfrac{1}{3}bx^3+\tfrac{1}{3}b^3x\right]_0^a = ab+\tfrac{1}{3}a^3b+\tfrac{1}{3}ab^3 = \tfrac{1}{3}ab(3+a^2+b^2),$

$M_y = \iint_D x\rho(x,y)\,dA = \int_0^a\int_0^b (x+x^3+xy^2)\,dy\,dx = \int_0^a \left[xy+x^3y+\tfrac{1}{3}xy^3\right]_{y=0}^{y=b} dx = \int_0^a \left(bx+bx^3+\tfrac{1}{3}b^3x\right)dx$

$= \left[\tfrac{1}{2}bx^2+\tfrac{1}{4}bx^4+\tfrac{1}{6}b^3x^2\right]_0^a = \tfrac{1}{2}a^2b+\tfrac{1}{4}a^4b+\tfrac{1}{6}a^2b^3 = \tfrac{1}{12}a^2b(6+3a^2+2b^2)$, and

$M_x = \iint_D y\rho(x,y)\,dA = \int_0^a\int_0^b (y+x^2y+y^3)\,dy\,dx = \int_0^a \left[\tfrac{1}{2}y^2+\tfrac{1}{2}x^2y^2+\tfrac{1}{4}y^4\right]_{y=0}^{y=b} dx = \int_0^a \left(\tfrac{1}{2}b^2+\tfrac{1}{2}b^2x^2+\tfrac{1}{4}b^4\right)dx$

$= \left[\tfrac{1}{2}b^2x+\tfrac{1}{6}b^2x^3+\tfrac{1}{4}b^4x\right]_0^a = \tfrac{1}{2}ab^2+\tfrac{1}{6}a^3b^2+\tfrac{1}{4}ab^4 = \tfrac{1}{12}ab^2(6+2a^2+3b^2).$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Hence, $(\overline{x},\overline{y}) = \left(\dfrac{M_y}{m},\dfrac{M_x}{m}\right) = \left(\dfrac{\frac{1}{12}a^2b(6+3a^2+2b^2)}{\frac{1}{3}ab(3+a^2+b^2)},\dfrac{\frac{1}{12}ab^2(6+2a^2+3b^2)}{\frac{1}{3}ab(3+a^2+b^2)}\right)$

$= \left(\dfrac{a(6+3a^2+2b^2)}{4(3+a^2+b^2)},\dfrac{b(6+2a^2+3b^2)}{4(3+a^2+b^2)}\right).$

5. $m = \int_0^2\int_{x/2}^{3-x}(x+y)\,dy\,dx = \int_0^2\left[xy+\frac{1}{2}y^2\right]_{y=x/2}^{y=3-x}dx = \int_0^2\left[x(3-x)+\frac{1}{2}(3-x)^2-\frac{1}{2}x^2-\frac{1}{8}x^2\right]dx$

$= \int_0^2\left(-\frac{9}{8}x^2+\frac{9}{2}\right)dx = \left[-\frac{9}{8}\left(\frac{1}{3}x^3\right)+\frac{9}{2}x\right]_0^2 = 6,$

$M_y = \int_0^2\int_{x/2}^{3-x}(x^2+xy)\,dy\,dx = \int_0^2\left[x^2y+\frac{1}{2}xy^2\right]_{y=x/2}^{y=3-x}dx = \int_0^2\left(\frac{9}{2}x-\frac{9}{8}x^3\right)dx = \frac{9}{2},$

$M_x = \int_0^2\int_{x/2}^{3-y}(xy+y^2)\,dy\,dx = \int_0^2\left[\frac{1}{2}xy^2+\frac{1}{3}y^3\right]_{y=x/2}^{y=3-x}dx = \int_0^2\left(9-\frac{9}{2}x\right)dx = 9.$

Hence $m=6$, $(\overline{x},\overline{y}) = \left(\dfrac{M_y}{m},\dfrac{M_x}{m}\right) = \left(\dfrac{3}{4},\dfrac{3}{2}\right).$

6. Here $D = \{(x,y) \mid 0 \le y \le \frac{2}{5},\ y/2 \le x \le 1-2y\}.$

$m = \int_0^{2/5}\int_{y/2}^{1-2y}x\,dx\,dy = \int_0^{2/5}\left[\frac{1}{2}x^2\right]_{x=y/2}^{x=1-2y}dy = \frac{1}{2}\int_0^{2/5}\left[(1-2y)^2-\left(\frac{1}{2}y\right)^2\right]dy$

$= \frac{1}{2}\int_0^{2/5}\left(\frac{15}{4}y^2-4y+1\right)dy = \frac{1}{2}\left[\frac{5}{4}y^3-2y^2+y\right]_0^{2/5} = \frac{1}{2}\left[\frac{2}{25}-\frac{8}{25}+\frac{2}{5}\right] = \frac{2}{25},$

$M_y = \int_0^{2/5}\int_{y/2}^{1-2y}x\cdot x\,dx\,dy = \int_0^{2/5}\left[\frac{1}{3}x^3\right]_{x=y/2}^{x=1-2y}dy = \frac{1}{3}\int_0^{2/5}\left[(1-2y)^3-\left(\frac{1}{2}y\right)^3\right]dy$

$= \frac{1}{3}\int_0^{2/5}\left(-\frac{65}{8}y^3+12y^2-6y+1\right)dy = \frac{1}{3}\left[-\frac{65}{32}y^4+4y^3-3y^2+y\right]_0^{2/5} = \frac{1}{3}\left[-\frac{13}{250}+\frac{32}{125}-\frac{12}{25}+\frac{2}{5}\right] = \frac{31}{750},$

$M_x = \int_0^{2/5}\int_{y/2}^{1-2y}y\cdot x\,dx\,dy = \int_0^{2/5}y\left[\frac{1}{2}x^2\right]_{x=y/2}^{x=1-2y}dy = \frac{1}{2}\int_0^{2/5}y\left(\frac{15}{4}y^2-4y+1\right)dy$

$= \frac{1}{2}\int_0^{2/5}\left(\frac{15}{4}y^3-4y^2+y\right)dy = \frac{1}{2}\left[\frac{15}{16}y^4-\frac{4}{3}y^3+\frac{1}{2}y^2\right]_0^{2/5} = \frac{1}{2}\left[\frac{3}{125}-\frac{32}{375}+\frac{2}{25}\right] = \frac{7}{750}.$

Hence $m=\frac{2}{25}$, $(\overline{x},\overline{y}) = \left(\frac{31/750}{2/25},\frac{7/750}{2/25}\right) = \left(\frac{31}{60},\frac{7}{60}\right).$

7. $m = \int_{-1}^1\int_0^{1-x^2}ky\,dy\,dx = k\int_{-1}^1\left[\frac{1}{2}y^2\right]_{y=0}^{y=1-x^2}dx = \frac{1}{2}k\int_{-1}^1(1-x^2)^2\,dx = \frac{1}{2}k\int_{-1}^1(1-2x^2+x^4)\,dx$

$= \frac{1}{2}k\left[x-\frac{2}{3}x^3+\frac{1}{5}x^5\right]_{-1}^1 = \frac{1}{2}k\left(1-\frac{2}{3}+\frac{1}{5}+1-\frac{2}{3}+\frac{1}{5}\right) = \frac{8}{15}k,$

$M_y = \int_{-1}^1\int_0^{1-x^2}kxy\,dy\,dx = k\int_{-1}^1\left[\frac{1}{2}xy^2\right]_{y=0}^{y=1-x^2}dx = \frac{1}{2}k\int_{-1}^1x(1-x^2)^2\,dx = \frac{1}{2}k\int_{-1}^1(x-2x^3+x^5)\,dx$

$= \frac{1}{2}k\left[\frac{1}{2}x^2-\frac{1}{2}x^4+\frac{1}{6}x^6\right]_{-1}^1 = \frac{1}{2}k\left(\frac{1}{2}-\frac{1}{2}+\frac{1}{6}-\frac{1}{2}+\frac{1}{2}-\frac{1}{6}\right) = 0,$

$M_x = \int_{-1}^1\int_0^{1-x^2}ky^2\,dy\,dx = k\int_{-1}^1\left[\frac{1}{3}y^3\right]_{y=0}^{y=1-x^2}dx = \frac{1}{3}k\int_{-1}^1(1-x^2)^3\,dx = \frac{1}{3}k\int_{-1}^1(1-3x^2+3x^4-x^6)\,dx$

$= \frac{1}{3}k\left[x-x^3+\frac{3}{5}x^5-\frac{1}{7}x^7\right]_{-1}^1 = \frac{1}{3}k\left(1-1+\frac{3}{5}-\frac{1}{7}+1-1+\frac{3}{5}-\frac{1}{7}\right) = \frac{32}{105}k.$

Hence $m=\frac{8}{15}k$, $(\overline{x},\overline{y}) = \left(0,\frac{32k/105}{8k/15}\right) = \left(0,\frac{4}{7}\right).$

8. The boundary curves intersect when $x+2=x^2 \iff x^2-x-2=0 \iff x=-1, x=2$. Thus here $D = \{(x,y) \mid -1 \le x \le 2,\ x^2 \le y \le x+2\}.$

$m = \int_{-1}^2\int_{x^2}^{x+2}kx^2\,dy\,dx = k\int_{-1}^2x^2\left[y\right]_{y=x^2}^{y=x+2}dx = k\int_{-1}^2(x^3+2x^2-x^4)\,dx$

$= k\left[\frac{1}{4}x^4+\frac{2}{3}x^3-\frac{1}{5}x^5\right]_{-1}^2 = k\left(\frac{44}{15}+\frac{13}{60}\right) = \frac{63}{20}k,$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$M_y = \int_{-1}^{2}\int_{x^2}^{x+2} kx^3\,dy\,dx = k\int_{-1}^{2} x^3\left[y\right]_{y=x^2}^{y=x+2}\,dx = k\int_{-1}^{2}(x^4+2x^3-x^5)\,dx$$

$$= k\left[\tfrac{1}{5}x^5+\tfrac{1}{2}x^4-\tfrac{1}{6}x^6\right]_{-1}^{2} = k\left(\tfrac{56}{15}-\tfrac{2}{15}\right) = \tfrac{18}{5}k,$$

$$M_x = \int_{-1}^{2}\int_{x^2}^{x+2} kx^2y\,dy\,dx = k\int_{-1}^{2} x^2\left[\tfrac{1}{2}y^2\right]_{y=x^2}^{y=x+2}\,dx = \tfrac{1}{2}k\int_{-1}^{2} x^2\left(x^2+4x+4-x^4\right)dx$$

$$= \tfrac{1}{2}k\int_{-1}^{2}(x^4+4x^3+4x^2-x^6)\,dx = \tfrac{1}{2}k\left[\tfrac{1}{5}x^5+x^4+\tfrac{4}{3}x^3-\tfrac{1}{7}x^7\right]_{-1}^{2} = \tfrac{1}{2}k\left(\tfrac{1552}{105}+\tfrac{41}{105}\right) = \tfrac{531}{70}k.$$

Hence $m = \frac{63}{20}k$, $(\overline{x},\overline{y}) = \left(\frac{18k/5}{63k/20}, \frac{531k/70}{63k/20}\right) = \left(\frac{8}{7}, \frac{118}{49}\right)$.

9. $m = \int_0^1\int_0^{e^{-x}} xy\,dy\,dx = \int_0^1 x\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=e^{-x}}dx = \tfrac{1}{2}\int_0^1 x\left(e^{-x}\right)^2 dx = \tfrac{1}{2}\int_0^1 xe^{-2x}\,dx$ [integrate by parts with $u = x$, $dv = e^{-2x}\,dx$]

$= \tfrac{1}{2}\left[-\tfrac{1}{4}(2x+1)e^{-2x}\right]_0^1 = -\tfrac{1}{8}\left(3e^{-2}-1\right) = \tfrac{1}{8} - \tfrac{3}{8}e^{-2}$.

$M_y = \int_0^1\int_0^{e^{-x}} x^2y\,dy\,dx = \int_0^1 x^2\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=e^{-x}}dx = \tfrac{1}{2}\int_0^1 x^2e^{-2x}\,dx$ [integrate by parts twice]

$= \tfrac{1}{2}\left[-\tfrac{1}{4}\left(2x^2+2x+1\right)e^{-2x}\right]_0^1 = -\tfrac{1}{8}\left(5e^{-2}-1\right) = \tfrac{1}{8} - \tfrac{5}{8}e^{-2}$.

$M_x = \int_0^1\int_0^{e^{-x}} xy^2\,dy\,dx = \int_0^1 x\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=e^{-x}}dx = \tfrac{1}{3}\int_0^1 xe^{-3x}\,dx$

$= \tfrac{1}{3}\left[-\tfrac{1}{9}(3x+1)e^{-3x}\right]_0^1 = -\tfrac{1}{27}\left(4e^{-3}-1\right) = \tfrac{1}{27} - \tfrac{4}{27}e^{-3}$.

Hence $m = \frac{1}{8}\left(1-3e^{-2}\right)$, $(\overline{x},\overline{y}) = \left(\frac{\frac{1}{8}\left(1-5e^{-2}\right)}{\frac{1}{8}\left(1-3e^{-2}\right)}, \frac{\frac{1}{27}\left(1-4e^{-3}\right)}{\frac{1}{8}\left(1-3e^{-2}\right)}\right) = \left(\frac{e^2-5}{e^2-3}, \frac{8\left(e^3-4\right)}{27\left(e^3-3e\right)}\right)$.

10. Note that $\cos x \geq 0$ for $-\pi/2 \leq x \leq \pi/2$.

$m = \int_{-\pi/2}^{\pi/2}\int_0^{\cos x} y\,dy\,dx = \int_{-\pi/2}^{\pi/2}\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=\cos x}dx = \tfrac{1}{2}\int_{-\pi/2}^{\pi/2}\cos^2 x\,dx = \tfrac{1}{2}\left[\tfrac{1}{2}x+\tfrac{1}{4}\sin 2x\right]_{-\pi/2}^{\pi/2} = \tfrac{\pi}{4}$,

$M_y = \int_{-\pi/2}^{\pi/2}\int_0^{\cos x} xy\,dy\,dx = \int_{-\pi/2}^{\pi/2} x\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=\cos x}dx = \tfrac{1}{2}\int_{-\pi/2}^{\pi/2} x\cos^2 x\,dx$ [integrate by parts with $u = x$, $dv = \cos^2 x\,dx$]

$= \tfrac{1}{2}\left[x\left(\tfrac{1}{2}x+\tfrac{1}{4}\sin 2x\right)\Big|_{-\pi/2}^{\pi/2} - \int_{-\pi/2}^{\pi/2}\left(\tfrac{1}{2}x+\tfrac{1}{4}\sin 2x\right)dx\right]$

$= \tfrac{1}{2}\left(\tfrac{1}{8}\pi^2 - \tfrac{1}{8}\pi^2 - \left[\tfrac{1}{4}x^2-\tfrac{1}{8}\cos 2x\right]_{-\pi/2}^{\pi/2}\right) = \tfrac{1}{2}\left(0-\left[\tfrac{1}{16}\pi^2+\tfrac{1}{8}-\tfrac{1}{16}\pi^2-\tfrac{1}{8}\right]\right) = 0$,

$M_x = \int_{-\pi/2}^{\pi/2}\int_0^{\cos x} y^2\,dy\,dx = \int_{-\pi/2}^{\pi/2}\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=\cos x}dx = \tfrac{1}{3}\int_{-\pi/2}^{\pi/2}\cos^3 x\,dx = \tfrac{1}{3}\int_{-\pi/2}^{\pi/2}(1-\sin^2 x)\cos x\,dx$

[substitute $u = \sin x \quad \Rightarrow \quad du = \cos x\,dx$]

$= \tfrac{1}{3}\left[\sin x - \tfrac{1}{3}\sin^3 x\right]_{-\pi/2}^{\pi/2} = \tfrac{1}{3}\left(1-\tfrac{1}{3}+1-\tfrac{1}{3}\right) = \tfrac{4}{9}$.

Hence $m = \frac{\pi}{4}$, $(\overline{x},\overline{y}) = \left(0, \frac{4/9}{\pi/4}\right) = \left(0, \frac{16}{9\pi}\right)$.

11. $\rho(x,y) = ky$, $m = \iint_D ky\,dA = \int_0^{\pi/2}\int_0^1 k(r\sin\theta)\,r\,dr\,d\theta = k\int_0^{\pi/2}\sin\theta\,d\theta\int_0^1 r^2\,dr$

$= k\left[-\cos\theta\right]_0^{\pi/2}\left[\tfrac{1}{3}r^3\right]_0^1 = k(1)\left(\tfrac{1}{3}\right) = \tfrac{1}{3}k$,

$M_y = \iint_D x\cdot ky\,dA = \int_0^{\pi/2}\int_0^1 k(r\cos\theta)(r\sin\theta)\,r\,dr\,d\theta = k\int_0^{\pi/2}\sin\theta\cos\theta\,d\theta\int_0^1 r^3\,dr$

$= k\left[\tfrac{1}{2}\sin^2\theta\right]_0^{\pi/2}\left[\tfrac{1}{4}r^4\right]_0^1 = k\left(\tfrac{1}{2}\right)\left(\tfrac{1}{4}\right) = \tfrac{1}{8}k$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$M_x = \iint_D y \cdot ky\,dA = \int_0^{\pi/2}\int_0^1 k(r\sin\theta)^2\, r\,dr\,d\theta = k\int_0^{\pi/2}\sin^2\theta\,d\theta\int_0^1 r^3\,dr$

$= k\left[\frac{1}{2}\theta - \frac{1}{4}\sin 2\theta\right]_0^{\pi/2}\left[\frac{1}{4}r^4\right]_0^1 = k\left(\frac{\pi}{4}\right)\left(\frac{1}{4}\right) = \frac{\pi}{16}k.$

Hence $(\overline{x},\overline{y}) = \left(\frac{k/8}{k/3}, \frac{k\pi/16}{k/3}\right) = \left(\frac{3}{8}, \frac{3\pi}{16}\right)$.

12. $\rho(x,y) = k(x^2+y^2) = kr^2$, $m = \int_0^{\pi/2}\int_0^1 kr^3\,dr\,d\theta = \frac{\pi}{8}k$,

$M_y = \int_0^{\pi/2}\int_0^1 kr^4\cos\theta\,dr\,d\theta = \frac{1}{5}k\int_0^{\pi/2}\cos\theta\,d\theta = \frac{1}{5}k\left[\sin\theta\right]_0^{\pi/2} = \frac{1}{5}k$,

$M_x = \int_0^{\pi/2}\int_0^1 kr^4\sin\theta\,dr\,d\theta = \frac{1}{5}k\int_0^{\pi/2}\sin\theta\,d\theta = \frac{1}{5}k\left[-\cos\theta\right]_0^{\pi/2} = \frac{1}{5}k$.

Hence $(\overline{x},\overline{y}) = \left(\frac{8}{5\pi}, \frac{8}{5\pi}\right)$.

13.



$\rho(x,y) = k\sqrt{x^2+y^2} = kr$,

$m = \iint_D \rho(x,y)dA = \int_0^{\pi}\int_1^2 kr\cdot r\,dr\,d\theta$

$= k\int_0^{\pi} d\theta\int_1^2 r^2\,dr = k(\pi)\left[\frac{1}{3}r^3\right]_1^2 = \frac{7}{3}\pi k,$

$M_y = \iint_D x\rho(x,y)dA = \int_0^{\pi}\int_1^2 (r\cos\theta)(kr)\,r\,dr\,d\theta = k\int_0^{\pi}\cos\theta\,d\theta\int_1^2 r^3\,dr$

$= k\left[\sin\theta\right]_0^{\pi}\left[\frac{1}{4}r^4\right]_1^2 = k(0)\left(\frac{15}{4}\right) = 0$ [this is to be expected as the region and density function are symmetric about the y-axis]

$M_x = \iint_D y\rho(x,y)dA = \int_0^{\pi}\int_1^2 (r\sin\theta)(kr)\,r\,dr\,d\theta = k\int_0^{\pi}\sin\theta\,d\theta\int_1^2 r^3\,dr$

$= k\left[-\cos\theta\right]_0^{\pi}\left[\frac{1}{4}r^4\right]_1^2 = k(1+1)\left(\frac{15}{4}\right) = \frac{15}{2}k.$

Hence $(\overline{x},\overline{y}) = \left(0, \frac{15k/2}{7\pi k/3}\right) = \left(0, \frac{45}{14\pi}\right)$.

14. Now $\rho(x,y) = k/\sqrt{x^2+y^2} = k/r$, so

$m = \iint_D \rho(x,y)dA = \int_0^{\pi}\int_1^2 (k/r)\,r\,dr\,d\theta = k\int_0^{\pi} d\theta\int_1^2 dr = k(\pi)(1) = \pi k,$

$M_y = \iint_D x\rho(x,y)dA = \int_0^{\pi}\int_1^2 (r\cos\theta)(k/r)\,r\,dr\,d\theta = k\int_0^{\pi}\cos\theta\,d\theta\int_1^2 r\,dr$

$= k\left[\sin\theta\right]_0^{\pi}\left[\frac{1}{2}r^2\right]_1^2 = k(0)\left(\frac{3}{2}\right) = 0,$

$M_x = \iint_D y\rho(x,y)dA = \int_0^{\pi}\int_1^2 (r\sin\theta)(k/r)\,r\,dr\,d\theta = k\int_0^{\pi}\sin\theta\,d\theta\int_1^2 r\,dr$

$= k\left[-\cos\theta\right]_0^{\pi}\left[\frac{1}{2}r^2\right]_1^2 = k(1+1)\left(\frac{3}{2}\right) = 3k.$

Hence $(\overline{x},\overline{y}) = \left(0, \frac{3k}{\pi k}\right) = \left(0, \frac{3}{\pi}\right)$.

15. Placing the vertex opposite the hypotenuse at $(0,0)$, $\rho(x,y) = k(x^2+y^2)$. Then

$m = \int_0^a\int_0^{a-x} k\left(x^2+y^2\right)dy\,dx = k\int_0^a\left[ax^2 - x^3 + \frac{1}{3}(a-x)^3\right]dx = k\left[\frac{1}{3}ax^3 - \frac{1}{4}x^4 - \frac{1}{12}(a-x)^4\right]_0^a = \frac{1}{6}ka^4.$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

By symmetry, $M_y = M_x = \int_0^a \int_0^{a-x} ky(x^2+y^2)\,dy\,dx = k\int_0^a \left[\frac{1}{2}(a-x)^2x^2 + \frac{1}{4}(a-x)^4\right] dx$

$= k\left[\frac{1}{6}a^2x^3 - \frac{1}{4}ax^4 + \frac{1}{10}x^5 - \frac{1}{20}(a-x)^5\right]_0^a = \frac{1}{15}ka^5$

Hence $(\overline{x}, \overline{y}) = \left(\frac{2}{5}a, \frac{2}{5}a\right)$.

16. $\rho(x,y) = k/\sqrt{x^2+y^2} = k/r$.



$$m = \int_{\pi/6}^{5\pi/6}\int_1^{2\sin\theta} \frac{k}{r}\, r\, dr\, d\theta = k\int_{\pi/6}^{5\pi/6} [(2\sin\theta) - 1]\, d\theta$$

$$= k\left[-2\cos\theta - \theta\right]_{\pi/6}^{5\pi/6} = 2k\left(\sqrt{3} - \frac{\pi}{3}\right)$$

By symmetry of D and $f(x) = x$, $M_y = 0$, and

$$M_x = \int_{\pi/6}^{5\pi/6}\int_1^{2\sin\theta} kr\sin\theta\, dr\, d\theta = \frac{1}{2}k\int_{\pi/6}^{5\pi/6}(4\sin^3\theta - \sin\theta)\, d\theta$$

$$= \frac{1}{2}k\left[-3\cos\theta + \frac{4}{3}\cos^3\theta\right]_{\pi/6}^{5\pi/6} = \sqrt{3}\,k$$

Hence $(\overline{x}, \overline{y}) = \left(0, \frac{3\sqrt{3}}{2(3\sqrt{3}-\pi)}\right)$.

17. $I_x = \iint_D y^2\rho(x,y)dA = \int_1^3\int_1^4 y^2\cdot ky^2\,dy\,dx = k\int_1^3 dx\int_1^4 y^4\,dy = k\,[x]_1^3\left[\frac{1}{5}y^5\right]_1^4 = k(2)\left(\frac{1023}{5}\right) = 409.2k,$

$I_y = \iint_D x^2\rho(x,y)\,dA = \int_1^3\int_1^4 x^2\cdot ky^2\,dy\,dx = k\int_1^3 x^2\,dx\int_1^4 y^2\,dy = k\left[\frac{1}{3}x^3\right]_1^3\left[\frac{1}{3}y^3\right]_1^4 = k\left(\frac{26}{3}\right)(21) = 182k,$

and $I_0 = I_x + I_y = 409.2k + 182k = 591.2k$.

18. $I_x = \iint_D y^2\rho(x,y)dA = \int_0^{2/5}\int_{y/2}^{1-2y} y^2\cdot x\,dx\,dy = \int_0^{2/5} y^2\left[\frac{1}{2}x^2\right]_{x=y/2}^{x=1-2y} dy = \frac{1}{2}\int_0^{2/5} y^2\left(\frac{15}{4}y^2 - 4y + 1\right) dy$

$= \frac{1}{2}\int_0^{2/5}\left(\frac{15}{4}y^4 - 4y^3 + y^2\right) dy = \frac{1}{2}\left[\frac{3}{4}y^5 - y^4 + \frac{1}{3}y^3\right]_0^{2/5} = \frac{16}{9375},$

$I_y = \iint_D x^2\rho(x,y)\,dA = \int_0^{2/5}\int_{y/2}^{1-2y} x^2\cdot x\,dx\,dy = \int_0^{2/5}\left[\frac{1}{4}x^4\right]_{x=y/2}^{x=1-2y} dy = \frac{1}{4}\int_0^{2/5}\left[(1-2y)^4 - \frac{1}{16}y^4\right] dy$

$= \frac{1}{4}\int_0^{2/5}\left(\frac{255}{16}y^4 - 32y^3 + 24y^2 - 8y + 1\right) dy = \frac{1}{4}\left[\frac{51}{16}y^5 - 8y^4 + 8y^3 - 4y^2 + y\right]_0^{2/5} = \frac{78}{3125},$

and $I_0 = I_x + I_y = \frac{16}{9375} + \frac{78}{3125} = \frac{2}{75}$.

19. As in Exercise 15, we place the vertex opposite the hypotenuse at $(0,0)$ and the equal sides along the positive axes.

$I_x = \int_0^a\int_0^{a-x} y^2k(x^2+y^2)\,dy\,dx = k\int_0^a\int_0^{a-x}(x^2y^2 + y^4)\,dy\,dx = k\int_0^a\left[\frac{1}{3}x^2y^3 + \frac{1}{5}y^5\right]_{y=0}^{y=a-x} dx$

$= k\int_0^a\left[\frac{1}{3}x^2(a-x)^3 + \frac{1}{5}(a-x)^5\right] dx = k\left[\frac{1}{3}\left(\frac{1}{3}a^3x^3 - \frac{3}{4}a^2x^4 + \frac{3}{5}ax^5 - \frac{1}{6}x^6\right) - \frac{1}{30}(a-x)^6\right]_0^a = \frac{7}{180}ka^6,$

$I_y = \int_0^a\int_0^{a-x} x^2k(x^2+y^2)\,dy\,dx = k\int_0^a\int_0^{a-x}(x^4 + x^2y^2)\,dy\,dx = k\int_0^a\left[x^4y + \frac{1}{3}x^2y^3\right]_{y=0}^{y=a-x} dx$

$= k\int_0^a\left[x^4(a-x) + \frac{1}{3}x^2(a-x)^3\right] dx = k\left[\frac{1}{5}ax^5 - \frac{1}{6}x^6 + \frac{1}{3}\left(\frac{1}{3}a^3x^3 - \frac{3}{4}a^2x^4 + \frac{3}{5}ax^5 - \frac{1}{6}x^6\right)\right]_0^a = \frac{7}{180}ka^6,$

and $I_0 = I_x + I_y = \frac{7}{90}ka^6$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. If we find the moments of inertia about the x- and y-axes, we can determine in which direction rotation will be more difficult. (See the explanation following Example 4.) The moment of inertia about the x-axis is given by

$$I_x = \iint_D y^2\rho(x,y)\,dA = \int_0^2\int_0^2 y^2(1+0.1x)\,dy\,dx = \int_0^2 (1+0.1x)\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=2} dx$$

$$= \tfrac{8}{3}\int_0^2 (1+0.1x)\,dx = \tfrac{8}{3}\left[x+0.1\cdot\tfrac{1}{2}x^2\right]_0^2 = \tfrac{8}{3}(2.2) \approx 5.87$$

Similarly, the moment of inertia about the y-axis is given by

$$I_y = \iint_D x^2\rho(x,y)\,dA = \int_0^2\int_0^2 x^2(1+0.1x)\,dy\,dx = \int_0^2 x^2(1+0.1x)\left[y\right]_{y=0}^{y=2} dx$$

$$= 2\int_0^2 (x^2+0.1x^3)\,dx = 2\left[\tfrac{1}{3}x^3+0.1\cdot\tfrac{1}{4}x^4\right]_0^2 = 2\left(\tfrac{8}{3}+0.4\right) \approx 6.13$$

Since $I_y > I_x$, more force is required to rotate the fan blade about the y-axis.

21. $I_x = \iint_D y^2\rho(x,y)dA = \int_0^h\int_0^b \rho y^2\,dx\,dy = \rho\int_0^b dx\,\int_0^h y^2\,dy = \rho\left[x\right]_0^b\,\left[\tfrac{1}{3}y^3\right]_0^h = \rho b\left(\tfrac{1}{3}h^3\right) = \tfrac{1}{3}\rho bh^3$,

$I_y = \iint_D x^2\rho(x,y)dA = \int_0^h\int_0^b \rho x^2\,dx\,dy = \rho\int_0^b x^2\,dx\,\int_0^h dy = \rho\left[\tfrac{1}{3}x^3\right]_0^b\,[y]_0^h = \tfrac{1}{3}\rho b^3h$,

and $m = \rho$(area of rectangle) $= \rho bh$ since the lamina is homogeneous. Hence $\overline{\overline{x}}^2 = \dfrac{I_y}{m} = \dfrac{\frac{1}{3}\rho b^3h}{\rho bh} = \dfrac{b^2}{3} \;\Rightarrow\; \overline{\overline{x}} = \dfrac{b}{\sqrt{3}}$

and $\overline{\overline{y}}^2 = \dfrac{I_x}{m} = \dfrac{\frac{1}{3}\rho bh^3}{\rho bh} = \dfrac{h^2}{3} \;\Rightarrow\; \overline{\overline{y}} = \dfrac{h}{\sqrt{3}}$.

22. Here we assume $b > 0$, $h > 0$ but note that we arrive at the same results if $b < 0$ or $h < 0$. We have

$D = \left\{(x,y) \mid 0 \le x \le b, 0 \le y \le h - \frac{h}{b}x\right\}$, so

$I_x = \int_0^b\int_0^{h-hx/b} y^2\rho\,dy\,dx = \rho\int_0^b\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=h-hx/b} dx = \tfrac{1}{3}\rho\int_0^b\left(h-\tfrac{h}{b}x\right)^3 dx$

$= \tfrac{1}{3}\rho\left[-\tfrac{b}{h}\left(\tfrac{1}{4}\right)\left(h-\tfrac{h}{b}x\right)^4\right]_0^b = -\tfrac{b}{12h}\rho(0-h^4) = \tfrac{1}{12}\rho bh^3$,

$I_y = \int_0^b\int_0^{h-hx/b} x^2\rho\,dy\,dx = \rho\int_0^b x^2\left(h-\tfrac{h}{b}x\right)dx = \rho\int_0^b\left(hx^2-\tfrac{h}{b}x^3\right)dx$

$= \rho\left[\tfrac{h}{3}x^3-\tfrac{h}{4b}x^4\right]_0^b = \rho(\tfrac{hb^3}{3}-\tfrac{hb^3}{4}) = \tfrac{1}{12}\rho b^3h$,

and $m = \int_0^b\int_0^{h-hx/b}\rho\,dy\,dx = \rho\int_0^b\left(h-\tfrac{h}{b}x\right)dx = \rho\left[hx-\tfrac{h}{2b}x^2\right]_0^b = \tfrac{1}{2}\rho bh$. Hence $\overline{\overline{x}}^2 = \dfrac{I_y}{m} = \dfrac{\frac{1}{12}\rho b^3h}{\frac{1}{2}\rho bh} = \dfrac{b^2}{6} \;\Rightarrow$

$\overline{\overline{x}} = \dfrac{b}{\sqrt{6}}$ and $\overline{\overline{y}}^2 = \dfrac{I_x}{m} = \dfrac{\frac{1}{12}\rho bh^3}{\frac{1}{2}\rho bh} = \dfrac{h^2}{6} \;\Rightarrow\; \overline{\overline{y}} = \dfrac{h}{\sqrt{6}}$.

23. In polar coordinates, the region is $D = \left\{(r,\theta) \mid 0 \le r \le a, 0 \le \theta \le \frac{\pi}{2}\right\}$, so

$I_x = \iint_D y^2\rho\,dA = \int_0^{\pi/2}\int_0^a \rho(r\sin\theta)^2\,r\,dr\,d\theta = \rho\int_0^{\pi/2}\sin^2 d\theta\,\int_0^a r^3\,dr$

$= \rho\left[\tfrac{1}{2}\theta-\tfrac{1}{4}\sin 2\theta\right]_0^{\pi/2}\,\left[\tfrac{1}{4}r^4\right]_0^a = \rho\left(\tfrac{\pi}{4}\right)\left(\tfrac{1}{4}a^4\right) = \tfrac{1}{16}\rho a^4\pi$,

$I_y = \iint_D x^2\rho\,dA = \int_0^{\pi/2}\int_0^a \rho(r\cos\theta)^2\,r\,dr\,d\theta = \rho\int_0^{\pi/2}\cos^2 d\theta\,\int_0^a r^3\,dr$

$= \rho\left[\tfrac{1}{2}\theta+\tfrac{1}{4}\sin 2\theta\right]_0^{\pi/2}\,\left[\tfrac{1}{4}r^4\right]_0^a = \rho\left(\tfrac{\pi}{4}\right)\left(\tfrac{1}{4}a^4\right) = \tfrac{1}{16}\rho a^4\pi$,

and $m = \rho\cdot A(D) = \rho\cdot\frac{1}{4}\pi a^2$ since the lamina is homogeneous. Hence $\overline{\overline{x}}^2 = \overline{\overline{y}}^2 = \dfrac{\frac{1}{16}\rho a^4\pi}{\frac{1}{4}\rho a^2\pi} = \dfrac{a^2}{4} \;\Rightarrow\; \overline{\overline{x}} = \overline{\overline{y}} = \dfrac{a}{2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $m = \int_0^{\pi}\int_0^{\sin x} \rho\, dy\, dx = \rho\int_0^{\pi} \sin x\, dx = \rho\big[-\cos x\big]_0^{\pi} = 2\rho$,

$I_x = \int_0^{\pi}\int_0^{\sin x} \rho y^2\, dy\, dx = \frac{1}{3}\rho\int_0^{\pi} \sin^3 x\, dx = \frac{1}{3}\rho\int_0^{\pi}(1-\cos^2 x)\sin x\, dx = \frac{1}{3}\rho\big[-\cos x + \frac{1}{3}\cos^3 x\big]_0^{\pi} = \frac{4}{9}\rho$,

$I_y = \int_0^{\pi}\int_0^{\sin x} \rho x^2\, dy\, dx = \rho\int_0^{\pi} x^2 \sin x\, dx = \rho\big[-x^2\cos x + 2x\sin x + 2\cos x\big]_0^{\pi}$ [by integrating by parts twice]

$= \rho(\pi^2 - 4)$.

Then $\overline{\overline{y}}^2 = \dfrac{I_x}{m} = \dfrac{2}{9}$, so $\overline{\overline{y}} = \dfrac{\sqrt{2}}{3}$ and $\overline{\overline{x}}^2 = \dfrac{I_y}{m} = \dfrac{\pi^2-4}{2}$, so $\overline{\overline{x}} = \sqrt{\dfrac{\pi^2-4}{2}}$.

25. The right loop of the curve is given by $D = \{(r,\theta) \mid 0 \le r \le \cos 2\theta,\ -\pi/4 \le \theta \le \pi/4\}$. Using a CAS, we find $m = \iint_D \rho(x,y)\, dA = \iint_D (x^2+y^2)\, dA = \int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} r^2\, r\, dr\, d\theta = \dfrac{3\pi}{64}$. Then

$$\overline{x} = \frac{1}{m}\iint_D x\rho(x,y)\, dA = \frac{64}{3\pi}\int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} (r\cos\theta)\, r^2\, r\, dr\, d\theta = \frac{64}{3\pi}\int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} r^4\cos\theta\, dr\, d\theta = \frac{16384\sqrt{2}}{10395\pi} \text{ and}$$

$$\overline{y} = \frac{1}{m}\iint_D y\rho(x,y)\, dA = \frac{64}{3\pi}\int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} (r\sin\theta)\, r^2\, r\, dr\, d\theta = \frac{64}{3\pi}\int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} r^4\sin\theta\, dr\, d\theta = 0, \text{ so}$$

$$(\overline{x},\overline{y}) = \left(\frac{16384\sqrt{2}}{10395\pi}, 0\right).$$

The moments of inertia are

$$I_x = \iint_D y^2\rho(x,y)\, dA = \int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} (r\sin\theta)^2\, r^2\, r\, dr\, d\theta = \int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} r^5\sin^2\theta\, dr\, d\theta = \frac{5\pi}{384} - \frac{4}{105},$$

$$I_y = \iint_D x^2\rho(x,y)\, dA = \int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} (r\cos\theta)^2\, r^2\, r\, dr\, d\theta = \int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} r^5\cos^2\theta\, dr\, d\theta = \frac{5\pi}{384} + \frac{4}{105}, \text{ and}$$

$$I_0 = I_x + I_y = \frac{5\pi}{192}.$$

26. Using a CAS, we find $m = \iint_D \rho(x,y)\, dA = \int_0^2\int_0^{xe^{-x}} x^2y^2\, dy\, dx = \frac{8}{729}(5 - 899e^{-6})$. Then

$$\overline{x} = \frac{1}{m}\iint_D x\rho(x,y)\, dA = \frac{729}{8(5-899e^{-6})}\int_0^2\int_0^{xe^{-x}} x^3y^2\, dy\, dx = \frac{2(5e^6-1223)}{5e^6-899} \text{ and}$$

$$\overline{y} = \frac{1}{m}\iint_D y\rho(x,y)\, dA = \frac{729}{8(5-899e^{-6})}\int_0^2\int_0^{xe^{-x}} x^2y^3\, dy\, dx = \frac{729(45e^6-42037e^{-2})}{32768(5e^6-899)}, \text{ so}$$

$$(\overline{x},\overline{y}) = \left(\frac{2(5e^6-1223)}{5e^6-899}, \frac{729(45e^6-42037e^{-2})}{32768(5e^6-899)}\right).$$

The moments of inertia are $I_x = \iint_D y^2\rho(x,y)\, dA = \int_0^2\int_0^{xe^{-x}} x^2y^4\, dy\, dx = \frac{16}{390625}(63 - 305593e^{-10})$,

$I_y = \iint_D x^2\rho(x,y)\, dA = \int_0^2\int_0^{xe^{-x}} x^4y^2\, dy\, dx = \frac{80}{2187}(7 - 2101e^{-6})$, and

$I_0 = I_x + I_y = \frac{16}{854296875}(13809656 - 4103515625e^{-6} - 668331891e^{-10})$.

27. (a) $f(x,y)$ is a joint density function, so we know $\iint_{\mathbb{R}^2} f(x,y)\, dA = 1$. Since $f(x,y) = 0$ outside the rectangle $[0,1]\times[0,2]$, we can say

$$\iint_{\mathbb{R}^2} f(x,y)\, dA = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y)\, dy\, dx = \int_0^1\int_0^2 Cx(1+y)\, dy\, dx$$
$$= C\int_0^1 x\big[y + \tfrac{1}{2}y^2\big]_{y=0}^{y=2}\, dx = C\int_0^1 4x\, dx = C\big[2x^2\big]_0^1 = 2C$$

Then $2C = 1 \Rightarrow C = \frac{1}{2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $P(X \le 1, Y \le 1) = \int_{-\infty}^{1}\int_{-\infty}^{1} f(x,y)\,dy\,dx = \int_0^1\int_0^1 \frac{1}{2}x(1+y)\,dy\,dx$

$= \int_0^1 \frac{1}{2}x\left[y + \frac{1}{2}y^2\right]_{y=0}^{y=1} dx = \int_0^1 \frac{1}{2}x\left(\frac{3}{2}\right) dx = \frac{3}{4}\left[\frac{1}{2}x^2\right]_0^1 = \frac{3}{8}$ or 0.375

(c) $P(X+Y \le 1) = P((X,Y) \in D)$ where D is the triangular region shown in the figure. Thus



$$P(X+Y \le 1) = \iint_D f(x,y)\,dA = \int_0^1\int_0^{1-x} \frac{1}{2}x(1+y)\,dy\,dx$$

$$= \int_0^1 \frac{1}{2}x\left[y + \frac{1}{2}y^2\right]_{y=0}^{y=1-x} dx = \int_0^1 \frac{1}{2}x\left(\frac{1}{2}x^2 - 2x + \frac{3}{2}\right) dx$$

$$= \frac{1}{4}\int_0^1 (x^3 - 4x^2 + 3x)\,dx = \frac{1}{4}\left[\frac{x^4}{4} - 4\frac{x^3}{3} + 3\frac{x^2}{2}\right]_0^1$$

$$= \frac{5}{48} \approx 0.1042$$

28. (a) $f(x,y) \ge 0$, so f is a joint density function if $\iint_{\mathbb{R}^2} f(x,y)\,dA = 1$. Here, $f(x,y) = 0$ outside the square $[0,1] \times [0,1]$, so $\iint_{\mathbb{R}^2} f(x,y)\,dA = \int_0^1\int_0^1 4xy\,dy\,dx = \int_0^1 \left[2xy^2\right]_{y=0}^{y=1} dx = \int_0^1 2x\,dx = x^2\big]_0^1 = 1$.

Thus, $f(x,y)$ is a joint density function.

(b) (i) No restriction is placed on Y, so

$$P(X \ge \tfrac{1}{2}) = \int_{1/2}^{\infty}\int_{-\infty}^{\infty} f(x,y)\,dy\,dx = \int_{1/2}^1\int_0^1 4xy\,dy\,dx = \int_{1/2}^1 \left[2xy^2\right]_{y=0}^{y=1} dx = \int_{1/2}^1 2x\,dx = x^2\big]_{1/2}^1 = \frac{3}{4}.$$

(ii) $P(X \ge \frac{1}{2}, Y \le \frac{1}{2}) = \int_{1/2}^{\infty}\int_{\infty}^{1/2} f(x,y)\,dy\,dx = \int_{1/2}^1\int_0^{1/2} 4xy\,dy\,dx$

$$= \int_{1/2}^1 \left[2xy^2\right]_{y=0}^{y=1/2} dx = \int_{1/2}^1 \frac{1}{2}x\,dx = \frac{1}{2}\cdot\frac{1}{2}x^2\big]_{1/2}^1 = \frac{3}{16}$$

(c) The expected value of X is given by

$$\mu_1 = \iint_{\mathbb{R}^2} x\,f(x,y)\,dA = \int_0^1\int_0^1 x(4xy)\,dy\,dx = \int_0^1 2x^2\left[y^2\right]_{y=0}^{y=1} dx = 2\int_0^1 x^2\,dx = 2\left[\tfrac{1}{3}x^3\right]_0^1 = \frac{2}{3}$$

The expected value of Y is

$$\mu_2 = \iint_{\mathbb{R}^2} y\,f(x,y)\,dA = \int_0^1\int_0^1 y(4xy)\,dy\,dx = \int_0^1 4x\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=1} dx = \frac{4}{3}\int_0^1 x\,dx = \frac{4}{3}\left[\tfrac{1}{2}x^2\right]_0^1 = \frac{2}{3}$$

29. (a) $f(x,y) \ge 0$, so f is a joint density function if $\iint_{\mathbb{R}^2} f(x,y)\,dA = 1$. Here, $f(x,y) = 0$ outside the first quadrant, so

$$\iint_{\mathbb{R}^2} f(x,y)\,dA = \int_0^{\infty}\int_0^{\infty} 0.1e^{-(0.5x+0.2y)}\,dy\,dx = 0.1\int_0^{\infty}\int_0^{\infty} e^{-0.5x}e^{-0.2y}\,dy\,dx = 0.1\int_0^{\infty} e^{-0.5x}\,dx \int_0^{\infty} e^{-0.2y}\,dy$$

$$= 0.1\lim_{t\to\infty}\int_0^t e^{-0.5x}\,dx \lim_{t\to\infty}\int_0^t e^{-0.2y}\,dy = 0.1\lim_{t\to\infty}\left[-2e^{-0.5x}\right]_0^t \lim_{t\to\infty}\left[-5e^{-0.2y}\right]_0^t$$

$$= 0.1\lim_{t\to\infty}\left[-2(e^{-0.5t}-1)\right]\lim_{t\to\infty}\left[-5(e^{-0.2t}-1)\right] = (0.1)\cdot(-2)(0-1)\cdot(-5)(0-1) = 1$$

Thus $f(x,y)$ is a joint density function.

(b) (i) No restriction is placed on X, so

$$P(Y \ge 1) = \int_{-\infty}^{\infty}\int_1^{\infty} f(x,y)\,dy\,dx = \int_0^{\infty}\int_1^{\infty} 0.1e^{-(0.5x+0.2y)}\,dy\,dx$$

$$= 0.1\int_0^{\infty} e^{-0.5x}\,dx \int_1^{\infty} e^{-0.2y}\,dy = 0.1\lim_{t\to\infty}\int_0^t e^{-0.5x}\,dx \lim_{t\to\infty}\int_1^t e^{-0.2y}\,dy$$

$$= 0.1\lim_{t\to\infty}\left[-2e^{-0.5x}\right]_0^t \lim_{t\to\infty}\left[-5e^{-0.2y}\right]_1^t = 0.1\lim_{t\to\infty}\left[-2(e^{-0.5t}-1)\right]\lim_{t\to\infty}\left[-5(e^{-0.2t}-e^{-0.2})\right]$$

$$(0.1)\cdot(-2)(0-1)\cdot(-5)(0-e^{-0.2}) = e^{-0.2} \approx 0.8187$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(ii) $P(X \le 2, Y \le 4) = \int_{-\infty}^{2}\int_{-\infty}^{4} f(x,y)\,dy\,dx = \int_0^2\int_0^4 0.1e^{-(0.5x+0.2y)}\,dy\,dx$

$= 0.1\int_0^2 e^{-0.5x}\,dx \int_0^4 e^{-0.2y}\,dy = 0.1\left[-2e^{-0.5x}\right]_0^2\left[-5e^{-0.2y}\right]_0^4$

$= (0.1)\cdot(-2)(e^{-1}-1)\cdot(-5)(e^{-0.8}-1)$

$= (e^{-1}-1)(e^{-0.8}-1) = 1 + e^{-1.8} - e^{-0.8} - e^{-1} \approx 0.3481$

(c) The expected value of X is given by

$$\mu_1 = \iint_{\mathbb{R}^2} x\,f(x,y)\,dA = \int_0^\infty\int_0^\infty x\left[0.1e^{-(0.5x+0.2y)}\right]dy\,dx$$

$$= 0.1\int_0^\infty xe^{-0.5x}\,dx \int_0^\infty e^{-0.2y}\,dy = 0.1\lim_{t\to\infty}\int_0^t xe^{-0.5x}\,dx \lim_{t\to\infty}\int_0^t e^{-0.2y}\,dy$$

To evaluate the first integral, we integrate by parts with $u = x$ and $dv = e^{-0.5x}\,dx$ (or we can use Formula 96 in the Table of Integrals): $\int xe^{-0.5x}\,dx = -2xe^{-0.5x} - \int -2e^{-0.5x}\,dx = -2xe^{-0.5x} - 4e^{-0.5x} = -2(x+2)e^{-0.5x}$.

Thus

$$\mu_1 = 0.1\lim_{t\to\infty}\left[-2(x+2)e^{-0.5x}\right]_0^t \lim_{t\to\infty}\left[-5e^{-0.2y}\right]_0^t$$

$$= 0.1\lim_{t\to\infty}(-2)\left[(t+2)e^{-0.5t} - 2\right]\lim_{t\to\infty}(-5)\left[e^{-0.2t}-1\right]$$

$$= 0.1(-2)\left(\lim_{t\to\infty}\frac{t+2}{e^{0.5t}} - 2\right)(-5)(-1) = 2 \qquad \text{[by l'Hospital's Rule]}$$

The expected value of Y is given by

$$\mu_2 = \iint_{\mathbb{R}^2} y\,f(x,y)\,dA = \int_0^\infty\int_0^\infty y\left[0.1e^{-(0.5+0.2y)}\right]dy\,dx$$

$$= 0.1\int_0^\infty e^{-0.5x}\,dx \int_0^\infty ye^{-0.2y}\,dy = 0.1\lim_{t\to\infty}\int_0^t e^{-0.5x}\,dx \lim_{t\to\infty}\int_0^t ye^{-0.2y}\,dy$$

To evaluate the second integral, we integrate by parts with $u = y$ and $dv = e^{-0.2y}\,dy$ (or again we can use Formula 96 in the Table of Integrals) which gives $\int ye^{-0.2y}\,dy = -5ye^{-0.2y} + \int 5e^{-0.2y}\,dy = -5(y+5)e^{-0.2y}$. Then

$$\mu_2 = 0.1\lim_{t\to\infty}\left[-2e^{-0.5x}\right]_0^t \lim_{t\to\infty}\left[-5(y+5)e^{-0.2y}\right]_0^t$$

$$= 0.1\lim_{t\to\infty}\left[-2(e^{-0.5t}-1)\right]\lim_{t\to\infty}\left(-5\left[(t+5)e^{-0.2t} - 5\right]\right)$$

$$= 0.1(-2)(-1)\cdot(-5)\left(\lim_{t\to\infty}\frac{t+5}{e^{0.2t}} - 5\right) = 5 \qquad \text{[by l'Hospital's Rule]}$$

30. (a) The lifetime of each bulb has exponential density function

$$f(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{1}{1000}e^{-t/1000} & \text{if } t \ge 0 \end{cases}$$

If X and Y are the lifetimes of the individual bulbs, then X and Y are independent, so the joint density function is the product of the individual density functions:

$$f(x,y) = \begin{cases} 10^{-6}e^{-(x+y)/1000} & \text{if } x \ge 0, y \ge 0 \\ 0 & \text{otherwise} \end{cases}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The probability that both of the bulbs fail within 1000 hours is

$$\begin{aligned} P(X \le 1000, Y \le 1000) &= \int_{-\infty}^{1000}\int_{-\infty}^{1000} f(x,y)\,dy\,dx = \int_0^{1000}\int_0^{1000} 10^{-6}e^{-(x+y)/1000}\,dy\,dx \\ &= 10^{-6}\int_0^{1000} e^{-x/1000}\,dx \int_0^{1000} e^{-y/1000}\,dy \\ &= 10^{-6}\left[-1000e^{-x/1000}\right]_0^{1000}\left[-1000e^{-y/1000}\right]_0^{1000} \\ &= \left(e^{-1}-1\right)^2 \approx 0.3996 \end{aligned}$$

(b) Now we are asked for the probability that the combined lifetimes of both bulbs is 1000 hours or less. Thus we want to find $P(X+Y \le 1000)$, or equivalently $P((X,Y) \in D)$ where D is the triangular region shown in the figure. Then



$$\begin{aligned} P(X+Y \le 1000) &= \iint_D f(x,y)\,dA \\ &= \int_0^{1000}\int_0^{1000-x} 10^{-6}e^{-(x+y)/1000}\,dy\,dx \\ &= 10^{-6}\int_0^{1000}\left[-1000e^{-(x+y)/1000}\right]_{y=0}^{y=1000-x} dx = -10^{-3}\int_0^{1000}\left(e^{-1}-e^{-x/1000}\right)dx \\ &= -10^{-3}\left[e^{-1}x + 1000e^{-x/1000}\right]_0^{1000} = 1-2e^{-1} \approx 0.2642 \end{aligned}$$

31. (a) The random variables X and Y are normally distributed with $\mu_1 = 45$, $\mu_2 = 20$, $\sigma_1 = 0.5$, and $\sigma_2 = 0.1$.

The individual density functions for X and Y, then, are $f_1(x) = \dfrac{1}{0.5\sqrt{2\pi}}\,e^{-(x-45)^2/0.5}$ and

$f_2(y) = \dfrac{1}{0.1\sqrt{2\pi}}\,e^{-(y-20)^2/0.02}$. Since X and Y are independent, the joint density function is the product

$$f(x,y) = f_1(x)f_2(y) = \frac{1}{0.5\sqrt{2\pi}}\,e^{-(x-45)^2/0.5}\,\frac{1}{0.1\sqrt{2\pi}}\,e^{-(y-20)^2/0.02} = \frac{10}{\pi}e^{-2(x-45)^2-50(y-20)^2}.$$

Then $P(40 \le X \le 50, 20 \le Y \le 25) = \int_{40}^{50}\int_{20}^{25} f(x,y)\,dy\,dx = \frac{10}{\pi}\int_{40}^{50}\int_{20}^{25} e^{-2(x-45)^2-50(y-20)^2}\,dy\,dx.$

Using a CAS or calculator to evaluate the integral, we get $P(40 \le X \le 50, 20 \le Y \le 25) \approx 0.500$.

(b) $P(4(X-45)^2 + 100(Y-20)^2 \le 2) = \iint_D \frac{10}{\pi}e^{-2(x-45)^2-50(y-20)^2}\,dA$, where D is the region enclosed by the ellipse $4(x-45)^2 + 100(y-20)^2 = 2$. Solving for y gives $y = 20 \pm \frac{1}{10}\sqrt{2-4(x-45)^2}$, the upper and lower halves of the ellipse, and these two halves meet where $y = 20$ [since the ellipse is centered at $(45,20)$] $\Rightarrow$ $4(x-45)^2 = 2$ $\Rightarrow$ $x = 45 \pm \frac{1}{\sqrt{2}}$. Thus

$$\iint_D \frac{10}{\pi}e^{-2(x-45)^2-50(y-20)^2}\,dA = \frac{10}{\pi}\int_{45-1/\sqrt{2}}^{45+1/\sqrt{2}}\ \int_{20-\frac{1}{10}\sqrt{2-4(x-45)^2}}^{20+\frac{1}{10}\sqrt{2-4(x-45)^2}} e^{-2(x-45)^2-50(y-20)^2}\,dy\,dx.$$

Using a CAS or calculator to evaluate the integral, we get $P(4(X-45)^2 + 100(Y-20)^2 \le 2) \approx 0.632$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

32. Because X and Y are independent, the joint density function for Xavier's and Yolanda's arrival times is the product of the individual density functions:

$$f(x,y) = f_1(x)f_2(y) = \begin{cases} \frac{1}{50}e^{-x}y & \text{if } x \geq 0,\ 0 \leq y \leq 10 \\ 0 & \text{otherwise} \end{cases}$$

Since Xavier won't wait for Yolanda, they won't meet unless $X \geq Y$. Additionally, Yolanda will wait up to half an hour but no longer, so they won't meet unless $X - Y \leq 30$. Thus the probability that they meet is $P((X,Y) \in D)$ where D is the parallelogram shown in the figure. The integral is simpler to evaluate if we consider D as a type II region, so



$$\begin{aligned} P((X,Y) \in D) &= \iint_D f(x,y)\,dx\,dy = \int_0^{10}\int_y^{y+30} \tfrac{1}{50}e^{-x}y\,dx\,dy \\ &= \tfrac{1}{50}\int_0^{10} y\left[-e^{-x}\right]_{x=y}^{x=y+30} dy = \tfrac{1}{50}\int_0^{10} y(-e^{-(y+30)} + e^{-y})\,dy \\ &= \tfrac{1}{50}(1-e^{-30})\int_0^{10} ye^{-y}\,dy \end{aligned}$$

By integration by parts (or Formula 96 in the Table of Integrals), this is $\frac{1}{50}(1-e^{-30})\left[-(y+1)e^{-y}\right]_0^{10} = \frac{1}{50}(1-e^{-30})(1-11e^{-10}) \approx 0.020$. Thus there is only about a 2% chance they will meet. Such is student life!

33. (a) If $f(P,A)$ is the probability that an individual at A will be infected by an individual at P, and $k\,dA$ is the number of infected individuals in an element of area dA, then $f(P,A)k\,dA$ is the number of infections that should result from exposure of the individual at A to infected people in the element of area dA. Integration over D gives the number of infections of the person at A due to all the infected people in D. In rectangular coordinates (with the origin at the city's center), the exposure of a person at A is

$$E = \iint_D kf(P,A)\,dA = k\iint_D \tfrac{1}{20}\left[20 - d(P,A)\right]\,dA = k\iint_D \left[1 - \tfrac{1}{20}\sqrt{(x-x_0)^2+(y-y_0)^2}\right]\,dA$$

(b) If $A = (0,0)$, then

$$\begin{aligned} E &= k\iint_D \left[1 - \tfrac{1}{20}\sqrt{x^2+y^2}\right]\,dA \\ &= k\int_0^{2\pi}\int_0^{10} \left(1 - \tfrac{1}{20}r\right)r\,dr\,d\theta = 2\pi k\left[\tfrac{1}{2}r^2 - \tfrac{1}{60}r^3\right]_0^{10} \\ &= 2\pi k\left(50 - \tfrac{50}{3}\right) = \tfrac{200}{3}\pi k \approx 209k \end{aligned}$$



For A at the edge of the city, it is convenient to use a polar coordinate system centered at A. Then the polar equation for the circular boundary of the city becomes $r = 20\cos\theta$ instead of $r = 10$, and the distance from A to a point P in the city

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

is again r (see the figure). So

$$E = k\int_{-\pi/2}^{\pi/2}\int_0^{20\cos\theta}\left(1-\tfrac{1}{20}r\right)r\,dr\,d\theta = k\int_{-\pi/2}^{\pi/2}\left[\tfrac{1}{2}r^2-\tfrac{1}{60}r^3\right]_{r=0}^{r=20\cos\theta}d\theta$$
$$= k\int_{-\pi/2}^{\pi/2}\left(200\cos^2\theta-\tfrac{400}{3}\cos^3\theta\right)d\theta = 200k\int_{-\pi/2}^{\pi/2}\left[\tfrac{1}{2}+\tfrac{1}{2}\cos 2\theta-\tfrac{2}{3}\left(1-\sin^2\theta\right)\cos\theta\right]d\theta$$
$$= 200k\left[\tfrac{1}{2}\theta+\tfrac{1}{4}\sin 2\theta-\tfrac{2}{3}\sin\theta+\tfrac{2}{3}\cdot\tfrac{1}{3}\sin^3\theta\right]_{-\pi/2}^{\pi/2} = 200k\left[\tfrac{\pi}{4}+0-\tfrac{2}{3}+\tfrac{2}{9}+\tfrac{\pi}{4}+0-\tfrac{2}{3}+\tfrac{2}{9}\right]$$
$$= 200k\left(\tfrac{\pi}{2}-\tfrac{8}{9}\right)\approx 136k$$

Therefore the risk of infection is much lower at the edge of the city than in the middle, so it is better to live at the edge.

15.5 Surface Area

1. Here $z = f(x,y) = 5x+3y+6$ and D is the rectangle $[1,4]\times[2,6]$, so by Formula 2 the area of the surface is

$$A(S) = \iint_D \sqrt{[f_x(x,y)]^2+[f_y(x,y)]^2+1}\,dA = \iint_D \sqrt{5^2+3^2+1}\,dA = \sqrt{35}\iint_D dA$$
$$= \sqrt{35}\,A(D) = \sqrt{35}\,(3)(4) = 12\sqrt{35}$$

2. $z = f(x,y) = \frac{1}{2}-3x-2y$ and D is the disk $x^2+y^2\le 25$, so by Formula 2

$$A(S) = \iint_D \sqrt{(-3)^2+(-2)^2+1}\,dA = \sqrt{14}\iint_D dA = \sqrt{14}\,A(D) = \sqrt{14}\,(\pi\cdot 5^2) = 25\sqrt{14}\,\pi$$

3. The surface S is given by $z = f(x,y) = 6-3x-2y$ which intersects the xy-plane in the line $3x+2y=6$, so D is the triangular region given by $\left\{(x,y)\mid 0\le x\le 2,\ 0\le y\le 3-\frac{3}{2}x\right\}$. By Formula 2, the surface area of S is

$$A(S) = \iint_D \sqrt{(-3)^2+(-2)^2+1}\,dA = \sqrt{14}\iint_D dA = \sqrt{14}\,A(D) = \sqrt{14}\left(\tfrac{1}{2}\cdot 2\cdot 3\right) = 3\sqrt{14}$$

4. $z = f(x,y) = \frac{1}{4}x^2-\frac{1}{2}y+\frac{5}{4}$, and D is the triangular region given by $\{(x,y)\mid 0\le x\le 2,\ 0\le y\le 2x\}$. By Formula 2,

$$A(S) = \iint_D \sqrt{\left(\tfrac{1}{2}x\right)^2+\left(-\tfrac{1}{2}\right)^2+1}\,dA = \int_0^2\int_0^{2x}\sqrt{\tfrac{1}{4}x^2+\tfrac{5}{4}}\,dy\,dx = \int_0^2 \tfrac{1}{2}\sqrt{x^2+5}\,\left[y\right]_{y=0}^{y=2x}dx$$
$$= \tfrac{1}{2}\int_0^2 2x\sqrt{x^2+5}\,dx = \tfrac{1}{2}\cdot\tfrac{2}{3}(x^2+5)^{3/2}\Big]_0^2 = \tfrac{1}{3}(9^{3/2}-5^{3/2}) = 9-\tfrac{5}{3}\sqrt{5}$$

5. The paraboloid intersects the plane $z=-2$ when $1-x^2-y^2=-2 \iff x^2+y^2=3$, so $D=\left\{(x,y)\mid x^2+y^2\le 3\right\}$.

Here $z = f(x,y) = 1-x^2-y^2 \quad\Rightarrow\quad f_x=-2x,\ f_y=-2y$ and

$$A(S) = \iint_D \sqrt{(-2x)^2+(-2y)^2+1}\,dA = \iint_D \sqrt{4(x^2+y^2)+1}\,dA = \int_0^{2\pi}\int_0^{\sqrt{3}}\sqrt{4r^2+1}\,r\,dr\,d\theta$$
$$= \int_0^{2\pi}d\theta\int_0^{\sqrt{3}} r\sqrt{4r^2+1}\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{1}{12}(4r^2+1)^{3/2}\right]_0^{\sqrt{3}} = 2\pi\cdot\tfrac{1}{12}\left(13^{3/2}-1\right) = \tfrac{\pi}{6}\left(13\sqrt{13}-1\right)$$

6. $x^2+z^2=4 \quad\Rightarrow\quad z=\sqrt{4-x^2}$ (since $z\ge 0$), so $f_x = -x(4-x^2)^{-1/2}$, $f_y=0$ and

$$A(S) = \int_0^1\int_0^1\sqrt{\left[-x(4-x^2)^{-1/2}\right]^2+0^2+1}\,dy\,dx = \int_0^1\int_0^1\sqrt{\frac{x^2}{4-x^2}+1}\,dy\,dx$$
$$= \int_0^1\frac{2}{\sqrt{4-x^2}}\,dx\int_0^1 dy = \left[2\sin^{-1}\frac{x}{2}\right]_0^1\left[y\right]_0^1 = \left(2\cdot\tfrac{\pi}{6}-0\right)(1) = \tfrac{\pi}{3}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

7. $z = f(x,y) = y^2 - x^2$ with $1 \le x^2 + y^2 \le 4$. Then

$$A(S) = \iint_D \sqrt{4x^2 + 4y^2 + 1}\, dA = \int_0^{2\pi}\int_1^2 \sqrt{4r^2+1}\, r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_1^2 r\sqrt{4r^2+1}\, dr$$

$$= \left[\theta\right]_0^{2\pi}\left[\tfrac{1}{12}(4r^2+1)^{3/2}\right]_1^2 = \tfrac{\pi}{6}\left(17\sqrt{17} - 5\sqrt{5}\right)$$

8. $z = f(x,y) = \frac{2}{3}(x^{3/2} + y^{3/2})$ and $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le 1\}$. Then $f_x = x^{1/2}$, $f_y = y^{1/2}$ and

$$A(S) = \iint_D \sqrt{\left(\sqrt{x}\right)^2 + \left(\sqrt{y}\right)^2 + 1}\, dA = \int_0^1\int_0^1 \sqrt{x+y+1}\, dy\, dx = \int_0^1 \left[\tfrac{2}{3}(x+y+1)^{3/2}\right]_{y=0}^{y=1} dx$$

$$= \tfrac{2}{3}\int_0^1 \left[(x+2)^{3/2} - (x+1)^{3/2}\right] dx = \tfrac{2}{3}\left[\tfrac{2}{5}(x+2)^{5/2} - \tfrac{2}{5}(x+1)^{5/2}\right]_0^1$$

$$= \tfrac{4}{15}(3^{5/2} - 2^{5/2} - 2^{5/2} + 1) = \tfrac{4}{15}(3^{5/2} - 2^{7/2} + 1)$$

9. $z = f(x,y) = xy$ with $x^2 + y^2 \le 1$, so $f_x = y$, $f_y = x$ $\Rightarrow$

$$A(S) = \iint_D \sqrt{y^2 + x^2 + 1}\, dA = \int_0^{2\pi}\int_0^1 \sqrt{r^2+1}\, r\, dr\, d\theta = \int_0^{2\pi}\left[\tfrac{1}{3}(r^2+1)^{3/2}\right]_{r=0}^{r=1} d\theta$$

$$= \int_0^{2\pi} \tfrac{1}{3}\left(2\sqrt{2} - 1\right) d\theta = \tfrac{2\pi}{3}\left(2\sqrt{2} - 1\right)$$

10. Given the sphere $x^2 + y^2 + z^2 = 4$, when $z = 1$, we get $x^2 + y^2 = 3$ so $D = \left\{(x,y) \mid x^2 + y^2 \le 3\right\}$ and $z = f(x,y) = \sqrt{4 - x^2 - y^2}$. Thus

$$A(S) = \iint_D \sqrt{[(-x)(4-x^2-y^2)^{-1/2}]^2 + [(-y)(4-x^2-y^2)^{-1/2}]^2 + 1}\, dA$$

$$= \int_0^{2\pi}\int_0^{\sqrt{3}} \sqrt{\frac{r^2}{4-r^2} + 1}\, r\, dr\, d\theta = \int_0^{2\pi}\int_0^{\sqrt{3}} \sqrt{\frac{r^2 + 4 - r^2}{4-r^2}}\, r\, dr\, d\theta$$

$$= \int_0^{2\pi}\int_0^{\sqrt{3}} \frac{2r}{\sqrt{4-r^2}}\, dr\, d\theta$$

$$= \int_0^{2\pi}\left[-2(4-r^2)^{1/2}\right]_{r=0}^{r=\sqrt{3}} d\theta = \int_0^{2\pi}(-2+4)\, d\theta = 2\theta\Big]_0^{2\pi} = 4\pi$$

11. $z = \sqrt{a^2 - x^2 - y^2}$, $z_x = -x(a^2 - x^2 - y^2)^{-1/2}$, $z_y = -y(a^2 - x^2 - y^2)^{-1/2}$,

$$A(S) = \iint_D \sqrt{\frac{x^2+y^2}{a^2-x^2-y^2} + 1}\, dA$$

$$= \int_{-\pi/2}^{\pi/2}\int_0^{a\cos\theta} \sqrt{\frac{r^2}{a^2-r^2} + 1}\, r\, dr\, d\theta$$

$$= \int_{-\pi/2}^{\pi/2}\int_0^{a\cos\theta} \frac{ar}{\sqrt{a^2-r^2}}\, dr\, d\theta$$

$$= \int_{-\pi/2}^{\pi/2}\left[-a\sqrt{a^2-r^2}\right]_{r=0}^{r=a\cos\theta} d\theta$$

$$= \int_{-\pi/2}^{\pi/2} -a\left(\sqrt{a^2 - a^2\cos^2\theta} - a\right) d\theta = 2a^2\int_0^{\pi/2}\left(1 - \sqrt{1-\cos^2\theta}\right) d\theta$$

$$= 2a^2\int_0^{\pi/2} d\theta - 2a^2\int_0^{\pi/2}\sqrt{\sin^2\theta}\, d\theta = a^2\pi - 2a^2\int_0^{\pi/2}\sin\theta\, d\theta = a^2(\pi - 2)$$



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. To find the region D: $z = x^2 + y^2$ implies $z + z^2 = 4z$ or $z^2 - 3z = 0$. Thus $z = 0$ or $z = 3$ are the planes where the surfaces intersect. But $x^2 + y^2 + z^2 = 4z$ implies $x^2 + y^2 + (z-2)^2 = 4$, so $z = 3$ intersects the upper hemisphere. Thus $(z-2)^2 = 4 - x^2 - y^2$ or $z = 2 + \sqrt{4 - x^2 - y^2}$. Therefore D is the region inside the circle $x^2 + y^2 + (3-2)^2 = 4$, that is, $D = \{(x, y) \mid x^2 + y^2 \le 3\}$.

$$\begin{aligned} A(S) &= \iint_D \sqrt{[(-x)(4 - x^2 - y^2)^{-1/2}]^2 + [(-y)(4 - x^2 - y^2)^{-1/2}]^2 + 1}\, dA \\ &= \int_0^{2\pi} \int_0^{\sqrt{3}} \sqrt{\frac{r^2}{4 - r^2} + 1}\, r\, dr\, d\theta = \int_0^{2\pi} \int_0^{\sqrt{3}} \frac{2r\, dr}{\sqrt{4 - r^2}}\, d\theta = \int_0^{2\pi} \left[-2(4 - r^2)^{1/2}\right]_{r=0}^{r=\sqrt{3}} d\theta \\ &= \int_0^{2\pi} (-2 + 4)\, d\theta = 2\theta \Big]_0^{2\pi} = 4\pi \end{aligned}$$

13. $z = f(x, y) = (1 + x^2 + y^2)^{-1}$, $f_x = -2x(1 + x^2 + y^2)^{-2}$, $f_y = -2y(1 + x^2 + y^2)^{-2}$. Then

$$\begin{aligned} A(S) &= \iint_{x^2 + y^2 \le 1} \sqrt{[-2x(1 + x^2 + y^2)^{-2}]^2 + [-2y(1 + x^2 + y^2)^{-2}]^2 + 1}\, dA \\ &= \iint_{x^2 + y^2 \le 1} \sqrt{4(x^2 + y^2)(1 + x^2 + y^2)^{-4} + 1}\, dA \end{aligned}$$

Converting to polar coordinates we have

$$\begin{aligned} A(S) &= \int_0^{2\pi} \int_0^1 \sqrt{4r^2(1 + r^2)^{-4} + 1}\, r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_0^1 r \sqrt{4r^2(1 + r^2)^{-4} + 1}\, dr \\ &= 2\pi \int_0^1 r \sqrt{4r^2(1 + r^2)^{-4} + 1}\, dr \approx 3.6258 \text{ using a calculator.} \end{aligned}$$

14. $z = f(x, y) = \cos(x^2 + y^2)$, $f_x = -2x \sin(x^2 + y^2)$, $f_y = -2y \sin(x^2 + y^2)$.

$$A(S) = \iint_{x^2 + y^2 \le 1} \sqrt{4x^2 \sin^2(x^2 + y^2) + 4y^2 \sin^2(x^2 + y^2) + 1}\, dA = \iint_{x^2 + y^2 \le 1} \sqrt{4(x^2 + y^2) \sin^2(x^2 + y^2) + 1}\, dA.$$

Converting to polar coordinates gives

$$\begin{aligned} A(S) &= \int_0^{2\pi} \int_0^1 \sqrt{4r^2 \sin^2(r^2) + 1}\, r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_0^1 r \sqrt{4r^2 \sin^2(r^2) + 1}\, dr \\ &= 2\pi \int_0^1 r \sqrt{4r^2 \sin^2(r^2) + 1}\, dr \approx 4.1073 \quad \text{using a calculator.} \end{aligned}$$

15. (a) The midpoints of the four squares are $\left(\frac{1}{4}, \frac{1}{4}\right)$, $\left(\frac{1}{4}, \frac{3}{4}\right)$, $\left(\frac{3}{4}, \frac{1}{4}\right)$, and $\left(\frac{3}{4}, \frac{3}{4}\right)$. Here $f(x, y) = x^2 + y^2$, so the Midpoint Rule gives

$$\begin{aligned} A(S) &= \iint_D \sqrt{[f_x(x, y)]^2 + [f_y(x, y)]^2 + 1}\, dA = \iint_D \sqrt{(2x)^2 + (2y)^2 + 1}\, dA \\ &\approx \tfrac{1}{4}\Bigg(\sqrt{\left[2\left(\tfrac{1}{4}\right)\right]^2 + \left[2\left(\tfrac{1}{4}\right)\right]^2 + 1} + \sqrt{\left[2\left(\tfrac{1}{4}\right)\right]^2 + \left[2\left(\tfrac{3}{4}\right)\right]^2 + 1} \\ &\qquad + \sqrt{\left[2\left(\tfrac{3}{4}\right)\right]^2 + \left[2\left(\tfrac{1}{4}\right)\right]^2 + 1} + \sqrt{\left[2\left(\tfrac{3}{4}\right)\right]^2 + \left[2\left(\tfrac{3}{4}\right)\right]^2 + 1}\Bigg) \\ &= \tfrac{1}{4}\left(\sqrt{\tfrac{3}{2}} + 2\sqrt{\tfrac{7}{2}} + \sqrt{\tfrac{11}{2}}\right) \approx 1.8279 \end{aligned}$$

(b) A CAS estimates the integral to be $A(S) = \iint_D \sqrt{1 + (2x)^2 + (2y)^2}\, dA = \int_0^1 \int_0^1 \sqrt{1 + 4x^2 + 4y^2}\, dy\, dx \approx 1.8616$.

This agrees with the Midpoint estimate only in the first decimal place.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

16. (a) With $m = n = 2$ we have four squares with midpoints $\left(\frac{1}{2}, \frac{1}{2}\right)$, $\left(\frac{1}{2}, \frac{3}{2}\right)$, $\left(\frac{3}{2}, \frac{1}{2}\right)$, and $\left(\frac{3}{2}, \frac{3}{2}\right)$. Since $z = xy + x^2 + y^2$, the Midpoint Rule gives

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\, dA = \iint_D \sqrt{1 + (y + 2x)^2 + (x + 2y)^2}\, dA \\ &\approx 1\left(\sqrt{1 + \left(\tfrac{3}{2}\right)^2 + \left(\tfrac{3}{2}\right)^2} + \sqrt{1 + \left(\tfrac{5}{2}\right)^2 + \left(\tfrac{7}{2}\right)^2} + \sqrt{1 + \left(\tfrac{7}{2}\right)^2 + \left(\tfrac{5}{2}\right)^2} + \sqrt{1 + \left(\tfrac{9}{2}\right)^2 + \left(\tfrac{9}{2}\right)^2}\right) \\ &= \tfrac{\sqrt{22}}{2} + \tfrac{\sqrt{78}}{2} + \tfrac{\sqrt{78}}{2} + \tfrac{\sqrt{166}}{2} \approx 17.619 \end{aligned}$$

(b) Using a CAS, we have

$A(S) = \iint_D \sqrt{1 + (y + 2x)^2 + (x + 2y)^2}\, dA = \int_0^2 \int_0^2 \sqrt{1 + (y + 2x)^2 + (x + 2y)^2}\, dy\, dx \approx 17.7165$. This is within about 0.1 of the Midpoint Rule estimate.

17. $z = 1 + 2x + 3y + 4y^2$, so

$$A(S) = \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\, dA = \int_1^4 \int_0^1 \sqrt{1 + 4 + (3 + 8y)^2}\, dy\, dx = \int_1^4 \int_0^1 \sqrt{14 + 48y + 64y^2}\, dy\, dx.$$

Using a CAS, we have $\int_1^4 \int_0^1 \sqrt{14 + 48y + 64y^2}\, dy\, dx = \frac{45}{8}\sqrt{14} + \frac{15}{16}\ln(11\sqrt{5} + 3\sqrt{14}\sqrt{5}) - \frac{15}{16}\ln(3\sqrt{5} + \sqrt{14}\sqrt{5})$ or $\frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\dfrac{11\sqrt{5} + 3\sqrt{70}}{3\sqrt{5} + \sqrt{70}}$.

18. $f(x, y) = 1 + x + y + x^2 \;\Rightarrow\; f_x = 1 + 2x$, $f_y = 1$. We use a CAS to calculate the integral

$$\begin{aligned} A(S) &= \int_{-2}^{1} \int_{-1}^{1} \sqrt{f_x^2 + f_y^2 + 1}\, dy\, dx \\ &= \int_{-2}^{1} \int_{-1}^{1} \sqrt{(1 + 2x)^2 + 2}\, dy\, dx = 2\int_{-2}^{1} \sqrt{4x^2 + 4x + 3}\, dx \end{aligned}$$

and find that $A(S) = 3\sqrt{11} + 2\sinh^{-1}\left(\frac{3\sqrt{2}}{2}\right)$ or

$A(S) = 3\sqrt{11} + \ln(10 + 3\sqrt{11})$.



19. $f(x, y) = 1 + x^2y^2 \;\Rightarrow\; f_x = 2xy^2$, $f_y = 2x^2y$. We use a CAS (with precision reduced to five significant digits, to speed up the calculation) to estimate the integral

$A(S) = \displaystyle\int_{-1}^{1} \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \sqrt{f_x^2 + f_y^2 + 1}\, dy\, dx = \int_{-1}^{1} \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \sqrt{4x^2y^4 + 4x^4y^2 + 1}\, dy\, dx$, and find that $A(S) \approx 3.3213$.

20. Let $f(x, y) = \dfrac{1 + x^2}{1 + y^2}$. Then $f_x = \dfrac{2x}{1 + y^2}$.

$f_y = (1 + x^2)\left[-\dfrac{2y}{(1 + y^2)^2}\right] = -\dfrac{2y(1 + x^2)}{(1 + y^2)^2}$. We use a CAS to estimate $\int_{-1}^{1} \int_{-(1-|x|)}^{1-|x|} \sqrt{f_x^2 + f_y^2 + 1}\, dy\, dx \approx 2.6959$. In order to graph only the part of the surface above the square, we use $-(1 - |x|) \le y \le 1 - |x|$ as the y-range in our plot command.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. Here $z = f(x,y) = ax + by + c$, $f_x(x,y) = a$, $f_y(x,y) = b$, so

$A(S) = \iint_D \sqrt{a^2+b^2+1}\,dA = \sqrt{a^2+b^2+1}\iint_D dA = \sqrt{a^2+b^2+1}\,A(D)$.

22. Let S be the upper hemisphere. Then $z = f(x,y) = \sqrt{a^2-x^2-y^2}$, so

$$\begin{aligned} A(S) &= \iint_D \sqrt{[-x(a^2-x^2-y^2)^{-1/2}]^2 + [-y(a^2-x^2-y^2)^{-1/2}]^2 + 1}\,dA \\ &= \iint_D \sqrt{\frac{x^2+y^2}{a^2-x^2-y^2}+1}\,dA = \lim_{t\to a^-}\int_0^{2\pi}\int_0^t \sqrt{\frac{r^2}{a^2-r^2}+1}\,r\,dr\,d\theta \\ &= \lim_{t\to a^-}\int_0^{2\pi}\int_0^t \frac{ar}{\sqrt{a^2-r^2}}\,dr\,d\theta = 2\pi\lim_{t\to a}\left[-a\sqrt{a^2-r^2}\right]_0^t = 2\pi\lim_{t\to a^-} -a\left[\sqrt{a^2-t^2}-a\right] \\ &= 2\pi(-a)(-a) = 2\pi a^2. \text{ Thus the surface area of the entire sphere is } 4\pi a^2. \end{aligned}$$

23. If we project the surface onto the xz-plane, then the surface lies "above" the disk $x^2+z^2 \le 25$ in the xz-plane.

We have $y = f(x,z) = x^2+z^2$ and, adapting Formula 2, the area of the surface is

$$A(S) = \iint_{x^2+z^2\le 25} \sqrt{[f_x(x,z)]^2 + [f_z(x,z)]^2 + 1}\,dA = \iint_{x^2+z^2\le 25} \sqrt{4x^2+4z^2+1}\,dA$$

Converting to polar coordinates $x = r\cos\theta$, $z = r\sin\theta$ we have

$$A(S) = \int_0^{2\pi}\int_0^5 \sqrt{4r^2+1}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^5 r(4r^2+1)^{1/2}\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{1}{12}(4r^2+1)^{3/2}\right]_0^5 = \tfrac{\pi}{6}\left(101\sqrt{101}-1\right)$$

24. First we find the area of the face of the surface that intersects the positive y-axis. As in Exercise 23, we can project the face onto the xz-plane, so the surface lies "above" the disk $x^2+z^2 \le 1$. Then $y = f(x,z) = \sqrt{1-z^2}$ and the area is

$$\begin{aligned} A(S) &= \iint_{x^2+z^2\le 1} \sqrt{[f_x(x,z)]^2 + [f_z(x,z)]^2 + 1}\,dA = \iint_{x^2+z^2\le 1} \sqrt{0 + \left(\frac{-z}{\sqrt{1-z^2}}\right)^2 + 1}\,dA \\ &= \iint_{x^2+z^2\le 1} \sqrt{\frac{z^2}{1-z^2}+1}\,dA = \int_{-1}^1 \int_{-\sqrt{1-z^2}}^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz \\ &= 4\int_0^1 \int_0^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz \qquad \text{[by the symmetry of the surface]} \end{aligned}$$

This integral is improper (when $z = 1$), so

$$A(S) = \lim_{t\to 1^-} 4\int_0^t \int_0^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz = \lim_{t\to 1^-} 4\int_0^t \frac{\sqrt{1-z^2}}{\sqrt{1-z^2}}\,dz = \lim_{t\to 1^-} 4\int_0^t dz = \lim_{t\to 1^-} 4t = 4.$$

Since the complete surface consists of four congruent faces, the total surface area is $4(4) = 16$.

15.6 Triple Integrals

1. $\iiint_B xyz^2\,dV = \int_0^1\int_0^3\int_{-1}^2 xyz^2\,dy\,dz\,dx = \int_0^1\int_0^3 \left[\tfrac{1}{2}xy^2z^2\right]_{y=-1}^{y=2} dz\,dx = \int_0^1\int_0^3 \tfrac{3}{2}xz^2\,dz\,dx$

$= \int_0^1 \left[\tfrac{1}{2}xz^3\right]_{z=0}^{z=3} dx = \int_0^1 \tfrac{27}{2}\,x\,dx = \tfrac{27}{4}x^2\Big]_0^1 = \tfrac{27}{4}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

2. There are six different possible orders of integration.

$$\iiint_E (xy+z^2)\,dV = \int_0^2\int_0^1\int_0^3 (xy+z^2)\,dz\,dy\,dx = \int_0^2\int_0^1 \left[xyz+\tfrac{1}{3}z^3\right]_{z=0}^{z=3} dy\,dx = \int_0^2\int_0^1 (3xy+9)\,dy\,dx$$
$$= \int_0^2 \left[\tfrac{3}{2}xy^2+9y\right]_{y=0}^{y=1} dx = \int_0^2 \left(\tfrac{3}{2}x+9\right) dx = \left[\tfrac{3}{4}x^2+9x\right]_0^2 = 21$$

$$\iiint_E (xy+z^2)\,dV = \int_0^1\int_0^2\int_0^3 (xy+z^2)\,dz\,dx\,dy = \int_0^1\int_0^2 \left[xyz+\tfrac{1}{3}z^3\right]_{z=0}^{z=3} dx\,dy = \int_0^1\int_0^2 (3xy+9)\,dx\,dy$$
$$= \int_0^1 \left[\tfrac{3}{2}x^2y+9x\right]_{x=0}^{x=2} dy = \int_0^1 (6y+18)\,dy = \left[3y^2+18y\right]_0^1 = 21$$

$$\iiint_E (xy+z^2)\,dV = \int_0^2\int_0^3\int_0^1 (xy+z^2)\,dy\,dz\,dx = \int_0^2\int_0^3 \left[\tfrac{1}{2}xy^2+yz^2\right]_{y=0}^{y=1} dz\,dx = \int_0^2\int_0^3 \left(\tfrac{1}{2}x+z^2\right) dz\,dx$$
$$= \int_0^2 \left[\tfrac{1}{2}xz+\tfrac{1}{3}z^3\right]_{z=0}^{z=3} dx = \int_0^2 \left(\tfrac{3}{2}x+9\right) dx = \left[\tfrac{3}{4}x^2+9x\right]_0^2 = 21$$

$$\iiint_E (xy+z^2)\,dV = \int_0^3\int_0^2\int_0^1 (xy+z^2)\,dy\,dx\,dz = \int_0^3\int_0^2 \left[\tfrac{1}{2}xy^2+yz^2\right]_{y=0}^{y=1} dx\,dz = \int_0^3\int_0^2 \left(\tfrac{1}{2}x+z^2\right) dx\,dz$$
$$= \int_0^3 \left[\tfrac{1}{4}x^2+xz^2\right]_{x=0}^{x=2} dz = \int_0^3 \left(1+2z^2\right) dz = \left[z+\tfrac{2}{3}z^3\right]_0^3 = 21$$

$$\iiint_E (xy+z^2)\,dV = \int_0^1\int_0^3\int_0^2 (xy+z^2)\,dx\,dz\,dy = \int_0^1\int_0^3 \left[\tfrac{1}{2}x^2y+xz^2\right]_{x=0}^{x=2} dz\,dy = \int_0^1\int_0^3 \left(2y+2z^2\right) dz\,dy$$
$$= \int_0^1 \left[2yz+\tfrac{2}{3}z^3\right]_{z=0}^{z=3} dy = \int_0^1 (6y+18)\,dy = \left[3y^2+18y\right]_0^1 = 21$$

$$\iiint_E (xy+z^2)\,dV = \int_0^3\int_0^1\int_0^2 (xy+z^2)\,dx\,dy\,dz = \int_0^3\int_0^1 \left[\tfrac{1}{2}x^2y+xz^2\right]_{x=0}^{x=2} dy\,dz = \int_0^3\int_0^1 \left(2y+2z^2\right) dy\,dz$$
$$= \int_0^3 \left[y^2+2yz^2\right]_{y=0}^{y=1} dz = \int_0^3 \left(1+2z^2\right) dz = \left[z+\tfrac{2}{3}z^3\right]_0^3 = 21$$

3. $\int_0^2\int_0^{z^2}\int_0^{y-z}(2x-y)\,dx\,dy\,dz = \int_0^2\int_0^{z^2}\left[x^2-xy\right]_{x=0}^{x=y-z} dy\,dz = \int_0^2\int_0^{z^2}\left[(y-z)^2-(y-z)y\right] dy\,dz$

$$= \int_0^2\int_0^{z^2}\left(z^2-yz\right) dy\,dz = \int_0^2\left[yz^2-\tfrac{1}{2}y^2z\right]_{y=0}^{y=z^2} dz = \int_0^2\left(z^4-\tfrac{1}{2}z^5\right) dz$$
$$= \left[\tfrac{1}{5}z^5 \quad \tfrac{1}{12}z^6\right]_0^2 = \tfrac{32}{5}-\tfrac{64}{12} = \tfrac{16}{15}$$

4. $\int_0^1\int_y^{2y}\int_0^{x+y} 6xy\,dz\,dx\,dy = \int_0^1\int_y^{2y}\left[6xyz\right]_{z=0}^{z=x+y} dx\,dy = \int_0^1\int_y^{2y} 6xy(x+y)\,dx\,dy = \int_0^1\int_y^{2y}\left(6x^2y+6xy^2\right) dx\,dy$

$$= \int_0^1 \left[2x^3y+3x^2y^2\right]_{x=y}^{x=2y} dy = \int_0^1 23y^4\,dy = \tfrac{23}{5}y^5\Big]_0^1 = \tfrac{23}{5}$$

5. $\int_1^2\int_0^{2z}\int_0^{\ln x} xe^{-y}\,dy\,dx\,dz = \int_1^2\int_0^{2z}\left[-xe^{-y}\right]_{y=0}^{y=\ln x} dx\,dz = \int_1^2\int_0^{2z}\left(-xe^{-\ln x}+xe^0\right) dx\,dz$

$$= \int_1^2\int_0^{2z}(-1+x)\,dx\,dz = \int_1^2\left[-x+\tfrac{1}{2}x^2\right]_{x=0}^{x=2z} dz$$
$$= \int_1^2\left(-2z+2z^2\right) dz = \left[-z^2+\tfrac{2}{3}z^3\right]_1^2 = -4+\tfrac{16}{3}+1-\tfrac{2}{3} = \tfrac{5}{3}$$

6. $\displaystyle\int_0^1\int_0^1\int_0^{\sqrt{1-z^2}} \frac{z}{y+1}\,dx\,dz\,dy = \int_0^1\int_0^1\left[\frac{z}{y+1}\cdot x\right]_{x=0}^{x=\sqrt{1-z^2}} dz\,dy = \int_0^1\int_0^1 \frac{z\sqrt{1-z^2}}{y+1}\,dz\,dy$

$$= \int_0^1 \left[\frac{-\frac{1}{3}(1-z^2)^{3/2}}{y+1}\right]_{z=0}^{z=1} dy = \frac{1}{3}\int_0^1 \frac{1}{y+1}\,dy = \frac{1}{3}\ln(y+1)\Big]_0^1$$
$$= \tfrac{1}{3}(\ln 2-\ln 1) = \tfrac{1}{3}\ln 2$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $\int_0^{\pi}\int_0^1\int_0^{\sqrt{1-z^2}} z\sin x\,dy\,dz\,dx = \int_0^{\pi}\int_0^1 \left[yz\sin x\right]_{y=0}^{y=\sqrt{1-z^2}} dz\,dx = \int_0^{\pi}\int_0^1 z\sqrt{1-z^2}\,\sin x\,dz\,dx$

$= \int_0^{\pi}\sin x\left[-\tfrac{1}{3}(1-z^2)^{3/2}\right]_{z=0}^{z=1} dx = \int_0^{\pi}\tfrac{1}{3}\sin x\,dx = -\tfrac{1}{3}\cos x\Big]_0^{\pi} = -\tfrac{1}{3}(-1-1) = \tfrac{2}{3}$

8. $\int_0^1\int_0^1\int_0^{2-x^2-y^2} xye^z\,dz\,dy\,dx = \int_0^1\int_0^1\left[xye^z\right]_{z=0}^{z=2-x^2-y^2} dy\,dx = \int_0^1\int_0^1 (xye^{2-x^2-y^2} - xy)\,dy\,dx$

$= \int_0^1\left[-\tfrac{1}{2}xe^{2-x^2-y^2} - \tfrac{1}{2}xy^2\right]_{y=0}^{y=1} dx = \int_0^1\left(-\tfrac{1}{2}xe^{1-x^2} - \tfrac{1}{2}x + \tfrac{1}{2}xe^{2-x^2}\right)dx$

$= \left[\tfrac{1}{4}e^{1-x^2} - \tfrac{1}{4}x^2 - \tfrac{1}{4}e^{2-x^2}\right]_0^1 = \tfrac{1}{4} - \tfrac{1}{4} - \tfrac{1}{4}e - \tfrac{1}{4}e + 0 + \tfrac{1}{4}e^2 = \tfrac{1}{4}e^2 - \tfrac{1}{2}e$

9. $\iiint_E y\,dV = \int_0^3\int_0^x\int_{x-y}^{x+y} y\,dz\,dy\,dx = \int_0^3\int_0^x\left[yz\right]_{z=x-y}^{z=x+y} dy\,dx = \int_0^3\int_0^x 2y^2\,dy\,dx$

$= \int_0^3\left[\tfrac{2}{3}y^3\right]_{y=0}^{y=x} dx = \int_0^3\tfrac{2}{3}x^3\,dx = \tfrac{1}{6}x^4\Big]_0^3 = \tfrac{81}{6} = \tfrac{27}{2}$

10. $\iiint_E e^{z/y}\,dV = \int_0^1\int_y^1\int_0^{xy} e^{z/y}\,dz\,dx\,dy = \int_0^1\int_y^1\left[ye^{z/y}\right]_{z=0}^{z=xy} dx\,dy$

$= \int_0^1\int_y^1 (ye^x - y)\,dx\,dy = \int_0^1\left[ye^x - xy\right]_{x=y}^{x=1} dy = \int_0^1\left(ey - y - ye^y + y^2\right)dy$

$= \left[\tfrac{1}{2}ey^2 - \tfrac{1}{2}y^2 - (y-1)e^y + \tfrac{1}{3}y^3\right]_0^1$ [integrate by parts]

$= \tfrac{1}{2}e - \tfrac{1}{2} + \tfrac{1}{3} - 1 = \tfrac{1}{2}e - \tfrac{7}{6}$

11. $\iiint_E \frac{z}{x^2+z^2}\,dV = \int_1^4\int_y^4\int_0^z \frac{z}{x^2+z^2}\,dx\,dz\,dy = \int_1^4\int_y^4\left[z\cdot\frac{1}{z}\tan^{-1}\frac{x}{z}\right]_{x=0}^{x=z} dz\,dy$

$= \int_1^4\int_y^4\left[\tan^{-1}(1) - \tan^{-1}(0)\right]dz\,dy = \int_1^4\int_y^4\left(\tfrac{\pi}{4} - 0\right)dz\,dy = \tfrac{\pi}{4}\int_1^4\left[z\right]_{z=y}^{z=4} dy$

$= \tfrac{\pi}{4}\int_1^4(4-y)\,dy = \tfrac{\pi}{4}\left[4y - \tfrac{1}{2}y^2\right]_1^4 = \tfrac{\pi}{4}\left(16 - 8 - 4 + \tfrac{1}{2}\right) = \tfrac{9\pi}{8}$

12. Here $E = \{(x,y,z) \mid 0 \le x \le \pi, 0 \le y \le \pi - x, 0 \le z \le x\}$, so

$\iiint_E \sin y\,dV = \int_0^{\pi}\int_0^{\pi-x}\int_0^x \sin y\,dz\,dy\,dx = \int_0^{\pi}\int_0^{\pi-x}\left[z\sin y\right]_{z=0}^{z=x} dy\,dx = \int_0^{\pi}\int_0^{\pi-x} x\sin y\,dy\,dx$

$= \int_0^{\pi}\left[-x\cos y\right]_{y=0}^{y=\pi-x} dx = \int_0^{\pi}\left[-x\cos(\pi-x) + x\right]dx$

$= \left[x\sin(\pi-x) - \cos(\pi-x) + \tfrac{1}{2}x^2\right]_0^{\pi}$ [integrate by parts]

$= 0 - 1 + \tfrac{1}{2}\pi^2 - 0 - 1 - 0 = \tfrac{1}{2}\pi^2 - 2$

13. Here $E = \{(x,y,z) \mid 0 \le x \le 1, 0 \le y \le \sqrt{x}, 0 \le z \le 1 + x + y\}$, so

$\iiint_E 6xy\,dV = \int_0^1\int_0^{\sqrt{x}}\int_0^{1+x+y} 6xy\,dz\,dy\,dx = \int_0^1\int_0^{\sqrt{x}}\left[6xyz\right]_{z=0}^{z=1+x+y} dy\,dx$

$= \int_0^1\int_0^{\sqrt{x}} 6xy(1+x+y)\,dy\,dx = \int_0^1\left[3xy^2 + 3x^2y^2 + 2xy^3\right]_{y=0}^{y=\sqrt{x}} dx$

$= \int_0^1\left(3x^2 + 3x^3 + 2x^{5/2}\right)dx = \left[x^3 + \tfrac{3}{4}x^4 + \tfrac{4}{7}x^{7/2}\right]_0^1 = \tfrac{65}{28}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

14.



Here $E = \{(x, y, z) \mid -1 \le x \le 1,\ 0 \le y \le 2,\ x^2 - 1 \le z \le 1 - x^2\}$.

Thus,

$$\begin{aligned}\iiint_E (x - y)\,dV &= \int_{-1}^{1}\int_{0}^{2}\int_{x^2-1}^{1-x^2} (x - y)\,dz\,dy\,dx \\ &= \int_{-1}^{1}\int_{0}^{2} (x - y)(1 - x^2 - (x^2 - 1))\,dy\,dx \\ &= \int_{-1}^{1}\int_{0}^{2} (2x - 2x^3 - 2y + 2x^2 y)\,dy\,dx \\ &= \int_{-1}^{1} \left[2xy - 2x^3 y - y^2 + x^2 y^2\right]_{y=0}^{y=2} dx \\ &= \int_{-1}^{1} (4x - 4x^3 - 4 + 4x^2)\,dx \\ &= \left[2x^2 - x^4 - 4x + \tfrac{4}{3}x^3\right]_{-1}^{1} = -\tfrac{5}{3} - \tfrac{11}{3} = -\tfrac{16}{3}\end{aligned}$$

15.



Here $T = \{(x, y, z) \mid 0 \le x \le 2,\ 0 \le y \le 2 - x,\ 0 \le z \le 2 - x - y\}$.

Thus,

$$\begin{aligned}\iiint_T y^2\,dV &= \int_{0}^{2}\int_{0}^{2-x}\int_{0}^{2-x-y} y^2\,dz\,dy\,dx \\ &= \int_{0}^{2}\int_{0}^{2-x} y^2(2 - x - y)\,dy\,dx \\ &= \int_{0}^{2}\int_{0}^{2-x} \left[(2 - x)y^2 - y^3\right] dy\,dx \\ &= \int_{0}^{2} \left[(2 - x)\left(\tfrac{1}{3}y^3\right) - \tfrac{1}{4}y^4\right]_{y=0}^{y=2-x} dx \\ &= \int_{0}^{2} \left[\tfrac{1}{3}(2 - x)^4 - \tfrac{1}{4}(2 - x)^4\right] dx = \int_{0}^{2} \tfrac{1}{12}(2 - x)^4\,dx \\ &= \left[\tfrac{1}{12}\left(-\tfrac{1}{5}\right)(2 - x)^5\right]_{0}^{2} = -\tfrac{1}{60}(0 - 32) = \tfrac{8}{15}\end{aligned}$$

16.



The projection of T onto the xz-plane is the triangle bounded by the lines $z = x$, $x = 0$, and $z = 1$. Then

$T = \{(x, y, z) \mid 0 \le x \le 1,\ x \le z \le 1,\ 0 \le y \le z - x\}$, and

$$\begin{aligned}\iiint_T xz\,dV &= \int_{0}^{1}\int_{x}^{1}\int_{0}^{z-x} xz\,dy\,dz\,dx = \int_{0}^{1}\int_{x}^{1} xz(z - x)\,dz\,dx \\ &= \int_{0}^{1}\int_{x}^{1} (xz^2 - x^2 z)\,dz\,dx = \int_{0}^{1} \left[\tfrac{1}{3}xz^3 - \tfrac{1}{2}x^2 z^2\right]_{z=x}^{z=1} dx \\ &= \int_{0}^{1} \left(\tfrac{1}{3}x - \tfrac{1}{2}x^2 - \tfrac{1}{3}x^4 + \tfrac{1}{2}x^4\right) dx \\ &= \left[\tfrac{1}{6}x^2 - \tfrac{1}{6}x^3 + \tfrac{1}{30}x^5\right]_{0}^{1} = \tfrac{1}{6} - \tfrac{1}{6} + \tfrac{1}{30} = \tfrac{1}{30}\end{aligned}$$

17.



The projection of E onto the yz-plane is the disk $y^2 + z^2 \le 1$. Using polar coordinates $y = r\cos\theta$ and $z = r\sin\theta$, we get

$$\begin{aligned}\iiint_E x\,dV &= \iint_D \left[\int_{4y^2+4z^2}^{4} x\,dx\right] dA = \tfrac{1}{2}\iint_D \left[4^2 - (4y^2 + 4z^2)^2\right] dA \\ &= 8\int_{0}^{2\pi}\int_{0}^{1} (1 - r^4)\,r\,dr\,d\theta = 8\int_{0}^{2\pi} d\theta \int_{0}^{1} (r - r^5)\,dr \\ &= 8(2\pi)\left[\tfrac{1}{2}r^2 - \tfrac{1}{6}r^6\right]_{0}^{1} = \tfrac{16\pi}{3}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18.



$$\int_0^1 \int_{3x}^3 \int_0^{\sqrt{9-y^2}} z\,dz\,dy\,dx = \int_0^1 \int_{3x}^3 \tfrac{1}{2}(9-y^2)\,dy\,dx$$
$$= \int_0^1 \left[\tfrac{9}{2}y - \tfrac{1}{6}y^3\right]_{y=3x}^{y=3} dx$$
$$= \int_0^1 \left[9 - \tfrac{27}{2}x + \tfrac{9}{2}x^3\right] dx$$
$$= \left[9x - \tfrac{27}{4}x^2 + \tfrac{9}{8}x^4\right]_0^1 = \tfrac{27}{8}$$

19. The plane $2x + y + z = 4$ intersects the xy-plane when

$2x + y + 0 = 4 \;\Rightarrow\; y = 4 - 2x$, so

$E = \{(x, y, z) \mid 0 \le x \le 2, 0 \le y \le 4 - 2x, 0 \le z \le 4 - 2x - y\}$ and



$$V = \int_0^2 \int_0^{4-2x} \int_0^{4-2x-y} dz\,dy\,dx = \int_0^2 \int_0^{4-2x} (4 - 2x - y)\,dy\,dx$$
$$= \int_0^2 \left[4y - 2xy - \tfrac{1}{2}y^2\right]_{y=0}^{y=4-2x} dx$$
$$= \int_0^2 \left[4(4-2x) - 2x(4-2x) - \tfrac{1}{2}(4-2x)^2\right] dx$$
$$= \int_0^2 (2x^2 - 8x + 8)\,dx = \left[\tfrac{2}{3}x^3 - 4x^2 + 8x\right]_0^2 = \tfrac{16}{3}$$

20. The paraboloids intersect when $x^2 + z^2 = 8 - x^2 - z^2 \;\Leftrightarrow\; x^2 + z^2 = 4$, thus the intersection is the circle $x^2 + z^2 = 4$, $y = 4$. The projection of E onto the xz-plane is the disk $x^2 + z^2 \le 4$, so

$E = \left\{(x, y, z) \mid x^2 + z^2 \le y \le 8 - x^2 - z^2, x^2 + z^2 \le 4\right\}$. Let

$D = \left\{(x, z) \mid x^2 + z^2 \le 4\right\}$. Then using polar coordinates $x = r\cos\theta$ and $z = r\sin\theta$, we have



$$V = \iiint_E dV = \iint_D \left(\int_{x^2+z^2}^{8-x^2-z^2} dy\right) dA = \iint_D (8 - 2x^2 - 2z^2)\,dA$$
$$= \int_0^{2\pi} \int_0^2 (8 - 2r^2)\,r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^2 (8r - 2r^3)\,dr$$
$$= \left[\theta\right]_0^{2\pi} \left[4r^2 - \tfrac{1}{2}r^4\right]_0^2 = 2\pi(16 - 8) = 16\pi$$

21. The plane $y + z = 1$ intersects the xy-plane in the line $y = 1$, so

$E = \left\{(x, y, z) \mid -1 \le x \le 1, x^2 \le y \le 1, 0 \le z \le 1 - y\right\}$ and



$$V = \iiint_E dV = \int_{-1}^1 \int_{x^2}^1 \int_0^{1-y} dz\,dy\,dx = \int_{-1}^1 \int_{x^2}^1 (1 - y)\,dy\,dx$$
$$= \int_{-1}^1 \left[y - \tfrac{1}{2}y^2\right]_{y=x^2}^{y=1} dx = \int_{-1}^1 \left(\tfrac{1}{2} - x^2 + \tfrac{1}{2}x^4\right) dx$$
$$= \left[\tfrac{1}{2}x - \tfrac{1}{3}x^3 + \tfrac{1}{10}x^5\right]_{-1}^1 = \tfrac{1}{2} - \tfrac{1}{3} + \tfrac{1}{10} + \tfrac{1}{2} - \tfrac{1}{3} + \tfrac{1}{10} = \tfrac{8}{15}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

22. Here $E = \{(x, y, z) \mid -1 \le y \le 4 - z, x^2 + z^2 \le 4\}$, so

$$V = \int_{-2}^{2} \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{-1}^{4-z} dy\, dz\, dx = \int_{-2}^{2} \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} (4 - z + 1)\, dz\, dx$$

$$= \int_{-2}^{2} \left[5z - \tfrac{1}{2}z^2\right]_{z=-\sqrt{4-x^2}}^{z=\sqrt{4-x^2}} dx = \int_{-2}^{2} 10\sqrt{4-x^2}\, dx$$

$$= 10\left[\tfrac{x}{2}\sqrt{4-x^2} + 2\sin^{-1}\left(\tfrac{x}{2}\right)\right]_{-2}^{2} \qquad \left[\begin{array}{l}\text{using trigonometric substitution or}\\ \text{Formula 30 in the Table of Integrals}\end{array}\right]$$

$$= 10\left[2\sin^{-1}(1) - 2\sin^{-1}(-1)\right] = 20\left(\tfrac{\pi}{2} - \left(-\tfrac{\pi}{2}\right)\right) = 20\pi$$



Alternatively, use polar coordinates to evaluate the double integral:

$$\int_{-2}^{2} \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} (5 - z)\, dz\, dx = \int_{0}^{2\pi} \int_{0}^{2} (5 - r\sin\theta)\, r\, dr\, d\theta$$

$$= \int_0^{2\pi} \left[\tfrac{5}{2}r^2 - \tfrac{1}{3}r^3\sin\theta\right]_{r=0}^{r=2} d\theta$$

$$= \int_0^{2\pi} \left(10 - \tfrac{8}{3}\sin\theta\right) d\theta$$

$$= 10\theta + \tfrac{8}{3}\cos\theta\Big]_0^{2\pi} = 20\pi$$

23. (a) The wedge can be described as the region

$$D = \{(x, y, z) \mid y^2 + z^2 \le 1, 0 \le x \le 1, 0 \le y \le x\}$$

$$= \left\{(x, y, z) \mid 0 \le x \le 1, 0 \le y \le x, 0 \le z \le \sqrt{1 - y^2}\right\}$$

So the integral expressing the volume of the wedge is

$$\iiint_D dV = \int_0^1 \int_0^x \int_0^{\sqrt{1-y^2}} dz\, dy\, dx.$$



(b) A CAS gives $\int_0^1 \int_0^x \int_0^{\sqrt{1-y^2}} dz\, dy\, dx = \frac{\pi}{4} - \frac{1}{3}$.

(Or use Formulas 30 and 87 from the Table of Integrals.)

24. (a) Divide B into 8 cubes of size $\Delta V = 8$. With $f(x, y, z) = \sqrt{x^2 + y^2 + z^2}$, the Midpoint Rule gives

$$\iiint_B \sqrt{x^2 + y^2 + z^2}\, dV \approx \sum_{i=1}^{2} \sum_{j=1}^{2} \sum_{k=1}^{2} f(\overline{x}_i, \overline{y}_j, \overline{z}_k)\, \Delta V$$

$$= 8[f(1,1,1) + f(1,1,3) + f(1,3,1) + f(1,3,3) + f(3,1,1)$$
$$+ f(3,1,3) + f(3,3,1) + f(3,3,3)]$$

$$\approx 239.64$$

(b) Using a CAS we have $\iiint_B \sqrt{x^2 + y^2 + z^2}\, dV = \int_0^1 \int_0^1 \int_0^1 \sqrt{x^2 + y^2 + z^2}\, dz\, dy\, dx \approx 245.91$. This differs from the estimate in part (a) by about 2.5%.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. Here $f(x,y,z) = \cos(xyz)$ and $\Delta V = \frac{1}{2}\cdot\frac{1}{2}\cdot\frac{1}{2} = \frac{1}{8}$, so the Midpoint Rule gives

$$\begin{aligned}\iiint_B f(x,y,z)\,dV &\approx \sum_{i=1}^{l}\sum_{j=1}^{m}\sum_{k=1}^{n} f(\overline{x}_i,\overline{y}_j,\overline{z}_k)\,\Delta V\\ &= \tfrac{1}{8}\left[f\left(\tfrac{1}{4},\tfrac{1}{4},\tfrac{1}{4}\right)+f\left(\tfrac{1}{4},\tfrac{1}{4},\tfrac{3}{4}\right)+f\left(\tfrac{1}{4},\tfrac{3}{4},\tfrac{1}{4}\right)+f\left(\tfrac{1}{4},\tfrac{3}{4},\tfrac{3}{4}\right)\right.\\ &\qquad \left.+f\left(\tfrac{3}{4},\tfrac{1}{4},\tfrac{1}{4}\right)+f\left(\tfrac{3}{4},\tfrac{1}{4},\tfrac{3}{4}\right)+f\left(\tfrac{3}{4},\tfrac{3}{4},\tfrac{1}{4}\right)+f\left(\tfrac{3}{4},\tfrac{3}{4},\tfrac{3}{4}\right)\right]\\ &= \tfrac{1}{8}\left[\cos\tfrac{1}{64}+\cos\tfrac{3}{64}+\cos\tfrac{3}{64}+\cos\tfrac{9}{64}+\cos\tfrac{3}{64}+\cos\tfrac{9}{64}+\cos\tfrac{9}{64}+\cos\tfrac{27}{64}\right]\approx 0.985\end{aligned}$$

26. Here $f(x,y,z) = \sqrt{x}\,e^{xyz}$ and $\Delta V = 2\cdot\frac{1}{2}\cdot 1 = 1$, so the Midpoint Rule gives

$$\begin{aligned}\iiint_B f(x,y,z)\,dV &\approx \sum_{i=1}^{l}\sum_{j=1}^{m}\sum_{k=1}^{n} f(\overline{x}_i,\overline{y}_j,\overline{z}_k)\,\Delta V\\ &= 1\left[f\left(1,\tfrac{1}{4},\tfrac{1}{2}\right)+f\left(1,\tfrac{1}{4},\tfrac{3}{2}\right)+f\left(1,\tfrac{3}{4},\tfrac{1}{2}\right)+f\left(1,\tfrac{3}{4},\tfrac{3}{2}\right)\right.\\ &\qquad \left.+f\left(3,\tfrac{1}{4},\tfrac{1}{2}\right)+f\left(3,\tfrac{1}{4},\tfrac{3}{2}\right)+f\left(3,\tfrac{3}{4},\tfrac{1}{2}\right)+f\left(3,\tfrac{3}{4},\tfrac{3}{2}\right)\right]\\ &= e^{1/8}+e^{3/8}+e^{3/8}+e^{9/8}+\sqrt{3}e^{3/8}+\sqrt{3}e^{9/8}+\sqrt{3}e^{9/8}+\sqrt{3}e^{27/8}\approx 70.932\end{aligned}$$

27. $E = \{(x,y,z) \mid 0\le x\le 1, 0\le z\le 1-x, 0\le y\le 2-2z\}$, the solid bounded by the three coordinate planes and the planes $z = 1-x$, $y = 2-2z$.



28. $E = \{(x,y,z) \mid 0\le y\le 2, 0\le z\le 2-y, 0\le x\le 4-y^2\}$, the solid bounded by the three coordinate planes, the plane $z = 2-y$, and the cylindrical surface $x = 4-y^2$.



29.









[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

If D_1, D_2, D_3 are the projections of E on the xy-, yz-, and xz-planes, then

$$D_1 = \{(x,y) \mid -2 \le x \le 2, 0 \le y \le 4 - x^2\} = \{(x,y) \mid 0 \le y \le 4, -\sqrt{4-y} \le x \le \sqrt{4-y}\}$$

$$D_2 = \{(y,z) \mid 0 \le y \le 4, -\tfrac{1}{2}\sqrt{4-y} \le z \le \tfrac{1}{2}\sqrt{4-y}\} = \{(y,z) \mid -1 \le z \le 1, 0 \le y \le 4 - 4z^2\}$$

$$D_3 = \{(x,z) \mid x^2 + 4z^2 \le 4\}$$

Therefore

$$\begin{aligned}
E &= \left\{(x,y,z) \mid -2 \le x \le 2, 0 \le y \le 4 - x^2, -\tfrac{1}{2}\sqrt{4 - x^2 - y} \le z \le \tfrac{1}{2}\sqrt{4 - x^2 - y}\right\} \\
&= \left\{(x,y,z) \mid 0 \le y \le 4, -\sqrt{4-y} \le x \le \sqrt{4-y}, -\tfrac{1}{2}\sqrt{4 - x^2 - y} \le z \le \tfrac{1}{2}\sqrt{4 - x^2 - y}\right\} \\
&= \left\{(x,y,z) \mid -1 \le z \le 1, 0 \le y \le 4 - 4z^2, -\sqrt{4 - y - 4z^2} \le x \le \sqrt{4 - y - 4z^2}\right\} \\
&= \left\{(x,y,z) \mid 0 \le y \le 4, -\tfrac{1}{2}\sqrt{4-y} \le z \le \tfrac{1}{2}\sqrt{4-y}, -\sqrt{4 - y - 4z^2} \le x \le \sqrt{4 - y - 4z^2}\right\} \\
&= \left\{(x,y,z) \mid -2 \le x \le 2, -\tfrac{1}{2}\sqrt{4 - x^2} \le z \le \tfrac{1}{2}\sqrt{4 - x^2}, 0 \le y \le 4 - x^2 - 4z^2\right\} \\
&= \left\{(x,y,z) \mid -1 \le z \le 1, -\sqrt{4 - 4z^2} \le x \le \sqrt{4 - 4z^2}, 0 \le y \le 4 - x^2 - 4z^2\right\}
\end{aligned}$$

Then

$$\begin{aligned}
\iiint_E f(x,y,z)\,dV &= \int_{-2}^{2}\int_{0}^{4-x^2}\int_{\sqrt{4-x^2-y}/2}^{\sqrt{4-x^2-y}/2} f(x,y,z)\,dz\,dy\,dx = \int_{0}^{4}\int_{-\sqrt{4-y}}^{\sqrt{4-y}}\int_{-\sqrt{4-x^2-y}/2}^{\sqrt{4-x^2-y}/2} f(x,y,z)\,dz\,dx\,dy \\
&= \int_{-1}^{1}\int_{0}^{4-4z^2}\int_{-\sqrt{4-y-4z^2}}^{\sqrt{4-y-4z^2}} f(x,y,z)\,dx\,dy\,dz = \int_{0}^{4}\int_{-\sqrt{4-y}/2}^{\sqrt{4-y}/2}\int_{-\sqrt{4-y-4z^2}}^{\sqrt{4-y-4z^2}} f(x,y,z)\,dx\,dz\,dy \\
&= \int_{-2}^{2}\int_{-\sqrt{4-x^2}/2}^{\sqrt{4-x^2}/2}\int_{0}^{4-x^2-4z^2} f(x,y,z)\,dy\,dz\,dx = \int_{-1}^{1}\int_{-\sqrt{4-4z^2}}^{\sqrt{4-4z^2}}\int_{0}^{4-x^2-4z^2} f(x,y,z)\,dy\,dx\,dz
\end{aligned}$$

30.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

If D_1, D_2, D_3 are the projections of E on the xy-, yz-, and xz-planes, then

$$D_1 = \{(x, y) \mid -2 \le x \le 2,\ -3 \le y \le 3\}$$
$$D_2 = \{(y, z) \mid y^2 + z^2 \le 9\}$$
$$D_3 = \{(x, z) \mid -2 \le x \le 2,\ -3 \le z \le 3\}$$

Therefore

$$E = \left\{(x, y, z) \mid -2 \le x \le 2,\ -3 \le y \le 3,\ -\sqrt{9 - y^2} \le z \le \sqrt{9 - y^2}\right\}$$
$$= \left\{(x, y, z) \mid -3 \le y \le 3,\ -\sqrt{9 - y^2} \le z \le \sqrt{9 - y^2},\ -2 \le x \le 2\right\}$$
$$= \left\{(x, y, z) \mid -3 \le z \le 3,\ -\sqrt{9 - z^2} \le y \le \sqrt{9 - z^2},\ -2 \le x \le 2\right\}$$
$$= \left\{(x, y, z) \mid -2 \le x \le 2,\ -3 \le z \le 3,\ -\sqrt{9 - z^2} \le y \le \sqrt{9 - z^2}\right\}$$

and

$$\iiint_E f(x, y, z)\, dV = \int_{-2}^{2} \int_{-3}^{3} \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} f(x, y, z)\, dz\, dy\, dx = \int_{-3}^{3} \int_{-2}^{2} \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} f(x, y, z)\, dz\, dx\, dy$$
$$= \int_{-3}^{3} \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} \int_{-2}^{2} f(x, y, z)\, dx\, dz\, dy = \int_{-3}^{3} \int_{-\sqrt{9-z^2}}^{\sqrt{9-z^2}} \int_{-2}^{2} f(x, y, z)\, dx\, dy\, dz$$
$$= \int_{-2}^{2} \int_{-3}^{3} \int_{-\sqrt{9-z^2}}^{\sqrt{9-z^2}} f(x, y, z)\, dy\, dz\, dx = \int_{-3}^{3} \int_{-2}^{2} \int_{-\sqrt{9-z^2}}^{\sqrt{9-z^2}} f(x, y, z)\, dy\, dx\, dz$$

31.









If D_1, D_2, and D_3 are the projections of E on the xy-, yz-, and xz-planes, then

$$D_1 = \left\{(x, y) \mid -2 \le x \le 2, x^2 \le y \le 4\right\} = \left\{(x, y) \mid 0 \le y \le 4, -\sqrt{y} \le x \le \sqrt{y}\right\},$$
$$D_2 = \left\{(y, z) \mid 0 \le y \le 4, 0 \le z \le 2 - \tfrac{1}{2}y\right\} = \left\{(y, z) \mid 0 \le z \le 2, 0 \le y \le 4 - 2z\right\}, \text{ and}$$
$$D_3 = \left\{(x, z) \mid -2 \le x \le 2, 0 \le z \le 2 - \tfrac{1}{2}x^2\right\} = \left\{(x, z) \mid 0 \le z \le 2, -\sqrt{4 - 2z} \le x \le \sqrt{4 - 2z}\right\}$$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Therefore
$$\begin{aligned} E &= \left\{(x,y,z) \mid -2 \le x \le 2, x^2 \le y \le 4, 0 \le z \le 2 - \tfrac{1}{2}y\right\} \\ &= \left\{(x,y,z) \mid 0 \le y \le 4, \ -\sqrt{y} \le x \le \sqrt{y}, 0 \le z \le 2 - \tfrac{1}{2}y\right\} \\ &= \left\{(x,y,z) \mid 0 \le y \le 4, 0 \le z \le 2 - \tfrac{1}{2}y, \ -\sqrt{y} \le x \le \sqrt{y}\right\} \\ &= \left\{(x,y,z) \mid 0 \le z \le 2, 0 \le y \le 4 - 2z, \ -\sqrt{y} \le x \le \sqrt{y}\right\} \\ &= \left\{(x,y,z) \mid -2 \le x \le 2, 0 \le z \le 2 - \tfrac{1}{2}x^2, x^2 \le y \le 4 - 2z\right\} \\ &= \left\{(x,y,z) \mid 0 \le z \le 2, \ -\sqrt{4-2z} \le x \le \sqrt{4-2z}, x^2 \le y \le 4 - 2z\right\} \end{aligned}$$

Then
$$\begin{aligned} \iiint_E f(x,y,z)\,dV &= \int_{-2}^{2}\int_{x^2}^{4}\int_{0}^{2-y/2} f(x,y,z)\,dz\,dy\,dx = \int_{0}^{4}\int_{-\sqrt{y}}^{\sqrt{y}}\int_{0}^{2-y/2} f(x,y,z)\,dz\,dx\,dy \\ &= \int_{0}^{4}\int_{0}^{2-y/2}\int_{-\sqrt{y}}^{\sqrt{y}} f(x,y,z)\,dx\,dz\,dy = \int_{0}^{2}\int_{0}^{4-2z}\int_{-\sqrt{y}}^{\sqrt{y}} f(x,y,z)\,dx\,dy\,dz \\ &= \int_{-2}^{2}\int_{0}^{2-x^2/2}\int_{x^2}^{4-2z} f(x,y,z)\,dy\,dz\,dx = \int_{0}^{2}\int_{-\sqrt{4-2z}}^{\sqrt{4-2z}}\int_{x^2}^{4-2z} f(x,y,z)\,dy\,dx\,dz \end{aligned}$$

32.



If D_1, D_2, and D_3 are the projections of E on the xy-, yz-, and xz-planes, then

$$D_1 = \{(x,y) \mid 0 \le x \le 2, 2 - x \le y \le 2\} = \{(x,y) \mid 0 \le y \le 2, 2 - y \le x \le 2\},$$
$$D_2 = \left\{(y,z) \mid 0 \le y \le 2, 0 \le z \le \tfrac{1}{2}y\right\} = \{(y,z) \mid 0 \le z \le 1, 2z \le y \le 2\}, \text{ and}$$
$$D_3 = \left\{(x,z) \mid 0 \le x \le 2, 0 \le z \le \tfrac{1}{2}x\right\} = \{(x,z) \mid 0 \le z \le 1, 2z \le x \le 2\}$$

Therefore
$$\begin{aligned} E &= \left\{(x,y,z) \mid 0 \le x \le 2, 2 - x \le y \le 2, 0 \le z \le \tfrac{1}{2}(x+y-2)\right\} \\ &= \left\{(x,y,z) \mid 0 \le y \le 2, 2 - y \le x \le 2, 0 \le z \le \tfrac{1}{2}(x+y-2)\right\} \\ &= \left\{(x,y,z) \mid 0 \le y \le 2, 0 \le z \le \tfrac{1}{2}y, 2 - y + 2z \le x \le 2\right\} \\ &= \{(x,y,z) \mid 0 \le z \le 1, 2z \le y \le 2, 2 - y + 2z \le x \le 2\} \\ &= \left\{(x,y,z) \mid 0 \le x \le 2, 0 \le z \le \tfrac{1}{2}x, 2 - x + 2z \le y \le 2\right\} \\ &= \{(x,y,z) \mid 0 \le z \le 1, 2z \le x \le 2, 2 - x + 2z \le y \le 2\} \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Then $\iiint_E f(x,y,z)\,dV = \int_0^2 \int_{2-x}^2 \int_0^{(x+y-2)/2} f(x,y,z)\,dz\,dy\,dx = \int_0^2 \int_{2-y}^2 \int_0^{(x+y-2)/2} f(x,y,z)\,dz\,dx\,dy$

$= \int_0^2 \int_0^{y/2} \int_{2-y+2z}^2 f(x,y,z)\,dx\,dz\,dy = \int_0^1 \int_{2z}^2 \int_{2-y+2z}^2 f(x,y,z)\,dx\,dy\,dz$

$= \int_0^2 \int_0^{x/2} \int_{2-x+2z}^2 f(x,y,z)\,dy\,dz\,dx = \int_0^1 \int_{2z}^2 \int_{2-x+2z}^2 f(x,y,z)\,dy\,dx\,dz$

33.



The diagrams show the projections of E onto the xy-, yz-, and xz-planes. Therefore

$\int_0^1 \int_{\sqrt{x}}^1 \int_0^{1-y} f(x,y,z)\,dz\,dy\,dx = \int_0^1 \int_0^{y^2} \int_0^{1-y} f(x,y,z)\,dz\,dx\,dy = \int_0^1 \int_0^{1-z} \int_0^{y^2} f(x,y,z)\,dx\,dy\,dz$

$= \int_0^1 \int_0^{1-y} \int_0^{y^2} f(x,y,z)\,dx\,dz\,dy = \int_0^1 \int_0^{1-\sqrt{x}} \int_{\sqrt{x}}^{1-z} f(x,y,z)\,dy\,dz\,dx$

$= \int_0^1 \int_0^{(1-z)^2} \int_{\sqrt{x}}^{1-z} f(x,y,z)\,dy\,dx\,dz$

34.



The projections of E onto the xy- and xz-planes are as in the first two diagrams and so

$\int_0^1 \int_0^{1-x^2} \int_0^{1-x} f(x,y,z)\,dy\,dz\,dx = \int_0^1 \int_0^{\sqrt{1-z}} \int_0^{1-x} f(x,y,z)\,dy\,dx\,dz$

$= \int_0^1 \int_0^{1-y} \int_0^{1-x^2} f(x,y,z)\,dz\,dx\,dy = \int_0^1 \int_0^{1-x} \int_0^{1-x^2} f(x,y,z)\,dz\,dy\,dx$

Now the surface $z = 1 - x^2$ intersects the plane $y = 1 - x$ in a curve whose projection in the yz-plane is $z = 1 - (1-y)^2$ or $z = 2y - y^2$. So we must split up the projection of E on the yz-plane into two regions as in the third diagram. For (y,z) in R_1, $0 \le x \le 1 - y$ and for (y,z) in R_2, $0 \le x \le \sqrt{1-z}$, and so the given integral is also equal to

$\int_0^1 \int_0^{1-\sqrt{1-z}} \int_0^{\sqrt{1-z}} f(x,y,z)\,dx\,dy\,dz + \int_0^1 \int_{1-\sqrt{1-z}}^1 \int_0^{1-y} f(x,y,z)\,dx\,dy\,dz$

$= \int_0^1 \int_0^{2y-y^2} \int_0^{1-y} f(x,y,z)\,dx\,dz\,dy + \int_0^1 \int_{2y-y^2}^1 \int_0^{\sqrt{1-z}} f(x,y,z)\,dx\,dz\,dy.$

35.



$\int_0^1 \int_y^1 \int_0^y f(x,y,z)\,dz\,dx\,dy = \iiint_E f(x,y,z)\,dV$ where $E = \{(x,y,z) \mid 0 \le z \le y,\ y \le x \le 1,\ 0 \le y \le 1\}$.

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

If D_1, D_2, and D_3 are the projections of E onto the xy-, yz- and xz-planes then

$$D_1 = \{(x,y) \mid 0 \le y \le 1, y \le x \le 1\} = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le x\},$$

$$D_2 = \{(y,z) \mid 0 \le y \le 1, 0 \le z \le y\} = \{(y,z) \mid 0 \le z \le 1, z \le y \le 1\}, \text{ and}$$

$$D_3 = \{(x,z) \mid 0 \le x \le 1, 0 \le z \le x\} = \{(x,z) \mid 0 \le z \le 1, z \le x \le 1\}.$$

Thus we also have

$$E = \{(x,y,z) \mid 0 \le x \le 1, 0 \le y \le x, 0 \le z \le y\} = \{(x,y,z) \mid 0 \le y \le 1, 0 \le z \le y, y \le x \le 1\}$$
$$= \{(x,y,z) \mid 0 \le z \le 1, z \le y \le 1, y \le x \le 1\} = \{(x,y,z) \mid 0 \le x \le 1, 0 \le z \le x, z \le y \le x\}$$
$$= \{(x,y,z) \mid 0 \le z \le 1, z \le x \le 1, z \le y \le x\}.$$

Then

$$\int_0^1 \int_y^1 \int_0^y f(x,y,z)\,dz\,dx\,dy = \int_0^1 \int_0^x \int_0^y f(x,y,z)\,dz\,dy\,dx = \int_0^1 \int_0^y \int_y^1 f(x,y,z)\,dx\,dz\,dy$$
$$= \int_0^1 \int_z^1 \int_y^1 f(x,y,z)\,dx\,dy\,dz = \int_0^1 \int_0^x \int_z^x f(x,y,z)\,dy\,dz\,dx$$
$$= \int_0^1 \int_z^1 \int_z^x f(x,y,z)\,dy\,dx\,dz$$

36.









$\int_0^1 \int_y^1 \int_0^z f(x,y,z)\,dx\,dz\,dy = \iiint_E f(x,y,z)\,dV$ where $E = \{(x,y,z) \mid 0 \le x \le z, y \le z \le 1, 0 \le y \le 1\}$.

Notice that E is bounded below by two different surfaces, so we must split the projection of E onto the xy-plane into two regions as in the second diagram. If D_1, D_2, and D_3 are the projections of E on the xy-, yz- and xz-planes then

$$D_1 = R_1 \cup R_2 = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le x\} \cup \{(x,y) \mid 0 \le x \le 1, x \le y \le 1\}$$
$$= \{(x,y) \mid 0 \le y \le 1, y \le x \le 1\} \cup \{(x,y) \mid 0 \le y \le 1, 0 \le x \le y\},$$

$$D_2 = \{(y,z) \mid 0 \le y \le 1, y \le z \le 1\} = \{(y,z) \mid 0 \le z \le 1, 0 \le y \le z\}, \text{ and}$$

$$D_3 = \{(x,z) \mid 0 \le x \le 1, x \le z \le 1\} = \{(x,z) \mid 0 \le z \le 1, 0 \le x \le z\}.$$

Thus we also have

$$E = \{(x,y,z) \mid 0 \le x \le 1, 0 \le y \le x, x \le z \le 1\} \cup \{(x,y,z) \mid 0 \le x \le 1, x \le y \le 1, y \le z \le 1\}$$
$$= \{(x,y,z) \mid 0 \le y \le 1, y \le x \le 1, x \le z \le 1\} \cup \{(x,y,z) \mid 0 \le y \le 1, 0 \le x \le y, y \le z \le 1\}$$
$$= \{(x,y,z) \mid 0 \le z \le 1, 0 \le y \le z, 0 \le x \le z\} = \{(x,y,z) \mid 0 \le x \le 1, x \le z \le 1, 0 \le y \le z\}$$
$$= \{(x,y,z) \mid 0 \le z \le 1, 0 \le x \le z, 0 \le y \le z\}.$$

Then

$$\int_0^1 \int_y^1 \int_0^z f(x,y,z)\,dx\,dz\,dy = \int_0^1 \int_0^x \int_x^1 f(x,y,z)\,dz\,dy\,dx + \int_0^1 \int_x^1 \int_y^1 f(x,y,z)\,dz\,dy\,dx$$
$$= \int_0^1 \int_y^1 \int_x^1 f(x,y,z)\,dz\,dx\,dy + \int_0^1 \int_0^y \int_y^1 f(x,y,z)\,dz\,dx\,dy$$
$$= \int_0^1 \int_0^z \int_0^z f(x,y,z)\,dx\,dy\,dz = \int_0^1 \int_x^1 \int_0^z f(x,y,z)\,dy\,dz\,dx$$
$$= \int_0^1 \int_0^z \int_0^z f(x,y,z)\,dy\,dx\,dz$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. The region C is the solid bounded by a circular cylinder of radius 2 with axis the z-axis for $-2 \le z \le 2$. We can write $\iiint_C (4 + 5x^2yz^2)\,dV = \iiint_C 4\,dV + \iiint_C 5x^2yz^2\,dV$, but $f(x, y, z) = 5x^2yz^2$ is an odd function with respect to y. Since C is symmetrical about the xz-plane, we have $\iiint_C 5x^2yz^2\,dV = 0$. Thus $\iiint_C (4 + 5x^2yz^2)\,dV = \iiint_C 4\,dV = 4 \cdot V(E) = 4 \cdot \pi(2)^2(4) = 64\pi$.

38. We can write $\iiint_B (z^3 + \sin y + 3)\,dV = \iiint_B z^3\,dV + \iiint_B \sin y\,dV + \iiint_B 3\,dV$. But z^3 is an odd function with respect to z and the region B is symmetric about the xy-plane, so $\iiint_B z^3\,dV = 0$. Similarly, $\sin y$ is an odd function with respect to y and B is symmetric about the xz-plane, so $\iiint_B \sin y\,dV = 0$. Thus $\iiint_B (z^3 + \sin y + 3)\,dV = \iiint_B 3\,dV = 3 \cdot V(B) = 3 \cdot \frac{4}{3}\pi(1)^3 = 4\pi$.

39. The projection of E onto the xy-plane is the disk $D = \{(x, y) \mid x^2 + y^2 \le 1\}$.

$$m = \iiint_E \rho(x, y, z)\,dV = \iint_D \left[\int_0^{1-x^2-y^2} 3\,dz\right] dA = \iint_D 3(1 - x^2 - y^2)\,dA$$
$$= 3\int_0^1 \int_0^{2\pi} (1 - r^2)\,r\,dr\,d\theta = 3\int_0^{2\pi} d\theta \int_0^1 (r - r^3)\,dr$$
$$= 3\left[\theta\right]_0^{2\pi} \left[\tfrac{1}{2}r^2 - \tfrac{1}{4}r^4\right]_0^1 = 3\,(2\pi)\left(\tfrac{1}{2} - \tfrac{1}{4}\right) = \tfrac{3}{2}\pi$$

$$M_{yz} = \iiint_E x\rho(x, y, z)\,dV = \iint_D \left[\int_0^{1-x^2-y^2} 3x\,dz\right] dA = \iint_D 3x(1 - x^2 - y^2)\,dA$$
$$= 3\int_0^1 \int_0^{2\pi} (r\cos\theta)(1 - r^2)\,r\,dr\,d\theta = 3\int_0^{2\pi} \cos\theta\,d\theta \int_0^1 (r^2 - r^4)\,dr$$
$$= 3\left[\sin\theta\right]_0^{2\pi} \left[\tfrac{1}{3}r^3 - \tfrac{1}{5}r^5\right]_0^1 = 3\,(0)\left(\tfrac{1}{3} - \tfrac{1}{5}\right) = 0$$

$$M_{xz} = \iiint_E y\rho(x, y, z)\,dV = \iint_D \left[\int_0^{1-x^2-y^2} 3y\,dz\right] dA = \iint_D 3y(1 - x^2 - y^2)\,dA$$
$$= 3\int_0^1 \int_0^{2\pi} (r\sin\theta)(1 - r^2)\,r\,dr\,d\theta = 3\int_0^{2\pi} \sin\theta\,d\theta \int_0^1 (r^2 - r^4)\,dr$$
$$= 3\left[-\cos\theta\right]_0^{2\pi} \left[\tfrac{1}{3}r^3 - \tfrac{1}{5}r^5\right]_0^1 = 3\,(0)\left(\tfrac{1}{3} - \tfrac{1}{5}\right) = 0$$

$$M_{xy} = \iiint_E z\rho(x, y, z)\,dV = \iint_D \left[\int_0^{1-x^2-y^2} 3z\,dz\right] dA = \iint_D \left[\tfrac{3}{2}z^2\right]_{z=0}^{z=1-x^2-y^2} dA$$
$$= \tfrac{3}{2}\iint_D (1 - x^2 - y^2)^2\,dA = \tfrac{3}{2}\int_0^1 \int_0^{2\pi} (1 - r^2)^2\,r\,dr\,d\theta$$
$$= \tfrac{3}{2}\int_0^{2\pi} d\theta \int_0^1 (r - 2r^3 + r^5)\,dr = \tfrac{3}{2}\left[\theta\right]_0^{2\pi} \left[\tfrac{1}{2}r^2 - \tfrac{1}{2}r^4 + \tfrac{1}{6}r^6\right]_0^1$$
$$= \tfrac{3}{2}\,(2\pi)\left(\tfrac{1}{2} - \tfrac{1}{2} + \tfrac{1}{6}\right) = \tfrac{1}{2}\pi$$

Thus the mass is $\frac{3}{2}\pi$ and the center of mass is $(\overline{x}, \overline{y}, \overline{z}) = \left(\dfrac{M_{yz}}{m}, \dfrac{M_{xz}}{m}, \dfrac{M_{xy}}{m}\right) = \left(0, 0, \dfrac{1}{3}\right)$.

40. $m = \int_{-1}^1 \int_0^{1-y^2} \int_0^{1-z} 4\,dx\,dz\,dy = 4\int_{-1}^1 \int_0^{1-y^2} (1 - z)\,dz\,dy = 4\int_{-1}^1 \left[z - \frac{1}{2}z^2\right]_{z=0}^{z=1-y^2} dy = 2\int_{-1}^1 (1 - y^4)\,dy = \frac{16}{5}$,

$$M_{yz} = \int_{-1}^1 \int_0^{1-y^2} \int_0^{1-z} 4x\,dx\,dz\,dy = 2\int_{-1}^1 \int_0^{1-y^2} (1 - z)^2\,dz\,dy = 2\int_{-1}^1 \left[-\tfrac{1}{3}(1 - z)^3\right]_{z=0}^{z=1-y^2} dy$$
$$= \tfrac{2}{3}\int_{-1}^1 (1 - y^6)\,dy = \left(\tfrac{4}{3}\right)\left(\tfrac{6}{7}\right) = \tfrac{24}{21}$$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$M_{xz} = \int_{-1}^{1}\int_{0}^{1-y^2}\int_{0}^{1-z} 4y\,dx\,dz\,dy = \int_{-1}^{1}\int_{0}^{1-y^2} 4y(1-z)\,dz\,dy$

$= \int_{-1}^{1}\left[4y(1-y^2) - 2y(1-y^2)^2\right]dy = \int_{-1}^{1}(2y-2y^5)\,dy = 0$ [the integrand is odd]

$M_{xy} = \int_{-1}^{1}\int_{0}^{1-y^2}\int_{0}^{1-z} 4z\,dx\,dz\,dy = \int_{-1}^{1}\int_{0}^{1-y^2}(4z-4z^2)\,dz\,dy = 2\int_{-1}^{1}\left[(1-y^2)^2 - \frac{2}{3}(1-y^2)^3\right]dy$

$= 2\int_{-1}^{1}\left[\frac{1}{3} - y^4 + \frac{2}{3}y^6\right]dy = \left[\frac{4}{3}y - \frac{4}{5}y^5 + \frac{8}{21}y^7\right]_0^1 = \frac{96}{105} = \frac{32}{35}$

Thus, $(\overline{x},\overline{y},\overline{z}) = \left(\frac{5}{14}, 0, \frac{2}{7}\right)$

41. $m = \int_0^a\int_0^a\int_0^a (x^2+y^2+z^2)\,dx\,dy\,dz = \int_0^a\int_0^a\left[\frac{1}{3}x^3 + xy^2 + xz^2\right]_{x=0}^{x=a} dy\,dz = \int_0^a\int_0^a\left(\frac{1}{3}a^3 + ay^2 + az^2\right)dy\,dz$

$= \int_0^a\left[\frac{1}{3}a^3y + \frac{1}{3}ay^3 + ayz^2\right]_{y=0}^{y=a} dz = \int_0^a\left(\frac{2}{3}a^4 + a^2z^2\right)dz = \left[\frac{2}{3}a^4z + \frac{1}{3}a^2z^3\right]_0^a = \frac{2}{3}a^5 + \frac{1}{3}a^5 = a^5$

$M_{yz} = \int_0^a\int_0^a\int_0^a \left[x^3 + x(y^2+z^2)\right]dx\,dy\,dz = \int_0^a\int_0^a\left[\frac{1}{4}a^4 + \frac{1}{2}a^2(y^2+z^2)\right]dy\,dz$

$= \int_0^a\left(\frac{1}{4}a^5 + \frac{1}{6}a^5 + \frac{1}{2}a^3z^2\right)dz = \frac{1}{4}a^6 + \frac{1}{3}a^6 = \frac{7}{12}a^6 = M_{xz} = M_{xy}$ by symmetry of E and $\rho(x,y,z)$

Hence $(\overline{x},\overline{y},\overline{z}) = \left(\frac{7}{12}a, \frac{7}{12}a, \frac{7}{12}a\right)$.

42. $m = \int_0^1\int_0^{1-x}\int_0^{1-x-y} y\,dz\,dy\,dx = \int_0^1\int_0^{1-x}\left[(1-x)y - y^2\right]dy\,dx$

$= \int_0^1\left[\frac{1}{2}(1-x)^3 - \frac{1}{3}(1-x)^3\right]dx = \frac{1}{6}\int_0^1(1-x)^3\,dx = \frac{1}{24}$

$M_{yz} = \int_0^1\int_0^{1-x}\int_0^{1-x-y} xy\,dz\,dy\,dx = \int_0^1\int_0^{1-x}\left[(x-x^2)y - xy^2\right]dy\,dx$

$= \int_0^1\left[\frac{1}{2}x(1-x)^3 - \frac{1}{3}x(1-x)^3\right]dx = \frac{1}{6}\int_0^1\left(x - 3x^2 + 3x^3 - x^4\right)dx = \frac{1}{6}\left(\frac{1}{2} - 1 + \frac{3}{4} - \frac{1}{5}\right) = \frac{1}{120}$

$M_{xz} = \int_0^1\int_0^{1-x}\int_0^{1-x-y} y^2\,dz\,dy\,dx = \int_0^1\int_0^{1-x}\left[(1-x)y^2 - y^3\right]dy\,dx$

$= \int_0^1\left[\frac{1}{3}(1-x)^4 - \frac{1}{4}(1-x)^4\right]dx = \frac{1}{12}\left[-\frac{1}{5}(1-x)^5\right]_0^1 = \frac{1}{60}$

$M_{xy} = \int_0^1\int_0^{1-x}\int_0^{1-x-y} yz\,dz\,dy\,dx = \int_0^1\int_0^{1-x}\left[\frac{1}{2}y(1-x-y)^2\right]dy\,dx$

$= \frac{1}{2}\int_0^1\int_0^{1-x}\left[(1-x)^2y - 2(1-x)y^2 + y^3\right]dy\,dx = \frac{1}{2}\int_0^1\left[\frac{1}{2}(1-x)^4 - \frac{2}{3}(1-x)^4 + \frac{1}{4}(1-x)^4\right]dx$

$= \frac{1}{24}\int_0^1(1-x)^4\,dx = -\frac{1}{24}\left[\frac{1}{5}(1-x)^5\right]_0^1 = \frac{1}{120}$

Hence $(\overline{x},\overline{y},\overline{z}) = \left(\frac{1}{5}, \frac{2}{5}, \frac{1}{5}\right)$.

43. $I_x = \int_0^L\int_0^L\int_0^L k(y^2+z^2)\,dz\,dy\,dx = k\int_0^L\int_0^L\left(Ly^2 + \frac{1}{3}L^3\right)dy\,dx = k\int_0^L\frac{2}{3}L^4\,dx = \frac{2}{3}kL^5$

By symmetry, $I_x = I_y = I_z = \frac{2}{3}kL^5$.

44. $I_x = \int_{-c/2}^{c/2}\int_{-b/2}^{b/2}\int_{-a/2}^{a/2} k(y^2+z^2)\,dx\,dy\,dz = ka\int_{-c/2}^{c/2}\int_{-b/2}^{b/2}(y^2+z^2)\,dy\,dz$

$= ak\int_{-c/2}^{c/2}\left[\frac{1}{3}y^3 + z^2y\right]_{y=-b/2}^{y=b/2} dz = ak\int_{-c/2}^{c/2}\left(\frac{1}{12}b^3 + bz^2\right)dz = ak\left[\frac{1}{12}b^3z + \frac{1}{3}bz^3\right]_{-c/2}^{c/2}$

$= ak\left(\frac{1}{12}b^3c + \frac{1}{12}bc^3\right) = \frac{1}{12}kabc(b^2+c^2)$

By symmetry, $I_y = \frac{1}{12}kabc(a^2+c^2)$ and $I_z = \frac{1}{12}kabc(a^2+b^2)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. $I_z = \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV = \iint\limits_{x^2+y^2\le a^2} \left[\int_0^h k(x^2+y^2)\,dz\right] dA = \iint\limits_{x^2+y^2\le a^2} k(x^2+y^2)h\,dA$

$= kh\int_0^{2\pi}\int_0^a (r^2)\,r\,dr\,d\theta = kh\int_0^{2\pi} d\theta \int_0^a r^3\,dr = kh(2\pi)\left[\tfrac{1}{4}r^4\right]_0^a = 2\pi kh\cdot\tfrac{1}{4}a^4 = \tfrac{1}{2}\pi kha^4$

46. $I_z = \iiint_E (x^2+y^2)\rho(x,y,z)\,dV = \iint\limits_{x^2+y^2\le h^2} \left[\int_{\sqrt{x^2+y^2}}^h k(x^2+y^2)\,dz\right] dA$

$= \iint\limits_{x^2+y^2\le h^2} k(x^2+y^2)\left(h-\sqrt{x^2+y^2}\right) dA = k\int_0^{2\pi}\int_0^h r^2(h-r)\,r\,dr\,d\theta$

$= k\int_0^{2\pi} d\theta \int_0^h \left(r^3h - r^4\right) dr = k(2\pi)\left[\tfrac{1}{4}r^4h - \tfrac{1}{5}r^5\right]_0^h = 2\pi k\left(\tfrac{1}{4}h^5 - \tfrac{1}{5}h^5\right) = \tfrac{1}{10}\pi kh^5$

47. (a) $m = \int_{-1}^1\int_{x^2}^1\int_0^{1-y} \sqrt{x^2+y^2}\,dz\,dy\,dx$

(b) $(\overline{x},\overline{y},\overline{z})$ where $\overline{x} = \frac{1}{m}\int_{-1}^1\int_{x^2}^1\int_0^{1-y} x\sqrt{x^2+y^2}\,dz\,dy\,dx$, $\overline{y} = \frac{1}{m}\int_{-1}^1\int_{x^2}^1\int_0^{1-y} y\sqrt{x^2+y^2}\,dz\,dy\,dx$, and

$\overline{z} = \frac{1}{m}\int_{-1}^1\int_{x^2}^1\int_0^{1-y} z\sqrt{x^2+y^2}\,dz\,dy\,dx$.

(c) $I_z = \int_{-1}^1\int_{x^2}^1\int_0^{1-y}(x^2+y^2)\sqrt{x^2+y^2}\,dz\,dy\,dx = \int_{-1}^1\int_{x^2}^1\int_0^{1-y}(x^2+y^2)^{3/2}\,dz\,dy\,dx$

48. (a) $m = \int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}} \sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$

(b) $(\overline{x},\overline{y},\overline{z})$ where $\overline{x} = m^{-1}\int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}} x\sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$,

$\overline{y} = m^{-1}\int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}} y\sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$,

$\overline{z} = m^{-1}\int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}} z\sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$

(c) $I_z = \int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}}(x^2+y^2)(1+x+y+z)\,dz\,dx\,dy$

49. (a) $m = \int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y (1+x+y+z)\,dz\,dy\,dx = \frac{3\pi}{32} + \frac{11}{24}$

(b) $(\overline{x},\overline{y},\overline{z}) = \Big(m^{-1}\int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y x(1+x+y+z)\,dz\,dy\,dx,$

$m^{-1}\int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y y(1+x+y+z)\,dz\,dy\,dx,$

$m^{-1}\int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y z(1+x+y+z)\,dz\,dy\,dx\Big)$

$= \left(\dfrac{28}{9\pi+44}, \dfrac{30\pi+128}{45\pi+220}, \dfrac{45\pi+208}{135\pi+660}\right)$

(c) $I_z = \displaystyle\int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y (x^2+y^2)(1+x+y+z)\,dz\,dy\,dx = \frac{68+15\pi}{240}$

50. (a) $m = \int_0^1\int_{3x}^3\int_0^{\sqrt{9-y^2}}(x^2+y^2)\,dz\,dy\,dx = \frac{56}{5} = 11.2$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) $(\overline{x},\overline{y},\overline{z})$ where $\overline{x} = m^{-1}\int_0^1\int_{3x}^3\int_0^{\sqrt{9-y^2}} x(x^2+y^2)\,dz\,dy\,dx \approx 0.375$,

$\overline{y} = m^{-1}\int_0^1\int_{3x}^3\int_0^{\sqrt{9-y^2}} y(x^2+y^2)\,dz\,dy\,dx = \frac{45\pi}{64} \approx 2.209$,

$\overline{z} = m^{-1}\int_0^1\int_{3x}^3\int_0^{\sqrt{9-y^2}} z(x^2+y^2)\,dz\,dy\,dx = \frac{15}{16} = 0.9375$.

(c) $I_z = \int_0^1\int_{3x}^3\int_0^{\sqrt{9-y^2}} (x^2+y^2)^2\,dz\,dy\,dx = \frac{10{,}464}{175} \approx 59.79$

51. (a) $f(x,y,z)$ is a joint density function, so we know $\iiint_{\mathbb{R}^3} f(x,y,z)\,dV = 1$. Here we have

$$\begin{aligned}\iiint_{\mathbb{R}^3} f(x,y,z)\,dV &= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y,z)\,dz\,dy\,dx = \int_0^2\int_0^2\int_0^2 Cxyz\,dz\,dy\,dx\\ &= C\int_0^2 x\,dx\int_0^2 y\,dy\int_0^2 z\,dz = C\left[\tfrac12 x^2\right]_0^2\left[\tfrac12 y^2\right]_0^2\left[\tfrac12 z^2\right]_0^2 = 8C\end{aligned}$$

Then we must have $8C = 1 \;\Rightarrow\; C = \frac18$.

(b) $P(X\le 1, Y\le 1, Z\le 1) = \int_{-\infty}^1\int_{-\infty}^1\int_{-\infty}^1 f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^1\int_0^1 \tfrac18 xyz\,dz\,dy\,dx$

$$= \tfrac18\int_0^1 x\,dx\int_0^1 y\,dy\int_0^1 z\,dz = \tfrac18\left[\tfrac12 x^2\right]_0^1\left[\tfrac12 y^2\right]_0^1\left[\tfrac12 z^2\right]_0^1 = \tfrac18\left(\tfrac12\right)^3 = \tfrac{1}{64}$$

(c) $P(X+Y+Z\le 1) = P((X,Y,Z)\in E)$ where E is the solid region in the first octant bounded by the coordinate planes and the plane $x+y+z=1$. The plane $x+y+z=1$ meets the xy-plane in the line $x+y=1$, so we have

$$\begin{aligned}P(X+Y+Z\le 1) &= \iiint_E f(x,y,z)\,dV = \int_0^1\int_0^{1-x}\int_0^{1-x-y} \tfrac18 xyz\,dz\,dy\,dx\\ &= \tfrac18\int_0^1\int_0^{1-x} xy\left[\tfrac12 z^2\right]_{z=0}^{z=1-x-y} dy\,dx = \tfrac{1}{16}\int_0^1\int_0^{1-x} xy(1-x-y)^2\,dy\,dx\\ &= \tfrac{1}{16}\int_0^1\int_0^{1-x}[(x^3-2x^2+x)y + (2x^2-2x)y^2 + xy^3]\,dy\,dx\\ &= \tfrac{1}{16}\int_0^1\left[(x^3-2x^2+x)\tfrac12 y^2 + (2x^2-2x)\tfrac13 y^3 + x\left(\tfrac14 y^4\right)\right]_{y=0}^{y=1-x} dx\\ &= \tfrac{1}{192}\int_0^1 (x-4x^2+6x^3-4x^4+x^5)\,dx = \tfrac{1}{192}\left(\tfrac{1}{30}\right) = \tfrac{1}{5760}\end{aligned}$$

52. (a) $f(x,y,z)$ is a joint density function, so we know $\iiint_{\mathbb{R}^3} f(x,y,z)\,dV = 1$. Here we have

$$\begin{aligned}\iiint_{\mathbb{R}^3} f(x,y,z)\,dV &= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y,z)\,dz\,dy\,dx = \int_0^{\infty}\int_0^{\infty}\int_0^{\infty} Ce^{-(0.5x+0.2y+0.1z)}\,dz\,dy\,dx\\ &= C\int_0^{\infty} e^{-0.5x}\,dx\int_0^{\infty} e^{-0.2y}\,dy\int_0^{\infty} e^{-0.1z}\,dz\\ &= C\lim_{t\to\infty}\int_0^t e^{-0.5x}\,dx\lim_{t\to\infty}\int_0^t e^{-0.2y}\,dy\lim_{t\to\infty}\int_0^t e^{-0.1z}\,dz\\ &= C\lim_{t\to\infty}\left[-2e^{-0.5x}\right]_0^t\lim_{t\to\infty}\left[-5e^{-0.2y}\right]_0^t\lim_{t\to\infty}\left[-10e^{-0.1z}\right]_0^t\\ &= C\lim_{t\to\infty}\left[-2(e^{-0.5t}-1)\right]\lim_{t\to\infty}\left[-5(e^{-0.2t}-1)\right]\lim_{t\to\infty}\left[-10(e^{-0.1t}-1)\right]\\ &= C\cdot(-2)(0-1)\cdot(-5)(0-1)\cdot(-10)(0-1) = 100C\end{aligned}$$

So we must have $100C = 1 \;\Rightarrow\; C = \frac{1}{100}$.

(b) We have no restriction on Z, so

$$\begin{aligned}P(X\le 1, Y\le 1) &= \int_{-\infty}^1\int_{-\infty}^1\int_{-\infty}^{\infty} f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^1\int_0^{\infty}\tfrac{1}{100}e^{-(0.5x+0.2y+0.1z)}\,dz\,dy\,dx\\ &= \tfrac{1}{100}\int_0^1 e^{-0.5x}\,dx\int_0^1 e^{-0.2y}\,dy\int_0^{\infty} e^{-0.1z}\,dz\\ &= \tfrac{1}{100}\left[-2e^{-0.5x}\right]_0^1\left[-5e^{-0.2y}\right]_0^1\lim_{t\to\infty}\left[-10e^{-0.1z}\right]_0^t \qquad \text{[by part (a)]}\\ &= \tfrac{1}{100}(2-2e^{-0.5})(5-5e^{-0.2})(10) = (1-e^{-0.5})(1-e^{-0.2}) \approx 0.07132\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $P(X \le 1, Y \le 1, Z \le 1) = \int_{-\infty}^{1}\int_{-\infty}^{1}\int_{-\infty}^{1} f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^1\int_0^1 \frac{1}{100}e^{-(0.5x+0.2y+0.1z)}\,dz\,dy\,dx$

$$= \tfrac{1}{100}\int_0^1 e^{-0.5x}\,dx \int_0^1 e^{-0.2y}\,dy \int_0^1 e^{-0.1z}\,dz$$

$$= \tfrac{1}{100}\left[-2e^{-0.5x}\right]_0^1\left[-5e^{-0.2y}\right]_0^1\left[-10e^{-0.1z}\right]_0^1$$

$$= (1-e^{-0.5})(1-e^{-0.2})(1-e^{-0.1}) \approx 0.006787$$

53. $V(E) = L^3 \quad\Rightarrow\quad f_{\text{ave}} = \dfrac{1}{L^3}\displaystyle\int_0^L\int_0^L\int_0^L xyz\,dx\,dy\,dz = \frac{1}{L^3}\int_0^L x\,dx \int_0^L y\,dy \int_0^L z\,dz$

$$= \frac{1}{L^3}\left[\frac{x^2}{2}\right]_0^L\left[\frac{y^2}{2}\right]_0^L\left[\frac{z^2}{2}\right]_0^L = \frac{1}{L^3}\frac{L^2}{2}\frac{L^2}{2}\frac{L^2}{2} = \frac{L^3}{8}$$

54. The height of each point is given by its z-coordinate, so the average height of the points in $E = \{(x,y,z) \mid x^2+y^2+z^2 \le 1,\ z \ge 0\}$ is

$$\frac{1}{V(E)}\iiint_E z\,dV$$

Here $V(E) = \frac{1}{2}\cdot\frac{4}{3}\pi(1)^3 = \frac{2}{3}\pi$ [half the volume of a sphere], so

$$\tfrac{1}{V(E)}\iiint_E z\,dV = \tfrac{1}{2\pi/3}\int_{-1}^1\int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}}\int_0^{\sqrt{1-x^2-y^2}} z\,dz\,dy\,dx = \tfrac{3}{2\pi}\int_{-1}^1\int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}}\left[\tfrac{1}{2}z^2\right]_{z=0}^{z=\sqrt{1-x^2-y^2}}dy\,dx$$

$$= \tfrac{3}{2\pi}\cdot\tfrac{1}{2}\int_{-1}^1\int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}}(1-x^2-y^2)\,dy\,dx = \tfrac{3}{4\pi}\int_0^{2\pi}\int_0^1 (1-r^2)\,r\,dr\,d\theta$$

$$= \tfrac{3}{4\pi}\int_0^{2\pi} d\theta \int_0^1 (r-r^3)\,dr = \tfrac{3}{4\pi}(2\pi)\left[\tfrac{1}{2}r^2 - \tfrac{1}{4}r^4\right]_0^1 = \tfrac{3}{2}\left(\tfrac{1}{4}\right) = \tfrac{3}{8}$$

55. (a) The triple integral will attain its maximum when the integrand $1-x^2-2y^2-3z^2$ is positive in the region E and negative everywhere else. For if E contains some region F where the integrand is negative, the integral could be increased by excluding F from E, and if E fails to contain some part G of the region where the integrand is positive, the integral could be increased by including G in E. So we require that $x^2+2y^2+3z^2 \le 1$. This describes the region bounded by the ellipsoid $x^2+2y^2+3z^2 = 1$.

(b) The maximum value of $\iiint_E (1-x^2-2y^2-3z^2)\,dV$ occurs when E is the solid region bounded by the ellipsoid $x^2+2y^2+3z^2 = 1$. The projection of E on the xy-plane is the planar region bounded by the ellipse $x^2+2y^2 = 1$, so

$$E = \left\{(x,y,z) \mid -1 \le x \le 1, -\sqrt{\tfrac{1}{2}(1-x^2)} \le y \le \sqrt{\tfrac{1}{2}(1-x^2)}, -\sqrt{\tfrac{1}{3}(1-x^2-2y^2)} \le z \le \sqrt{\tfrac{1}{3}(1-x^2-2y^2)}\right\}$$

and

$$\iiint_E (1-x^2-2y^2-3z^2)\,dV = \int_{-1}^{1}\int_{-\sqrt{\frac{1}{2}(1-x^2)}}^{\sqrt{\frac{1}{2}(1-x^2)}}\int_{-\sqrt{\frac{1}{3}(1-x^2-2y^2)}}^{\sqrt{\frac{1}{3}(1-x^2-2y^2)}} (1-x^2-2y^2-3z^2)\,dz\,dy\,dx = \frac{4\sqrt{6}}{45}\pi$$

using a CAS.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

DISCOVERY PROJECT Volumes of Hyperspheres

In this project we use V_n to denote the n-dimensional volume of an n-dimensional hypersphere.

1. The interior of the circle is the set of points $\left\{(x,y) \mid -r \le y \le r,\ -\sqrt{r^2-y^2} \le x \le \sqrt{r^2-y^2}\right\}$. So, substituting $y = r\sin\theta$ and then using Formula 64 to evaluate the integral, we get

$$V_2 = \int_{-r}^{r}\int_{-\sqrt{r^2-y^2}}^{\sqrt{r^2-y^2}} dx\,dy = \int_{-r}^{r} 2\sqrt{r^2-y^2}\,dy = \int_{-\pi/2}^{\pi/2} 2r\sqrt{1-\sin^2\theta}\,(r\cos\theta\,d\theta)$$

$$= 2r^2\int_{-\pi/2}^{\pi/2}\cos^2\theta\,d\theta = 2r^2\left[\tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta\right]_{-\pi/2}^{\pi/2} = 2r^2\left(\tfrac{\pi}{2}\right) = \pi r^2$$

2. The region of integration is $\left\{(x,y,z) \mid -r \le z \le r, -\sqrt{r^2-z^2} \le y \le \sqrt{r^2-z^2}, -\sqrt{r^2-z^2-y^2} \le x \le \sqrt{r^2-z^2-y^2}\right\}$. Substituting $y = \sqrt{r^2-z^2}\sin\theta$ and using Formula 64 to integrate $\cos^2\theta$, we get

$$V_3 = \int_{-r}^{r}\int_{-\sqrt{r^2-z^2}}^{\sqrt{r^2-z^2}}\int_{-\sqrt{r^2-z^2-y^2}}^{\sqrt{r^2-z^2-y^2}} dx\,dy\,dz = \int_{-r}^{r}\int_{-\sqrt{r^2-z^2}}^{\sqrt{r^2-z^2}} 2\sqrt{r^2-z^2-y^2}\,dy\,dz$$

$$= \int_{-r}^{r}\int_{-\pi/2}^{\pi/2} 2\sqrt{r^2-z^2}\sqrt{1-\sin^2\theta}\left(\sqrt{r^2-z^2}\cos\theta\,d\theta\right)dz$$

$$= 2\left[\int_{-r}^{r}(r^2-z^2)\,dz\right]\left[\int_{-\pi/2}^{\pi/2}\cos^2\theta\,d\theta\right] = 2\left(\frac{4r^3}{3}\right)\left(\frac{\pi}{2}\right) = \frac{4\pi r^3}{3}$$

3. Here we substitute $y = \sqrt{r^2 - w^2 - z^2}\sin\theta$ and, later, $w = r\sin\phi$. Because $\int_{-\pi/2}^{\pi/2}\cos^p\theta\,d\theta$ seems to occur frequently in these calculations, it is useful to find a general formula for that integral. From Exercises 7.1.49-50, we have

$$\int_0^{\pi/2}\sin^{2k}x\,dx = \frac{1\cdot3\cdot5\cdot\cdots\cdot(2k-1)}{2\cdot4\cdot6\cdot\cdots\cdot2k}\frac{\pi}{2} \qquad \text{and} \qquad \int_0^{\pi/2}\sin^{2k+1}x\,dx = \frac{2\cdot4\cdot6\cdot\cdots\cdot2k}{1\cdot3\cdot5\cdot\cdots\cdot(2k+1)}$$

and from the symmetry of the sine and cosine functions, we can conclude that

$$\int_{-\pi/2}^{\pi/2}\cos^{2k}x\,dx = 2\int_0^{\pi/2}\sin^{2k}x\,dx = \frac{1\cdot3\cdot5\cdot\cdots\cdot(2k-1)\pi}{2\cdot4\cdot6\cdot\cdots\cdot2k} \tag{1}$$

$$\int_{-\pi/2}^{\pi/2}\cos^{2k+1}x\,dx = 2\int_0^{\pi/2}\sin^{2k+1}x\,dx = \frac{2\cdot2\cdot4\cdot6\cdot\cdots\cdot2k}{1\cdot3\cdot5\cdot\cdots\cdot(2k+1)} \tag{2}$$

Thus $V_4 = \int_{-r}^{r}\int_{-\sqrt{r^2-w^2}}^{\sqrt{r^2-w^2}}\int_{-\sqrt{r^2-w^2-z^2}}^{\sqrt{r^2-w^2-z^2}}\int_{-\sqrt{r^2-w^2-z^2-y^2}}^{\sqrt{r^2-w^2-z^2-y^2}} dx\,dy\,dz\,dw$

$$= 2\int_{-r}^{r}\int_{-\sqrt{r^2-w^2}}^{\sqrt{r^2-w^2}}\int_{-\sqrt{r^2-w^2-z^2}}^{\sqrt{r^2-w^2-z^2}}\sqrt{r^2-w^2-z^2-y^2}\,dy\,dz\,dw$$

$$= 2\int_{-r}^{r}\int_{-\sqrt{r^2-w^2}}^{\sqrt{r^2-w^2}}\int_{-\pi/2}^{\pi/2}(r^2-w^2-z^2)\cos^2\theta\,d\theta\,dz\,dw$$

$$= 2\left[\int_{-r}^{r}\int_{-\sqrt{r^2-w^2}}^{\sqrt{r^2-w^2}}(r^2-w^2-z^2)\,dz\,dw\right]\left[\int_{-\pi/2}^{\pi/2}\cos^2\theta\,d\theta\right]$$

$$= 2\left(\tfrac{\pi}{2}\right)\left[\int_{-r}^{r}\tfrac{4}{3}(r^2-w^2)^{3/2}\,dw\right] = \pi\left(\tfrac{4}{3}\right)\int_{-\pi/2}^{\pi/2} r^4\cos^4\phi\,d\phi = \frac{4\pi}{3}r^4\cdot\frac{1\cdot3\cdot\pi}{2\cdot4} = \frac{\pi^2r^4}{2}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. By using the substitutions $x_i = \sqrt{r^2 - x_n^2 - x_{n-1}^2 - \cdots - x_{i+1}^2}\cos\theta_i$ and then applying Formulas 1 and 2 from Problem 3, we can write

$$V_n = \int_{-r}^{r}\int_{-\sqrt{r^2-x_n^2}}^{\sqrt{r^2-x_n^2}}\cdots\int_{-\sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_3^2}}^{\sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_3^2}}\int_{-\sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_3^2-x_2^2}}^{\sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_3^2-x_2^2}} dx_1\,dx_2\cdots dx_{n-1}\,dx_n$$

$$= 2\left[\int_{-\pi/2}^{\pi/2}\cos^2\theta_2\,d\theta_2\right]\left[\int_{-\pi/2}^{\pi/2}\cos^3\theta_3\,d\theta_3\right]\cdots\left[\int_{-\pi/2}^{\pi/2}\cos^{n-1}\theta_{n-1}\,d\theta_{n-1}\right]\left[\int_{-\pi/2}^{\pi/2}\cos^n\theta_n\,d\theta_n\right]r^n$$

$$= \begin{cases} \left[2\cdot\dfrac{\pi}{2}\right]\left[\dfrac{2\cdot2}{1\cdot3}\cdot\dfrac{1\cdot3\pi}{2\cdot4}\right]\left[\dfrac{2\cdot2\cdot4}{1\cdot3\cdot5}\cdot\dfrac{1\cdot3\cdot5\pi}{2\cdot4\cdot6}\right]\cdots\left[\dfrac{2\cdot\cdots\cdot(n-2)}{1\cdot\cdots\cdot(n-1)}\cdot\dfrac{1\cdot\cdots\cdot(n-1)\,\pi}{2\cdot\cdots\cdot n}\right]r^n & n \text{ even} \\ 2\left[\dfrac{\pi}{2}\cdot\dfrac{2\cdot2}{1\cdot3}\right]\left[\dfrac{1\cdot3\pi}{2\cdot4}\cdot\dfrac{2\cdot2\cdot4}{1\cdot3\cdot5}\right]\cdots\left[\dfrac{1\cdot\cdots\cdot(n-2)\,\pi}{2\cdot\cdots\cdot(n-1)}\cdot\dfrac{2\cdot\cdots\cdot(n-1)}{1\cdot\cdots\cdot n}\right]r^n & n \text{ odd} \end{cases}$$

By canceling within each set of brackets, we find that

$$V_n = \begin{cases} \dfrac{2\pi}{2}\cdot\dfrac{2\pi}{4}\cdot\dfrac{2\pi}{6}\cdot\cdots\cdot\dfrac{2\pi}{n}r^n = \dfrac{(2\pi)^{n/2}}{2\cdot4\cdot6\cdot\cdots\cdot n}r^n = \dfrac{\pi^{n/2}}{\left(\frac{1}{2}n\right)!}r^n & n \text{ even} \\ 2\cdot\dfrac{2\pi}{3}\cdot\dfrac{2\pi}{5}\cdot\dfrac{2\pi}{7}\cdot\cdots\cdot\dfrac{2\pi}{n}r^n = \dfrac{2(2\pi)^{(n-1)/2}}{3\cdot5\cdot7\cdot\cdots\cdot n}r^n = \dfrac{2^n\left[\frac{1}{2}(n-1)\right]!\,\pi^{(n-1)/2}}{n!}r^n & n \text{ odd} \end{cases}$$

15.7 Triple Integrals in Cylindrical Coordinates

1. (a)



From Equations 1, $x = r\cos\theta = 4\cos\dfrac{\pi}{3} = 4\cdot\frac{1}{2} = 2$,

$y = r\sin\theta = 4\sin\dfrac{\pi}{3} = 4\cdot\dfrac{\sqrt{3}}{2} = 2\sqrt{3}$, $z = -2$, so the point is $(2, 2\sqrt{3}, -2)$ in rectangular coordinates.

(b)



$x = 2\cos\left(-\frac{\pi}{2}\right) = 0$, $y = 2\sin\left(-\frac{\pi}{2}\right) = -2$,

and $z = 1$, so the point is $(0, -2, 1)$ in rectangular coordinates.

2. (a)



$x = \sqrt{2}\cos\dfrac{3\pi}{4} = \sqrt{2}\left(-\frac{\sqrt{2}}{2}\right) = -1$,

$y = \sqrt{2}\sin\dfrac{3\pi}{4} = \sqrt{2}\left(\frac{\sqrt{2}}{2}\right) = 1$, and $z = 2$,

so the point is $(-1, 1, 2)$ in rectangular coordinates.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) 

$x = 1\cos 1 = \cos 1$, $y = 1\sin 1 = \sin 1$, and $z = 1$, so the point is $(\cos 1, \sin 1, 1) \approx (0.54, 0.84, 1)$ in rectangular coordinates.

3. (a) From Equations 2 we have $r^2 = (-1)^2 + 1^2 = 2$ so $r = \sqrt{2}$; $\tan\theta = \frac{1}{-1} = -1$ and the point $(-1, 1)$ is in the second quadrant of the xy-plane, so $\theta = \frac{3\pi}{4} + 2n\pi$; $z = 1$. Thus, one set of cylindrical coordinates is $\left(\sqrt{2}, \frac{3\pi}{4}, 1\right)$.

(b) $r^2 = (-2)^2 + \left(2\sqrt{3}\right)^2 = 16$ so $r = 4$; $\tan\theta = \frac{2\sqrt{3}}{-2} = -\sqrt{3}$ and the point $\left(-2, 2\sqrt{3}\right)$ is in the second quadrant of the xy-plane, so $\theta = \frac{2\pi}{3} + 2n\pi$; $z = 3$. Thus, one set of cylindrical coordinates is $\left(4, \frac{2\pi}{3}, 3\right)$.

4. (a) $r^2 = \left(-\sqrt{2}\right)^2 + \left(\sqrt{2}\right)^2 = 4$ so $r = 2$; $\tan\theta = \frac{\sqrt{2}}{-\sqrt{2}} = -1$ and the point $\left(-\sqrt{2}, \sqrt{2}\right)$ is in the second quadrant of the xy-plane, so $\theta = \frac{3\pi}{4} + 2n\pi$; $z = 1$. Thus, one set of cylindrical coordinates is $\left(2, \frac{3\pi}{4}, 1\right)$.

(b) $r^2 = 2^2 + 2^2 = 8$ so $r = \sqrt{8} = 2\sqrt{2}$; $\tan\theta = \frac{2}{2} = 1$ and the point $(2, 2)$ is in the first quadrant of the xy-plane, so $\theta = \frac{\pi}{4} + 2n\pi$; $z = 2$. Thus, one set of cylindrical coordinates is $\left(2\sqrt{2}, \frac{\pi}{4}, 2\right)$.

5. Since $r = 2$, the distance from any point to the z-axis is 2. Because θ and z may vary, the surface is a circular cylinder with radius 2 and axis the z-axis. (See Figure 4.)

Also, $x^2 + y^2 = r^2 = 4$, which we recognize as an equation of this cylinder.

6. Since $\theta = \frac{\pi}{6}$ but r and z may vary, the surface is a vertical plane including the z-axis and intersecting the xy-plane in the line $y = \frac{1}{\sqrt{3}}x$. (Here we are assuming that r can be negative; if we restrict $r \geq 0$, then we get a half-plane.)

7. Since $r^2 + z^2 = 4$ and $r^2 = x^2 + y^2$, we have $x^2 + y^2 + z^2 = 4$, a sphere centered at the origin with radius 2.

8. $r = 2\sin\theta \quad \Rightarrow \quad r^2 = 2r\sin\theta \quad \Rightarrow \quad x^2 + y^2 = 2y \quad \Leftrightarrow \quad x^2 + (y-1)^2 = 1$. z doesn't appear in the equation, so any horizontal trace in $z = k$ is the circle $x^2 + (y-1)^2 = 1$, $z = k$, which has center $(0, 1, k)$ and radius 1. Thus the surface is a circular cylinder with radius 1 and axis the vertical line $x = 0$, $y = 1$.

9. (a) Substituting $x^2 + y^2 = r^2$ and $x = r\cos\theta$, the equation $x^2 - x + y^2 + z^2 = 1$ becomes $r^2 - r\cos\theta + z^2 = 1$ or $z^2 = 1 + r\cos\theta - r^2$.

(b) Substituting $x = r\cos\theta$ and $y = r\sin\theta$, the equation $z = x^2 - y^2$ becomes $z = (r\cos\theta)^2 - (r\sin\theta)^2 = r^2(\cos^2\theta - \sin^2\theta)$ or $z = r^2\cos 2\theta$.

10. (a) The equation $2x^2 + 2y^2 - z^2 = 4$ can be written as $2(x^2 + y^2) - z^2 = 4$ which becomes $2r^2 - z^2 = 4$ or $z^2 = 2r^2 - 4$ in cylindrical coordinates.

(b) Substituting $x = r\cos\theta$ and $y = r\sin\theta$, the equation $2x - y + z = 1$ becomes $2r\cos\theta - r\sin\theta + z = 1$ or $z = 1 + r(\sin\theta - 2\cos\theta)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11.



$z = r^2 \Leftrightarrow z = x^2 + y^2$, a circular paraboloid opening upward with vertex the origin, and $z = 8 - r^2 \Leftrightarrow z = 8 - (x^2 + y^2)$, a circular paraboloid opening downward with vertex $(0, 0, 8)$. The paraboloids intersect when $r^2 = 8 - r^2 \Leftrightarrow r^2 = 4$. Thus $r^2 \le z \le 8 - r^2$ describes the solid above the paraboloid $z = x^2 + y^2$ and below the paraboloid $z = 8 - x^2 - y^2$ for $x^2 + y^2 \le 4$.

12.



$z = r = \sqrt{x^2 + y^2}$ is a cone that opens upward. Thus $r \le z \le 2$ is the region above this cone and beneath the horizontal plane $z = 2$. $0 \le \theta \le \frac{\pi}{2}$ restricts the solid to that part of this region in the first octant.

13. We can position the cylindrical shell vertically so that its axis coincides with the z-axis and its base lies in the xy-plane. If we use centimeters as the unit of measurement, then cylindrical coordinates conveniently describe the shell as $6 \le r \le 7$, $0 \le \theta \le 2\pi$, $0 \le z \le 20$.

14. In cylindrical coordinates, the equations are $z = r^2$ and $z = 5 - r^2$. The curve of intersection is $r^2 = 5 - r^2$ or $r = \sqrt{5/2}$. So we graph the surfaces in cylindrical coordinates, with $0 \le r \le \sqrt{5/2}$. In Maple, we can use the `coords=cylindrical` option in a regular `plot3d` command. In Mathematica, we can use `RevolutionPlot3D` or `ParametricPlot3D`.



15.



The region of integration is given in cylindrical coordinates by $E = \{(r, \theta, z) \mid -\pi/2 \le \theta \le \pi/2, 0 \le r \le 2, 0 \le z \le r^2\}$. This represents the solid region above quadrants I and IV of the xy-plane enclosed by the circular cylinder $r = 2$, bounded above by the circular paraboloid $z = r^2$ ($z = x^2 + y^2$), and bounded below by the xy-plane ($z = 0$).

$$\int_{-\pi/2}^{\pi/2} \int_0^2 \int_0^{r^2} r \, dz \, dr \, d\theta = \int_{-\pi/2}^{\pi/2} \int_0^2 \left[rz\right]_{z=0}^{z=r^2} dr \, d\theta = \int_{-\pi/2}^{\pi/2} \int_0^2 r^3 \, dr \, d\theta$$

$$= \int_{-\pi/2}^{\pi/2} d\theta \int_0^2 r^3 \, dr = \left[\theta\right]_{-\pi/2}^{\pi/2} \left[\tfrac{1}{4} r^4\right]_0^2$$

$$= \pi (4 - 0) = 4\pi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

16.



The region of integration is given in cylindrical coordinates by $E = \{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 2, 0 \le z \le r\}$. This represents the solid region enclosed by the circular cylinder $r = 2$, bounded above by the cone $z = r$, and bounded below by the xy-plane.

$$\int_0^2 \int_0^{2\pi} \int_0^r r\,dz\,d\theta\,dr = \int_0^2 \int_0^{2\pi} \left[rz\right]_{z=0}^{z=r} d\theta\,dr = \int_0^2 \int_0^{2\pi} r^2\,d\theta\,dr$$
$$= \int_0^2 r^2\,dr \int_0^{2\pi} d\theta = \left[\tfrac{1}{3}r^3\right]_0^2 \left[\theta\right]_0^{2\pi} = \tfrac{8}{3} \cdot 2\pi = \tfrac{16}{3}\pi$$

17. In cylindrical coordinates, E is given by $\{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 4, -5 \le z \le 4\}$. So

$$\iiint_E \sqrt{x^2 + y^2}\,dV = \int_0^{2\pi} \int_0^4 \int_{-5}^4 \sqrt{r^2}\, r\,dz\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^4 r^2\,dr \int_{-5}^4 dz$$
$$= \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{3}r^3\right]_0^4 \left[z\right]_{-5}^4 = (2\pi)\left(\tfrac{64}{3}\right)(9) = 384\pi$$

18. The paraboloid $z = x^2 + y^2 = r^2$ intersects the plane $z = 4$ in the circle $x^2 + y^2 = 4$ or $r^2 = 4 \;\Rightarrow\; r = 2$, so in cylindrical coordinates, E is given by $\left\{(r, \theta, z) \,\middle|\, 0 \le \theta \le 2\pi, 0 \le r \le 2, r^2 \le z \le 4\right\}$. Thus

$$\iiint_E z\,dV = \int_0^{2\pi} \int_0^2 \int_{r^2}^4 (z)\, r\,dz\,dr\,d\theta = \int_0^{2\pi} \int_0^2 \left[\tfrac{1}{2}rz^2\right]_{z=r^2}^{z=4} dr\,d\theta$$
$$= \int_0^{2\pi} \int_0^2 \left(8r - \tfrac{1}{2}r^5\right) dr\,d\theta = \int_0^{2\pi} d\theta \int_0^2 \left(8r - \tfrac{1}{2}r^5\right) dr = 2\pi\left[4r^2 - \tfrac{1}{12}r^6\right]_0^2$$
$$= 2\pi\left(16 - \tfrac{16}{3}\right) = \tfrac{64}{3}\pi$$

19. The paraboloid $z = 4 - x^2 - y^2 = 4 - r^2$ intersects the xy-plane in the circle $x^2 + y^2 = 4$ or $r^2 = 4 \;\Rightarrow\; r = 2$, so in cylindrical coordinates, E is given by $\left\{(r, \theta, z) \,\middle|\, 0 \le \theta \le \pi/2, 0 \le r \le 2, 0 \le z \le 4 - r^2\right\}$. Thus

$$\iiint_E (x + y + z)\,dV = \int_0^{\pi/2} \int_0^2 \int_0^{4-r^2} (r\cos\theta + r\sin\theta + z)\, r\,dz\,dr\,d\theta$$
$$= \int_0^{\pi/2} \int_0^2 \left[r^2(\cos\theta + \sin\theta)z + \tfrac{1}{2}rz^2\right]_{z=0}^{z=4-r^2} dr\,d\theta$$
$$= \int_0^{\pi/2} \int_0^2 \left[(4r^2 - r^4)(\cos\theta + \sin\theta) + \tfrac{1}{2}r(4 - r^2)^2\right] dr\,d\theta$$
$$= \int_0^{\pi/2} \left[\left(\tfrac{4}{3}r^3 - \tfrac{1}{5}r^5\right)(\cos\theta + \sin\theta) - \tfrac{1}{12}(4 - r^2)^3\right]_{r=0}^{r=2} d\theta$$
$$= \int_0^{\pi/2} \left[\tfrac{64}{15}(\cos\theta + \sin\theta) + \tfrac{16}{3}\right] d\theta = \left[\tfrac{64}{15}(\sin\theta - \cos\theta) + \tfrac{16}{3}\theta\right]_0^{\pi/2}$$
$$= \tfrac{64}{15}(1 - 0) + \tfrac{16}{3} \cdot \tfrac{\pi}{2} - \tfrac{64}{15}(0 - 1) - 0 = \tfrac{8}{3}\pi + \tfrac{128}{15}$$

20. In cylindrical coordinates E is bounded by the planes $z = 0$, $z = r\sin\theta + 4$ and the cylinders $r = 1$ and $r = 4$, so E is given by $\{(r, \theta, z) \mid 0 \le \theta \le 2\pi,\ 1 \le r \le 4,\ 0 \le z \le r\sin\theta + 4\}$. Thus

$$\iiint_E (x - y)\,dV = \int_0^{2\pi} \int_1^4 \int_0^{r\sin\theta+4} (r\cos\theta - r\sin\theta)\, r\,dz\,dr\,d\theta = \int_0^{2\pi} \int_1^4 (r^2\cos\theta - r^2\sin\theta)\left[z\right]_{z=0}^{z=r\sin\theta+4} dr\,d\theta$$
$$= \int_0^{2\pi} \int_1^4 (r^2\cos\theta - r^2\sin\theta)(r\sin\theta + 4)\,dr\,d\theta$$
$$= \int_0^{2\pi} \int_1^4 \left[r^3(\sin\theta\cos\theta - \sin^2\theta) + 4r^2(\cos\theta - \sin\theta)\right] dr\,d\theta$$
$$= \int_0^{2\pi} \left[\tfrac{1}{4}r^4(\sin\theta\cos\theta - \sin^2\theta) + \tfrac{4}{3}r^3(\cos\theta - \sin\theta)\right]_{r=1}^{r=4} d\theta$$
$$= \int_0^{2\pi} \left[\left(64 - \tfrac{1}{4}\right)(\sin\theta\cos\theta - \sin^2\theta) + \left(\tfrac{256}{3} - \tfrac{4}{3}\right)(\cos\theta - \sin\theta)\right] d\theta$$
$$= \int_0^{2\pi} \left[\tfrac{255}{4}(\sin\theta\cos\theta - \sin^2\theta) + 84(\cos\theta - \sin\theta)\right] d\theta$$
$$= \left[\tfrac{255}{4}\left(\tfrac{1}{2}\sin^2\theta - \left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right)\right) + 84\left(\sin\theta + \cos\theta\right)\right]_0^{2\pi} = \tfrac{255}{4}(-\pi) + 84(1) - 0 - 84(1) = -\tfrac{255}{4}\pi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. In cylindrical coordinates, E is bounded by the cylinder $r = 1$, the plane $z = 0$, and the cone $z = 2r$. So $E = \{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 1, 0 \le z \le 2r\}$ and

$$\begin{aligned}\iiint_E x^2\,dV &= \int_0^{2\pi}\int_0^1\int_0^{2r} r^2\cos^2\theta\, r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^1 \left[r^3\cos^2\theta\, z\right]_{z=0}^{z=2r} dr\,d\theta = \int_0^{2\pi}\int_0^1 2r^4\cos^2\theta\,dr\,d\theta \\ &= \int_0^{2\pi}\left[\tfrac{2}{5}r^5\cos^2\theta\right]_{r=0}^{r=1} d\theta = \tfrac{2}{5}\int_0^{2\pi}\cos^2\theta\,d\theta = \tfrac{2}{5}\int_0^{2\pi}\tfrac{1}{2}(1+\cos 2\theta)\,d\theta = \tfrac{1}{5}\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{2\pi} = \tfrac{2\pi}{5}\end{aligned}$$

22. In cylindrical coordinates E is the solid region within the cylinder $r = 1$ bounded above and below by the sphere $r^2 + z^2 = 4$, so $E = \left\{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 1, -\sqrt{4-r^2} \le z \le \sqrt{4-r^2}\right\}$. Thus the volume is

$$\begin{aligned}\iiint_E dV &= \int_0^{2\pi}\int_0^1\int_{-\sqrt{4-r^2}}^{\sqrt{4-r^2}} r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^1 2r\sqrt{4-r^2}\,dr\,d\theta \\ &= \int_0^{2\pi} d\theta\int_0^1 2r\sqrt{4-r^2}\,dr = 2\pi\left[-\tfrac{2}{3}(4-r^2)^{3/2}\right]_0^1 = \tfrac{4}{3}\pi(8-3^{3/2})\end{aligned}$$

23. In cylindrical coordinates, E is bounded below by the cone $z = r$ and above by the sphere $r^2 + z^2 = 2$ or $z = \sqrt{2-r^2}$. The cone and the sphere intersect when $2r^2 = 2 \quad \Rightarrow \quad r = 1$, so $E = \left\{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 1, r \le z \le \sqrt{2-r^2}\right\}$ and the volume is

$$\begin{aligned}\iiint_E dV &= \int_0^{2\pi}\int_0^1\int_r^{\sqrt{2-r^2}} r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^1 \left[rz\right]_{z=r}^{z=\sqrt{2-r^2}} dr\,d\theta = \int_0^{2\pi}\int_0^1 \left(r\sqrt{2-r^2} - r^2\right) dr\,d\theta \\ &= \int_0^{2\pi} d\theta\int_0^1 \left(r\sqrt{2-r^2} - r^2\right) dr = 2\pi\left[-\tfrac{1}{3}(2-r^2)^{3/2} - \tfrac{1}{3}r^3\right]_0^1 \\ &= 2\pi\left(-\tfrac{1}{3}\right)\left(1 + 1 - 2^{3/2}\right) = -\tfrac{2}{3}\pi\left(2 - 2\sqrt{2}\right) = \tfrac{4}{3}\pi\left(\sqrt{2}-1\right)\end{aligned}$$

24. In cylindrical coordinates, E is bounded below by the paraboloid $z = r^2$ and above by the sphere $r^2 + z^2 = 2$ or $z = \sqrt{2-r^2}$. The paraboloid and the sphere intersect when $r^2 + r^4 = 2 \quad \Rightarrow \quad (r^2+2)(r^2-1) = 0 \quad \Rightarrow \quad r = 1$, so $E = \left\{(r, \theta, z) \mid 0 \le \theta \le 2\pi,\ 0 \le r \le 1,\ r^2 \le z \le \sqrt{2-r^2}\right\}$ and the volume is

$$\begin{aligned}\iiint_E dV &= \int_0^{2\pi}\int_0^1\int_{r^2}^{\sqrt{2-r^2}} r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^1 \left[rz\right]_{z=r^2}^{z=\sqrt{2-r^2}} dr\,d\theta = \int_0^{2\pi}\int_0^1 \left(r\sqrt{2-r^2} - r^3\right) dr\,d\theta \\ &= \int_0^{2\pi} d\theta\int_0^1 \left(r\sqrt{2-r^2} - r^3\right) dr = 2\pi\left[-\tfrac{1}{3}(2-r^2)^{3/2} - \tfrac{1}{4}r^4\right]_0^1 \\ &= 2\pi\left(-\tfrac{1}{3} - \tfrac{1}{4} + \tfrac{1}{3}\cdot 2^{3/2} - 0\right) = 2\pi\left(-\tfrac{7}{12} + \tfrac{2}{3}\sqrt{2}\right) = \left(-\tfrac{7}{6} + \tfrac{4}{3}\sqrt{2}\right)\pi\end{aligned}$$

25. (a) In cylindrical coordinates, E is bounded above by the paraboloid $z = 24 - r^2$ and below by the cone $z = 2\sqrt{r^2}$ or $z = 2r$ $(r \ge 0)$. The surfaces intersect when

$24 - r^2 = 2r \quad \Rightarrow \quad r^2 + 2r - 24 = 0 \quad \Rightarrow \quad (r+6)(r-4) = 0 \quad \Rightarrow \quad r = 4$, so

$E = \left\{(r, \theta, z) \mid 2r \le z \le 24 - r^2, 0 \le r \le 4, 0 \le \theta \le 2\pi\right\}$ and the volume is

$$\begin{aligned}\iiint_E dV &= \int_0^{2\pi}\int_0^4\int_{2r}^{24-r^2} r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^4 r\left(24 - r^2 - 2r\right) dr\,d\theta = \int_0^{2\pi} d\theta\int_0^4 \left(24r - r^3 - 2r^2\right) dr \\ &= 2\pi\left[12r^2 - \tfrac{1}{4}r^4 - \tfrac{2}{3}r^3\right]_0^4 = 2\pi\left(192 - 64 - \tfrac{128}{3}\right) = \tfrac{512}{3}\pi\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b) For constant density K, $m = KV = \frac{512}{3}\pi K$ from part (a). Since the region is homogeneous and symmetric, $M_{yz} = M_{xz} = 0$ and

$$M_{xy} = \int_0^{2\pi}\int_0^4\int_{2r}^{24-r^2}(zK)\,r\,dz\,dr\,d\theta = K\int_0^{2\pi}\int_0^4 r\left[\tfrac{1}{2}z^2\right]_{z=2r}^{z=24-r^2}dr\,d\theta$$

$$= \tfrac{K}{2}\int_0^{2\pi}\int_0^4 r[(24-r^2)^2-4r^2]\,dr\,d\theta = \tfrac{K}{2}\int_0^{2\pi}d\theta\int_0^4(576r-52r^3+r^5)\,dr$$

$$= \tfrac{K}{2}(2\pi)\left[288r^2-13r^4+\tfrac{1}{6}r^6\right]_0^4 = \pi K\left(4608-3328+\tfrac{2048}{3}\right) = \tfrac{5888}{3}\pi K$$

Thus $(\overline{x},\overline{y},\overline{z}) = \left(\dfrac{M_{yz}}{m},\dfrac{M_{xz}}{m},\dfrac{M_{xy}}{m}\right) = \left(0,0,\dfrac{5888\pi K/3}{512\pi K/3}\right) = \left(0,0,\tfrac{23}{2}\right)$.

26. (a) $V = \int_{-\pi/2}^{\pi/2}\int_0^{a\cos\theta}\int_{-\sqrt{a^2-r^2}}^{\sqrt{a^2-r^2}} r\,dz\,dr\,d\theta$

$$= 4\int_0^{\pi/2}\int_0^{a\cos\theta}\int_0^{\sqrt{a^2-r^2}} r\,dz\,dr\,d\theta$$

$$= 4\int_0^{\pi/2}\int_0^{a\cos\theta} r\sqrt{a^2-r^2}\,dr\,d\theta$$

$$= -\tfrac{4}{3}\int_0^{\pi/2}\left[(a^2-r^2)^{3/2}\right]_{r=0}^{r=a\cos\theta}d\theta$$

$$= -\tfrac{4}{3}\int_0^{\pi/2}\left[(a^2-a^2\cos^2\theta)^{3/2}-a^3\right]d\theta$$

$$= -\tfrac{4}{3}\int_0^{\pi/2}\left[(a^2\sin^2\theta)^{3/2}-a^3\right]d\theta$$

$$= -\tfrac{4}{3}\int_0^{\pi/2}(a^3\sin^3\theta-a^3)\,d\theta$$

$$= -\frac{4a^3}{3}\int_0^{\pi/2}\left[\sin\theta\,(1-\cos^2\theta)-1\right]d\theta$$

$$= -\frac{4a^3}{3}\left[-\cos\theta+\tfrac{1}{3}\cos^3\theta-\theta\right]_0^{\pi/2} = -\frac{4a^3}{3}\left(-\tfrac{\pi}{2}+\tfrac{2}{3}\right) = \tfrac{2}{9}a^3(3\pi-4)$$

(b)



To plot the cylinder and the sphere on the same screen in Maple, we can use the sequence of commands

```
sphere:=plot3d(1,theta=0..2*Pi,phi=0..Pi,coords=spherical):
cylinder:=plot3d(cos(theta),theta=-Pi/2..Pi/2,z=-1..1,coords=cylindrical):
with(plots):
display3d({sphere,cylinder});
```

In Mathematica, we can use

```
sphere=SphericalPlot3D[1,{phi,0,Pi},{theta,0,2Pi}]
cylinder=ParametricPlot3D[{(Cos[theta])^2,Cos[theta]*Sin[theta],z},
                          {theta,-Pi/2,Pi/2},{z,-1,1}]
Show[sphere,cylinder]
```

27. The paraboloid $z = 4x^2+4y^2$ intersects the plane $z = a$ when $a = 4x^2+4y^2$ or $x^2+y^2 = \frac{1}{4}a$. So, in cylindrical coordinates, $E = \left\{(r,\theta,z) \mid 0 \le r \le \frac{1}{2}\sqrt{a}, 0 \le \theta \le 2\pi, 4r^2 \le z \le a\right\}$. Thus

$$m = \int_0^{2\pi}\int_0^{\sqrt{a}/2}\int_{4r^2}^{a} Kr\,dz\,dr\,d\theta = K\int_0^{2\pi}\int_0^{\sqrt{a}/2}(ar-4r^3)\,dr\,d\theta$$

$$= K\int_0^{2\pi}\left[\tfrac{1}{2}ar^2-r^4\right]_{r=0}^{r=\sqrt{a}/2}d\theta = K\int_0^{2\pi}\tfrac{1}{16}a^2\,d\theta = \tfrac{1}{8}a^2\pi K$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Since the region is homogeneous and symmetric, $M_{yz} = M_{xz} = 0$ and

$$M_{xy} = \int_0^{2\pi}\int_0^{\sqrt{a}/2}\int_{4r^2}^{a} Krz\,dz\,dr\,d\theta = K\int_0^{2\pi}\int_0^{\sqrt{a}/2}\left(\tfrac{1}{2}a^2r - 8r^5\right)dr\,d\theta$$

$$= K\int_0^{2\pi}\left[\tfrac{1}{4}a^2r^2 - \tfrac{4}{3}r^6\right]_{r=0}^{r=\sqrt{a}/2} d\theta = K\int_0^{2\pi}\tfrac{1}{24}a^3\,d\theta = \tfrac{1}{12}a^3\pi K$$

Hence $(\overline{x}, \overline{y}, \overline{z}) = \left(0, 0, \tfrac{2}{3}a\right)$.

28. Since density is proportional to the distance from the z-axis, we can say $\rho(x, y, z) = K\sqrt{x^2+y^2}$. Then

$$m = 2\int_0^{2\pi}\int_0^{a}\int_0^{\sqrt{a^2-r^2}} Kr^2\,dz\,dr\,d\theta = 2K\int_0^{2\pi}\int_0^{a} r^2\sqrt{a^2-r^2}\,dr\,d\theta$$

$$= 2K\int_0^{2\pi}\left[\tfrac{1}{8}r(2r^2-a^2)\sqrt{a^2-r^2} + \tfrac{1}{8}a^4\sin^{-1}(r/a)\right]_{r=0}^{r=a} d\theta = 2K\int_0^{2\pi}\left[\left(\tfrac{1}{8}a^4\right)\left(\tfrac{\pi}{2}\right)\right]d\theta = \tfrac{1}{4}a^4\pi^2K$$

29. The region of integration is the region above the cone $z = \sqrt{x^2+y^2}$, or $z = r$, and below the plane $z = 2$. Also, we have $-2 \le y \le 2$ with $-\sqrt{4-y^2} \le x \le \sqrt{4-y^2}$ which describes a circle of radius 2 in the xy-plane centered at $(0, 0)$. Thus,

$$\int_{-2}^{2}\int_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}}\int_{\sqrt{x^2+y^2}}^{2} xz\,dz\,dx\,dy = \int_0^{2\pi}\int_0^{2}\int_r^{2}(r\cos\theta)\,z\,r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^{2}\int_r^{2} r^2(\cos\theta)\,z\,dz\,dr\,d\theta$$

$$= \int_0^{2\pi}\int_0^{2} r^2(\cos\theta)\left[\tfrac{1}{2}z^2\right]_{z=r}^{z=2} dr\,d\theta = \tfrac{1}{2}\int_0^{2\pi}\int_0^{2} r^2(\cos\theta)\left(4-r^2\right)dr\,d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi}\cos\theta\,d\theta\int_0^{2}\left(4r^2-r^4\right)dr = \tfrac{1}{2}\left[\sin\theta\right]_0^{2\pi}\left[\tfrac{4}{3}r^3 - \tfrac{1}{5}r^5\right]_0^{2} = 0$$

30. The region of integration is the region above the plane $z = 0$ and below the paraboloid $z = 9 - x^2 - y^2$. Also, we have $-3 \le x \le 3$ with $0 \le y \le \sqrt{9-x^2}$ which describes the upper half of a circle of radius 3 in the xy-plane centered at $(0, 0)$. Thus,

$$\int_{-3}^{3}\int_0^{\sqrt{9-x^2}}\int_0^{9-x^2-y^2}\sqrt{x^2+y^2}\,dz\,dy\,dx = \int_0^{\pi}\int_0^{3}\int_0^{9-r^2}\sqrt{r^2}\,r\,dz\,dr\,d\theta = \int_0^{\pi}\int_0^{3}\int_0^{9-r^2} r^2\,dz\,dr\,d\theta$$

$$= \int_0^{\pi}\int_0^{3} r^2\left(9-r^2\right)dr\,d\theta = \int_0^{\pi}d\theta\int_0^{3}\left(9r^2-r^4\right)dr$$

$$= \left[\theta\right]_0^{\pi}\left[3r^3 - \tfrac{1}{5}r^5\right]_0^{3} = \pi\left(81 - \tfrac{243}{5}\right) = \tfrac{162}{5}\pi$$

31. (a) The mountain comprises a solid conical region C. The work done in lifting a small volume of material ΔV with density $g(P)$ to a height $h(P)$ above sea level is $h(P)g(P)\,\Delta V$. Summing over the whole mountain we get $W = \iiint_C h(P)g(P)\,dV$.

(b) Here C is a solid right circular cone with radius $R = 62{,}000$ ft, height $H = 12{,}400$ ft, and density $g(P) = 200$ lb/ft^3 at all points P in C. We use cylindrical coordinates:

$$W = \int_0^{2\pi}\int_0^{H}\int_0^{R(1-z/H)} z\cdot 200r\,dr\,dz\,d\theta = 2\pi\int_0^{H} 200z\left[\tfrac{1}{2}r^2\right]_{r=0}^{r=R(1-z/H)} dz$$

$$= 400\pi\int_0^{H} z\,\frac{R^2}{2}\left(1-\frac{z}{H}\right)^2 dz = 200\pi R^2\int_0^{H}\left(z - \frac{2z^2}{H} + \frac{z^3}{H^2}\right)dz$$

$$= 200\pi R^2\left[\frac{z^2}{2} - \frac{2z^3}{3H} + \frac{z^4}{4H^2}\right]_0^{H} = 200\pi R^2\left(\frac{H^2}{2} - \frac{2H^2}{3} + \frac{H^2}{4}\right)$$

$$= \tfrac{50}{3}\pi R^2H^2 = \tfrac{50}{3}\pi(62{,}000)^2(12{,}400)^2 \approx 3.1\times 10^{19}\text{ ft-lb}$$



$$\frac{r}{R} = \frac{H-z}{H} = 1 - \frac{z}{H}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

DISCOVERY PROJECT The Intersection of Three Cylinders

1. The three cylinders in the illustration in the text can be visualized as representing the surfaces $x^2 + y^2 = 1$, $x^2 + z^2 = 1$, and $y^2 + z^2 = 1$. Then we sketch the solid of intersection with the coordinate axes and equations indicated. To be more precise, we start by finding the bounding curves of the solid (shown in the first graph below) enclosed by the two cylinders $x^2 + z^2 = 1$ and $y^2 + z^2 = 1$: $x = \pm y = \pm\sqrt{1 - z^2}$ are the symmetric



equations, and these can be expressed parametrically as $x = s$, $y = \pm s$, $z = \pm\sqrt{1 - s^2}$, $-1 \le s \le 1$. Now the cylinder $x^2 + y^2 = 1$ intersects these curves at the eight points $\left(\pm\frac{1}{\sqrt{2}}, \pm\frac{1}{\sqrt{2}}, \pm\frac{1}{\sqrt{2}}\right)$. The resulting solid has twelve curved faces bounded by "edges" which are arcs of circles, as shown in the third diagram. Each cylinder defines four of the twelve faces.



2. To find the volume, we split the solid into sixteen congruent pieces, one of which lies in the part of the first octant with $0 \le \theta \le \frac{\pi}{4}$. (Naturally, we use cylindrical coordinates!) This piece is described by $\{(r, \theta, z) \mid 0 \le r \le 1, 0 \le \theta \le \frac{\pi}{4}, 0 \le z \le \sqrt{1 - x^2}\}$, and so, substituting $x = r\cos\theta$, the volume of the entire solid is



$$V = 16\int_0^{\pi/4}\int_0^1\int_0^{\sqrt{1-x^2}} r\,dz\,dr\,d\theta$$

$$= 16\int_0^{\pi/4}\int_0^1 r\sqrt{1 - r^2\cos^2\theta}\,dr\,d\theta$$

$$= 16 - 8\sqrt{2} \approx 4.6863$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. To graph the edges of the solid, we use parametrized curves similar to those found in Problem 1 for the intersection of two cylinders. We must restrict the parameter intervals so that each arc extends exactly to the desired vertex. One possible set of parametric equations (with all sign choices allowed) is

$x = r,\ y = \pm r,\ z = \pm\sqrt{1 - r^2},\ -\frac{1}{\sqrt{2}} \le r \le \frac{1}{\sqrt{2}}$;

$x = \pm s,\ y = \pm\sqrt{1 - s^2},\ z = s,\ -\frac{1}{\sqrt{2}} \le s \le \frac{1}{\sqrt{2}}$;

$x = \pm\sqrt{1 - t^2},\ y = t,\ z = \pm t,\ -\frac{1}{\sqrt{2}} \le t \le \frac{1}{\sqrt{2}}$.



4. Let the three cylinders be $x^2 + y^2 = a^2$, $x^2 + z^2 = 1$, and $y^2 + z^2 = 1$.

If $a < 1$, then the four faces defined by the cylinder $x^2 + y^2 = 1$ in Problem 1 collapse into a single face, as in the first graph. If $1 < a < \sqrt{2}$, then each pair of vertically opposed faces, defined by one of the other two cylinders, collapse into a single face, as in the second graph. If $a \ge \sqrt{2}$, then the vertical cylinder encloses the solid of intersection of the other two cylinders completely, so the solid of intersection coincides with the solid of intersection of the two cylinders $x^2 + z^2 = 1$ and $y^2 + z^2 = 1$, as illustrated in Problem 1.

If we were to vary b or c instead of a, we would get solids with the same shape, but differently oriented.



$a = 0.95,\ b = c = 1$



$a = 1.1,\ b = c = 1$

5. If $a < 1$, the solid looks similar to the first graph in Problem 4. As in Problem 2, we split the solid into sixteen congruent pieces, one of which can be described as the solid above the polar region $\{(r, \theta) \mid 0 \le r \le a,\ 0 \le \theta \le \frac{\pi}{4}\}$ in the xy-plane and below the surface $z = \sqrt{1 - x^2} = \sqrt{1 - r^2\cos^2\theta}$. Thus, the total volume is $V = 16\int_0^{\pi/4}\int_0^a \sqrt{1 - r^2\cos^2\theta}\, r\, dr\, d\theta$.

If $a > 1$ and $a < \sqrt{2}$, we have a solid similar to the second graph in Problem 4. Its intersection with the xy-plane is graphed at the right. Again we split the solid into sixteen congruent pieces, one of which is the solid above the region shown in the second figure and below the surface $z = \sqrt{1 - x^2} = \sqrt{1 - r^2\cos^2\theta}$.





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

We split the region of integration where the outside boundary changes from the vertical line $x = 1$ to the circle $x^2 + y^2 = a^2$ or $r = a$. R_1 is a right triangle, so $\cos\theta = \frac{1}{a}$. Thus, the boundary between R_1 and R_2 is $\theta = \cos^{-1}\left(\frac{1}{a}\right)$ in polar coordinates, or $y = \sqrt{a^2 - 1}\,x$ in rectangular coordinates. Using rectangular coordinates for the region R_1 and polar coordinates for R_2, we find the total volume of the solid to be

$$V = 16\left[\int_0^1 \int_0^{\sqrt{a^2-1}\,x} \sqrt{1-x^2}\,dy\,dx + \int_{\cos^{-1}(1/a)}^{\pi/4} \int_0^a \sqrt{1 - r^2\cos^2\theta}\,r\,dr\,d\theta\right]$$

If $a \geq \sqrt{2}$, the cylinder $x^2 + y^2 = 1$ completely encloses the intersection of the other two cylinders, so the solid of intersection of the three cylinders coincides with the intersection of $x^2 + z^2 = 1$ and $y^2 + z^2 = 1$ as illustrated in Exercise 15.5.24. Its volume is $V = 16\int_0^1 \int_0^x \sqrt{1-x^2}\,dy\,dx$.

15.8 Triple Integrals in Spherical Coordinates

1. (a)



From Equations 1, $x = \rho\sin\phi\cos\theta = 6\sin\frac{\pi}{6}\cos\frac{\pi}{3} = 6\cdot\frac{1}{2}\cdot\frac{1}{2} = \frac{3}{2}$,

$y = \rho\sin\phi\sin\theta = 6\sin\frac{\pi}{6}\sin\frac{\pi}{3} = 6\cdot\frac{1}{2}\cdot\frac{\sqrt{3}}{2} = \frac{3\sqrt{3}}{2}$, and

$z = \rho\cos\phi = 6\cos\frac{\pi}{6} = 6\cdot\frac{\sqrt{3}}{2} = 3\sqrt{3}$, so the point is $\left(\frac{3}{2}, \frac{3\sqrt{3}}{2}, 3\sqrt{3}\right)$ in rectangular coordinates.

(b)



$x = 3\sin\frac{3\pi}{4}\cos\frac{\pi}{2} = 3\cdot\frac{\sqrt{2}}{2}\cdot 0 = 0$,

$y = 3\sin\frac{3\pi}{4}\sin\frac{\pi}{2} = 3\cdot\frac{\sqrt{2}}{2}\cdot 1 = \frac{3\sqrt{2}}{2}$, and

$z = 3\cos\frac{3\pi}{4} = 3\left(-\frac{\sqrt{2}}{2}\right) = -\frac{3\sqrt{2}}{2}$, so the point is $\left(0, \frac{3\sqrt{2}}{2}, -\frac{3\sqrt{2}}{2}\right)$ in rectangular coordinates.

2. (a)



$x = 2\sin\frac{\pi}{2}\cos\frac{\pi}{2} = 2\cdot 1\cdot 0 = 0$, $y = 2\sin\frac{\pi}{2}\sin\frac{\pi}{2} = 2\cdot 1\cdot 1 = 2$,

$z = 2\cos\frac{\pi}{2} = 2\cdot 0 = 0$ so the point is $(0, 2, 0)$ in rectangular coordinates.

(b)



$x = 4\sin\frac{\pi}{3}\cos\left(-\frac{\pi}{4}\right) = 4\cdot\frac{\sqrt{3}}{2}\cdot\frac{\sqrt{2}}{2} = \sqrt{6}$,

$y = 4\sin\frac{\pi}{3}\sin\left(-\frac{\pi}{4}\right) = 4\left(\frac{\sqrt{3}}{2}\right)\left(-\frac{\sqrt{2}}{2}\right) = -\sqrt{6}$,

$z = 4\cos\frac{\pi}{3} = 4\cdot\frac{1}{2} = 2$ so the point is $\left(\sqrt{6}, -\sqrt{6}, 2\right)$ in rectangular coordinates.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a) From Equations 1 and 2, $\rho = \sqrt{x^2+y^2+z^2} = \sqrt{0^2+(-2)^2+0^2} = 2$, $\cos\phi = \dfrac{z}{\rho} = \dfrac{0}{2} = 0 \;\Rightarrow\; \phi = \dfrac{\pi}{2}$, and

$\cos\theta = \dfrac{x}{\rho\sin\phi} = \dfrac{0}{2\sin(\pi/2)} = 0 \;\Rightarrow\; \theta = \dfrac{3\pi}{2}$ [since $y < 0$]. Thus spherical coordinates are $\left(2, \dfrac{3\pi}{2}, \dfrac{\pi}{2}\right)$.

(b) $\rho = \sqrt{1+1+2} = 2$, $\cos\phi = \dfrac{z}{\rho} = \dfrac{-\sqrt{2}}{2} \;\Rightarrow\; \phi = \dfrac{3\pi}{4}$, and

$\cos\theta = \dfrac{x}{\rho\sin\phi} = \dfrac{-1}{2\sin(3\pi/4)} = \dfrac{-1}{2\left(\sqrt{2}/2\right)} = -\dfrac{1}{\sqrt{2}} \;\Rightarrow\; \theta = \dfrac{3\pi}{4}$ [since $y > 0$]. Thus spherical coordinates are $\left(2, \dfrac{3\pi}{4}, \dfrac{3\pi}{4}\right)$.

4. (a) $\rho = \sqrt{x^2+y^2+z^2} = \sqrt{1+0+3} = 2$, $\cos\phi = \dfrac{z}{\rho} = \dfrac{\sqrt{3}}{2} \;\Rightarrow\; \phi = \dfrac{\pi}{6}$, and $\cos\theta = \dfrac{x}{\rho\sin\phi} = \dfrac{1}{2\sin(\pi/6)} = 1 \;\Rightarrow$

$\theta = 0$. Thus spherical coordinates are $\left(2, 0, \dfrac{\pi}{6}\right)$.

(b) $\rho = \sqrt{3+1+12} = 4$, $\cos\phi = \dfrac{z}{\rho} = \dfrac{2\sqrt{3}}{4} = \dfrac{\sqrt{3}}{2} \;\Rightarrow\; \phi = \dfrac{\pi}{6}$, and $\cos\theta = \dfrac{x}{\rho\sin\phi} = \dfrac{\sqrt{3}}{4\sin(\pi/6)} = \dfrac{\sqrt{3}}{2} \;\Rightarrow$

$\theta = \dfrac{11\pi}{6}$ [since $y < 0$]. Thus spherical coordinates are $\left(4, \dfrac{11\pi}{6}, \dfrac{\pi}{6}\right)$.

5. Since $\phi = \frac{\pi}{3}$ but ρ and θ can vary, the surface is the top half of a right circular cone with vertex at the origin and axis the positive z-axis. (See Figure 4.)

6. $\rho^2 - 3\rho + 2 = 0 \;\Rightarrow\; (\rho-1)(\rho-2) = 0 \;\Rightarrow\; \rho = 1$ or $\rho = 2$. Thus the equation represents two surfaces. In the case $\rho = 1$, the distance from any point to the origin is 1. Because θ and ϕ can vary, the surface is a sphere centered at the origin with radius 1. (See Figure 2.) Similarly, $\rho = 2$ is a sphere centered at the origin with radius 2.

Also, $\rho = 1 \;\Rightarrow\; \rho^2 = 1 \;\Rightarrow\; x^2+y^2+z^2 = 1$ which we recognize as the equation of the unit sphere, and similarly, $\rho = 2 \;\Rightarrow\; \rho^2 = 4 \;\Rightarrow\; x^2+y^2+z^2 = 4$.

7. From Equations 1 we have $z = \rho\cos\phi$, so $\rho\cos\phi = 1 \;\Leftrightarrow\; z = 1$, and the surface is the horizontal plane $z = 1$.

8. $\rho = \cos\phi \;\Rightarrow\; \rho^2 = \rho\cos\phi \;\Leftrightarrow\; x^2+y^2+z^2 = z \;\Leftrightarrow\; x^2+y^2+z^2-z+\frac{1}{4} = \frac{1}{4} \;\Leftrightarrow\; x^2+y^2+(z-\frac{1}{2})^2 = \frac{1}{4}$. Therefore, the surface is a sphere of radius $\frac{1}{2}$ centered at $\left(0, 0, \frac{1}{2}\right)$.

9. (a) From Equation 2 we have $\rho^2 = x^2+y^2+z^2$, so $x^2+y^2+z^2 = 9 \;\Leftrightarrow\; \rho^2 = 9 \;\Rightarrow\; \rho = 3$ (since $\rho \geq 0$).

(b) From Equations 1 we have $x = \rho\sin\phi\cos\theta$, $y = \rho\sin\phi\sin\theta$, and $z = \rho\cos\phi$, so the equation $x^2 - y^2 - z^2 = 1$ becomes $(\rho\sin\phi\cos\theta)^2 - (\rho\sin\phi\sin\theta)^2 - (\rho\cos\phi)^2 = 1 \;\Leftrightarrow\; (\rho^2\sin^2\phi)(\cos^2\theta - \sin^2\theta) - \rho^2\cos^2\phi = 1 \;\Leftrightarrow$ $\rho^2(\sin^2\phi\cos 2\theta - \cos^2\phi) = 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

10. (a) $x = \rho\sin\phi\cos\theta$, $y = \rho\sin\phi\sin\theta$, and $z = \rho\cos\phi$, so the equation $z = x^2 + y^2$ becomes

$\rho\cos\phi = (\rho\sin\phi\cos\theta)^2 + (\rho\sin\phi\sin\theta)^2$ or $\rho\cos\phi = \rho^2\sin^2\phi$. If $\rho \neq 0$, this becomes $\cos\phi = \rho\sin^2\phi$ or $\rho = \cos\phi\csc^2\phi$ or $\rho = \cot\phi\csc\phi$. ($\rho = 0$ corresponds to the origin which is included in the surface.)

(b) The equation $z = x^2 - y^2$ becomes $\rho\cos\phi = (\rho\sin\phi\cos\theta)^2 - (\rho\sin\phi\sin\theta)^2$ or $\rho\cos\phi = \rho^2(\sin^2\phi)(\cos^2\theta - \sin^2\theta) \Leftrightarrow \rho\cos\phi = \rho^2\sin^2\phi\cos 2\theta$. If $\rho \neq 0$, this becomes $\cos\phi = \rho\sin^2\phi\cos 2\theta$. ($\rho = 0$ corresponds to the origin which is included in the surface.)

11. $\rho \leq 1$ represents the (solid) unit ball. $0 \leq \phi \leq \frac{\pi}{6}$ restricts the solid to that portion on or above the cone $\phi = \frac{\pi}{6}$, and $0 \leq \theta \leq \pi$ further restricts the solid to that portion on or to the right of the xz-plane.



12. $1 \leq \rho \leq 2$ represents the solid region between and including the spheres of radii 1 and 2, centered at the origin. $\frac{\pi}{2} \leq \phi \leq \pi$ restricts the solid to that portion on or below the xy-plane.



13. $2 \leq \rho \leq 4$ represents the solid region between and including the spheres of radii 2 and 4, centered at the origin. $0 \leq \phi \leq \frac{\pi}{3}$ restricts the solid to that portion on or above the cone $\phi = \frac{\pi}{3}$, and $0 \leq \theta \leq \pi$ further restricts the solid to that portion on or to the right of the xz-plane.

14. $\rho \leq 2$ represents the solid sphere of radius 2 centered at the origin. Notice that $x^2 + y^2 = (\rho\sin\phi\cos\theta)^2 + (\rho\sin\phi\sin\theta)^2 = \rho^2\sin^2\phi$. Then $\rho = \csc\phi \Rightarrow \rho\sin\phi = 1 \Rightarrow \rho^2\sin^2\phi = x^2 + y^2 = 1$, so $\rho \leq \csc\phi$ restricts the solid to that portion on or inside the circular cylinder $x^2 + y^2 = 1$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $z \geq \sqrt{x^2+y^2}$ because the solid lies above the cone. Squaring both sides of this inequality gives $z^2 \geq x^2+y^2 \;\Rightarrow\; 2z^2 \geq x^2+y^2+z^2=\rho^2 \;\Rightarrow\; z^2=\rho^2\cos^2\phi \geq \frac{1}{2}\rho^2 \;\Rightarrow\; \cos^2\phi \geq \frac{1}{2}$. The cone opens upward so that the inequality is $\cos\phi \geq \frac{1}{\sqrt{2}}$, or equivalently $0 \leq \phi \leq \frac{\pi}{4}$. In spherical coordinates the sphere $z=x^2+y^2+z^2$ is $\rho\cos\phi=\rho^2 \;\Rightarrow\; \rho=\cos\phi$. $0 \leq \rho \leq \cos\phi$ because the solid lies below the sphere. The solid can therefore be described as the region in spherical coordinates satisfying $0 \leq \rho \leq \cos\phi,\ 0 \leq \phi \leq \frac{\pi}{4}$.

16. (a) The hollow ball is a spherical shell with outer radius 15 cm and inner radius 14.5 cm. If we center the ball at the origin of the coordinate system and use centimeters as the unit of measurement, then spherical coordinates conveniently describe the hollow ball as $14.5 \leq \rho \leq 15,\ 0 \leq \theta \leq 2\pi,\ 0 \leq \phi \leq \pi$.

(b) If we position the ball as in part (a), one possibility is to take the half of the ball that is above the xy-plane which is described by $14.5 \leq \rho \leq 15,\ 0 \leq \theta \leq 2\pi,\ 0 \leq \phi \leq \pi/2$.

17.



The region of integration is given in spherical coordinates by $E=\{(\rho,\theta,\phi) \mid 0 \leq \rho \leq 3,\ 0 \leq \theta \leq \pi/2,\ 0 \leq \phi \leq \pi/6\}$. This represents the solid region in the first octant bounded above by the sphere $\rho=3$ and below by the cone $\phi=\pi/6$.

$$\begin{aligned}\int_0^{\pi/6}\int_0^{\pi/2}\int_0^3 \rho^2\sin\phi\,d\rho\,d\theta\,d\phi &= \int_0^{\pi/6}\sin\phi\,d\phi\int_0^{\pi/2}d\theta\int_0^3\rho^2\,d\rho\\ &= \left[-\cos\phi\right]_0^{\pi/6}\left[\theta\right]_0^{\pi/2}\left[\tfrac{1}{3}\rho^3\right]_0^3\\ &= \left(1-\frac{\sqrt{3}}{2}\right)\left(\frac{\pi}{2}\right)(9)=\frac{9\pi}{4}\left(2-\sqrt{3}\right)\end{aligned}$$

18. The region of integration is given in spherical coordinates by $E=\{(\rho,\theta,\phi) \mid 0 \leq \rho \leq \sec\phi,\ 0 \leq \theta \leq 2\pi,\ 0 \leq \phi \leq \pi/4\}$.

$\rho=\sec\phi \;\Leftrightarrow\; \rho\cos\phi=1 \;\Leftrightarrow\; z=1$, so E is the solid region above the cone $\phi=\pi/4$ and below the plane $z=1$.



$$\begin{aligned}\int_0^{\pi/4}\int_0^{2\pi}\int_0^{\sec\phi}\rho^2\sin\phi\,d\rho\,d\theta\,d\phi &= \int_0^{\pi/4}\int_0^{2\pi}\left[\tfrac{1}{3}\rho^3\sin\phi\right]_{\rho=0}^{\rho=\sec\phi}d\theta\,d\phi\\ &= \int_0^{\pi/4}\int_0^{2\pi}\tfrac{1}{3}\sec^3\phi\sin\phi\,d\theta\,d\phi = \tfrac{1}{3}\int_0^{\pi/4}\sec^3\phi\sin\phi\,d\phi\int_0^{2\pi}d\theta\\ &= \tfrac{1}{3}\int_0^{\pi/4}\tan\phi\sec^2\phi\,d\phi\int_0^{2\pi}d\theta = \tfrac{1}{3}\left[\tfrac{1}{2}\tan^2\phi\right]_0^{\pi/4}\left[\theta\right]_0^{2\pi}\\ &= \tfrac{1}{3}\left(\tfrac{1}{2}-0\right)(2\pi)=\tfrac{\pi}{3}\end{aligned}$$

19. The solid E is most conveniently described if we use cylindrical coordinates:

$E=\{(r,\theta,z) \mid 0 \leq \theta \leq \frac{\pi}{2},\ 0 \leq r \leq 3,\ 0 \leq z \leq 2\}$. Then

$$\iiint_E f(x,y,z)\,dV=\int_0^{\pi/2}\int_0^3\int_0^2 f(r\cos\theta,r\sin\theta,z)\,r\,dz\,dr\,d\theta.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

20. The solid E is most conveniently described if we use spherical coordinates:

$E = \{(\rho, \theta, \phi) \mid 1 \le \rho \le 2,\ \frac{\pi}{2} \le \theta \le 2\pi,\ 0 \le \phi \le \frac{\pi}{2}\}$. Then

$\iiint_E f(x, y, z)\, dV = \int_0^{\pi/2} \int_{\pi/2}^{2\pi} \int_1^2 f(\rho \sin\phi \cos\theta, \rho \sin\phi \sin\theta, \rho \cos\phi)\, \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi.$

21. In spherical coordinates, B is represented by $\{(\rho, \theta, \phi) \mid 0 \le \rho \le 5, 0 \le \theta \le 2\pi, 0 \le \phi \le \pi\}$. Thus

$$\begin{aligned}\iiint_B (x^2 + y^2 + z^2)^2\, dV &= \int_0^{\pi} \int_0^{2\pi} \int_0^5 (\rho^2)^2 \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi = \int_0^{\pi} \sin\phi\, d\phi \int_0^{2\pi} d\theta \int_0^5 \rho^6\, d\rho \\ &= \left[-\cos\phi\right]_0^{\pi} \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{7}\rho^7\right]_0^5 = (2)(2\pi)\left(\tfrac{78{,}125}{7}\right) \\ &= \tfrac{312{,}500}{7}\pi \approx 140{,}249.7\end{aligned}$$

22. In spherical coordinates, E is represented by $\{(\rho, \theta, \phi) \mid 0 \le \rho \le 1,\ 0 \le \theta \le 2\pi,\ 0 \le \phi \le \frac{\pi}{3}\}$. Thus

$$\begin{aligned}\iiint_E y^2 z^2\, dV &= \int_0^{\pi/3} \int_0^{2\pi} \int_0^1 (\rho \sin\phi \sin\theta)^2 (\rho \cos\phi)^2\, \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi \\ &= \int_0^{\pi/3} \sin^3\phi \cos^2\phi\, d\phi \int_0^{2\pi} \sin^2\theta\, d\theta \int_0^1 \rho^6\, d\rho \\ &= \int_0^{\pi/3} (1 - \cos^2\phi) \cos^2\phi \sin\phi\, d\phi \int_0^{2\pi} \tfrac{1}{2}(1 - \cos 2\theta)\, d\theta \int_0^1 \rho^6\, d\rho \\ &= \left[\tfrac{1}{5}\cos^5\phi - \tfrac{1}{3}\cos^3\phi\right]_0^{\pi/3} \left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi} \left[\tfrac{1}{7}\rho^7\right]_0^1 \\ &= \left[\tfrac{1}{5}\left(\tfrac{1}{2}\right)^5 - \tfrac{1}{3}\left(\tfrac{1}{2}\right)^3 - \tfrac{1}{5} + \tfrac{1}{3}\right](\pi - 0)\left(\tfrac{1}{7} - 0\right) = \tfrac{47}{480} \cdot \pi \cdot \tfrac{1}{7} = \tfrac{47}{3360}\pi\end{aligned}$$

23. In spherical coordinates, E is represented by $\{(\rho, \theta, \phi) \mid 2 \le \rho \le 3, 0 \le \theta \le 2\pi, 0 \le \phi \le \pi\}$ and

$x^2 + y^2 = \rho^2 \sin^2\phi \cos^2\theta + \rho^2 \sin^2\phi \sin^2\theta = \rho^2 \sin^2\phi\,(\cos^2\theta + \sin^2\theta) = \rho^2 \sin^2\phi$. Thus

$$\begin{aligned}\iiint_E (x^2 + y^2)\, dV &= \int_0^{\pi} \int_0^{2\pi} \int_2^3 (\rho^2 \sin^2\phi)\, \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi = \int_0^{\pi} \sin^3\phi\, d\phi \int_0^{2\pi} d\theta \int_2^3 \rho^4\, d\rho \\ &= \int_0^{\pi} (1 - \cos^2\phi)\, \sin\phi\, d\phi \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{5}\rho^5\right]_2^3 = \left[-\cos\phi + \tfrac{1}{3}\cos^3\phi\right]_0^{\pi} (2\pi) \cdot \tfrac{1}{5}(243 - 32) \\ &= \left(1 - \tfrac{1}{3} + 1 - \tfrac{1}{3}\right)(2\pi)\left(\tfrac{211}{5}\right) = \tfrac{1688\pi}{15}\end{aligned}$$

24. In spherical coordinates, E is represented by $\{(\rho, \theta, \phi) \mid 0 \le \rho \le 3, 0 \le \theta \le \pi, 0 \le \phi \le \pi\}$. Thus

$$\begin{aligned}\iiint_E y^2\, dV &= \int_0^{\pi} \int_0^{\pi} \int_0^3 (\rho \sin\phi \sin\theta)^2\, \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi = \int_0^{\pi} \sin^3\phi\, d\phi \int_0^{\pi} \sin^2\theta\, d\theta \int_0^3 \rho^4\, d\rho \\ &= \int_0^{\pi} (1 - \cos^2\phi)\, \sin\phi\, d\phi \int_0^{\pi} \tfrac{1}{2}(1 - \cos 2\theta)\, d\theta \int_0^3 \rho^4\, d\rho \\ &= \left[-\cos\phi + \tfrac{1}{3}\cos^3\phi\right]_0^{\pi} \left[\tfrac{1}{2}\left(\theta - \tfrac{1}{2}\sin 2\theta\right)\right]_0^{\pi} \left[\tfrac{1}{5}\rho^5\right]_0^3 \\ &= \left(\tfrac{2}{3} + \tfrac{2}{3}\right)\left(\tfrac{1}{2}\pi\right)\left(\tfrac{1}{5}(243)\right) = \left(\tfrac{4}{3}\right)\left(\tfrac{\pi}{2}\right)\left(\tfrac{243}{5}\right) = \tfrac{162\pi}{5}\end{aligned}$$

25. In spherical coordinates, E is represented by $\{(\rho, \theta, \phi) \mid 0 \le \rho \le 1, 0 \le \theta \le \frac{\pi}{2}, 0 \le \phi \le \frac{\pi}{2}\}$. Thus

$$\begin{aligned}\iiint_E x e^{x^2+y^2+z^2}\, dV &= \int_0^{\pi/2} \int_0^{\pi/2} \int_0^1 (\rho \sin\phi \cos\theta) e^{\rho^2}\, \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi = \int_0^{\pi/2} \sin^2\phi\, d\phi \int_0^{\pi/2} \cos\theta\, d\theta \int_0^1 \rho^3 e^{\rho^2}\, d\rho \\ &= \int_0^{\pi/2} \tfrac{1}{2}(1 - \cos 2\phi)\, d\phi \int_0^{\pi/2} \cos\theta\, d\theta \left(\left[\tfrac{1}{2}\rho^2 e^{\rho^2}\right]_0^1 - \int_0^1 \rho e^{\rho^2}\, d\rho\right) \\ &\qquad \left[\text{integrate by parts with } u = \rho^2,\ dv = \rho e^{\rho^2} d\rho\right] \\ &= \left[\tfrac{1}{2}\phi - \tfrac{1}{4}\sin 2\phi\right]_0^{\pi/2} \left[\sin\theta\right]_0^{\pi/2} \left[\tfrac{1}{2}\rho^2 e^{\rho^2} - \tfrac{1}{2}e^{\rho^2}\right]_0^1 = \left(\tfrac{\pi}{4} - 0\right)(1 - 0)\left(0 + \tfrac{1}{2}\right) = \tfrac{\pi}{8}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. In spherical coordinates, the cone $z = \sqrt{x^2 + y^2}$ is equivalent to $\varphi = \pi/4$ (as in Example 4) and E is represented by $\{(\rho, \theta, \phi) \mid 1 \le \rho \le 2,\ 0 \le \theta \le 2\pi,\ 0 \le \phi \le \pi/4\}$. Also $\sqrt{x^2 + y^2 + z^2} = \sqrt{\rho^2} = \rho$, so

$$\iiint_E \sqrt{x^2 + y^2 + z^2}\, dV = \int_0^{\pi/4} \int_0^{2\pi} \int_1^2 \rho \cdot \rho^2 \sin\phi \, d\rho \, d\theta \, d\phi = \int_0^{\pi/4} \sin\phi \, d\phi \int_0^{2\pi} d\theta \int_1^2 \rho^3 \, d\rho$$

$$= \left[-\cos\phi\right]_0^{\pi/4} \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{4}\rho^4\right]_1^2 = \left(-\tfrac{\sqrt{2}}{2} + 1\right)(2\pi) \cdot \tfrac{1}{4}(16 - 1) = \tfrac{15}{2}\pi\left(1 - \tfrac{\sqrt{2}}{2}\right)$$

27. The solid region is given by $E = \left\{(\rho, \theta, \phi) \mid 0 \le \rho \le a, 0 \le \theta \le 2\pi, \tfrac{\pi}{6} \le \phi \le \tfrac{\pi}{3}\right\}$ and its volume is

$$V = \iiint_E dV = \int_{\pi/6}^{\pi/3} \int_0^{2\pi} \int_0^a \rho^2 \sin\phi \, d\rho \, d\theta \, d\phi = \int_{\pi/6}^{\pi/3} \sin\phi \, d\phi \int_0^{2\pi} d\theta \int_0^a \rho^2 \, d\rho$$

$$= \left[-\cos\phi\right]_{\pi/6}^{\pi/3} \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{3}\rho^3\right]_0^a = \left(-\tfrac{1}{2} + \tfrac{\sqrt{3}}{2}\right)(2\pi)\left(\tfrac{1}{3}a^3\right) = \tfrac{\sqrt{3}-1}{3}\pi a^3$$

28. If we center the ball at the origin, then the ball is given by $B = \{(\rho, \theta, \phi) \mid 0 \le \rho \le a, 0 \le \theta \le 2\pi, 0 \le \phi \le \pi\}$ and the distance from any point (x, y, z) in the ball to the center $(0, 0, 0)$ is $\sqrt{x^2 + y^2 + z^2} = \rho$. Thus the average distance is

$$\frac{1}{V(B)} \iiint_B \rho \, dV = \frac{1}{\frac{4}{3}\pi a^3} \int_0^{\pi} \int_0^{2\pi} \int_0^a \rho \cdot \rho^2 \sin\phi \, d\rho \, d\theta \, d\phi = \frac{3}{4\pi a^3} \int_0^{\pi} \sin\phi \, d\phi \int_0^{2\pi} d\theta \int_0^a \rho^3 \, d\rho$$

$$= \frac{3}{4\pi a^3} \left[-\cos\phi\right]_0^{\pi} \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{4}\rho^4\right]_0^a = \frac{3}{4\pi a^3}(2)(2\pi)\left(\tfrac{1}{4}a^4\right) = \tfrac{3}{4}a$$

29. (a) Since $\rho = 4\cos\phi$ implies $\rho^2 = 4\rho\cos\phi \iff x^2 + y^2 + z^2 = 4z \iff x^2 + y^2 + (z - 2)^2 = 4$, the equation is that of a sphere of radius 2 with center at $(0, 0, 2)$. Thus

$$V = \int_0^{2\pi} \int_0^{\pi/3} \int_0^{4\cos\phi} \rho^2 \sin\phi \, d\rho \, d\phi \, d\theta = \int_0^{2\pi} \int_0^{\pi/3} \left[\tfrac{1}{3}\rho^3\right]_{\rho=0}^{\rho=4\cos\phi} \sin\phi \, d\phi \, d\theta = \int_0^{2\pi} \int_0^{\pi/3} \left(\tfrac{64}{3}\cos^3\phi\right) \sin\phi \, d\phi \, d\theta$$

$$= \int_0^{2\pi} \left[-\tfrac{16}{3}\cos^4\phi\right]_{\phi=0}^{\phi=\pi/3} d\theta = \int_0^{2\pi} -\tfrac{16}{3}\left(\tfrac{1}{16} - 1\right) d\theta = 5\theta\Big]_0^{2\pi} = 10\pi$$

(b) By the symmetry of the problem $M_{yz} = M_{xz} = 0$. Then

$$M_{xy} = \int_0^{2\pi} \int_0^{\pi/3} \int_0^{4\cos\phi} \rho^3 \cos\phi \sin\phi \, d\rho \, d\phi \, d\theta = \int_0^{2\pi} \int_0^{\pi/3} \cos\phi \sin\phi \left(64\cos^4\phi\right) d\phi \, d\theta$$

$$= \int_0^{2\pi} 64 \left[-\tfrac{1}{6}\cos^6\phi\right]_{\phi=0}^{\phi=\pi/3} d\theta = \int_0^{2\pi} \tfrac{21}{2}\, d\theta = 21\pi$$

Hence $(\overline{x}, \overline{y}, \overline{z}) = (0, 0, 21\pi/(10\pi)) = (0, 0, 2.1)$.

30. In spherical coordinates, the sphere $x^2 + y^2 + z^2 = 4$ is equivalent to $\rho = 2$ and the cone $z = \sqrt{x^2 + y^2}$ is represented by $\phi = \tfrac{\pi}{4}$ (as in Example 4). Thus, the solid is given by $\left\{(\rho, \theta, \phi) \mid 0 \le \rho \le 2,\ 0 \le \theta \le 2\pi,\ \tfrac{\pi}{4} \le \phi \le \tfrac{\pi}{2}\right\}$ and

$$V = \int_{\pi/4}^{\pi/2} \int_0^{2\pi} \int_0^2 \rho^2 \sin\phi \, d\rho \, d\theta \, d\phi = \int_{\pi/4}^{\pi/2} \sin\phi \, d\phi \int_0^{2\pi} d\theta \int_0^2 \rho^2 \, d\rho$$

$$= \left[-\cos\phi\right]_{\pi/4}^{\pi/2} \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{3}\rho^3\right]_0^2 = \left(\tfrac{\sqrt{2}}{2}\right)(2\pi)\left(\tfrac{8}{3}\right) = \tfrac{8\sqrt{2}\,\pi}{3}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

31. (a) By the symmetry of the region, $M_{yz} = 0$ and $M_{xz} = 0$. Assuming constant density K, $m = \iiint_E K\,dV = K\iiint_E dV = \frac{\pi}{8}K$ (from Example 4). Then

$$M_{xy} = \iiint_E z\,K\,dV = K\int_0^{2\pi}\int_0^{\pi/4}\int_0^{\cos\phi}(\rho\cos\phi)\,\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = K\int_0^{2\pi}\int_0^{\pi/4}\sin\phi\cos\phi\left[\tfrac{1}{4}\rho^4\right]_{\rho=0}^{\rho=\cos\phi}d\phi\,d\theta$$

$$= \tfrac{1}{4}K\int_0^{2\pi}\int_0^{\pi/4}\sin\phi\cos\phi\left(\cos^4\phi\right)d\phi\,d\theta = \tfrac{1}{4}K\int_0^{2\pi}d\theta\int_0^{\pi/4}\cos^5\phi\sin\phi\,d\phi$$

$$= \tfrac{1}{4}K\left[\theta\right]_0^{2\pi}\left[-\tfrac{1}{6}\cos^6\phi\right]_0^{\pi/4} = \tfrac{1}{4}K(2\pi)\left(-\tfrac{1}{6}\right)\left[\left(\frac{\sqrt{2}}{2}\right)^6 - 1\right] = -\frac{\pi}{12}K\left(-\frac{7}{8}\right) = \frac{7\pi}{96}K$$

Thus the centroid is $(\overline{x},\overline{y},\overline{z}) = \left(\dfrac{M_{yz}}{m},\dfrac{M_{xz}}{m},\dfrac{M_{xy}}{m}\right) = \left(0,0,\dfrac{7\pi K/96}{\pi K/8}\right) = \left(0,0,\frac{7}{12}\right)$.

(b) As in Exercise 23, $x^2+y^2 = \rho^2\sin^2\phi$ and

$$I_z = \iiint_E (x^2+y^2)\,K\,dV = K\int_0^{2\pi}\int_0^{\pi/4}\int_0^{\cos\phi}(\rho^2\sin^2\phi)\,\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = K\int_0^{2\pi}\int_0^{\pi/4}\sin^3\phi\left[\tfrac{1}{5}\rho^5\right]_{\rho=0}^{\rho=\cos\phi}d\phi\,d\theta$$

$$= \tfrac{1}{5}K\int_0^{2\pi}\int_0^{\pi/4}\sin^3\phi\cos^5\phi\,d\phi\,d\theta = \tfrac{1}{5}K\int_0^{2\pi}d\theta\int_0^{\pi/4}\cos^5\phi\left(1-\cos^2\phi\right)\sin\phi\,d\phi$$

$$= \tfrac{1}{5}K\left[\theta\right]_0^{2\pi}\left[-\tfrac{1}{6}\cos^6\phi + \tfrac{1}{8}\cos^8\phi\right]_0^{\pi/4}$$

$$= \tfrac{1}{5}K(2\pi)\left[-\tfrac{1}{6}\left(\frac{\sqrt{2}}{2}\right)^6 + \tfrac{1}{8}\left(\frac{\sqrt{2}}{2}\right)^8 + \tfrac{1}{6} - \tfrac{1}{8}\right] = \tfrac{2\pi}{5}K\left(\tfrac{11}{384}\right) = \tfrac{11\pi}{960}K$$

32. (a) Placing the center of the base at $(0,0,0)$, $\rho(x,y,z) = K\sqrt{x^2+y^2+z^2}$ is the density function. So

$$m = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a K\rho^3\sin\phi\,d\rho\,d\phi\,d\theta = K\int_0^{2\pi}d\theta\int_0^{\pi/2}\sin\phi\,d\phi\int_0^a\rho^3\,d\rho$$

$$= K\left[\theta\right]_0^{2\pi}\left[-\cos\phi\right]_0^{\pi/2}\left[\tfrac{1}{4}\rho^4\right]_0^a = K(2\pi)(1)\left(\tfrac{1}{4}a^4\right) = \tfrac{1}{2}\pi Ka^4$$

(b) By the symmetry of the problem $M_{yz} = M_{xz} = 0$. Then

$$M_{xy} = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a K\rho^4\sin\phi\cos\phi\,d\rho\,d\phi\,d\theta = K\int_0^{2\pi}d\theta\int_0^{\pi/2}\sin\phi\cos\phi\,d\phi\int_0^a\rho^4\,d\rho$$

$$= K\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{2}\sin^2\phi\right]_0^{\pi/2}\left[\tfrac{1}{5}\rho^5\right]_0^a = K(2\pi)\left(\tfrac{1}{2}\right)\left(\tfrac{1}{5}a^5\right) = \tfrac{1}{5}\pi Ka^5$$

Hence $(\overline{x},\overline{y},\overline{z}) = \left(0,0,\frac{2}{5}a\right)$.

(c) $I_z = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a (K\rho^3\sin\phi)(\rho^2\sin^2\phi)\,d\rho\,d\phi\,d\theta = K\int_0^{2\pi}d\theta\int_0^{\pi/2}\sin^3\phi\,d\phi\int_0^a\rho^5\,d\rho$

$$= K\left[\theta\right]_0^{2\pi}\left[-\cos\phi + \tfrac{1}{3}\cos^3\phi\right]_0^{\pi/2}\left[\tfrac{1}{6}\rho^6\right]_0^a = K(2\pi)\left(\tfrac{2}{3}\right)\left(\tfrac{1}{6}a^6\right) = \tfrac{2}{9}\pi Ka^6$$

33. (a) The density function is $\rho(x,y,z) = K$, a constant, and by the symmetry of the problem $M_{xz} = M_{yz} = 0$. Then $M_{xy} = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a K\rho^3\sin\phi\,\cos\phi\,d\rho\,d\phi\,d\theta = \frac{1}{2}\pi Ka^4\int_0^{\pi/2}\sin\phi\,\cos\phi\,d\phi = \frac{1}{8}\pi Ka^4$. But the mass is K(volume of the hemisphere) $= \frac{2}{3}\pi Ka^3$, so the centroid is $\left(0,0,\frac{3}{8}a\right)$.

(b) Place the center of the base at $(0,0,0)$; the density function is $\rho(x,y,z) = K$. By symmetry, the moments of inertia about any two such diameters will be equal, so we just need to find I_x:

$$I_x = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a (K\rho^2\sin\phi)\,\rho^2\left(\sin^2\phi\,\sin^2\theta + \cos^2\phi\right)d\rho\,d\phi\,d\theta$$

$$= K\int_0^{2\pi}\int_0^{\pi/2}(\sin^3\phi\,\sin^2\theta + \sin\phi\,\cos^2\phi)\left(\tfrac{1}{5}a^5\right)d\phi\,d\theta$$

$$= \tfrac{1}{5}Ka^5\int_0^{2\pi}\left[\sin^2\theta\left(-\cos\phi + \tfrac{1}{3}\cos^3\phi\right) + \left(-\tfrac{1}{3}\cos^3\phi\right)\right]_{\phi=0}^{\phi=\pi/2}d\theta = \tfrac{1}{5}Ka^5\int_0^{2\pi}\left[\tfrac{2}{3}\sin^2\theta + \tfrac{1}{3}\right]d\theta$$

$$= \tfrac{1}{5}Ka^5\left[\tfrac{2}{3}\left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right) + \tfrac{1}{3}\theta\right]_0^{2\pi} = \tfrac{1}{5}Ka^5\left[\tfrac{2}{3}(\pi - 0) + \tfrac{1}{3}(2\pi - 0)\right] = \tfrac{4}{15}Ka^5\pi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. Place the center of the base at $(0,0,0)$, then the density is $\rho(x,y,z) = Kz$, K a constant. Then

$m = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a (K\rho\cos\phi)\,\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = 2\pi K\int_0^{\pi/2}\cos\phi\sin\phi\cdot\frac{1}{4}a^4\,d\phi = \frac{1}{2}\pi Ka^4\left[-\frac{1}{4}\cos 2\phi\right]_0^{\pi/2} = \frac{\pi}{4}Ka^4.$

By the symmetry of the problem $M_{xz} = M_{yz} = 0$, and

$M_{xy} = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a K\rho^4\cos^2\phi\sin\phi\,d\rho\,d\phi\,d\theta = \frac{2}{5}\pi Ka^5\int_0^{\pi/2}\cos^2\phi\sin\phi\,d\phi = \frac{2}{5}\pi Ka^5\left[-\frac{1}{3}\cos^3\theta\right]_0^{\pi/2} = \frac{2}{15}\pi Ka^5.$

Hence $(\overline{x},\overline{y},\overline{z}) = \left(0,0,\frac{8}{15}a\right)$.

35. In spherical coordinates $z = \sqrt{x^2+y^2}$ becomes $\phi = \frac{\pi}{4}$ (as in Example 4). Then

$V = \int_0^{2\pi}\int_0^{\pi/4}\int_0^1 \rho^2\sin\phi\,d\rho\,d\phi\,d\theta = \int_0^{2\pi}d\theta\int_0^{\pi/4}\sin\phi\,d\phi\int_0^1\rho^2\,d\rho = 2\pi\left(-\frac{\sqrt{2}}{2}+1\right)\left(\frac{1}{3}\right) = \frac{1}{3}\pi\left(2-\sqrt{2}\right),$

$M_{xy} = \int_0^{2\pi}\int_0^{\pi/4}\int_0^1 \rho^3\sin\phi\cos\phi\,d\rho\,d\phi\,d\theta = 2\pi\left[-\frac{1}{4}\cos 2\phi\right]_0^{\pi/4}\left(\frac{1}{4}\right) = \frac{\pi}{8}$ and by symmetry $M_{yz} = M_{xz} = 0$.

Hence $(\overline{x},\overline{y},\overline{z}) = \left(0,0,\dfrac{3}{8\left(2-\sqrt{2}\right)}\right)$.

36. Place the center of the sphere at $(0,0,0)$, let the diameter of intersection be along the z-axis, one of the planes be the xz-plane and the other be the plane whose angle with the xz-plane is $\theta = \frac{\pi}{6}$. Then in spherical coordinates the volume is given by

$V = \int_0^{\pi/6}\int_0^{\pi}\int_0^a \rho^2\sin\phi\,d\rho\,d\phi\,d\theta = \int_0^{\pi/6}d\theta\int_0^{\pi}\sin\phi\,d\phi\int_0^a\rho^2\,d\rho = \frac{\pi}{6}(2)\left(\frac{1}{3}a^3\right) = \frac{1}{9}\pi a^3.$

37. (a) If we orient the cylinder so that its axis is the z-axis and its base lies in the xy-plane, then the cylinder is described, in cylindrical coordinates, by $E = \{(r,\theta,z) \mid 0 \le r \le a,\ 0 \le \theta \le 2\pi,\ 0 \le z \le h\}$. Assuming constant density K, the moment of inertia about its axis (the z-axis) is

$$I_z = \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^a\int_0^h K(r^2)\,r\,dz\,dr\,d\theta = K\int_0^{2\pi}d\theta\int_0^a r^3\,dr\int_0^h dz$$
$$= K\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^a\left[z\right]_0^h = K(2\pi)\left(\tfrac{1}{4}a^4\right)(h) = \tfrac{1}{2}\pi Ka^4h$$

(b) By symmetry, the moments of inertia about any two diameters of the base will be equal, and one of the diameters lies on the x-axis, so we compute:

$$I_x = \iiint_E (y^2+z^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^a\int_0^h K(r^2\sin^2\theta + z^2)\,r\,dz\,dr\,d\theta$$
$$= K\int_0^{2\pi}\int_0^a\int_0^h r^3\sin^2\theta\,dz\,dr\,d\theta + K\int_0^{2\pi}\int_0^a\int_0^h rz^2\,dz\,dr\,d\theta$$
$$= K\int_0^{2\pi}\sin^2\theta\,d\theta\int_0^a r^3\,dr\int_0^h dz + K\int_0^{2\pi}d\theta\int_0^a r\,dr\int_0^h z^2\,dz$$
$$= K\left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^a\left[z\right]_0^h + K\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{2}r^2\right]_0^a\left[\tfrac{1}{3}z^3\right]_0^h$$
$$= K(\pi)\left(\tfrac{1}{4}a^4\right)(h) + K(2\pi)\left(\tfrac{1}{2}a^2\right)\left(\tfrac{1}{3}h^3\right) = \tfrac{1}{12}\pi Ka^2h(3a^2+4h^2)$$

38.



Orient the cone so that its axis is the z-axis and its base lies in the xy-plane, as shown in the figure. (Then the z-axis is the axis of the cone and the x-axis contains a diameter of the base.) A right circular cone with axis the z-axis and vertex at the origin has equation $z^2 = c^2(x^2+y^2)$. Here we have the bottom frustum, shifted upward h units, and with $c^2 = h^2/a^2$ so that the cone includes the point $(a,0,0)$. Thus an equation of the cone in rectangular coordinates is $z = h - \frac{h}{a}\sqrt{x^2+y^2}$, $0 \le z \le h$. In cylindrical

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

coordinates, the cone is described by

$$E = \{(r,\theta,z) \mid 0 \le r \le a,\ 0 \le \theta \le 2\pi,\ 0 \le z \le h\left(1 - \tfrac{1}{a}r\right)\}$$

(a) Assuming constant density K, the moment of inertia about its axis (the z-axis) is

$$\begin{aligned} I_z &= \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^a \int_0^{h(1-r/a)} K(r^2)\,r\,dz\,dr\,d\theta \\ &= K\int_0^{2\pi}\int_0^a \left[r^3 z\right]_{z=0}^{z=h(1-r/a)}\,dr\,d\theta = K\int_0^{2\pi}\int_0^a r^3 h\left(1-\tfrac{1}{a}r\right)\,dr\,d\theta \\ &= Kh\int_0^{2\pi} d\theta \int_0^a \left(r^3 - \tfrac{1}{a}r^4\right)\,dr = Kh\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4 - \tfrac{1}{5a}r^5\right]_0^a \\ &= Kh\,(2\pi)\left(\tfrac{1}{4}a^4 - \tfrac{1}{5}a^4\right) = \tfrac{1}{10}\pi K a^4 h \end{aligned}$$

(b) By symmetry, the moments of intertia about any two diameters of the base will be equal, and one of the diameters lies on the x-axis, so we compute:

$$\begin{aligned} I_x &= \iiint_E (y^2+z^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^a \int_0^{h(1-r/a)} K(r^2\sin^2\theta + z^2)\,r\,dz\,dr\,d\theta \\ &= K\int_0^{2\pi}\int_0^a \left[(r^3\sin^2\theta)z + \tfrac{1}{3}rz^3\right]_{z=0}^{z=h(1-r/a)}\,dr\,d\theta \\ &= K\int_0^{2\pi}\int_0^a \left[(r^3\sin^2\theta)\left(h\left(1-\tfrac{1}{a}r\right)\right) + \tfrac{1}{3}r\left(h\left(1-\tfrac{1}{a}r\right)\right)^3\right]\,dr\,d\theta \\ &= Kh\int_0^{2\pi}\int_0^a (r^3\sin^2\theta)\left(1-\tfrac{1}{a}r\right)\,dr\,d\theta + Kh^3\int_0^{2\pi}\int_0^a \tfrac{1}{3}r\left(1-\tfrac{1}{a}r\right)^3\,dr\,d\theta \\ &= Kh\int_0^{2\pi}\sin^2\theta\,d\theta\int_0^a \left(r^3 - \tfrac{1}{a}r^4\right)\,dr + \tfrac{1}{3}Kh^3\int_0^{2\pi} d\theta\int_0^a (r - \tfrac{3}{a}r^2 + \tfrac{3}{a^2}r^3 - \tfrac{1}{a^3}r^4)\,dr \\ &= Kh\left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4 - \tfrac{1}{5a}r^5\right]_0^a + \tfrac{1}{3}Kh^3\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{2}r^2 - \tfrac{1}{a}r^3 + \tfrac{3}{4a^2}r^4 - \tfrac{1}{5a^3}r^5\right]_0^a \\ &= Kh\,(\pi)\left(\tfrac{1}{4}a^4 - \tfrac{1}{5}a^4\right) + \tfrac{1}{3}Kh^3\,(2\pi)\left(\tfrac{1}{2}a^2 - a^2 + \tfrac{3}{4}a^2 - \tfrac{1}{5}a^2\right) \\ &= \pi K h\left(\tfrac{1}{20}a^4\right) + \tfrac{2}{3}\pi K h^3\left(\tfrac{1}{20}a^2\right) = \pi K a^2 h\left(\tfrac{1}{20}a^2 + \tfrac{1}{30}h^2\right) \end{aligned}$$

39. In cylindrical coordinates the paraboloid is given by $z = r^2$ and the plane by $z = 2r\sin\theta$ and the projection of the intersection onto the xy-plane is the circle $r = 2\sin\theta$. Then $\iiint_E z\,dV = \int_0^{\pi}\int_0^{2\sin\theta}\int_{r^2}^{2r\sin\theta} rz\,dz\,dr\,d\theta = \frac{5\pi}{6}$ [using a CAS].

40. (a) The region enclosed by the torus is $\{(\rho,\theta,\phi) \mid 0 \le \theta \le 2\pi, 0 \le \phi \le \pi, 0 \le \rho \le \sin\phi\}$, so its volume is

$$V = \int_0^{2\pi}\int_0^{\pi}\int_0^{\sin\phi} \rho^2\sin\phi\,d\rho\,d\phi\,d\theta = 2\pi\int_0^{\pi}\tfrac{1}{3}\sin^4\phi\,d\phi = \tfrac{2}{3}\pi\left[\tfrac{3}{8}\phi - \tfrac{1}{4}\sin 2\phi + \tfrac{1}{16}\sin 4\phi\right]_0^{\pi} = \tfrac{1}{4}\pi^2.$$

(b) In Maple, we can plot the torus using the command

```
plot3d(sin(phi),theta=0..2*Pi,
       phi=0..Pi,coords=spherical);.
```

In Mathematica, use

```
SphericalPlot3D[Sin[phi],{phi,0,Pi},{theta,0,2Pi}].
```



41. The region E of integration is the region above the cone $z = \sqrt{x^2+y^2}$ and below the sphere $x^2+y^2+z^2 = 2$ in the first octant. Because E is in the first octant we have $0 \le \theta \le \frac{\pi}{2}$. The cone has equation $\phi = \frac{\pi}{4}$ (as in Example 4), so $0 \le \phi \le \frac{\pi}{4}$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and $0 \le \rho \le \sqrt{2}$. Then the integral becomes

$$\int_0^{\pi/4}\int_0^{\pi/2}\int_0^{\sqrt{2}} (\rho\sin\phi\cos\theta)\,(\rho\sin\phi\sin\theta)\,\rho^2\sin\phi\,d\rho\,d\theta\,d\phi$$

$$= \int_0^{\pi/4}\sin^3\phi\,d\phi\int_0^{\pi/2}\sin\theta\cos\theta\,d\theta\int_0^{\sqrt{2}}\rho^4\,d\rho = \left(\int_0^{\pi/4}\left(1-\cos^2\phi\right)\sin\phi\,d\phi\right)\left[\tfrac{1}{2}\sin^2\theta\right]_0^{\pi/2}\left[\tfrac{1}{5}\rho^5\right]_0^{\sqrt{2}}$$

$$= \left[\tfrac{1}{3}\cos^3\phi-\cos\phi\right]_0^{\pi/4}\cdot\tfrac{1}{2}\cdot\tfrac{1}{5}\left(\sqrt{2}\right)^5 = \left[\tfrac{\sqrt{2}}{12}-\tfrac{\sqrt{2}}{2}-\left(\tfrac{1}{3}-1\right)\right]\cdot\tfrac{2\sqrt{2}}{5} = \tfrac{4\sqrt{2}-5}{15}$$

42. The region of integration is the solid sphere $x^2+y^2+z^2 \le a^2$, so $0 \le \theta \le 2\pi$, $0 \le \phi \le \pi$, and $0 \le \rho \le a$. Also $x^2z+y^2z+z^3 = (x^2+y^2+z^2)z = \rho^2 z = \rho^3\cos\phi$, so the integral becomes

$$\int_0^{\pi}\int_0^{2\pi}\int_0^{a}\left(\rho^3\cos\phi\right)\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \int_0^{\pi}\sin\phi\cos\phi\,d\phi\int_0^{2\pi}d\theta\int_0^{a}\rho^5\,d\rho = \left[\tfrac{1}{2}\sin^2\phi\right]_0^{\pi}\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{6}\rho^6\right]_0^{a} = 0$$

43. The region of integration is the solid sphere $x^2+y^2+(z-2)^2 \le 4$ or equivalently

$\rho^2\sin^2\phi+(\rho\cos\phi-2)^2 = \rho^2-4\rho\cos\phi+4 \le 4 \quad\Rightarrow\quad \rho \le 4\cos\phi$, so $0 \le \theta \le 2\pi$, $0 \le \phi \le \frac{\pi}{2}$, and $0 \le \rho \le 4\cos\phi$. Also $(x^2+y^2+z^2)^{3/2} = (\rho^2)^{3/2} = \rho^3$, so the integral becomes

$$\int_0^{\pi/2}\int_0^{2\pi}\int_0^{4\cos\phi}\left(\rho^3\right)\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \int_0^{\pi/2}\int_0^{2\pi}\sin\phi\left[\tfrac{1}{6}\rho^6\right]_{\rho=0}^{\rho=4\cos\phi}d\theta\,d\phi$$

$$= \tfrac{1}{6}\int_0^{\pi/2}\int_0^{2\pi}\sin\phi\left(4096\cos^6\phi\right)d\theta\,d\phi$$

$$= \tfrac{1}{6}(4096)\int_0^{\pi/2}\cos^6\phi\sin\phi\,d\phi\int_0^{2\pi}d\theta = \tfrac{2048}{3}\left[-\tfrac{1}{7}\cos^7\phi\right]_0^{\pi/2}\left[\theta\right]_0^{2\pi}$$

$$= \tfrac{2048}{3}\left(\tfrac{1}{7}\right)(2\pi) = \tfrac{4096\pi}{21}$$

44. The solid region between the ground and an altitude of 5 km (5000 m) is given by

$$E = \left\{(\rho,\theta,\phi) \mid 6.370\times10^6 \le \rho \le 6.375\times10^6, 0 \le \theta \le 2\pi, 0 \le \phi \le \pi\right\}.$$

Then the mass of the atmosphere in this region is

$$m = \iiint_E \delta\,dV = \int_0^{2\pi}\int_0^{\pi}\int_{6.370\times10^6}^{6.375\times10^6}(619.09-0.000097\rho)\,\rho^2\sin\phi\,d\rho\,d\phi\,d\theta$$

$$= \int_0^{2\pi}d\theta\int_0^{\pi}\sin\phi\,d\phi\int_{6.370\times10^6}^{6.375\times10^6}\left(619.09\rho^2-0.000097\rho^3\right)d\rho$$

$$= \left[\theta\right]_0^{2\pi}\left[-\cos\phi\right]_0^{\pi}\left[\tfrac{619.09}{3}\rho^3-\tfrac{0.000097}{4}\rho^4\right]_{6.370\times10^6}^{6.375\times10^6}$$

$$= (2\pi)(2)\left[\tfrac{619.09}{3}\left((6.375\times10^6)^3-(6.370\times10^6)^3\right)-\tfrac{0.000097}{4}\left((6.375\times10^6)^4-(6.370\times10^6)^4\right)\right]$$

$$\approx 4\pi\left(1.944\times10^{17}\right) \approx 2.44\times10^{18}\text{ kg}$$

45. In cylindrical coordinates, the equation of the cylinder is $r = 3$, $0 \le z \le 10$.

The hemisphere is the upper part of the sphere radius 3, center $(0,0,10)$, equation $r^2+(z-10)^2 = 3^2$, $z \ge 10$. In Maple, we can use the `coords=cylindrical` option in a regular `plot3d` command. In Mathematica, we can use `ParametricPlot3D`.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

46. We begin by finding the positions of Los Angeles and Montréal in spherical coordinates, using the method described in the exercise:

Montréal	Los Angeles
$\rho = 3960$ mi	$\rho = 3960$ mi
$\theta = 360° - 73.60° = 286.40°$	$\theta = 360° - 118.25° = 241.75°$
$\phi = 90° - 45.50° = 44.50°$	$\phi = 90° - 34.06° = 55.94°$

Now we change the above to Cartesian coordinates using $x = \rho\cos\theta\sin\phi$, $y = \rho\sin\theta\sin\phi$ and $z = \rho\cos\phi$ to get two position vectors of length 3960 mi (since both cities must lie on the surface of the earth). In particular:

$$\text{Montréal: } \langle 783.67, -2662.67, 2824.47\rangle \qquad \text{Los Angeles: } \langle -1552.80, -2889.91, 2217.84\rangle$$

To find the angle γ between these two vectors we use the dot product:

$\langle 783.67, -2662.67, 2824.47\rangle \cdot \langle -1552.80, -2889.91, 2217.84\rangle = (3960)^2\cos\gamma \;\Rightarrow\; \cos\gamma \approx 0.8126 \;\Rightarrow$

$\gamma \approx 0.6223$ rad. The great circle distance between the cities is $s = \rho\gamma \approx 3960(0.6223) \approx 2464$ mi.

47. If E is the solid enclosed by the surface $\rho = 1 + \frac{1}{5}\sin 6\theta\,\sin 5\phi$, it can be described in spherical coordinates as $E = \{(\rho,\theta,\phi) \mid 0 \le \rho \le 1 + \frac{1}{5}\sin 6\theta\sin 5\phi, 0 \le \theta \le 2\pi, 0 \le \phi \le \pi\}$. Its volume is given by

$V(E) = \iiint_E dV = \int_0^{\pi}\int_0^{2\pi}\int_0^{1+(\sin 6\theta\sin 5\phi)/5} \rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \frac{136\pi}{99}$ [using a CAS].

48. The given integral is equal to $\lim\limits_{R\to\infty}\int_0^{2\pi}\int_0^{\pi}\int_0^{R} \rho e^{-\rho^2}\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = \lim\limits_{R\to\infty}\int_0^{2\pi} d\theta\int_0^{\pi}\sin\phi\,d\phi\int_0^{R}\rho^3 e^{-\rho^2}\,d\rho$. Now use integration by parts with $u = \rho^2$, $dv = \rho e^{-\rho^2}\,d\rho$ to get

$$\lim_{R\to\infty} 2\pi(2)\left(\rho^2\left(-\tfrac{1}{2}\right)e^{-\rho^2}\Big]_0^R - \int_0^R 2\rho\left(-\tfrac{1}{2}\right)e^{-\rho^2}\,d\rho\right) = \lim_{R\to\infty} 4\pi\left(-\tfrac{1}{2}R^2e^{-R^2} + \left[-\tfrac{1}{2}e^{-\rho^2}\right]_0^R\right)$$

$$= 4\pi\lim_{R\to\infty}\left[-\tfrac{1}{2}R^2e^{-R^2} - \tfrac{1}{2}e^{-R^2} + \tfrac{1}{2}\right] = 4\pi\left(\tfrac{1}{2}\right) = 2\pi$$

(Note that $R^2e^{-R^2} \to 0$ as $R \to \infty$ by l'Hospital's Rule.)

49. (a) From the diagram, $z = r\cot\phi_0$ to $z = \sqrt{a^2 - r^2}$, $r = 0$ to $r = a\sin\phi_0$ (or use $a^2 - r^2 = r^2\cot^2\phi_0$). Thus



$$V = \int_0^{2\pi}\int_0^{a\sin\phi_0}\int_{r\cot\phi_0}^{\sqrt{a^2-r^2}} r\,dz\,dr\,d\theta$$

$$= 2\pi\int_0^{a\sin\phi_0}\left(r\sqrt{a^2-r^2} - r^2\cot\phi_0\right)dr$$

$$= \tfrac{2\pi}{3}\left[-(a^2-r^2)^{3/2} - r^3\cot\phi_0\right]_0^{a\sin\phi_0}$$

$$= \tfrac{2\pi}{3}\left[-\left(a^2 - a^2\sin^2\phi_0\right)^{3/2} - a^3\sin^3\phi_0\cot\phi_0 + a^3\right]$$

$$= \tfrac{2}{3}\pi a^3\left[1 - \left(\cos^3\phi_0 + \sin^2\phi_0\cos\phi_0\right)\right] = \tfrac{2}{3}\pi a^3(1 - \cos\phi_0)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The wedge in question is the shaded area rotated from $\theta = \theta_1$ to $\theta = \theta_2$.



Letting

V_{ij} = volume of the region bounded by the sphere of radius ρ_i and the cone with angle ϕ_j ($\theta = \theta_1$ to θ_2)

and letting V be the volume of the wedge, we have

$$\begin{aligned} V &= (V_{22} - V_{21}) - (V_{12} - V_{11}) \\ &= \tfrac{1}{3}(\theta_2 - \theta_1)\left[\rho_2^3(1 - \cos\phi_2) - \rho_2^3(1 - \cos\phi_1) - \rho_1^3(1 - \cos\phi_2) + \rho_1^3(1 - \cos\phi_1)\right] \\ &= \tfrac{1}{3}(\theta_2 - \theta_1)\left[(\rho_2^3 - \rho_1^3)(1 - \cos\phi_2) - (\rho_2^3 - \rho_1^3)(1 - \cos\phi_1)\right] = \tfrac{1}{3}(\theta_2 - \theta_1)\left[(\rho_2^3 - \rho_1^3)(\cos\phi_1 - \cos\phi_2)\right] \end{aligned}$$

Or: Show that $V = \displaystyle\int_{\theta_1}^{\theta_2} \int_{\rho_1 \sin\phi_1}^{\rho_2 \sin\phi_2} \int_{r\cot\phi_2}^{r\cot\phi_1} r\, dz\, dr\, d\theta$.

(c) By the Mean Value Theorem with $f(\rho) = \rho^3$ there exists some $\tilde{\rho}$ with $\rho_1 \le \tilde{\rho} \le \rho_2$ such that $f(\rho_2) - f(\rho_1) = f'(\tilde{\rho})(\rho_2 - \rho_1)$ or $\rho_1^3 - \rho_2^3 = 3\tilde{\rho}^2 \Delta\rho$. Similarly there exists ϕ with $\phi_1 \le \tilde{\phi} \le \phi_2$ such that $\cos\phi_2 - \cos\phi_1 = \left(-\sin\tilde{\phi}\right)\Delta\phi$. Substituting into the result from (b) gives

$\Delta V = (\tilde{\rho}^2\, \Delta\rho)(\theta_2 - \theta_1)(\sin\tilde{\phi})\, \Delta\phi = \tilde{\rho}^2 \sin\tilde{\phi}\, \Delta\rho\, \Delta\phi\, \Delta\theta$.

APPLIED PROJECT Roller Derby

1. $mgh = \frac{1}{2}mv^2 + \frac{1}{2}I\omega^2 = \frac{1}{2}(m + I/r^2)v^2$, so $v^2 = \dfrac{2mgh}{m + I/r^2} = \dfrac{2gh}{1 + I^*}$.

2. The vertical component of the speed is $v \sin\alpha$, so

$$\frac{dy}{dt} = \sqrt{\frac{2gy}{1 + I^*}}\sin\alpha = \sqrt{\frac{2g}{1 + I^*}}\sin\alpha\,\sqrt{y}.$$



3. Solving the separable differential equation, we get $\dfrac{dy}{\sqrt{y}} = \sqrt{\dfrac{2g}{1 + I^*}}\sin\alpha\, dt \quad \Rightarrow \quad 2\sqrt{y} = \sqrt{\dfrac{2g}{1 + I^*}}\,(\sin\alpha)t + C.$

But $y = 0$ when $t = 0$, so $C = 0$ and we have $2\sqrt{y} = \sqrt{\dfrac{2g}{1 + I^*}}\,(\sin\alpha)t$. Solving for t when $y = h$ gives

$$T = \frac{2\sqrt{h}}{\sin\alpha}\sqrt{\frac{1 + I^*}{2g}} = \sqrt{\frac{2h(1 + I^*)}{g\sin^2\alpha}}.$$

4. Assume that the length of each cylinder is ℓ. Then the density of the solid cylinder is $\dfrac{m}{\pi r^2 \ell}$, and from Formulas 15.6.16, its moment of inertia (using cylindrical coordinates) is

$$I_z = \iiint \frac{m}{\pi r^2 \ell}\,(x^2 + y^2)\, dV = \int_0^{\ell}\int_0^{2\pi}\int_0^{r} \frac{m}{\pi r^2 \ell}\, R^2 R\, dR\, d\theta\, dz = \frac{m}{\pi r^2 \ell}\, 2\pi\ell\left[\tfrac{1}{4}R^4\right]_0^r = \frac{mr^2}{2}$$

and so $I^* = \dfrac{I_z}{mr^2} = \dfrac{1}{2}$. [continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

For the hollow cylinder, we consider its entire mass to lie a distance r from the axis of rotation, so $x^2+y^2=r^2$ is a constant. We express the density in terms of mass per unit area as $\rho=\dfrac{m}{2\pi r\ell}$, and then the moment of inertia is calculated as a double integral: $I_z=\iint (x^2+y^2)\,\dfrac{m}{2\pi r\ell}\,dA=\dfrac{mr^2}{2\pi r\ell}\iint dA=mr^2$, so $I^*=\dfrac{I_z}{mr^2}=1$.

5. The volume of such a ball is $\frac{4}{3}\pi(r^3-a^3)=\frac{4}{3}\pi r(1-b^3)$, and so its density is $\dfrac{m}{\frac{4}{3}\pi r^3(1-b^3)}$. Using Formula 15.8.3, we get

$$\begin{aligned}
I_z &= \iiint (x^2+y^2)\,\frac{m}{\frac{4}{3}\pi r^3(1-b^3)}\,dV \\
&= \frac{m}{\frac{4}{3}\pi r^3(1-b^3)}\int_a^r\int_0^{2\pi}\int_0^{\pi}(\rho^2\sin^2\phi)(\rho^2\sin\phi)\,d\phi\,d\theta\,d\rho \\
&= \frac{m}{\frac{4}{3}\pi r^3(1-b^3)}\cdot 2\pi\left[-\frac{(2+\sin^2\phi)\cos\phi}{3}\right]_0^{\pi}\left[\frac{\rho^5}{5}\right]_a^r \qquad \text{[from the Table of Integrals]} \\
&= \frac{m}{\frac{4}{3}\pi r^3(1-b^3)}\cdot 2\pi\cdot\frac{4}{3}\cdot\frac{r^5-a^5}{5}=\frac{2mr^5(1-b^5)}{5r^3(1-b^3)}=\frac{2(1-b^5)mr^2}{5(1-b^3)}
\end{aligned}$$

Therefore $I^*=\dfrac{2(1-b^5)}{5(1-b^3)}$. Since a represents the inner radius, $a\to 0$ corresponds to a solid ball, and $a\to r$ corresponds to a hollow ball.

6. For a solid ball, $a\to 0 \;\Rightarrow\; b\to 0$, so $I^*=\lim\limits_{b\to 0}\dfrac{2(1-b^5)}{5(1-b^3)}=\dfrac{2}{5}$. For a hollow ball, $a\to r \;\Rightarrow\; b\to 1$, so

$$I^*=\lim_{b\to 1}\frac{2(1-b^5)}{5(1-b^3)}=\frac{2}{5}\lim_{b\to 1}\frac{-5b^4}{-3b^2}=\frac{2}{5}\left(\frac{5}{3}\right)=\frac{2}{3} \qquad \text{[by l'Hospital's Rule]}$$

Note: We could instead have calculated $I^*=\lim\limits_{b\to 1}\dfrac{2(1-b)(1+b+b^2+b^3+b^4)}{5(1-b)(1+b+b^2)}=\dfrac{2\cdot 5}{5\cdot 3}=\dfrac{2}{3}$.

Thus the objects finish in the following order: solid ball $\left(I^*=\frac{2}{5}\right)$, solid cylinder $\left(I^*=\frac{1}{2}\right)$, hollow ball $\left(I^*=\frac{2}{3}\right)$, hollow cylinder $(I^*=1)$.

15.9 Change of Variables in Multiple Integrals

1. $x=2u+v,\ y=4u-v$.

The Jacobian is $\dfrac{\partial(x,y)}{\partial(u,v)}=\begin{vmatrix}\partial x/\partial u & \partial x/\partial v\\ \partial y/\partial u & \partial y/\partial v\end{vmatrix}=\begin{vmatrix}2 & 1\\ 4 & -1\end{vmatrix}=(2)(-1)-(1)(4)=-6$.

2. $x=u^2+uv,\ y=uv^2$.

$$\frac{\partial(x,y)}{\partial(u,v)}=\begin{vmatrix}\partial x/\partial u & \partial x/\partial v\\ \partial y/\partial u & \partial y/\partial v\end{vmatrix}=\begin{vmatrix}2u+v & u\\ v^2 & 2uv\end{vmatrix}=(2u+v)(2uv)-u(v^2)=4u^2v+2uv^2-uv^2=4u^2v+uv^2$$

3. $x=s\cos t,\ y=s\sin t$.

$$\frac{\partial(x,y)}{\partial(s,t)}=\begin{vmatrix}\partial x/\partial s & \partial x/\partial t\\ \partial y/\partial s & \partial y/\partial t\end{vmatrix}=\begin{vmatrix}\cos t & -s\sin t\\ \sin t & s\cos t\end{vmatrix}=s\cos^2 t-(-s\sin^2 t)=s(\cos^2 t+\sin^2 t)=s$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $x = pe^q,\ y = qe^p$.

$$\frac{\partial(x,y)}{\partial(p,q)} = \begin{vmatrix} \partial x/\partial p & \partial x/\partial q \\ \partial y/\partial p & \partial y/\partial q \end{vmatrix} = \begin{vmatrix} e^q & pe^q \\ qe^p & e^p \end{vmatrix} = e^q e^p - pe^q \cdot qe^p = e^{p+q} - pqe^{p+q} = (1-pq)e^{p+q}$$

5. $x = uv,\ y = vw,\ z = wu$.

$$\frac{\partial(x,y,z)}{\partial(u,v,w)} = \begin{vmatrix} \partial x/\partial u & \partial x/\partial v & \partial x/\partial w \\ \partial y/\partial u & \partial y/\partial v & \partial y/\partial w \\ \partial z/\partial u & \partial z/\partial v & \partial z/\partial w \end{vmatrix} = \begin{vmatrix} v & u & 0 \\ 0 & w & v \\ w & 0 & u \end{vmatrix} = v\begin{vmatrix} w & v \\ 0 & u \end{vmatrix} - u\begin{vmatrix} 0 & v \\ w & u \end{vmatrix} + 0\begin{vmatrix} 0 & w \\ w & 0 \end{vmatrix}$$

$$= v(uw - 0) - u(0 - vw) + 0 = uvw + uvw = 2uvw$$

6. $x = u + vw,\ y = v + wu,\ z = w + uv$.

$$\frac{\partial(x,y,z)}{\partial(u,v,w)} = \begin{vmatrix} 1 & w & v \\ w & 1 & u \\ v & u & 1 \end{vmatrix} = 1\begin{vmatrix} 1 & u \\ u & 1 \end{vmatrix} - w\begin{vmatrix} w & u \\ v & 1 \end{vmatrix} + v\begin{vmatrix} w & 1 \\ v & u \end{vmatrix} = 1(1-u^2) - w\,(w - uv) + v\,(uw - v)$$

$$= 1 - u^2 - w^2 + uvw + uvw - v^2 = 1 + 2uvw - u^2 - v^2 - w^2$$

7. The transformation maps the boundary of S to the boundary of the image R, so we first look at side S_1 in the uv-plane. S_1 is described by $v = 0,\ 0 \le u \le 3$, so $x = 2u + 3v = 2u$ and $y = u - v = u$. Eliminating u, we have $x = 2y,\ 0 \le x \le 6$. S_2 is the line segment $u = 3,\ 0 \le v \le 2$, so $x = 6 + 3v$ and $y = 3 - v$. Then $v = 3 - y \ \Rightarrow\ x = 6 + 3(3 - y) = 15 - 3y$, $6 \le x \le 12$. S_3 is the line segment $v = 2,\ 0 \le u \le 3$, so $x = 2u + 6$ and $y = u - 2$, giving $u = y + 2 \ \Rightarrow\ x = 2y + 10$, $6 \le x \le 12$. Finally, S_4 is the segment $u = 0,\ 0 \le v \le 2$, so $x = 3v$ and $y = -v \ \Rightarrow\ x = -3y,\ 0 \le x \le 6$.

The image of set S is the region R shown in the xy-plane, a parallelogram bounded by these four segments.



8. S_1 is the line segment $v = 0,\ 0 \le u \le 1$, so $x = v = 0$ and $y = u(1 + v^2) = u$. Since $0 \le u \le 1$, the image is the line segment $x = 0,\ 0 \le y \le 1$. S_2 is the segment $u = 1,\ 0 \le v \le 1$, so $x = v$ and $y = u(1 + v^2) = 1 + x^2$. Thus the image is the portion of the parabola $y = 1 + x^2$ for $0 \le x \le 1$. S_3 is the segment $v = 1,\ 0 \le u \le 1$, so $x = 1$ and $y = 2u$. The image is the segment $x = 1,\ 0 \le y \le 2$. S_4 is described by $u = 0,\ 0 \le v \le 1$, so $0 \le x = v \le 1$ and $y = u(1 + v^2) = 0$. The image is the line segment $y = 0,\ 0 \le x \le 1$. Thus, the image of S is the region R bounded by the parabola $y = 1 + x^2$, the x-axis, and the lines $x = 0,\ x = 1$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

9. S_1 is the line segment $u = v$, $0 \le u \le 1$, so $y = v = u$ and $x = u^2 = y^2$. Since $0 \le u \le 1$, the image is the portion of the parabola $x = y^2$, $0 \le y \le 1$. S_2 is the segment $v = 1$, $0 \le u \le 1$, thus $y = v = 1$ and $x = u^2$, so $0 \le x \le 1$. The image is the line segment $y = 1$, $0 \le x \le 1$. S_3 is the segment $u = 0$, $0 \le v \le 1$, so $x = u^2 = 0$ and $y = v \;\Rightarrow\; 0 \le y \le 1$. The image is the segment $x = 0$, $0 \le y \le 1$. Thus, the image of S is the region R in the first quadrant bounded by the parabola $x = y^2$, the y-axis, and the line $y = 1$.



10. Substituting $u = \dfrac{x}{a}$, $v = \dfrac{y}{b}$ into $u^2 + v^2 \le 1$ gives $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} \le 1$, so the image of $u^2 + v^2 \le 1$ is the elliptical region $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} \le 1$.



11. R is a parallelogram enclosed by the parallel lines $y = 2x - 1$, $y = 2x + 1$ and the parallel lines $y = 1 - x$, $y = 3 - x$. The first pair of equations can be written as $y - 2x = -1$, $y - 2x = 1$. If we let $u = y - 2x$ then these lines are mapped to the vertical lines $u = -1$, $u = 1$ in the uv-plane. Similarly, the second pair of equations can be written as $x + y = 1$, $x + y = 3$, and setting $v = x + y$ maps these lines to the horizontal lines $v = 1$, $v = 3$ in the uv-plane. Boundary curves are mapped to boundary curves under a transformation, so here the equations $u = y - 2x$, $v = x + y$ define a transformation T^{-1} that maps R in the xy-plane to the square S enclosed by the lines $u = -1$, $u = 1$, $v = 1$, $v = 3$ in the uv-plane. To find the transformation T that maps S to R we solve $u = y - 2x$, $v = x + y$ for x, y: Subtracting the first equation from the second gives $v - u = 3x \;\Rightarrow\; x = \frac{1}{3}(v - u)$ and adding twice the second equation to the first gives $u + 2v = 3y \;\Rightarrow$ $y = \frac{1}{3}(u + 2v)$. Thus one possible transformation T (there are many) is given by $x = \frac{1}{3}(v - u)$, $y = \frac{1}{3}(u + 2v)$.



12. The boundaries of the parallelogram R are the lines $y = \frac{3}{4}x$ or $4y - 3x = 0$, $y = \frac{3}{4}x + \frac{5}{2}$ or $4y - 3x = 10$, $y = -\frac{1}{2}x$ or $x + 2y = 0$, $y = -\frac{1}{2}x + 5$ or $x + 2y = 10$. Setting $u = 4y - 3x$ and $v = x + 2y$ defines a transformation T^{-1} that maps R

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

in the xy-plane to the square S enclosed by the lines $u = 0$, $u = 10$, $v = 0$, $v = 10$ in the uv-plane. Solving $u = 4y - 3x$, $v = x + 2y$ for x and y gives $2v - u = 5x \Rightarrow x = \frac{1}{5}(2v - u)$, $u + 3v = 10y \Rightarrow y = \frac{1}{10}(u + 3v)$. Thus one possible transformation T is given by $x = \frac{1}{5}(2v - u)$, $y = \frac{1}{10}(u + 3v)$.



13. R is a portion of an annular region (see the figure) that is easily described in polar coordinates as $R = \{(r, \theta) \mid 1 \le r \le \sqrt{2},\ 0 \le \theta \le \pi/2\}$. If we converted a double integral over R to polar coordinates the resulting region of integration is a rectangle (in the $r\theta$-plane), so we can create a transformation T here by letting u play the role of r and v the role of θ. Thus T is defined by $x = u\cos v$, $y = u\sin v$ and T maps the rectangle $S = \{(u, v) \mid 1 \le u \le \sqrt{2},\ 0 \le v \le \pi/2\}$ in the uv-plane to R in the xy-plane.



14. The boundaries of the region R are the curves $y = 1/x$ or $xy = 1$, $y = 4/x$ or $xy = 4$, $y = x$ or $y/x = 1$, $y = 4x$ or $y/x = 4$. Setting $u = xy$ and $v = y/x$ defines a transformation T^{-1} that maps R in the xy-plane to the square S enclosed by the lines $u = 1$, $u = 4$, $v = 1$, $v = 4$ in the uv-plane. Solving $u = xy$, $v = y/x$ for x and y gives $x^2 = u/v \Rightarrow x = \sqrt{u/v}$ [since x, y, u, v are all positive], $y^2 = uv \Rightarrow y = \sqrt{uv}$. Thus one possible transformation T is given by $x = \sqrt{u/v}$, $y = \sqrt{uv}$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

15. $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} = 3$ and $x - 3y = (2u+v) - 3(u+2v) = -u - 5v$. To find the region S in the uv-plane that corresponds to R we first find the corresponding boundary under the given transformation. The line through $(0,0)$ and $(2,1)$ is $y = \frac{1}{2}x$ which is the image of $u + 2v = \frac{1}{2}(2u+v) \;\Rightarrow\; v = 0$; the line through $(2,1)$ and $(1,2)$ is $x + y = 3$ which is the image of $(2u+v) + (u+2v) = 3 \;\Rightarrow\; u + v = 1$; the line through $(0,0)$ and $(1,2)$ is $y = 2x$ which is the image of $u + 2v = 2(2u+v) \;\Rightarrow\; u = 0$. Thus S is the triangle $0 \le v \le 1 - u$, $0 \le u \le 1$ in the uv-plane and

$$\iint_R (x-3y)\,dA = \int_0^1 \int_0^{1-u} (-u-5v)\,|3|\,dv\,du = -3\int_0^1 \left[uv + \tfrac{5}{2}v^2\right]_{v=0}^{v=1-u} du$$
$$= -3\int_0^1 \left(u - u^2 + \tfrac{5}{2}(1-u)^2\right) du = -3\left[\tfrac{1}{2}u^2 - \tfrac{1}{3}u^3 - \tfrac{5}{6}(1-u)^3\right]_0^1 = -3\left(\tfrac{1}{2} - \tfrac{1}{3} + \tfrac{5}{6}\right) = -3$$

16. $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/4 & 1/4 \\ -3/4 & 1/4 \end{vmatrix} = \frac{1}{4}$, $4x + 8y = 4 \cdot \frac{1}{4}(u+v) + 8 \cdot \frac{1}{4}(v - 3u) = 3v - 5u$. R is a parallelogram bounded by the lines $x - y = -4$, $x - y = 4$, $3x + y = 0$, $3x + y = 8$. Since $u = x - y$ and $v = 3x + y$, R is the image of the rectangle enclosed by the lines $u = -4$, $u = 4$, $v = 0$, and $v = 8$. Thus

$$\iint_R (4x+8y)\,dA = \int_{-4}^4 \int_0^8 (3v-5u)\left|\tfrac{1}{4}\right| dv\,du = \tfrac{1}{4}\int_{-4}^4 \left[\tfrac{3}{2}v^2 - 5uv\right]_{v=0}^{v=8} du$$
$$= \tfrac{1}{4}\int_{-4}^4 (96 - 40u)\,du = \tfrac{1}{4}\left[96u - 20u^2\right]_{-4}^4 = 192$$

17. $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 2 & 0 \\ 0 & 3 \end{vmatrix} = 6$, $x^2 = 4u^2$ and the planar ellipse $9x^2 + 4y^2 \le 36$ is the image of the disk $u^2 + v^2 \le 1$. Thus

$$\iint_R x^2\,dA = \iint_{u^2+v^2\le 1} (4u^2)(6)\,du\,dv = \int_0^{2\pi}\int_0^1 (24r^2\cos^2\theta)\,r\,dr\,d\theta = 24\int_0^{2\pi}\cos^2\theta\,d\theta\int_0^1 r^3\,dr$$
$$= 24\left[\tfrac{1}{2}x + \tfrac{1}{4}\sin 2x\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^1 = 24(\pi)\left(\tfrac{1}{4}\right) = 6\pi$$

18. $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \sqrt{2} & -\sqrt{2/3} \\ \sqrt{2} & \sqrt{2/3} \end{vmatrix} = \dfrac{4}{\sqrt{3}}$, $x^2 - xy + y^2 = 2u^2 + 2v^2$ and the planar ellipse $x^2 - xy + y^2 \le 2$ is the image of the disk $u^2 + v^2 \le 1$. Thus

$$\iint_R (x^2 - xy + y^2)\,dA = \iint_{u^2+v^2\le 1} (2u^2+2v^2)\left(\tfrac{4}{\sqrt{3}}\,du\,dv\right) = \int_0^{2\pi}\int_0^1 \tfrac{8}{\sqrt{3}}r^3\,dr\,d\theta = \tfrac{4\pi}{\sqrt{3}}$$

19. $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/v & -u/v^2 \\ 0 & 1 \end{vmatrix} = \dfrac{1}{v}$, $xy = u$, $y = x$ is the image of the parabola $v^2 = u$, $y = 3x$ is the image of the parabola $v^2 = 3u$, and the hyperbolas $xy = 1$, $xy = 3$ are the images of the lines $u = 1$ and $u = 3$ respectively. Thus

$$\iint_R xy\,dA = \int_1^3 \int_{\sqrt{u}}^{\sqrt{3u}} u\left(\frac{1}{v}\right) dv\,du = \int_1^3 u\left(\ln\sqrt{3u} - \ln\sqrt{u}\right) du = \int_1^3 u\ln\sqrt{3}\,du = 4\ln\sqrt{3} = 2\ln 3.$$

20. Here $y = \dfrac{v}{u}$, $x = \dfrac{u^2}{v}$ so $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 2u/v & -u^2/v^2 \\ -v/u^2 & 1/u \end{vmatrix} = \dfrac{1}{v}$ and R is the image of the square with vertices $(1,1)$, $(2,1)$, $(2,2)$, and $(1,2)$. So



$$\iint_R y^2\,dA = \int_1^2 \int_1^2 \frac{v^2}{u^2}\left(\frac{1}{v}\right) du\,dv = \int_1^2 \frac{v}{2}\,dv = \frac{3}{1}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. (a) $\dfrac{\partial(x,y,z)}{\partial(u,v,w)} = \begin{vmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{vmatrix} = abc$ and since $u = \dfrac{x}{a}$, $v = \dfrac{y}{b}$, $w = \dfrac{z}{c}$ the solid enclosed by the ellipsoid is the image of the ball $u^2 + v^2 + w^2 \le 1$. So

$$\iiint_E dV = \iiint_{u^2+v^2+w^2 \le 1} abc\, du\, dv\, dw = (abc)(\text{volume of the ball}) = \tfrac{4}{3}\pi abc$$

(b) If we approximate the surface of the earth by the ellipsoid $\dfrac{x^2}{6378^2} + \dfrac{y^2}{6378^2} + \dfrac{z^2}{6356^2} = 1$, then we can estimate the volume of the earth by finding the volume of the solid E enclosed by the ellipsoid. From part (a), this is $\iiint_E dV = \frac{4}{3}\pi(6378)(6378)(6356) \approx 1.083 \times 10^{12}$ km^3.

(c) The moment of intertia about the z-axis is $I_z = \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV$, where E is the solid enclosed by $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} + \dfrac{z^2}{c^2} = 1$. As in part (a), we use the transformation $x = au$, $y = bv$, $z = cw$, so $\left|\dfrac{\partial(x,y,z)}{\partial(u,v,w)}\right| = abc$ and

$$\begin{aligned} I_z &= \iiint_E (x^2+y^2)\,k\,dV = \iiint_{u^2+v^2+w^2 \le 1} k(a^2u^2 + b^2v^2)(abc)\,du\,dv\,dw \\ &= abck \int_0^{\pi}\int_0^{2\pi}\int_0^1 (a^2\rho^2\sin^2\phi\cos^2\theta + b^2\rho^2\sin^2\phi\sin^2\theta)\,\rho^2\sin\phi\,d\rho\,d\theta\,d\phi \\ &= abck\left[a^2\int_0^{\pi}\int_0^{2\pi}\int_0^1 (\rho^2\sin^2\phi\cos^2\theta)\,\rho^2\sin\phi\,d\rho\,d\theta\,d\phi + b^2\int_0^{\pi}\int_0^{2\pi}\int_0^1 (\rho^2\sin^2\phi\sin^2\theta)\,\rho^2\sin\phi\,d\rho\,d\theta\,d\phi\right] \\ &= a^3bck\int_0^{\pi}\sin^3\phi\,d\phi\int_0^{2\pi}\cos^2\theta\,d\theta\int_0^1\rho^4\,d\rho + ab^3ck\int_0^{\pi}\sin^3\phi\,d\phi\int_0^{2\pi}\sin^2\theta\,d\theta\int_0^1\rho^4\,d\rho \\ &= a^3bck\left[\tfrac{1}{3}\cos^3\phi - \cos\phi\right]_0^{\pi}\left[\tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{5}\rho^5\right]_0^1 + ab^3ck\left[\tfrac{1}{3}\cos^3\phi - \cos\phi\right]_0^{\pi}\left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{5}\rho^5\right]_0^1 \\ &= a^3bck\left(\tfrac{4}{3}\right)(\pi)\left(\tfrac{1}{5}\right) + ab^3ck\left(\tfrac{4}{3}\right)(\pi)\left(\tfrac{1}{5}\right) = \tfrac{4}{15}\pi(a^2+b^2)abck \end{aligned}$$

22. R is the region enclosed by the curves $xy = a$, $xy = b$, $xy^{1.4} = c$, and $xy^{1.4} = d$, so if we let $u = xy$ and $v = xy^{1.4}$ then R is the image of the rectangle enclosed by the lines $u = a$, $u = b$ $(a < b)$ and $v = c$, $v = d$ $(c < d)$. Now

$x = u/y \quad \Rightarrow \quad v = (u/y)y^{1.4} = uy^{0.4} \quad \Rightarrow \quad y^{0.4} = u^{-1}v \quad \Rightarrow \quad y = (u^{-1}v)^{1/0.4} = u^{-2.5}v^{2.5}$ and

$x = uy^{-1} = u(u^{-2.5}v^{2.5})^{-1} = u^{3.5}v^{-2.5}$, so

$\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 3.5u^{2.5}v^{-2.5} & -2.5u^{3.5}v^{-3.5} \\ -2.5u^{-3.5}v^{2.5} & 2.5u^{-2.5}v^{1.5} \end{vmatrix} = 8.75v^{-1} - 6.25v^{-1} = 2.5v^{-1}$. Thus the area of R, and the work done by the engine, is

$$\iint_R dA = \int_a^b\int_c^d \left|2.5v^{-1}\right| dv\,du = 2.5\int_a^b du\int_c^d (1/v)\,dv = 2.5\big[u\big]_a^b\big[\ln|v|\big]_c^d = 2.5(b-a)(\ln d - \ln c) = 2.5(b-a)\ln\frac{d}{c}.$$

23. Letting $u = x - 2y$ and $v = 3x - y$, we have $x = \frac{1}{5}(2v - u)$ and $y = \frac{1}{5}(v - 3u)$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} -1/5 & 2/5 \\ -3/5 & 1/5 \end{vmatrix} = \dfrac{1}{5}$ and R is the image of the rectangle enclosed by the lines $u = 0$, $u = 4$, $v = 1$, and $v = 8$. Thus

$$\iint_R \frac{x-2y}{3x-y}\,dA = \int_0^4\int_1^8 \frac{u}{v}\left|\frac{1}{5}\right| dv\,du = \frac{1}{5}\int_0^4 u\,du\int_1^8 \frac{1}{v}\,dv = \tfrac{1}{5}\big[\tfrac{1}{2}u^2\big]_0^4\big[\ln|v|\big]_1^8 = \tfrac{8}{5}\ln 8.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

24. Letting $u = x + y$ and $v = x - y$, we have $x = \frac{1}{2}(u+v)$ and $y = \frac{1}{2}(u-v)$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/2 & 1/2 \\ 1/2 & -1/2 \end{vmatrix} = -\dfrac{1}{2}$ and R is the image of the rectangle enclosed by the lines $u = 0$, $u = 3$, $v = 0$, and $v = 2$. Thus

$$\iint_R (x+y)\,e^{x^2-y^2}\,dA = \int_0^3 \int_0^2 ue^{uv}\left|-\tfrac{1}{2}\right| dv\,du = \tfrac{1}{2}\int_0^3 \left[e^{uv}\right]_{v=0}^{v=2} du = \tfrac{1}{2}\int_0^3 (e^{2u}-1)\,du$$
$$= \tfrac{1}{2}\left[\tfrac{1}{2}e^{2u} - u\right]_0^3 = \tfrac{1}{2}\left(\tfrac{1}{2}e^6 - 3 - \tfrac{1}{2}\right) = \tfrac{1}{4}(e^6 - 7)$$

25. Letting $u = y - x$, $v = y + x$, we have $y = \frac{1}{2}(u+v)$, $x = \frac{1}{2}(v-u)$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} -1/2 & 1/2 \\ 1/2 & 1/2 \end{vmatrix} = -\dfrac{1}{2}$ and R is the image of the trapezoidal region with vertices $(-1,1)$, $(-2,2)$, $(2,2)$, and $(1,1)$. Thus

$$\iint_R \cos\left(\frac{y-x}{y+x}\right) dA = \int_1^2 \int_{-v}^{v} \cos\frac{u}{v}\left|-\frac{1}{2}\right| du\,dv = \frac{1}{2}\int_1^2 \left[v\sin\frac{u}{v}\right]_{u=-v}^{u=v} dv = \frac{1}{2}\int_1^2 2v\sin(1)\,dv = \tfrac{3}{2}\sin 1$$

26. Letting $u = 3x$, $v = 2y$, we have $9x^2 + 4y^2 = u^2 + v^2$, $x = \frac{1}{3}u$, and $y = \frac{1}{2}v$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \dfrac{1}{6}$ and R is the image of the quarter-disk D given by $u^2 + v^2 \le 1$, $u \ge 0$, $v \ge 0$. Thus

$$\iint_R \sin(9x^2+4y^2)\,dA = \iint_D \tfrac{1}{6}\sin(u^2+v^2)\,du\,dv = \int_0^{\pi/2}\int_0^1 \tfrac{1}{6}\sin(r^2)\,r\,dr\,d\theta = \tfrac{\pi}{12}\left[-\tfrac{1}{2}\cos r^2\right]_0^1 = \tfrac{\pi}{24}(1-\cos 1)$$

27. Let $u = x + y$ and $v = -x + y$. Then $u + v = 2y \;\Rightarrow\; y = \frac{1}{2}(u+v)$ and $u - v = 2x \;\Rightarrow\; x = \frac{1}{2}(u-v)$.

$\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/2 & -1/2 \\ 1/2 & 1/2 \end{vmatrix} = \dfrac{1}{2}$. Now $|u| = |x+y| \le |x| + |y| \le 1 \;\Rightarrow\; -1 \le u \le 1$,



and $|v| = |-x+y| \le |x| + |y| \le 1 \;\Rightarrow\; -1 \le v \le 1$. R is the image of the square region with vertices $(1,1)$, $(1,-1)$, $(-1,-1)$, and $(-1,1)$.

So $\iint_R e^{x+y}\,dA = \frac{1}{2}\int_{-1}^1\int_{-1}^1 e^u\,du\,dv = \frac{1}{2}\left[e^u\right]_{-1}^1\left[v\right]_{-1}^1 = e - e^{-1}$.

28. Let $u = x + y$ and $v = y$, then $x = u - v$, $y = v$, $\dfrac{\partial(x,y)}{\partial(u,v)} = 1$ and R is the image under T of the triangular region with vertices $(0,0)$, $(1,0)$ and $(1,1)$. Thus

$$\iint_R f(x+y)\,dA = \int_0^1\int_0^u (1)\,f(u)\,dv\,du = \int_0^1 f(u)\left[\,v\,\right]_{v=0}^{v=u} du = \int_0^1 uf(u)\,du \quad \text{as desired.}$$

15 Review

TRUE-FALSE QUIZ

1. This is true by Fubini's Theorem.

2. False. $\int_0^1\int_0^x \sqrt{x+y^2}\,dy\,dx$ describes the region of integration as a Type I region. To reverse the order of integration, we must consider the region as a Type II region: $\int_0^1\int_y^1 \sqrt{x+y^2}\,dx\,dy$.

3. True by Equation 15.1.11.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $\int_{-1}^{1}\int_{0}^{1} e^{x^2+y^2}\sin y\,dx\,dy = \left(\int_{0}^{1} e^{x^2}\,dx\right)\left(\int_{-1}^{1} e^{y^2}\sin y\,dy\right) = \left(\int_{0}^{1} e^{x^2}\,dx\right)(0) = 0$, since $e^{y^2}\sin y$ is an odd function. Therefore the statement is true.

5. True. By Equation 15.1.11 we can write $\int_0^1\int_0^1 f(x)\,f(y)\,dy\,dx = \int_0^1 f(x)\,dx\int_0^1 f(y)\,dy$. But $\int_0^1 f(y)\,dy = \int_0^1 f(x)\,dx$ so this becomes $\int_0^1 f(x)\,dx\int_0^1 f(x)\,dx = \left[\int_0^1 f(x)\,dx\right]^2$.

6. This statement is true because in the given region, $\left(x^2+\sqrt{y}\,\right)\sin(x^2y^2) \le (1+2)(1) = 3$, so $\int_1^4\int_0^1 \left(x^2+\sqrt{y}\,\right)\sin(x^2y^2)\,dx\,dy \le \int_1^4\int_0^1 3\,dA = 3A(D) = 3(3) = 9$.

7. True: $\iint_D \sqrt{4-x^2-y^2}\,dA =$ the volume under the surface $x^2+y^2+z^2=4$ and above the xy-plane
$= \frac{1}{2}$ (the volume of the sphere $x^2+y^2+z^2=4$) $= \frac{1}{2}\cdot\frac{4}{3}\pi(2)^3 = \frac{16}{3}\pi$

8. True. The moment of inertia about the z-axis of a solid E with constant density k is $I_z = \iiint_E (x^2+y^2)\rho(x,y,z)\,dV = \iiint_E (kr^2)\,r\,dz\,dr\,d\theta = \iiint_E kr^3\,dz\,dr\,d\theta$.

9. The volume enclosed by the cone $z=\sqrt{x^2+y^2}$ and the plane $z=2$ is, in cylindrical coordinates, $V = \int_0^{2\pi}\int_0^2\int_r^2 r\,dz\,dr\,d\theta \ne \int_0^{2\pi}\int_0^2\int_r^2 dz\,dr\,d\theta$, so the assertion is false.

EXERCISES

1. As shown in the contour map, we divide R into 9 equally sized subsquares, each with area $\Delta A = 1$. Then we approximate $\iint_R f(x,y)\,dA$ by a Riemann sum with $m=n=3$ and the sample points the upper right corners of each square, so

$$\begin{aligned}\iint_R f(x,y)\,dA &\approx \sum_{i=1}^{3}\sum_{j=1}^{3} f(x_i,y_j)\,\Delta A\\ &= \Delta A\,[f(1,1)+f(1,2)+f(1,3)+f(2,1)+f(2,2)+f(2,3)+f(3,1)+f(3,2)+f(3,3)]\end{aligned}$$

Using the contour lines to estimate the function values, we have

$$\iint_R f(x,y)\,dA \approx 1[2.7+4.7+8.0+4.7+6.7+10.0+6.7+8.6+11.9] \approx 64.0$$

2. As in Exercise 1, we have $m=n=3$ and $\Delta A = 1$. Using the contour map to estimate the value of f at the center of each subsquare, we have

$$\begin{aligned}\iint_R f(x,y)\,dA &\approx \sum_{i=1}^{3}\sum_{j=1}^{3} f(\overline{x}_i,\overline{y}_j)\,\Delta A\\ &= \Delta A\,[f(0.5,0.5)+(0.5,1.5)+(0.5,2.5)+(1.5,0.5)+f(1.5,1.5)\\ &\qquad +f(1.5,2.5)+(2.5,0.5)+f(2.5,1.5)+f(2.5,2.5)]\\ &\approx 1[1.2+2.5+5.0+3.2+4.5+7.1+5.2+6.5+9.0] = 44.2\end{aligned}$$

3. $\int_1^2\int_0^2 (y+2xe^y)\,dx\,dy = \int_1^2 \left[xy+x^2e^y\right]_{x=0}^{x=2} dy = \int_1^2 (2y+4e^y)\,dy = \left[y^2+4e^y\right]_1^2$
$= 4+4e^2-1-4e = 4e^2-4e+3$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

4. $\int_0^1 \int_0^1 ye^{xy}\,dx\,dy = \int_0^1 \left[e^{xy}\right]_{x=0}^{x=1} dy = \int_0^1 (e^y - 1)\,dy = \left[e^y - y\right]_0^1 = e - 2$

5. $\int_0^1 \int_0^x \cos(x^2)\,dy\,dx = \int_0^1 \left[\cos(x^2)y\right]_{y=0}^{y=x} dx = \int_0^1 x\cos(x^2)\,dx = \frac{1}{2}\sin(x^2)\Big]_0^1 = \frac{1}{2}\sin 1$

6. $\int_0^1 \int_x^{e^x} 3xy^2\,dy\,dx = \int_0^1 \left[xy^3\right]_{y=x}^{y=e^x} dx = \int_0^1 (xe^{3x} - x^4)\,dx = \frac{1}{3}xe^{3x}\Big]_0^1 - \int_0^1 \frac{1}{3}e^{3x}\,dx - \left[\frac{1}{5}x^5\right]_0^1$ [integrate by parts in the first term]

$= \frac{1}{3}e^3 - \left[\frac{1}{9}e^{3x}\right]_0^1 - \frac{1}{5} = \frac{2}{9}e^3 - \frac{4}{45}$

7. $\int_0^\pi \int_0^1 \int_0^{\sqrt{1-y^2}} y\sin x\,dz\,dy\,dx = \int_0^\pi \int_0^1 \left[(y\sin x)z\right]_{z=0}^{z=\sqrt{1-y^2}} dy\,dx = \int_0^\pi \int_0^1 y\sqrt{1-y^2}\sin x\,dy\,dx$

$= \int_0^\pi \left[-\frac{1}{3}(1-y^2)^{3/2}\sin x\right]_{y=0}^{y=1} dx = \int_0^\pi \frac{1}{3}\sin x\,dx = -\frac{1}{3}\cos x\Big]_0^\pi = \frac{2}{3}$

8. $\int_0^1 \int_0^y \int_x^1 6xyz\,dz\,dx\,dy = \int_0^1 \int_0^y \left[3xyz^2\right]_{z=x}^{z=1} dx\,dy = \int_0^1 \int_0^y (3xy - 3x^3y)\,dx\,dy$

$= \int_0^1 \left[\frac{3}{2}x^2y - \frac{3}{4}x^4y\right]_{x=0}^{x=y} dy = \int_0^1 \left(\frac{3}{2}y^3 - \frac{3}{4}y^5\right) dy = \left[\frac{3}{8}y^4 - \frac{1}{8}y^6\right]_0^1 = \frac{1}{4}$

9. The region R is more easily described by polar coordinates: $R = \{(r,\theta) \mid 2 \le r \le 4, 0 \le \theta \le \pi\}$. Thus $\iint_R f(x,y)\,dA = \int_0^\pi \int_2^4 f(r\cos\theta, r\sin\theta)\,r\,dr\,d\theta$.

10. The region R is a type II region that can be described as the region enclosed by the lines $y = 4 - x$, $y = 4 + x$, and the x-axis. So using rectangular coordinates, we can say $R = \{(x,y) \mid 0 \le y \le 4, y - 4 \le x \le 4 - y\}$ and $\iint_R f(x,y)\,dA = \int_0^4 \int_{y-4}^{4-y} f(x,y)\,dx\,dy$.

11. $x = r\cos\theta = 2\sqrt{3}\cos\frac{\pi}{3} = 2\sqrt{3}\cdot\frac{1}{2} = \sqrt{3}$, $y = r\sin\theta = 2\sqrt{3}\sin\frac{\pi}{3} = 2\sqrt{3}\cdot\frac{\sqrt{3}}{2} = 3$, $z = 2$, so in rectangular coordinates the point is $\left(\sqrt{3}, 3, 2\right)$. $\rho = \sqrt{r^2 + z^2} = \sqrt{12 + 4} = 4$, $\theta = \frac{\pi}{3}$, and $\cos\phi = z/\rho = \frac{1}{2}$, so $\phi = \frac{\pi}{3}$ and spherical coordinates are $\left(4, \frac{\pi}{3}, \frac{\pi}{3}\right)$.

12. $r = \sqrt{4+4} = 2\sqrt{2}$; $z = -1$; $\tan\theta = \frac{2}{2} = 1$ and the point $(2,2)$ is in the first quadrant of the xy-plane, so $\theta = \frac{\pi}{4}$. Thus in cylindrical coordinates the point is $\left(2\sqrt{2}, \frac{\pi}{4}, -1\right)$. $\rho = \sqrt{4+4+1} = 3$, $\cos\phi = z/\rho = -\frac{1}{3}$, so the spherical coordinates are $\left(3, \frac{\pi}{4}, \cos^{-1}\left(-\frac{1}{3}\right)\right)$.

13. $x = \rho\sin\phi\cos\theta = 8\sin\frac{\pi}{6}\cos\frac{\pi}{4} = 8\cdot\frac{1}{2}\cdot\frac{\sqrt{2}}{2} = 2\sqrt{2}$, $y = \rho\sin\phi\sin\theta = 8\sin\frac{\pi}{6}\sin\frac{\pi}{4} = 2\sqrt{2}$, and $z = \rho\cos\phi = 8\cos\frac{\pi}{6} = 8\cdot\frac{\sqrt{3}}{2} = 4\sqrt{3}$. Thus rectangular coordinates for the point are $\left(2\sqrt{2}, 2\sqrt{2}, 4\sqrt{3}\right)$.

$r^2 = x^2 + y^2 = 8 + 8 = 16 \quad \Rightarrow \quad r = 4$, $\theta = \frac{\pi}{4}$, and $z = 4\sqrt{3}$, so cylindrical coordinates are $\left(4, \frac{\pi}{4}, 4\sqrt{3}\right)$.

14. (a) $\theta = \frac{\pi}{4}$. In cylindrical coordinates (assuming that r can be negative), this is a vertical plane that includes the z-axis and intersects the xy-plane in the line $y = x$. In spherical coordinates, because $\rho \ge 0$ and $0 \le \phi \le \pi$, we get a vertical half-plane that includes the z-axis and intersects the xy-plane in the half-line $y = x$, $x \ge 0$.

(b) $\phi = \frac{\pi}{4}$. In spherical coordinates, this is one frustum of a circular cone with vertex the origin and axis the positive z-axis.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. (a) $x^2 + y^2 + z^2 = 4$. In cylindrical coordinates, this becomes $r^2 + z^2 = 4$. In spherical coordinates, it becomes $\rho^2 = 4$ or $\rho = 2$.

(b) $x^2 + y^2 = 4$. In cylindrical coordinates: $r^2 = 4$ or $r = 2$. In spherical coordinates: $\rho^2 - z^2 = 4$ or $\rho^2 - \rho^2\cos^2\phi = 4$ or $\rho^2\sin^2\phi = 4$ or $\rho\sin\phi = 2$.

16. $\rho = 2\cos\phi \;\Rightarrow\; \rho^2 = 2\rho\cos\phi \;\Rightarrow\; x^2 + y^2 + z^2 = 2z \;\Rightarrow$ $x^2 + y^2 + (z-1)^2 = 1$. This is the equation of a sphere with radius 1, centered at $(0, 0, 1)$. Therefore, $0 \le \rho \le 2\cos\phi$ is the solid ball whose boundary is this sphere. $0 \le \theta \le \frac{\pi}{2}$ and $0 \le \phi \le \frac{\pi}{6}$ restrict the solid to the section of this ball that lies above the cone $\phi = \frac{\pi}{6}$ and is in the first octant.



17.



The region whose area is given by $\int_0^{\pi/2}\int_0^{\sin 2\theta} r\,dr\,d\theta$ is $\{(r,\theta) \mid 0 \le \theta \le \frac{\pi}{2}, 0 \le r \le \sin 2\theta\}$, which is the region contained in the loop in the first quadrant of the four-leaved rose $r = \sin 2\theta$.

18. The solid is $\{(\rho,\theta,\phi) \mid 1 \le \rho \le 2,\ 0 \le \theta \le \frac{\pi}{2},\ 0 \le \phi \le \frac{\pi}{2}\}$ which is the region in the first octant on or between the two spheres $\rho = 1$ and $\rho = 2$.

19.



$$\int_0^1\int_x^1 \cos(y^2)\,dy\,dx = \int_0^1\int_0^y \cos(y^2)\,dx\,dy$$
$$= \int_0^1 \cos(y^2)\big[x\big]_{x=0}^{x=y}\,dy = \int_0^1 y\cos(y^2)\,dy$$
$$= \big[\tfrac{1}{2}\sin(y^2)\big]_0^1 = \tfrac{1}{2}\sin 1$$

20.



$$\int_0^1\int_{\sqrt{y}}^1 \frac{ye^{x^2}}{x^3}\,dx\,dy = \int_0^1\int_0^{x^2} \frac{ye^{x^2}}{x^3}\,dy\,dx = \int_0^1 \frac{e^{x^2}}{x^3}\big[\tfrac{1}{2}y^2\big]_{y=0}^{y=x^2}\,dx$$
$$= \int_0^1 \tfrac{1}{2}xe^{x^2}\,dx = \tfrac{1}{4}e^{x^2}\Big]_0^1 = \tfrac{1}{4}(e \quad 1)$$

21. $\iint_R ye^{xy}\,dA = \int_0^3\int_0^2 ye^{xy}\,dx\,dy = \int_0^3 \big[e^{xy}\big]_{x=0}^{x=2}\,dy = \int_0^3 (e^{2y} - 1)\,dy = \big[\tfrac{1}{2}e^{2y} - y\big]_0^3 = \tfrac{1}{2}e^6 - 3 - \tfrac{1}{2} = \tfrac{1}{2}e^6 \quad \tfrac{7}{2}$

22. $\iint_D xy\,dA = \int_0^1\int_{y^2}^{y+2} xy\,dx\,dy = \int_0^1 y\big[\tfrac{1}{2}x^2\big]_{x=y^2}^{x=y+2}\,dy = \tfrac{1}{2}\int_0^1 y\big[(y+2)^2 - y^4\big]\,dy$

$$= \tfrac{1}{2}\int_0^1 (y^3 + 4y^2 + 4y - y^5)\,dy = \tfrac{1}{2}\big[\tfrac{1}{4}y^4 + \tfrac{4}{3}y^3 + 2y^2 - \tfrac{1}{6}y^6\big]_0^1 = \tfrac{41}{24}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

23.



$$\iint_D \frac{y}{1+x^2}\,dA = \int_0^1 \int_0^{\sqrt{x}} \frac{y}{1+x^2}\,dy\,dx = \int_0^1 \frac{1}{1+x^2}\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=\sqrt{x}}\,dx$$
$$= \tfrac{1}{2}\int_0^1 \frac{x}{1+x^2}\,dx = \left[\tfrac{1}{4}\ln(1+x^2)\right]_0^1 = \tfrac{1}{4}\ln 2$$

24. $$\iint_D \frac{1}{1+x^2}\,dA = \int_0^1 \int_x^1 \frac{1}{1+x^2}\,dy\,dx = \int_0^1 \frac{1}{1+x^2}\left[y\right]_{y=x}^{y=1}\,dx = \int_0^1 \frac{1-x}{1+x^2}\,dx = \int_0^1 \left(\frac{1}{1+x^2} - \frac{x}{1+x^2}\right)dx$$
$$= \left[\tan^{-1}x - \tfrac{1}{2}\ln(1+x^2)\right]_0^1 = \tan^{-1}1 - \tfrac{1}{2}\ln 2 - \left(\tan^{-1}0 - \tfrac{1}{2}\ln 1\right) = \tfrac{\pi}{4} - \tfrac{1}{2}\ln 2$$

25.



$$\iint_D y\,dA = \int_0^2 \int_{y^2}^{8-y^2} y\,dx\,dy$$
$$= \int_0^2 y\left[x\right]_{x=y^2}^{x=8-y^2}\,dy = \int_0^2 y(8-y^2-y^2)\,dy$$
$$= \int_0^2 (8y-2y^3)\,dy = \left[4y^2 - \tfrac{1}{2}y^4\right]_0^2 = 8$$

26.



$$\iint_D y\,dA = \int_1^2 \int_{1/y}^{y} y\,dx\,dy = \int_1^2 y\left(y - \frac{1}{y}\right)dy$$
$$= \int_1^2 (y^2-1)\,dy = \left[\tfrac{1}{3}y^3 - y\right]_1^2$$
$$= \left(\tfrac{8}{3} - 2\right) - \left(\tfrac{1}{3} - 1\right) = \tfrac{4}{3}$$

27.



$$\iint_D (x^2+y^2)^{3/2}\,dA = \int_0^{\pi/3} \int_0^3 (r^2)^{3/2} r\,dr\,d\theta$$
$$= \int_0^{\pi/3} d\theta \int_0^3 r^4\,dr = \left[\theta\right]_0^{\pi/3}\left[\tfrac{1}{5}r^5\right]_0^3$$
$$= \frac{\pi}{3}\frac{3^5}{5} = \frac{81\pi}{5}$$

28. $$\iint_D x\,dA = \int_0^{\pi/2}\int_1^{\sqrt{2}} (r\cos\theta)\,r\,dr\,d\theta = \int_0^{\pi/2}\cos\theta\,d\theta \int_1^{\sqrt{2}} r^2\,dr = \left[\sin\theta\right]_0^{\pi/2}\left[\tfrac{1}{3}r^3\right]_1^{\sqrt{2}}$$
$$= 1\cdot\tfrac{1}{3}(2^{3/2}-1) = \tfrac{1}{3}(2^{3/2}-1)$$

29. $$\iiint_E xy\,dV = \int_0^3\int_0^x\int_0^{x+y} xy\,dz\,dy\,dx = \int_0^3\int_0^x xy\left[z\right]_{z=0}^{z=x+y}\,dy\,dx = \int_0^3\int_0^x xy(x+y)\,dy\,dx$$
$$= \int_0^3\int_0^x (x^2y + xy^2)\,dy\,dx = \int_0^3 \left[\tfrac{1}{2}x^2y^2 + \tfrac{1}{3}xy^3\right]_{y=0}^{y=x}\,dx = \int_0^3 \left(\tfrac{1}{2}x^4 + \tfrac{1}{3}x^4\right)dx$$
$$= \tfrac{5}{6}\int_0^3 x^4\,dx = \left[\tfrac{1}{6}x^5\right]_0^3 = \tfrac{81}{2} = 40.5$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $\iiint_T xy\,dV = \int_0^{1/3}\int_0^{1-3x}\int_0^{1-3x-y} xy\,dz\,dy\,dx = \int_0^{1/3}\int_0^{1-3x} xy(1-3x-y)\,dy\,dx$

$= \int_0^{1/3}\int_0^{1-3x}(xy - 3x^2y - xy^2)\,dy\,dx$

$= \int_0^{1/3}\left[\frac{1}{2}xy^2 - \frac{3}{2}x^2y^2 - \frac{1}{3}xy^3\right]_{y=0}^{y=1-3x} dx$

$= \int_0^{1/3}\left[\frac{1}{2}x(1-3x)^2 - \frac{3}{2}x^2(1-3x)^2 - \frac{1}{3}x(1-3x)^3\right] dx$

$= \int_0^{1/3}\left(\frac{1}{6}x - \frac{3}{2}x^2 + \frac{9}{2}x^3 - \frac{9}{2}x^4\right) dx$

$= \frac{1}{12}x^2 - \frac{1}{2}x^3 + \frac{9}{8}x^4 - \frac{9}{10}x^5\Big]_0^{1/3} = \frac{1}{1080}$



31. $\iiint_E y^2z^2\,dV = \int_{-1}^{1}\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{1-y^2-z^2} y^2z^2\,dx\,dz\,dy = \int_{-1}^{1}\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} y^2z^2(1-y^2-z^2)\,dz\,dy$

$= \int_0^{2\pi}\int_0^1 (r^2\cos^2\theta)(r^2\sin^2\theta)(1-r^2)\,r\,dr\,d\theta = \int_0^{2\pi}\left(\frac{1}{2}\sin 2\theta\right)^2 d\theta \int_0^1 (r^5 - r^7)\,dr$

$= \int_0^{2\pi}\frac{1}{4}\left[\frac{1}{2}(1-\cos 4\theta)\right] d\theta \int_0^1 (r^5 - r^7)\,dr = \frac{1}{8}\left[\theta - \frac{1}{4}\sin 4\theta\right]_0^{2\pi}\left[\frac{1}{6}r^6 - \frac{1}{8}r^8\right]_0^1$

$= \frac{1}{8}(2\pi)\left(\frac{1}{6} - \frac{1}{8}\right) = \frac{\pi}{4}\cdot\frac{1}{24} = \frac{\pi}{96}$

32. $\iiint_E z\,dV = \int_0^1\int_0^{\sqrt{1-y^2}}\int_0^{2-y} z\,dx\,dz\,dy = \int_0^1\int_0^{\sqrt{1-y^2}}(2-y)z\,dz\,dy = \int_0^1 \frac{1}{2}(2-y)(1-y^2)\,dy$

$= \int_0^1 \frac{1}{2}(2 - y - 2y^2 + y^3)\,dy = \frac{13}{24}$

33. $\iiint_E yz\,dV = \int_{-2}^{2}\int_0^{\sqrt{4-x^2}}\int_0^y yz\,dz\,dy\,dx = \int_{-2}^{2}\int_0^{\sqrt{4-x^2}}\left[\frac{1}{2}yz^2\right]_{z=0}^{z=y} dy\,dx = \frac{1}{2}\int_{-2}^{2}\int_0^{\sqrt{4-x^2}} y^3\,dy\,dx$

$= \frac{1}{2}\int_0^{\pi}\int_0^2 (r\sin\theta)^3\,r\,dr\,d\theta = \frac{1}{2}\int_0^{\pi}\sin^3\theta\,d\theta \int_0^2 r^4\,dr = \frac{1}{2}\int_0^{\pi}(1-\cos^2\theta)\sin\theta\,d\theta \int_0^2 r^4\,dr$

$= \frac{1}{2}\left[-\cos\theta + \frac{1}{3}\cos^3\theta\right]_0^{\pi}\left[\frac{1}{5}r^5\right]_0^2 = \frac{1}{2}\left(\frac{2}{3} + \frac{2}{3}\right)\left(\frac{32}{5}\right) = \frac{64}{15}$

34. $\iiint_H z^3\sqrt{x^2+y^2+z^2}\,dV = \int_0^{2\pi}\int_0^{\pi/2}\int_0^1 (\rho^3\cos^3\phi)\rho(\rho^2\sin\phi)\,d\rho\,d\phi\,d\theta$

$= \int_0^{2\pi} d\theta \int_0^{\pi/2}\cos^3\phi\sin\phi\,d\phi \int_0^1 \rho^6\,d\rho = 2\pi\left[-\frac{1}{4}\cos^4\phi\right]_0^{\pi/2}\left(\frac{1}{7}\right) = \frac{\pi}{14}$

35. $V = \int_0^2\int_1^4 (x^2 + 4y^2)\,dy\,dx = \int_0^2\left[x^2y + \frac{4}{3}y^3\right]_{y=1}^{y=4} dx = \int_0^2 (3x^2 + 84)\,dx = x^3 + 84x\Big]_0^2 = 176$

36.



$V = \int_0^1\int_{y+1}^{4-2y}\int_0^{x^2y} dz\,dx\,dy = \int_0^1\int_{y+1}^{4-2y} x^2y\,dx\,dy$

$= \int_0^1 \frac{1}{3}\left[(4-2y)^3y - (y+1)^3y\right] dy$

$= \int_0^1 3(-y^4 + 5y^3 - 11y^2 + 7y)\,dy = 3\left(-\frac{1}{5} + \frac{5}{4} - \frac{11}{3} + \frac{7}{2}\right) = \frac{53}{20}$

37.



$V = \int_0^2\int_0^y\int_0^{(2-y)/2} dz\,dx\,dy = \int_0^2\int_0^y \left(1 - \frac{1}{2}y\right) dx\,dy$

$= \int_0^2 \left(y - \frac{1}{2}y^2\right) dy = \frac{1}{2}y^2 - \frac{1}{6}y^3\Big]_0^2 = \frac{2}{3}$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

38. $V = \int_0^{2\pi}\int_0^2\int_0^{3-r\sin\theta} r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^2 (3r - r^2\sin\theta)\,dr\,d\theta = \int_0^{2\pi}\left[6 - \frac{8}{3}\sin\theta\right]d\theta = 6\theta + \frac{8}{3}\cos\theta\Big]_0^{2\pi} = 12\pi$

39. Using the wedge above the plane $z = 0$ and below the plane $z = mx$ and noting that we have the same volume for $m < 0$ as for $m > 0$ (so use $m > 0$), we have

$V = 2\int_0^{a/3}\int_0^{\sqrt{a^2-9y^2}} mx\,dx\,dy = 2\int_0^{a/3} \frac{1}{2}m(a^2 - 9y^2)\,dy = m\big[a^2y - 3y^3\big]_0^{a/3} = m\big(\frac{1}{3}a^3 - \frac{1}{9}a^3\big) = \frac{2}{9}ma^3.$

40. The paraboloid and the half-cone intersect when $x^2 + y^2 = \sqrt{x^2+y^2}$, that is when $x^2 + y^2 = 1$ or 0. So

$V = \iint\limits_{x^2+y^2\le 1}\int_{x^2+y^2}^{\sqrt{x^2+y^2}} dz\,dA = \int_0^{2\pi}\int_0^1\int_{r^2}^{r} r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^1 (r^2 - r^3)\,dr\,d\theta = \int_0^{2\pi}\big(\frac{1}{3} - \frac{1}{4}\big)\,d\theta = \frac{1}{12}(2\pi) = \frac{\pi}{6}.$

41. (a) $m = \int_0^1\int_0^{1-y^2} y\,dx\,dy = \int_0^1 (y - y^3)\,dy = \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$

(b) $M_y = \int_0^1\int_0^{1-y^2} xy\,dx\,dy = \int_0^1 \frac{1}{2}y(1-y^2)^2\,dy = -\frac{1}{12}(1-y^2)^3\big]_0^1 = \frac{1}{12}$,

$M_x = \int_0^1\int_0^{1-y^2} y^2\,dx\,dy = \int_0^1 (y^2 - y^4)\,dy = \frac{2}{15}$. Hence $(\overline{x},\overline{y}) = \big(\frac{1}{3}, \frac{8}{15}\big)$.

(c) $I_x = \int_0^1\int_0^{1-y^2} y^3\,dx\,dy = \int_0^1 (y^3 - y^5)\,dy = \frac{1}{12}$,

$I_y = \int_0^1\int_0^{1-y^2} yx^2\,dx\,dy = \int_0^1 \frac{1}{3}y(1-y^2)^3\,dy = -\frac{1}{24}(1-y^2)^4\big]_0^1 = \frac{1}{24}$,

$\overline{\overline{y}}^2 = I_x/m = \frac{1/12}{1/4} = \frac{1}{3} \;\Rightarrow\; \overline{\overline{y}} = \frac{1}{\sqrt{3}}$, and $\overline{\overline{x}}^2 = I_y/m = \frac{1/24}{1/4} = \frac{1}{6} \;\Rightarrow\; \overline{\overline{x}} = \frac{1}{\sqrt{6}}$.

42. (a) In polar coordinates, the lamina occupies the region $D = \{(r,\theta) \mid 0 \le r \le a, 0 \le \theta \le \pi/2\}$. Assuming constant density K, then $m = K\,A(D) = K \cdot \frac{1}{4}\pi a^2 = \frac{1}{4}\pi K a^2$,

$M_y = \iint_D Kx\,dA = K\int_0^{\pi/2}\int_0^a (r\cos\theta)\,r\,dr\,d\theta = K\int_0^{\pi/2}\cos\theta\,d\theta\int_0^a r^2\,dr = K\,[\sin\theta]_0^{\pi/2}\,\big[\frac{1}{3}r^3\big]_0^a = \frac{1}{3}Ka^3$, and

$M_x = \iint_D Ky\,dA = K\int_0^{\pi/2}\sin\theta\,d\theta\int_0^a r^2\,dr = K\,[-\cos\theta]_0^{\pi/2}\,\big[\frac{1}{3}r^3\big]_0^a = \frac{1}{3}Ka^3$ [by symmetry $M_y = M_x$].

Thus the centroid is $(\overline{x},\overline{y}) = (M_y/m, M_x/m) = \big(\frac{4}{3\pi}a, \frac{4}{3\pi}a\big)$.

(b) $m = \iint_D \rho(x,y)\,dA = \iint_D xy^2\,dA = \int_0^{\pi/2}\int_0^a (r\cos\theta)(r\sin\theta)^2\,r\,dr\,d\theta$

$= \int_0^{\pi/2}\sin^2\theta\,\cos\theta\,d\theta\int_0^a r^4\,dr = \big[\frac{1}{3}\sin^3\theta\big]_0^{\pi/2}\big[\frac{1}{5}r^5\big]_0^a = \frac{1}{15}a^5$,

$M_y = \int_0^{\pi/2}\int_0^a r^5\cos^2\theta\sin^2\theta\,dr\,d\theta = \frac{1}{8}\big[\theta - \frac{1}{4}\sin 4\theta\big]_0^{\pi/2}\big[\frac{1}{6}r^6\big]_0^a = \frac{1}{96}\pi a^6$, and

$M_x = \int_0^{\pi/2}\int_0^a r^5\cos\theta\sin^3\theta\,dr\,d\theta = \big[\frac{1}{4}\sin^4\theta\big]_0^{\pi/2}\big[\frac{1}{6}r^6\big]_0^a = \frac{1}{24}a^6$. Hence $(\overline{x},\overline{y}) = \big(\frac{5}{32}\pi a, \frac{5}{8}a\big)$.

43. (a) A right circular cone with axis the z-axis and vertex at the origin has equation $z^2 = c^2(x^2 + y^2)$. Here we have the bottom frustum, shifted upward h units, and with $c^2 = h^2/a^2$ so that the cone includes the point $(a, 0, 0)$. Thus an equation of the cone in rectangular coordinates is $z = h - \frac{h}{a}\sqrt{x^2+y^2}$, $0 \le z \le h$. In cylindrical coordinates, the cone is described by $E = \big\{(r,\theta,z) \mid 0 \le r \le a,\ 0 \le \theta \le 2\pi,\ 0 \le z \le h\,(1 - \frac{1}{a}r)\big\}$, and its volume is $V = \frac{1}{3}\pi a^2 h$. By symmetry

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$M_{yz} = M_{xz} = 0$, and

$$M_{xy} = \int_0^{2\pi}\int_0^a\int_0^{h(1-r/a)} z \cdot r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^a \left[\frac{1}{2}rz^2\right]_{z=0}^{z=h(1-r/a)} dr\,d\theta$$

$$= \frac{1}{2}\int_0^{2\pi}\int_0^a rh^2\left(1-\frac{r}{a}\right)^2 dr\,d\theta = \frac{1}{2}h^2\int_0^{2\pi} d\theta \int_0^a \left(r - \frac{2}{a}r^2 + \frac{1}{a^2}r^3\right) dr$$

$$= \tfrac{1}{2}h^2\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{2}r^2 - \tfrac{2}{3a}r^3 + \tfrac{1}{4a^2}r^4\right]_0^a = \tfrac{1}{2}h^2(2\pi)\left(\tfrac{1}{2}a^2 - \tfrac{2}{3}a^2 + \tfrac{1}{4}a^2\right)$$

$$= \pi h^2\left(\tfrac{1}{12}a^2\right) = \tfrac{1}{12}\pi a^2h^2$$

Hence the centroid is $(\overline{x}, \overline{y}, \overline{z}) = \left(0, 0, [\pi a^2h^2/12]/[\pi a^2h/3]\right) = \left(0, 0, \tfrac{1}{4}h\right)$.

(b) The density function is $\rho = \sqrt{x^2+y^2} = \sqrt{r^2} = r$, so the moment of inertia about the cone's axis (the z-axis) is

$$I_z = \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^a\int_0^{h(1-r/a)} (r^2)(r)\,r\,dz\,dr\,d\theta$$

$$= \int_0^{2\pi}\int_0^a \left[r^4z\right]_{z=0}^{z=h(1-r/a)} dr\,d\theta = \int_0^{2\pi}\int_0^a r^4h\left(1-\tfrac{1}{a}r\right) dr\,d\theta$$

$$= h\int_0^{2\pi} d\theta \int_0^a \left(r^4 - \tfrac{1}{a}r^5\right) dr = h\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{5}r^5 - \tfrac{1}{6a}r^6\right]_0^a$$

$$= h(2\pi)\left(\tfrac{1}{5}a^5 - \tfrac{1}{6}a^5\right) = \tfrac{1}{15}\pi a^5h$$

44. $1 \le z^2 \le 4 \;\Rightarrow\; 1/a^2 \le x^2+y^2 \le 4/a^2$. Let $D = \left\{(x,y) \mid 1/a^2 \le x^2+y^2 \le 4/a^2\right\}$. $z = f(x,y) = a\sqrt{x^2+y^2}$, so $f_x(x,y) = ax(x^2+y^2)^{-1/2}$, $f_y(x,y) = ay(x^2+y^2)^{-1/2}$, and

$$A(S) = \iint_D \sqrt{\frac{a^2x^2+a^2y^2}{x^2+y^2} + 1}\,dA = \iint_D \sqrt{a^2+1}\,dA = \sqrt{a^2+1}\,A(D)$$

$$= \sqrt{a^2+1}\left[\pi\left(\frac{2}{a}\right)^2 - \pi\left(\frac{1}{a}\right)^2\right] = \frac{3\pi}{a^2}\sqrt{a^2+1}$$

45. Let D represent the given triangle; then D can be described as the area enclosed by the x- and y-axes and the line $y = 2-2x$, or equivalently $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le 2-2x\}$. We want to find the surface area of the part of the graph of $z = x^2+y$ that lies over D, so using Equation 15.5.3 we have

$$A(S) = \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\,dA = \iint_D \sqrt{1+(2x)^2+(1)^2}\,dA = \int_0^1\int_0^{2-2x} \sqrt{2+4x^2}\,dy\,dx$$

$$= \int_0^1 \sqrt{2+4x^2}\left[y\right]_{y=0}^{y=2-2x} dx = \int_0^1 (2-2x)\sqrt{2+4x^2}\,dx = \int_0^1 2\sqrt{2+4x^2}\,dx - \int_0^1 2x\sqrt{2+4x^2}\,dx$$

Using Formula 21 in the Table of Integrals with $a = \sqrt{2}$, $u = 2x$, and $du = 2\,dx$, we have $\int 2\sqrt{2+4x^2}\,dx = x\sqrt{2+4x^2} + \ln\left(2x+\sqrt{2+4x^2}\right)$. If we substitute $u = 2+4x^2$ in the second integral, then $du = 8x\,dx$ and $\int 2x\sqrt{2+4x^2}\,dx = \tfrac{1}{4}\int \sqrt{u}\,du = \tfrac{1}{4}\cdot\tfrac{2}{3}u^{3/2} = \tfrac{1}{6}(2+4x^2)^{3/2}$. Thus

$$A(S) = \left[x\sqrt{2+4x^2} + \ln\left(2x+\sqrt{2+4x^2}\right) - \tfrac{1}{6}(2+4x^2)^{3/2}\right]_0^1$$

$$= \sqrt{6} + \ln\left(2+\sqrt{6}\right) - \tfrac{1}{6}(6)^{3/2} - \ln\sqrt{2} + \tfrac{\sqrt{2}}{3} = \ln\tfrac{2+\sqrt{6}}{\sqrt{2}} + \tfrac{\sqrt{2}}{3}$$

$$= \ln\left(\sqrt{2}+\sqrt{3}\right) + \tfrac{\sqrt{2}}{3} \approx 1.6176$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

46. Using Formula 15.5.3 with $\partial z/\partial x = \sin y$, $\partial z/\partial y = x\cos y$, we get

$S = \int_{-\pi}^{\pi}\int_{-3}^{3}\sqrt{1+\sin^2 y + x^2\cos^2 y}\,dx\,dy \approx 62.9714.$



47.



$$\int_0^3\int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}}(x^3+xy^2)\,dy\,dx = \int_0^3\int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} x(x^2+y^2)\,dy\,dx$$
$$= \int_{-\pi/2}^{\pi/2}\int_0^3 (r\cos\theta)(r^2)\,r\,dr\,d\theta$$
$$= \int_{-\pi/2}^{\pi/2}\cos\theta\,d\theta\int_0^3 r^4\,dr$$
$$= \left[\sin\theta\right]_{-\pi/2}^{\pi/2}\left[\tfrac{1}{5}r^5\right]_0^3 = 2\cdot\tfrac{1}{5}(243) = \tfrac{486}{5} = 97.2$$

48. The region of integration is the solid hemisphere $x^2+y^2+z^2 \le 4$, $x \ge 0$.

$$\int_{-2}^{2}\int_0^{\sqrt{4-y^2}}\int_{-\sqrt{4-x^2-y^2}}^{\sqrt{4-x^2-y^2}} y^2\sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$$
$$= \int_{-\pi/2}^{\pi/2}\int_0^{\pi}\int_0^2 (\rho\sin\phi\sin\theta)^2\left(\sqrt{\rho^2}\right)\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = \int_{-\pi/2}^{\pi/2}\sin^2\theta\,d\theta\int_0^{\pi}\sin^3\phi\,d\phi\int_0^2\rho^5\,d\rho$$
$$= \left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_{-\pi/2}^{\pi/2}\left[-\cos\phi + \tfrac{1}{3}\cos^3\phi\right]_0^{\pi}\left[\tfrac{1}{6}\rho^6\right]_0^2 = \left(\tfrac{\pi}{2}\right)\left(\tfrac{2}{3}+\tfrac{2}{3}\right)\left(\tfrac{32}{3}\right) = \tfrac{64}{9}\pi$$

49. From the graph, it appears that $1-x^2 = e^x$ at $x \approx -0.71$ and at $x = 0$, with $1-x^2 > e^x$ on $(-0.71, 0)$. So the desired integral is

$$\iint_D y^2\,dA \approx \int_{-0.71}^0\int_{e^x}^{1-x^2} y^2\,dy\,dx$$
$$= \tfrac{1}{3}\int_{-0.71}^0\left[(1-x^2)^3 - e^{3x}\right]dx$$
$$= \tfrac{1}{3}\left[x - x^3 + \tfrac{3}{5}x^5 - \tfrac{1}{7}x^7 - \tfrac{1}{3}e^{3x}\right]_{-0.71}^0 \approx 0.0512$$



50. Let the tetrahedron be called T. The front face of T is given by the plane $x + \frac{1}{2}y + \frac{1}{3}z = 1$, or $z = 3 - 3x - \frac{3}{2}y$, which intersects the xy-plane in the line $y = 2-2x$. So the total mass is

$m = \iiint_T \rho(x,y,z)\,dV = \int_0^1\int_0^{2-2x}\int_0^{3-3x-3y/2}(x^2+y^2+z^2)\,dz\,dy\,dx = \frac{7}{5}$. The center of mass is

$(\overline{x},\overline{y},\overline{z}) = \left(m^{-1}\iiint_T x\rho(x,y,z)\,dV, m^{-1}\iiint_T y\rho(x,y,z)\,dV, m^{-1}\iiint_T z\rho(x,y,z)\,dV\right) = \left(\frac{4}{21},\frac{11}{21},\frac{8}{7}\right)$.

51. (a) $f(x,y)$ is a joint density function, so we know that $\iint_{\mathbb{R}^2} f(x,y)\,dA = 1$. Since $f(x,y) = 0$ outside the rectangle $[0,3]\times[0,2]$, we can say

$$\iint_{\mathbb{R}^2} f(x,y)\,dA = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y)\,dy\,dx = \int_0^3\int_0^2 C(x+y)\,dy\,dx$$
$$= C\int_0^3\left[xy + \tfrac{1}{2}y^2\right]_{y=0}^{y=2}dx = C\int_0^3(2x+2)\,dx = C\left[x^2+2x\right]_0^3 = 15C$$

Then $15C = 1 \;\Rightarrow\; C = \frac{1}{15}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $P(X \le 2, Y \ge 1) = \int_{-\infty}^{2} \int_{1}^{\infty} f(x,y)\,dy\,dx = \int_0^2 \int_1^2 \frac{1}{15}(x,y)\,dy\,dx = \frac{1}{15}\int_0^2 \left[xy + \frac{1}{2}y^2\right]_{y=1}^{y=2} dx$

$= \frac{1}{15}\int_0^2 \left(x + \frac{3}{2}\right) dx = \frac{1}{15}\left[\frac{1}{2}x^2 + \frac{3}{2}x\right]_0^2 = \frac{1}{3}$

(c) $P(X+Y \le 1) = P((X,Y) \in D)$ where D is the triangular region shown in the figure. Thus



$$P(X+Y \le 1) = \iint_D f(x,y)\,dA = \int_0^1 \int_0^{1-x} \frac{1}{15}(x+y)\,dy\,dx$$

$$= \frac{1}{15}\int_0^1 \left[xy + \frac{1}{2}y^2\right]_{y=0}^{y=1-x} dx$$

$$= \frac{1}{15}\int_0^1 \left[x(1-x) + \frac{1}{2}(1-x)^2\right] dx$$

$$= \frac{1}{30}\int_0^1 (1-x^2)\,dx = \frac{1}{30}\left[x - \frac{1}{3}x^3\right]_0^1 = \frac{1}{45}$$

52. Each lamp has exponential density function

$$f(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{1}{800}e^{-t/800} & \text{if } t \ge 0 \end{cases}$$

If X, Y, and Z are the lifetimes of the individual bulbs, then X, Y, and Z are independent, so the joint density function is the product of the individual density functions:

$$f(x,y,z) = \begin{cases} \frac{1}{800^3}e^{-(x+y+z)/800} & \text{if } x \ge 0, y \ge 0, z \ge 0 \\ 0 & \text{otherwise} \end{cases}$$

The probability that all three bulbs fail within a total of 1000 hours is $P(X+Y+Z \le 1000)$, or equivalently $P((X,Y,Z) \in E)$ where E is the solid region in the first octant bounded by the coordinate planes and the plane $x+y+z = 1000$. The plane $x+y+z = 1000$ meets the xy-plane in the line $x+y = 1000$, so we have

$$P(X+Y+Z \le 1000) = \iiint_E f(x,y,z)\,dV = \int_0^{1000}\int_0^{1000-x}\int_0^{1000-x-y} \frac{1}{800^3}e^{-(x+y+z)/800}\,dz\,dy\,dx$$

$$= \frac{1}{800^3}\int_0^{1000}\int_0^{1000-x} -800\left[e^{-(x+y+z)/800}\right]_{z=0}^{z=1000-x-y} dy\,dx$$

$$= \frac{-1}{800^2}\int_0^{1000}\int_0^{1000-x}\left[e^{-5/4} - e^{-(x+y)/800}\right] dy\,dx$$

$$= \frac{-1}{800^2}\int_0^{1000}\left[e^{-5/4}y + 800e^{-(x+y)/800}\right]_{y=0}^{y=1000-x} dx$$

$$= \frac{-1}{800^2}\int_0^{1000}\left[e^{-5/4}(1800-x) - 800e^{-x/800}\right] dx$$

$$= \frac{-1}{800^2}\left[-\frac{1}{2}e^{-5/4}(1800-x)^2 + 800^2 e^{-x/800}\right]_0^{1000}$$

$$= \frac{-1}{800^2}\left[-\frac{1}{2}e^{-5/4}(800)^2 + 800^2 e^{-5/4} + \frac{1}{2}e^{-5/4}(1800)^2 - 800^2\right]$$

$$= 1 - \frac{97}{32}e^{-5/4} \approx 0.1315$$

53.





$$\int_{-1}^{1}\int_{x^2}^{1}\int_0^{1-y} f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^{1-z}\int_{-\sqrt{y}}^{\sqrt{y}} f(x,y,z)\,dx\,dy\,dz$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

54.









$\int_0^2\int_0^{y^3}\int_0^{y^2} f(x,y,z)\,dz\,dx\,dy = \iiint_E f(x,y,z)\,dV$ where $E = \{(x,y,z) \mid 0 \le y \le 2, 0 \le x \le y^3, 0 \le z \le y^2\}$.

If D_1, D_2, and D_3 are the projections of E onto the xy-, yz-, and xz-planes, then

$D_1 = \{(x,y) \mid 0 \le y \le 2, 0 \le x \le y^3\} = \{(x,y) \mid 0 \le x \le 8, \sqrt[3]{x} \le y \le 2\}$,

$D_2 = \{(y,z) \mid 0 \le z \le 4, \sqrt{z} \le y \le 2\} = \{(y,z) \mid 0 \le y \le 2, 0 \le z \le y^2\}$, $D_3 = \{(x,z) \mid 0 \le x \le 8, 0 \le z \le 4\}$.

Therefore we have

$$\begin{aligned}\int_0^2\int_0^{y^3}\int_0^{y^2} f(x,y,z)\,dz\,dx\,dy &= \int_0^8\int_{\sqrt[3]{x}}^2\int_0^{y^2} f(x,y,z)\,dz\,dy\,dx = \int_0^4\int_{\sqrt{z}}^2\int_0^{y^3} f(x,y,z)\,dx\,dy\,dz\\ &= \int_0^2\int_0^{y^2}\int_0^{y^3} f(x,y,z)\,dx\,dz\,dy\\ &= \int_0^8\int_0^{x^{2/3}}\int_{\sqrt[3]{x}}^2 f(x,y,z)\,dy\,dz\,dx + \int_0^8\int_{x^{2/3}}^4\int_{\sqrt{z}}^2 f(x,y,z)\,dy\,dz\,dx\\ &= \int_0^4\int_0^{z^{3/2}}\int_{\sqrt{z}}^2 f(x,y,z)\,dy\,dx\,dz + \int_0^4\int_{z^{3/2}}^8\int_{\sqrt[3]{x}}^2 f(x,y,z)\,dy\,dx\,dz\end{aligned}$$

55. Since $u = x - y$ and $v = x + y$, $x = \frac{1}{2}(u+v)$ and $y = \frac{1}{2}(v-u)$. Thus $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/2 & 1/2 \\ -1/2 & 1/2 \end{vmatrix} = \frac{1}{2}$.

R is the image under this transformation of the square with vertices $(u,v) = (-2,2)$, $(0,2)$, $(0,4)$, and $(-2,4)$. So

$$\begin{aligned}\iint_R \frac{x-y}{x+y}\,dA &= \int_2^4\int_{-2}^0 \frac{u}{v}\left(\frac{1}{2}\right)du\,dv = \frac{1}{2}\int_2^4\left[\frac{u^2}{2v}\right]_{u=-2}^{u=0} dv = \frac{1}{2}\int_2^4\left(-\frac{2}{v}\right)dv\\ &= -\ln v\Big]_2^4 = -\ln 4 + \ln 2 = -2\ln 2 + \ln 2 = -\ln 2.\end{aligned}$$

56. $\dfrac{\partial(x,y,z)}{\partial(u,v,w)} = \begin{vmatrix} 2u & 0 & 0 \\ 0 & 2v & 0 \\ 0 & 0 & 2w \end{vmatrix} = 8uvw$, so

$$\begin{aligned}V &= \iiint_E dV = \int_0^1\int_0^{1-u}\int_0^{1-u-v} 8uvw\,dw\,dv\,du = \int_0^1\int_0^{1-u} 4uv(1-u-v)^2\,du\\ &= \int_0^1\int_0^{1-u}\left[4u(1-u)^2v - 8u(1-u)v^2 + 4uv^3\right]dv\,du\\ &= \int_0^1\left[2u(1-u)^4 - \tfrac{8}{3}u(1-u)^4 + u(1-u)^4\right]du = \int_0^1 \tfrac{1}{3}u(1-u)^4\,du\\ &= \int_0^1 \tfrac{1}{3}\left[(1-u)^4 - (1-u)^5\right]du = \tfrac{1}{3}\left[-\tfrac{1}{5}(1-u)^5 + \tfrac{1}{6}(1-u)^6\right]_0^1 = \tfrac{1}{3}\left(-\tfrac{1}{6} + \tfrac{1}{5}\right) = \tfrac{1}{90}\end{aligned}$$

57. Let $u = y - x$ and $v = y + x$ so $x = y - u = (v - x) - u \;\Rightarrow\; x = \frac{1}{2}(v-u)$ and $y = v - \frac{1}{2}(v-u) = \frac{1}{2}(v+u)$.

$\left|\dfrac{\partial(x,y)}{\partial(u,v)}\right| = \left|\dfrac{\partial x}{\partial u}\dfrac{\partial y}{\partial v} - \dfrac{\partial x}{\partial v}\dfrac{\partial y}{\partial u}\right| = \left|\left(-\tfrac{1}{2}\right)\left(\tfrac{1}{2}\right) - \left(\tfrac{1}{2}\right)\left(\tfrac{1}{2}\right)\right| = \left|-\tfrac{1}{2}\right| = \tfrac{1}{2}$. R is the image under this transformation of the square

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

with vertices $(u, v) = (0, 0)$, $(-2, 0)$, $(0, 2)$, and $(-2, 2)$. So

$$\iint_R xy\,dA = \int_0^2 \int_{-2}^0 \frac{v^2 - u^2}{4}\left(\frac{1}{2}\right) du\,dv = \tfrac{1}{8}\int_0^2 \left[v^2 u - \tfrac{1}{3}u^3\right]_{u=-2}^{u=0} dv = \tfrac{1}{8}\int_0^2 \left(2v^2 - \tfrac{8}{3}\right) dv = \tfrac{1}{8}\left[\tfrac{2}{3}v^3 - \tfrac{8}{3}v\right]_0^2 = 0$$

This result could have been anticipated by symmetry, since the integrand is an odd function of y and R is symmetric about the x-axis.

58. By the Extreme Value Theorem (14.7.8), f has an absolute minimum value m and an absolute maximum value M in D. Then by Property 15.2.11, $mA(D) \le \iint_D f(x, y)\,dA \le MA(D)$. Dividing through by the positive number $A(D)$, we get $m \le \dfrac{1}{A(D)}\iint_D f(x, y)\,dA \le M$. This says that the average value of f over D lies between m and M. But f is continuous on D and takes on the values m and M, and so by the Intermediate Value Theorem must take on all values between m and M. Specifically, there exists a point (x_0, y_0) in D such that $f(x_0, y_0) = \dfrac{1}{A(D)}\iint_D f(x, y)\,dA$ or equivalently $\iint_D f(x, y)\,dA = f(x_0, y_0)\,A(D)$.

59. For each r such that D_r lies within the domain, $A(D_r) = \pi r^2$, and by the Mean Value Theorem for Double Integrals there exists (x_r, y_r) in D_r such that $f(x_r, y_r) = \dfrac{1}{\pi r^2}\iint_{D_r} f(x, y)\,dA$. But $\lim\limits_{r \to 0^+} (x_r, y_r) = (a, b)$, so $\lim\limits_{r \to 0^+} \dfrac{1}{\pi r^2}\iint_{D_r} f(x, y)\,dA = \lim\limits_{r \to 0^+} f(x_r, y_r) = f(a, b)$ by the continuity of f.

60. (a) $$\iint_D \frac{1}{(x^2 + y^2)^{n/2}}\,dA = \int_0^{2\pi}\int_r^R \frac{1}{(t^2)^{n/2}} t\,dt\,d\theta = 2\pi\int_r^R t^{1-n}\,dt$$

$$= \begin{cases} \dfrac{2\pi}{2-n}t^{2-n}\Big]_r^R = \dfrac{2\pi}{2-n}\left(R^{2-n} - r^{2-n}\right) & \text{if } n \ne 2 \\ 2\pi \ln(R/r) & \text{if } n = 2 \end{cases}$$

(b) The integral in part (a) has a limit as $r \to 0^+$ for all values of n such that $2 - n > 0 \iff n < 2$.

(c) $$\iiint_E \frac{1}{(x^2 + y^2 + z^2)^{n/2}}\,dV = \int_r^R \int_0^{\pi}\int_0^{2\pi} \frac{1}{(\rho^2)^{n/2}}\rho^2 \sin\phi\,d\theta\,d\phi\,d\rho = 2\pi\int_r^R \int_0^{\pi} \rho^{2-n}\sin\phi\,d\phi\,d\rho$$

$$= \begin{cases} \dfrac{4\pi}{3-n}\rho^{3-n}\Big]_r^R = \dfrac{4\pi}{3-n}\left(R^{3-n} - r^{3-n}\right) & \text{if } n \ne 3 \\ 4\pi \ln(R/r) & \text{if } n = 3 \end{cases}$$

(d) As $r \to 0^+$, the above integral has a limit, provided that $3 - n > 0 \iff n < 3$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

PROBLEMS PLUS

1.



Let $R = \bigcup_{i=1}^{5} R_i$, where

$R_i = \{(x, y) \mid x + y \geq i + 2, x + y < i + 3, 1 \leq x \leq 3, 2 \leq y \leq 5\}$.

$\iint_R [\![x + y]\!]\, dA = \sum_{i=1}^{5} \iint_{R_i} [\![x + y]\!]\, dA = \sum_{i=1}^{5} [\![x + y]\!] \iint_{R_i} dA$, since

$[\![x + y]\!] = \text{constant} = i + 2$ for $(x, y) \in R_i$. Therefore

$$\begin{aligned} \iint_R [\![x + y]\!]\, dA &= \textstyle\sum_{i=1}^{5} (i + 2)\,[A(R_i)] \\ &= 3A(R_1) + 4A(R_2) + 5A(R_3) + 6A(R_4) + 7A(R_5) \\ &= 3\left(\tfrac{1}{2}\right) + 4\left(\tfrac{3}{2}\right) + 5(2) + 6\left(\tfrac{3}{2}\right) + 7\left(\tfrac{1}{2}\right) = 30 \end{aligned}$$

2.



Let $R = \{(x, y) \mid 0 \leq x, y \leq 1\}$. For $x, y \in R$, $\max\{x^2, y^2\} = x^2$ if $x \geq y$, and $\max\{x^2, y^2\} = y^2$ if $x \leq y$. Therefore we divide R into two regions: $R = R_1 \cup R_2$, where $R_1 = \{(x, y) \mid 0 \leq x \leq 1, 0 \leq y \leq x\}$ and $R_2 = \{(x, y) \mid 0 \leq y \leq 1, 0 \leq x \leq y\}$. Now $\max\{x^2, y^2\} = x^2$ for $(x, y) \in R_1$, and $\max\{x^2, y^2\} = y^2$ for $(x, y) \in R_2 \Rightarrow$

$$\begin{aligned} \int_0^1 \int_0^1 e^{\max\{x^2, y^2\}}\, dy\, dx &= \iint_R e^{\max\{x^2, y^2\}}\, dA = \iint_{R_1} e^{\max\{x^2, y^2\}}\, dA + \iint_{R_2} e^{\max\{x^2, y^2\}}\, dA \\ &= \int_0^1 \int_0^x e^{x^2}\, dy\, dx + \int_0^1 \int_0^y e^{y^2}\, dx\, dy = \int_0^1 x e^{x^2}\, dx + \int_0^1 y e^{y^2}\, dy = e^{x^2}\Big]_0^1 = e - 1 \end{aligned}$$

3. $f_{\text{ave}} = \dfrac{1}{b - a} \displaystyle\int_a^b f(x)\, dx = \frac{1}{1 - 0} \int_0^1 \left[\int_x^1 \cos(t^2)\, dt \right] dx$

$= \int_0^1 \int_x^1 \cos(t^2)\, dt\, dx = \int_0^1 \int_0^t \cos(t^2)\, dx\, dt$ [changing the order of integration]

$= \int_0^1 t \cos(t^2)\, dt = \frac{1}{2} \sin(t^2)\Big]_0^1 = \frac{1}{2} \sin 1$



4. Let $u = \mathbf{a} \cdot \mathbf{r}$, $v = \mathbf{b} \cdot \mathbf{r}$, $w = \mathbf{c} \cdot \mathbf{r}$, where $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$, $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$, $\mathbf{c} = \langle c_1, c_2, c_3 \rangle$. Under this change of variables, E corresponds to the rectangular box $0 \leq u \leq \alpha$, $0 \leq v \leq \beta$, $0 \leq w \leq \gamma$. So, by Formula 15.9.13,

$\displaystyle\int_0^\gamma \int_0^\beta \int_0^\alpha uvw\, du\, dv\, dw = \iiint_E (\mathbf{a} \cdot \mathbf{r})(\mathbf{b} \cdot \mathbf{r})(\mathbf{c} \cdot \mathbf{r}) \left| \frac{\partial(u, v, w)}{\partial(x, y, z)} \right| dV$. But

$$\left| \frac{\partial(u, v, w)}{\partial(x, y, z)} \right| = \left| \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \right| = |\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})| \quad \Rightarrow$$

$$\begin{aligned} \iiint_E (\mathbf{a} \cdot \mathbf{r})(\mathbf{b} \cdot \mathbf{r})(\mathbf{c} \cdot \mathbf{r})\, dV &= \frac{1}{|\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|} \int_0^\gamma \int_0^\beta \int_0^\alpha uvw\, du\, dv\, dw \\ &= \frac{1}{|\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|} \left(\frac{\alpha^2}{2} \right) \left(\frac{\beta^2}{2} \right) \left(\frac{\gamma^2}{2} \right) = \frac{(\alpha\beta\gamma)^2}{8\,|\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|} \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. Since $|xy| < 1$, except at $(1, 1)$, the formula for the sum of a geometric series gives $\dfrac{1}{1-xy} = \sum_{n=0}^{\infty} (xy)^n$, so

$$\int_0^1 \int_0^1 \tfrac{1}{1-xy}\, dx\, dy = \int_0^1 \int_0^1 \sum_{n=0}^{\infty} (xy)^n\, dx\, dy = \sum_{n=0}^{\infty} \int_0^1 \int_0^1 (xy)^n\, dx\, dy = \sum_{n=0}^{\infty} \left[\int_0^1 x^n\, dx\right]\left[\int_0^1 y^n\, dy\right]$$

$$= \sum_{n=0}^{\infty} \tfrac{1}{n+1} \cdot \tfrac{1}{n+1} = \sum_{n=0}^{\infty} \tfrac{1}{(n+1)^2} = \tfrac{1}{1^2} + \tfrac{1}{2^2} + \tfrac{1}{3^2} + \cdots = \textstyle\sum_{n=1}^{\infty} \tfrac{1}{n^2}$$

6. Let $x = \dfrac{u-v}{\sqrt{2}}$ and $y = \dfrac{u+v}{\sqrt{2}}$. We know the region of integration in the xy-plane, so to find its image in the uv-plane we get u and v in terms of x and y, and then use the methods of Section 15.9. $x + y = \dfrac{u-v}{\sqrt{2}} + \dfrac{u+v}{\sqrt{2}} = \sqrt{2}\,u$, so $u = \dfrac{x+y}{\sqrt{2}}$, and similarly $v = \dfrac{y-x}{\sqrt{2}}$. S_1 is given by $y = 0$, $0 \le x \le 1$, so from the equations derived above, the image of S_1 is S_1': $u = \frac{1}{\sqrt{2}}x$, $v = -\frac{1}{\sqrt{2}}x$, $0 \le x \le 1$, that is, $v = -u$, $0 \le u \le \frac{1}{\sqrt{2}}$. Similarly, the image of S_2 is S_2': $v = u - \sqrt{2}$, $\frac{1}{\sqrt{2}} \le u \le \sqrt{2}$, the image of S_3 is S_3': $v = \sqrt{2} - u$, $\frac{1}{\sqrt{2}} \le u \le \sqrt{2}$, and the image of S_4 is S_4': $v - u$, $0 \le u \le \frac{1}{\sqrt{2}}$.



The Jacobian of the transformation is $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \partial x/\partial u & \partial x/\partial v \\ \partial y/\partial u & \partial y/\partial v \end{vmatrix} = \begin{vmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{vmatrix} = 1$. From the diagram, we see that we must evaluate two integrals: one over the region $\left\{(u,v) \mid 0 \le u \le \frac{1}{\sqrt{2}},\ -u \le v \le u\right\}$ and the other over $\left\{(u,v) \mid \frac{1}{\sqrt{2}} \le u \le \sqrt{2},\ -\sqrt{2}+u \le v \le \sqrt{2}-u\right\}$. So

$$\int_0^1 \int_0^1 \frac{dx\, dy}{1-xy} = \int_0^{\sqrt{2}/2} \int_{-u}^{u} \frac{dv\, du}{1 - \left[\frac{1}{\sqrt{2}}(u+v)\right]\left[\frac{1}{\sqrt{2}}(u-v)\right]} + \int_{\sqrt{2}/2}^{\sqrt{2}} \int_{-\sqrt{2}+u}^{\sqrt{2}-u} \frac{dv\, du}{1 - \left[\frac{1}{\sqrt{2}}(u+v)\right]\left[\frac{1}{\sqrt{2}}(u-v)\right]}$$

$$= \int_0^{\sqrt{2}/2} \int_{-u}^{u} \frac{2\, dv\, du}{2 - u^2 + v^2} + \int_{\sqrt{2}/2}^{\sqrt{2}} \int_{-\sqrt{2}+u}^{\sqrt{2}-u} \frac{2\, dv\, du}{2 - u^2 + v^2}$$

$$= 2\left[\int_0^{\sqrt{2}/2} \frac{1}{\sqrt{2-u^2}} \left[\arctan \frac{v}{\sqrt{2-u^2}}\right]_{-u}^{u} du + \int_{\sqrt{2}/2}^{\sqrt{2}} \frac{1}{\sqrt{2-u^2}} \left[\arctan \frac{v}{\sqrt{2-u^2}}\right]_{-\sqrt{2}+u}^{\sqrt{2}-u} du\right]$$

$$= 4\left[\int_0^{\sqrt{2}/2} \frac{1}{\sqrt{2-u^2}} \arctan \frac{u}{\sqrt{2-u^2}}\, du + \int_{\sqrt{2}/2}^{\sqrt{2}} \frac{1}{\sqrt{2-u^2}} \arctan \frac{\sqrt{2}-u}{\sqrt{2-u^2}}\, du\right]$$

Now let $u = \sqrt{2}\sin\theta$, so $du = \sqrt{2}\cos\theta\, d\theta$ and the limits change to 0 and $\frac{\pi}{6}$ (in the first integral) and $\frac{\pi}{6}$ and $\frac{\pi}{2}$ (in the

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

second integral). Continuing:

$$\int_0^1\int_0^1 \frac{dx\,dy}{1-xy} = 4\left[\int_0^{\pi/6} \frac{1}{\sqrt{2-2\sin^2\theta}} \arctan\left(\frac{\sqrt{2}\sin\theta}{\sqrt{2-2\sin^2\theta}}\right)\left(\sqrt{2}\cos\theta\,d\theta\right) + \int_{\pi/6}^{\pi/2} \frac{1}{\sqrt{2-2\sin^2\theta}} \arctan\left(\frac{\sqrt{2}-\sqrt{2}\sin\theta}{\sqrt{2-2\sin^2\theta}}\right)\left(\sqrt{2}\cos\theta\,d\theta\right)\right]$$

$$= 4\left[\int_0^{\pi/6} \frac{\sqrt{2}\cos\theta}{\sqrt{2}\cos\theta} \arctan\left(\frac{\sqrt{2}\sin\theta}{\sqrt{2}\cos\theta}\right) d\theta + \int_{\pi/6}^{\pi/2} \frac{\sqrt{2}\cos\theta}{\sqrt{2}\cos\theta} \arctan\left(\frac{\sqrt{2}\,(1-\sin\theta)}{\sqrt{2}\cos\theta}\right) d\theta\right]$$

$$= 4\left[\int_0^{\pi/6} \arctan(\tan\theta)\,d\theta + \int_{\pi/6}^{\pi/2} \arctan\left(\frac{1-\sin\theta}{\cos\theta}\right) d\theta\right]$$

But (following the hint)

$$\frac{1-\sin\theta}{\cos\theta} = \frac{1-\cos\left(\frac{\pi}{2}-\theta\right)}{\sin\left(\frac{\pi}{2}-\theta\right)} = \frac{1-\left[1-2\sin^2\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)\right]}{2\sin\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)\cos\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)} \quad \text{[half-angle formulas]}$$

$$= \frac{2\sin^2\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)}{2\sin\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)\cos\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)} = \tan\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)$$

Continuing:

$$\int_0^1\int_0^1 \frac{dx\,dy}{1-xy} = 4\left[\int_0^{\pi/6} \arctan(\tan\theta)\,d\theta + \int_{\pi/6}^{\pi/2} \arctan\left(\tan\left(\tfrac{1}{2}\left(\tfrac{\pi}{2}-\theta\right)\right)\right) d\theta\right]$$

$$= 4\left[\int_0^{\pi/6} \theta\,d\theta + \int_{\pi/6}^{\pi/2} \left[\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right] d\theta\right] = 4\left(\left[\frac{\theta^2}{2}\right]_0^{\pi/6} + \left[\frac{\pi\theta}{4}-\frac{\theta^2}{4}\right]_{\pi/6}^{\pi/2}\right) = 4\left(\frac{3\pi^2}{72}\right) = \frac{\pi^2}{6}$$

7. (a) Since $|xyz| < 1$ except at $(1,1,1)$, the formula for the sum of a geometric series gives $\dfrac{1}{1-xyz} = \sum_{n=0}^{\infty} (xyz)^n$, so

$$\int_0^1\int_0^1\int_0^1 \frac{1}{1-xyz}\,dx\,dy\,dz = \int_0^1\int_0^1\int_0^1 \sum_{n=0}^{\infty} (xyz)^n\,dx\,dy\,dz = \sum_{n=0}^{\infty}\int_0^1\int_0^1\int_0^1 (xyz)^n\,dx\,dy\,dz$$

$$= \sum_{n=0}^{\infty} \left[\int_0^1 x^n\,dx\right]\left[\int_0^1 y^n\,dy\right]\left[\int_0^1 z^n\,dz\right] = \sum_{n=0}^{\infty} \frac{1}{n+1}\cdot\frac{1}{n+1}\cdot\frac{1}{n+1}$$

$$= \sum_{n=0}^{\infty} \frac{1}{(n+1)^3} = \frac{1}{1^3}+\frac{1}{2^3}+\frac{1}{3^3}+\cdots = \sum_{n=1}^{\infty}\frac{1}{n^3}$$

(b) Since $|-xyz| < 1$, except at $(1,1,1)$, the formula for the sum of a geometric series gives $\dfrac{1}{1+xyz} = \sum_{n=0}^{\infty} (-xyz)^n$, so

$$\int_0^1\int_0^1\int_0^1 \frac{1}{1+xyz}\,dx\,dy\,dz = \int_0^1\int_0^1\int_0^1 \sum_{n=0}^{\infty} (-xyz)^n\,dx\,dy\,dz = \sum_{n=0}^{\infty}\int_0^1\int_0^1\int_0^1 (-xyz)^n\,dx\,dy\,dz$$

$$= \sum_{n=0}^{\infty} (-1)^n \left[\int_0^1 x^n\,dx\right]\left[\int_0^1 y^n\,dy\right]\left[\int_0^1 z^n\,dz\right] = \sum_{n=0}^{\infty} (-1)^n \frac{1}{n+1}\cdot\frac{1}{n+1}\cdot\frac{1}{n+1}$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{(n+1)^3} = \frac{1}{1^3}-\frac{1}{2^3}+\frac{1}{3^3}-\cdots = \sum_{n=0}^{\infty}\frac{(-1)^{n-1}}{n^3}$$

To evaluate this sum, we first write out a few terms: $s = 1-\dfrac{1}{2^3}+\dfrac{1}{3^3}-\dfrac{1}{4^3}+\dfrac{1}{5^3}-\dfrac{1}{6^3} \approx 0.8998$. Notice that $a_7 = \dfrac{1}{7^3} < 0.003$. By the Alternating Series Estimation Theorem from Section 11.5, we have $|s - s_6| \le a_7 < 0.003$. This error of 0.003 will not affect the second decimal place, so we have $s \approx 0.90$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

8. $\displaystyle\int_0^\infty \frac{\arctan \pi x - \arctan x}{x}\,dx = \int_0^\infty \left[\frac{\arctan yx}{x}\right]_{y=1}^{y=\pi} dx = \int_0^\infty \int_1^\pi \frac{1}{1+y^2x^2}\,dy\,dx = \int_1^\pi \int_0^\infty \frac{1}{1+y^2x^2}\,dx\,dy$

$$= \int_1^\pi \lim_{t\to\infty} \left[\frac{\arctan yx}{y}\right]_{x=0}^{x=t} dy = \int_1^\pi \frac{\pi}{2y}\,dy = \frac{\pi}{2}\left[\ln y\right]_1^\pi = \frac{\pi}{2}\ln \pi$$

9. (a) $x = r\cos\theta,\ y = r\sin\theta,\ z = z$. Then $\dfrac{\partial u}{\partial r} = \dfrac{\partial u}{\partial x}\dfrac{\partial x}{\partial r} + \dfrac{\partial u}{\partial y}\dfrac{\partial y}{\partial r} + \dfrac{\partial u}{\partial z}\dfrac{\partial z}{\partial r} = \dfrac{\partial u}{\partial x}\cos\theta + \dfrac{\partial u}{\partial y}\sin\theta$ and

$$\frac{\partial^2 u}{\partial r^2} = \cos\theta\left[\frac{\partial^2 u}{\partial x^2}\frac{\partial x}{\partial r} + \frac{\partial^2 u}{\partial y\,\partial x}\frac{\partial y}{\partial r} + \frac{\partial^2 u}{\partial z\,\partial x}\frac{\partial z}{\partial r}\right] + \sin\theta\left[\frac{\partial^2 u}{\partial y^2}\frac{\partial y}{\partial r} + \frac{\partial^2 u}{\partial x\,\partial y}\frac{\partial x}{\partial r} + \frac{\partial^2 u}{\partial z\,\partial y}\frac{\partial z}{\partial r}\right]$$

$$= \frac{\partial^2 u}{\partial x^2}\cos^2\theta + \frac{\partial^2 u}{\partial y^2}\sin^2\theta + 2\frac{\partial^2 u}{\partial y\,\partial x}\cos\theta\,\sin\theta$$

Similarly $\dfrac{\partial u}{\partial \theta} = -\dfrac{\partial u}{\partial x}r\sin\theta + \dfrac{\partial u}{\partial y}r\cos\theta$ and

$\dfrac{\partial^2 u}{\partial \theta^2} = \dfrac{\partial^2 u}{\partial x^2}r^2\sin^2\theta + \dfrac{\partial^2 u}{\partial y^2}r^2\cos^2\theta - 2\dfrac{\partial^2 u}{\partial y\,\partial x}r^2\sin\theta\,\cos\theta - \dfrac{\partial u}{\partial x}r\cos\theta - \dfrac{\partial u}{\partial y}r\sin\theta$. So

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r}\frac{\partial u}{\partial r} + \frac{1}{r^2}\frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2} = \frac{\partial^2 u}{\partial x^2}\cos^2\theta + \frac{\partial^2 u}{\partial y^2}\sin^2\theta + 2\frac{\partial^2 u}{\partial y\,\partial x}\cos\theta\,\sin\theta + \frac{\partial u}{\partial x}\frac{\cos\theta}{r} + \frac{\partial u}{\partial y}\frac{\sin\theta}{r}$$

$$+ \frac{\partial^2 u}{\partial x^2}\sin^2\theta + \frac{\partial^2 u}{\partial y^2}\cos^2\theta - 2\frac{\partial^2 u}{\partial y\,\partial x}\sin\theta\,\cos\theta$$

$$- \frac{\partial u}{\partial x}\frac{\cos\theta}{r} - \frac{\partial u}{\partial y}\frac{\sin\theta}{r} + \frac{\partial^2 u}{\partial z^2}$$

$$= \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2}$$

(b) $x = \rho\sin\phi\cos\theta,\ y = \rho\sin\phi\sin\theta,\ z = \rho\cos\phi$. Then

$\dfrac{\partial u}{\partial \rho} = \dfrac{\partial u}{\partial x}\dfrac{\partial x}{\partial \rho} + \dfrac{\partial u}{\partial y}\dfrac{\partial y}{\partial \rho} + \dfrac{\partial u}{\partial z}\dfrac{\partial z}{\partial \rho} = \dfrac{\partial u}{\partial x}\sin\phi\,\cos\theta + \dfrac{\partial u}{\partial y}\sin\phi\,\sin\theta + \dfrac{\partial u}{\partial z}\cos\phi$, and

$$\frac{\partial^2 u}{\partial \rho^2} = \sin\phi\cos\theta\left[\frac{\partial^2 u}{\partial x^2}\frac{\partial x}{\partial \rho} + \frac{\partial^2 u}{\partial y\,\partial x}\frac{\partial y}{\partial \rho} + \frac{\partial^2 u}{\partial z\,\partial x}\frac{\partial z}{\partial \rho}\right]$$

$$+ \sin\phi\sin\theta\left[\frac{\partial^2 u}{\partial y^2}\frac{\partial y}{\partial \rho} + \frac{\partial^2 u}{\partial x\,\partial y}\frac{\partial x}{\partial \rho} + \frac{\partial^2 u}{\partial z\,\partial y}\frac{\partial z}{\partial \rho}\right]$$

$$+ \cos\phi\left[\frac{\partial^2 u}{\partial z^2}\frac{\partial z}{\partial \rho} + \frac{\partial^2 u}{\partial x\,\partial z}\frac{\partial x}{\partial \rho} + \frac{\partial^2 u}{\partial y\,\partial z}\frac{\partial y}{\partial \rho}\right]$$

$$= 2\frac{\partial^2 u}{\partial y\,\partial x}\sin^2\phi\,\sin\theta\,\cos\theta + 2\frac{\partial^2 u}{\partial z\,\partial x}\sin\phi\,\cos\phi\,\cos\theta + 2\frac{\partial^2 u}{\partial y\,\partial z}\sin\phi\,\cos\phi\,\sin\theta$$

$$+ \frac{\partial^2 u}{\partial x^2}\sin^2\phi\cos^2\theta + \frac{\partial^2 u}{\partial y^2}\sin^2\phi\sin^2\theta + \frac{\partial^2 u}{\partial z^2}\cos^2\phi$$

Similarly $\dfrac{\partial u}{\partial \phi} = \dfrac{\partial u}{\partial x}\rho\cos\phi\cos\theta + \dfrac{\partial u}{\partial y}\rho\cos\phi\sin\theta - \dfrac{\partial u}{\partial z}\rho\sin\phi$, and

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\frac{\partial^2 u}{\partial \phi^2} = 2\frac{\partial^2 u}{\partial y\,\partial x}\rho^2\cos^2\phi\,\sin\theta\,\cos\theta - 2\frac{\partial^2 u}{\partial x\,\partial z}\rho^2\sin\phi\,\cos\phi\,\cos\theta$$
$$- 2\frac{\partial^2 u}{\partial y\,\partial z}\rho^2\sin\phi\,\cos\phi\,\sin\theta + \frac{\partial^2 u}{\partial x^2}\rho^2\cos^2\phi\,\cos^2\theta + \frac{\partial^2 u}{\partial y^2}\rho^2\cos^2\phi\,\sin^2\theta$$
$$+ \frac{\partial^2 u}{\partial z^2}\rho^2\sin^2\phi - \frac{\partial u}{\partial x}\rho\sin\phi\,\cos\theta - \frac{\partial u}{\partial y}\rho\sin\phi\,\sin\theta - \frac{\partial u}{\partial z}\rho\cos\phi$$

And $\dfrac{\partial u}{\partial \theta} = -\dfrac{\partial u}{\partial x}\rho\sin\phi\sin\theta + \dfrac{\partial u}{\partial y}\rho\sin\phi\cos\theta$, while

$$\frac{\partial^2 u}{\partial \theta^2} = -2\frac{\partial^2 u}{\partial y\,\partial x}\rho^2\sin^2\phi\,\cos\theta\,\sin\theta + \frac{\partial^2 u}{\partial x^2}\rho^2\sin^2\phi\,\sin^2\theta$$
$$+ \frac{\partial^2 u}{\partial y^2}\rho^2\sin^2\phi\cos^2\theta - \frac{\partial u}{\partial x}\rho\sin\phi\,\cos\theta - \frac{\partial u}{\partial y}\rho\sin\phi\,\sin\theta$$

Therefore

$$\frac{\partial^2 u}{\partial \rho^2} + \frac{2}{\rho}\frac{\partial u}{\partial \rho} + \frac{\cot\phi}{\rho^2}\frac{\partial u}{\partial \phi} + \frac{1}{\rho^2}\frac{\partial^2 u}{\partial \phi^2} + \frac{1}{\rho^2\sin^2\phi}\frac{\partial^2 u}{\partial \theta^2}$$
$$= \frac{\partial^2 u}{\partial x^2}\left[(\sin^2\phi\,\cos^2\theta) + (\cos^2\phi\,\cos^2\theta) + \sin^2\theta\right]$$
$$+ \frac{\partial^2 u}{\partial y^2}\left[(\sin^2\phi\,\sin^2\theta) + (\cos^2\phi\,\sin^2\theta) + \cos^2\theta\right] + \frac{\partial^2 u}{\partial z^2}\left[\cos^2\phi + \sin^2\phi\right]$$
$$+ \frac{\partial u}{\partial x}\left[\frac{2\sin^2\phi\,\cos\theta + \cos^2\phi\,\cos\theta - \sin^2\phi\,\cos\theta - \cos\theta}{\rho\sin\phi}\right]$$
$$+ \frac{\partial u}{\partial y}\left[\frac{2\sin^2\phi\,\sin\theta + \cos^2\phi\,\sin\theta - \sin^2\phi\,\sin\theta - \sin\theta}{\rho\sin\phi}\right]$$

But $2\sin^2\phi\cos\theta + \cos^2\phi\cos\theta - \sin^2\phi\cos\theta - \cos\theta = (\sin^2\phi + \cos^2\phi - 1)\,\cos\theta = 0$ and similarly the coefficient of $\partial u/\partial y$ is 0. Also $\sin^2\phi\cos^2\theta + \cos^2\phi\cos^2\theta + \sin^2\theta = \cos^2\theta\,(\sin^2\phi + \cos^2\phi) + \sin^2\theta = 1$, and similarly the coefficient of $\partial^2 u/\partial y^2$ is 1. So Laplace's Equation in spherical coordinates is as stated.

10. (a) Consider a polar division of the disk, similar to that in Figure 15.3.4, where $0 = \theta_0 < \theta_1 < \theta_2 < \cdots < \theta_n = 2\pi$, $0 = r_1 < r_2 < \cdots < r_m = R$, and where the polar subrectangle R_{ij}, as well as r_i^*, θ_j^*, Δr and $\Delta\theta$ are the same as in that figure. Thus $\Delta A_i = r_i^*\,\Delta r\,\Delta\theta$. The mass of R_{ij} is $\rho\,\Delta A_i$, and its distance from m is $s_{ij} \approx \sqrt{(r_i^*)^2 + d^2}$. According to Newton's Law of Gravitation, the force of attraction experienced by m due to this polar subrectangle is in the direction from m towards R_{ij} and has magnitude $\dfrac{Gm\rho\,\Delta A_i}{s_{ij}^2}$. The symmetry of the lamina with respect to the x- and y-axes and the position of m are such that all horizontal components of the gravitational force cancel, so that the total force is simply in the z-direction. Thus, we need only be concerned with the components of this vertical force; that is, $\dfrac{Gm\rho\,\Delta A_i}{s_{ij}^2}\sin\alpha$, where α is the angle between the origin, r_i^* and the mass m. Thus $\sin\alpha = \dfrac{d}{s_{ij}}$ and the previous result becomes

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$\dfrac{Gm\rho d\,\Delta A_i}{s_{ij}^3}$. The total attractive force is just the Riemann sum $\displaystyle\sum_{i=1}^{m}\sum_{j=1}^{n}\frac{Gm\rho d\,\Delta A_i}{s_{ij}^3}=\sum_{i=1}^{m}\sum_{j=1}^{n}\frac{Gm\rho d(r_i^*)\,\Delta r\,\Delta\theta}{\left[(r_i^*)^2+d^2\right]^{3/2}}$

which becomes $\displaystyle\int_0^R\int_0^{2\pi}\frac{Gm\rho d}{(r^2+d^2)^{3/2}}r\,d\theta\,dr$ as $m\to\infty$ and $n\to\infty$. Therefore,

$$F=2\pi Gm\rho d\int_0^R\frac{r}{(r^2+d^2)^{3/2}}\,dr=2\pi Gm\rho d\left[-\frac{1}{\sqrt{r^2+d^2}}\right]_0^R=2\pi Gm\rho d\left(\frac{1}{d}-\frac{1}{\sqrt{R^2+d^2}}\right)$$

(b) This is just the result of part (a) in the limit as $R\to\infty$. In this case $\dfrac{1}{\sqrt{R^2+d^2}}\to 0$, and we are left with

$$F=2\pi Gm\rho d\left(\frac{1}{d}-0\right)=2\pi Gm\rho.$$

11. $\int_0^x\int_0^y\int_0^z f(t)\,dt\,dz\,dy=\iiint_E f(t)\,dV$, where

$E=\{(t,z,y)\mid 0\le t\le z,0\le z\le y,0\le y\le x\}$.

If we let D be the projection of E on the yt-plane then

$D=\{(y,t)\mid 0\le t\le x,t\le y\le x\}$. And we see from the diagram

that $E=\{(t,z,y)\mid t\le z\le y,t\le y\le x,0\le t\le x\}$. So



$$\begin{aligned}\int_0^x\int_0^y\int_0^z f(t)\,dt\,dz\,dy&=\int_0^x\int_t^x\int_t^y f(t)\,dz\,dy\,dt=\int_0^x\left[\int_t^x(y-t)\,f(t)\,dy\right]dt\\&=\int_0^x\left[\left(\tfrac12y^2-ty\right)f(t)\right]_{y=t}^{y=x}dt=\int_0^x\left[\tfrac12x^2-tx-\tfrac12t^2+t^2\right]f(t)\,dt\\&=\int_0^x\left[\tfrac12x^2-tx+\tfrac12t^2\right]f(t)\,dt=\int_0^x\left(\tfrac12x^2-2tx+t^2\right)f(t)\,dt\\&=\tfrac12\int_0^x(x-t)^2\,f(t)\,dt\end{aligned}$$

12. $n^{-2}\displaystyle\sum_{i=1}^{n}\sum_{j=1}^{n^2}\frac{1}{\sqrt{n^2+ni+j}}=\sum_{i=1}^{n}\sum_{j=1}^{n^2}\frac{1}{\frac1n\sqrt{n^2+ni+j}}\cdot\frac{1}{n^3}=\sum_{i=1}^{n}\sum_{j=1}^{n^2}\frac{1}{\sqrt{1+\frac{i}{n}+\frac{j}{n^2}}}\cdot\frac{1}{n^3}$ can be considered a double

Riemann sum of the function $f(x,y)=\dfrac{1}{\sqrt{1+x+y}}$ where the square region $R=\{(x,y)\mid 0\le x\le 1,0\le y\le 1\}$ is

divided into subrectangles by dividing the interval $[0,1]$ on the x-axis into n subintervals, each of width $\frac1n$, and $[0,1]$ on the y-axis is divided into n^2 subintervals, each of width $\frac{1}{n^2}$. Then the area of each subrectangle is $\Delta A=\frac{1}{n^3}$, and if we take the upper right corners of the subrectangles as sample points, we have $(x_{ij}^*,y_{ij}^*)=\left(\frac{i}{n},\frac{j}{n^2}\right)$. Finally, note that $n^2\to\infty$ as $n\to\infty$, so

$$\lim_{n\to\infty}n^{-2}\sum_{i=1}^{n}\sum_{j=1}^{n^2}\frac{1}{\sqrt{n^2+ni+j}}=\lim_{n,n^2\to\infty}\sum_{i=1}^{n}\sum_{j=1}^{n^2}\frac{1}{\sqrt{1+\frac{i}{n}+\frac{j}{n^2}}}\cdot\frac{1}{n^3}=\lim_{n,n^2\to\infty}\sum_{i=1}^{n}\sum_{j=1}^{n^2}f(x_{ij}^*,y_{ij}^*)\,\Delta A$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

But by Definition 15.1.5 this is equal to $\iint_R f(x,y)\,dA$, so

$$\lim_{n\to\infty} n^{-2} \sum_{i=1}^{n} \sum_{j=1}^{n^2} \frac{1}{\sqrt{n^2+ni+j}} = \iint_R f(x,y)\,dA = \int_0^1 \int_0^1 \frac{1}{\sqrt{1+x+y}}\,dy\,dx$$

$$= \int_0^1 \left[2(1+x+y)^{1/2}\right]_{y=0}^{y=1} dx = 2\int_0^1 \left(\sqrt{2+x} - \sqrt{1+x}\right) dx$$

$$= 2\left[\tfrac{2}{3}(2+x)^{3/2} - \tfrac{2}{3}(1+x)^{3/2}\right]_0^1 = \tfrac{4}{3}(3^{3/2} - 2^{3/2} - 2^{3/2} + 1)$$

$$= \tfrac{4}{3}(3\sqrt{3} - 4\sqrt{2} + 1) = 4\sqrt{3} - \tfrac{16}{3}\sqrt{2} + \tfrac{4}{3}$$

13. The volume is $V = \iiint_R dV$ where R is the solid region given. From Exercise 15.9.21(a), the transformation $x = au$, $y = bv$, $z = cw$ maps the unit ball $u^2 + v^2 + w^2 \le 1$ to the solid ellipsoid $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} + \dfrac{z^2}{c^2} \le 1$ with $\dfrac{\partial(x,y,z)}{\partial(u,v,w)} = abc$. The same transformation maps the plane $u + v + w = 1$ to $\dfrac{x}{a} + \dfrac{y}{b} + \dfrac{z}{c} = 1$. Thus the region R in xyz-space corresponds to the region S in uvw-space consisting of the smaller piece of the unit ball cut off by the plane $u + v + w = 1$, a "cap of a sphere" (see the figure).



We will need to compute the volume of S, but first consider the general case where a horizontal plane slices the upper portion of a sphere of radius r to produce a cap of height h. We use spherical coordinates. From the figure, a line through the origin at angle ϕ from the z-axis intersects the plane when $\cos\phi = (r-h)/a \;\Rightarrow\; a = (r-h)/\cos\phi$, and the line passes through the outer rim of the cap when $a = r \;\Rightarrow\; \cos\phi = (r-h)/r \;\Rightarrow\; \phi = \cos^{-1}\left((r-h)/r\right)$. Thus the cap is described by $\left\{(\rho,\theta,\phi) \mid (r-h)/\cos\phi \le \rho \le r, 0 \le \theta \le 2\pi, 0 \le \phi \le \cos^{-1}\left((r-h)/r\right)\right\}$ and its volume is



$$V = \int_0^{2\pi} \int_0^{\cos^{-1}((r-h)/r)} \int_{(r-h)/\cos\phi}^{r} \rho^2 \sin\phi\,d\rho\,d\phi\,d\theta$$

$$= \int_0^{2\pi} \int_0^{\cos^{-1}((r-h)/r)} \left[\tfrac{1}{3}\rho^3 \sin\phi\right]_{\rho=(r-h)/\cos\phi}^{\rho=r} d\phi\,d\theta$$

$$= \frac{1}{3}\int_0^{2\pi} \int_0^{\cos^{-1}((r-h)/r)} \left[r^3\sin\phi - \frac{(r-h)^3}{\cos^3\phi}\sin\phi\right] d\phi\,d\theta$$

$$= \tfrac{1}{3}\int_0^{2\pi} \left[-r^3\cos\phi - \tfrac{1}{2}(r-h)^3\cos^{-2}\phi\right]_{\phi=0}^{\phi=\cos^{-1}((r-h)/r)} d\theta$$

$$= \frac{1}{3}\int_0^{2\pi} \left[-r^3\left(\frac{r-h}{r}\right) - \frac{1}{2}(r-h)^3\left(\frac{r-h}{r}\right)^{-2} + r^3 + \frac{1}{2}(r-h)^3\right] d\theta$$

$$= \tfrac{1}{3}\int_0^{2\pi} (\tfrac{3}{2}rh^2 - \tfrac{1}{2}h^3)\,d\theta = \tfrac{1}{3}(\tfrac{3}{2}rh^2 - \tfrac{1}{2}h^3)(2\pi) = \pi h^2(r - \tfrac{1}{3}h)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(This volume can also be computed by treating the cap as a solid of revolution and using the single variable disk method; see Exercise 5.2.49 [ET 6.2.49].)

To determine the height h of the cap cut from the unit ball by the plane $u + v + w = 1$, note that the line $u = v = w$ passes through the origin with direction vector $\langle 1, 1, 1 \rangle$ which is perpendicular to the plane. Therefore this line coincides with a radius of the sphere that passes through the center of the cap and h is measured along this line. The line intersects the plane at $\left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\right)$ and the sphere at $\left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right)$. (See the figure.)



The distance between these points is $h = \sqrt{3\left(\frac{1}{\sqrt{3}} - \frac{1}{3}\right)^2} = \sqrt{3}\left(\frac{1}{\sqrt{3}} - \frac{1}{3}\right) = 1 - \frac{1}{\sqrt{3}}$. Thus the volume of R is

$$V = \iiint_R dV = \iiint_S \left|\frac{\partial(x, y, z)}{\partial(u, v, w)}\right| dV = abc \iiint_S dV = abc\, V(S)$$

$$= abc \cdot \pi h^2 (r - \tfrac{1}{3}h) = abc \cdot \pi\left(1 - \tfrac{1}{\sqrt{3}}\right)^2 \left[1 - \tfrac{1}{3}\left(1 - \tfrac{1}{\sqrt{3}}\right)\right]$$

$$= abc\pi \left(\tfrac{4}{3} - \tfrac{2}{\sqrt{3}}\right)\left(\tfrac{2}{3} + \tfrac{1}{3\sqrt{3}}\right) = abc\pi \left(\tfrac{2}{3} - \tfrac{8}{9\sqrt{3}}\right) \approx 0.482abc$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16 □ VECTOR CALCULUS

16.1 Vector Fields

1. $\mathbf{F}(x, y) = 0.3\,\mathbf{i} - 0.4\,\mathbf{j}$

All vectors in this field are identical, with length 0.5 and parallel to $\langle 3, -4 \rangle$.



2. $\mathbf{F}(x, y) = \frac{1}{2}x\,\mathbf{i} + y\,\mathbf{j}$

The length of the vector $\frac{1}{2}x\,\mathbf{i} + y\,\mathbf{j}$ is $\sqrt{\frac{1}{4}x^2 + y^2}$.

Vectors point roughly away from the origin and vectors farther from the origin are longer.



3. $\mathbf{F}(x, y) = -\frac{1}{2}\,\mathbf{i} + (y - x)\,\mathbf{j}$

The length of the vector $-\frac{1}{2}\,\mathbf{i} + (y - x)\,\mathbf{j}$ is $\sqrt{\frac{1}{4} + (y - x)^2}$. Vectors along the line $y = x$ are horizontal with length $\frac{1}{2}$.



4. $\mathbf{F}(x, y) = y\,\mathbf{i} + (x + y)\,\mathbf{j}$

The length of the vector $y\,\mathbf{i} + (x + y)\,\mathbf{j}$ is $\sqrt{y^2 + (x + y)^2}$. Vectors along the x-axis are vertical, and vectors along the line $y = -x$ are horizontal with length $|y|$.



5. $\mathbf{F}(x, y) = \dfrac{y\,\mathbf{i} + x\,\mathbf{j}}{\sqrt{x^2 + y^2}}$

The length of the vector $\dfrac{y\,\mathbf{i} + x\,\mathbf{j}}{\sqrt{x^2 + y^2}}$ is 1.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

6. $\mathbf{F}(x, y) = \dfrac{y\,\mathbf{i} - x\,\mathbf{j}}{\sqrt{x^2 + y^2}}$

All the vectors $\mathbf{F}(x, y)$ are unit vectors tangent to circles centered at the origin with radius $\sqrt{x^2 + y^2}$.



7. $\mathbf{F}(x, y) = \mathbf{i}$

All vectors in this field are identical, with length 1 and pointing in the direction of the positive x-axis.



8. $\mathbf{F}(x, y, z) = z\,\mathbf{i}$

At each point (x, y, z), $\mathbf{F}(x, y, z)$ is a vector of length $|z|$. For $z > 0$, all point in the direction of the positive x-axis, while for $z < 0$, all are in the direction of the negative x-axis. In each plane $z = k$, all the vectors are identical.



9. $\mathbf{F}(x, y, z) = -y\,\mathbf{i}$

At each point (x, y, z), $\mathbf{F}(x, y, z)$ is a vector of length $|y|$. For $y > 0$, all point in the direction of the negative x-axis, while for $y < 0$, all are in the direction of the positive x-axis. In each plane $y = k$, all the vectors are identical.



10. $\mathbf{F}(x, y, z) = \mathbf{i} + \mathbf{k}$

All vectors in this field have length $\sqrt{2}$ and point in the same direction, parallel to the xz-plane.



11. $\mathbf{F}(x, y) = \langle x, -y \rangle$ corresponds to graph IV. In the first quadrant all the vectors have positive x-components and negative y-components, in the second quadrant all vectors have negative x- and y-components, in the third quadrant all vectors have negative x-components and positive y-components, and in the fourth quadrant all vectors have positive x- and y-components. In addition, the vectors get shorter as we approach the origin.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $\mathbf{F}(x, y) = \langle y, x - y \rangle$ corresponds to graph III. All vectors in quadrants I and II have positive x-components while all vectors in quadrants III and IV have negative x-components. In addition, vectors along the line $y = x$ are horizontal, and vectors get shorter as we approach the origin.

13. $\mathbf{F}(x, y) = \langle y, y + 2 \rangle$ corresponds to graph I. As in Exercise 12, all vectors in quadrants I and II have positive x-components while all vectors in quadrants III and IV have negative x-components. Vectors along the line $y = -2$ are horizontal, and the vectors are independent of x (vectors along horizontal lines are identical).

14. $\mathbf{F}(x, y) = \langle \cos(x + y), x \rangle$ corresponds to graph II. All vectors in quadrants I and IV have positive y-components while all vectors in quadrants II and III have negative y-components. Also, the y-components of vectors along any vertical line remain constant while the x-component oscillates.

15. $\mathbf{F}(x, y, z) = \mathbf{i} + 2\,\mathbf{j} + 3\,\mathbf{k}$ corresponds to graph IV, since all vectors have identical length and direction.

16. $\mathbf{F}(x, y, z) = \mathbf{i} + 2\,\mathbf{j} + z\,\mathbf{k}$ corresponds to graph I, since the horizontal vector components remain constant, but the vectors above the xy-plane point generally upward while the vectors below the xy-plane point generally downward.

17. $\mathbf{F}(x, y, z) = x\,\mathbf{i} + y\,\mathbf{j} + 3\,\mathbf{k}$ corresponds to graph III; the projection of each vector onto the xy-plane is $x\,\mathbf{i} + y\,\mathbf{j}$, which points away from the origin, and the vectors point generally upward because their z-components are all 3.

18. $\mathbf{F}(x, y, z) = x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$ corresponds to graph II; each vector $\mathbf{F}(x, y, z)$ has the same length and direction as the position vector of the point (x, y, z), and therefore the vectors all point directly away from the origin.

19.



The vector field seems to have very short vectors near the line $y = 2x$. For $\mathbf{F}(x, y) = \langle 0, 0 \rangle$ we must have $y^2 - 2xy = 0$ and $3xy - 6x^2 = 0$. The first equation holds if $y = 0$ or $y = 2x$, and the second holds if $x = 0$ or $y = 2x$. So both equations hold [and thus $\mathbf{F}(x, y) = \mathbf{0}$] along the line $y = 2x$.

20.



From the graph, it appears that all of the vectors in the field lie on lines through the origin, and that the vectors have very small magnitudes near the circle $|\mathbf{x}| = 2$ and near the origin. Note that $\mathbf{F}(\mathbf{x}) = \mathbf{0} \Leftrightarrow r(r - 2) = 0 \Leftrightarrow r = 0$ or 2, so as we suspected, $\mathbf{F}(\mathbf{x}) = \mathbf{0}$ for $|\mathbf{x}| = 2$ and for $|\mathbf{x}| = 0$. Note that where $r^2 - r < 0$, the vectors point towards the origin, and where $r^2 - r > 0$, they point away from the origin.

21. $f(x, y) = y \sin(xy) \Rightarrow$

$$\begin{aligned}\nabla f(x, y) &= f_x(x, y)\,\mathbf{i} + f_y(x, y)\,\mathbf{j} = (y \cos(xy) \cdot y)\,\mathbf{i} + [y \cdot x \cos(xy) + \sin(xy) \cdot 1]\,\mathbf{j} \\ &= y^2 \cos(xy)\,\mathbf{i} + [xy \cos(xy) + \sin(xy)]\,\mathbf{j}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

22. $f(s,t) = \sqrt{2s+3t} \;\Rightarrow$

$$\nabla f(s,t) = f_s(s,t)\,\mathbf{i} + f_t(s,t)\,\mathbf{j} = \left[\tfrac{1}{2}(2s+3t)^{-1/2}\cdot 2\right]\mathbf{i} + \left[\tfrac{1}{2}(2s+3t)^{-1/2}\cdot 3\right]\mathbf{j} = \frac{1}{\sqrt{2s+3t}}\,\mathbf{i} + \frac{3}{2\sqrt{2s+3t}}\,\mathbf{j}$$

23. $f(x,y,z) = \sqrt{x^2+y^2+z^2} \;\Rightarrow$

$$\begin{aligned}\nabla f(x,y,z) &= f_x(x,y,z)\,\mathbf{i} + f_y(x,y,z)\,\mathbf{j} + f_z(x,y,z)\,\mathbf{k}\\ &= \tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2x)\,\mathbf{i} + \tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2y)\,\mathbf{j} + \tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2z)\,\mathbf{k}\\ &= \frac{x}{\sqrt{x^2+y^2+z^2}}\,\mathbf{i} + \frac{y}{\sqrt{x^2+y^2+z^2}}\,\mathbf{j} + \frac{z}{\sqrt{x^2+y^2+z^2}}\,\mathbf{k}\end{aligned}$$

24. $f(x,y,z) = x^2 y e^{y/z} \;\Rightarrow$

$$\begin{aligned}\nabla f(x,y,z) &= f_x(x,y,z)\,\mathbf{i} + f_y(x,y,z)\,\mathbf{j} + f_z(x,y,z)\,\mathbf{k}\\ &= 2xye^{y/z}\,\mathbf{i} + x^2\left[y\cdot e^{yz}(1/z) + e^{y/z}\cdot 1\right]\mathbf{j} + \left[x^2 y e^{y/z}(-y/z^2)\right]\mathbf{k}\\ &= 2xye^{y/z}\,\mathbf{i} + x^2 e^{y/z}\left(\frac{y}{z}+1\right)\mathbf{j} - \frac{x^2y^2}{z^2}e^{y/z}\,\mathbf{k}\end{aligned}$$

25. $f(x,y) = \tfrac{1}{2}(x-y)^2 \;\Rightarrow$

$\nabla f(x,y) = (x-y)(1)\,\mathbf{i} + (x-y)(-1)\,\mathbf{j} = (x-y)\,\mathbf{i} + (y-x)\,\mathbf{j}$.

The length of $\nabla f(x,y)$ is $\sqrt{(x-y)^2+(y-x)^2} = \sqrt{2}\,|x-y|$.

The vectors are $\mathbf{0}$ along the line $y = x$. Elsewhere the vectors point away from the line $y = x$ with length that increases as the distance from the line increases.



26. $f(x,y) = \tfrac{1}{2}(x^2-y^2) \;\Rightarrow\; \nabla f(x,y) = x\,\mathbf{i} - y\,\mathbf{j}$.

The length of $\nabla f(x,y)$ is $\sqrt{x^2+y^2}$. The lengths of the vectors increase as the distance from the origin increases, and the terminal point of each vector lies on the x-axis.



27. We graph $\nabla f(x,y) = \dfrac{2x}{1+x^2+2y^2}\,\mathbf{i} + \dfrac{4y}{1+x^2+2y^2}\,\mathbf{j}$ along with a contour map of f.

The graph shows that the gradient vectors are perpendicular to the level curves. Also, the gradient vectors point in the direction in which f is increasing and are longer where the level curves are closer together.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. We graph $\nabla f(x, y) = -\sin x\,\mathbf{i} - 2\cos y\,\mathbf{j}$ along with a contour map of f.



The graph shows that the gradient vectors are perpendicular to the level curves. Also, the gradient vectors point in the direction in which f is increasing and are longer where the level curves are closer together.

29. $f(x, y) = x^2 + y^2 \;\Rightarrow\; \nabla f(x, y) = 2x\,\mathbf{i} + 2y\,\mathbf{j}$. Thus, each vector $\nabla f(x, y)$ has the same direction and twice the length of the position vector of the point (x, y), so the vectors all point directly away from the origin and their lengths increase as we move away from the origin. Hence, ∇f is graph III.

30. $f(x, y) = x(x + y) = x^2 + xy \;\Rightarrow\; \nabla f(x, y) = (2x + y)\,\mathbf{i} + x\,\mathbf{j}$. The y-component of each vector is x, so the vectors point upward in quadrants I and IV and downward in quadrants II and III. Also, the x-component of each vector is 0 along the line $y = -2x$ so the vectors are vertical there. Thus, ∇f is graph IV.

31. $f(x, y) = (x + y)^2 \;\Rightarrow\; \nabla f(x, y) = 2(x + y)\,\mathbf{i} + 2(x + y)\,\mathbf{j}$. The x- and y-components of each vector are equal, so all vectors are parallel to the line $y = x$. The vectors are $\mathbf{0}$ along the line $y = -x$ and their length increases as the distance from this line increases. Thus, ∇f is graph II.

32. $f(x, y) = \sin\sqrt{x^2 + y^2} \;\Rightarrow$

$$\nabla f(x, y) = \left[\cos\sqrt{x^2 + y^2} \cdot \tfrac{1}{2}(x^2 + y^2)^{-1/2}(2x)\right]\mathbf{i} + \left[\cos\sqrt{x^2 + y^2} \cdot \tfrac{1}{2}(x^2 + y^2)^{-1/2}(2y)\right]\mathbf{j}$$

$$= \frac{\cos\sqrt{x^2 + y^2}}{\sqrt{x^2 + y^2}}\,x\,\mathbf{i} + \frac{\cos\sqrt{x^2 + y^2}}{\sqrt{x^2 + y^2}}\,y\,\mathbf{j} \text{ or } \frac{\cos\sqrt{x^2 + y^2}}{\sqrt{x^2 + y^2}}\,(x\,\mathbf{i} + y\,\mathbf{j})$$

Thus each vector is a scalar multiple of its position vector, so the vectors point toward or away from the origin with length that changes in a periodic fashion as we move away from the origin. ∇f is graph I.

33. At $t = 3$ the particle is at $(2, 1)$ so its velocity is $\mathbf{V}(2, 1) = \langle 4, 3\rangle$. After 0.01 units of time, the particle's change in location should be approximately $0.01\,\mathbf{V}(2, 1) = 0.01\,\langle 4, 3\rangle = \langle 0.04, 0.03\rangle$, so the particle should be approximately at the point $(2.04, 1.03)$.

34. At $t = 1$ the particle is at $(1, 3)$ so its velocity is $\mathbf{F}(1, 3) = \langle 1, -1\rangle$. After 0.05 units of time, the particle's change in location should be approximately $0.05\,\mathbf{F}(1, 3) = 0.05\,\langle 1, -1\rangle = \langle 0.05, -0.05\rangle$, so the particle should be approximately at the point $(1.05, 2.95)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

35. (a) We sketch the vector field $\mathbf{F}(x, y) = x\,\mathbf{i} - y\,\mathbf{j}$ along with several approximate flow lines. The flow lines appear to be hyperbolas with shape similar to the graph of $y = \pm 1/x$, so we might guess that the flow lines have equations $y = C/x$.



(b) If $x = x(t)$ and $y = y(t)$ are parametric equations of a flow line, then the velocity vector of the flow line at the point (x, y) is $x'(t)\,\mathbf{i} + y'(t)\,\mathbf{j}$. Since the velocity vectors coincide with the vectors in the vector field, we have $x'(t)\,\mathbf{i} + y'(t)\,\mathbf{j} = x\,\mathbf{i} - y\,\mathbf{j} \;\Rightarrow\; dx/dt = x,\ dy/dt = -y$. To solve these differential equations, we know $dx/dt = x \;\Rightarrow\; dx/x = dt \;\Rightarrow\; \ln|x| = t + C \;\Rightarrow\; x = \pm e^{t+C} = Ae^t$ for some constant A, and $dy/dt = -y \;\Rightarrow\; dy/y = -dt \;\Rightarrow\; \ln|y| = -t + K \;\Rightarrow\; y = \pm e^{-t+K} = Be^{-t}$ for some constant B. Therefore $xy = Ae^t Be^{-t} = AB =$ constant. If the flow line passes through $(1, 1)$ then $(1)\,(1) =$ constant $= 1 \;\Rightarrow\; xy = 1 \;\Rightarrow\; y = 1/x,\ x > 0$.

36. (a) We sketch the vector field $\mathbf{F}(x, y) = \mathbf{i} + x\,\mathbf{j}$ along with several approximate flow lines. The flow lines appear to be parabolas.



(b) If $x = x(t)$ and $y = y(t)$ are parametric equations of a flow line, then the velocity vector of the flow line at the point (x, y) is $x'(t)\,\mathbf{i} + y'(t)\,\mathbf{j}$. Since the velocity vectors coincide with the vectors in the vector field, we have $x'(t)\,\mathbf{i} + y'(t)\,\mathbf{j} = \mathbf{i} + x\,\mathbf{j} \;\Rightarrow\; \dfrac{dx}{dt} = 1,\ \dfrac{dy}{dt} = x$. Thus $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{x}{1} = x$.

(c) From part (b), $dy/dx = x$. Integrating, we have $y = \frac{1}{2}x^2 + c$. Since the particle starts at the origin, we know $(0, 0)$ is on the curve, so $0 = 0 + c \;\Rightarrow\; c = 0$ and the path the particle follows is $y = \frac{1}{2}x^2$.

16.2 Line Integrals

1. $x = t^2$ and $y = 2t$, $0 \le t \le 3$, so by Formula 3

$$\int_C y\,ds = \int_0^3 2t\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt = \int_0^3 2t\sqrt{(2t)^2 + (2)^2}\,dt = \int_0^3 2t\sqrt{4t^2 + 4}\,dt$$

$$= \int_0^3 4t\sqrt{t^2 + 1}\,dt = 2 \cdot \tfrac{2}{3}\left(t^2 + 1\right)^{3/2}\Big]_0^3 = \tfrac{4}{3}(10^{3/2} - 1) \text{ or } \tfrac{4}{3}(10\sqrt{10} - 1)$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. $x = t^3$ and $y = t^4$, $1 \le t \le 2$, so by Formula 3

$$\begin{aligned}\int_C (x/y)\,ds &= \int_1^2 (t^3/t^4)\sqrt{(3t^2)^2 + (4t^3)^2}\,dt = \int_1^2 (1/t)\cdot t^2\sqrt{9+16t^2}\,dt = \int_1^2 t\sqrt{9+16t^2}\,dt \\ &= \tfrac{1}{32}\cdot\tfrac{2}{3}\left(9+16t^2\right)^{3/2}\Big]_1^2 = \tfrac{1}{48}(73^{3/2} - 25^{3/2}) \text{ or } \tfrac{1}{48}(73\sqrt{73} - 125)\end{aligned}$$

3. Parametric equations for C are $x = 4\cos t,\ y = 4\sin t,\ -\frac{\pi}{2} \le t \le \frac{\pi}{2}$. Then

$$\begin{aligned}\int_C xy^4\,ds &= \int_{-\pi/2}^{\pi/2}(4\cos t)(4\sin t)^4\sqrt{(-4\sin t)^2 + (4\cos t)^2}\,dt = \int_{-\pi/2}^{\pi/2} 4^5\cos t\,\sin^4 t\,\sqrt{16(\sin^2 t + \cos^2 t)}\,dt \\ &= 4^5\int_{-\pi/2}^{\pi/2}(\sin^4 t\cos t)(4)\,dt = (4)^6\left[\tfrac{1}{5}\sin^5 t\right]_{-\pi/2}^{\pi/2} = 4^6\cdot\tfrac{2}{5} = 1638.4\end{aligned}$$

4. Parametric equations for C are $x = 2+3t,\ y = 4t,\ 0 \le t \le 1$. Then

$$\int_C xe^y\,ds = \int_0^1 (2+3t)\,e^{4t}\sqrt{3^2+4^2}\,dt = 5\int_0^1 (2+3t)\,e^{4t}\,dt$$

Integrating by parts with $u = 2+3t \ \Rightarrow\ du = 3\,dt,\ dv = e^{4t}\,dt \ \Rightarrow\ v = \frac{1}{4}e^{4t}$ gives

$$\int_C xe^y\,ds = 5\left[\tfrac{1}{4}(2+3t)e^{4t} - \tfrac{3}{16}e^{4t}\right]_0^1 = 5\left[\tfrac{5}{4}e^4 - \tfrac{3}{16}e^4 - \tfrac{1}{2} + \tfrac{3}{16}\right] = \tfrac{85}{16}e^4 - \tfrac{25}{16}$$

5. If we choose x as the parameter, parametric equations for C are $x = x,\ y = x^2$ for $0 \le x \le \pi$ and by Equations 7

$$\begin{aligned}\int_C \left(x^2y + \sin x\right)dy &= \int_0^\pi \left[x^2(x^2) + \sin x\right]\cdot 2x\,dx = 2\int_0^\pi \left(x^5 + x\sin x\right)dx \\ &= 2\left[\tfrac{1}{6}x^6 - x\cos x + \sin x\right]_0^\pi \quad \left[\begin{array}{c}\text{where we integrated by parts} \\ \text{in the second term}\end{array}\right] \\ &= 2\left[\tfrac{1}{6}\pi^6 + \pi + 0 - 0\right] = \tfrac{1}{3}\pi^6 + 2\pi\end{aligned}$$

6. Choosing y as the parameter, we have $x = y^3,\ y = y,\ -1 \le y \le 1$. Then

$$\int_C e^x\,dx = \int_{-1}^1 e^{y^3}\cdot 3y^2\,dy = e^{y^3}\Big]_{-1}^1 = e^1 - e^{-1} = e - \tfrac{1}{e}.$$

7.



$C = C_1 + C_2$

On C_1: $x = x, y = \frac{1}{2}x \ \Rightarrow\ dy = \frac{1}{2}\,dx,\ 0 \le x \le 2.$

On C_2: $x = x, y = 3 - x \ \Rightarrow\ dy = -dx,\ 2 \le x \le 3.$

Then

$$\begin{aligned}\int_C (x+2y)\,dx + x^2\,dy &= \int_{C_1}(x+2y)\,dx + x^2\,dy + \int_{C_2}(x+2y)\,dx + x^2\,dy \\ &= \int_0^2 \left[x + 2\left(\tfrac{1}{2}x\right) + x^2\left(\tfrac{1}{2}\right)\right]dx + \int_2^3 \left[x + 2(3-x) + x^2(-1)\right]dx \\ &= \int_0^2 \left(2x + \tfrac{1}{2}x^2\right)dx + \int_2^3 \left(6 - x - x^2\right)dx \\ &= \left[x^2 + \tfrac{1}{6}x^3\right]_0^2 + \left[6x - \tfrac{1}{2}x^2 - \tfrac{1}{3}x^3\right]_2^3 = \tfrac{16}{3} - 0 + \tfrac{9}{2} - \tfrac{22}{3} = \tfrac{5}{2}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

8.



$C = C_1 + C_2$

On C_1: $x = 2\cos t \;\Rightarrow\; dx = -2\sin t\,dt$, $y = 2\sin t \;\Rightarrow$
$dy = 2\cos t\;dt$, $0 \le t \le \frac{\pi}{2}$.

On C_2: $x = 4t \;\Rightarrow\; dx = 4\,dt$, $y = 2 + t \;\Rightarrow$
$dy = dt$, $0 \le t \le 1$.

Then

$$\begin{aligned}\int_C x^2\,dx + y^2\,dy &= \int_{C_1} x^2\,dx + y^2\,dy + \int_{C_2} x^2\,dx + y^2\,dy \\ &= \int_0^{\pi/2} (2\cos t)^2(-2\sin t\,dt) + (2\sin t)^2(2\cos t\,dt) + \int_0^1 (4t)^2(4\,dt) + (2+t)^2\,dt \\ &= 8\int_0^{\pi/2} (-\cos^2 t\sin t + \sin^2 t\cos t)\,dt + \int_0^1 (65t^2 + 4t + 4)\,dt \\ &= 8\left[\tfrac{1}{3}\cos^3 t + \tfrac{1}{3}\sin^3 t\right]_0^{\pi/2} + \left[\tfrac{65}{3}t^3 + 2t^2 + 4t\right]_0^1 = 8\left(\tfrac{1}{3} - \tfrac{1}{3}\right) + \tfrac{65}{3} + 2 + 4 = \tfrac{83}{3}\end{aligned}$$

9. $x = \cos t$, $y = \sin t$, $z = t$, $0 \le t \le \pi/2$. Then by Formula 9,

$$\begin{aligned}\int_C x^2 y\,ds &= \int_0^{\pi/2} (\cos t)^2(\sin t)\sqrt{\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2 + \left(\tfrac{dz}{dt}\right)^2}\,dt \\ &= \int_0^{\pi/2} \cos^2 t\sin t\,\sqrt{(-\sin t)^2 + (\cos t)^2 + (1)^2}\,dt = \int_0^{\pi/2} \cos^2 t\sin t\,\sqrt{\sin^2 t + \cos^2 t + 1}\,dt \\ &= \sqrt{2}\int_0^{\pi/2} \cos^2 t\sin t\,dt = \sqrt{2}\left[-\tfrac{1}{3}\cos^3 t\right]_0^{\pi/2} = \sqrt{2}\left(0 + \tfrac{1}{3}\right) = \tfrac{\sqrt{2}}{3}\end{aligned}$$

10. Parametric equations for C are $x = 3 - 2t$, $y = 1 + t$, $z = 2 + 3t$, $0 \le t \le 1$. Then

$$\begin{aligned}\int_C y^2 z\,ds &= \int_0^1 (1+t)^2(2+3t)\sqrt{(-2)^2 + 1^2 + 3^2}\,dt = \sqrt{14}\int_0^1 (3t^3 + 8t^2 + 7t + 2)\,dt \\ &= \sqrt{14}\left[\tfrac{3}{4}t^4 + \tfrac{8}{3}t^3 + \tfrac{7}{2}t^2 + 2t\right]_0^1 = \sqrt{14}\left(\tfrac{3}{4} + \tfrac{8}{3} + \tfrac{7}{2} + 2\right) = \tfrac{107}{12}\sqrt{14}\end{aligned}$$

11. Parametric equations for C are $x = t$, $y = 2t$, $z = 3t$, $0 \le t \le 1$. Then

$$\int_C xe^{yz}\,ds = \int_0^1 te^{(2t)(3t)}\sqrt{1^2 + 2^2 + 3^2}\,dt = \sqrt{14}\int_0^1 te^{6t^2}\,dt = \sqrt{14}\left[\tfrac{1}{12}e^{6t^2}\right]_0^1 = \tfrac{\sqrt{14}}{12}(e^6 - 1).$$

12. $\sqrt{(dx/dt)^2 + (dy/dt)^2 + (dz/dt)^2} = \sqrt{1^2 + (-2\sin 2t)^2 + (2\cos 2t)^2} = \sqrt{1 + 4(\sin^2 2t + \cos^2 2t)} = \sqrt{5}$. Then

$$\begin{aligned}\int_C (x^2 + y^2 + z^2)\,ds &= \int_0^{2\pi} (t^2 + \cos^2 2t + \sin^2 2t)\sqrt{5}\,dt = \sqrt{5}\int_0^{2\pi} (t^2 + 1)\,dt \\ &= \sqrt{5}\left[\tfrac{1}{3}t^3 + t\right]_0^{2\pi} = \sqrt{5}\left[\tfrac{1}{3}(8\pi^3) + 2\pi\right] = \sqrt{5}\left(\tfrac{8}{3}\pi^3 + 2\pi\right)\end{aligned}$$

13. $\int_C xye^{yz}\,dy = \int_0^1 (t)(t^2)e^{(t^2)(t^3)}\cdot 2t\,dt = \int_0^1 2t^4 e^{t^5}\,dt = \tfrac{2}{5}e^{t^5}\Big]_0^1 = \tfrac{2}{5}(e^1 - e^0) = \tfrac{2}{5}(e - 1)$

14. $\int_C y\,dx + z\,dy + x\,dz = \int_1^4 t\cdot\tfrac{1}{2}t^{-1/2}\,dt + t^2\cdot dt + \sqrt{t}\cdot 2t\,dt = \int_1^4 \left(\tfrac{1}{2}t^{1/2} + t^2 + 2t^{3/2}\right)dt$

$$= \left[\tfrac{1}{3}t^{3/2} + \tfrac{1}{3}t^3 + \tfrac{4}{5}t^{5/2}\right]_1^4 = \tfrac{8}{3} + \tfrac{64}{3} + \tfrac{128}{5} - \tfrac{1}{3} - \tfrac{1}{3} - \tfrac{4}{5} = \tfrac{722}{15}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. Parametric equations for C are $x = 1 + 3t,\ \ y = t,\ \ z = 2t,\ 0 \le t \le 1$. Then

$$\begin{aligned}\int_C z^2\,dx + x^2\,dy + y^2\,dz &= \int_0^1 (2t)^2 \cdot 3\,dt + (1+3t)^2\,dt + t^2 \cdot 2\,dt = \int_0^1 \left(23t^2 + 6t + 1\right) dt \\ &= \left[\tfrac{23}{3}t^3 + 3t^2 + t\right]_0^1 = \tfrac{23}{3} + 3 + 1 = \tfrac{35}{3}\end{aligned}$$

16.



On C_1: $x = t \ \Rightarrow\ dx = dt,\ y = 0 \ \Rightarrow$
$dy = 0\,dt,\ z = t \ \Rightarrow\ dz = dt,\ 0 \le t \le 1.$

On C_2: $x = 1 - t \ \Rightarrow\ dx = -dt,\ y = t \ \Rightarrow$
$dy = dt,\ z = 1 + t \ \Rightarrow\ dz = dt,\ 0 \le t \le 1.$

Then

$$\begin{aligned}&\int_C (y+z)\,dx + (x+z)\,dy + (x+y)\,dz \\ &\quad= \int_{C_1} (y+z)\,dx + (x+z)\,dy + (x+y)\,dz + \int_{C_2} (y+z)\,dx + (x+z)\,dy + (x+y)\,dz \\ &\quad= \int_0^1 (0+t)\,dt + (t+t)\cdot 0\,dt + (t+0)\,dt + \int_0^1 (t+1+t)(-dt) + (1-t+1+t)\,dt + (1-t+t)\,dt \\ &\quad= \int_0^1 2t\,dt + \int_0^1 (-2t+2)\,dt = \left[t^2\right]_0^1 + \left[-t^2+2t\right]_0^1 = 1 + 1 = 2\end{aligned}$$

17. (a) Along the line $x = -3$, the vectors of $\mathbf{F}$ have positive y-components, so since the path goes upward, the integrand $\mathbf{F} \cdot \mathbf{T}$ is always positive. Therefore $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \mathbf{F} \cdot \mathbf{T}\,ds$ is positive.

(b) All of the (nonzero) field vectors along the circle with radius 3 are pointed in the clockwise direction, that is, opposite the direction to the path. So $\mathbf{F} \cdot \mathbf{T}$ is negative, and therefore $\int_{C_2} \mathbf{F} \cdot d\mathbf{r} = \int_{C_2} \mathbf{F} \cdot \mathbf{T}\,ds$ is negative.

18. Vectors starting on C_1 point in roughly the same direction as C_1, so the tangential component $\mathbf{F} \cdot \mathbf{T}$ is positive. Then $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \mathbf{F} \cdot \mathbf{T}\,ds$ is positive. On the other hand, no vectors starting on C_2 point in the same direction as C_2, while some vectors point in roughly the opposite direction, so we would expect $\int_{C_2} \mathbf{F} \cdot d\mathbf{r} = \int_{C_2} \mathbf{F} \cdot \mathbf{T}\,ds$ to be negative.

19. $\mathbf{r}(t) = t^3\,\mathbf{i} + t^2\,\mathbf{j}$, so $\mathbf{F}(\mathbf{r}(t)) = (t^3)(t^2)^2\,\mathbf{i} - (t^3)^2\,\mathbf{j} = t^7\,\mathbf{i} - t^6\,\mathbf{j}$ and $\mathbf{r}'(t) = 3t^2\,\mathbf{i} + 2t\,\mathbf{j}$. Then

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\,dt = \int_0^1 (t^7 \cdot 3t^2 - t^6 \cdot 2t)\,dt = \int_0^1 (3t^9 - 2t^7)\,dt = \left[\tfrac{3}{10}t^{10} - \tfrac{1}{4}t^8\right]_0^1 = \tfrac{3}{10} - \tfrac{1}{4} = \tfrac{1}{20}.$$

20. $\mathbf{F}(\mathbf{r}(t)) = \left(t^2 + (t^3)^2\right)\mathbf{i} + (t^2)(-2t)\,\mathbf{j} + (t^3 - 2t)\,\mathbf{k} = (t^2 + t^6)\,\mathbf{i} - 2t^3\,\mathbf{j} + (t^3 - 2t)\,\mathbf{k}$, $\mathbf{r}'(t) = 2t\,\mathbf{i} + 3t^2\,\mathbf{j} - 2\,\mathbf{k}$. Then

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^2 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\,dt = \int_0^2 (2t^3 + 2t^7 - 6t^5 - 2t^3 + 4t)\,dt = \int_0^2 (2t^7 - 6t^5 + 4t)\,dt \\ &= \left[\tfrac{1}{4}t^8 - t^6 + 2t^2\right]_0^2 = 64 - 64 + 8 = 8\end{aligned}$$

21. $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^1 \langle \sin t^3, \cos(-t^2), t^4 \rangle \cdot \langle 3t^2, -2t, 1 \rangle\,dt$

$$= \int_0^1 (3t^2 \sin t^3 - 2t\cos t^2 + t^4)\,dt = \left[-\cos t^3 - \sin t^2 + \tfrac{1}{5}t^5\right]_0^1 = \tfrac{6}{5} - \cos 1 - \sin 1$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

22. $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{\pi} \langle \cos t, \sin t, \cos t \sin t\rangle \cdot \langle -\sin t, \cos t, 1\rangle \, dt = \int_0^{\pi} \sin t \cos t \, dt = \frac{1}{2}\sin^2 t\big]_0^{\pi} = 0$

23. $\mathbf{F}(\mathbf{r}(t)) = \sqrt{\sin^2 t + \sin t \cos t}\,\mathbf{i} + \left[(\sin t \cos t)/\sin^2 t\right]\mathbf{j} = \sqrt{\sin^2 t + \sin t \cos t}\,\mathbf{i} + \cot t\,\mathbf{j}$,

$\mathbf{r}'(t) = 2\sin t \cos t\,\mathbf{i} + (\cos^2 t - \sin^2 t)\,\mathbf{j}$. Then

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_{\pi/6}^{\pi/3} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_{\pi/6}^{\pi/3} \left[2\sin t\cos t\sqrt{\sin^2 t + \sin t\cos t} + (\cot t)(\cos^2 t - \sin^2 t)\right] dt$$
$$\approx 0.5424$$

24. $\mathbf{F}(\mathbf{r}(t)) = (\cos t \tan t)e^{\sin t}\,\mathbf{i} + (\tan t \sin t)e^{\cos t}\,\mathbf{j} + (\sin t \cos t)e^{\tan t}\,\mathbf{k}$

$= (\sin t)e^{\sin t}\,\mathbf{i} + (\tan t \sin t)e^{\cos t}\,\mathbf{j} + (\sin t \cos t)e^{\tan t}\,\mathbf{k}$,

$\mathbf{r}'(t) = \cos t\,\mathbf{i} - \sin t\,\mathbf{j} + \sec^2 t\,\mathbf{k}$. Then

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{\pi/4} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^{\pi/4} \left[(\sin t\cos t)e^{\sin t} - (\tan t \sin^2 t)e^{\cos t} + (\tan t)e^{\tan t}\right] dt \approx 0.8527$$

25. $x = t^2$, $y = t^3$, $z = \sqrt{t}$ so by Formula 9,

$$\int_C xy\arctan z\,ds = \int_1^2 (t^2)(t^3)\arctan\sqrt{t}\cdot\sqrt{(2t)^2 + (3t^2)^2 + \left[1/(2\sqrt{t}\,)\right]^2}\,dt$$
$$= \int_1^2 t^5\sqrt{4t^2 + 9t^4 + 1/(4t)}\,\arctan\sqrt{t}\,dt \approx 94.8231$$

26. $x = 1 + 3t$, $y = 2 + t^2$, $z = t^4$ so by Formula 9,

$$\int_C z\ln(x+y)\,ds = \int_{-1}^1 t^4\ln(1 + 3t + 2 + t^2)\cdot\sqrt{(3)^2 + (2t)^2 + (4t^3)^2}\,dt$$
$$= \int_{-1}^1 t^4\sqrt{9 + 4t^2 + 16t^6}\,\ln(3 + 3t + t^2)\,dt \approx 1.7260$$

27. We graph $\mathbf{F}(x, y) = (x - y)\,\mathbf{i} + xy\,\mathbf{j}$ and the curve C. We see that most of the vectors starting on C point in roughly the same direction as C, so for these portions of C the tangential component $\mathbf{F}\cdot\mathbf{T}$ is positive. Although some vectors in the third quadrant which start on C point in roughly the opposite direction, and hence give negative tangential components, it seems reasonable that the effect of these portions of C is outweighed by the positive tangential components. Thus, we would expect $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \mathbf{F}\cdot\mathbf{T}\,ds$ to be positive.

To verify, we evaluate $\int_C \mathbf{F}\cdot d\mathbf{r}$. The curve C can be represented by $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j}$, $0 \le t \le \frac{3\pi}{2}$, so $\mathbf{F}(\mathbf{r}(t)) = (2\cos t - 2\sin t)\,\mathbf{i} + 4\cos t\sin t\,\mathbf{j}$ and $\mathbf{r}'(t) = -2\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$. Then



$$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{3\pi/2} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt$$
$$= \int_0^{3\pi/2}[-2\sin t(2\cos t - 2\sin t) + 2\cos t(4\cos t\sin t)]\,dt$$
$$= 4\int_0^{3\pi/2}(\sin^2 t - \sin t\cos t + 2\sin t\cos^2 t)\,dt$$
$$3\pi + \tfrac{2}{3} \qquad \text{[using a CAS]}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. We graph $\mathbf{F}(x,y) = \dfrac{x}{\sqrt{x^2+y^2}}\,\mathbf{i} + \dfrac{y}{\sqrt{x^2+y^2}}\,\mathbf{j}$ and the curve C. In the first quadrant, each vector starting on C points in roughly the same direction as C, so the tangential component $\mathbf{F}\cdot\mathbf{T}$ is positive. In the second quadrant, each vector starting on C points in roughly the direction opposite to C, so $\mathbf{F}\cdot\mathbf{T}$ is negative. Here, it appears that the tangential components in the first and second quadrants counteract each other, so it seems reasonable to guess that $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \mathbf{F}\cdot\mathbf{T}\,ds$ is zero. To verify, we evaluate $\int_C \mathbf{F}\cdot d\mathbf{r}$. The curve C can be represented by $\mathbf{r}(t) = t\,\mathbf{i} + (1+t^2)\,\mathbf{j}$, $-1 \le t \le 1$, so $\mathbf{F}(\mathbf{r}(t)) = \dfrac{t}{\sqrt{t^2+(1+t^2)^2}}\,\mathbf{i} + \dfrac{1+t^2}{\sqrt{t^2+(1+t^2)^2}}\,\mathbf{j}$ and $\mathbf{r}'(t) = \mathbf{i} + 2t\,\mathbf{j}$. Then



$$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_{-1}^{1} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_{-1}^{1}\left(\frac{t}{\sqrt{t^2+(1+t^2)^2}} + \frac{2t(1+t^2)}{\sqrt{t^2+(1+t^2)^2}}\right)dt$$

$$= \int_{-1}^{1} \frac{t(3+2t^2)}{\sqrt{t^4+3t^2+1}}\,dt = 0 \qquad \text{[since the integrand is an odd function]}$$

29. (a) $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 \left\langle e^{t^2-1}, t^5\right\rangle \cdot \left\langle 2t, 3t^2\right\rangle dt = \int_0^1 \left(2te^{t^2-1} + 3t^7\right) dt = \left[e^{t^2-1} + \frac{3}{8}t^8\right]_0^1 = \frac{11}{8} - 1/e$

(b) $\mathbf{r}(0) = \mathbf{0}$, $\mathbf{F}(\mathbf{r}(0)) = \langle e^{-1}, 0\rangle$;

$\mathbf{r}\left(\frac{1}{\sqrt{2}}\right) = \left\langle \frac{1}{2}, \frac{1}{2\sqrt{2}}\right\rangle$, $\mathbf{F}\left(\mathbf{r}\left(\frac{1}{\sqrt{2}}\right)\right) = \left\langle e^{-1/2}, \frac{1}{4\sqrt{2}}\right\rangle$;

$\mathbf{r}(1) = \langle 1, 1\rangle$, $\mathbf{F}(\mathbf{r}(1)) = \langle 1, 1\rangle$.



In order to generate the graph with Maple, we use the `line` command in the `plottools` package to define each of the vectors. For example,

```
v1:=line([0,0],[exp(-1),0]):
```

generates the vector from the vector field at the point $(0,0)$ (but without an arrowhead) and gives it the name `v1`. To show everything on the same screen, we use the `display` command. In Mathematica, we use `ListPlot` (with the `PlotJoined -> True` option) to generate the vectors, and then `Show` to show everything on the same screen.

30. (a) $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_{-1}^{1} \langle 2t, t^2, 3t\rangle \cdot \langle 2, 3, -2t\rangle\,dt = \int_{-1}^{1} (4t + 3t^2 - 6t^2)\,dt = \left[2t^2 - t^3\right]_{-1}^{1} = -2$

(b) Now $\mathbf{F}(\mathbf{r}(t)) = \langle 2t, t^2, 3t\rangle$, so $\mathbf{F}(\mathbf{r}(-1)) = \langle -2, 1, -3\rangle$, $\mathbf{F}\left(\mathbf{r}\left(-\frac{1}{2}\right)\right) = \left\langle -1, \frac{1}{4}, -\frac{3}{2}\right\rangle$, $\mathbf{F}\left(\mathbf{r}\left(\frac{1}{2}\right)\right) = \left\langle 1, \frac{1}{4}, \frac{3}{2}\right\rangle$, and $\mathbf{F}(\mathbf{r}(1)) = \langle 2, 1, 3\rangle$.





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

31. $x = e^{-t}\cos 4t,\ \ y = e^{-t}\sin 4t,\ \ z = e^{-t},\ \ 0 \le t \le 2\pi$.

Then $\dfrac{dx}{dt} = e^{-t}(-\sin 4t)(4) - e^{-t}\cos 4t = -e^{-t}(4\sin 4t + \cos 4t)$,

$\dfrac{dy}{dt} = e^{-t}(\cos 4t)(4) - e^{-t}\sin 4t = -e^{-t}(-4\cos 4t + \sin 4t)$, and $\dfrac{dz}{dt} = -e^{-t}$, so

$$\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} = \sqrt{(-e^{-t})^2[(4\sin 4t + \cos 4t)^2 + (-4\cos 4t + \sin 4t)^2 + 1]}$$

$$= e^{-t}\sqrt{16(\sin^2 4t + \cos^2 4t) + \sin^2 4t + \cos^2 4t + 1} = 3\sqrt{2}\,e^{-t}$$

Therefore
$$\int_C x^3y^2z\,ds = \int_0^{2\pi}(e^{-t}\cos 4t)^3(e^{-t}\sin 4t)^2(e^{-t})\,(3\sqrt{2}\,e^{-t})\,dt$$

$$= \int_0^{2\pi} 3\sqrt{2}\,e^{-7t}\cos^3 4t\sin^2 4t\,dt = \tfrac{172{,}704}{5{,}632{,}705}\sqrt{2}\,(1 - e^{-14\pi})$$

32. (a) We parametrize the circle C as $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j},\ \ 0 \le t \le 2\pi$. So $\mathbf{F}(\mathbf{r}(t)) = \langle 4\cos^2 t, 4\cos t\sin t\rangle$, $\mathbf{r}'(t) = \langle -2\sin t, 2\cos t\rangle$, and $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi}(-8\cos^2 t\sin t + 8\cos^2 t\sin t)\,dt = 0$.

(b)



From the graph, we see that all of the vectors in the field are perpendicular to the path. This indicates that the field does no work on the particle, since the field never pulls the particle in the direction in which it is going. In other words, at any point along C, $\mathbf{F}\cdot\mathbf{T} = 0$, and so certainly $\int_C \mathbf{F}\cdot d\mathbf{r} = 0$.

33. We use the parametrization $x = 2\cos t,\ y = 2\sin t,\ -\frac{\pi}{2} \le t \le \frac{\pi}{2}$. Then

$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt = \sqrt{(-2\sin t)^2 + (2\cos t)^2}\,dt = 2\,dt$, so $m = \int_C k\,ds = 2k\int_{-\pi/2}^{\pi/2} dt = 2k(\pi)$,

$\overline{x} = \frac{1}{2\pi k}\int_C xk\,ds = \frac{1}{2\pi}\int_{-\pi/2}^{\pi/2}(2\cos t)2\,dt = \frac{1}{2\pi}\left[4\sin t\right]_{-\pi/2}^{\pi/2} = \frac{4}{\pi}$, $\overline{y} = \frac{1}{2\pi k}\int_C yk\,ds = \frac{1}{2\pi}\int_{-\pi/2}^{\pi/2}(2\sin t)2\,dt = 0$.

Hence $(\overline{x}, \overline{y}) = \left(\frac{4}{\pi}, 0\right)$.

34. We use the parametrization $x = a\cos t,\ \ y = a\sin t,\ \ 0 \le t \le \frac{\pi}{2}$. Then

$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt = \sqrt{(-a\sin t)^2 + (a\cos t)^2}\,dt = a\,dt$, so

$$m = \int_C \rho(x,y)\,ds = \int_C kxy\,ds = \int_0^{\pi/2} k(a\cos t)(a\sin t)\,a\,dt = ka^3\int_0^{\pi/2}\cos t\sin t\,dt = ka^3\left[\tfrac{1}{2}\sin^2 t\right]_0^{\pi/2} = \tfrac{1}{2}ka^3,$$

$$\overline{x} = \frac{1}{ka^3/2}\int_C x(kxy)\,ds = \frac{2}{ka^3}\int_0^{\pi/2} k(a\cos t)^2(a\sin t)a\,dt = \frac{2}{ka^3}\cdot ka^4\int_0^{\pi/2}\cos^2 t\,\sin t\,dt$$

$2a\left[-\frac{1}{3}\cos^3 t\right]_0^{\pi/2} = 2a\left(0 + \frac{1}{3}\right) = \frac{2}{3}a$, and

$$\overline{y} = \frac{1}{ka^3/2}\int_C y(kxy)\,ds = \frac{2}{ka^3}\int_0^{\pi/2} k(a\cos t)(a\sin t)^2 a\,dt = \frac{2}{ka^3}\cdot ka^4\int_0^{\pi/2}\sin^2 t\,\cos t\,dt$$

$= 2a\left[\frac{1}{3}\sin^3 t\right]_0^{\pi/2} - 2a\left(\frac{1}{3} \quad 0\right) - \frac{2}{3}a$.

Therefore the mass is $\frac{1}{2}ka^3$ and the center of mass is $(\overline{x}, \overline{y})$ $\quad \left(\frac{2}{3}a, \frac{2}{3}a\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. (a) $\overline{x} = \dfrac{1}{m}\displaystyle\int_C x\rho(x,y,z)\,ds$, $\overline{y} = \dfrac{1}{m}\displaystyle\int_C y\rho(x,y,z)\,ds$, $\overline{z} = \dfrac{1}{m}\displaystyle\int_C z\rho(x,y,z)\,ds$ where $m = \int_C \rho(x,y,z)\,ds$.

(b) $m = \int_C k\,ds = k\int_0^{2\pi}\sqrt{4\sin^2 t + 4\cos^2 t + 9}\,dt = k\sqrt{13}\int_0^{2\pi} dt = 2\pi k\sqrt{13}$,

$$\overline{x} = \frac{1}{2\pi k\sqrt{13}}\int_0^{2\pi} 2k\sqrt{13}\sin t\,dt = 0,\ \overline{y} = \frac{1}{2\pi k\sqrt{13}}\int_0^{2\pi} 2k\sqrt{13}\cos t\,dt = 0,$$

$$\overline{z} = \frac{1}{2\pi k\sqrt{13}}\int_0^{2\pi}\left(k\sqrt{13}\right)(3t)\,dt = \frac{3}{2\pi}(2\pi^2) = 3\pi.\ \text{Hence } (\overline{x},\overline{y},\overline{z}) = (0,0,3\pi).$$

36. $m = \int_C (x^2+y^2+z^2)\,ds = \int_0^{2\pi}(t^2+1)\sqrt{(1)^2+(-\sin t)^2+(\cos t)^2}\,dt = \int_0^{2\pi}(t^2+1)\sqrt{2}\,dt = \sqrt{2}\left(\frac{8}{3}\pi^3 + 2\pi\right)$,

$$\overline{x} = \frac{1}{\sqrt{2}\left(\frac{8}{3}\pi^3+2\pi\right)}\int_0^{2\pi}\sqrt{2}\,(t^3+t)\,dt = \frac{4\pi^4+2\pi^2}{\frac{8}{3}\pi^3+2\pi} = \frac{3\pi(2\pi^2+1)}{4\pi^2+3},$$

$$\overline{y} = \frac{3}{2\sqrt{2}\,\pi(4\pi^2+3)}\int_0^{2\pi}\left(\sqrt{2}\cos t\right)(t^2+1)\,dt = 0,\ \text{and}$$

$$\overline{z} = \frac{3}{2\sqrt{2}\,\pi(4\pi^2+3)}\int_0^{2\pi}\left(\sqrt{2}\sin t\right)(t^2+1)\,dt = 0.\ \text{Hence } (\overline{x},\overline{y},\overline{z}) = \left(\frac{3\pi(2\pi^2+1)}{4\pi^2+3}, 0, 0\right).$$

37. From Example 3, $\rho(x,y) = k(1-y)$, $x = \cos t$, $y = \sin t$, and $ds = dt$, $0 \le t \le \pi$ $\Rightarrow$

$$I_x = \int_C y^2\rho(x,y)\,ds = \int_0^{\pi}\sin^2 t\,[k(1-\sin t)]\,dt = k\int_0^{\pi}(\sin^2 t - \sin^3 t)\,dt$$

$$= \tfrac{1}{2}k\int_0^{\pi}(1-\cos 2t)\,dt - k\int_0^{\pi}(1-\cos^2 t)\sin t\,dt \quad \left[\begin{array}{c}\text{Let } u = \cos t,\ du = -\sin t\,dt \\ \text{in the second integral}\end{array}\right]$$

$$= k\left[\tfrac{\pi}{2} + \int_1^{-1}(1-u^2)\,du\right] = k\left(\tfrac{\pi}{2} - \tfrac{4}{3}\right)$$

$$I_y = \int_C x^2\rho(x,y)\,ds = k\int_0^{\pi}\cos^2 t\,(1-\sin t)\,dt = \tfrac{k}{2}\int_0^{\pi}(1+\cos 2t)\,dt - k\int_0^{\pi}\cos^2 t\sin t\,dt$$

$= k\left(\frac{\pi}{2} - \frac{2}{3}\right)$, using the same substitution as above.

38. The wire is given as $x = 2\sin t$, $y = 2\cos t$, $z = 3t$, $0 \le t \le 2\pi$ with $\rho(x,y,z) = k$. Then

$ds = \sqrt{(2\cos t)^2 + (-2\sin t)^2 + 3^2}\,dt = \sqrt{4(\cos^2 t + \sin^2 t) + 9}\,dt = \sqrt{13}\,dt$ and

$$I_x = \int_C (y^2+z^2)\rho(x,y,z)\,ds = \int_0^{2\pi}(4\cos^2 t + 9t^2)(k)\sqrt{13}\,dt = \sqrt{13}\,k\left[4\left(\tfrac{1}{2}t + \tfrac{1}{4}\sin 2t\right) + 3t^3\right]_0^{2\pi}$$

$$= \sqrt{13}\,k(4\pi + 24\pi^3) = 4\sqrt{13}\,\pi k(1+6\pi^2)$$

$$I_y = \int_C (x^2+z^2)\rho(x,y,z)\,ds = \int_0^{2\pi}\left(4\sin^2 t + 9t^2\right)(k)\sqrt{13}\,dt = \sqrt{13}\,k\left[4\left(\tfrac{1}{2}t - \tfrac{1}{4}\sin 2t\right) + 3t^3\right]_0^{2\pi}$$

$$= \sqrt{13}\,k(4\pi + 24\pi^3) = 4\sqrt{13}\,\pi k(1+6\pi^2)$$

$$I_z = \int_C (x^2+y^2)\rho(x,y,z)\,ds = \int_0^{2\pi}(4\sin^2 t + 4\cos^2 t)(k)\sqrt{13}\,dt = 4\sqrt{13}\,k\int_0^{2\pi} dt = 8\pi\sqrt{13}\,k$$

39. $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi}\langle t - \sin t, 3 - \cos t\rangle \cdot \langle 1 - \cos t, \sin t\rangle\,dt$

$$= \int_0^{2\pi}(t - t\cos t - \sin t + \sin t\cos t + 3\sin t - \sin t\cos t)\,dt$$

$$= \int_0^{2\pi}(t - t\cos t + 2\sin t)\,dt = \left[\tfrac{1}{2}t^2 - (t\sin t + \cos t) - 2\cos t\right]_0^{2\pi} \quad \left[\begin{array}{c}\text{integrate by parts} \\ \text{in the second term}\end{array}\right]$$

$$= 2\pi^2$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. Choosing y as the parameter, the curve C is parametrized by $x = y^2 + 1$, $y = y$, $0 \le y \le 1$. Then

$$W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^1 \left\langle \left(y^2+1\right)^2, ye^{y^2+1} \right\rangle \cdot \langle 2y, 1\rangle \, dy = \int_0^1 \left[2y\left(y^2+1\right)^2 + ye^{y^2+1}\right] dy$$

$$= \left[\tfrac{1}{3}\left(y^2+1\right)^3 + \tfrac{1}{2}e^{y^2+1}\right]_0^1 = \tfrac{8}{3} + \tfrac{1}{2}e^2 - \tfrac{1}{3} - \tfrac{1}{2}e = \tfrac{1}{2}e^2 - \tfrac{1}{2}e + \tfrac{7}{3}$$

41. $\mathbf{r}(t) = \langle 2t, t, 1-t\rangle$, $0 \le t \le 1$.

$$W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^1 \left\langle 2t - t^2, t - (1-t)^2, 1 - t - (2t)^2 \right\rangle \cdot \langle 2, 1, -1\rangle \, dt$$

$$= \int_0^1 \left(4t - 2t^2 + t - 1 + 2t - t^2 - 1 + t + 4t^2\right) dt = \int_0^1 \left(t^2 + 8t - 2\right) dt = \left[\tfrac{1}{3}t^3 + 4t^2 - 2t\right]_0^1 = \tfrac{7}{3}$$

42. $\mathbf{r}(t) = 2\,\mathbf{i} + t\,\mathbf{j} + 5t\,\mathbf{k}$, $0 \le t \le 1$. Therefore

$$W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^1 \frac{K\langle 2, t, 5t\rangle}{(4+26t^2)^{3/2}} \cdot \langle 0, 1, 5\rangle \, dt = K\int_0^1 \frac{26t}{(4+26t^2)^{3/2}} \, dt = K\left[-(4+26t^2)^{-1/2}\right]_0^1 = K\left(\tfrac{1}{2} - \tfrac{1}{\sqrt{30}}\right).$$

43. (a) $\mathbf{r}(t) = at^2\,\mathbf{i} + bt^3\,\mathbf{j} \;\Rightarrow\; \mathbf{v}(t) = \mathbf{r}'(t) = 2at\,\mathbf{i} + 3bt^2\,\mathbf{j} \;\Rightarrow\; \mathbf{a}(t) = \mathbf{v}'(t) = 2a\,\mathbf{i} + 6bt\,\mathbf{j}$, and force is mass times acceleration: $\mathbf{F}(t) = m\,\mathbf{a}(t) = 2ma\,\mathbf{i} + 6mbt\,\mathbf{j}$.

(b) $W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^1 (2ma\,\mathbf{i} + 6mbt\,\mathbf{j}) \cdot (2at\,\mathbf{i} + 3bt^2\,\mathbf{j})\, dt = \int_0^1 \left(4ma^2t + 18mb^2t^3\right) dt$

$= \left[2ma^2t^2 + \tfrac{9}{2}mb^2t^4\right]_0^1 = 2ma^2 + \tfrac{9}{2}mb^2$

44. $\mathbf{r}(t) = a\sin t\,\mathbf{i} + b\cos t\,\mathbf{j} + ct\,\mathbf{k} \;\Rightarrow\; \mathbf{v}(t) = \mathbf{r}'(t) = a\cos t\,\mathbf{i} - b\sin t\,\mathbf{j} + c\,\mathbf{k} \;\Rightarrow\; \mathbf{a}(t) = \mathbf{v}'(t) = -a\sin t\,\mathbf{i} - b\cos t\,\mathbf{j}$ and $\mathbf{F}(t) = m\,\mathbf{a}(t) = -ma\sin t\,\mathbf{i} - mb\cos t\,\mathbf{j}$. Thus

$$W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{\pi/2} (-ma\sin t\,\mathbf{i} - mb\cos t\,\mathbf{j}) \cdot (a\cos t\,\mathbf{i} - b\sin t\,\mathbf{j} + c\,\mathbf{k})\, dt$$

$$= \int_0^{\pi/2} \left(-ma^2\sin t\cos t + mb^2\sin t\cos t\right) dt = m(b^2 - a^2)\left[\tfrac{1}{2}\sin^2 t\right]_0^{\pi/2} = \tfrac{1}{2}m(b^2 - a^2)$$

45. The combined weight of the man and the paint is 185 lb, so the force exerted (equal and opposite to that exerted by gravity) is $\mathbf{F} = 185\,\mathbf{k}$. To parametrize the staircase, let $x = 20\cos t$, $y = 20\sin t$, $z = \frac{90}{6\pi}t = \frac{15}{\pi}t$, $0 \le t \le 6\pi$. Then the work done is

$$W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{6\pi} \langle 0, 0, 185\rangle \cdot \left\langle -20\sin t, 20\cos t, \tfrac{15}{\pi}\right\rangle dt = (185)\tfrac{15}{\pi}\int_0^{6\pi} dt = (185)\left(\tfrac{15}{\pi}\right)(6\pi) \approx 1.67 \times 10^4 \text{ ft-lb}$$

46. This time m is a function of t: $m = 185 - \frac{9}{6\pi}t = 185 - \frac{3}{2\pi}t$. So let $\mathbf{F} = \left(185 - \frac{3}{2\pi}t\right)\mathbf{k}$. To parametrize the staircase, let $x = 20\cos t$, $y = 20\sin t$, $z = \frac{90}{6\pi}t = \frac{15}{\pi}t$, $0 \le t \le 6\pi$. Therefore

$$W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{6\pi} \left\langle 0, 0, 185 - \tfrac{3}{2\pi}t\right\rangle \cdot \left\langle -20\sin t, 20\cos t, \tfrac{15}{\pi}\right\rangle dt = \tfrac{15}{\pi}\int_0^{6\pi} \left(185 - \tfrac{3}{2\pi}t\right) dt$$

$$= \tfrac{15}{\pi}\left[185t - \tfrac{3}{4\pi}t^2\right]_0^{6\pi} = 90\left(185 - \tfrac{9}{2}\right) \approx 1.62 \times 10^4 \text{ ft-lb}$$

47. (a) $\mathbf{r}(t) = \langle \cos t, \sin t\rangle$, $0 \le t \le 2\pi$, and let $\mathbf{F} = \langle a, b\rangle$. Then

$$W = \int_C \mathbf{F} \cdot d\,\mathbf{r} = \int_0^{2\pi} \langle a, b\rangle \cdot \langle -\sin t, \cos t\rangle \, dt = \int_0^{2\pi} (-a\sin t + b\cos t)\, dt = \left[a\cos t + b\sin t\right]_0^{2\pi}$$

$$= a + 0 - a + 0 = 0$$

(b) Yes. $\mathbf{F}(x, y) = k\,\mathbf{x} = \langle kx, ky\rangle$ and

$$W = \int_C \mathbf{F} \cdot d\,\mathbf{r} = \int_0^{2\pi} \langle k\cos t, k\sin t\rangle \cdot \langle -\sin t, \cos t\rangle \, dt = \int_0^{2\pi} (-k\sin t\,\cos t + k\sin t\,\cos t)\, dt = \int_0^{2\pi} 0\, dt = 0.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. Consider the base of the fence in the xy-plane, centered at the origin, with the height given by $z = h(x, y)$. To graph the fence, observe that the fence is highest when $y = 0$ (where the height is 5 m) and lowest when $x = 0$ (a height of 3 m). When $y = \pm x$, the height is 4 m.

Also, the fence can be graphed using parametric equations (see Section 16.6): $x = 10\cos u,\ \ y = 10\sin u,$

$$\begin{aligned} z &= v\big[4 + 0.01((10\cos u)^2 - (10\sin u)^2)\big] = v(4 + \cos^2 u - \sin^2 u) \\ &= v(4 + \cos 2u),\ 0 \le u \le 2\pi,\ 0 \le v \le 1. \end{aligned}$$



The area of the fence is $\int_C h(x, y)\, ds$ where C, the base of the fence, is given by $x = 10\cos t,\ \ y = 10\sin t,\ \ 0 \le t \le 2\pi$. Then

$$\begin{aligned} \int_C h(x, y)\, ds &= \int_0^{2\pi} \left[4 + 0.01((10\cos t)^2 - (10\sin t)^2)\right] \sqrt{(-10\sin t)^2 + (10\cos t)^2}\, dt \\ &= \int_0^{2\pi} (4 + \cos 2t)\sqrt{100}\, dt = 10\big[4t + \tfrac{1}{2}\sin 2t\big]_0^{2\pi} = 10(8\pi) = 80\pi \text{ m}^2 \end{aligned}$$

If we paint both sides of the fence, the total surface area to cover is 160π m^2, and since 1 L of paint covers 100 m^2, we require $\frac{160\pi}{100} = 1.6\pi \approx 5.03$ L of paint.

49. Let $\mathbf{r}(t) = \langle x(t), y(t), z(t)\rangle$ and $\mathbf{v} = \langle v_1, v_2, v_3\rangle$. Then

$$\begin{aligned} \int_C \mathbf{v}\cdot d\mathbf{r} &= \int_a^b \langle v_1, v_2, v_3\rangle \cdot \langle x'(t), y'(t), z'(t)\rangle\, dt = \int_a^b [v_1\, x'(t) + v_2\, y'(t) + v_3\, z'(t)]\, dt \\ &= \big[v_1\, x(t) + v_2\, y(t) + v_3\, z(t)\big]_a^b = [v_1\, x(b) + v_2\, y(b) + v_3\, z(b)] - [v_1\, x(a) + v_2\, y(a) + v_3\, z(a)] \\ &= v_1\,[x(b) - x(a)] + v_2\,[y(b) - y(a)] + v_3\,[z(b) - z(a)] \\ &= \langle v_1, v_2, v_3\rangle \cdot \langle x(b) - x(a), y(b) - y(a), z(b) - z(a)\rangle \\ &= \langle v_1, v_2, v_3\rangle \cdot [\langle x(b), y(b), z(b)\rangle - \langle x(a), y(a), z(a)\rangle] = \mathbf{v}\cdot[\mathbf{r}(b) - \mathbf{r}(a)] \end{aligned}$$

50. If $\mathbf{r}(t) = \langle x(t), y(t), z(t)\rangle$ then

$$\begin{aligned} \int_C \mathbf{r}\cdot d\mathbf{r} &= \int_a^b \langle x(t), y(t), z(t)\rangle \cdot \langle x'(t), y'(t), z'(t)\rangle\, dt = \int_a^b [x(t)\, x'(t) + y(t)\, y'(t) + z(t)\, z'(t)]\, dt \\ &= \big[\tfrac{1}{2}[x(t)]^2 + \tfrac{1}{2}[y(t)]^2 + \tfrac{1}{2}[z(t)]^2\big]_a^b \\ &= \tfrac{1}{2}\left\{\left([x(b)]^2 + [y(b)]^2 + [z(b)]^2\right) - \left([x(a)]^2 + [y(a)]^2 + [z(a)]^2\right)\right\} \\ &= \tfrac{1}{2}\left[|\mathbf{r}(b)|^2 - |\mathbf{r}(a)|^2\right] \end{aligned}$$

51. The work done in moving the object is $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \mathbf{F}\cdot\mathbf{T}\, ds$. We can approximate this integral by dividing C into 7 segments of equal length $\Delta s = 2$ and approximating $\mathbf{F}\cdot\mathbf{T}$, that is, the tangential component of force, at a point (x_i^*, y_i^*) on each segment. Since C is composed of straight line segments, $\mathbf{F}\cdot\mathbf{T}$ is the scalar projection of each force vector onto C. If we choose (x_i^*, y_i^*) to be the point on the segment closest to the origin, then the work done is

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$\int_C \mathbf{F}\cdot\mathbf{T}\,ds \approx \sum_{i=1}^{7}[\mathbf{F}(x_i^*, y_i^*)\cdot\mathbf{T}(x_i^*, y_i^*)]\,\Delta s = [2+2+2+2+1+1+1](2) = 22$. Thus, we estimate the work done to be approximately 22 J.

52. Use the orientation pictured in the figure. Then since $\mathbf{B}$ is tangent to any circle that lies in the plane perpendicular to the wire, $\mathbf{B} = |\mathbf{B}|\,\mathbf{T}$ where $\mathbf{T}$ is the unit tangent to the circle C: $x = r\cos\theta,\ \ y = r\sin\theta$. Thus $\mathbf{B} = |\mathbf{B}|\,\langle -\sin\theta, \cos\theta\rangle$. Then $\int_C \mathbf{B}\cdot d\mathbf{r} = \int_0^{2\pi} |\mathbf{B}|\,\langle -\sin\theta, \cos\theta\rangle \cdot \langle -r\sin\theta, r\cos\theta\rangle\, d\theta = \int_0^{2\pi} |\mathbf{B}|\, r\, d\theta = 2\pi r\,|\mathbf{B}|$. (Note that $|\mathbf{B}|$ here is the magnitude of the field at a distance r from the wire's center.) But by Ampere's Law $\int_C \mathbf{B}\cdot d\mathbf{r} = \mu_0 I$. Hence $|\mathbf{B}| = \mu_0 I/(2\pi r)$.

16.3 The Fundamental Theorem for Line Integrals

1. C appears to be a smooth curve, and since ∇f is continuous, we know f is differentiable. Then Theorem 2 says that the value of $\int_C \nabla f\cdot d\mathbf{r}$ is simply the difference of the values of f at the terminal and initial points of C. From the graph, this is $50 - 10 = 40$.

2. C is represented by the vector function $\mathbf{r}(t) = (t^2+1)\,\mathbf{i} + (t^3+t)\,\mathbf{j}$, $0 \le t \le 1$, so $\mathbf{r}'(t) = 2t\,\mathbf{i} + (3t^2+1)\,\mathbf{j}$. Since $3t^2 + 1 \neq 0$, we have $\mathbf{r}'(t) \neq \mathbf{0}$, thus C is a smooth curve. ∇f is continuous, and hence f is differentiable, so by Theorem 2 we have $\int_C \nabla f\cdot d\mathbf{r} = f(\mathbf{r}(1)) - f(\mathbf{r}(0)) = f(2,2) - f(1,0) = 9 - 3 = 6$.

3. Let $P(x,y) = xy + y^2$ and $Q(x,y) = x^2 + 2xy$. Then $\partial P/\partial y = x + 2y$ and $\partial Q/\partial x = 2x + 2y$. Since $\partial P/\partial y \neq \partial Q/\partial x$, $\mathbf{F}(x,y) = P\,\mathbf{i} + Q\,\mathbf{j}$ is not conservative by Theorem 5.

4. $\partial(y^2 - 2x)/\partial y = 2y = \partial(2xy)/\partial x$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$ which is open and simply-connected, so $\mathbf{F}$ is conservative by Theorem 6. Thus, there exists a function f such that $\nabla f = \mathbf{F}$, that is, $f_x(x,y) = y^2 - 2x$ and $f_y(x,y) = 2xy$. But $f_x(x,y) = y^2 - 2x$ implies $f(x,y) = xy^2 \quad x^2 + g(y)$ and differentiating both sides of this equation with respect to y gives $f_y(x,y) = 2xy + g'(y)$. Thus $2xy = 2xy + g'(y)$ so $g'(y) = 0$ and $g(y) = K$ where K is a constant. Hence $f(x,y) = xy^2 - x^2 + K$ is a potential function for $\mathbf{F}$.

5. $\dfrac{\partial}{\partial y}\left(y^2 e^{xy}\right) = y^2\cdot xe^{xy} + 2ye^{xy} = (xy^2 + 2y)e^{xy}$,

$\dfrac{\partial}{\partial x}\left[(1+xy)e^{xy}\right] = (1+xy)\cdot ye^{xy} + ye^{xy} = ye^{xy} + xy^2e^{xy} + ye^{xy} = (xy^2 + 2y)e^{xy}$.

Since these partial derivatives are equal and the domain of $\mathbf{F}$ is $\mathbb{R}^2$ which is open and simply-connected, $\mathbf{F}$ is conservative by Theorem 6. Thus, there exists a function f such that $\nabla f = \mathbf{F}$, that is, $f_x(x,y) = y^2e^{xy}$ and $f_y(x,y) = (1+xy)e^{xy}$. But $f_x(x,y) = y^2e^{xy}$ implies $f(x,y) = ye^{xy} + g(y)$ and differentiating both sides of this equation with respect to y gives $f_y(x,y) = (1+xy)e^{xy} + g'(y)$. Thus $(1+xy)e^{xy} = (1+xy)e^{xy} + g'(y)$ so $g'(y) = 0$ and $g(y) = K$ where K is a constant. Hence $f(x,y) = ye^{xy} + K$ is a potential function for $\mathbf{F}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $\partial(ye^x)/\partial y = e^x = \partial(e^x + e^y)/\partial x$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$ which is open and simply-connected, so $\mathbf{F}$ is conservative. Hence there exists a function f such that $\nabla f = \mathbf{F}$. Here $f_x(x, y) = ye^x$ implies $f(x, y) = ye^x + g(y)$ and then $f_y(x, y) = e^x + g'(y)$. But $f_y(x, y) = e^x + e^y$ so $g'(y) = e^y \quad \Rightarrow \quad g(y) = e^y + K$ and $f(x, y) = ye^x + e^y + K$ is a potential function for $\mathbf{F}$.

7. $\partial(ye^x + \sin y)/\partial y = e^x + \cos y = \partial(e^x + x\cos y)/\partial x$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$. Hence $\mathbf{F}$ is conservative so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x, y) = ye^x + \sin y$ implies $f(x, y) = ye^x + x\sin y + g(y)$ and $f_y(x, y) = e^x + x\cos y + g'(y)$. But $f_y(x, y) = e^x + x\cos y$ so $g(y) = K$ and $f(x, y) = ye^x + x\sin y + K$ is a potential function for $\mathbf{F}$.

8. $\partial(2xy + y^{-2})/\partial y = 2x - 2y^{-3} = \partial(x^2 - 2xy^{-3})/\partial x$ and the domain of $\mathbf{F}$ is $\{(x, y) \mid y > 0\}$ which is open and simply-connected. Hence $\mathbf{F}$ is conservative, so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x, y) = 2xy + y^{-2}$ implies $f(x, y) = x^2y + xy^{-2} + g(y)$ and $f_y(x, y) = x^2 - 2xy^{-3} + g'(y)$. But $f_y(x, y) = x^2 - 2xy^{-3}$ so $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$. Then $f(x, y) = x^2y + xy^{-2} + K$ is a potential function for $\mathbf{F}$.

9. $\partial(y^2\cos x + \cos y)/\partial y = 2y\cos x - \sin y = \partial(2y\sin x - x\sin y)/\partial x$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$ which is open and simply connected. Hence $\mathbf{F}$ is conservative so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x, y) = y^2\cos x + \cos y$ implies $f(x, y) = y^2\sin x + x\cos y + g(y)$ and $f_y(x, y) = 2y\sin x - x\sin y + g'(y)$. But $f_y(x, y) = 2y\sin x - x\sin y$ so $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$ and $f(x, y) = y^2\sin x + x\cos y + K$ is a potential function for $\mathbf{F}$.

10. $\partial(\ln y + y/x)/\partial y = 1/y + 1/x = \partial(\ln x + x/y)/\partial x$ and the domain of $\mathbf{F}$ is $\{(x, y) \mid x > 0,\ y > 0\}$ which is open and simply connected. Hence $\mathbf{F}$ is conservative so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x, y) = \ln y + y/x$ implies $f(x, y) = x\ln y + y\ln x + g(y)$ and $f_y(x, y) = x/y + \ln x + g'(y)$. But $f_y(x, y) = \ln x + x/y$ so $g'(y) = 0 \quad \Rightarrow$ $g(y) = K$ and $f(x, y) = x\ln y + y\ln x + K$ is a potential function for $\mathbf{F}$.

11. (a) $\mathbf{F}$ has continuous first-order partial derivatives and $\dfrac{\partial}{\partial y}(2xy) = 2x = \dfrac{\partial}{\partial x}(x^2)$ on $\mathbb{R}^2$, which is open and simply-connected. Thus, $\mathbf{F}$ is conservative by Theorem 6. Then we know that the line integral of $\mathbf{F}$ is independent of path; in particular, the value of $\int_C \mathbf{F}\cdot d\mathbf{r}$ depends only on the endpoints of C. Since all three curves have the same initial and terminal points, $\int_C \mathbf{F}\cdot d\mathbf{r}$ will have the same value for each curve

(b) We first find a potential function f, so that $\nabla f = \mathbf{F}$. We know $f_x(x, y) = 2xy$ and $f_y(x, y) = x^2$. Integrating $f_x(x, y)$ with respect to x, we have $f(x, y) = x^2y + g(y)$. Differentiating both sides with respect to y gives $f_y(x, y) = x^2 + g'(y)$, so we must have $x^2 + g'(y) = x^2 \quad \Rightarrow \quad g'(y) = 0 \quad \Rightarrow \quad g(y) = K$, a constant. Thus $f(x, y) = x^2y + K$, and we can take $K = 0$. All three curves start at $(1, 2)$ and end at $(3, 2)$, so by Theorem 2, $\int_C \mathbf{F}\cdot d\mathbf{r} = f(3, 2) - f(1, 2) = 18 - 2 = 16$ for each curve.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

12. (a) If $\mathbf{F} = \nabla f$ then $f_x(x, y) = 3 + 2xy^2$ and $f_y(x, y) = 2x^2y$.

$f_x(x, y) = 3 + 2xy^2$ implies $f(x, y) = 3x + x^2y^2 + g(y)$ and $f_y(x, y) = 2x^2y + g'(y)$. But $f_y(x, y) = 2x^2y$ so $g'(y) = 0 \Rightarrow g(y) = K$. We can take $K = 0$, so $f(x, y) = 3x + x^2y^2$.

(b) C is a smooth curve with initial point $(1, 1)$ and terminal point $\left(4, \frac{1}{4}\right)$, so by Theorem 2

$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f\left(4, \frac{1}{4}\right) - f(1, 1) = (12 + 1) - (3 + 1) = 9.$

13. (a) If $\mathbf{F} = \nabla f$ then $f_x(x, y) = x^2y^3$ and $f_y(x, y) = x^3y^2$.

$f_x(x, y) = x^2y^3$ implies $f(x, y) = \frac{1}{3}x^3y^3 + g(y)$ and $f_y(x, y) = x^3y^2 + g'(y)$. But $f_y(x, y) = x^3y^2$ so $g'(y) = 0 \Rightarrow g(y) = K$, a constant. We can take $K = 0$, so $f(x, y) = \frac{1}{3}x^3y^3$.

(b) C is a smooth curve with initial point $\mathbf{r}(0) = (0, 0)$ and terminal point $\mathbf{r}(1) = (-1, 3)$, so by Theorem 2

$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(-1, 3) - f(0, 0) = -9 - 0 = -9.$

14. (a) $f_y(x, y) = x^2e^{xy}$ implies $f(x, y) = xe^{xy} + g(x) \Rightarrow f_x(x, y) = xye^{xy} + e^{xy} + g'(x) = (1 + xy)e^{xy} + g'(x)$. But $f_x(x, y) = (1 + xy)e^{xy}$ so $g'(x) = 0 \Rightarrow g(x) = K$. We can take $K = 0$, so $f(x, y) = xe^{xy}$.

(b) The initial point of C is $\mathbf{r}(0) = (1, 0)$ and the terminal point is $\mathbf{r}(\pi/2) = (0, 2)$, so

$\int_C \mathbf{F} \cdot d\mathbf{r} = f(0, 2) - f(1, 0) = 0 - e^0 = -1.$

15. (a) $f_x(x, y, z) = yz$ implies $f(x, y, z) = xyz + g(y, z)$ and so $f_y(x, y, z) = xz + g_y(y, z)$. But $f_y(x, y, z) = xz$ so $g_y(y, z) = 0 \Rightarrow g(y, z) = h(z)$. Thus $f(x, y, z) = xyz + h(z)$ and $f_z(x, y, z) = xy + h'(z)$. But $f_z(x, y, z) = xy + 2z$, so $h'(z) = 2z \Rightarrow h(z) = z^2 + K$. Hence $f(x, y, z) = xyz + z^2$ (taking $K = 0$).

(b) $\int_C \mathbf{F} \cdot d\mathbf{r} = f(4, 6, 3) - f(1, 0, -2) = 81 - 4 = 77.$

16. (a) $f_x(x, y, z) = y^2z + 2xz^2$ implies $f(x, y, z) = xy^2z + x^2z^2 + g(y, z)$ and so $f_y(x, y, z) = 2xyz + g_y(y, z)$. But $f_y(x, y, z) = 2xyz$ so $g_y(y, z) = 0 \Rightarrow g(y, z) = h(z)$. Thus $f(x, y, z) = xy^2z + x^2z^2 + h(z)$ and $f_z(x, y, z) = xy^2 + 2x^2z + h'(z)$. But $f_z(x, y, z) = xy^2 + 2x^2z$, so $h'(z) = 0 \Rightarrow h(z) = K$. Hence $f(x, y, z) = xy^2z + x^2z^2$ (taking $K = 0$).

(b) $t = 0$ corresponds to the point $(0, 1, 0)$ and $t = 1$ corresponds to $(1, 2, 1)$, so

$\int_C \mathbf{F} \cdot d\mathbf{r} = f(1, 2, 1) - f(0, 1, 0) = 5 - 0 = 5.$

17. (a) $f_x(x, y, z) = yze^{xz}$ implies $f(x, y, z) = ye^{xz} + g(y, z)$ and so $f_y(x, y, z) = e^{xz} + g_y(y, z)$. But $f_y(x, y, z) = e^{xz}$ so $g_y(y, z) = 0 \Rightarrow g(y, z) = h(z)$. Thus $f(x, y, z) = ye^{xz} + h(z)$ and $f_z(x, y, z) = xye^{xz} + h'(z)$. But $f_z(x, y, z) = xye^{xz}$, so $h'(z) = 0 \Rightarrow h(z) = K$. Hence $f(x, y, z) = ye^{xz}$ (taking $K = 0$).

(b) $\mathbf{r}(0) = \langle 1, -1, 0\rangle$, $\mathbf{r}(2) = \langle 5, 3, 0\rangle$ so $\int_C \mathbf{F} \cdot d\mathbf{r} = f(5, 3, 0) - f(1, -1, 0) = 3e^0 + e^0 = 4.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. (a) $f_x(x,y,z) = \sin y$ implies $f(x,y,z) = x\sin y + g(y,z)$ and so $f_y(x,y,z) = x\cos y + g_y(y,z)$. But $f_y(x,y,z) = x\cos y + \cos z$ so $g_y(y,z) = \cos z \;\Rightarrow\; g(y,z) = y\cos z + h(z)$. Thus $f(x,y,z) = x\sin y + y\cos z + h(z)$ and $f_z(x,y,z) = -y\sin z + h'(z)$. But $f_z(x,y,z) = -y\sin z$, so $h'(z) = 0 \;\Rightarrow\; h(z) = K$. Hence $f(x,y,z) = x\sin y + y\cos z$ (taking $K = 0$).

(b) $\mathbf{r}(0) = \langle 0,0,0\rangle$, $\mathbf{r}(\pi/2) = \langle 1, \pi/2, \pi\rangle$ so $\int_C \mathbf{F}\cdot d\mathbf{r} = f(1,\pi/2,\pi) - f(0,0,0) = 1 - \frac{\pi}{2} \quad 0 = 1 - \frac{\pi}{2}$.

19. The functions $2xe^{-y}$ and $2y - x^2e^{-y}$ have continuous first-order derivatives on $\mathbb{R}^2$ and $\dfrac{\partial}{\partial y}\left(2xe^{-y}\right) = -2xe^{-y} = \dfrac{\partial}{\partial x}\left(2y - x^2e^{-y}\right)$, so $\mathbf{F}(x,y) = 2xe^{-y}\,\mathbf{i} + \left(2y - x^2e^{-y}\right)\mathbf{j}$ is a conservative vector field by Theorem 6 and hence the line integral is independent of path. Thus a potential function f exists, and $f_x(x,y) = 2xe^{-y}$ implies $f(x,y) = x^2e^{-y} + g(y)$ and $f_y(x,y) = -x^2e^{-y} + g'(y)$. But $f_y(x,y) = 2y - x^2e^{-y}$ so $g'(y) = 2y \;\Rightarrow\; g(y) = y^2 + K$. We can take $K = 0$, so $f(x,y) = x^2e^{-y} + y^2$. Then $\int_C 2xe^{-y}\,dx + (2y - x^2e^{-y})\,dy = f(2,1) - f(1,0) = 4e^{-1} + 1 - 1 = 4/e$.

20. The functions $\sin y$ and $x\cos y - \sin y$ have continuous first-order derivatives on $\mathbb{R}^2$ and $\dfrac{\partial}{\partial y}(\sin y) = \cos y = \dfrac{\partial}{\partial x}(x\cos y - \sin y)$, so $\mathbf{F}(x,y) = \sin y\,\mathbf{i} + (x\cos y - \sin y)\,\mathbf{j}$ is a conservative vector field by Theorem 6 and hence the line integral is independent of path. Thus a potential function f exists, and $f_x(x,y) = \sin y$ implies $f(x,y) = x\sin y + g(y)$ and $f_y(x,y) = x\cos y + g'(y)$. But $f_y(x,y) = x\cos y - \sin y$ so $g'(y) = -\sin y \;\Rightarrow\; g(y) = \cos y + K$. We can take $K = 0$, so $f(x,y) = x\sin y + \cos y$. Then $\int_C \sin y\,dx + (x\cos y - \sin y)\,dy = f(1,\pi) - f(2,0) = -1 - 1 = -2$.

21. If $\mathbf{F}$ is conservative, then $\int_C \mathbf{F}\cdot d\mathbf{r}$ is independent of path. This means that the work done along all piecewise-smooth curves that have the described initial and terminal points is the same. Your reply: It doesn't matter which curve is chosen.

22. The curves C_1 and C_2 connect the same two points but $\int_{C_1}\mathbf{F}\cdot d\mathbf{r} \neq \int_{C_2}\mathbf{F}\cdot d\mathbf{r}$. Thus $\mathbf{F}$ is not independent of path, and therefore is not conservative.

23. $\mathbf{F}(x,y) = x^3\,\mathbf{i} + y^3\,\mathbf{j}$, $W = \int_C \mathbf{F}\cdot d\mathbf{r}$. Since $\partial(x^3)/\partial y = 0 = \partial(y^3)/\partial x$, there exists a function f such that $\nabla f = \mathbf{F}$. In fact, $f_x(x,y) = x^3 \;\Rightarrow\; f(x,y) = \frac{1}{4}x^4 + g(y) \;\Rightarrow\; f_y(x,y) = 0 + g'(y)$. But $f_y(x,y) = y^3$ so $g'(y) = y^3 \;\Rightarrow\; g(y) = \frac{1}{4}y^4 + K$. We can take $K = 0 \;\Rightarrow\; f(x,y) = \frac{1}{4}x^4 + \frac{1}{4}y^4$. Thus $W = \int_C \mathbf{F}\cdot d\mathbf{r} = f(2,2) - f(1,0) = (4+4) - \left(\frac{1}{4} + 0\right) = \frac{31}{4}$.

24. $\mathbf{F}(x,y) = (2x+y)\,\mathbf{i} + x\,\mathbf{j}$, $W = \int_C \mathbf{F}\cdot d\mathbf{r}$. Since $\partial(2x+y)/\partial y = 1 = \partial(x)/\partial x$, there exists a function f such that $\nabla f = \mathbf{F}$. In fact, $f_x(x,y) = 2x + y \;\Rightarrow\; f(x,y) = x^2 + xy + g(y) \;\Rightarrow\; f_y(x,y) = x + g'(y)$. But $f_y(x,y) = x$ so $g'(y) = 0 \;\Rightarrow\; g(y) = K$. We can take $K = 0 \;\Rightarrow\; f(x,y) = x^2 + xy$. Thus $W = \int_C \mathbf{F}\cdot d\mathbf{r} = f(4,3) - f(1,1) = (16+12) - (1+1) = 26$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

25. We know that if the vector field (call it $\mathbf{F}$) is conservative, then around any closed path C, $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$. But take C to be a circle centered at the origin, oriented counterclockwise. All of the field vectors that start on C are roughly in the direction of motion along C, so the integral around C will be positive. Therefore the field is not conservative.

26. If a vector field $\mathbf{F}$ is conservative, then around any closed path C, $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$. For any closed path we draw in the field, it appears that some vectors on the curve point in approximately the same direction as the curve and a similar number point in roughly the opposite direction. (Some appear perpendicular to the curve as well.) Therefore it is plausible that $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$ for every closed curve C which means $\mathbf{F}$ is conservative.

27.



From the graph, it appears that $\mathbf{F}$ is conservative, since around all closed paths, the number and size of the field vectors pointing in directions similar to that of the path seem to be roughly the same as the number and size of the vectors pointing in the opposite direction. To check, we calculate $\dfrac{\partial}{\partial y}(\sin y) = \cos y = \dfrac{\partial}{\partial x}(1 + x\cos y)$. Thus $\mathbf{F}$ is conservative, by Theorem 6.

28. $\nabla f(x, y) = \cos(x - 2y)\,\mathbf{i} - 2\cos(x - 2y)\,\mathbf{j}$

(a) We use Theorem 2: $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \nabla f \cdot d\mathbf{r} = f(\mathbf{r}(b)) - f(\mathbf{r}(a))$ where C_1 starts at $t = a$ and ends at $t = b$. So because $f(0, 0) = \sin 0 = 0$ and $f(\pi, \pi) = \sin(\pi - 2\pi) = 0$, one possible curve C_1 is the straight line from $(0, 0)$ to (π, π); that is, $\mathbf{r}(t) = \pi t\,\mathbf{i} + \pi t\,\mathbf{j},\ 0 \le t \le 1$.

(b) From (a), $\int_{C_2} \mathbf{F} \cdot d\mathbf{r} = f(\mathbf{r}(b)) - f(\mathbf{r}(a))$. So because $f(0, 0) = \sin 0 = 0$ and $f\left(\frac{\pi}{2}, 0\right) = 1$, one possible curve C_2 is $\mathbf{r}(t) = \frac{\pi}{2} t\,\mathbf{i}$, $0 \le t \le 1$, the straight line from $(0, 0)$ to $\left(\frac{\pi}{2}, 0\right)$.

29. Since $\mathbf{F}$ is conservative, there exists a function f such that $\mathbf{F} = \nabla f$, that is, $P = f_x$, $Q = f_y$, and $R = f_z$. Since P, Q, and R have continuous first order partial derivatives, Clairaut's Theorem says that $\partial P/\partial y = f_{xy} = f_{yx} = \partial Q/\partial x$, $\partial P/\partial z = f_{xz} = f_{zx} = \partial R/\partial x$, and $\partial Q/\partial z = f_{yz} = f_{zy} = \partial R/\partial y$.

30. Here $\mathbf{F}(x, y, z) = y\,\mathbf{i} + x\,\mathbf{j} + xyz\,\mathbf{k}$. Then using the notation of Exercise 29, $\partial P/\partial z = 0$ while $\partial R/\partial x = yz$. Since these aren't equal, $\mathbf{F}$ is not conservative. Thus by Theorem 4, the line integral of $\mathbf{F}$ is not independent of path.

31. $D = \{(x, y) \mid 0 < y < 3\}$ consists of those points between, but not on, the horizontal lines $y = 0$ and $y = 3$.



(a) Since D does not include any of its boundary points, it is open. More formally, at any point in D there is a disk centered at that point that lies entirely in D.

(b) Any two points chosen in D can always be joined by a path that lies entirely in D, so D is connected. (D consists of just one "piece.")

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) D is connected and it has no holes, so it's simply-connected. (Every simple closed curve in D encloses only points that are in D.)

32. $D = \{(x, y) \mid 1 < |x| < 2\}$ consists of those points between, but not on, the vertical lines $x = 1$ and $x = 2$, together with the points between the vertical lines $x = -1$ and $x = -2$.



(a) The region does not include any of its boundary points, so it is open.

(b) D consists of two separate pieces, so it is not connected. [For instance, both the points $(-1.5, 0)$ and $(1.5, 0)$ lie in D but they cannot be joined by a path that lies entirely in D.]

(c) Because D is not connected, it's not simply-connected.

33. $D = \{(x, y) \mid 1 \le x^2 + y^2 \le 4,\ y \ge 0\}$ is the semiannular region in the upper half-plane between circles centered at the origin of radii 1 and 2 (including all boundary points).



(a) D includes boundary points, so it is not open. [Note that at any boundary point, $(1, 0)$ for instance, any disk centered there cannot lie entirely in D.]

(b) The region consists of one piece, so it's connected.

(c) D is connected and has no holes, so it's simply-connected.

34. $D = \{(x, y) \mid (x, y) \ne (2, 3)\}$ consists of all points in the xy-plane except for $(2, 3)$.



(a) D has only one boundary point, namely $(2, 3)$, which is not included, so the region is open.

(b) D is connected, as it consists of only one piece.

(c) D is not simply-connected, as it has a hole at $(2, 3)$. Thus any simple closed curve that encloses $(2, 3)$ lies in D but includes a point that is not in D.

35. (a) $P = -\dfrac{y}{x^2 + y^2}$, $\dfrac{\partial P}{\partial y} = \dfrac{y^2 - x^2}{(x^2 + y^2)^2}$ and $Q = \dfrac{x}{x^2 + y^2}$, $\dfrac{\partial Q}{\partial x} = \dfrac{y^2 - x^2}{(x^2 + y^2)^2}$. Thus $\dfrac{\partial P}{\partial y} = \dfrac{\partial Q}{\partial x}$.

(b) C_1: $x = \cos t$, $y = \sin t$, $0 \le t \le \pi$, C_2: $x = \cos t$, $y = \sin t$, $t = 2\pi$ to $t = \pi$. Then

$$\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_0^{\pi} \frac{(-\sin t)(-\sin t) + (\cos t)(\cos t)}{\cos^2 t + \sin^2 t}\, dt = \int_0^{\pi} dt = \pi \text{ and } \int_{C_2} \mathbf{F} \cdot d\mathbf{r} = \int_{2\pi}^{\pi} dt = -\pi$$

Since these aren't equal, the line integral of $\mathbf{F}$ isn't independent of path. (Or notice that $\int_{C_3} \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} dt = 2\pi$ where C_3 is the circle $x^2 + y^2 = 1$, and apply the contrapositive of Theorem 3.) This doesn't contradict Theorem 6, since the domain of $\mathbf{F}$, which is $\mathbb{R}^2$ except the origin, isn't simply-connected.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

36. (a) Here $\mathbf{F}(\mathbf{r}) = c\mathbf{r}/|\mathbf{r}|^3$ and $\mathbf{r} = x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$. Then $f(\mathbf{r}) = -c/|\mathbf{r}|$ is a potential function for $\mathbf{F}$, that is, $\nabla f = \mathbf{F}$. (See the discussion of gradient fields in Section 16.1.) Hence $\mathbf{F}$ is conservative and its line integral is independent of path. Let $P_1 = (x_1, y_1, z_1)$ and $P_2 = (x_2, y_2, z_2)$.

$$W = \int_C \mathbf{F}\cdot d\mathbf{r} = f(P_2) - f(P_1) = -\frac{c}{(x_2^2+y_2^2+z_2^2)^{1/2}} + \frac{c}{(x_1^2+y_1^2+z_1^2)^{1/2}} = c\left(\frac{1}{d_1} - \frac{1}{d_2}\right).$$

(b) In this case, $c = -(mMG) \Rightarrow$

$$W = -mMG\left(\frac{1}{1.52\times10^{11}} - \frac{1}{1.47\times10^{11}}\right)$$
$$= -(5.97\times10^{24})(1.99\times10^{30})(6.67\times10^{-11})(-2.2377\times10^{-13}) \approx 1.77\times10^{32}\ \text{J}$$

(c) In this case, $c = \epsilon qQ \Rightarrow$

$$W = \epsilon qQ\left(\frac{1}{10^{-12}} - \frac{1}{5\times10^{-13}}\right) = (8.985\times10^{9})(1)(-1.6\times10^{-19})(-10^{12}) \approx 1400\ \text{J}.$$

16.4 Green's Theorem

1. (a)



C_1: $x = t \Rightarrow dx = dt$, $y = 0 \Rightarrow dy = 0\,dt$, $0 \le t \le 5$.

C_2: $x = 5 \Rightarrow dx = 0\,dt$, $y = t \Rightarrow dy = dt$, $0 \le t \le 4$.

C_3: $x = 5 - t \Rightarrow dx = -dt$, $y = 4 \Rightarrow dy = 0\,dt$, $0 \le t \le 5$.

C_4: $x = 0 \Rightarrow dx = 0\,dt$, $y = 4 - t \Rightarrow dy = -dt$, $0 \le t \le 4$

Thus $\oint_C y^2\,dx + x^2y\,dy = \oint_{C_1+C_2+C_3+C_4} y^2\,dx + x^2y\,dy = \int_0^5 0\,dt + \int_0^4 25t\,dt + \int_0^5(-16+0)\,dt + \int_0^4 0\,dt$

$$= 0 + \left[\tfrac{25}{2}t^2\right]_0^4 + [-16t]_0^5 + 0 = 200 + (-80) = 120$$

(b) Note that C as given in part (a) is a positively oriented, piecewise-smooth, simple closed curve. Then by Green's Theorem,

$$\oint_C y^2\,dx + x^2y\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(x^2y) - \tfrac{\partial}{\partial y}(y^2)\right] dA = \int_0^5\int_0^4 (2xy - 2y)\,dy\,dx = \int_0^5 \left[xy^2 - y^2\right]_{y=0}^{y=4} dx$$
$$= \int_0^5 (16x - 16)\,dx = \left[8x^2 - 16x\right]_0^5 = 200 - 80 = 120$$

2. (a) Parametric equations for C are $x = 4\cos t$, $y = 4\sin t$, $0 \le t \le 2\pi$. Then $dx = -4\sin t\,dt$, $dy = 4\cos t\,dt$ and

$$\oint_C y\,dx - x\,dy = \int_0^{2\pi}[(4\sin t)(-4\sin t) - (4\cos t)(4\cos t)]\,dt$$
$$= -16\int_0^{2\pi}(\sin^2 t + \cos^2 t)\,dt = -16\int_0^{2\pi} 1\,dt = -16(2\pi) = -32\pi$$

(b) Note that C as given in part (a) is a positively oriented, smooth, simple closed curve. Then by Green's Theorem,

$$\oint_C y\,dx - x\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(-x) - \tfrac{\partial}{\partial y}(y)\right] dA = \iint_D (-1-1)\,dA = -2\iint_D dA$$
$$-2(\text{area of } D) = -2\cdot\pi(4)^2 = -32\pi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a)



C_1: $x = t \Rightarrow dx = dt,\ y = 0 \Rightarrow dy = 0\,dt,\ 0 \le t \le 1.$

C_2: $x = 1 \Rightarrow dx = 0\,dt,\ y = t \Rightarrow dy = dt,\ 0 \le t \le 2.$

C_3: $x = 1 - t \Rightarrow dx = -dt,\ y = 2 - 2t \Rightarrow dy = -2\,dt,\ 0 \le t \le 1.$

Thus

$$\begin{aligned}\oint_C xy\,dx + x^2y^3\,dy &= \oint_{C_1+C_2+C_3} xy\,dx + x^2y^3\,dy \\ &= \int_0^1 0\,dt + \int_0^2 t^3\,dt + \int_0^1 \left[-(1-t)(2-2t) - 2(1-t)^2(2-2t)^3\right] dt \\ &= 0 + \left[\tfrac{1}{4}t^4\right]_0^2 + \int_0^1 \left[-2(1-t)^2 - 16(1-t)^5\right] dt \\ &= 4 + \left[\tfrac{2}{3}(1-t)^3 + \tfrac{8}{3}(1-t)^6\right]_0^1 = 4 + 0 - \tfrac{10}{3} = \tfrac{2}{3}\end{aligned}$$

(b) $\oint_C xy\,dx + x^2y^3\,dy = \iint_D \left[\frac{\partial}{\partial x}(x^2y^3) - \frac{\partial}{\partial y}(xy)\right] dA = \int_0^1 \int_0^{2x} (2xy^3 - x)\,dy\,dx$

$$= \int_0^1 \left[\tfrac{1}{2}xy^4 - xy\right]_{y=0}^{y=2x} dx = \int_0^1 (8x^5 - 2x^2)\,dx = \tfrac{4}{3} - \tfrac{2}{3} = \tfrac{2}{3}$$

4. (a)



C_1: $x = t \Rightarrow dx = dt,\ y = t^2 \Rightarrow dy = 2t\,dt,\ 0 \le t \le 1$

C_2: $x = 1 - t \Rightarrow dx = -dt,\ y = 1 \Rightarrow dy = 0\,dt,\ 0 \le t \le 1$

C_3: $x = 0 \Rightarrow dx = 0\,dt,\ y = 1 - t \Rightarrow dy = -dt,\ 0 \le t \le 1$

Thus

$$\begin{aligned}\oint_C x^2y^2\,dx + xy\,dy &= \oint_{C_1+C_2+C_3} x^2y^2\,dx + xy\,dy \\ &= \int_0^1 \left[t^2(t^2)^2\,dt + t(t^2)(2t\,dt)\right] + \int_0^1 \left[(1-t)^2(1)^2(-dt) + (1-t)(1)(0\,dt)\right] \\ &\qquad + \int_0^1 \left[(0)^2(1-t)^2(0\,dt) + (0)(1-t)(-dt)\right] \\ &= \int_0^1 \left(t^6 + 2t^4\right) dt + \int_0^1 \left(-1 + 2t - t^2\right) dt + \int_0^1 0\,dt \\ &= \left[\tfrac{1}{7}t^7 + \tfrac{2}{5}t^5\right]_0^1 + \left[-t + t^2 - \tfrac{1}{3}t^3\right]_0^1 + 0 = \left(\tfrac{1}{7} + \tfrac{2}{5}\right) + \left(-1 + 1 - \tfrac{1}{3}\right) = \tfrac{22}{105}\end{aligned}$$

(b) $\oint_C x^2y^2\,dx + xy\,dy = \iint_D \left[\frac{\partial}{\partial x}(xy) - \frac{\partial}{\partial y}(x^2y^2)\right] dA = \int_0^1 \int_{x^2}^1 (y - 2x^2y)\,dy\,dx$

$$= \int_0^1 \left[\tfrac{1}{2}y^2 - x^2y^2\right]_{y=x^2}^{y=1} dx = \int_0^1 \left(\tfrac{1}{2} - x^2 - \tfrac{1}{2}x^4 + x^6\right) dx$$

$$= \left[\tfrac{1}{2}x - \tfrac{1}{3}x^3 - \tfrac{1}{10}x^5 + \tfrac{1}{7}x^7\right]_0^1 = \tfrac{1}{2} - \tfrac{1}{3} - \tfrac{1}{10} + \tfrac{1}{7} = \tfrac{22}{105}$$

5. The region D enclosed by C is $[0, 3] \times [0, 4]$, so

$$\begin{aligned}\int_C ye^x\,dx + 2e^x\,dy &= \iint_D \left[\frac{\partial}{\partial x}(2e^x) - \frac{\partial}{\partial y}(ye^x)\right] dA = \int_0^3 \int_0^4 (2e^x - e^x)\,dy\,dx \\ &= \int_0^3 e^x\,dx \int_0^4 dy = \left[e^x\right]_0^3 \left[y\right]_0^4 = (e^3 - e^0)(4 - 0) = 4(e^3 - 1)\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

6.



The region D enclosed by C is given by $\{(x, y) \mid 0 \le y \le 1,\ 0 \le x \le 2y\}$, so

$$\begin{aligned}\int_C (x^2+y^2)\,dx + (x^2-y^2)\,dy &= \iint_D \left[\tfrac{\partial}{\partial x}(x^2-y^2) - \tfrac{\partial}{\partial y}(x^2+y^2)\right] dA \\ &= \int_0^1 \int_0^{2y} (2x-2y)\,dx\,dy \\ &= \int_0^1 \left[x^2 - 2xy\right]_{x=0}^{x=2y} dy \\ &= \int_0^1 (4y^2-4y^2)\,dy = \int_0^1 0\,dy = 0\end{aligned}$$

7. $\int_C \left(y + e^{\sqrt{x}}\right) dx + (2x + \cos y^2)\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(2x+\cos y^2) - \tfrac{\partial}{\partial y}\left(y + e^{\sqrt{x}}\right)\right] dA$

$$= \int_0^1 \int_{x^2}^{\sqrt{x}} (2-1)\,dy\,dx = \int_0^1 (\sqrt{x} - x^2)\,dx = \left[\tfrac{2}{3}x^{3/2} - \tfrac{1}{3}x^3\right]_0^1 = \tfrac{1}{3}$$

8. $\int_C y^4\,dx + 2xy^3\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(2xy^3) - \tfrac{\partial}{\partial y}(y^4)\right] dA = \iint_D (2y^3 - 4y^3)\,dA$

$$= -2\iint_D y^3\,dA = 0$$

because $f(x, y) = y^3$ is an odd function with respect to y and D is symmetric about the x-axis.

9. $\int_C y^3\,dx - x^3\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(-x^3) - \tfrac{\partial}{\partial y}(y^3)\right] dA = \iint_D (-3x^2 - 3y^2)\,dA = \int_0^{2\pi} \int_0^2 (-3r^2)\, r\,dr\,d\theta$

$$= -3\int_0^{2\pi} d\theta \int_0^2 r^3\,dr = -3\left[\theta\right]_0^{2\pi} \left[\tfrac{1}{4}r^4\right]_0^2 = -3(2\pi)(4) = -24\pi$$

10. $\int_C (1-y^3)\,dx + (x^3 + e^{y^2})\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(x^3 + e^{y^2}) - \tfrac{\partial}{\partial y}(1-y^3)\right] dA = \iint_D (3x^2 + 3y^2)\,dA$

$$= \int_0^{2\pi} \int_2^3 (3r^2)\, r\,dr\,d\theta = 3\int_0^{2\pi} d\theta \int_2^3 r^3\,dr$$

$$= 3\left[\theta\right]_0^{2\pi} \left[\tfrac{1}{4}r^4\right]_2^3 = 3(2\pi)\cdot\tfrac{1}{4}(81-16) = \tfrac{195}{2}\pi$$

11. $\mathbf{F}(x, y) = \langle y\cos x - xy\sin x,\ xy + x\cos x\rangle$ and the region D enclosed by C is given by $\{(x, y) \mid 0 \le x \le 2,\ 0 \le y \le 4 - 2x\}$. C is traversed clockwise, so $-C$ gives the positive orientation.

$$\int_C \mathbf{F}\cdot d\mathbf{r} = -\int_{-C} (y\cos x - xy\sin x)\,dx + (xy + x\cos x)\,dy = -\iint_D \left[\tfrac{\partial}{\partial x}(xy + x\cos x) - \tfrac{\partial}{\partial y}(y\cos x - xy\sin x)\right] dA$$

$$= -\iint_D (y - x\sin x + \cos x - \cos x + x\sin x)\,dA = -\int_0^2 \int_0^{4-2x} y\,dy\,dx$$

$$= -\int_0^2 \left[\tfrac{1}{2}y^2\right]_{y=0}^{y=4-2x} dx = -\int_0^2 \tfrac{1}{2}(4-2x)^2\,dx = -\int_0^2 (8 - 8x + 2x^2)\,dx = -\left[8x - 4x^2 + \tfrac{2}{3}x^3\right]_0^2$$

$$= -\left(16 - 16 + \tfrac{16}{3} - 0\right) = -\tfrac{16}{3}$$

12. $\mathbf{F}(x, y) = \left\langle e^{-x} + y^2,\ e^{-y} + x^2\right\rangle$ and the region D enclosed by C is given by $\{(x, y) \mid -\pi/2 \le x \le \pi/2, 0 \le y \le \cos x\}$. C is traversed clockwise, so $-C$ gives the positive orientation.

$$\int_C \mathbf{F}\cdot d\mathbf{r} = -\int_{-C} \left(e^{-x} + y^2\right) dx + \left(e^{-y} + x^2\right) dy = -\iint_D \left[\tfrac{\partial}{\partial x}\left(e^{-y} + x^2\right) - \tfrac{\partial}{\partial y}\left(e^{-x} + y^2\right)\right] dA$$

$$= -\int_{-\pi/2}^{\pi/2} \int_0^{\cos x} (2x - 2y)\,dy\,dx = -\int_{-\pi/2}^{\pi/2} \left[2xy - y^2\right]_{y=0}^{y=\cos x} dx$$

$$= -\int_{-\pi/2}^{\pi/2} (2x\cos x - \cos^2 x)\,dx = -\int_{-\pi/2}^{\pi/2} \left[2x\cos x - \tfrac{1}{2}(1 + \cos 2x)\right] dx$$

$$= -\left[2x\sin x + 2\cos x - \tfrac{1}{2}\left(x + \tfrac{1}{2}\sin 2x\right)\right]_{-\pi/2}^{\pi/2} \qquad \text{[integrate by parts in the first term]}$$

$$= -\left(\pi - \tfrac{1}{4}\pi - \pi - \tfrac{1}{4}\pi\right) = \tfrac{1}{2}\pi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $\mathbf{F}(x, y) = \langle y - \cos y, x \sin y \rangle$ and the region D enclosed by C is the disk with radius 2 centered at $(3, -4)$.

C is traversed clockwise, so $-C$ gives the positive orientation.

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= -\int_{-C} (y - \cos y)\,dx + (x \sin y)\,dy = -\iint_D \left[\tfrac{\partial}{\partial x}(x \sin y) - \tfrac{\partial}{\partial y}(y - \cos y)\right] dA \\ &= -\iint_D (\sin y - 1 - \sin y)\,dA = \iint_D dA = \text{area of } D = \pi(2)^2 = 4\pi\end{aligned}$$

14. $\mathbf{F}(x, y) = \left\langle \sqrt{x^2 + 1}, \tan^{-1} x \right\rangle$ and the region D enclosed by C is given by $\{(x, y) \mid 0 \le x \le 1, x \le y \le 1\}$.

C is oriented positively, so

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_C \sqrt{x^2 + 1}\,dx + \tan^{-1} x\,dy = \iint_D \left[\frac{\partial}{\partial x}\left(\tan^{-1} x\right) - \frac{\partial}{\partial y}\left(\sqrt{x^2 + 1}\right)\right] dA \\ &= \int_0^1 \int_x^1 \left(\frac{1}{1 + x^2} - 0\right) dy\,dx = \int_0^1 \frac{1}{1 + x^2}\left[y\right]_{y=x}^{y=1} dx = \int_0^1 \frac{1}{1 + x^2}(1 - x)\,dx \\ &= \int_0^1 \left(\frac{1}{1 + x^2} - \frac{x}{1 + x^2}\right) dx = \left[\tan^{-1} x - \frac{1}{2}\ln(1 + x^2)\right]_0^1 = \frac{\pi}{4} - \frac{1}{2}\ln 2\end{aligned}$$

15. Here $C = C_1 + C_2$ where

C_1 can be parametrized as $x = t,\ y = 0,\ -\pi/2 \le t \le \pi/2$, and

C_2 is given by $x = -t,\ y = \cos t,\ -\pi/2 \le t \le \pi/2$.



Then the line integral is

$$\begin{aligned}\oint_{C_1 + C_2} x^3y^4\,dx + x^5y^4\,dy &= \int_{-\pi/2}^{\pi/2}(0 + 0)\,dt + \int_{-\pi/2}^{\pi/2}[(-t)^3(\cos t)^4(-1) + (-t)^5(\cos t)^4(-\sin t)]\,dt \\ &= 0 + \int_{-\pi/2}^{\pi/2}(t^3\cos^4 t + t^5\cos^4 t \sin t)\,dt = \tfrac{1}{15}\pi^4 - \tfrac{4144}{1125}\pi^2 + \tfrac{7{,}578{,}368}{253{,}125} \approx 0.0779\end{aligned}$$

according to a CAS. The double integral is

$$\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA = \int_{-\pi/2}^{\pi/2}\int_0^{\cos x} (5x^4y^4 - 4x^3y^3)\,dy\,dx = \tfrac{1}{15}\pi^4 - \tfrac{4144}{1125}\pi^2 + \tfrac{7{,}578{,}368}{253{,}125} \approx 0.0779,$$ verifying Green's

Theorem in this case.

16. We can parametrize C as $x = \cos\theta,\ y = 2\sin\theta,\ 0 \le \theta \le 2\pi$. Then the line integral is

$$\begin{aligned}\oint_C P\,dx + Q\,dy &= \int_0^{2\pi}\left[2\cos\theta - (\cos\theta)^3(2\sin\theta)^5\right](-\sin\theta)\,d\theta + \int_0^{2\pi}(\cos\theta)^3(2\sin\theta)^8 \cdot 2\cos\theta\,d\theta \\ &= \int_0^{2\pi}(-2\cos\theta\sin\theta + 32\cos^3\theta\sin^6\theta + 512\cos^4\theta\sin^8\theta)\,d\theta = 7\pi,\end{aligned}$$

according to a CAS. The double integral is $\displaystyle\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA = \int_{-1}^{1}\int_{-\sqrt{4 - 4x^2}}^{\sqrt{4 - 4x^2}} (3x^2y^8 + 5x^3y^4)\,dy\,dx = 7\pi.$

17. By Green's Theorem, $W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C x(x + y)\,dx + xy^2\,dy = \iint_D (y^2 - x)\,dA$ where C is the path described in the question and D is the triangle bounded by C. So

$$\begin{aligned}W &= \int_0^1\int_0^{1-x}(y^2 - x)\,dy\,dx = \int_0^1\left[\tfrac{1}{3}y^3 - xy\right]_{y=0}^{y=1-x} dx = \int_0^1\left(\tfrac{1}{3}(1 - x)^3 - x(1 - x)\right) dx \\ &= \left[-\tfrac{1}{12}(1 - x)^4 - \tfrac{1}{2}x^2 + \tfrac{1}{3}x^3\right]_0^1 = \left(-\tfrac{1}{2} + \tfrac{1}{3}\right) - \left(-\tfrac{1}{12}\right) = -\tfrac{1}{12}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

18. By Green's Theorem, $W = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \sin x\,dx + \left(\sin y + xy^2 + \frac{1}{3}x^3\right) dy = \iint_D (y^2 + x^2 - 0)\,dA$, where D is the region (a quarter-disk) bounded by C. Converting to polar coordinates, we have

$W = \int_0^{\pi/2} \int_0^5 r^2 \cdot r\,dr\,d\theta = \left[\theta\right]_0^{\pi/2} \left[\frac{1}{4}r^4\right]_0^5 = \frac{1}{2}\pi\left(\frac{625}{4}\right) = \frac{625}{8}\pi.$

19. Let C_1 be the arch of the cycloid from $(0, 0)$ to $(2\pi, 0)$, which corresponds to $0 \le t \le 2\pi$, and let C_2 be the segment from $(2\pi, 0)$ to $(0, 0)$, so C_2 is given by $x = 2\pi - t$, $y = 0$, $0 \le t \le 2\pi$. Then $C = C_1 \cup C_2$ is traversed clockwise, so $-C$ is oriented positively. Thus $-C$ encloses the area under one arch of the cycloid and from (5) we have

$$\begin{aligned} A = -\oint_{-C} y\,dx &= \int_{C_1} y\,dx + \int_{C_2} y\,dx = \int_0^{2\pi} (1 - \cos t)(1 - \cos t)\,dt + \int_0^{2\pi} 0\,(-dt) \\ &= \int_0^{2\pi} (1 - 2\cos t + \cos^2 t)\,dt + 0 = \left[t - 2\sin t + \tfrac{1}{2}t + \tfrac{1}{4}\sin 2t\right]_0^{2\pi} = 3\pi \end{aligned}$$

20.



$$\begin{aligned} A = \oint_C x\,dy &= \int_0^{2\pi} (5\cos t - \cos 5t)(5\cos t - 5\cos 5t)\,dt \\ &= \int_0^{2\pi} (25\cos^2 t - 30\cos t\cos 5t + 5\cos^2 5t)\,dt \\ &= \left[25\left(\tfrac{1}{2}t + \tfrac{1}{4}\sin 2t\right) - 30\left(\tfrac{1}{8}\sin 4t + \tfrac{1}{12}\sin 6t\right) + 5\left(\tfrac{1}{2}t + \tfrac{1}{20}\sin 10t\right)\right]_0^{2\pi} \end{aligned}$$

[Use Formula 80 in the Table of Integrals]

$$= 30\pi$$

21. (a) Using Equation 16.2.8, we write parametric equations of the line segment as $x = (1 - t)x_1 + tx_2$, $y = (1 - t)y_1 + ty_2$, $0 \le t \le 1$. Then $dx = (x_2 - x_1)\,dt$ and $dy = (y_2 - y_1)\,dt$, so

$$\begin{aligned} \int_C x\,dy - y\,dx &= \int_0^1 [(1 - t)x_1 + tx_2](y_2 - y_1)\,dt + [(1 - t)y_1 + ty_2](x_2 - x_1)\,dt \\ &= \int_0^1 \left(x_1(y_2 - y_1) - y_1(x_2 - x_1) + t[(y_2 - y_1)(x_2 - x_1) - (x_2 - x_1)(y_2 - y_1)]\right) dt \\ &= \int_0^1 (x_1y_2 - x_2y_1)\,dt = x_1y_2 - x_2y_1 \end{aligned}$$

(b) We apply Green's Theorem to the path $C = C_1 \cup C_2 \cup \cdots \cup C_n$, where C_i is the line segment that joins (x_i, y_i) to (x_{i+1}, y_{i+1}) for $i = 1, 2, \ldots, n - 1$, and C_n is the line segment that joins (x_n, y_n) to (x_1, y_1). From (5), $\frac{1}{2}\int_C x\,dy - y\,dx = \iint_D dA$, where D is the polygon bounded by C. Therefore

$$\begin{aligned} \text{area of polygon} = A(D) &= \iint_D dA = \tfrac{1}{2}\int_C x\,dy - y\,dx \\ &= \tfrac{1}{2}\left(\int_{C_1} x\,dy - y\,dx + \int_{C_2} x\,dy - y\,dx + \cdots + \int_{C_{n-1}} x\,dy - y\,dx + \int_{C_n} x\,dy - y\,dx\right) \end{aligned}$$

To evaluate these integrals we use the formula from (a) to get

$A(D) = \frac{1}{2}[(x_1y_2 - x_2y_1) + (x_2y_3 - x_3y_2) + \cdots + (x_{n-1}y_n - x_ny_{n-1}) + (x_ny_1 - x_1y_n)]$.

(c) $A = \frac{1}{2}[(0 \cdot 1 - 2 \cdot 0) + (2 \cdot 3 - 1 \cdot 1) + (1 \cdot 2 - 0 \cdot 3) + (0 \cdot 1 - (-1) \cdot 2) + (-1 \cdot 0 - 0 \cdot 1)]$

$= \frac{1}{2}(0 + 5 + 2 + 2) = \frac{9}{2}$

22. By Green's Theorem, $\frac{1}{2A}\oint_C x^2\,dy = \frac{1}{2A}\iint_D 2x\,dA = \frac{1}{A}\iint_D x\,dA = \overline{x}$ and

$-\frac{1}{2A}\oint_C y^2\,dx = -\frac{1}{2A}\iint_D (-2y)\,dA = \frac{1}{A}\iint_D y\,dA = \overline{y}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

23. We orient the quarter-circular region as shown in the figure.



$A = \frac{1}{4}\pi a^2$ so $\overline{x} = \dfrac{1}{\pi a^2/2}\oint_C x^2\,dy$ and $\overline{y} = -\dfrac{1}{\pi a^2/2}\oint_C y^2 dx$.

Here $C = C_1 + C_2 + C_3$ where C_1: $x = t,\ \ y = 0,\ \ 0 \le t \le a$;

C_2: $x = a\cos t,\ \ y = a\sin t,\ \ 0 \le t \le \frac{\pi}{2}$; and

C_3: $x = 0, y = a - t, 0 \le t \le a$. Then

$$\oint_C x^2\,dy = \int_{C_1} x^2\,dy + \int_{C_2} x^2\,dy + \int_{C_3} x^2\,dy = \int_0^a 0\,dt + \int_0^{\pi/2}(a\cos t)^2(a\cos t)\,dt + \int_0^a 0\,dt$$
$$= \int_0^{\pi/2} a^3\cos^3 t\,dt = a^3\int_0^{\pi/2}(1-\sin^2 t)\cos t\,dt = a^3\left[\sin t - \tfrac{1}{3}\sin^3 t\right]_0^{\pi/2} = \tfrac{2}{3}a^3$$

so $\overline{x} = \dfrac{1}{\pi a^2/2}\oint_C x^2\,dy = \dfrac{4a}{3\pi}$.

$$\oint_C y^2 dx = \int_{C_1} y^2\,dx + \int_{C_2} y^2\,dx + \int_{C_3} y^2\,dx = \int_0^a 0\,dt + \int_0^{\pi/2}(a\sin t)^2(-a\sin t)\,dt + \int_0^a 0\,dt$$
$$= \int_0^{\pi/2}(-a^3\sin^3 t)\,dt = -a^3\int_0^{\pi/2}(1-\cos^2 t)\sin t\,dt = -a^3\left[\tfrac{1}{3}\cos^3 t - \cos t\right]_0^{\pi/2} = -\tfrac{2}{3}a^3.$$

so $\overline{y} = -\dfrac{1}{\pi a^2/2}\oint_C y^2 dx = \dfrac{4a}{3\pi}$. Thus $(\overline{x},\overline{y}) = \left(\dfrac{4a}{3\pi}, \dfrac{4a}{3\pi}\right)$.

24. Here $A = \frac{1}{2}ab$ and $C = C_1 + C_2 + C_3$, where C_1: $x = x, y = 0, 0 \le x \le a$;

C_2: $x = a, y = y, 0 \le y \le b$; and C_3: $x = x, y = \frac{b}{a}x$, $x = a$ to $x = 0$. Then

$$\oint_C x^2\,dy = \int_{C_1} x^2\,dy + \int_{C_2} x^2\,dy + \int_{C_3} x^2\,dy = 0 + \int_0^b a^2\,dy + \int_a^0 (x^2)\left(\tfrac{b}{a}\,dx\right)$$
$$= a^2 b + \tfrac{b}{a}\left[\tfrac{1}{3}x^3\right]_a^0 = a^2 b - \tfrac{1}{3}a^2 b = \tfrac{2}{3}a^2 b.$$

Similarly, $\oint_C y^2\,dx = \int_{C_1} y^2\,dx + \int_{C_2} y^2\,dx + \int_{C_3} y^2\,dx = 0 + 0 + \int_a^0 \left(\frac{b}{a}x\right)^2 dx = \frac{b^2}{a^2}\cdot\frac{1}{3}x^3\Big]_a^0 = -\frac{1}{3}ab^2$. Thus

$\overline{x} = \frac{1}{2A}\oint_C x^2\,dy = \frac{1}{ab}\cdot\frac{2}{3}a^2 b = \frac{2}{3}a$ and $\overline{y} = -\frac{1}{2A}\oint_C y^2\,dx = -\frac{1}{ab}\left(-\frac{1}{3}ab^2\right) = \frac{1}{3}b$, so $(\overline{x},\overline{y}) = \left(\frac{2}{3}a, \frac{1}{3}b\right)$.

25. By Green's Theorem, $-\frac{1}{3}\rho\oint_C y^3\,dx = -\frac{1}{3}\rho\iint_D(-3y^2)\,dA = \iint_D y^2\rho\,dA = I_x$ and

$\frac{1}{3}\rho\oint_C x^3\,dy = \frac{1}{3}\rho\iint_D(3x^2)\,dA = \iint_D x^2\rho\,dA = I_y$.

26. By symmetry the moments of inertia about any two diameters are equal. Centering the disk at the origin, the moment of inertia about a diameter equals

$$I_y = \tfrac{1}{3}\rho\oint_C x^3\,dy = \tfrac{1}{3}\rho\int_0^{2\pi}(a^4\cos^4 t)\,dt = \tfrac{1}{3}a^4\rho\int_0^{2\pi}\left[\tfrac{3}{8} + \tfrac{1}{2}\cos 2t + \tfrac{1}{8}\cos 4t\right]dt = \tfrac{1}{3}a^4\rho\cdot\tfrac{3(2\pi)}{8} = \tfrac{1}{4}\pi a^4\rho$$

27. As in Example 5, let C' be a counterclockwise-oriented circle with center the origin and radius a, where a is chosen to be small enough so that C' lies inside C, and D the region bounded by C and C'. Here

$$P = \frac{2xy}{(x^2+y^2)^2} \quad\Rightarrow\quad \frac{\partial P}{\partial y} = \frac{2x(x^2+y^2)^2 - 2xy\cdot 2(x^2+y^2)\cdot 2y}{(x^2+y^2)^4} = \frac{2x^3 - 6xy^2}{(x^2+y^2)^3} \text{ and}$$

$$Q = \frac{y^2-x^2}{(x^2+y^2)^2} \quad\Rightarrow\quad \frac{\partial Q}{\partial x} = \frac{-2x(x^2+y^2)^2 - (y^2-x^2)\cdot 2(x^2+y^2)\cdot 2x}{(x^2+y^2)^4} = \frac{2x^3 - 6xy^2}{(x^2+y^2)^3}.$$ Thus, as in the example,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$$\int_{C'} P\,dx + Q\,dy + \int_{-C''} P\,dx + Q\,dy = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA = \iint_D 0\,dA = 0$$

and $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_{C'} \mathbf{F}\cdot d\mathbf{r}$. We parametrize C' as $\mathbf{r}(t) = a\cos t\,\mathbf{i} + a\sin t\,\mathbf{j}$, $0 \le t \le 2\pi$. Then

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \int_{C'} \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi} \frac{2(a\cos t)(a\sin t)\,\mathbf{i} + (a^2\sin^2 t - a^2\cos^2 t)\,\mathbf{j}}{(a^2\cos^2 t + a^2\sin^2 t)^2}\cdot\left(-a\sin t\,\mathbf{i} + a\cos t\,\mathbf{j}\right) dt \\ &= \frac{1}{a}\int_0^{2\pi} \left(-\cos t\sin^2 t - \cos^3 t\right) dt = \frac{1}{a}\int_0^{2\pi} \left(-\cos t\sin^2 t - \cos t\left(1-\sin^2 t\right)\right) dt \\ &= -\frac{1}{a}\int_0^{2\pi} \cos t\,dt = -\frac{1}{a}\sin t\Big]_0^{2\pi} = 0\end{aligned}$$

28. P and Q have continuous partial derivatives on $\mathbb{R}^2$, so by Green's Theorem we have

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA = \iint_D (3-1)\,dA = 2\iint_D dA = 2\cdot A(D) = 2\cdot 6 = 12$$

29. Since C is a simple closed path which doesn't pass through or enclose the origin, there exists an open region that doesn't contain the origin but does contain D. Thus $P = -y/(x^2+y^2)$ and $Q = x/(x^2+y^2)$ have continuous partial derivatives on this open region containing D and we can apply Green's Theorem. But by Exercise 16.3.35(a), $\partial P/\partial y = \partial Q/\partial x$, so $\oint_C \mathbf{F}\cdot d\mathbf{r} = \iint_D 0\,dA = 0$.

30. We express D as a type II region: $D = \{(x,y) \mid f_1(y) \le x \le f_2(y),\ c \le y \le d\}$ where f_1 and f_2 are continuous functions. Then $\iint_D \frac{\partial Q}{\partial x}\,dA = \int_c^d \int_{f_1(y)}^{f_2(y)} \frac{\partial Q}{\partial x}\,dx\,dy = \int_c^d [Q(f_2(y),y) - Q(f_1(y),y)]\,dy$ by the Fundamental Theorem of Calculus. But referring to the figure, $\oint_C Q\,dy = \oint_{C_1+C_2+C_3+C_4} Q\,dy$.



Then $\int_{C_1} Q\,dy = \int_d^c Q(f_1(y),y)\,dy$, $\int_{C_2} Q\,dy = \int_{C_4} Q\,dy = 0$, and $\int_{C_3} Q\,dy = \int_c^d Q(f_2(y),y)\,dy$. Hence $\oint_C Q\,dy = \int_c^d [Q(f_2(y),y) - Q(f_1(y),y)]\,dy = \iint_D (\partial Q/\partial x)\,dA$.

31. Using the first part of (5), we have that $\iint_R dx\,dy = A(R) = \int_{\partial R} x\,dy$. But $x = g(u,v)$, and $dy = \frac{\partial h}{\partial u}\,du + \frac{\partial h}{\partial v}\,dv$, and we orient ∂S by taking the positive direction to be that which corresponds, under the mapping, to the positive direction along ∂R, so

$$\begin{aligned}\int_{\partial R} x\,dy &= \int_{\partial S} g(u,v)\left(\frac{\partial h}{\partial u}\,du + \frac{\partial h}{\partial v}\,dv\right) = \int_{\partial S} g(u,v)\frac{\partial h}{\partial u}\,du + g(u,v)\frac{\partial h}{\partial v}\,dv \\ &= \pm\iint_S \left[\tfrac{\partial}{\partial u}\left(g(u,v)\tfrac{\partial h}{\partial v}\right) - \tfrac{\partial}{\partial v}\left(g(u,v)\tfrac{\partial h}{\partial u}\right)\right] dA \qquad \text{[using Green's Theorem in the } uv\text{-plane]} \\ &= \pm\iint_S \left(\tfrac{\partial g}{\partial u}\tfrac{\partial h}{\partial v} + g(u,v)\tfrac{\partial^2 h}{\partial u\,\partial v} - \tfrac{\partial g}{\partial v}\tfrac{\partial h}{\partial u} - g(u,v)\tfrac{\partial^2 h}{\partial v\,\partial u}\right) dA \qquad \text{[using the Chain Rule]} \\ &= \pm\iint_S \left(\tfrac{\partial x}{\partial u}\tfrac{\partial y}{\partial v} - \tfrac{\partial x}{\partial v}\tfrac{\partial y}{\partial u}\right) dA \quad \text{[by the equality of mixed partials]} \quad = \pm\iint_S \tfrac{\partial(x,y)}{\partial(u,v)}\,du\,dv\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The sign is chosen to be positive if the orientation that we gave to ∂S corresponds to the usual positive orientation, and it is negative otherwise. In either case, since $A(R)$ is positive, the sign chosen must be the same as the sign of $\dfrac{\partial(x,y)}{\partial(u,v)}$.

Therefore $A(R) = \iint_R dx\,dy = \iint_S \left|\dfrac{\partial(x,y)}{\partial(u,v)}\right| du\,dv$.

16.5 Curl and Divergence

1. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ xy^2z^2 & x^2yz^2 & x^2y^2z \end{vmatrix}$

$$= \left[\frac{\partial}{\partial y}(x^2y^2z) - \frac{\partial}{\partial z}(x^2yz^2)\right]\mathbf{i} - \left[\frac{\partial}{\partial x}(x^2y^2z) - \frac{\partial}{\partial z}(xy^2z^2)\right]\mathbf{j} + \left[\frac{\partial}{\partial x}(x^2yz^2) - \frac{\partial}{\partial y}(xy^2z^2)\right]\mathbf{k}$$
$$= (2x^2yz - 2x^2yz)\,\mathbf{i} - (2xy^2z - 2xy^2z)\,\mathbf{j} + (2xyz^2 - 2xyz^2)\,\mathbf{k} = \mathbf{0}$$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}(xy^2z^2) + \dfrac{\partial}{\partial y}(x^2yz^2) + \dfrac{\partial}{\partial z}(x^2y^2z) = y^2z^2 + x^2z^2 + x^2y^2$

2. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ 0 & x^3yz^2 & y^4z^3 \end{vmatrix}$

$$= \left[\frac{\partial}{\partial y}(y^4z^3) - \frac{\partial}{\partial z}(x^3yz^2)\right]\mathbf{i} - \left[\frac{\partial}{\partial x}(y^4z^3) - \frac{\partial}{\partial z}(0)\right]\mathbf{j} + \left[\frac{\partial}{\partial x}(x^3yz^2) - \frac{\partial}{\partial y}(0)\right]\mathbf{k}$$
$$= (4y^3z^3 - 2x^3yz)\,\mathbf{i} - (0-0)\,\mathbf{j} + (3x^2yz^2 - 0)\,\mathbf{k} = (4y^3z^3 - 2x^3yz)\,\mathbf{i} + 3x^2yz^2\,\mathbf{k}$$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}(0) + \dfrac{\partial}{\partial y}(x^3yz^2) + \dfrac{\partial}{\partial z}(y^4z^3) = 0 + x^3z^2 + 3y^4z^2 = x^3z^2 + 3y^4z^2$

3. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ xye^z & 0 & yze^x \end{vmatrix} = (ze^x - 0)\,\mathbf{i} - (yze^x - xye^z)\,\mathbf{j} + (0 - xe^z)\,\mathbf{k}$

$$= ze^x\,\mathbf{i} + (xye^z - yze^x)\,\mathbf{j} - xe^z\,\mathbf{k}$$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}(xye^z) + \dfrac{\partial}{\partial y}(0) + \dfrac{\partial}{\partial z}(yze^x) = ye^z + 0 + ye^x = y(e^z + e^x)$

4. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \sin yz & \sin zx & \sin xy \end{vmatrix}$

$$= (x\cos xy - x\cos zx)\,\mathbf{i} - (y\cos xy - y\cos yz)\,\mathbf{j} + (z\cos zx - z\cos yz)\,\mathbf{k}$$
$$= x(\cos xy - \cos zx)\,\mathbf{i} + y(\cos yz - \cos xy)\,\mathbf{j} + z(\cos zx - \cos yz)\,\mathbf{k}$$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}(\sin yz) + \dfrac{\partial}{\partial y}(\sin zx) + \dfrac{\partial}{\partial z}(\sin xy) = 0 + 0 + 0 - 0$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

5. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \dfrac{\sqrt{x}}{1+z} & \dfrac{\sqrt{y}}{1+x} & \dfrac{\sqrt{z}}{1+y} \end{vmatrix}$

$= \left[\sqrt{z}\,(-1)(1+y)^{-2} - 0\right]\mathbf{i} - \left[0 - \sqrt{x}(-1)(1+z)^{-2}\right]\mathbf{j} + \left[\sqrt{y}\,(-1)(1+x)^{-2} - 0\right]\mathbf{k}$

$= -\dfrac{\sqrt{z}}{(1+y)^2}\,\mathbf{i} - \dfrac{\sqrt{x}}{(1+z)^2}\,\mathbf{j} - \dfrac{\sqrt{y}}{(1+x)^2}\,\mathbf{k}$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}\left(\dfrac{\sqrt{x}}{1+z}\right) + \dfrac{\partial}{\partial y}\left(\dfrac{\sqrt{y}}{1+x}\right) + \dfrac{\partial}{\partial z}\left(\dfrac{\sqrt{z}}{1+y}\right)$

$= \dfrac{1}{2\sqrt{x}\,(1+z)} + \dfrac{1}{2\sqrt{y}\,(1+x)} + \dfrac{1}{2\sqrt{z}\,(1+y)}$

6. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \ln(2y+3z) & \ln(x+3z) & \ln(x+2y) \end{vmatrix}$

$= \left(\dfrac{2}{x+2y} - \dfrac{3}{x+3z}\right)\mathbf{i} - \left(\dfrac{1}{x+2y} - \dfrac{3}{2y+3z}\right)\mathbf{j} + \left(\dfrac{1}{x+3z} - \dfrac{2}{2y+3z}\right)\mathbf{k}$

$= \left(\dfrac{2}{x+2y} - \dfrac{3}{x+3z}\right)\mathbf{i} + \left(\dfrac{3}{2y+3z} - \dfrac{1}{x+2y}\right)\mathbf{j} + \left(\dfrac{1}{x+3z} - \dfrac{2}{2y+3z}\right)\mathbf{k}$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}\left[\ln(2y+3z)\right] + \dfrac{\partial}{\partial y}\left[\ln(x+3z)\right] + \dfrac{\partial}{\partial z}\left[\ln(x+2y)\right] = 0+0+0 = 0$

7. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ e^x\sin y & e^y\sin z & e^z\sin x \end{vmatrix} = (0 - e^y\cos z)\,\mathbf{i} - (e^z\cos x - 0)\,\mathbf{j} + (0 - e^x\cos y)\,\mathbf{k}$

$= \langle -e^y\cos z, -e^z\cos x, -e^x\cos y\rangle$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}\left(e^x\sin y\right) + \dfrac{\partial}{\partial y}\left(e^y\sin z\right) + \dfrac{\partial}{\partial z}\left(e^z\sin x\right) = e^x\sin y + e^y\sin z + e^z\sin x$

8. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \arctan(xy) & \arctan(yz) & \arctan(zx) \end{vmatrix}$

$= \left(0 - \dfrac{y}{1+(yz)^2}\right)\mathbf{i} - \left(\dfrac{z}{1+(zx)^2} - 0\right)\mathbf{j} + \left(0 - \dfrac{x}{1+(xy)^2}\right)\mathbf{k}$

$= \left\langle -\dfrac{y}{1+y^2z^2}, -\dfrac{z}{1+x^2z^2}, -\dfrac{x}{1+x^2y^2}\right\rangle$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}\left[\arctan(xy)\right] + \dfrac{\partial}{\partial y}\left[\arctan(yz)\right] + \dfrac{\partial}{\partial z}\left[\arctan(zx)\right] = \dfrac{y}{1+x^2y^2} + \dfrac{z}{1+y^2z^2} + \dfrac{x}{1+x^2z^2}$

9. If the vector field is $\mathbf{F} = P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k}$, then we know $R = 0$. In addition, the x-component of each vector of $\mathbf{F}$ is 0, so $P = 0$, hence $\dfrac{\partial P}{\partial x} = \dfrac{\partial P}{\partial y} = \dfrac{\partial P}{\partial z} = \dfrac{\partial R}{\partial x} = \dfrac{\partial R}{\partial y} = \dfrac{\partial R}{\partial z} = 0$. Q decreases as y increases, so $\dfrac{\partial Q}{\partial y} < 0$, but Q doesn't change

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

in the x- or z-directions, so $\dfrac{\partial Q}{\partial x} = \dfrac{\partial Q}{\partial z} = 0$.

(a) $\operatorname{div} \mathbf{F} = \dfrac{\partial P}{\partial x} + \dfrac{\partial Q}{\partial y} + \dfrac{\partial R}{\partial z} = 0 + \dfrac{\partial Q}{\partial y} + 0 < 0$

(b) $\operatorname{curl} \mathbf{F} = \left(\dfrac{\partial R}{\partial y} - \dfrac{\partial Q}{\partial z}\right)\mathbf{i} + \left(\dfrac{\partial P}{\partial z} - \dfrac{\partial R}{\partial x}\right)\mathbf{j} + \left(\dfrac{\partial Q}{\partial x} - \dfrac{\partial P}{\partial y}\right)\mathbf{k} = (0-0)\,\mathbf{i} + (0-0)\,\mathbf{j} + (0-0)\,\mathbf{k} = \mathbf{0}$

10. If the vector field is $\mathbf{F} = P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k}$, then we know $R = 0$. In addition, P and Q don't vary in the z-direction, so $\dfrac{\partial R}{\partial x} = \dfrac{\partial R}{\partial y} = \dfrac{\partial R}{\partial z} = \dfrac{\partial P}{\partial z} = \dfrac{\partial Q}{\partial z} = 0$. As x increases, the x-component of each vector of $\mathbf{F}$ increases while the y-component remains constant, so $\dfrac{\partial P}{\partial x} > 0$ and $\dfrac{\partial Q}{\partial x} = 0$. Similarly, as y increases, the y-component of each vector increases while the x-component remains constant, so $\dfrac{\partial Q}{\partial y} > 0$ and $\dfrac{\partial P}{\partial y} = 0$.

(a) $\operatorname{div} \mathbf{F} = \dfrac{\partial P}{\partial x} + \dfrac{\partial Q}{\partial y} + \dfrac{\partial R}{\partial z} = \dfrac{\partial P}{\partial x} + \dfrac{\partial Q}{\partial y} + 0 > 0$

(b) $\operatorname{curl} \mathbf{F} = \left(\dfrac{\partial R}{\partial y} - \dfrac{\partial Q}{\partial z}\right)\mathbf{i} + \left(\dfrac{\partial P}{\partial z} - \dfrac{\partial R}{\partial x}\right)\mathbf{j} + \left(\dfrac{\partial Q}{\partial x} - \dfrac{\partial P}{\partial y}\right)\mathbf{k} = (0-0)\,\mathbf{i} + (0-0)\,\mathbf{j} + (0-0)\,\mathbf{k} = \mathbf{0}$

11. If the vector field is $\mathbf{F} = P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k}$, then we know $R = 0$. In addition, the y-component of each vector of $\mathbf{F}$ is 0, so $Q = 0$, hence $\dfrac{\partial Q}{\partial x} = \dfrac{\partial Q}{\partial y} = \dfrac{\partial Q}{\partial z} = \dfrac{\partial R}{\partial x} = \dfrac{\partial R}{\partial y} = \dfrac{\partial R}{\partial z} = 0$. P increases as y increases, so $\dfrac{\partial P}{\partial y} > 0$, but P doesn't change in the x- or z-directions, so $\dfrac{\partial P}{\partial x} = \dfrac{\partial P}{\partial z} = 0$.

(a) $\operatorname{div} \mathbf{F} = \dfrac{\partial P}{\partial x} + \dfrac{\partial Q}{\partial y} + \dfrac{\partial R}{\partial z} = 0 + 0 + 0 = 0$

(b) $\operatorname{curl} \mathbf{F} = \left(\dfrac{\partial R}{\partial y} - \dfrac{\partial Q}{\partial z}\right)\mathbf{i} + \left(\dfrac{\partial P}{\partial z} - \dfrac{\partial R}{\partial x}\right)\mathbf{j} + \left(\dfrac{\partial Q}{\partial x} - \dfrac{\partial P}{\partial y}\right)\mathbf{k} = (0-0)\,\mathbf{i} + (0-0)\,\mathbf{j} + \left(0 - \dfrac{\partial P}{\partial y}\right)\mathbf{k} = -\dfrac{\partial P}{\partial y}\mathbf{k}$

Since $\dfrac{\partial P}{\partial y} > 0$, $-\dfrac{\partial P}{\partial y}\mathbf{k}$ is a vector pointing in the negative z-direction.

12. (a) $\operatorname{curl} f = \nabla \times f$ is meaningless because f is a scalar field.

(b) $\operatorname{grad} f$ is a vector field.

(c) $\operatorname{div} \mathbf{F}$ is a scalar field.

(d) $\operatorname{curl}(\operatorname{grad} f)$ is a vector field.

(e) $\operatorname{grad} \mathbf{F}$ is meaningless because $\mathbf{F}$ is not a scalar field.

(f) $\operatorname{grad}(\operatorname{div} \mathbf{F})$ is a vector field.

(g) $\operatorname{div}(\operatorname{grad} f)$ is a scalar field.

(h) $\operatorname{grad}(\operatorname{div} f)$ is meaningless because f is a scalar field.

(i) $\operatorname{curl}(\operatorname{curl} \mathbf{F})$ is a vector field.

(j) $\operatorname{div}(\operatorname{div} \mathbf{F})$ is meaningless because $\operatorname{div} \mathbf{F}$ is a scalar field.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(k) $(\operatorname{grad} f) \times (\operatorname{div} \mathbf{F})$ is meaningless because $\operatorname{div} \mathbf{F}$ is a scalar field.

(l) $\operatorname{div}(\operatorname{curl}(\operatorname{grad} f))$ is a scalar field.

13. $\operatorname{curl} \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ y^2z^3 & 2xyz^3 & 3xy^2z^2 \end{vmatrix} = (6xyz^2 - 6xyz^2)\,\mathbf{i} - (3y^2z^2 - 3y^2z^2)\,\mathbf{j} + (2yz^3 - 2yz^3)\,\mathbf{k} = \mathbf{0}$

and $\mathbf{F}$ is defined on all of $\mathbb{R}^3$ with component functions which have continuous partial derivatives, so by Theorem 4, $\mathbf{F}$ is conservative. Thus, there exists a function f such that $\mathbf{F} = \nabla f$. Then $f_x(x, y, z) = y^2z^3$ implies $f(x, y, z) = xy^2z^3 + g(y, z)$ and $f_y(x, y, z) = 2xyz^3 + g_y(y, z)$. But $f_y(x, y, z) = 2xyz^3$, so $g(y, z) = h(z)$ and $f(x, y, z) = xy^2z^3 + h(z)$. Thus $f_z(x, y, z) = 3xy^2z^2 + h'(z)$ but $f_z(x, y, z) = 3xy^2z^2$ so $h(z) = K$, a constant. Hence a potential function for $\mathbf{F}$ is $f(x, y, z) = xy^2z^3 + K$.

14. $\operatorname{curl} \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ xyz^4 & x^2z^4 & 4x^2yz^3 \end{vmatrix} = (4x^2z^3 - 4x^2z^3)\,\mathbf{i} - (8xyz^3 - 4xyz^3)\,\mathbf{j} + (2xz^4 - xz^4)\,\mathbf{k} \neq \mathbf{0},$

so $\mathbf{F}$ is not conservative.

15. $\operatorname{curl} \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ z\cos y & xz\sin y & x\cos y \end{vmatrix}$

$= (-x\sin y - x\sin y)\,\mathbf{i} - (\cos y - \cos y)\,\mathbf{j} + [z\sin y - (-z\sin y)]\,\mathbf{k} = -2x\sin y\,\mathbf{i} + 2z\sin y\,\mathbf{k} \neq \mathbf{0},$

so $\mathbf{F}$ is not conservative.

16. $\operatorname{curl} \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ 1 & \sin z & y\cos z \end{vmatrix} = (\cos z - \cos z)\,\mathbf{i} - (0 - 0)\,\mathbf{j} + (0 - 0)\,\mathbf{k} = \mathbf{0}$, $\mathbf{F}$ is defined on all of $\mathbb{R}^3$, and the partial derivatives of the component functions are continuous, so $\mathbf{F}$ is conservative. Thus there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x, y, z) = 1$ implies $f(x, y, z) = x + g(y, z)$ and $f_y(x, y, z) = g_y(y, z)$. But $f_y(x, y, z) = \sin z$, so $g(y, z) = y\sin z + h(z)$ and $f(x, y, z) = x + y\sin z + h(z)$. Thus $f_z(x, y, z) = y\cos z + h'(z)$ but $f_z(x, y, z) = y\cos z$ so $h(z) = K$ and $f(x, y, z) = x + y\sin z + K$.

17. $\operatorname{curl} \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ e^{yz} & xze^{yz} & xye^{yz} \end{vmatrix}$

$= [xyze^{yz} + xe^{yz} - (xyze^{yz} + xe^{yz})]\,\mathbf{i} - (ye^{yz} - ye^{yz})\,\mathbf{j} + (ze^{yz} - ze^{yz})\,\mathbf{k} = \mathbf{0}$

$\mathbf{F}$ is defined on all of $\mathbb{R}^3$, and the partial derivatives of the component functions are continuous, so $\mathbf{F}$ is conservative. Thus there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x, y, z) = e^{yz}$ implies $f(x, y, z) = xe^{yz} + g(y, z) \;\Rightarrow$ $f_y(x, y, z) = xze^{yz} + g_y(y, z)$. But $f_y(x, y, z) = xze^{yz}$, so $g(y, z) = h(z)$ and $f(x, y, z) = xe^{yz} + h(z)$. Thus $f_z(x, y, z) = xye^{yz} + h'(z)$ but $f_z(x, y, z) = xye^{yz}$ so $h(z) = K$ and a potential function for $\mathbf{F}$ is $f(x, y, z) = xe^{yz} + K$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ e^x \sin yz & ze^x \cos yz & ye^x \cos yz \end{vmatrix}$

$= [-yze^x \sin yz + e^x \cos yz - (-yze^x \sin yz + e^x \cos yz)]\,\mathbf{i} - (ye^x \cos yz - ye^x \cos yz)\,\mathbf{j}$
$+ (ze^x \cos yz - ze^x \cos yz)\,\mathbf{k} = \mathbf{0}$

$\mathbf{F}$ is defined on all of $\mathbb{R}^3$, and the partial derivatives of the component functions are continuous, so $\mathbf{F}$ is conservative. Thus there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x, y, z) = e^x \sin yz$ implies $f(x, y, z) = e^x \sin yz + g(y, z) \Rightarrow f_y(x, y, z) = ze^x \cos yz + g_y(y, z)$. But $f_y(x, y, z) = ze^x \cos yz$, so $g(y, z) = h(z)$ and $f(x, y, z) = e^x \sin yz + h(z)$. Thus $f_z(x, y, z) = ye^x \cos yz + h'(z)$ but $f_z(x, y, z) = ye^x \cos yz$ so $h(z) = K$ and a potential function for $\mathbf{F}$ is $f(x, y, z) = e^x \sin yz + K$.

19. No. Assume there is such a $\mathbf{G}$. Then $\operatorname{div}(\operatorname{curl}\mathbf{G}) = \dfrac{\partial}{\partial x}(x \sin y) + \dfrac{\partial}{\partial y}(\cos y) + \dfrac{\partial}{\partial z}(z - xy) = \sin y - \sin y + 1 \neq 0$, which contradicts Theorem 11.

20. No. Assume there is such a $\mathbf{G}$. Then $\operatorname{div}(\operatorname{curl}\mathbf{G}) = \dfrac{\partial}{\partial x}(x) + \dfrac{\partial}{\partial y}(y) + \dfrac{\partial}{\partial z}(z) = 1 + 1 + 1 \neq 0$ which contradicts Theorem 11.

21. $\operatorname{curl}\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ f(x) & g(y) & h(z) \end{vmatrix} = (0 - 0)\,\mathbf{i} + (0 - 0)\,\mathbf{j} + (0 - 0)\,\mathbf{k} = \mathbf{0}$. Hence $\mathbf{F} = f(x)\,\mathbf{i} + g(y)\,\mathbf{j} + h(z)\,\mathbf{k}$ is irrotational.

22. $\operatorname{div}\mathbf{F} = \dfrac{\partial}{\partial x}(f(y, z)) + \dfrac{\partial}{\partial y}(g(x, z)) + \dfrac{\partial}{\partial z}(h(x, y)) = 0$ so $\mathbf{F}$ is incompressible.

For Exercises 23–29, let $\mathbf{F}(x, y, z) = P_1\,\mathbf{i} + Q_1\,\mathbf{j} + R_1\,\mathbf{k}$ and $\mathbf{G}(x, y, z) = P_2\,\mathbf{i} + Q_2\,\mathbf{j} + R_2\,\mathbf{k}$.

23. $\operatorname{div}(\mathbf{F} + \mathbf{G}) = \operatorname{div}\langle P_1 + P_2, Q_1 + Q_2, R_1 + R_2\rangle = \dfrac{\partial(P_1 + P_2)}{\partial x} + \dfrac{\partial(Q_1 + Q_2)}{\partial y} + \dfrac{\partial(R_1 + R_2)}{\partial z}$

$= \dfrac{\partial P_1}{\partial x} + \dfrac{\partial P_2}{\partial x} + \dfrac{\partial Q_1}{\partial y} + \dfrac{\partial Q_2}{\partial y} + \dfrac{\partial R_1}{\partial z} + \dfrac{\partial R_2}{\partial z} = \left(\dfrac{\partial P_1}{\partial x} + \dfrac{\partial Q_1}{\partial y} + \dfrac{\partial R_1}{\partial z}\right) + \left(\dfrac{\partial P_2}{\partial x} + \dfrac{\partial Q_2}{\partial y} + \dfrac{\partial R_2}{\partial z}\right)$

$= \operatorname{div}\langle P_1, Q_1, R_1\rangle + \operatorname{div}\langle P_2, Q_2, R_2\rangle = \operatorname{div}\mathbf{F} + \operatorname{div}\mathbf{G}$

24. $\operatorname{curl}\mathbf{F} + \operatorname{curl}\mathbf{G} = \left[\left(\dfrac{\partial R_1}{\partial y} - \dfrac{\partial Q_1}{\partial z}\right)\mathbf{i} + \left(\dfrac{\partial P_1}{\partial z} - \dfrac{\partial R_1}{\partial x}\right)\mathbf{j} + \left(\dfrac{\partial Q_1}{\partial x} - \dfrac{\partial P_1}{\partial y}\right)\mathbf{k}\right]$
$+ \left[\left(\dfrac{\partial R_2}{\partial y} - \dfrac{\partial Q_2}{\partial z}\right)\mathbf{i} + \left(\dfrac{\partial P_2}{\partial z} - \dfrac{\partial R_2}{\partial x}\right)\mathbf{j} + \left(\dfrac{\partial Q_2}{\partial x} - \dfrac{\partial P_2}{\partial y}\right)\mathbf{k}\right]$

$= \left[\dfrac{\partial(R_1 + R_2)}{\partial y} - \dfrac{\partial(Q_1 + Q_2)}{\partial z}\right]\mathbf{i} + \left[\dfrac{\partial(P_1 + P_2)}{\partial z} - \dfrac{\partial(R_1 + R_2)}{\partial x}\right]\mathbf{j}$
$+ \left[\dfrac{\partial(Q_1 + Q_2)}{\partial x} - \dfrac{\partial(P_1 + P_2)}{\partial y}\right]\mathbf{k} = \operatorname{curl}(\mathbf{F} + \mathbf{G})$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

25. $\operatorname{div}(f\mathbf{F}) = \operatorname{div}(f\langle P_1, Q_1, R_1\rangle) = \operatorname{div}\langle fP_1, fQ_1, fR_1\rangle = \dfrac{\partial(fP_1)}{\partial x} + \dfrac{\partial(fQ_1)}{\partial y} + \dfrac{\partial(fR_1)}{\partial z}$

$$= \left(f\frac{\partial P_1}{\partial x} + P_1\frac{\partial f}{\partial x}\right) + \left(f\frac{\partial Q_1}{\partial y} + Q_1\frac{\partial f}{\partial y}\right) + \left(f\frac{\partial R_1}{\partial z} + R_1\frac{\partial f}{\partial z}\right)$$

$$= f\left(\frac{\partial P_1}{\partial x} + \frac{\partial Q_1}{\partial y} + \frac{\partial R_1}{\partial z}\right) + \langle P_1, Q_1, R_1\rangle \cdot \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}\right\rangle = f\operatorname{div}\mathbf{F} + \mathbf{F}\cdot\nabla f$$

26. $\operatorname{curl}(f\mathbf{F}) = \left[\dfrac{\partial(fR_1)}{\partial y} - \dfrac{\partial(fQ_1)}{\partial z}\right]\mathbf{i} + \left[\dfrac{\partial(fP_1)}{\partial z} - \dfrac{\partial(fR_1)}{\partial x}\right]\mathbf{j} + \left[\dfrac{\partial(fQ_1)}{\partial x} - \dfrac{\partial(fP_1)}{\partial y}\right]\mathbf{k}$

$$= \left[f\frac{\partial R_1}{\partial y} + R_1\frac{\partial f}{\partial y} - f\frac{\partial Q_1}{\partial z} - Q_1\frac{\partial f}{\partial z}\right]\mathbf{i} + \left[f\frac{\partial P_1}{\partial z} + P_1\frac{\partial f}{\partial z} - f\frac{\partial R_1}{\partial x} - R_1\frac{\partial f}{\partial x}\right]\mathbf{j}$$
$$+ \left[f\frac{\partial Q_1}{\partial x} + Q_1\frac{\partial f}{\partial x} - f\frac{\partial P_1}{\partial y} - P_1\frac{\partial f}{\partial y}\right]\mathbf{k}$$

$$= f\left[\frac{\partial R_1}{\partial y} - \frac{\partial Q_1}{\partial z}\right]\mathbf{i} + f\left[\frac{\partial P_1}{\partial z} - \frac{\partial R_1}{\partial x}\right]\mathbf{j} + f\left[\frac{\partial Q_1}{\partial x} - \frac{\partial P_1}{\partial y}\right]\mathbf{k}$$
$$+ \left[R_1\frac{\partial f}{\partial y} - Q_1\frac{\partial f}{\partial z}\right]\mathbf{i} + \left[P_1\frac{\partial f}{\partial z} - R_1\frac{\partial f}{\partial x}\right]\mathbf{j} + \left[Q_1\frac{\partial f}{\partial x} - P_1\frac{\partial f}{\partial y}\right]\mathbf{k}$$

$$= f\operatorname{curl}\mathbf{F} + (\nabla f)\times\mathbf{F}$$

27. $\operatorname{div}(\mathbf{F}\times\mathbf{G}) = \nabla\cdot(\mathbf{F}\times\mathbf{G}) = \begin{vmatrix} \partial/\partial x & \partial/\partial y & \partial/\partial z \\ P_1 & Q_1 & R_1 \\ P_2 & Q_2 & R_2 \end{vmatrix} = \dfrac{\partial}{\partial x}\begin{vmatrix} Q_1 & R_1 \\ Q_2 & R_2 \end{vmatrix} - \dfrac{\partial}{\partial y}\begin{vmatrix} P_1 & R_1 \\ P_2 & R_2 \end{vmatrix} + \dfrac{\partial}{\partial z}\begin{vmatrix} P_1 & Q_1 \\ P_2 & Q_2 \end{vmatrix}$

$$= \left[Q_1\frac{\partial R_2}{\partial x} + R_2\frac{\partial Q_1}{\partial x} - Q_2\frac{\partial R_1}{\partial x} - R_1\frac{\partial Q_2}{\partial x}\right] - \left[P_1\frac{\partial R_2}{\partial y} + R_2\frac{\partial P_1}{\partial y} - P_2\frac{\partial R_1}{\partial y} - R_1\frac{\partial P_2}{\partial y}\right]$$
$$+ \left[P_1\frac{\partial Q_2}{\partial z} + Q_2\frac{\partial P_1}{\partial z} - P_2\frac{\partial Q_1}{\partial z} - Q_1\frac{\partial P_2}{\partial z}\right]$$

$$= \left[P_2\left(\frac{\partial R_1}{\partial y} - \frac{\partial Q_1}{\partial z}\right) + Q_2\left(\frac{\partial P_1}{\partial z} - \frac{\partial R_1}{\partial x}\right) + R_2\left(\frac{\partial Q_1}{\partial x} - \frac{\partial P_1}{\partial y}\right)\right]$$
$$- \left[P_1\left(\frac{\partial R_2}{\partial y} - \frac{\partial Q_2}{\partial z}\right) + Q_1\left(\frac{\partial P_2}{\partial z} - \frac{\partial R_2}{\partial x}\right) + R_1\left(\frac{\partial Q_2}{\partial x} - \frac{\partial P_2}{\partial y}\right)\right]$$

$$= \mathbf{G}\cdot\operatorname{curl}\mathbf{F} - \mathbf{F}\cdot\operatorname{curl}\mathbf{G}$$

28. $\operatorname{div}(\nabla f\times\nabla g) = \nabla g\cdot\operatorname{curl}(\nabla f) - \nabla f\cdot\operatorname{curl}(\nabla g)$ [by Exercise 27] $= 0$ [by Theorem 3]

29. $\operatorname{curl}(\operatorname{curl}\mathbf{F}) = \nabla\times(\nabla\times\mathbf{F}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \partial R_1/\partial y - \partial Q_1/\partial z & \partial P_1/\partial z - \partial R_1/\partial x & \partial Q_1/\partial x - \partial P_1/\partial y \end{vmatrix}$

$$= \left(\frac{\partial^2 Q_1}{\partial y\partial x} - \frac{\partial^2 P_1}{\partial y^2} - \frac{\partial^2 P_1}{\partial z^2} + \frac{\partial^2 R_1}{\partial z\partial x}\right)\mathbf{i} + \left(\frac{\partial^2 R_1}{\partial z\partial y} - \frac{\partial^2 Q_1}{\partial z^2} - \frac{\partial^2 Q_1}{\partial x^2} + \frac{\partial^2 P_1}{\partial x\partial y}\right)\mathbf{j}$$
$$+ \left(\frac{\partial^2 P_1}{\partial x\partial z} - \frac{\partial^2 R_1}{\partial x^2} - \frac{\partial^2 R_1}{\partial y^2} + \frac{\partial^2 Q_1}{\partial y\partial z}\right)\mathbf{k}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now let's consider $\operatorname{grad}(\operatorname{div}\mathbf{F}) - \nabla^2\mathbf{F}$ and compare with the above.

(Note that $\nabla^2\mathbf{F}$ is defined on page 1147 [ET 1107].)

$$\operatorname{grad}(\operatorname{div}\mathbf{F}) - \nabla^2\mathbf{F} = \left[\left(\frac{\partial^2 P_1}{\partial x^2} + \frac{\partial^2 Q_1}{\partial x\partial y} + \frac{\partial^2 R_1}{\partial x\partial z}\right)\mathbf{i} + \left(\frac{\partial^2 P_1}{\partial y\partial x} + \frac{\partial^2 Q_1}{\partial y^2} + \frac{\partial^2 R_1}{\partial y\partial z}\right)\mathbf{j} + \left(\frac{\partial^2 P_1}{\partial z\partial x} + \frac{\partial^2 Q_1}{\partial z\partial y} + \frac{\partial^2 R_1}{\partial z^2}\right)\mathbf{k}\right]$$

$$\left[\left(\frac{\partial^2 P_1}{\partial x^2} + \frac{\partial^2 P_1}{\partial y^2} + \frac{\partial^2 P_1}{\partial z^2}\right)\mathbf{i} + \left(\frac{\partial^2 Q_1}{\partial x^2} + \frac{\partial^2 Q_1}{\partial y^2} + \frac{\partial^2 Q_1}{\partial z^2}\right)\mathbf{j} + \left(\frac{\partial^2 R_1}{\partial x^2} + \frac{\partial^2 R_1}{\partial y^2} + \frac{\partial^2 R_1}{\partial z^2}\right)\mathbf{k}\right]$$

$$= \left(\frac{\partial^2 Q_1}{\partial x\partial y} + \frac{\partial^2 R_1}{\partial x\partial z} - \frac{\partial^2 P_1}{\partial y^2} - \frac{\partial^2 P_1}{\partial z^2}\right)\mathbf{i} + \left(\frac{\partial^2 P_1}{\partial y\partial x} + \frac{\partial^2 R_1}{\partial y\partial z} - \frac{\partial^2 Q_1}{\partial x^2} - \frac{\partial^2 Q_1}{\partial z^2}\right)\mathbf{j}$$

$$+ \left(\frac{\partial^2 P_1}{\partial z\partial x} + \frac{\partial^2 Q_1}{\partial z\partial y} - \frac{\partial^2 R_1}{\partial x^2} - \frac{\partial^2 R_2}{\partial y^2}\right)\mathbf{k}$$

Then applying Clairaut's Theorem to reverse the order of differentiation in the second partial derivatives as needed and comparing, we have $\operatorname{curl}\operatorname{curl}\mathbf{F} = \operatorname{grad}\operatorname{div}\mathbf{F} - \nabla^2\mathbf{F}$ as desired.

30. (a) $\nabla\cdot\mathbf{r} = \left(\dfrac{\partial}{\partial x}\mathbf{i} + \dfrac{\partial}{\partial y}\mathbf{j} + \dfrac{\partial}{\partial z}\mathbf{k}\right)\cdot(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}) = 1 + 1 + 1 = 3$

(b) $\nabla\cdot(r\mathbf{r}) = \nabla\cdot\sqrt{x^2+y^2+z^2}\,(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k})$

$$= \left(\frac{x^2}{\sqrt{x^2+y^2+z^2}} + \sqrt{x^2+y^2+z^2}\right) + \left(\frac{y^2}{\sqrt{x^2+y^2+z^2}} + \sqrt{x^2+y^2+z^2}\right)$$

$$+ \left(\frac{z^2}{\sqrt{x^2+y^2+z^2}} + \sqrt{x^2+y^2+z^2}\right)$$

$$= \frac{1}{\sqrt{x^2+y^2+z^2}}\,(4x^2+4y^2+4z^2) = 4\sqrt{x^2+y^2+z^2} = 4r$$

Another method:

By Exercise 25, $\nabla\cdot(r\mathbf{r}) = \operatorname{div}(r\mathbf{r}) = r\operatorname{div}\mathbf{r} + \mathbf{r}\cdot\nabla r = 3r + \mathbf{r}\cdot\dfrac{\mathbf{r}}{r}$ [see Exercise 31(a) below] $= 4r$.

(c) $\nabla^2 r^3 = \nabla^2\left(x^2+y^2+z^2\right)^{3/2}$

$$= \tfrac{\partial}{\partial x}\left[\tfrac{3}{2}(x^2+y^2+z^2)^{1/2}(2x)\right] + \tfrac{\partial}{\partial y}\left[\tfrac{3}{2}(x^2+y^2+z^2)^{1/2}(2y)\right] + \tfrac{\partial}{\partial z}\left[\tfrac{3}{2}(x^2+y^2+z^2)^{1/2}(2z)\right]$$

$$= 3\left[\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2x)(x) + (x^2+y^2+z^2)^{1/2}\right]$$

$$+ 3\left[\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2y)(y) + (x^2+y^2+z^2)^{1/2}\right]$$

$$+ 3\left[\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2z)(z) + (x^2+y^2+z^2)^{1/2}\right]$$

$$= 3(x^2+y^2+z^2)^{-1/2}(4x^2+4y^2+4z^2) = 12(x^2+y^2+z^2)^{1/2} = 12r$$

Another method: $\frac{\partial}{\partial x}\left(x^2+y^2+z^2\right)^{3/2} = 3x\sqrt{x^2+y^2+z^2} \;\Rightarrow\; \nabla r^3 = 3r(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}) = 3r\,\mathbf{r}$,

so $\nabla^2 r^3 = \nabla\cdot\nabla r^3 = \nabla\cdot(3r\,\mathbf{r}) = 3(4r) = 12r$ by part (b).

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

31. (a) $\nabla r = \nabla\sqrt{x^2+y^2+z^2} = \dfrac{x}{\sqrt{x^2+y^2+z^2}}\,\mathbf{i} + \dfrac{y}{\sqrt{x^2+y^2+z^2}}\,\mathbf{j} + \dfrac{z}{\sqrt{x^2+y^2+z^2}}\,\mathbf{k} = \dfrac{x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}}{\sqrt{x^2+y^2+z^2}} = \dfrac{\mathbf{r}}{r}$

(b) $\nabla \times \mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \dfrac{\partial}{\partial x} & \dfrac{\partial}{\partial y} & \dfrac{\partial}{\partial z} \\ x & y & z \end{vmatrix} = \left[\dfrac{\partial}{\partial y}(z) - \dfrac{\partial}{\partial z}(y)\right]\mathbf{i} + \left[\dfrac{\partial}{\partial z}(x) - \dfrac{\partial}{\partial x}(z)\right]\mathbf{j} + \left[\dfrac{\partial}{\partial x}(y) - \dfrac{\partial}{\partial y}(x)\right]\mathbf{k} = \mathbf{0}$

(c) $\nabla\left(\dfrac{1}{r}\right) = \nabla\left(\dfrac{1}{\sqrt{x^2+y^2+z^2}}\right)$

$$= \frac{\frac{1}{2\sqrt{x^2+y^2+z^2}}(2x)}{x^2+y^2+z^2}\,\mathbf{i} - \frac{\frac{1}{2\sqrt{x^2+y^2+z^2}}(2y)}{x^2+y^2+z^2}\,\mathbf{j} - \frac{\frac{1}{2\sqrt{x^2+y^2+z^2}}(2z)}{x^2+y^2+z^2}\,\mathbf{k}$$

$$= -\frac{x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}}{(x^2+y^2+z^2)^{3/2}} = -\frac{\mathbf{r}}{r^3}$$

(d) $\nabla \ln r = \nabla \ln(x^2+y^2+z^2)^{1/2} = \frac{1}{2}\nabla \ln(x^2+y^2+z^2)$

$$= \frac{x}{x^2+y^2+z^2}\,\mathbf{i} + \frac{y}{x^2+y^2+z^2}\,\mathbf{j} + \frac{z}{x^2+y^2+z^2}\,\mathbf{k} = \frac{x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}}{x^2+y^2+z^2} = \frac{\mathbf{r}}{r^2}$$

32. $\mathbf{r} = x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k} \;\Rightarrow\; r = |\mathbf{r}| = \sqrt{x^2+y^2+z^2}$, so

$$\mathbf{F} = \frac{\mathbf{r}}{r^p} = \frac{x}{(x^2+y^2+z^2)^{p/2}}\,\mathbf{i} + \frac{y}{(x^2+y^2+z^2)^{p/2}}\,\mathbf{j} + \frac{z}{(x^2+y^2+z^2)^{p/2}}\,\mathbf{k}$$

Then $\dfrac{\partial}{\partial x}\dfrac{x}{(x^2+y^2+z^2)^{p/2}} = \dfrac{(x^2+y^2+z^2)-px^2}{(x^2+y^2+z^2)^{1+p/2}} = \dfrac{r^2-px^2}{r^{p+2}}$. Similarly,

$\dfrac{\partial}{\partial y}\dfrac{y}{(x^2+y^2+z^2)^{p/2}} = \dfrac{r^2-py^2}{r^{p+2}}$ and $\dfrac{\partial}{\partial z}\dfrac{z}{(x^2+y^2+z^2)^{p/2}} = \dfrac{r^2-pz^2}{r^{p+2}}$. Thus

$$\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \frac{r^2-px^2}{r^{p+2}} + \frac{r^2-py^2}{r^{p+2}} + \frac{r^2-pz^2}{r^{p+2}} = \frac{3r^2-px^2-py^2-pz^2}{r^{p+2}}$$

$$= \frac{3r^2-p(x^2+y^2+z^2)}{r^{p+2}} = \frac{3r^2-pr^2}{r^{p+2}} = \frac{3-p}{r^p}$$

Consequently, if $p = 3$ we have $\operatorname{div}\mathbf{F} = 0$.

33. By (13), $\oint_C f(\nabla g)\cdot\mathbf{n}\,ds = \iint_D \operatorname{div}(f\nabla g)\,dA = \iint_D [f\operatorname{div}(\nabla g) + \nabla g\cdot\nabla f]\,dA$ by Exercise 25. But $\operatorname{div}(\nabla g) = \nabla^2 g$. Hence $\iint_D f\nabla^2 g\,dA = \oint_C f(\nabla g)\cdot\mathbf{n}\,ds - \iint_D \nabla g\cdot\nabla f\,dA$.

34. By Exercise 33, $\iint_D f\nabla^2 g\,dA = \oint_C f(\nabla g)\cdot\mathbf{n}\,ds - \iint_D \nabla g\cdot\nabla f\,dA$ and

$\iint_D g\nabla^2 f\,dA = \oint_C g(\nabla f)\cdot\mathbf{n}\,ds - \iint_D \nabla f\cdot\nabla g\,dA$. Hence

$\iint_D (f\nabla^2 g - g\nabla^2 f)\,dA = \oint_C [f(\nabla g)\cdot\mathbf{n} - g(\nabla f)\cdot\mathbf{n}]\,ds + \iint_D (\nabla f\cdot\nabla g - \nabla g\cdot\nabla f)\,dA = \oint_C [f\nabla g - g\nabla f]\cdot\mathbf{n}\,ds$.

35. Let $f(x,y) = 1$. Then $\nabla f = \mathbf{0}$ and Green's first identity (see Exercise 33) says

$\iint_D \nabla^2 g\,dA = \oint_C (\nabla g)\cdot\mathbf{n}\,ds - \iint_D \mathbf{0}\cdot\nabla g\,dA \;\Rightarrow\; \iint_D \nabla^2 g\,dA = \oint_C \nabla g\cdot\mathbf{n}\,ds$. But g is harmonic on D, so

$\nabla^2 g = 0 \;\Rightarrow\; \oint_C \nabla g\cdot\mathbf{n}\,ds = 0$ and $\oint_C D_{\mathbf{n}} g\,ds = \oint_C (\nabla g\cdot\mathbf{n})\,ds = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. Let $g = f$. Then Green's first identity (see Exercise 33) says $\iint_D f\,\nabla^2 f\,dA = \oint_C (f)(\nabla f)\cdot \mathbf{n}\,ds - \iint_D \nabla f\cdot\nabla f\,dA$. But f is harmonic, so $\nabla^2 f = 0$, and $\nabla f\cdot\nabla f = |\nabla f|^2$, so we have $0 = \oint_C (f)(\nabla f)\cdot\mathbf{n}\,ds - \iint_D |\nabla f|^2\,dA \;\Rightarrow$ $\iint_D |\nabla f|^2\,dA = \oint_C (f)(\nabla f)\cdot\mathbf{n}\,ds = 0$ since $f(x,y) = 0$ on C.

37. (a) We know that $\omega = v/d$, and from the diagram $\sin\theta = d/r \;\Rightarrow\; v = d\omega = (\sin\theta) r\omega = |\mathbf{w}\times\mathbf{r}|$. But $\mathbf{v}$ is perpendicular to both $\mathbf{w}$ and $\mathbf{r}$, so that $\mathbf{v} = \mathbf{w}\times\mathbf{r}$.

(b) From (a), $\mathbf{v} = \mathbf{w}\times\mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & \omega \\ x & y & z \end{vmatrix} = (0\cdot z - \omega y)\,\mathbf{i} + (\omega x - 0\cdot z)\,\mathbf{j} + (0\cdot y - x\cdot 0)\,\mathbf{k} = -\omega y\,\mathbf{i} + \omega x\,\mathbf{j}$

(c) $\operatorname{curl}\mathbf{v} = \nabla\times\mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ -\omega y & \omega x & 0 \end{vmatrix}$

$$= \left[\frac{\partial}{\partial y}(0) - \frac{\partial}{\partial z}(\omega x)\right]\mathbf{i} + \left[\frac{\partial}{\partial z}(-\omega y) - \frac{\partial}{\partial x}(0)\right]\mathbf{j} + \left[\frac{\partial}{\partial x}(\omega x) - \frac{\partial}{\partial y}(-\omega y)\right]\mathbf{k}$$
$$= [\omega - (-\omega)]\,\mathbf{k} = 2\omega\,\mathbf{k} = 2\mathbf{w}$$

38. Let $\mathbf{H} = \langle h_1, h_2, h_3\rangle$ and $\mathbf{E} = \langle E_1, E_2, E_3\rangle$.

(a) $\nabla\times(\nabla\times\mathbf{E}) = \nabla\times(\operatorname{curl}\mathbf{E}) = \nabla\times\left(-\frac{1}{c}\frac{\partial\mathbf{H}}{\partial t}\right) = -\frac{1}{c}\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \partial h_1/\partial t & \partial h_2/\partial t & \partial h_3/\partial t \end{vmatrix}$

$$= -\frac{1}{c}\left[\left(\frac{\partial^2 h_3}{\partial y\,\partial t} - \frac{\partial^2 h_2}{\partial z\,\partial t}\right)\mathbf{i} + \left(\frac{\partial^2 h_1}{\partial z\,\partial t} - \frac{\partial^2 h_3}{\partial x\,\partial t}\right)\mathbf{j} + \left(\frac{\partial^2 h_2}{\partial x\,\partial t} - \frac{\partial^2 h_1}{\partial y\,\partial t}\right)\mathbf{k}\right]$$

$$= -\frac{1}{c}\frac{\partial}{\partial t}\left[\left(\frac{\partial h_3}{\partial y} - \frac{\partial h_2}{\partial z}\right)\mathbf{i} + \left(\frac{\partial h_1}{\partial z} - \frac{\partial h_3}{\partial x}\right)\mathbf{j} + \left(\frac{\partial h_2}{\partial x} - \frac{\partial h_1}{\partial y}\right)\mathbf{k}\right]$$

[assuming that the partial derivatives are continuous so that the order of differentiation does not matter]

$$= -\frac{1}{c}\frac{\partial}{\partial t}\operatorname{curl}\mathbf{H} = -\frac{1}{c}\frac{\partial}{\partial t}\left(\frac{1}{c}\frac{\partial\mathbf{E}}{\partial t}\right) = -\frac{1}{c^2}\frac{\partial^2\mathbf{E}}{\partial t^2}$$

(b) $\nabla\times(\nabla\times\mathbf{H}) = \nabla\times(\operatorname{curl}\mathbf{H}) = \nabla\times\left(\frac{1}{c}\frac{\partial\mathbf{E}}{\partial t}\right) = \frac{1}{c}\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \partial E_1/\partial t & \partial E_2/\partial t & \partial E_3/\partial t \end{vmatrix}$

$$= \frac{1}{c}\left[\left(\frac{\partial^2 E_3}{\partial y\,\partial t} - \frac{\partial^2 E_2}{\partial z\,\partial t}\right)\mathbf{i} + \left(\frac{\partial^2 E_1}{\partial z\,\partial t} - \frac{\partial^2 E_3}{\partial x\,\partial t}\right)\mathbf{j} + \left(\frac{\partial^2 E_2}{\partial x\,\partial t} - \frac{\partial^2 E_1}{\partial y\,\partial t}\right)\mathbf{k}\right]$$

$$= \frac{1}{c}\frac{\partial}{\partial t}\left[\left(\frac{\partial E_3}{\partial y} - \frac{\partial E_2}{\partial z}\right)\mathbf{i} + \left(\frac{\partial E_1}{\partial z} - \frac{\partial E_3}{\partial x}\right)\mathbf{j} + \left(\frac{\partial E_2}{\partial x} - \frac{\partial E_1}{\partial y}\right)\mathbf{k}\right]$$

[assuming that the partial derivatives are continuous so that the order of differentiation does not matter]

$$= \frac{1}{c}\frac{\partial}{\partial t}\operatorname{curl}\mathbf{E} = \frac{1}{c}\frac{\partial}{\partial t}\left(-\frac{1}{c}\frac{\partial\mathbf{H}}{\partial t}\right) = -\frac{1}{c^2}\frac{\partial^2\mathbf{H}}{\partial t^2}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c) Using Exercise 29, we have that $\operatorname{curl}\operatorname{curl}\mathbf{E} = \operatorname{grad}\operatorname{div}\mathbf{E} - \nabla^2\mathbf{E} \Rightarrow$

$$\nabla^2\mathbf{E} = \operatorname{grad}\operatorname{div}\mathbf{E} - \operatorname{curl}\operatorname{curl}\mathbf{E} = \operatorname{grad} 0 + \frac{1}{c^2}\frac{\partial^2\mathbf{E}}{\partial t^2} \quad [\text{from part (a)}] \quad = \frac{1}{c^2}\frac{\partial^2\mathbf{E}}{\partial t^2}.$$

(d) As in part (c), $\nabla^2\mathbf{H} = \operatorname{grad}\operatorname{div}\mathbf{H} - \operatorname{curl}\operatorname{curl}\mathbf{H} = \operatorname{grad} 0 + \dfrac{1}{c^2}\dfrac{\partial^2\mathbf{H}}{\partial t^2}$ [using part (b)] $= \dfrac{1}{c^2}\dfrac{\partial^2\mathbf{H}}{\partial t^2}$.

39. For any continuous function f on $\mathbb{R}^3$, define a vector field $\mathbf{G}(x, y, z) = \langle g(x, y, z), 0, 0\rangle$ where $g(x, y, z) = \int_0^x f(t, y, z)\, dt$. Then $\operatorname{div}\mathbf{G} = \dfrac{\partial}{\partial x}(g(x, y, z)) + \dfrac{\partial}{\partial y}(0) + \dfrac{\partial}{\partial z}(0) = \dfrac{\partial}{\partial x}\int_0^x f(t, y, z)\, dt = f(x, y, z)$ by the Fundamental Theorem of Calculus. Thus every continuous function f on $\mathbb{R}^3$ is the divergence of some vector field.

16.6 Parametric Surfaces and Their Areas

1. $P(4, -5, 1)$ lies on the parametric surface $\mathbf{r}(u, v) = \langle u + v, u - 2v, 3 + u - v\rangle$ if and only if there are values for u and v where $u + v = 4$, $u - 2v = -5$, and $3 + u - v = 1$. From the first equation we have $u = 4 - v$ and substituting into the second equation gives $4 - v - 2v = -5 \Leftrightarrow v = 3$. Then $u = 1$, and these values satisfy the third equation, so P does lie on the surface.

$Q(0, 4, 6)$ lies on $\mathbf{r}(u, v)$ if and only if $u + v = 0$, $u - 2v = 4$, and $3 + u - v = 6$, but solving the first two equations simultaneoulsy gives $u = \frac{4}{3}$, $v = -\frac{4}{3}$ and these values do not satisfy the third equation, so Q does not lie on the surface.

2. $P(1, 2, 1)$ lies on the parametric surface $\mathbf{r}(u, v) = \langle 1 + u - v, u + v^2, u^2 - v^2\rangle$ if and only if there are values for u and v where $1 + u - v = 1$, $u + v^2 = 2$, and $u^2 - v^2 = 1$. From the first equation we have $u = v$ and substituting into the third equation gives $0 = 1$, an impossibility, so P does not lie on the surface.

$Q(2, 3, 3)$ lies on $\mathbf{r}(u, v)$ if and only if $1 + u - v = 2$, $u + v^2 = 3$, and $u^2 - v^2 = 3$. From the first equation we have $u = v + 1$ and substituting into the second equation gives $v + 1 + v^2 = 3 \Leftrightarrow v^2 + v - 2 = 0 \Leftrightarrow (v + 2)(v - 1) = 0$, so $v = -2 \Rightarrow u = -1$ or $v = 1 \Rightarrow u = 2$. The third equation is satisfied by $u = 2$, $v = 1$ so Q does lie on the surface.

3. $\mathbf{r}(u, v) = (u + v)\,\mathbf{i} + (3 - v)\,\mathbf{j} + (1 + 4u + 5v)\,\mathbf{k} = \langle 0, 3, 1\rangle + u\langle 1, 0, 4\rangle + v\langle 1, -1, 5\rangle$. From Example 3, we recognize this as a vector equation of a plane through the point $(0, 3, 1)$ and containing vectors $\mathbf{a} = \langle 1, 0, 4\rangle$ and $\mathbf{b} = \langle 1, -1, 5\rangle$. If we wish to find a more conventional equation for the plane, a normal vector to the plane is $\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 4 \\ 1 & -1 & 5 \end{vmatrix} = 4\,\mathbf{i} - \mathbf{j} - \mathbf{k}$

and an equation of the plane is $4(x - 0) - (y - 3) - (z - 1) = 0$ or $4x - y - z = -4$.

4. $\mathbf{r}(u, v) = u^2\,\mathbf{i} + u\cos v\,\mathbf{j} + u\sin v\,\mathbf{k}$, so the corresponding parametric equations for the surface are $x = u^2$, $y = u\cos v$, $z = u\sin v$. For any point (x, y, z) on the surface, we have $y^2 + z^2 = u^2\cos^2 v + u^2\sin^2 v = u^2 = x$. Since no restrictions are placed on the parameters, the surface is $x = y^2 + z^2$, which we recognize as a circular paraboloid whose axis is the x-axis.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\mathbf{r}(s,t) = \langle s\cos t, s\sin t, s\rangle$, so the corresponding parametric equations for the surface are $x = s\cos t,\ \ y = s\sin t,\ \ z = s$. For any point (x, y, z) on the surface, we have $x^2 + y^2 = s^2\cos^2 t + s^2\sin^2 t = s^2 = z^2$. Since no restrictions are placed on the parameters, the surface is $z^2 = x^2 + y^2$, which we recognize as a circular cone with axis the z-axis.

6. $\mathbf{r}(s,t) = \langle 3\cos t, s, \sin t\rangle$, so the corresponding parametric equations for the surface are $x = 3\cos t,\ \ y = s,\ \ z = \sin t$. For any point (x, y, z) on the surface, we have $(x/3)^2 + z^2 = \cos^2 t + \sin^2 t = 1$, so vertical cross-sections parallel to the xz-plane are all identical ellipses. Since $y = s$ and $-1 \le s \le 1$, the surface is the portion of the elliptic cylinder $\frac{1}{9}x^2 + z^2 = 1$ corresponding to $-1 \le y \le 1$.

7. $\mathbf{r}(u,v) = \langle u^2, v^2, u+v\rangle,\ \ -1 \le u \le 1,\ \ -1 \le v \le 1$.

The surface has parametric equations $x = u^2$, $y = v^2$, $z = u + v$, $-1 \le u \le 1$, $-1 \le v \le 1$.

In Maple, the surface can be graphed by entering `plot3d([u^2,v^2,u+v],u=-1..1,v=-1..1);`.
In Mathematica we use the `ParametricPlot3D` command.
If we keep u constant at u_0, $x = u_0^2$, a constant, so the corresponding grid curves must be the curves parallel to the yz-plane. If v is constant, we have $y = v_0^2$, a constant, so these grid curves are the curves parallel to the xz-plane.



8. $\mathbf{r}(u,v) = \langle u, v^3, -v\rangle,\ \ -2 \le u \le 2,\ -2 \le v \le 2$.

The surface has parametric equations $x = u$, $y = v^3$, $z = -v$, $-2 \le u \le 2$, $-2 \le v \le 2$. If $u = u_0$ is constant, $x = u_0 =$ constant, so the corresponding grid curves are the curves parallel to the yz-plane. If $v = v_0$ is constant, $y = v_0^3 =$ constant, so the corresponding grid curves are the curves parallel to the xz-plane.



9. $\mathbf{r}(u,v) = \langle u^3, u\sin v, u\cos v\rangle,\ -1 \le u \le 1,\ 0 \le v \le 2\pi$

The surface has parametric equations $x = u^3,\ \ y = u\sin v$, $z = u\cos v,\ -1 \le u \le 1,\ 0 \le v \le 2\pi$. Note that if $u = u_0$ is constant then $x = u_0^3$ is constant and $y = u_0\sin v$, $z = u_0\cos v$ describe a circle in y, z of radius $|u_0|$, so the corresponding grid curves are circles parallel to the yz-plane. If $v = v_0$, a constant, the parametric equations become $x = u^3$, $y = u\sin v_0$, $z = u\cos v_0$. Then $y = (\tan v_0)z$, so these are the grid curves we see that lie in planes $y = kz$ that pass through the x-axis.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

10. $\mathbf{r}(u,v) = \langle u, \sin(u+v), \sin v \rangle,\ -\pi \le u \le \pi,\ -\pi \le v \le \pi.$

The surface has parametric equations $x = u$, $y = \sin(u+v)$, $z = \sin v$, $-\pi \le u \le \pi$, $-\pi \le v \le \pi$. If $u = u_0$ is constant, $x = u_0 =$ constant, so the corresponding grid curves are the curves parallel to the yz-plane. If $v = v_0$ is constant, $z = \sin v_0 =$ constant, so the corresponding grid curves are the curves parallel to the xy-plane.



11. $x = \sin v,\ y = \cos u \sin 4v,\ z = \sin 2u \sin 4v,\ 0 \le u \le 2\pi,\ -\frac{\pi}{2} \le v \le \frac{\pi}{2}.$

Note that if $v = v_0$ is constant, then $x = \sin v_0$ is constant, so the corresponding grid curves must be parallel to the yz-plane. These are the vertically oriented grid curves we see, each shaped like a "figure-eight." When $u = u_0$ is held constant, the parametric equations become $x = \sin v$, $y = \cos u_0 \sin 4v$, $z = \sin 2u_0 \sin 4v$. Since z is a constant multiple of y, the corresponding grid curves are the curves contained in planes $z = ky$ that pass through the x-axis.



12. $x = \cos u,\ y = \sin u \sin v,\ z = \cos v,\ 0 \le u \le 2\pi,\ 0 \le v \le 2\pi.$

If $u = u_0$ is constant, then $x = \cos u_0 =$ constant, so the corresponding grid curves are the curves parallel to the yz-plane. If $v = v_0$ is constant, then $z = \cos v_0 =$ constant, so the corresponding grid curves are the curves parallel to the xy-plane.



13. $\mathbf{r}(u,v) = u\cos v\,\mathbf{i} + u\sin v\,\mathbf{j} + v\,\mathbf{k}$. The parametric equations for the surface are $x = u\cos v$, $y = u\sin v$, $z = v$. We look at the grid curves first; if we fix v, then x and y parametrize a straight line in the plane $z = v$ which intersects the z-axis. If u is held constant, the projection onto the xy-plane is circular; with $z = v$, each grid curve is a helix. The surface is a spiraling ramp, graph IV.

14. $\mathbf{r}(u,v) = uv^2\,\mathbf{i} + u^2v\,\mathbf{j} + (u^2 - v^2)\,\mathbf{k}$. The parametric equations for the surface are $x = uv^2$, $y = u^2v$, $z = u^2 - v^2$. If $u = u_0$ is held constant, then $x = u_0v^2$, $y = u_0^2v$ so $x = u_0(y/u_0^2)^2 = (1/u_0^3)y^2$, and $z = u_0^2 - v^2 = u_0^2 - (1/u_0)x$. Thus each grid curve corresponding to $u = u_0$ lies in the plane $z = u_0^2 - (1/u_0)x$ and its projection onto the xy-plane is a parabola $x = ky^2$ with axis the x-axis. Similarly, if $v = v_0$ is held constant, then $x = uv_0^2$, $y = u^2v_0$ ⇒

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$y = (x/v_0^2)^2 v_0 = (1/v_0^3)x^2$, and $z = u^2 - v_0^2 = (1/v_0)y - v_0^2$. Each grid curve lies in the plane $z = (1/v_0)y - v_0^2$ and its projection onto the xy-plane is a parabola $y = kx^2$ with axis the y-axis. The surface is graph VI.

15. $\mathbf{r}(u, v) = (u^3 - u)\,\mathbf{i} + v^2\,\mathbf{j} + u^2\,\mathbf{k}$. The parametric equations for the surface are $x = u^3 - u,\ y = v^2,\ z = u^2$. If we fix u then x and z are constant so each corresponding grid curve is contained in a line parallel to the y-axis. (Since $y = v^2 \geq 0$, the grid curves are half-lines.) If v is held constant, then $y = v^2 =$ constant, so each grid curve is contained in a plane parallel to the xz-plane. Since x and z are functions of u only, the grid curves all have the same shape. The surface is the cylinder shown in graph I.

16. $x = (1 - u)(3 + \cos v)\cos 4\pi u,\ y = (1 - u)(3 + \cos v)\sin 4\pi u,\ z = 3u + (1 - u)\sin v$. These equations correspond to graph V: when $u = 0$, then $x = 3 + \cos v$, $y = 0$, and $z = \sin v$, which are equations of a circle with radius 1 in the xz-plane centered at $(3, 0, 0)$. When $u = \frac{1}{2}$, then $x = \frac{3}{2} + \frac{1}{2}\cos v$, $y = 0$, and $z = \frac{3}{2} + \frac{1}{2}\sin v$, which are equations of a circle with radius $\frac{1}{2}$ in the xz-plane centered at $\left(\frac{3}{2}, 0, \frac{3}{2}\right)$. When $u = 1$, then $x = y = 0$ and $z = 3$, giving the topmost point shown in the graph. This suggests that the grid curves with u constant are the vertically oriented circles visible on the surface. The spiralling grid curves correspond to keeping v constant.

17. $x = \cos^3 u \cos^3 v,\ y = \sin^3 u \cos^3 v,\ z = \sin^3 v$. If $v = v_0$ is held constant then $z = \sin^3 v_0$ is constant, so the corresponding grid curve lies in a horizontal plane. Several of the graphs exhibit horizontal grid curves, but the curves for this surface are neither ellipses nor straight lines, so graph III is the only possibility. (In fact, the horizontal grid curves here are members of the family $x = a\cos^3 u,\ y = a\sin^3 u$ and are called astroids.) The vertical grid curves we see on the surface correspond to $u = u_0$ held constant, as then we have $x = \cos^3 u_0 \cos^3 v,\ y = \sin^3 u_0 \cos^3 v$ so the corresponding grid curve lies in the vertical plane $y = (\tan^3 u_0)x$ through the z-axis.

18. $x = \sin u,\ y = \cos u \sin v,\ z = \sin v$. If $v = v_0$ is fixed, then $z = \sin v_0$ is constant, and $x = \sin u$, $y = (\sin v_0)\cos u$ describe an ellipse that is contained in the horizontal plane $z = \sin v_0$. If $u = u_0$ is fixed, then $x = \sin u_0$ is constant, and $y = (\cos u_0)\sin v$, $z = \sin v \quad \Rightarrow \quad y = (\cos u_0)z$, so the grid curves are portions of lines through the x-axis contained in the plane $x = \sin u_0$ (parallel to the yz-plane). The surface is graph II.

19. From Example 3, parametric equations for the plane through the point $(0, 0, 0)$ that contains the vectors $\mathbf{a} = \langle 1, -1, 0\rangle$ and $\mathbf{b} = \langle 0, 1, -1\rangle$ are $x = 0 + u(1) + v(0) = u,\ y = 0 + u(-1) + v(1) = v - u,\ z = 0 + u(0) + v(-1) = -v$.

20. From Example 3, parametric equations for the plane through the point $(0, -1, 5)$ that contains the vectors $\mathbf{a} = \langle 2, 1, 4\rangle$ and $\mathbf{b} = \langle -3, 2, 5\rangle$ are $x = 0 + u(2) + v(-3) = 2u - 3v,\ y = -1 + u(1) + v(2) = -1 + u + 2v$, $z = 5 + u(4) + v(5) = 5 + 4u + 5v$.

21. Solving the equation for x gives $x^2 = 1 + y^2 + \frac{1}{4}z^2 \quad \Rightarrow \quad x = \sqrt{1 + y^2 + \frac{1}{4}z^2}$. (We choose the positive root since we want the part of the hyperboloid that corresponds to $x \geq 0$.) If we let y and z be the parameters, parametric equations are $y = y$, $z = z,\ x = \sqrt{1 + y^2 + \frac{1}{4}z^2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. Solving the equation for y gives $y^2 = \frac{1}{2}(1 - x^2 - 3z^2) \Rightarrow y = -\sqrt{\frac{1}{2}(1 - x^2 - 3z^2)}$ (since we want the part of the ellipsoid that corresponds to $y \le 0$). If we let x and z be the parameters, parametric equations are $x = x,\ z = z,$ $y = -\sqrt{\frac{1}{2}(1 - x^2 - 3z^2)}$.

Alternate solution: The equation can be rewritten as $x^2 + \dfrac{y^2}{\left(1/\sqrt{2}\right)^2} + \dfrac{z^2}{(1/\sqrt{3})^2} = 1$, and if we let $x = u\cos v$ and $z = \dfrac{1}{\sqrt{3}}\, u \sin v$, then $y = -\sqrt{\frac{1}{2}(1 - x^2 - 3z^2)} = -\sqrt{\frac{1}{2}(1 - u^2\cos^2 v - u^2 \sin^2 v)} = -\sqrt{\frac{1}{2}(1 - u^2)}$, where $0 \le u \le 1$ and $0 \le v \le 2\pi$.

Second alternate solution: We can adapt the formulas for converting from spherical to rectangular coordinates as follows. We let $x = \sin\phi\cos\theta,\ y = \dfrac{1}{\sqrt{2}}\sin\phi\sin\theta,\ z = \dfrac{1}{\sqrt{3}}\cos\phi$; the surface is generated for $0 \le \phi \le \pi,\ \pi \le \theta \le 2\pi$.

23. Since the cone intersects the sphere in the circle $x^2 + y^2 = 2,\ z = \sqrt{2}$ and we want the portion of the sphere above this, we can parametrize the surface as $x = x,\ y = y,\ z = \sqrt{4 - x^2 - y^2}$ where $x^2 + y^2 \le 2$.

Alternate solution: Using spherical coordinates, $x = 2\sin\phi\cos\theta,\ y = 2\sin\phi\sin\theta,\ z = 2\cos\phi$ where $0 \le \phi \le \frac{\pi}{4}$ and $0 \le \theta \le 2\pi$.

24. We can parametrize the cylinder as $x = 3\cos\theta,\ y = y,\ z = 3\sin\theta$. To restrict the surface to that portion above the xy-plane and between the planes $y = -4$ and $y = 4$ we require $0 \le \theta \le \pi,\ -4 \le y \le 4$.

25. In spherical coordinates, parametric equations are $x = 6\sin\phi\cos\theta,\ y = 6\sin\phi\sin\theta,\ z = 6\cos\phi$. The intersection of the sphere with the plane $z = 3\sqrt{3}$ corresponds to $z = 6\cos\phi = 3\sqrt{3} \Rightarrow \cos\phi = \frac{\sqrt{3}}{2} \Rightarrow \phi = \frac{\pi}{6}$, and the plane $z = 0$ (the xy-plane) corresponds to $\phi = \frac{\pi}{2}$. Thus the surface is described by $\frac{\pi}{6} \le \phi \le \frac{\pi}{2},\ 0 \le \theta \le 2\pi$.

26. Using x and y as the parameters, $x = x,\ y = y,\ z = x + 3$ where $0 \le x^2 + y^2 \le 1$. Also, since the plane intersects the cylinder in an ellipse, the surface is a planar ellipse in the plane $z = x + 3$. Thus, parametrizing with respect to s and θ, we have $x = s\cos\theta,\ y = s\sin\theta,\ z = 3 + s\cos\theta$ where $0 \le s \le 1$ and $0 \le \theta \le 2\pi$.

27. The surface appears to be a portion of a circular cylinder of radius 3 with axis the x-axis. An equation of the cylinder is $y^2 + z^2 = 9$, and we can impose the restrictions $0 \le x \le 5,\ y \le 0$ to obtain the portion shown. To graph the surface on a CAS, we can use parametric equations $x = u,\ y = 3\cos v,\ z = 3\sin v$ with the parameter domain $0 \le u \le 5,\ \frac{\pi}{2} \le v \le \frac{3\pi}{2}$. Alternatively, we can regard x and z as parameters. Then parametric equations are $x = x,\ z = z,\ y = -\sqrt{9 - z^2}$, where $0 \le x \le 5$ and $-3 \le z \le 3$.

28. The surface appears to be a portion of a sphere of radius 1 centered at the origin. In spherical coordinates, the sphere has equation $\rho = 1$, and imposing the restrictions $\frac{\pi}{2} < \theta < 2\pi,\ \frac{\pi}{4} < \phi < \pi$ will give only the portion of the sphere shown. Thus, to graph the surface on a CAS we can either use spherical coordinates with the stated restrictions, or we can use parametric equations: $x = \sin\phi\cos\theta,\ y = \sin\phi\sin\theta,\ z = \cos\phi,\ \frac{\pi}{2} \le \theta \le 2\pi,\ \frac{\pi}{4} \le \phi \le \pi$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. Using Equations 3, we have the parametrization $x = x$,

$$y = \frac{1}{1+x^2}\cos\theta, \quad z = \frac{1}{1+x^2}\sin\theta, \quad -2 \le x \le 2, \quad 0 \le \theta \le 2\pi.$$



30. Letting θ be the angle of rotation about the y-axis (adapting Equations 3), we have the parametrization $x = (1/y)\cos\theta$, $y = y$, $z = (1/y)\sin\theta$, $y \ge 1$, $0 \le \theta \le 2\pi$.



31. (a) Replacing $\cos u$ by $\sin u$ and $\sin u$ by $\cos u$ gives parametric equations $x = (2 + \sin v)\sin u$, $y = (2 + \sin v)\cos u$, $z = u + \cos v$. From the graph, it appears that the direction of the spiral is reversed. We can verify this observation by noting that the projection of the spiral grid curves onto the xy-plane, given by $x = (2 + \sin v)\sin u$, $y = (2 + \sin v)\cos u$, $z = 0$, draws a circle in the clockwise direction for each value of v. The original equations, on the other hand, give circular projections drawn in the counterclockwise direction. The equation for z is identical in both surfaces, so as z increases, these grid curves spiral up in opposite directions for the two surfaces.



(b) Replacing $\cos u$ by $\cos 2u$ and $\sin u$ by $\sin 2u$ gives parametric equations $x = (2 + \sin v)\cos 2u$, $y = (2 + \sin v)\sin 2u$, $z = u + \cos v$. From the graph, it appears that the number of coils in the surface doubles within the same parametric domain. We can verify this observation by noting that the projection of the spiral grid curves onto the xy-plane, given by $x = (2 + \sin v)\cos 2u$, $y = (2 + \sin v)\sin 2u$, $z = 0$ (where v is constant), complete circular revolutions for $0 \le u \le \pi$ while the original surface requires $0 \le u \le 2\pi$ for a complete revolution. Thus, the new surface winds around twice as fast as the original surface, and since the equation for z is identical in both surfaces, we observe twice as many circular coils in the same z-interval.



32. First we graph the surface as viewed from the front, then from two additional viewpoints.



The surface appears as a twisted sheet, and is unusual because it has only one side. (The Möbius strip is discussed in more detail in Section 16.7.)

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

33. $\mathbf{r}(u,v) = (u+v)\,\mathbf{i} + 3u^2\,\mathbf{j} + (u-v)\,\mathbf{k}$.

$\mathbf{r}_u = \mathbf{i} + 6u\,\mathbf{j} + \mathbf{k}$ and $\mathbf{r}_v = \mathbf{i} - \mathbf{k}$, so $\mathbf{r}_u \times \mathbf{r}_v = -6u\,\mathbf{i} + 2\,\mathbf{j} - 6u\,\mathbf{k}$. Since the point $(2,3,0)$ corresponds to $u=1$, $v=1$, a normal vector to the surface at $(2,3,0)$ is $-6\,\mathbf{i} + 2\,\mathbf{j} - 6\,\mathbf{k}$, and an equation of the tangent plane is $-6x + 2y - 6z = -6$ or $3x - y + 3z = 3$.

34. $\mathbf{r}(u,v) = (u^2+1)\,\mathbf{i} + (v^3+1)\,\mathbf{j} + (u+v)\,\mathbf{k}$.

$\mathbf{r}_u = 2u\,\mathbf{i} + \mathbf{k}$ and $\mathbf{r}_v = 3v^2\,\mathbf{j} + \mathbf{k}$, so $\mathbf{r}_u \times \mathbf{r}_v = -3v^2\,\mathbf{i} - 2u\,\mathbf{j} + 6uv^2\,\mathbf{k}$. Since the point $(5,2,3)$ corresponds to $u=2$, $v=1$, a normal vector to the surface at $(5,2,3)$ is $-3\,\mathbf{i} - 4\,\mathbf{j} + 12\,\mathbf{k}$, and an equation of the tangent plane is $-3(x-5) - 4(y-2) + 12(z-3) = 0$ or $3x + 4y - 12z = -13$.

35. $\mathbf{r}(u,v) = u\cos v\,\mathbf{i} + u\sin v\,\mathbf{j} + v\,\mathbf{k} \;\Rightarrow\; \mathbf{r}\left(1, \frac{\pi}{3}\right) = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}, \frac{\pi}{3}\right)$.

$\mathbf{r}_u = \cos v\,\mathbf{i} + \sin v\,\mathbf{j}$ and $\mathbf{r}_v = -u\sin v\,\mathbf{i} + u\cos v\,\mathbf{j} + \mathbf{k}$, so a normal vector to the surface at the point $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}, \frac{\pi}{3}\right)$ is $\mathbf{r}_u\left(1, \frac{\pi}{3}\right) \times \mathbf{r}_v\left(1, \frac{\pi}{3}\right) = \left(\frac{1}{2}\,\mathbf{i} + \frac{\sqrt{3}}{2}\,\mathbf{j}\right) \times \left(-\frac{\sqrt{3}}{2}\,\mathbf{i} + \frac{1}{2}\,\mathbf{j} + \mathbf{k}\right) = \frac{\sqrt{3}}{2}\,\mathbf{i} - \frac{1}{2}\,\mathbf{j} + \mathbf{k}$. Thus an equation of the tangent plane at $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}, \frac{\pi}{3}\right)$ is $\frac{\sqrt{3}}{2}\left(x - \frac{1}{2}\right) - \frac{1}{2}\left(y - \frac{\sqrt{3}}{2}\right) + 1\left(z - \frac{\pi}{3}\right) = 0$ or $\frac{\sqrt{3}}{2}x - \frac{1}{2}y + z = \frac{\pi}{3}$.

36. $\mathbf{r}(u,v) = \sin u\,\mathbf{i} + \cos u\,\sin v\,\mathbf{j} + \sin v\,\mathbf{k} \;\Rightarrow\; \mathbf{r}\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = \left(\frac{1}{2}, \frac{\sqrt{3}}{4}, \frac{1}{2}\right)$.

$\mathbf{r}_u = \cos u\,\mathbf{i} - \sin u\sin v\,\mathbf{j}$ and $\mathbf{r}_v = \cos u\cos v\,\mathbf{j} + \cos v\,\mathbf{k}$, so a normal vector to the surface at the point $\left(\frac{1}{2}, \frac{\sqrt{3}}{4}, \frac{1}{2}\right)$ is $\mathbf{r}_u\left(\frac{\pi}{6}, \frac{\pi}{6}\right) \times \mathbf{r}_v\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = \left(\frac{\sqrt{3}}{2}\,\mathbf{i} - \frac{1}{4}\,\mathbf{j}\right) \times \left(\frac{3}{4}\,\mathbf{j} + \frac{\sqrt{3}}{2}\,\mathbf{k}\right) = -\frac{\sqrt{3}}{8}\,\mathbf{i} - \frac{3}{4}\,\mathbf{j} + \frac{3\sqrt{3}}{8}\,\mathbf{k}$.

Thus an equation of the tangent plane at $\left(\frac{1}{2}, \frac{\sqrt{3}}{4}, \frac{1}{2}\right)$ is $-\frac{\sqrt{3}}{8}\left(x - \frac{1}{2}\right) - \frac{3}{4}\left(y - \frac{\sqrt{3}}{4}\right) + \frac{3\sqrt{3}}{8}\left(z - \frac{1}{2}\right) = 0$ or $\sqrt{3}\,x + 6y - 3\sqrt{3}\,z = \frac{\sqrt{3}}{2}$ or $2x + 4\sqrt{3}\,y - 6z = 1$.

37. $\mathbf{r}(u,v) = u^2\,\mathbf{i} + 2u\sin v\,\mathbf{j} + u\cos v\,\mathbf{k} \;\Rightarrow\; \mathbf{r}(1,0) = (1,0,1)$.

$\mathbf{r}_u = 2u\,\mathbf{i} + 2\sin v\,\mathbf{j} + \cos v\,\mathbf{k}$ and $\mathbf{r}_v = 2u\cos v\,\mathbf{j} - u\sin v\,\mathbf{k}$, so a normal vector to the surface at the point $(1,0,1)$ is $\mathbf{r}_u(1,0) \times \mathbf{r}_v(1,0) = (2\,\mathbf{i} + \mathbf{k}) \times (2\,\mathbf{j}) = -2\,\mathbf{i} + 4\,\mathbf{k}$.

Thus an equation of the tangent plane at $(1,0,1)$ is $-2(x-1) + 0(y-0) + 4(z-1) = 0$ or $-x + 2z = 1$.



38. $\mathbf{r}(u,v) = (1 - u^2 - v^2)\,\mathbf{i} - v\,\mathbf{j} - u\,\mathbf{k}$.

$\mathbf{r}_u = -2u\,\mathbf{i} - \mathbf{k}$ and $\mathbf{r}_v = -2v\,\mathbf{i} - \mathbf{j}$. Since the point $(-1,-1,-1)$ corresponds to $u=1$, $v=1$, a normal vector to the surface at $(-1,-1,-1)$ is $\mathbf{r}_u(1,1) \times \mathbf{r}_v(1,1) = (-2\,\mathbf{i} - \mathbf{k}) \times (-2\,\mathbf{i} - \mathbf{j}) = -\mathbf{i} + 2\,\mathbf{j} + 2\,\mathbf{k}$.



Thus an equation of the tangent plane is $-1(x+1) + 2(y+1) + 2(z+1) = 0$ or $-x + 2y + 2z = -3$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. The surface S is given by $z = f(x, y) = 6 - 3x - 2y$ which intersects the xy-plane in the line $3x + 2y = 6$, so D is the triangular region given by $\{(x, y) \mid 0 \le x \le 2,\ 0 \le y \le 3 - \frac{3}{2}x\}$. By Formula 9, the surface area of S is

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\, dA \\ &= \iint_D \sqrt{1 + (-3)^2 + (-2)^2}\, dA = \sqrt{14} \iint_D dA = \sqrt{14}\, A(D) = \sqrt{14}\left(\tfrac{1}{2} \cdot 2 \cdot 3\right) = 3\sqrt{14}. \end{aligned}$$

40. $\mathbf{r}(u, v) = \langle u + v, 2 - 3u, 1 + u - v\rangle \ \Rightarrow\ \mathbf{r}_u = \langle 1, -3, 1\rangle$, $\mathbf{r}_v = \langle 1, 0, -1\rangle$, and $\mathbf{r}_u \times \mathbf{r}_v = \langle 3, 2, 3\rangle$. Then by Definition 6,

$$A(S) = \iint_D |\, \mathbf{r}_u \times \mathbf{r}_v \,|\, dA = \int_0^2 \int_{-1}^1 |\, \langle 3, 2, 3\rangle \,|\, dv\, du = \sqrt{22} \int_0^2 du \int_{-1}^1 dv = \sqrt{22}\,(2)(2) = 4\sqrt{22}$$

41. Here we can write $z = f(x, y) = \frac{1}{3} - \frac{1}{3}x - \frac{2}{3}y$ and D is the disk $x^2 + y^2 \le 3$, so by Formula 9 the area of the surface is

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\, dA = \iint_D \sqrt{1 + \left(-\tfrac{1}{3}\right)^2 + \left(-\tfrac{2}{3}\right)^2}\, dA = \tfrac{\sqrt{14}}{3} \iint_D dA \\ &= \tfrac{\sqrt{14}}{3}\, A(D) = \tfrac{\sqrt{14}}{3} \cdot \pi\left(\sqrt{3}\right)^2 = \sqrt{14}\,\pi \end{aligned}$$

42. $z = f(x, y) = \sqrt{x^2 + y^2} \ \Rightarrow\ \dfrac{\partial z}{\partial x} = \dfrac{1}{2}\left(x^2 + y^2\right)^{-1/2} \cdot 2x = \dfrac{x}{\sqrt{x^2 + y^2}}$, $\dfrac{\partial z}{\partial y} = \dfrac{y}{\sqrt{x^2 + y^2}}$, and

$$\sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} = \sqrt{1 + \frac{x^2}{x^2 + y^2} + \frac{y^2}{x^2 + y^2}} = \sqrt{1 + \frac{x^2 + y^2}{x^2 + y^2}} = \sqrt{2}$$

Here D is given by $\{(x, y) \mid 0 \le x \le 1,\ x^2 \le y \le x\}$, so by Formula 9, the surface area of S is

$$A(S) = \iint_D \sqrt{2}\, dA = \int_0^1 \int_{x^2}^x \sqrt{2}\, dy\, dx = \sqrt{2} \int_0^1 (x - x^2)\, dx = \sqrt{2}\left[\tfrac{1}{2}x^2 - \tfrac{1}{3}x^3\right]_0^1 = \sqrt{2}\left(\tfrac{1}{2} - \tfrac{1}{3}\right) = \tfrac{\sqrt{2}}{6}$$

43. $z = f(x, y) = \frac{2}{3}(x^{3/2} + y^{3/2})$ and $D = \{(x, y) \mid 0 \le x \le 1,\ 0 \le y \le 1\}$. Then $f_x = x^{1/2}$, $f_y = y^{1/2}$ and

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\sqrt{x}\right)^2 + \left(\sqrt{y}\right)^2}\, dA = \int_0^1 \int_0^1 \sqrt{1 + x + y}\, dy\, dx \\ &= \int_0^1 \left[\tfrac{2}{3}(x + y + 1)^{3/2}\right]_{y=0}^{y=1} dx = \tfrac{2}{3} \int_0^1 \left[(x + 2)^{3/2} - (x + 1)^{3/2}\right] dx \\ &= \tfrac{2}{3}\left[\tfrac{2}{5}(x + 2)^{5/2} - \tfrac{2}{5}(x + 1)^{5/2}\right]_0^1 = \tfrac{4}{15}(3^{5/2} - 2^{5/2} - 2^{5/2} + 1) = \tfrac{4}{15}(3^{5/2} - 2^{7/2} + 1) \end{aligned}$$

44. $z = f(x, y) = 4 - 2x^2 + y$ and $D = \{(x, y) \mid 0 \le x \le 1,\ 0 \le y \le x\}$. Thus, by Formula 9,

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + (-4x)^2 + (1)^2}\, dA = \int_0^1 \int_0^x \sqrt{16x^2 + 2}\, dy\, dx = \int_0^1 x\sqrt{16x^2 + 2}\, dx \\ &= \tfrac{1}{32} \cdot \tfrac{2}{3}(16x^2 + 2)^{3/2}\Big]_0^1 = \tfrac{1}{48}(18^{3/2} - 2^{3/2}) = \tfrac{1}{48}(54\sqrt{2} - 2\sqrt{2}) = \tfrac{13}{12}\sqrt{2} \end{aligned}$$

45. $z = f(x, y) = xy$ with $x^2 + y^2 \le 1$, so $f_x = y$, $f_y = x$ $\Rightarrow$

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + y^2 + x^2}\, dA = \int_0^{2\pi} \int_0^1 \sqrt{r^2 + 1}\, r\, dr\, d\theta = \int_0^{2\pi} \left[\tfrac{1}{3}\left(r^2 + 1\right)^{3/2}\right]_{r=0}^{r=1} d\theta \\ &= \int_0^{2\pi} \tfrac{1}{3}\left(2\sqrt{2} - 1\right) d\theta = \tfrac{2\pi}{3}\left(2\sqrt{2} - 1\right) \end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

46. A parametric representation of the surface is $x = z^2 + y,\ y = y,\ z = z$ with $0 \le y \le 2,\ 0 \le z \le 2$.

Hence $\mathbf{r}_y \times \mathbf{r}_z = (\mathbf{i} + \mathbf{j}) \times (2z\,\mathbf{i} + \mathbf{k}) = \mathbf{i} - \mathbf{j} - 2z\,\mathbf{k}$.

Then

$$A(S) = \iint_D |\,\mathbf{r}_y \times \mathbf{r}_z\,|\,dA = \int_0^2 \int_0^2 \sqrt{1+1+4z^2}\,dy\,dz = \int_0^2 2\sqrt{2+4z^2}\,dz$$

$$= \left[2 \cdot \tfrac{1}{2}\left(z\sqrt{2+4z^2} + \ln\left(2z + \sqrt{2+4z^2}\right)\right)\right]_0^2 \quad \left[\begin{array}{c}\text{Use trigonometric substitution}\\ \text{or Formula 21 in the Table of Integrals}\end{array}\right]$$

$$= 6\sqrt{2} + \ln\left(4 + 3\sqrt{2}\right) - \ln\sqrt{2} \text{ or } 6\sqrt{2} + \ln\frac{4+3\sqrt{2}}{\sqrt{2}} = 6\sqrt{2} + \ln\left(2\sqrt{2}+3\right)$$

Note: In general, if $x = f(y,z)$ then $\mathbf{r}_y \times \mathbf{r}_z = \mathbf{i} - \dfrac{\partial f}{\partial y}\,\mathbf{j} - \dfrac{\partial f}{\partial z}\,\mathbf{k}$ and $A(S) = \displaystyle\iint_D \sqrt{1 + \left(\frac{\partial f}{\partial y}\right)^2 + \left(\frac{\partial f}{\partial z}\right)^2}\,dA$.

47. A parametric representation of the surface is $x = x,\ y = x^2 + z^2,\ z = z$ with $0 \le x^2 + z^2 \le 16$.

Hence $\mathbf{r}_x \times \mathbf{r}_z = (\mathbf{i} + 2x\,\mathbf{j}) \times (2z\,\mathbf{j} + \mathbf{k}) = 2x\,\mathbf{i} - \mathbf{j} + 2z\,\mathbf{k}$.

Note: In general, if $y = f(x,z)$ then $\mathbf{r}_x \times \mathbf{r}_z = \dfrac{\partial f}{\partial x}\,\mathbf{i} - \mathbf{j} + \dfrac{\partial f}{\partial z}\,\mathbf{k}$, and $A(S) = \displaystyle\iint_D \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial z}\right)^2}\,dA$. Then

$$A(S) = \iint\limits_{0 \le x^2 + z^2 \le 16} \sqrt{1 + 4x^2 + 4z^2}\,dA = \int_0^{2\pi}\int_0^4 \sqrt{1+4r^2}\;r\,dr\,d\theta$$

$$= \int_0^{2\pi} d\theta \int_0^4 r\sqrt{1+4r^2}\,dr = 2\pi\left[\tfrac{1}{12}(1+4r^2)^{3/2}\right]_0^4 = \tfrac{\pi}{6}\left(65^{3/2} - 1\right)$$

48. $\mathbf{r}_u = \langle \cos v, \sin v, 0\rangle$, $\mathbf{r}_v = \langle -u\sin v, u\cos v, 1\rangle$, and $\mathbf{r}_u \times \mathbf{r}_v = \langle \sin v, -\cos v, u\rangle$. Then

$$A(S) = \int_0^{\pi}\int_0^1 \sqrt{1+u^2}\,du\,dv = \int_0^{\pi} dv \int_0^1 \sqrt{1+u^2}\,du$$

$$= \pi\left[\tfrac{u}{2}\sqrt{u^2+1} + \tfrac{1}{2}\ln\left|u + \sqrt{u^2+1}\,\right|\right]_0^1 = \tfrac{\pi}{2}\left[\sqrt{2} + \ln\left(1+\sqrt{2}\right)\right]$$

49. $\mathbf{r}_u = \langle 2u, v, 0\rangle$, $\mathbf{r}_v = \langle 0, u, v\rangle$, and $\mathbf{r}_u \times \mathbf{r}_v = \langle v^2, -2uv, 2u^2\rangle$. Then

$$A(S) = \iint_D |\mathbf{r}_u \times \mathbf{r}_v|\,dA = \int_0^1\int_0^2 \sqrt{v^4 + 4u^2v^2 + 4u^4}\,dv\,du = \int_0^1\int_0^2 \sqrt{(v^2+2u^2)^2}\,dv\,du$$

$$= \int_0^1\int_0^2 (v^2 + 2u^2)\,dv\,du = \int_0^1 \left[\tfrac{1}{3}v^3 + 2u^2v\right]_{v=0}^{v=2} du = \int_0^1 \left(\tfrac{8}{3} + 4u^2\right) du = \left[\tfrac{8}{3}u + \tfrac{4}{3}u^3\right]_0^1 = 4$$

50. The cylinder encloses separate portions of the sphere in the upper and lower halves. The top half of the sphere is $z = f(x,y) = \sqrt{b^2 - x^2 - y^2}$ and D is given by $\{(x,y) \mid x^2 + y^2 \le a^2\}$. By Formula 9, the surface area of the upper enclosed portion is

$$A = \iint_D \sqrt{1 + \left(\frac{x}{\sqrt{b^2 - x^2 - y^2}}\right)^2 + \left(\frac{y}{\sqrt{b^2 - x^2 - y^2}}\right)^2}\,dA = \iint_D \sqrt{1 + \frac{x^2+y^2}{b^2 - x^2 - y^2}}\,dA$$

$$= \iint_D \sqrt{\frac{b^2}{b^2 - x^2 - y^2}}\,dA = \int_0^{2\pi}\int_0^a \frac{b}{\sqrt{b^2 - r^2}}\,r\,dr\,d\theta = b\int_0^{2\pi} d\theta \int_0^a \frac{r}{\sqrt{b^2 - r^2}}\,dr$$

$$= b\left[\theta\right]_0^{2\pi}\left[-\sqrt{b^2 - r^2}\,\right]_0^a = 2\pi b\left(-\sqrt{b^2 - a^2} + \sqrt{b^2 - 0}\right) = 2\pi b\left(b - \sqrt{b^2 - a^2}\right)$$

The lower portion of the sphere enclosed by the cylinder has identical shape, so the total area is $2A = 4\pi b\left(b - \sqrt{b^2 - a^2}\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. From Equation 9 we have $A(S) = \iint_D \sqrt{1 + (f_x)^2 + (f_y)^2}\, dA$. But if $|f_x| \le 1$ and $|f_y| \le 1$ then $0 \le (f_x)^2 \le 1$, $0 \le (f_y)^2 \le 1 \;\Rightarrow\; 1 \le 1 + (f_x)^2 + (f_y)^2 \le 3 \;\Rightarrow\; 1 \le \sqrt{1 + (f_x)^2 + (f_y)^2} \le \sqrt{3}$. By Property 15.2.11, $\iint_D 1\, dA \le \iint_D \sqrt{1 + (f_x)^2 + (f_y)^2}\, dA \le \iint_D \sqrt{3}\, dA \;\Rightarrow\; A(D) \le A(S) \le \sqrt{3}\, A(D) \;\Rightarrow$ $\pi R^2 \le A(S) \le \sqrt{3}\pi R^2$.

52. $z = f(x, y) = \cos(x^2 + y^2)$ with $x^2 + y^2 \le 1$.

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + (-2x\sin(x^2+y^2))^2 + (-2y\sin(x^2+y^2))^2}\, dA \\ &= \iint_D \sqrt{1 + 4x^2\sin^2(x^2+y^2) + 4y^2\sin^2(x^2+y^2)}\, dA = \iint_D \sqrt{1 + 4(x^2+y^2)\sin^2(x^2+y^2)}\, dA \\ &= \int_0^{2\pi}\int_0^1 \sqrt{1 + 4r^2\sin^2(r^2)}\, r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_0^1 r\sqrt{1 + 4r^2\sin^2(r^2)}\, dr \\ &= 2\pi \int_0^1 r\sqrt{1 + 4r^2\sin^2(r^2)}\, dr \approx 4.1073 \end{aligned}$$

53. $z = f(x, y) = \ln(x^2 + y^2 + 2)$ with $x^2 + y^2 \le 1$.

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\frac{2x}{x^2+y^2+2}\right)^2 + \left(\frac{2y}{x^2+y^2+2}\right)^2}\, dA = \iint_D \sqrt{1 + \frac{4x^2+4y^2}{(x^2+y^2+2)^2}}\, dA \\ &= \int_0^{2\pi}\int_0^1 \sqrt{1 + \frac{4r^2}{(r^2+2)^2}}\, r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_0^1 r\sqrt{\frac{(r^2+2)^2 + 4r^2}{(r^2+2)^2}}\, dr = 2\pi \int_0^1 \frac{r\sqrt{r^4+8r^2+4}}{r^2+2}\, dr \approx 3.5618 \end{aligned}$$

54. Let $f(x, y) = \dfrac{1 + x^2}{1 + y^2}$. Then $f_x = \dfrac{2x}{1 + y^2}$,

$f_y = (1 + x^2)\left[-\dfrac{2y}{(1 + y^2)^2}\right] = -\dfrac{2y(1 + x^2)}{(1 + y^2)^2}$.

We use a CAS to estimate

$\int_{-1}^{1}\int_{-(1-|x|)}^{1-|x|} \sqrt{1 + f_x^2 + f_y^2}\, dy\, dx \approx 2.6959$.

In order to graph only the part of the surface above the square, we use $-(1 - |x|) \le y \le 1 - |x|$ as the y-range in our plot command.



55. (a) $A(S) = \iint_D \sqrt{1 + \left(\dfrac{\partial z}{\partial x}\right)^2 + \left(\dfrac{\partial z}{\partial y}\right)^2}\, dA = \int_0^6\int_0^4 \sqrt{1 + \dfrac{4x^2+4y^2}{(1+x^2+y^2)^4}}\, dy\, dx.$

Using the Midpoint Rule with $f(x, y) = \sqrt{1 + \dfrac{4x^2+4y^2}{(1+x^2+y^2)^4}}$, $m = 3$, $n = 2$ we have

$$A(S) \approx \sum_{i=1}^{3}\sum_{j=1}^{2} f(\overline{x}_i, \overline{y}_j)\, \Delta A = 4\,[f(1,1) + f(1,3) + f(3,1) + f(3,3) + f(5,1) + f(5,3)] \approx 24.2055$$

(b) Using a CAS we have $A(S) = \int_0^6\int_0^4 \sqrt{1 + \dfrac{4x^2+4y^2}{(1+x^2+y^2)^4}}\, dy\, dx \approx 24.2476$. This agrees with the estimate in part (a) to the first decimal place.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

56. $\mathbf{r}(u,v) = \langle \cos^3 u \cos^3 v, \sin^3 u \cos^3 v, \sin^3 v \rangle$, so $\mathbf{r}_u = \langle -3\cos^2 u \sin u \cos^3 v, 3\sin^2 u \cos u \cos^3 v, 0 \rangle$,

$\mathbf{r}_v = \langle -3\cos^3 u \cos^2 v \sin v, -3\sin^3 u \cos^2 v \sin v, 3\sin^2 v \cos v \rangle$, and

$\mathbf{r}_u \times \mathbf{r}_v = \langle 9\cos u \sin^2 u \cos^4 v \sin^2 v, 9\cos^2 u \sin u \cos^4 v \sin^2 v, 9\cos^2 u \sin^2 u \cos^5 v \sin v \rangle$. Then

$$\begin{aligned} |\mathbf{r}_u \times \mathbf{r}_v| &= 9\sqrt{\cos^2 u \sin^4 u \cos^8 v \sin^4 v + \cos^4 u \sin^2 u \cos^8 v \sin^4 v + \cos^4 u \sin^4 u \cos^{10} v \sin^2 v} \\ &= 9\sqrt{\cos^2 u \sin^2 u \cos^8 v \sin^2 v (\sin^2 v + \cos^2 u \sin^2 u \cos^2 v)} \\ &= 9\cos^4 v\,|\cos u \sin u \sin v|\sqrt{\sin^2 v + \cos^2 u \sin^2 u \cos^2 v} \end{aligned}$$

Using a CAS, we have $A(S) = \int_0^{\pi}\int_0^{2\pi} 9\cos^4 v\,|\cos u \sin u \sin v|\sqrt{\sin^2 v + \cos^2 u \sin^2 u \cos^2 v}\,dv\,du \approx 4.4506$.

57. $z = 1 + 2x + 3y + 4y^2$, so

$$A(S) = \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\,dA = \int_1^4\int_0^1 \sqrt{1 + 4 + (3+8y)^2}\,dy\,dx = \int_1^4\int_0^1 \sqrt{14 + 48y + 64y^2}\,dy\,dx.$$

Using a CAS, we have

$\int_1^4\int_0^1 \sqrt{14 + 48y + 64y^2}\,dy\,dx = \frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\left(11\sqrt{5} + 3\sqrt{14}\sqrt{5}\right) - \frac{15}{16}\ln\left(3\sqrt{5} + \sqrt{14}\sqrt{5}\right)$

or $\frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\frac{11\sqrt{5}+3\sqrt{70}}{3\sqrt{5}+\sqrt{70}}$.

58. (a) $\mathbf{r}_u = a\cos v\,\mathbf{i} + b\sin v\,\mathbf{j} + 2u\,\mathbf{k}$, $\mathbf{r}_v = -au\sin v\,\mathbf{i} + bu\cos v\,\mathbf{j} + 0\,\mathbf{k}$, and

$\mathbf{r}_u \times \mathbf{r}_v = -2bu^2\cos v\,\mathbf{i} - 2au^2\sin v\,\mathbf{j} + abu\,\mathbf{k}$.

$A(S) = \int_0^{2\pi}\int_0^2 |\mathbf{r}_u \times \mathbf{r}_v|\,du\,dv = \int_0^{2\pi}\int_0^2 \sqrt{4b^2u^4\cos^2 v + 4a^2u^4\sin^2 v + a^2b^2u^2}\,du\,dv$

(b) $x^2 = a^2u^2\cos^2 v$, $y^2 = b^2u^2\sin^2 v$, $z = u^2 \;\Rightarrow\; x^2/a^2 + y^2/b^2 = u^2 = z$ which is an elliptic paraboloid. To find D, notice that $0 \le u \le 2 \;\Rightarrow\; 0 \le z \le 4 \;\Rightarrow\; 0 \le x^2/a^2 + y^2/b^2 \le 4$. Therefore, using Formula 9, we have

$$A(S) = \int_{-2a}^{2a}\int_{-b\sqrt{4-(x^2/a^2)}}^{b\sqrt{4-(x^2/a^2)}} \sqrt{1 + (2x/a^2)^2 + (2y/b^2)^2}\,dy\,dx.$$

(c)



(d) We substitute $a = 2$, $b = 3$ in the integral in part (a) to get $A(S) = \int_0^{2\pi}\int_0^2 2u\sqrt{9u^2\cos^2 v + 4u^2\sin^2 v + 9}\,du\,dv$. We use a CAS to estimate the integral accurate to four decimal places. To speed up the calculation, we can set `Digits:=7;` (in Maple) or use the approximation command `N` (in Mathematica). We find that $A(S) \approx 115.6596$.

59. (a) $x = a\sin u\cos v$, $y = b\sin u\sin v$, $z = c\cos u \;\Rightarrow$

$$\begin{aligned} \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} &= (\sin u\cos v)^2 + (\sin u\sin v)^2 + (\cos u)^2 \\ &= \sin^2 u + \cos^2 u = 1 \end{aligned}$$

and since the ranges of u and v are sufficient to generate the entire graph, the parametric equations represent an ellipsoid.

(b)



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) From the parametric equations (with $a = 1$, $b = 2$, and $c = 3$),

we calculate $\mathbf{r}_u = \cos u \cos v\,\mathbf{i} + 2\cos u \sin v\,\mathbf{j} - 3\sin u\,\mathbf{k}$ and

$\mathbf{r}_v = -\sin u \sin v\,\mathbf{i} + 2\sin u \cos v\,\mathbf{j}$. So $\mathbf{r}_u \times \mathbf{r}_v = 6\sin^2 u \cos v\,\mathbf{i} + 3\sin^2 u \sin v\,\mathbf{j} + 2\sin u \cos u\,\mathbf{k}$, and the surface area is given by $A(S) = \int_0^{2\pi}\int_0^{\pi} |\mathbf{r}_u \times \mathbf{r}_v|\,du\,dv = \int_0^{2\pi}\int_0^{\pi} \sqrt{36\sin^4 u\cos^2 v + 9\sin^4 u \sin^2 v + 4\cos^2 u \sin^2 u}\,du\,dv$

60. (a) $x = a\cosh u \cos v$, $y = b\cosh u \sin v$, $z = c\sinh u \;\Rightarrow$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = \cosh^2 u \cos^2 v + \cosh^2 u \sin^2 v - \sinh^2 u$$
$$= \cosh^2 u - \sinh^2 u = 1$$

and the parametric equations represent a hyperboloid of one sheet.

(b)



(c) $\mathbf{r}_u = \sinh u \cos v\,\mathbf{i} + 2\sinh u \sin v\,\mathbf{j} + 3\cosh u\,\mathbf{k}$ and

$\mathbf{r}_v = -\cosh u \sin v\,\mathbf{i} + 2\cosh u \cos v\,\mathbf{j}$, so $\mathbf{r}_u \times \mathbf{r}_v = -6\cosh^2 u \cos v\,\mathbf{i} - 3\cosh^2 u \sin v\,\mathbf{j} + 2\cosh u \sinh u\,\mathbf{k}$.

We integrate between $u = \sinh^{-1}(-1) = -\ln\left(1+\sqrt{2}\right)$ and $u = \sinh^{-1} 1 = \ln\left(1+\sqrt{2}\right)$, since then z varies between -3 and 3, as desired. So the surface area is

$$A(S) = \int_0^{2\pi}\int_{-\ln(1+\sqrt{2})}^{\ln(1+\sqrt{2})} |\mathbf{r}_u \times \mathbf{r}_v|\,du\,dv$$
$$= \int_0^{2\pi}\int_{-\ln(1+\sqrt{2})}^{\ln(1+\sqrt{2})} \sqrt{36\cosh^4 u \cos^2 v + 9\cosh^4 u \sin^2 v + 4\cosh^2 u \sinh^2 u}\,du\,dv$$

61. To find the region D: $z = x^2 + y^2$ implies $z + z^2 = 4z$ or $z^2 - 3z = 0$. Thus $z = 0$ or $z = 3$ are the planes where the surfaces intersect. But $x^2 + y^2 + z^2 = 4z$ implies $x^2 + y^2 + (z-2)^2 = 4$, so $z = 3$ intersects the upper hemisphere. Thus $(z-2)^2 = 4 - x^2 - y^2$ or $z = 2 + \sqrt{4 - x^2 - y^2}$. Therefore D is the region inside the circle $x^2 + y^2 + (3-2)^2 = 4$, that is, $D = \{(x,y) \mid x^2 + y^2 \le 3\}$.

$$A(S) = \iint_D \sqrt{1 + [(-x)(4-x^2-y^2)^{-1/2}]^2 + [(-y)(4-x^2-y^2)^{-1/2}]^2}\,dA$$
$$= \int_0^{2\pi}\int_0^{\sqrt{3}} \sqrt{1 + \frac{r^2}{4-r^2}}\,r\,dr\,d\theta = \int_0^{2\pi}\int_0^{\sqrt{3}} \frac{2r\,dr}{\sqrt{4-r^2}}\,d\theta = \int_0^{2\pi}\left[-2(4-r^2)^{1/2}\right]_{r=0}^{r=\sqrt{3}} d\theta$$
$$= \int_0^{2\pi}(-2+4)\,d\theta = 2\theta\Big]_0^{2\pi} = 4\pi$$

62. We first find the area of the face of the surface that intersects the positive y-axis. A parametric representation of the surface is $x = x$, $y = \sqrt{1-z^2}$, $z = z$ with $x^2 + z^2 \le 1$. Then $\mathbf{r}(x,z) = \langle x, \sqrt{1-z^2}, z\rangle \;\Rightarrow\; \mathbf{r}_x = \langle 1, 0, 0\rangle$,

$\mathbf{r}_z = \langle 0, -z/\sqrt{1-z^2}, 1\rangle$ and $\mathbf{r}_x \times \mathbf{r}_z = \langle 0, -1, -z/\sqrt{1-z^2}\,\rangle \;\Rightarrow\; |\mathbf{r}_x \times \mathbf{r}_z| = \sqrt{1 + \dfrac{z^2}{1-z^2}} = \dfrac{1}{\sqrt{1-z^2}}$.

$$A(S) = \iint_{x^2+z^2\le 1} |\mathbf{r}_x \times \mathbf{r}_z|\,dA = \int_{-1}^{1}\int_{-\sqrt{1-z^2}}^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz = 4\int_0^1\int_0^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz$$

[by the symmetry of the surface]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

This integral is improper [when $z = 1$], so

$$A(S) = \lim_{t\to 1^-} 4\int_0^t \int_0^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz = \lim_{t\to 1^-} 4\int_0^t \frac{\sqrt{1-z^2}}{\sqrt{1-z^2}}\,dz = \lim_{t\to 1^-} 4\int_0^t dz = \lim_{t\to 1^-} 4t = 4$$

Since the complete surface consists of four congruent faces, the total surface area is $4(4) = 16$.

Alternate solution: The face of the surface that intersects the positive y-axis can also be parametrized as $\mathbf{r}(x,\theta) = \langle x, \cos\theta, \sin\theta\rangle$ for $-\frac{\pi}{2} \le \theta \le \frac{\pi}{2}$ and $x^2 + z^2 \le 1 \quad\Leftrightarrow\quad x^2 + \sin^2\theta \le 1 \quad\Leftrightarrow$ $-\sqrt{1-\sin^2\theta} \le x \le \sqrt{1-\sin^2\theta} \quad\Leftrightarrow\quad -\cos\theta \le x \le \cos\theta$. Then $\mathbf{r}_x = \langle 1,0,0\rangle$, $\mathbf{r}_\theta = \langle 0, -\sin\theta, \cos\theta\rangle$ and $\mathbf{r}_x \times \mathbf{r}_\theta = \langle 0, -\cos\theta, -\sin\theta\rangle \quad\Rightarrow\quad |\,\mathbf{r}_x \times \mathbf{r}_\theta\,| = 1$, so $A(S) = \int_{-\pi/2}^{\pi/2}\int_{-\cos\theta}^{\cos\theta} 1\,dx\,d\theta = \int_{-\pi/2}^{\pi/2} 2\cos\theta\,d\theta = 2\sin\theta\big]_{-\pi/2}^{\pi/2} = 4$. Again, the area of the complete surface is $4(4) = 16$.

63. Let $A(S_1)$ be the surface area of that portion of the surface which lies above the plane $z = 0$. Then $A(S) = 2A(S_1)$. Following Example 10, a parametric representation of S_1 is $x = a\sin\phi\cos\theta$, $y = a\sin\phi\sin\theta$, $z = a\cos\phi$ and $|\mathbf{r}_\phi \times \mathbf{r}_\theta| = a^2\sin\phi$. For D, $0 \le \phi \le \frac{\pi}{2}$ and for each fixed ϕ, $\left(x - \frac{1}{2}a\right)^2 + y^2 \le \left(\frac{1}{2}a\right)^2$ or $\left[a\sin\phi\cos\theta - \frac{1}{2}a\right]^2 + a^2\sin^2\phi\sin^2\theta \le (a/2)^2$ implies $a^2\sin^2\phi - a^2\sin\phi\cos\theta \le 0$ or $\sin\phi\,(\sin\phi - \cos\theta) \le 0$. But $0 \le \phi \le \frac{\pi}{2}$, so $\cos\theta \ge \sin\phi$ or $\sin\left(\frac{\pi}{2} + \theta\right) \ge \sin\phi$ or $\phi - \frac{\pi}{2} \le \theta \le \frac{\pi}{2} - \phi$. Hence $D = \left\{(\phi,\theta) \mid 0 \le \phi \le \frac{\pi}{2},\ \phi - \frac{\pi}{2} \le \theta \le \frac{\pi}{2} - \phi\right\}$. Then

$$\begin{aligned} A(S_1) &= \int_0^{\pi/2}\int_{\phi-(\pi/2)}^{(\pi/2)-\phi} a^2\sin\phi\,d\theta\,d\phi = a^2\int_0^{\pi/2}(\pi - 2\phi)\sin\phi\,d\phi \\ &= a^2\left[(-\pi\cos\phi) - 2(-\phi\cos\phi + \sin\phi)\right]_0^{\pi/2} = a^2(\pi - 2) \end{aligned}$$

Thus $A(S) = 2a^2(\pi - 2)$.

Alternate solution: Working on S_1 we could parametrize the portion of the sphere by $x = x$, $y = y$, $z = \sqrt{a^2 - x^2 - y^2}$. Then $|\mathbf{r}_x \times \mathbf{r}_y| = \sqrt{1 + \frac{x^2}{a^2 - x^2 - y^2} + \frac{y^2}{a^2 - x^2 - y^2}} = \frac{a}{\sqrt{a^2 - x^2 - y^2}}$ and

$$A(S_1) = \iint_{0 \le (x-(a/2))^2 + y^2 \le (a/2)^2} \frac{a}{\sqrt{a^2 - x^2 - y^2}}\,dA = \int_{-\pi/2}^{\pi/2}\int_0^{a\cos\theta} \frac{a}{\sqrt{a^2 - r^2}}\,r\,dr\,d\theta$$

$$\begin{aligned} &= \int_{-\pi/2}^{\pi/2} -a(a^2 - r^2)^{1/2}\Big]_{r=0}^{r=a\cos\theta} d\theta = \int_{-\pi/2}^{\pi/2} a^2[1 - (1 - \cos^2\theta)^{1/2}]\,d\theta \\ &= \int_{-\pi/2}^{\pi/2} a^2(1 - |\sin\theta|)\,d\theta = 2a^2\int_0^{\pi/2}(1 - \sin\theta)\,d\theta = 2a^2\left(\tfrac{\pi}{2} - 1\right) \end{aligned}$$

Thus $A(S) = 4a^2\left(\frac{\pi}{2} - 1\right) = 2a^2(\pi - 2)$.

Notes:

(1) Perhaps working in spherical coordinates is the most obvious approach here. However, you must be careful in setting up D.

(2) In the alternate solution, you can avoid having to use $|\sin\theta|$ by working in the first octant and then multiplying by 4. However, if you set up S_1 as above and arrived at $A(S_1) = a^2\pi$, you now see your error.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. (a) Here $z = a\sin\alpha$, $y = |AB|$, and $x = |OA|$. But

$|OB| = |OC| + |CB| = b + a\cos\alpha$ and $\sin\theta = \dfrac{|AB|}{|OB|}$ so that

$y = |OB|\sin\theta = (b + a\cos\alpha)\sin\theta$. Similarly $\cos\theta = \dfrac{|OA|}{|OB|}$ so

$x = (b + a\cos\alpha)\cos\theta$. Hence a parametric representation for the torus is $x = b\cos\theta + a\cos\alpha\cos\theta$, $y = b\sin\theta + a\cos\alpha\sin\theta$, $z = a\sin\alpha$, where $0 \le \alpha \le 2\pi$, $0 \le \theta \le 2\pi$.



(b)



$a = 1, b = 8$



$a = 3, b = 8$



$a = 3, b = 4$

(c) $x = b\cos\theta + a\cos\alpha\cos\theta$, $y = b\sin\theta + a\cos\alpha\sin\theta$, $z = a\sin\alpha$, so $\mathbf{r}_\alpha = \langle -a\sin\alpha\cos\theta, -a\sin\alpha\sin\theta, a\cos\alpha\rangle$, $\mathbf{r}_\theta = \langle -(b + a\cos\alpha)\sin\theta, (b + a\cos\alpha)\cos\theta, 0\rangle$ and

$$\begin{aligned}\mathbf{r}_\alpha \times \mathbf{r}_\theta &= \left(-ab\cos\alpha\cos\theta - a^2\cos\alpha\cos^2\theta\right)\mathbf{i} + \left(-ab\sin\alpha\cos\theta - a^2\sin\alpha\cos^2\theta\right)\mathbf{j} \\ &\quad + \left(-ab\cos^2\alpha\sin\theta - a^2\cos^2\alpha\sin\theta\cos\theta - ab\sin^2\alpha\sin\theta - a^2\sin^2\alpha\sin\theta\cos\theta\right)\mathbf{k} \\ &= -a(b + a\cos\alpha)\left[(\cos\theta\cos\alpha)\mathbf{i} + (\sin\theta\cos\alpha)\mathbf{j} + (\sin\alpha)\mathbf{k}\right]\end{aligned}$$

Then $|\mathbf{r}_\alpha \times \mathbf{r}_\theta| = a(b + a\cos\alpha)\sqrt{\cos^2\theta\cos^2\alpha + \sin^2\theta\cos^2\alpha + \sin^2\alpha} = a(b + a\cos\alpha)$.

Note: $b > a$, $-1 \le \cos\alpha \le 1$ so $|b + a\cos\alpha| = b + a\cos\alpha$. Hence

$A(S) = \int_0^{2\pi}\int_0^{2\pi} a(b + a\cos\alpha)\,d\alpha\,d\theta = 2\pi\left[ab\alpha + a^2\sin\alpha\right]_0^{2\pi} = 4\pi^2 ab.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

16.7 Surface Integrals

1. The box is a cube where each face has surface area 4. The centers of the faces are $(\pm1, 0, 0)$, $(0, \pm1, 0)$, $(0, 0, \pm1)$. For each face we take the point P_{ij}^* to be the center of the face and $f(x, y, z) = \cos(x + 2y + 3z)$, so by Definition 1,

$$\begin{aligned}\iint_S f(x, y, z)\, dS &\approx [f(1, 0, 0)](4) + [f(-1, 0, 0)](4) + [f(0, 1, 0)](4) \\ &\quad + [f(0, -1, 0)](4) + [f(0, 0, 1)](4) + [f(0, 0, -1)](4) \\ &= 4\,[\cos 1 + \cos(-1) + \cos 2 + \cos(-2) + \cos 3 + \cos(-3)] \approx -6.93\end{aligned}$$

2. Each quarter-cylinder has surface area $\frac{1}{4}[2\pi(1)(2)] = \pi$, and the top and bottom disks have surface area $\pi(1)^2 = \pi$. We can take $(0, 0, 1)$ as a sample point in the top disk, $(0, 0, -1)$ in the bottom disk, and $(\pm1, 0, 0)$, $(0, \pm1, 0)$ in the four quarter-cylinders. Then $\iint_S f(x, y, z)\, dS$ can be approximated by the Riemann sum

$$\begin{aligned}&f(1, 0, 0)(\pi) + f(-1, 0, 0)(\pi) + f(0, 1, 0)\,(\pi) + f(0, -1, 0)(\pi) + f(0, 0, 1)(\pi) + f(0, 0, -1)(\pi) \\ &\qquad = (2 + 2 + 3 + 3 + 4 + 4)\pi = 18\pi \approx 56.5.\end{aligned}$$

3. We can use the xz- and yz-planes to divide H into four patches of equal size, each with surface area equal to $\frac{1}{8}$ the surface area of a sphere with radius $\sqrt{50}$, so $\Delta S = \frac{1}{8}(4)\pi\left(\sqrt{50}\right)^2 = 25\pi$. Then $(\pm3, \pm4, 5)$ are sample points in the four patches, and using a Riemann sum as in Definition 1, we have

$$\begin{aligned}\iint_H f(x, y, z)\, dS &\approx f(3, 4, 5)\,\Delta S + f(3, -4, 5)\,\Delta S + f(-3, 4, 5)\,\Delta S + f(-3, -4, 5)\,\Delta S \\ &= (7 + 8 + 9 + 12)(25\pi) = 900\pi \approx 2827\end{aligned}$$

4. On the surface, $f(x, y, z) = g\left(\sqrt{x^2 + y^2 + z^2}\right) = g(2) = -5$. So since the area of a sphere is $4\pi r^2$,

$$\iint_S f(x, y, z)\, dS = \iint_S g(2)\, dS = -5 \iint_S dS = -5[4\pi(2)^2] = -80\pi.$$

5. $\mathbf{r}(u, v) = (u + v)\,\mathbf{i} + (u - v)\,\mathbf{j} + (1 + 2u + v)\,\mathbf{k}$, $0 \le u \le 2$, $0 \le v \le 1$ and

$\mathbf{r}_u \times \mathbf{r}_v = (\mathbf{i} + \mathbf{j} + 2\,\mathbf{k}) \times (\mathbf{i} - \mathbf{j} + \mathbf{k}) = 3\,\mathbf{i} + \mathbf{j} - 2\,\mathbf{k} \quad \Rightarrow \quad |\mathbf{r}_u \times \mathbf{r}_v| = \sqrt{3^2 + 1^2 + (-2)^2} = \sqrt{14}$. Then by Formula 2,

$$\begin{aligned}\iint_S (x + y + z)\, dS &= \iint_D (u + v + u - v + 1 + 2u + v)\, |\mathbf{r}_u \times \mathbf{r}_v|\, dA = \int_0^1 \int_0^2 (4u + v + 1) \cdot \sqrt{14}\, du\, dv \\ &= \sqrt{14} \int_0^1 \left[2u^2 + uv + u\right]_{u=0}^{u=2} dv = \sqrt{14} \int_0^1 (2v + 10)\, dv = \sqrt{14}\left[v^2 + 10v\right]_0^1 = 11\sqrt{14}\end{aligned}$$

6. $\mathbf{r}(u, v) = u\cos v\,\mathbf{i} + u\sin v\,\mathbf{j} + u\,\mathbf{k}$, $0 \le u \le 1$, $0 \le v \le \pi/2$ and

$\mathbf{r}_u \times \mathbf{r}_v = (\cos v\,\mathbf{i} + \sin v\,\mathbf{j} + \mathbf{k}) \times (-u\sin v\,\mathbf{i} + u\cos v\,\mathbf{j}) = -u\cos v\,\mathbf{i} - u\sin v\,\mathbf{j} + u\,\mathbf{k} \quad \Rightarrow$

$|\mathbf{r}_u \times \mathbf{r}_v| = \sqrt{u^2\cos^2 v + u^2\sin^2 v + u^2} = \sqrt{2u^2} = \sqrt{2}\,u$ [since $u \ge 0$]. Then by Formula 2,

$$\begin{aligned}\iint_S xyz\, dS &= \iint_D (u\cos v)(u\sin v)(u)\, |\mathbf{r}_u \times \mathbf{r}_v|\, dA = \int_0^1 \int_0^{\pi/2} (u^3 \sin v\cos v) \cdot \sqrt{2}\,u\, dv\, du \\ &= \sqrt{2}\int_0^1 u^4\, du \int_0^{\pi/2} \sin v\cos v\, dv = \sqrt{2}\left[\tfrac{1}{5}u^5\right]_0^1 \left[\tfrac{1}{2}\sin^2 v\right]_0^{\pi/2} = \sqrt{2}\cdot\tfrac{1}{5}\cdot\tfrac{1}{2} = \tfrac{1}{10}\sqrt{2}\end{aligned}$$

7. $\mathbf{r}(u, v) = \langle u\cos v, u\sin v, v\rangle$, $0 \le u \le 1$, $0 \le v \le \pi$ and

$\mathbf{r}_u \times \mathbf{r}_v = \langle\cos v, \sin v, 0\rangle \times \langle -u\sin v, u\cos v, 1\rangle = \langle\sin v, -\cos v, u\rangle \quad \Rightarrow$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$|\mathbf{r}_u \times \mathbf{r}_v| = \sqrt{\sin^2 v + \cos^2 v + u^2} = \sqrt{u^2+1}$. Then

$$\iint_S y\,dS = \iint_D (u\sin v)\ |\mathbf{r}_u \times \mathbf{r}_v|\ dA = \int_0^1 \int_0^\pi (u\sin v)\cdot\sqrt{u^2+1}\,dv\,du = \int_0^1 u\sqrt{u^2+1}\,du\ \int_0^\pi \sin v\,dv$$
$$= \left[\tfrac{1}{3}(u^2+1)^{3/2}\right]_0^1\ [-\cos v]_0^\pi = \tfrac{1}{3}(2^{3/2}-1)\cdot 2 = \tfrac{2}{3}(2\sqrt{2}-1)$$

8. $\mathbf{r}(u,v) = \langle 2uv, u^2-v^2, u^2+v^2\rangle$, $u^2+v^2 \le 1$ and

$\mathbf{r}_u \times \mathbf{r}_v = \langle 2v, 2u, 2u\rangle \times \langle 2u, -2v, 2v\rangle = \langle 8uv, 4u^2-4v^2, -4u^2-4v^2\rangle$, so

$$|\mathbf{r}_u \times \mathbf{r}_v| = \sqrt{(8uv)^2 + (4u^2-4v^2)^2 + (-4u^2-4v^2)^2} = \sqrt{64u^2v^2+32u^4+32v^4}$$
$$= \sqrt{32(u^2+v^2)^2} = 4\sqrt{2}\,(u^2+v^2)$$

Then

$$\iint_S (x^2+y^2)\,dS = \iint_D \left[(2uv)^2 + (u^2-v^2)^2\right]\ |\mathbf{r}_u \times \mathbf{r}_v|\ dA = \iint_D (4u^2v^2+u^4-2u^2v^2+v^4)\cdot 4\sqrt{2}\,(u^2+v^2)\,dA$$
$$= 4\sqrt{2}\iint_D (u^4+2u^2v^2+v^4)\,(u^2+v^2)\,dA = 4\sqrt{2}\iint_D (u^2+v^2)^3\,dA = 4\sqrt{2}\int_0^{2\pi}\int_0^1 (r^2)^3\,r\,dr\,d\theta$$
$$= 4\sqrt{2}\int_0^{2\pi} d\theta \int_0^1 r^7\,dr = 4\sqrt{2}\ [\theta]_0^{2\pi}\ \left[\tfrac{1}{8}r^8\right]_0^1 = 4\sqrt{2}\cdot 2\pi\cdot\tfrac{1}{8} = \sqrt{2}\,\pi$$

9. $z = 1+2x+3y$ so $\dfrac{\partial z}{\partial x} = 2$ and $\dfrac{\partial z}{\partial y} = 3$. Then by Formula 4,

$$\iint_S x^2yz\,dS = \iint_D x^2yz\sqrt{\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 + 1}\,dA = \int_0^3\int_0^2 x^2y(1+2x+3y)\sqrt{4+9+1}\,dy\,dx$$
$$= \sqrt{14}\int_0^3\int_0^2 (x^2y+2x^3y+3x^2y^2)\,dy\,dx = \sqrt{14}\int_0^3 \left[\tfrac{1}{2}x^2y^2 + x^3y^2 + x^2y^3\right]_{y=0}^{y=2} dx$$
$$= \sqrt{14}\int_0^3 (10x^2+4x^3)\,dx = \sqrt{14}\left[\tfrac{10}{3}x^3 + x^4\right]_0^3 = 171\sqrt{14}$$

10. S is the part of the plane $z = 4-2x-2y$ over the region $D = \{(x,y) \mid 0 \le x \le 2,\ 0 \le y \le 2-x\}$. Thus

$$\iint_S xz\,dS = \iint_D x(4-2x-2y)\sqrt{(-2)^2+(-2)^2+1}\,dA = 3\int_0^2\int_0^{2-x} \left(4x-2x^2-2xy\right)dy\,dx$$
$$= 3\int_0^2 \left[4xy-2x^2y-xy^2\right]_{y=0}^{y=2-x} dx = 3\int_0^2 \left[4x(2-x)-2x^2(2-x)-x(2-x)^2\right]dx$$
$$= 3\int_0^2 \left(x^3-4x^2+4x\right)dx = 3\left[\tfrac{1}{4}x^4 - \tfrac{4}{3}x^3 + 2x^2\right]_0^2 = 3\left(4-\tfrac{32}{3}+8\right) = 4$$

11. An equation of the plane through the points $(1,0,0)$, $(0,-2,0)$, and $(0,0,4)$ is $4x-2y+z=4$, so S is the region in the plane $z = 4-4x+2y$ over $D = \{(x,y) \mid 0 \le x \le 1,\ 2x-2 \le y \le 0\}$. Thus by Formula 4,

$$\iint_S x\,dS = \iint_D x\sqrt{(-4)^2+(2)^2+1}\,dA = \sqrt{21}\int_0^1\int_{2x-2}^0 x\,dy\,dx = \sqrt{21}\int_0^1 [xy]_{y=2x-2}^{y=0}\,dx$$
$$= \sqrt{21}\int_0^1 (-2x^2+2x)\,dx = \sqrt{21}\left[-\tfrac{2}{3}x^3+x^2\right]_0^1 = \sqrt{21}\left(-\tfrac{2}{3}+1\right) = \tfrac{\sqrt{21}}{3}$$

12. $z = \frac{2}{3}(x^{3/2}+y^{3/2})$ and

$$\iint_S y\,dS = \iint_D y\sqrt{\left(\sqrt{x}\right)^2 + \left(\sqrt{y}\right)^2 + 1}\,dA = \int_0^1\int_0^1 y\sqrt{x+y+1}\,dx\,dy$$
$$= \int_0^1 y\left[\tfrac{2}{3}(x+y+1)^{3/2}\right]_{x=0}^{x=1} dy = \int_0^1 \tfrac{2}{3}y\left[(y+2)^{3/2}-(y+1)^{3/2}\right]dy$$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Substituting $u = y + 2$ in the first term and $t = y + 1$ in the second, we have

$$\begin{aligned}\iint_S y\,dS &= \tfrac{2}{3}\int_2^3 (u-2)u^{3/2}\,du - \tfrac{2}{3}\int_1^2 (t-1)t^{3/2}\,dt = \tfrac{2}{3}\left[\tfrac{2}{7}u^{7/2} - \tfrac{4}{5}u^{5/2}\right]_2^3 - \tfrac{2}{3}\left[\tfrac{2}{7}t^{7/2} - \tfrac{2}{5}t^{5/2}\right]_1^2 \\ &= \tfrac{2}{3}\left[\tfrac{2}{7}(3^{7/2} - 2^{7/2}) - \tfrac{4}{5}(3^{5/2} - 2^{5/2}) - \tfrac{2}{7}(2^{7/2} - 1) + \tfrac{2}{5}(2^{5/2} - 1)\right] \\ &= \tfrac{2}{3}\left(\tfrac{18}{35}\sqrt{3} + \tfrac{8}{35}\sqrt{2} - \tfrac{4}{35}\right) = \tfrac{4}{105}\left(9\sqrt{3} + 4\sqrt{2} - 2\right)\end{aligned}$$

13. Using y and z as parameters, we have $\mathbf{r}(y, z) = (y^2 + z^2)\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$, $y^2 + z^2 \le 1$. Then

$\mathbf{r}_y \times \mathbf{r}_z = (2y\,\mathbf{i} + \mathbf{j}) \times (2z\,\mathbf{i} + \mathbf{k}) = \mathbf{i} - 2y\,\mathbf{j} - 2z\,\mathbf{k}$ and $|\mathbf{r}_y \times \mathbf{r}_z| = \sqrt{1 + 4y^2 + 4z^2} = \sqrt{1 + 4(y^2 + z^2)}$. Thus

$$\begin{aligned}\iint_S z^2\,dS &= \iint_{y^2+z^2\le 1} z^2\sqrt{1 + 4(y^2 + z^2)}\,dA = \int_0^{2\pi}\int_0^1 (r\sin\theta)^2\sqrt{1 + 4r^2}\,r\,dr\,d\theta \\ &= \int_0^{2\pi}\sin^2\theta\,d\theta\int_0^1 r^3\sqrt{1 + 4r^2}\,dr \qquad [\text{let } u = 1 + 4r^2 \;\Rightarrow\; r^2 = \tfrac{1}{4}(u-1) \text{ and } r\,dr = \tfrac{1}{8}du] \\ &= \left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\int_1^5 \tfrac{1}{4}(u-1)\sqrt{u}\cdot\tfrac{1}{8}du = \pi\cdot\tfrac{1}{32}\int_1^5 (u^{3/2} - u^{1/2})\,du = \tfrac{1}{32}\pi\left[\tfrac{2}{5}u^{5/2} - \tfrac{2}{3}u^{3/2}\right]_1^5 \\ &= \tfrac{1}{32}\pi\left[\tfrac{2}{5}(5)^{5/2} - \tfrac{2}{3}(5)^{3/2} - \tfrac{2}{5} + \tfrac{2}{3}\right] = \tfrac{1}{32}\pi\left(\tfrac{20}{3}\sqrt{5} + \tfrac{4}{15}\right) = \tfrac{1}{120}\pi\left(25\sqrt{5} + 1\right)\end{aligned}$$

14. Using x and z as parameters, we have $\mathbf{r}(x, z) = x\,\mathbf{i} + \sqrt{x^2 + z^2}\,\mathbf{j} + z\,\mathbf{k}$, $x^2 + z^2 \le 25$. Then

$$\mathbf{r}_x \times \mathbf{r}_z = \left(\mathbf{i} + \frac{x}{\sqrt{x^2+z^2}}\,\mathbf{j}\right) \times \left(\frac{z}{\sqrt{x^2+z^2}}\,\mathbf{j} + \mathbf{k}\right) = \frac{x}{\sqrt{x^2+z^2}}\,\mathbf{i} - \mathbf{j} + \frac{z}{\sqrt{x^2+z^2}}\,\mathbf{k} \text{ and}$$

$|\mathbf{r}_x \times \mathbf{r}_z| = \sqrt{\dfrac{x^2}{x^2+z^2} + 1 + \dfrac{z^2}{x^2+z^2}} = \sqrt{\dfrac{x^2+z^2}{x^2+z^2} + 1} = \sqrt{2}$. Thus

$$\begin{aligned}\iint_S y^2z^2\,dS &= \iint_{x^2+z^2\le 25} (x^2 + z^2)z^2\sqrt{2}\,dA = \sqrt{2}\int_0^{2\pi}\int_0^5 r^2(r\sin\theta)^2\,r\,dr\,d\theta \\ &= \sqrt{2}\int_0^{2\pi}\sin^2\theta\,d\theta\int_0^5 r^5\,dr = \sqrt{2}\left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{6}r^6\right]_0^5 \\ &= \sqrt{2}\,(\pi)\cdot\tfrac{1}{6}(15{,}625 - 0) = \frac{15{,}625\sqrt{2}}{6}\pi\end{aligned}$$

15. Using x and z as parameters, we have $\mathbf{r}(x, z) = x\,\mathbf{i} + (x^2 + 4z)\,\mathbf{j} + z\,\mathbf{k}$, $0 \le x \le 1, 0 \le z \le 1$. Then

$\mathbf{r}_x \times \mathbf{r}_z = (\mathbf{i} + 2x\,\mathbf{j}) \times (4\,\mathbf{j} + \mathbf{k}) = 2x\,\mathbf{i} - \mathbf{j} + 4\,\mathbf{k}$ and $|\mathbf{r}_x \times \mathbf{r}_z| = \sqrt{4x^2 + 1 + 16} = \sqrt{4x^2 + 17}$. Thus

$$\begin{aligned}\iint_S x\,dS &= \int_0^1\int_0^1 x\sqrt{4x^2 + 17}\,dz\,dx = \int_0^1 x\sqrt{4x^2 + 17}\,dx = \left[\tfrac{1}{8}\cdot\tfrac{2}{3}(4x^2 + 17)^{3/2}\right]_0^1 \\ &= \tfrac{1}{12}(21^{3/2} - 17^{3/2}) = \tfrac{1}{12}\left(21\sqrt{21} - 17\sqrt{17}\right) = \tfrac{7}{4}\sqrt{21} - \tfrac{17}{12}\sqrt{17}\end{aligned}$$

16. The sphere intersects the cone in the circle $x^2 + y^2 = \frac{1}{2}$, $z = \frac{1}{\sqrt{2}}$, so S is the portion of the sphere where $z \ge \frac{1}{\sqrt{2}}$.

Using spherical coordinates to parametrize the sphere we have $\mathbf{r}(\phi, \theta) = \sin\phi\cos\theta\,\mathbf{i} + \sin\phi\sin\theta\,\mathbf{j} + \cos\phi\,\mathbf{k}$, and $|\mathbf{r}_\phi \times \mathbf{r}_\theta| = \sin\phi$ (as in Example 1). The portion where $z \ge \frac{1}{\sqrt{2}}$ corresponds to $0 \le \phi \le \frac{\pi}{4}$, $0 \le \theta \le 2\pi$ so

$$\begin{aligned}\iint_S y^2\,dS &= \int_0^{2\pi}\int_0^{\pi/4}(\sin\phi\sin\theta)^2(\sin\phi)\,d\phi\,d\theta = \int_0^{2\pi}\sin^2\theta\,d\theta\int_0^{\pi/4}\sin^3\phi\,d\phi = \int_0^{2\pi}\sin^2\theta\,d\theta\int_0^{\pi/4}(1 - \cos^2\phi)\sin\phi\,d\phi \\ &= \left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{3}\cos^3\phi - \cos\phi\right]_0^{\pi/4} = \pi\left(\tfrac{\sqrt{2}}{12} - \tfrac{\sqrt{2}}{2} - \tfrac{1}{3} + 1\right) = \left(\tfrac{2}{3} - \tfrac{5\sqrt{2}}{12}\right)\pi\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. Using spherical coordinates to parametrize the sphere we have $\mathbf{r}(\phi,\theta) = 2\sin\phi\cos\theta\,\mathbf{i} + 2\sin\phi\sin\theta\,\mathbf{j} + 2\cos\phi\,\mathbf{k}$ and $|\mathbf{r}_\phi \times \mathbf{r}_\theta| = 4\sin\phi$ (see Example 16.6.10). Here S is the portion of the sphere corresponding to $0 \le \phi \le \pi/2$, so

$$\begin{aligned}\iint_S (x^2z + y^2z)\,dS &= \iint_S (x^2+y^2)z\,dS = \int_0^{2\pi}\int_0^{\pi/2} (4\sin^2\phi)(2\cos\phi)(4\sin\phi)\,d\phi\,d\theta \\ &= 32\int_0^{2\pi} d\theta \int_0^{\pi/2} \sin^3\phi\cos\phi\,d\phi = 32\,(2\pi)\left[\tfrac{1}{4}\sin^4\phi\right]_0^{\pi/2} = 16\pi(1-0) = 16\pi\end{aligned}$$

18. S is given by $\mathbf{r}(u,v) = \cos v\,\mathbf{i} + u\,\mathbf{j} + \sin v\,\mathbf{k}$, $0 \le u \le 2$, $0 \le v \le \pi$. Then

$\mathbf{r}_u \times \mathbf{r}_v = \mathbf{j} \times (-\sin v\,\mathbf{i} + \cos v\,\mathbf{k}) = \cos v\,\mathbf{i} + \sin v\,\mathbf{k}$ and $|\mathbf{r}_u \times \mathbf{r}_v| = \sqrt{\cos^2 v + \sin^2 v} = 1$, so

$$\begin{aligned}\iint_S (x+y+z)\,dS &= \int_0^{\pi}\int_0^{2} (\cos v + u + \sin v)(1)\,du\,dv = \int_0^{\pi}\left[u(\cos v + \sin v) + \tfrac{1}{2}u^2\right]_{u=0}^{u=2} dv \\ &= \int_0^{\pi} (2\cos v + 2\sin v + 2)\,dv = [2\sin v - 2\cos v + 2v]_0^{\pi} = 2 + 2\pi + 2 = 4 + 2\pi\end{aligned}$$

19. Here S consists of three surfaces: S_1, the lateral surface of the cylinder; S_2, the front formed by the plane $x + y = 5$; and the back, S_3, in the plane $x = 0$.

On S_1: the surface is given by $\mathbf{r}(u,v) = u\,\mathbf{i} + 3\cos v\,\mathbf{j} + 3\sin v\,\mathbf{k}$, $0 \le v \le 2\pi$, and $0 \le x \le 5 - y \;\Rightarrow$ $0 \le u \le 5 - 3\cos v$. Then $\mathbf{r}_u \times \mathbf{r}_v = -3\cos v\,\mathbf{j} - 3\sin v\,\mathbf{k}$ and $|\mathbf{r}_u \times \mathbf{r}_v| = \sqrt{9\cos^2 v + 9\sin^2 v} = 3$, so

$$\begin{aligned}\iint_{S_1} xz\,dS &= \int_0^{2\pi}\int_0^{5-3\cos v} u(3\sin v)(3)\,du\,dv = 9\int_0^{2\pi}\left[\tfrac{1}{2}u^2\right]_{u=0}^{u=5-3\cos v}\sin v\,dv \\ &= \tfrac{9}{2}\int_0^{2\pi}(5-3\cos v)^2\sin v\,dv = \tfrac{9}{2}\left[\tfrac{1}{9}(5-3\cos v)^3\right]_0^{2\pi} = 0.\end{aligned}$$

On S_2: $\mathbf{r}(y,z) = (5-y)\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$ and $|\mathbf{r}_y \times \mathbf{r}_z| = |\mathbf{i} + \mathbf{j}| = \sqrt{2}$, where $y^2 + z^2 \le 9$ and

$$\begin{aligned}\iint_{S_2} xz\,dS &= \iint_{y^2+z^2\le 9} (5-y)z\sqrt{2}\,dA = \sqrt{2}\int_0^{2\pi}\int_0^3 (5 - r\cos\theta)(r\sin\theta)\,r\,dr\,d\theta \\ &= \sqrt{2}\int_0^{2\pi}\int_0^3 (5r^2 - r^3\cos\theta)(\sin\theta)\,dr\,d\theta = \sqrt{2}\int_0^{2\pi}\left[\tfrac{5}{3}r^3 - \tfrac{1}{4}r^4\cos\theta\right]_{r=0}^{r=3}\sin\theta\,d\theta \\ &= \sqrt{2}\int_0^{2\pi}\left(45 - \tfrac{81}{4}\cos\theta\right)\sin\theta\,d\theta = \sqrt{2}\left(\tfrac{4}{81}\right)\cdot\tfrac{1}{2}\left(45 - \tfrac{81}{4}\cos\theta\right)^2\Big]_0^{2\pi} = 0\end{aligned}$$

On S_3: $x = 0$ so $\iint_{S_3} xz\,dS = 0$. Hence $\iint_S xz\,dS = 0 + 0 + 0 = 0$.

20. Let S_1 be the lateral surface, S_2 the top disk, and S_3 the bottom disk.

On S_1: $\mathbf{r}(\theta,z) = 3\cos\theta\,\mathbf{i} + 3\sin\theta\,\mathbf{j} + z\,\mathbf{k}$, $0 \le \theta \le 2\pi$, $0 \le z \le 2$, $|\mathbf{r}_\theta \times \mathbf{r}_z| = 3$,

$\iint_{S_1}(x^2+y^2+z^2)\,dS = \int_0^{2\pi}\int_0^2 (9+z^2)\,3\,dz\,d\theta = 2\pi(54+8) = 124\pi$.

On S_2: $\mathbf{r}(\theta,r) = r\cos\theta\,\mathbf{i} + r\sin\theta\,\mathbf{j} + 2\,\mathbf{k}$, $0 \le r \le 3$, $0 \le \theta \le 2\pi$, $|\mathbf{r}_\theta \times \mathbf{r}_r| = r$,

$\iint_{S_2}(x^2+y^2+z^2)\,dS = \int_0^{2\pi}\int_0^3 (r^2+4)\,r\,dr\,d\theta = 2\pi\left(\frac{81}{4} + 18\right) = \frac{153}{2}\pi$.

On S_3: $\mathbf{r}(\theta,r) = r\cos\theta\,\mathbf{i} + r\sin\theta\,\mathbf{j}$, $0 \le r \le 3$, $0 \le \theta \le 2\pi$, $|\mathbf{r}_\theta \times \mathbf{r}_r| = r$,

$\iint_{S_3}(x^2+y^2+z^2)\,dS = \int_0^{2\pi}\int_0^3 (r^2+0)\,r\,dr\,d\theta = 2\pi\left(\frac{81}{4}\right) = \frac{81}{2}\pi$.

Hence $\iint_S (x^2+y^2+z^2)\,dS = 124\pi + \frac{153}{2}\pi + \frac{81}{2}\pi = 241\pi$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. From Exercise 5, $\mathbf{r}(u,v) = (u+v)\,\mathbf{i} + (u-v)\,\mathbf{j} + (1+2u+v)\,\mathbf{k}$, $0 \le u \le 2$, $0 \le v \le 1$, and $\mathbf{r}_u \times \mathbf{r}_v = 3\,\mathbf{i} + \mathbf{j} - 2\,\mathbf{k}$.

Then

$$\begin{aligned}\mathbf{F}(\mathbf{r}(u,v)) &= (1+2u+v)e^{(u+v)(u-v)}\,\mathbf{i} - 3(1+2u+v)e^{(u+v)(u-v)}\,\mathbf{j} + (u+v)(u-v)\,\mathbf{k} \\ &= (1+2u+v)e^{u^2-v^2}\,\mathbf{i} - 3(1+2u+v)e^{u^2-v^2}\,\mathbf{j} + (u^2-v^2)\,\mathbf{k}\end{aligned}$$

Because the z-component of $\mathbf{r}_u \times \mathbf{r}_v$ is negative we use $-(\mathbf{r}_u \times \mathbf{r}_v)$ in Formula 9 for the upward orientation:

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iint_D \mathbf{F}\cdot(-(\mathbf{r}_u\times\mathbf{r}_v))\,dA = \int_0^1\int_0^2 \left[-3(1+2u+v)e^{u^2-v^2} + 3(1+2u+v)e^{u^2-v^2} + 2(u^2-v^2)\right] du\,dv \\ &= \int_0^1\int_0^2 2(u^2-v^2)\,du\,dv = 2\int_0^1 \left[\tfrac{1}{3}u^3 - uv^2\right]_{u=0}^{u=2} dv = 2\int_0^1 \left(\tfrac{8}{3} - 2v^2\right) dv \\ &= 2\left[\tfrac{8}{3}v - \tfrac{2}{3}v^3\right]_0^1 = 2\left(\tfrac{8}{3} - \tfrac{2}{3}\right) = 4\end{aligned}$$

22. $\mathbf{r}(u,v) = \langle u\cos v, u\sin v, v\rangle$, $0 \le u \le 1$, $0 \le v \le \pi$ and

$\mathbf{r}_u \times \mathbf{r}_v = \langle \cos v, \sin v, 0\rangle \times \langle -u\sin v, u\cos v, 1\rangle = \langle \sin v, -\cos v, u\rangle$. Here $\mathbf{F}(\mathbf{r}(u,v)) = v\,\mathbf{i} + u\sin v\,\mathbf{j} + u\cos v\,\mathbf{k}$ and, by Formula 9,

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iint_D \mathbf{F}\cdot(\mathbf{r}_u\times\mathbf{r}_v)\,dA = \int_0^1\int_0^\pi (v\sin v - u\sin v\cos v + u^2\cos v)\,dv\,du \\ &= \int_0^1 \left[\sin v - v\cos v - \tfrac{1}{2}u\sin^2 v + u^2\sin v\right]_{v=0}^{v=\pi} du = \int_0^1 \pi\,du = \pi u\Big]_0^1 = \pi\end{aligned}$$

23. $\mathbf{F}(x,y,z) = xy\,\mathbf{i} + yz\,\mathbf{j} + zx\,\mathbf{k}$, $z = g(x,y) = 4 - x^2 - y^2$, and D is the square $[0,1]\times[0,1]$, so by Equation 10

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iint_D [-xy(-2x) - yz(-2y) + zx]\,dA = \int_0^1\int_0^1 [2x^2y + 2y^2(4-x^2-y^2) + x(4-x^2-y^2)]\,dy\,dx \\ &= \int_0^1 \left[x^2y^2 + \tfrac{8}{3}y^3 - \tfrac{2}{3}x^2y^3 - \tfrac{2}{5}y^5 + 4xy - x^3y - \tfrac{1}{3}xy^3\right]_{y=0}^{y=1} dx \\ &= \int_0^1 \left(\tfrac{1}{3}x^2 + \tfrac{11}{3}x - x^3 + \tfrac{34}{15}\right) dx = \left[\tfrac{1}{9}x^3 + \tfrac{11}{6}x^2 - \tfrac{1}{4}x^4 + \tfrac{34}{15}x\right]_0^1 = \tfrac{713}{180}\end{aligned}$$

24. $\mathbf{F}(x,y,z) = -x\,\mathbf{i} - y\,\mathbf{j} + z^3\,\mathbf{k}$, $z = g(x,y) = \sqrt{x^2+y^2}$, and D is the annular region $\{(x,y) \mid 1 \le x^2+y^2 \le 9\}$. Since S has downward orientation, we have

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= -\iint_D \left[-(-x)\left(\frac{x}{\sqrt{x^2+y^2}}\right) - (-y)\left(\frac{y}{\sqrt{x^2+y^2}}\right) + z^3\right] dA \\ &= \iint_D \left[\frac{x^2+y^2}{\sqrt{x^2+y^2}} + \left(\sqrt{x^2+y^2}\right)^3\right] dA = \int_0^{2\pi}\int_1^3 \left(\frac{r^2}{r} + r^3\right) r\,dr\,d\theta \\ &= \int_0^{2\pi} d\theta \int_1^3 (r^2+r^4)\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{1}{3}r^3 + \tfrac{1}{5}r^5\right]_1^3 \\ &= 2\pi\left(9 + \tfrac{243}{5} - \tfrac{1}{3} - \tfrac{1}{5}\right) = \tfrac{1712}{15}\pi\end{aligned}$$

25. $\mathbf{F}(x,y,z) = x\,\mathbf{i} + y\,\mathbf{j} + z^2\,\mathbf{k}$, and using spherical coordinates, S is given by $x = \sin\phi\cos\theta$, $y = \sin\phi\sin\theta$, $z = \cos\phi$, $0 \le \theta \le 2\pi$, $0 \le \phi \le \pi$. $\mathbf{F}(\mathbf{r}(\phi,\theta)) = (\sin\phi\cos\theta)\,\mathbf{i} + (\sin\phi\sin\theta)\,\mathbf{j} + (\cos^2\phi)\,\mathbf{k}$ and, from Example 4, $\mathbf{r}_\phi \times \mathbf{r}_\theta = \sin^2\phi\cos\theta\,\mathbf{i} + \sin^2\phi\sin\theta\,\mathbf{j} + \sin\phi\cos\phi\,\mathbf{k}$. Thus

$$\mathbf{F}(\mathbf{r}(\phi,\theta))\cdot(\mathbf{r}_\phi\times\mathbf{r}_\theta) = \sin^3\phi\cos^2\theta + \sin^3\phi\sin^2\theta + \sin\phi\cos^3\phi = \sin^3\phi + \sin\phi\cos^3\phi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iint_D [\mathbf{F}(\mathbf{r}(\phi,\theta))\cdot(\mathbf{r}_\phi\times\mathbf{r}_\theta)]\,dA = \int_0^{2\pi}\int_0^{\pi}(\sin^3\phi+\sin\phi\cos^3\phi)\,d\phi\,d\theta\\ &= \int_0^{2\pi} d\theta\int_0^{\pi}(1-\cos^2\phi+\cos^3\phi)\sin\phi\,d\phi = (2\pi)\left[-\cos\phi+\tfrac{1}{3}\cos^3\phi-\tfrac{1}{4}\cos^4\phi\right]_0^{\pi}\\ &= 2\pi\left(1-\tfrac{1}{3}-\tfrac{1}{4}+1-\tfrac{1}{3}+\tfrac{1}{4}\right) = \tfrac{8}{3}\pi\end{aligned}$$

26. $\mathbf{F}(x,y,z) = y\,\mathbf{i} - x\,\mathbf{j} + 2z\,\mathbf{k}$, $z = g(x,y) = \sqrt{4-x^2-y^2}$ and D is the disk $\{(x,y) \mid x^2+y^2 \le 4\}$. S has downward orientation, so by Equation 10,

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= -\iint_D[-y\cdot\tfrac{1}{2}(4-x^2-y^2)^{-1/2}(-2x)-(-x)\cdot\tfrac{1}{2}(4-x^2-y^2)^{-1/2}(-2y)+2z]\,dA\\ &= -\iint_D\left(\frac{xy}{\sqrt{4-x^2-y^2}}-\frac{xy}{\sqrt{4-x^2-y^2}}+2\sqrt{4-x^2-y^2}\right)dA\\ &= -\iint_D 2\sqrt{4-x^2-y^2}\,dA = -2\int_0^{2\pi}\int_0^2\sqrt{4-r^2}\,r\,dr\,d\theta = -2\int_0^{2\pi}d\theta\int_0^2 r\sqrt{4-r^2}\,dr\\ &= -2(2\pi)\left[-\tfrac{1}{2}\cdot\tfrac{2}{3}(4-r^2)^{3/2}\right]_0^2 = -4\pi\left[0+\tfrac{1}{3}(4)^{3/2}\right] = -4\pi\cdot\tfrac{8}{3} = -\tfrac{32}{3}\pi\end{aligned}$$

27. Let S_1 be the paraboloid $y = x^2+z^2$, $0 \le y \le 1$ and S_2 the disk $x^2+z^2 \le 1$, $y = 1$. Since S is a closed surface, we use the outward orientation.

On S_1: $\mathbf{F}(\mathbf{r}(x,z)) = (x^2+z^2)\,\mathbf{j} - z\,\mathbf{k}$ and $\mathbf{r}_x\times\mathbf{r}_z = 2x\,\mathbf{i} - \mathbf{j} + 2z\,\mathbf{k}$ (since the $\mathbf{j}$-component must be negative on S_1). Then

$$\begin{aligned}\iint_{S_1}\mathbf{F}\cdot d\mathbf{S} &= \iint_{x^2+z^2\le 1}[-(x^2+z^2)-2z^2]\,dA = -\int_0^{2\pi}\int_0^1(r^2+2r^2\sin^2\theta)\,r\,dr\,d\theta\\ &= -\int_0^{2\pi}\int_0^1 r^3(1+2\sin^2\theta)\,dr\,d\theta = -\int_0^{2\pi}(1+1-\cos 2\theta)\,d\theta\int_0^1 r^3\,dr\\ &= -\left[2\theta-\tfrac{1}{2}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^1 = -4\pi\cdot\tfrac{1}{4} = -\pi\end{aligned}$$

On S_2: $\mathbf{F}(\mathbf{r}(x,z)) = \mathbf{j} - z\,\mathbf{k}$ and $\mathbf{r}_z\times\mathbf{r}_x = \mathbf{j}$. Then $\iint_{S_2}\mathbf{F}\cdot d\mathbf{S} = \iint_{x^2+z^2\le 1}(1)\,dA = \pi$.

Hence $\iint_S \mathbf{F}\cdot d\mathbf{S} = -\pi+\pi = 0$.

28. $\mathbf{F}(x,y,z) = yz\,\mathbf{i} + zx\,\mathbf{j} + xy\,\mathbf{k}$, $z = g(x,y) = x\sin y$, and D is the rectangle $[0,2]\times[0,\pi]$, so by Equation 10

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iint_D[-yz(\sin y)-zx(x\cos y)+xy]\,dA = \int_0^{\pi}\int_0^2(-xy\sin^2 y - x^3\sin y\cos y + xy)\,dx\,dy\\ &= \int_0^{\pi}\left[-\tfrac{1}{2}x^2y\sin^2 y - \tfrac{1}{4}x^4\sin y\cos y + \tfrac{1}{2}x^2y\right]_{x=0}^{x=2}dy\\ &= \int_0^{\pi}\left(-2y\sin^2 y - 4\sin y\cos y + 2y\right)dy \qquad \text{[integrate by parts in the first term]}\\ &= \left[\left(-\tfrac{1}{2}y^2+\tfrac{1}{2}y\sin 2y+\tfrac{1}{4}\cos 2y\right)-2\sin^2 y+y^2\right]_0^{\pi} = -\tfrac{1}{2}\pi^2+\tfrac{1}{4}+\pi^2-\tfrac{1}{4} = \tfrac{1}{2}\pi^2\end{aligned}$$

29. Here S consists of the six faces of the cube as labeled in the figure. On S_1:



$\mathbf{F} = \mathbf{i} + 2y\,\mathbf{j} + 3z\,\mathbf{k}$, $\mathbf{r}_y\times\mathbf{r}_z = \mathbf{i}$ and $\iint_{S_1}\mathbf{F}\cdot d\mathbf{S} = \int_{-1}^1\int_{-1}^1 dy\,dz = 4$;

S_2: $\mathbf{F} = x\,\mathbf{i} + 2\,\mathbf{j} + 3z\,\mathbf{k}$, $\mathbf{r}_z\times\mathbf{r}_x = \mathbf{j}$ and $\iint_{S_2}\mathbf{F}\cdot d\mathbf{S} = \int_{-1}^1\int_{-1}^1 2\,dx\,dz = 8$;

S_3: $\mathbf{F} = x\,\mathbf{i} + 2y\,\mathbf{j} + 3\,\mathbf{k}$, $\mathbf{r}_x\times\mathbf{r}_y = \mathbf{k}$ and $\iint_{S_3}\mathbf{F}\cdot d\mathbf{S} = \int_{-1}^1\int_{-1}^1 3\,dx\,dy = 12$;

S_4: $\mathbf{F} = -\mathbf{i} + 2y\,\mathbf{j} + 3z\,\mathbf{k}$, $\mathbf{r}_z\times\mathbf{r}_y = -\mathbf{i}$ and $\iint_{S_4}\mathbf{F}\cdot d\mathbf{S} = 4$;

S_5: $\mathbf{F} = x\,\mathbf{i} - 2\,\mathbf{j} + 3z\,\mathbf{k}$, $\mathbf{r}_x\times\mathbf{r}_z = -\mathbf{j}$ and $\iint_{S_5}\mathbf{F}\cdot d\mathbf{S} = 8$;

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

S_6: $\mathbf{F} = x\,\mathbf{i} + 2y\,\mathbf{j} - 3\,\mathbf{k}$, $\mathbf{r}_y \times \mathbf{r}_x = -\mathbf{k}$ and $\iint_{S_6} \mathbf{F} \cdot d\mathbf{S} = \int_{-1}^{1}\int_{-1}^{1} 3\,dx\,dy = 12$.

Hence $\iint_S \mathbf{F} \cdot d\mathbf{S} = \sum_{i=1}^{6} \iint_{S_i} \mathbf{F} \cdot d\mathbf{S} = 48$.

30. Here S consists of three surfaces: S_1, the lateral surface of the cylinder; S_2, the front formed by the plane $x + y = 2$; and the back, S_3, in the plane $y = 0$.

On S_1: $\mathbf{F}(\mathbf{r}(\theta, y)) = \sin\theta\,\mathbf{i} + y\,\mathbf{j} + 5\,\mathbf{k}$ and $\mathbf{r}_\theta \times \mathbf{r}_y = \sin\theta\,\mathbf{i} + \cos\theta\,\mathbf{k}$ $\Rightarrow$

$$\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \int_0^{2\pi}\int_0^{2-\sin\theta} (\sin^2\theta + 5\cos\theta)\,dy\,d\theta$$

$$= \int_0^{2\pi} (2\sin^2\theta + 10\cos\theta - \sin^3\theta - 5\sin\theta\,\cos\theta)\,d\theta = 2\pi$$

On S_2: $\mathbf{F}(\mathbf{r}(x, z)) = x\,\mathbf{i} + (2 - x)\,\mathbf{j} + 5\,\mathbf{k}$ and $\mathbf{r}_z \times \mathbf{r}_x = \mathbf{i} + \mathbf{j}$.

$$\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \iint_{x^2 + z^2 < 1} [x + (2 - x)]\,dA = 2\pi$$

On S_3: $\mathbf{F}(\mathbf{r}(x, z)) = x\,\mathbf{i} + 5\,\mathbf{k}$ and $\mathbf{r}_x \times \mathbf{r}_z = -\mathbf{j}$ so $\iint_{S_3} \mathbf{F} \cdot d\mathbf{S} = 0$. Hence $\iint_S \mathbf{F} \cdot d\mathbf{S} = 4\pi$.

31. Here S consists of four surfaces: S_1, the top surface (a portion of the circular cylinder $y^2 + z^2 = 1$); S_2, the bottom surface (a portion of the xy-plane); S_3, the front half-disk in the plane $x = 2$, and S_4, the back half-disk in the plane $x = 0$.

On S_1: The surface is $z = \sqrt{1 - y^2}$ for $0 \le x \le 2$, $-1 \le y \le 1$ with upward orientation, so

$$\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \int_0^2 \int_{-1}^{1} \left[-x^2\,(0) - y^2\left(-\frac{y}{\sqrt{1 - y^2}}\right) + z^2\right] dy\,dx = \int_0^2 \int_{-1}^{1} \left(\frac{y^3}{\sqrt{1 - y^2}} + 1 - y^2\right) dy\,dx$$

$$= \int_0^2 \left[-\sqrt{1 - y^2} + \tfrac{1}{3}(1 - y^2)^{3/2} + y - \tfrac{1}{3}y^3\right]_{y=-1}^{y=1} dx = \int_0^2 \tfrac{4}{3}\,dx = \tfrac{8}{3}$$

On S_2: The surface is $z = 0$ with downward orientation, so

$$\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \int_0^2 \int_{-1}^{1} (-z^2)\,dy\,dx = \int_0^2 \int_{-1}^{1} (0)\,dy\,dx = 0$$

On S_3: The surface is $x = 2$ for $-1 \le y \le 1$, $0 \le z \le \sqrt{1 - y^2}$, oriented in the positive x-direction. Regarding y and z as parameters, we have $\mathbf{r}_y \times \mathbf{r}_z = \mathbf{i}$ and

$$\iint_{S_3} \mathbf{F} \cdot d\mathbf{S} = \int_{-1}^{1} \int_0^{\sqrt{1-y^2}} x^2\,dz\,dy = \int_{-1}^{1} \int_0^{\sqrt{1-y^2}} 4\,dz\,dy = 4A\,(S_3) = 2\pi$$

On S_4: The surface is $x = 0$ for $-1 \le y \le 1$, $0 \le z \le \sqrt{1 - y^2}$, oriented in the negative x-direction. Regarding y and z as parameters, we use $-(\mathbf{r}_y \times \mathbf{r}_z) = -\mathbf{i}$ and

$$\iint_{S_4} \mathbf{F} \cdot d\mathbf{S} = \int_{-1}^{1} \int_0^{\sqrt{1-y^2}} x^2\,dz\,dy = \int_{-1}^{1} \int_0^{\sqrt{1-y^2}} (0)\,dz\,dy = 0$$

Thus $\iint_S \mathbf{F} \cdot d\mathbf{S} = \frac{8}{3} + 0 + 2\pi + 0 = 2\pi + \frac{8}{3}$.

32. Here S consists of four surfaces: S_1, the triangular face with vertices $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$; S_2, the face of the tetrahedron in the xy-plane; S_3, the face in the xz-plane; and S_4, the face in the yz-plane.

On S_1: The face is the portion of the plane $z = 1 - x - y$ for $0 \le x \le 1$, $0 \le y \le 1 - x$ with upward orientation, so

$$\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \int_0^1 \int_0^{1-x} [-y\,(-1) - (z - y)\,(-1) + x]\,dy\,dx = \int_0^1 \int_0^{1-x} (z + x)\,dy\,dx = \int_0^1 \int_0^{1-x} (1 - y)\,dy\,dx$$

$$= \int_0^1 \left[y - \tfrac{1}{2}y^2\right]_{y=0}^{y=1-x} dx = \tfrac{1}{2}\int_0^1 (1 - x^2)\,dx = \tfrac{1}{2}\left[x - \tfrac{1}{3}x^3\right]_0^1 = \tfrac{1}{3}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

On S_2: The surface is $z = 0$ with downward orientation, so

$$\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \int_0^1 \int_0^{1-x} (-x)\, dy\, dx = -\int_0^1 x(1-x)\, dx = -\left[\tfrac{1}{2}x^2 - \tfrac{1}{3}x^3\right]_0^1 = -\tfrac{1}{6}$$

On S_3: The surface is $y = 0$ for $0 \le x \le 1, 0 \le z \le 1 - x$, oriented in the negative y-direction. Regarding x and z as parameters, we have $\mathbf{r}_x \times \mathbf{r}_z = -\mathbf{j}$ and

$$\iint_{S_3} \mathbf{F} \cdot d\mathbf{S} = \int_0^1 \int_0^{1-x} -(z-y)\, dz\, dx = -\int_0^1 \int_0^{1-x} z\, dz\, dx = -\int_0^1 \left[\tfrac{1}{2}z^2\right]_{z=0}^{z=1-x} dx$$

$$= -\tfrac{1}{2}\int_0^1 (1-x)^2\, dx = \tfrac{1}{6}\left[(1-x)^3\right]_0^1 = -\tfrac{1}{6}$$

On S_4: The surface is $x = 0$ for $0 \le y \le 1, 0 \le z \le 1 - y$, oriented in the negative x-direction. Regarding y and z as parameters, we have $\mathbf{r}_y \times \mathbf{r}_z = \mathbf{i}$ so we use $-(\mathbf{r}_y \times \mathbf{r}_z) = -\mathbf{i}$ and

$$\iint_{S_4} \mathbf{F} \cdot d\mathbf{S} = \int_0^1 \int_0^{1-y} (-y)\, dz\, dy = -\int_0^1 y(1-y)\, dy = -\left[\tfrac{1}{2}y^2 - \tfrac{1}{3}y^3\right]_0^1 = -\tfrac{1}{6}$$

Thus $\iint_S \mathbf{F} \cdot d\mathbf{S} = \frac{1}{3} - \frac{1}{6} - \frac{1}{6} - \frac{1}{6} = -\frac{1}{6}$.

33. $z = xe^y \;\Rightarrow\; \partial z/\partial x = e^y$, $\partial z/\partial y = xe^y$, so by Formula 4, a CAS gives

$$\iint_S (x^2 + y^2 + z^2)\, dS = \int_0^1 \int_0^1 (x^2 + y^2 + x^2e^{2y})\sqrt{e^{2y} + x^2e^{2y} + 1}\, dx\, dy \approx 4.5822.$$

34. $z = x^2y^2 \;\Rightarrow\; \partial z/\partial x = 2xy^2$, $\partial z/\partial y = 2x^2y$, so by Formula 4, a CAS gives

$$\iint_S xyz\, dS = \int_0^2 \int_0^1 xy(x^2y^2)\sqrt{(2xy^2)^2 + (2x^2y)^2 + 1}\, dx\, dy$$

$$= \int_0^2 \int_0^1 x^3y^3\sqrt{4x^2y^4 + 4x^4y^2 + 1}\, dx\, dy = -\tfrac{151}{33} - \tfrac{1}{220}\sqrt{3}\,\pi + \tfrac{1977}{176}\ln 7 - \tfrac{9891}{880}\ln 3 + \tfrac{3}{440}\sqrt{3}\tan^{-1}\tfrac{5}{\sqrt{3}}$$

35. We use Formula 4 with $z = 3 - 2x^2 - y^2 \;\Rightarrow\; \partial z/\partial x = -4x$, $\partial z/\partial y = -2y$. The boundaries of the region $3 - 2x^2 - y^2 \ge 0$ are $-\sqrt{\frac{3}{2}} \le x \le \sqrt{\frac{3}{2}}$ and $-\sqrt{3 - 2x^2} \le y \le \sqrt{3 - 2x^2}$, so we use a CAS (with precision reduced to seven or fewer digits; otherwise the calculation may take a long time) to calculate

$$\iint_S x^2y^2z^2\, dS = \int_{-\sqrt{3/2}}^{\sqrt{3/2}} \int_{-\sqrt{3-2x^2}}^{\sqrt{3-2x^2}} x^2y^2(3 - 2x^2 - y^2)^2\sqrt{16x^2 + 4y^2 + 1}\, dy\, dx \approx 3.4895$$

36. The flux of $\mathbf{F}$ across S is given by $\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot \mathbf{n}\, dS$. Now on S, $z = g(x, y) = 2\sqrt{1 - y^2}$, so $\partial g/\partial x = 0$ and $\partial g/\partial y = -2y(1 - y^2)^{-1/2}$. Therefore, by (10),

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \int_{-2}^2 \int_{-1}^1 \left(-x^2y\left[-2y(1-y^2)^{-1/2}\right] + \left[2\sqrt{1-y^2}\right]^2 e^{x/5}\right) dy\, dx = \tfrac{1}{3}(16\pi + 80e^{2/5} - 80e^{-2/5})$$





© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

37. If S is given by $y = h(x, z)$, then S is also the level surface $f(x, y, z) = y - h(x, z) = 0$.

$\mathbf{n} = \dfrac{\nabla f(x,y,z)}{|\nabla f(x,y,z)|} = \dfrac{-h_x\,\mathbf{i} + \mathbf{j} - h_z\,\mathbf{k}}{\sqrt{h_x^2 + 1 + h_z^2}}$, and $-\mathbf{n}$ is the unit normal that points to the left. Now we proceed as in the derivation of (10), using Formula 4 to evaluate

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_S \mathbf{F}\cdot\mathbf{n}\,dS = \iint_D (P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k})\,\frac{\dfrac{\partial h}{\partial x}\,\mathbf{i} - \mathbf{j} + \dfrac{\partial h}{\partial z}\,\mathbf{k}}{\sqrt{\left(\dfrac{\partial h}{\partial x}\right)^2 + 1 + \left(\dfrac{\partial h}{\partial z}\right)^2}}\sqrt{\left(\frac{\partial h}{\partial x}\right)^2 + 1 + \left(\frac{\partial h}{\partial z}\right)^2}\,dA$$

where D is the projection of S onto the xz-plane. Therefore $\displaystyle\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_D \left(P\,\frac{\partial h}{\partial x} - Q + R\,\frac{\partial h}{\partial z}\right) dA$.

38. If S is given by $x = k(y, z)$, then S is also the level surface $f(x, y, z) = x - k(y, z) = 0$.

$\mathbf{n} = \dfrac{\nabla f(x,y,z)}{|\nabla f(x,y,z)|} = \dfrac{\mathbf{i} - k_y\,\mathbf{j} - k_z\,\mathbf{k}}{\sqrt{1 + k_y^2 + k_z^2}}$, and since the x-component is positive this is the unit normal that points forward.

Now we proceed as in the derivation of (10), using Formula 4 for

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_S \mathbf{F}\cdot\mathbf{n}\,dS = \iint_D (P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k})\,\frac{\mathbf{i} - \dfrac{\partial k}{\partial y}\,\mathbf{j} - \dfrac{\partial k}{\partial z}\,\mathbf{k}}{\sqrt{1 + \left(\dfrac{\partial k}{\partial y}\right)^2 + \left(\dfrac{\partial k}{\partial z}\right)^2}}\sqrt{1 + \left(\frac{\partial k}{\partial y}\right)^2 + \left(\frac{\partial k}{\partial z}\right)^2}\,dA$$

where D is the projection of S onto the yz-plane. Therefore $\displaystyle\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_D \left(P - Q\,\frac{\partial k}{\partial y} - R\,\frac{\partial k}{\partial z}\right) dA$.

39. $m = \iint_S K\,dS = K\cdot 4\pi\left(\frac{1}{2}a^2\right) = 2\pi a^2 K$; by symmetry $M_{xz} = M_{yz} = 0$, and

$M_{xy} = \iint_S zK\,dS = K\int_0^{2\pi}\int_0^{\pi/2}(a\cos\phi)(a^2\sin\phi)\,d\phi\,d\theta = 2\pi K a^3\left[-\frac{1}{4}\cos 2\phi\right]_0^{\pi/2} = \pi K a^3$.

Hence $(\overline{x}, \overline{y}, \overline{z}) = \left(0, 0, \frac{1}{2}a\right)$.

40. S is given by $\mathbf{r}(x, y) = x\,\mathbf{i} + y\,\mathbf{j} + \sqrt{x^2 + y^2}\,\mathbf{k}$, $|\mathbf{r}_x \times \mathbf{r}_y| = \sqrt{1 + \dfrac{x^2 + y^2}{x^2 + y^2}} = \sqrt{2}$ so

$$m = \iint_S \left(10 - \sqrt{x^2 + y^2}\right) dS = \iint_{1\le x^2+y^2\le 16} \left(10 - \sqrt{x^2 + y^2}\right)\sqrt{2}\,dA$$

$$= \int_0^{2\pi}\int_1^4 \sqrt{2}\,(10 - r)\,r\,dr\,d\theta = 2\pi\sqrt{2}\left[5r^2 - \tfrac{1}{3}r^3\right]_1^4 = 108\sqrt{2}\,\pi$$

41. (a) $I_z = \iint_S (x^2 + y^2)\rho(x, y, z)\,dS$

(b) $I_z = \iint_S (x^2 + y^2)\left(10 - \sqrt{x^2 + y^2}\right) dS = \iint_{1\le x^2+y^2\le 16} (x^2 + y^2)\left(10 - \sqrt{x^2 + y^2}\right)\sqrt{2}\,dA$

$= \int_0^{2\pi}\int_1^4 \sqrt{2}\,(10r^3 - r^4)\,dr\,d\theta = 2\sqrt{2}\,\pi\left(\frac{4329}{10}\right) = \frac{4329}{5}\sqrt{2}\,\pi$

42. Using spherical coordinates to parametrize the sphere we have $\mathbf{r}(\phi, \theta) = 5\sin\phi\cos\theta\,\mathbf{i} + 5\sin\phi\sin\theta\,\mathbf{j} + 5\cos\phi\,\mathbf{k}$, and $|\mathbf{r}_\phi \times \mathbf{r}_\theta| = 25\sin\phi$ (see Example 16.6.10). S is the portion of the sphere where $z \ge 4$, so $0 \le \phi \le \tan^{-1}(3/4)$ and $0 \le \theta \le 2\pi$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(a) $m = \iint_S \rho(x,y,z)dS = \int_0^{2\pi}\int_0^{\tan^{-1}(3/4)} k(25\sin\phi)\,d\phi\,d\theta = 25k\int_0^{2\pi} d\theta \int_0^{\tan^{-1}(3/4)} \sin\phi\,d\phi$

$= 25k(2\pi)\left[-\cos\left(\tan^{-1}\tfrac{3}{4}\right)+1\right] = 50\pi k\left(-\tfrac{4}{5}+1\right) = 10\pi k.$

Because S has constant density, $\overline{x} = \overline{y} = 0$ by symmetry, and

$\overline{z} = \frac{1}{m}\iint_S z\rho(x,y,z)dS = \frac{1}{10\pi k}\int_0^{2\pi}\int_0^{\tan^{-1}(3/4)} k(5\cos\phi)(25\sin\phi)\,d\phi\,d\theta$

$= \frac{1}{10\pi k}(125k)\int_0^{2\pi} d\theta \int_0^{\tan^{-1}(3/4)} \sin\phi\cos\phi\,d\phi = \frac{1}{10\pi k}(125k)(2\pi)\left[\tfrac{1}{2}\sin^2\phi\right]_0^{\tan^{-1}(3/4)} = 25\cdot\tfrac{1}{2}\left(\tfrac{3}{5}\right)^2 = \tfrac{9}{2},$

so the center of mass is $(\overline{x},\overline{y},\overline{z}) = \left(0,0,\tfrac{9}{2}\right)$.

(b) $I_z = \iint_S (x^2+y^2)\rho(x,y,z)dS = \int_0^{2\pi}\int_0^{\tan^{-1}(3/4)} k(25\sin^2\phi)(25\sin\phi)\,d\phi\,d\theta$

$= 625k\int_0^{2\pi} d\theta \int_0^{\tan^{-1}(3/4)} \sin^3\phi\,d\phi = 625k(2\pi)\left[\tfrac{1}{3}\cos^3\phi - \cos\phi\right]_0^{\tan^{-1}(3/4)}$

$= 1250\pi k\left[\tfrac{1}{3}\left(\tfrac{4}{5}\right)^3 - \tfrac{4}{5} - \tfrac{1}{3} + 1\right] = 1250\pi k\left(\tfrac{14}{375}\right) = \tfrac{140}{3}\pi k$

43. The rate of flow through the cylinder is the flux $\iint_S \rho\mathbf{v}\cdot\mathbf{n}\,dS = \iint_S \rho\mathbf{v}\cdot d\mathbf{S}$. We use the parametric representation $\mathbf{r}(u,v) = 2\cos u\,\mathbf{i} + 2\sin u\,\mathbf{j} + v\,\mathbf{k}$ for S, where $0\le u\le 2\pi$, $0\le v\le 1$, so $\mathbf{r}_u = -2\sin u\,\mathbf{i} + 2\cos u\,\mathbf{j}$, $\mathbf{r}_v = \mathbf{k}$, and the outward orientation is given by $\mathbf{r}_u\times\mathbf{r}_v = 2\cos u\,\mathbf{i} + 2\sin u\,\mathbf{j}$. Then

$$\begin{aligned}\iint_S \rho\mathbf{v}\cdot d\mathbf{S} &= \rho\int_0^{2\pi}\int_0^1 \left(v\,\mathbf{i} + 4\sin^2 u\,\mathbf{j} + 4\cos^2 u\,\mathbf{k}\right)\cdot(2\cos u\,\mathbf{i} + 2\sin u\,\mathbf{j})\,dv\,du\\ &= \rho\int_0^{2\pi}\int_0^1 \left(2v\cos u + 8\sin^3 u\right)dv\,du = \rho\int_0^{2\pi}\left(\cos u + 8\sin^3 u\right)du\\ &= \rho\left[\sin u + 8\left(-\tfrac{1}{3}\right)(2+\sin^2 u)\cos u\right]_0^{2\pi} = 0\ \text{kg/s}\end{aligned}$$

44. A parametric representation for the hemisphere S is $\mathbf{r}(\phi,\theta) = 3\sin\phi\cos\theta\,\mathbf{i} + 3\sin\phi\sin\theta\,\mathbf{j} + 3\cos\phi\,\mathbf{k}$, $0\le\phi\le\pi/2$, $0\le\theta\le 2\pi$. Then $\mathbf{r}_\phi = 3\cos\phi\cos\theta\,\mathbf{i} + 3\cos\phi\sin\theta\,\mathbf{j} - 3\sin\phi\,\mathbf{k}$, $\mathbf{r}_\theta = -3\sin\phi\sin\theta\,\mathbf{i} + 3\sin\phi\cos\theta\,\mathbf{j}$, and the outward orientation is given by $\mathbf{r}_\phi\times\mathbf{r}_\theta = 9\sin^2\phi\cos\theta\,\mathbf{i} + 9\sin^2\phi\sin\theta\,\mathbf{j} + 9\sin\phi\cos\phi\,\mathbf{k}$. The rate of flow through S is

$$\begin{aligned}\iint_S \rho\mathbf{v}\cdot d\mathbf{S} &= \rho\int_0^{\pi/2}\int_0^{2\pi}(3\sin\phi\sin\theta\,\mathbf{i} + 3\sin\phi\cos\theta\,\mathbf{j})\cdot\left(9\sin^2\phi\cos\theta\,\mathbf{i} + 9\sin^2\phi\sin\theta\,\mathbf{j} + 9\sin\phi\cos\phi\,\mathbf{k}\right)d\theta\,d\phi\\ &= 27\rho\int_0^{\pi/2}\int_0^{2\pi}\left(\sin^3\phi\sin\theta\cos\theta + \sin^3\phi\sin\theta\cos\theta\right)d\theta\,d\phi = 54\rho\int_0^{\pi/2}\sin^3\phi\,d\phi\int_0^{2\pi}\sin\theta\cos\theta\,d\theta\\ &= 54\rho\left[-\tfrac{1}{3}(2+\sin^2\phi)\cos\phi\right]_0^{\pi/2}\left[\tfrac{1}{2}\sin^2\theta\right]_0^{2\pi} = 0\ \text{kg/s}\end{aligned}$$

45. S consists of the hemisphere S_1 given by $z = \sqrt{a^2-x^2-y^2}$ and the disk S_2 given by $0\le x^2+y^2\le a^2$, $z=0$.

On S_1: $\mathbf{E} = a\sin\phi\cos\theta\,\mathbf{i} + a\sin\phi\sin\theta\,\mathbf{j} + 2a\cos\phi\,\mathbf{k}$,

$\mathbf{T}_\phi\times\mathbf{T}_\theta = a^2\sin^2\phi\cos\theta\,\mathbf{i} + a^2\sin^2\phi\sin\theta\,\mathbf{j} + a^2\sin\phi\cos\phi\,\mathbf{k}$. Thus

$$\begin{aligned}\iint_{S_1}\mathbf{E}\cdot d\mathbf{S} &= \int_0^{2\pi}\int_0^{\pi/2}(a^3\sin^3\phi + 2a^3\sin\phi\cos^2\phi)\,d\phi\,d\theta\\ &= \int_0^{2\pi}\int_0^{\pi/2}(a^3\sin\phi + a^3\sin\phi\cos^2\phi)\,d\phi\,d\theta = (2\pi)a^3\left(1+\tfrac{1}{3}\right) = \tfrac{8}{3}\pi a^3\end{aligned}$$

On S_2: $\mathbf{E} = x\,\mathbf{i} + y\,\mathbf{j}$, and $\mathbf{r}_y\times\mathbf{r}_x = -\mathbf{k}$ so $\iint_{S_2}\mathbf{E}\cdot d\mathbf{S} = 0$. Hence the total charge is $q = \varepsilon_0\iint_S \mathbf{E}\cdot d\mathbf{S} = \frac{8}{3}\pi a^3\varepsilon_0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

46. Referring to the figure, on

S_1: $\mathbf{E} = \mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$, $\mathbf{r}_y \times \mathbf{r}_z = \mathbf{i}$ and $\iint_{S_1} \mathbf{E} \cdot d\mathbf{S} = \int_{-1}^{1}\int_{-1}^{1} dy\,dz = 4$;

S_2: $\mathbf{E} = x\,\mathbf{i} + \mathbf{j} + z\,\mathbf{k}$, $\mathbf{r}_z \times \mathbf{r}_x = \mathbf{j}$ and $\iint_{S_2} \mathbf{E} \cdot d\mathbf{S} = \int_{-1}^{1}\int_{-1}^{1} dx\,dz = 4$;

S_3: $\mathbf{E} = x\,\mathbf{i} + y\,\mathbf{j} + \mathbf{k}$, $\mathbf{r}_x \times \mathbf{r}_y = \mathbf{k}$ and $\iint_{S_3} \mathbf{E} \cdot d\mathbf{S} = \int_{-1}^{1}\int_{-1}^{1} dx\,dy = 4$;

S_4: $\mathbf{E} = -\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$, $\mathbf{r}_z \times \mathbf{r}_y = -\mathbf{i}$ and $\iint_{S_4} \mathbf{E} \cdot d\mathbf{S} = 4$.



Similarly $\iint_{S_5} \mathbf{E} \cdot d\mathbf{S} = \iint_{S_6} \mathbf{E} \cdot d\mathbf{S} = 4$. Hence $q = \varepsilon_0 \iint_S \mathbf{E} \cdot d\mathbf{S} = \varepsilon_0 \sum_{i=1}^{6} \iint_{S_i} \mathbf{E} \cdot d\mathbf{S} = 24\varepsilon_0$.

47. $K\nabla u = 6.5(4y\,\mathbf{j} + 4z\,\mathbf{k})$. S is given by $\mathbf{r}(x,\theta) = x\,\mathbf{i} + \sqrt{6}\cos\theta\,\mathbf{j} + \sqrt{6}\sin\theta\,\mathbf{k}$ and since we want the inward heat flow, we use $\mathbf{r}_x \times \mathbf{r}_\theta = -\sqrt{6}\cos\theta\,\mathbf{j} - \sqrt{6}\sin\theta\,\mathbf{k}$. Then the rate of heat flow inward is given by

$\iint_S (-K\,\nabla u) \cdot d\mathbf{S} = \int_0^{2\pi}\int_0^4 -(6.5)(-24)\,dx\,d\theta = (2\pi)(156)(4) = 1248\pi$.

48. $u(x,y,z) = c/\sqrt{x^2+y^2+z^2}$,

$$\mathbf{F} = -K\,\nabla u = -K\left[-\frac{cx}{(x^2+y^2+z^2)^{3/2}}\,\mathbf{i} - \frac{cy}{(x^2+y^2+z^2)^{3/2}}\,\mathbf{j} - \frac{cz}{(x^2+y^2+z^2)^{3/2}}\,\mathbf{k}\right]$$
$$= \frac{cK}{(x^2+y^2+z^2)^{3/2}}\,(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k})$$

and the outward unit normal is $\mathbf{n} = \dfrac{1}{a}\,(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k})$.

Thus $\mathbf{F} \cdot \mathbf{n} = \dfrac{cK}{a(x^2+y^2+z^2)^{3/2}}\,(x^2+y^2+z^2)$, but on S, $x^2+y^2+z^2 = a^2$ so $\mathbf{F} \cdot \mathbf{n} = \dfrac{cK}{a^2}$. Hence the rate of heat flow across S is $\displaystyle\iint_S \mathbf{F} \cdot d\mathbf{S} = \frac{cK}{a^2}\iint_S dS = \frac{cK}{a^2}\,(4\pi a^2) = 4\pi K c$.

49. Let S be a sphere of radius a centered at the origin. Then $|\mathbf{r}| = a$ and $\mathbf{F}(\mathbf{r}) = c\mathbf{r}/|\mathbf{r}|^3 = (c/a^3)\,(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k})$. A parametric representation for S is $\mathbf{r}(\phi,\theta) = a\sin\phi\cos\theta\,\mathbf{i} + a\sin\phi\sin\theta\,\mathbf{j} + a\cos\phi\,\mathbf{k}$, $0 \le \phi \le \pi$, $0 \le \theta \le 2\pi$. Then $\mathbf{r}_\phi = a\cos\phi\cos\theta\,\mathbf{i} + a\cos\phi\sin\theta\,\mathbf{j} - a\sin\phi\,\mathbf{k}$, $\mathbf{r}_\theta = -a\sin\phi\sin\theta\,\mathbf{i} + a\sin\phi\cos\theta\,\mathbf{j}$, and the outward orientation is given by $\mathbf{r}_\phi \times \mathbf{r}_\theta = a^2\sin^2\phi\cos\theta\,\mathbf{i} + a^2\sin^2\phi\sin\theta\,\mathbf{j} + a^2\sin\phi\cos\phi\,\mathbf{k}$. The flux of $\mathbf{F}$ across S is

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \int_0^{\pi}\int_0^{2\pi} \frac{c}{a^3}\,(a\sin\phi\cos\theta\,\mathbf{i} + a\sin\phi\sin\theta\,\mathbf{j} + a\cos\phi\,\mathbf{k})$$
$$\cdot\,(a^2\sin^2\phi\cos\theta\,\mathbf{i} + a^2\sin^2\phi\sin\theta\,\mathbf{j} + a^2\sin\phi\cos\phi\,\mathbf{k})\,d\theta\,d\phi$$
$$= \frac{c}{a^3}\int_0^{\pi}\int_0^{2\pi} a^3\,(\sin^3\phi + \sin\phi\cos^2\phi)\,d\theta\,d\phi = c\int_0^{\pi}\int_0^{2\pi}\sin\phi\,d\theta\,d\phi = 4\pi c$$

Thus the flux does not depend on the radius a.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16.8 Stokes' Theorem

1. Both H and P are oriented piecewise-smooth surfaces that are bounded by the simple, closed, smooth curve $x^2 + y^2 = 4$, $z = 0$ (which we can take to be oriented positively for both surfaces). Then H and P satisfy the hypotheses of Stokes' Theorem, so by (3) we know $\iint_H \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \int_C \mathbf{F}\cdot d\mathbf{r} = \iint_P \operatorname{curl}\mathbf{F}\cdot d\mathbf{S}$ (where C is the boundary curve).

2. The paraboloid $z = 1 - x^2 - y^2$ intersects the xy-plane in the circle $x^2 + y^2 = 1$, $z = 0$. This boundary curve C should be oriented in the counterclockwise direction when viewed from above, so a vector equation of C is $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j}$, $0 \le t \le 2\pi$. Then $\mathbf{r}'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j}$,

$$\mathbf{F}(\mathbf{r}(t)) = (\cos t)^2(\sin 0)\,\mathbf{i} + (\sin t)^2\,\mathbf{j} + (\cos t)(\sin t)\,\mathbf{k} = \sin^2 t\,\mathbf{j} + \sin t\,\cos t\,\mathbf{k},$$

and by Stokes' Theorem,

$$\begin{aligned}\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \int_C \mathbf{F}\cdot d\mathbf{r} &= \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^{2\pi}(\sin^2 t\,\mathbf{j} + \sin t\,\cos t\,\mathbf{k})\cdot(-\sin t\,\mathbf{i} + \cos t\,\mathbf{j})\,dt\\ &= \int_0^{2\pi}(0 + \sin^2 t\cos t + 0)\,dt = \left[\tfrac{1}{3}\sin^3 t\right]_0^{2\pi} = 0\end{aligned}$$

3. The boundary curve C is the circle $x^2 + z^2 = 16$, $y = 0$ where the hemisphere intersects the xz-plane. The curve should be oriented in the counterclockwise direction when viewed from the right (from the positive y-axis), so a vector equation of C is $\mathbf{r}(t) = 4\cos(-t)\,\mathbf{i} + 4\sin(-t)\,\mathbf{k} = 4\cos t\,\mathbf{i} - 4\sin t\,\mathbf{k}$, $0 \le t \le 2\pi$. Then $\mathbf{r}'(t) = -4\sin t\,\mathbf{i} - 4\cos t\,\mathbf{k}$ and $\mathbf{F}(\mathbf{r}(t)) = (-4\sin t)e^0\,\mathbf{i} + (4\cos t)(\cos 0)\,\mathbf{j} + (4\cos t)(-4\sin t)(\sin 0)\,\mathbf{k} = -4\sin t\,\mathbf{i} + 4\cos t\,\mathbf{j}$, and by Stokes' Theorem,

$$\begin{aligned}\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \int_C \mathbf{F}\cdot d\mathbf{r} &= \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^{2\pi}(-4\sin t\,\mathbf{i} + 4\cos t\,\mathbf{j})\cdot(-4\sin t\,\mathbf{i} - 4\cos t\,\mathbf{k})\,dt\\ &= \int_0^{2\pi}(16\sin^2 t + 0 + 0)\,dt = 16\left[\tfrac{1}{2}t - \tfrac{1}{4}\sin 2t\right]_0^{2\pi} = 16\pi\end{aligned}$$

4. The boundary curve C is the circle $y^2 + z^2 = 4$, $x = 2$ which should be oriented in the counterclockwise direction when viewed from the front, so a vector equation of C is $\mathbf{r}(t) = 2\,\mathbf{i} + 2\cos t\,\mathbf{j} + 2\sin t\,\mathbf{k}$, $0 \le t \le 2\pi$. Then $\mathbf{F}(\mathbf{r}(t)) = \tan^{-1}(32\cos t\sin^2 t)\,\mathbf{i} + 8\cos t\,\mathbf{j} + 16\sin^2 t\,\mathbf{k}$, $\mathbf{r}'(t) = -2\sin t\,\mathbf{j} + 2\cos t\,\mathbf{k}$, and $\mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t) = -16\sin t\cos t + 32\sin^2 t\cos t$. Thus

$$\begin{aligned}\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \oint_C \mathbf{F}\cdot d\mathbf{r} &= \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^{2\pi}(-16\sin t\cos t + 32\sin^2 t\cos t)\,dt\\ &= \left[-8\sin^2 t + \tfrac{32}{3}\sin^3 t\right]_0^{2\pi} = 0\end{aligned}$$

5. C is the square in the plane $z = -1$. Rather than evaluating a line integral around C we can use Equation 3: $\iint_{S_1} \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \oint_C \mathbf{F}\cdot d\mathbf{r} = \iint_{S_2} \operatorname{curl}\mathbf{F}\cdot d\mathbf{S}$ where S_1 is the original cube without the bottom and S_2 is the bottom face of the cube. $\operatorname{curl}\mathbf{F} = x^2 z\,\mathbf{i} + (xy - 2xyz)\,\mathbf{j} + (y - xz)\,\mathbf{k}$. For S_2, we choose $\mathbf{n} = \mathbf{k}$ so that C has the same orientation for both surfaces. Then $\operatorname{curl}\mathbf{F}\cdot\mathbf{n} = y - xz = x + y$ on S_2, where $z = -1$. Thus $\iint_{S_2} \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \int_{-1}^{1}\int_{-1}^{1}(x + y)\,dx\,dy = 0$ so $\iint_{S_1} \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

6. The boundary curve C is the circle $x^2 + z^2 = 1$, $y = 0$ which should be oriented in the counterclockwise direction when viewed from the right, so a vector equation of C is $\mathbf{r}(t) = \cos(-t)\,\mathbf{i} + \sin(-t)\,\mathbf{k} = \cos t\,\mathbf{i} - \sin t\,\mathbf{k}$, $0 \le t \le 2\pi$. Then $\mathbf{F}(\mathbf{r}(t)) = \mathbf{i} + e^{-\cos t \sin t}\,\mathbf{j} - \cos^2 t\,\sin t\,\mathbf{k}$, $\mathbf{r}'(t) = -\sin t\,\mathbf{i} - \cos t\,\mathbf{k}$, and $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = -\sin t + \cos^3 t\,\sin t$. Thus

$$\begin{aligned}\iint_S \operatorname{curl}\mathbf{F} \cdot d\mathbf{S} &= \oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\,dt = \int_0^{2\pi} (-\sin t + \cos^3 t\,\sin t)\,dt \\ &= \left[\cos t - \tfrac{1}{4}\cos^4 t\right]_0^{2\pi} = 0\end{aligned}$$

7. $\operatorname{curl}\mathbf{F} = -2z\,\mathbf{i} - 2x\,\mathbf{j} - 2y\,\mathbf{k}$ and we take the surface S to be the planar region enclosed by C, so S is the portion of the plane $x + y + z = 1$ over $D = \{(x, y) \mid 0 \le x \le 1, 0 \le y \le 1 - x\}$. Since C is oriented counterclockwise, we orient S upward. Using Equation 16.7.10, we have $z = g(x, y) = 1 - x - y$, $P = -2z$, $Q = -2x$, $R = -2y$, and

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F} \cdot d\mathbf{S} = \iint_D [-(-2z)(-1) - (-2x)(-1) + (-2y)]\,dA \\ &= \int_0^1 \int_0^{1-x} (-2)\,dy\,dx = -2\int_0^1 (1 - x)\,dx = -1\end{aligned}$$

8. $\operatorname{curl}\mathbf{F} = (x - y)\,\mathbf{i} - y\,\mathbf{j} + \mathbf{k}$ and S is the portion of the plane $3x + 2y + z = 1$ over $D = \left\{(x, y) \mid 0 \le x \le \tfrac{1}{3}, 0 \le y \le \tfrac{1}{2}(1 - 3x)\right\}$. We orient S upward and use Equation 16.7.10 with $z = g(x, y) = 1 - 3x - 2y$:

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F} \cdot d\mathbf{S} = \iint_D [-(x - y)(-3) - (-y)(-2) + 1]\,dA = \int_0^{1/3} \int_0^{(1-3x)/2} (1 + 3x - 5y)\,dy\,dx \\ &= \int_0^{1/3} \left[(1 + 3x)y - \tfrac{5}{2}y^2\right]_{y=0}^{y=(1-3x)/2} dx = \int_0^{1/3} \left[\tfrac{1}{2}(1 + 3x)(1 - 3x) - \tfrac{5}{2} \cdot \tfrac{1}{4}(1 - 3x)^2\right] dx \\ &= \int_0^{1/3} \left(-\tfrac{81}{8}x^2 + \tfrac{15}{4}x - \tfrac{1}{8}\right) dx = \left[-\tfrac{27}{8}x^3 + \tfrac{15}{8}x^2 - \tfrac{1}{8}x\right]_0^{1/3} = -\tfrac{1}{8} + \tfrac{5}{24} - \tfrac{1}{24} = \tfrac{1}{24}\end{aligned}$$

9. $\operatorname{curl}\mathbf{F} = -y\,\mathbf{i} - z\,\mathbf{j} - x\,\mathbf{k}$ and we take S to be the part of the paraboloid $z = 1 - x^2 - y^2$ in the first octant. Since C is oriented counterclockwise (from above), we orient S upward. Then using Equation 16.7.10 with $z = g(x, y) = 1 - x^2 - y^2$ we have

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F} \cdot d\mathbf{S} = \iint_D [-(-y)(-2x) - (-z)(-2y) + (-x)]\,dA = \iint_D \left[-2xy - 2y(1 - x^2 - y^2) - x\right] dA \\ &= \int_0^{\pi/2} \int_0^1 \left[-2(r\cos\theta)(r\sin\theta) - 2(r\sin\theta)(1 - r^2) - r\cos\theta\right] r\,dr\,d\theta \\ &= \int_0^{\pi/2} \int_0^1 \left[-2r^3\sin\theta\cos\theta - 2(r^2 - r^4)\sin\theta - r^2\cos\theta\right] dr\,d\theta \\ &= \int_0^{\pi/2} \left[-\tfrac{1}{2}r^4\sin\theta\cos\theta - 2\left(\tfrac{1}{3}r^3 - \tfrac{1}{5}r^5\right)\sin\theta - \tfrac{1}{3}r^3\cos\theta\right]_{r=0}^{r=1} d\theta \\ &= \int_0^{\pi/2} \left(-\tfrac{1}{2}\sin\theta\cos\theta - \tfrac{4}{15}\sin\theta - \tfrac{1}{3}\cos\theta\right) d\theta = \left[-\tfrac{1}{4}\sin^2\theta + \tfrac{4}{15}\cos\theta - \tfrac{1}{3}\sin\theta\right]_0^{\pi/2} \\ &= -\tfrac{1}{4} - \tfrac{4}{15} - \tfrac{1}{3} = -\tfrac{17}{20}\end{aligned}$$

10. The curve of intersection is an ellipse in the plane $z = y + 2$. $\operatorname{curl}\mathbf{F} = (1 - x)\,\mathbf{i} - \mathbf{j} + (z - 2)\,\mathbf{k}$ and we take the surface S to be the planar region enclosed by C with upward orientation. From Equation 16.7.10 with $z = g(x, y) = y + 2$ we have

$$\begin{aligned}\oint_C \mathbf{F} \cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F} \cdot d\mathbf{S} = \iint_{x^2 + y^2 \le 1} [-(1 - x)(0) - (-1)(1) + (y + 2 - 2)]\,dA \\ &= \iint_{x^2 + y^2 \le 1} (y + 1)\,dA = \int_0^{2\pi} \int_0^1 (r\sin\theta + 1)\,r\,dr\,d\theta = \int_0^{2\pi} \left[\tfrac{1}{3}r^3\sin\theta + \tfrac{1}{2}r^2\right]_{r=0}^{r=1} d\theta \\ &= \int_0^{2\pi} \left(\tfrac{1}{3}\sin\theta + \tfrac{1}{2}\right) d\theta = \left[-\tfrac{1}{3}\cos\theta + \tfrac{1}{2}\theta\right]_0^{2\pi} = \pi\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. (a) The curve of intersection is an ellipse in the plane $x + y + z = 1$ with unit normal $\mathbf{n} = \frac{1}{\sqrt{3}}(\mathbf{i} + \mathbf{j} + \mathbf{k})$, $\operatorname{curl} \mathbf{F} = x^2\,\mathbf{j} + y^2\,\mathbf{k}$, and $\operatorname{curl} \mathbf{F} \cdot \mathbf{n} = \frac{1}{\sqrt{3}}(x^2 + y^2)$. Then

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \tfrac{1}{\sqrt{3}}(x^2 + y^2)\,dS = \iint_{x^2+y^2 \le 9} (x^2 + y^2)\,dx\,dy = \int_0^{2\pi}\int_0^3 r^3\,dr\,d\theta = 2\pi\left(\tfrac{81}{4}\right) = \tfrac{81\pi}{2}$$

(b)



(c) One possible parametrization is $x = 3\cos t$, $y = 3\sin t$, $z = 1 - 3\cos t - 3\sin t$, $0 \le t \le 2\pi$.



12. (a) S is the part of the surface $z = y^2 - x^2$ that lies above the unit disk D. $\operatorname{curl} \mathbf{F} = x\,\mathbf{i} - y\,\mathbf{j} + (x^2 - x^2)\,\mathbf{k} = x\,\mathbf{i} - y\,\mathbf{j}$. Using Equation 16.7.10 with $g(x, y) = y^2 - x^2$, $P = x$, $Q = -y$, we have

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} = \iint_D [-x(-2x) - (-y)(2y)]\,dA = 2\iint_D (x^2 + y^2)\,dA \\ &= 2\int_0^{2\pi}\int_0^1 r^2 r\,dr\,d\theta = 2(2\pi)\left[\tfrac{1}{4}r^4\right]_0^1 = \pi\end{aligned}$$

(b)



(c) One possible set of parametric equations is $x = \cos t$, $y = \sin t$, $z = \sin^2 t - \cos^2 t$, $0 \le t \le 2\pi$.



13. The boundary curve C is the circle $x^2 + y^2 = 16$, $z = 4$ oriented in the clockwise direction as viewed from above (since S is oriented downward). We can parametrize C by $\mathbf{r}(t) = 4\cos t\,\mathbf{i} - 4\sin t\,\mathbf{j} + 4\,\mathbf{k}$, $0 \le t \le 2\pi$, and then $\mathbf{r}'(t) = -4\sin t\,\mathbf{i} - 4\cos t\,\mathbf{j}$. Thus $\mathbf{F}(\mathbf{r}(t)) = 4\sin t\,\mathbf{i} + 4\cos t\,\mathbf{j} - 2\,\mathbf{k}$, $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = -16\sin^2 t - 16\cos^2 t = -16$, and

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\,dt = \int_0^{2\pi}(-16)\,dt = -16\,(2\pi) = -32\pi$$

Now $\operatorname{curl} \mathbf{F} = 2\,\mathbf{k}$, and the projection D of S on the xy-plane is the disk $x^2 + y^2 \le 16$, so by Equation 16.7.10 with $z = g(x, y) = \sqrt{x^2 + y^2}$ [and multiplying by -1 for the downward orientation] we have

$$\iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} = -\iint_D (-0 - 0 + 2)\,dA = -2 \cdot A(D) = -2 \cdot \pi(4^2) = -32\pi$$

14. The paraboloid intersects the plane $z = 1$ when $1 = 5 - x^2 - y^2 \iff x^2 + y^2 = 4$, so the boundary curve C is the circle $x^2 + y^2 = 4$, $z = 1$ oriented in the counterclockwise direction as viewed from above. We can parametrize C by $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} + \mathbf{k}$, $0 \le t \le 2\pi$, and then $\mathbf{r}'(t) = -2\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$. Thus $\mathbf{F}(\mathbf{r}(t)) = -4\sin t\,\mathbf{i} + 2\sin t\,\mathbf{j} + 6\cos t\,\mathbf{k}$, $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = 8\sin^2 t + 4\sin t\cos t$, and

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi}(8\sin^2 t + 4\sin t\cos t)\,dt = 8\left(\tfrac{1}{2}t - \tfrac{1}{4}\sin 2t\right) + 2\sin^2 t\Big]_0^{2\pi} = 8\pi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

Now curl $\mathbf{F} = (-3-2y)\,\mathbf{j} + 2z\,\mathbf{k}$, and the projection D of S on the xy-plane is the disk $x^2+y^2 \le 4$, so by Equation 16.7.10 with $z = g(x,y) = 5-x^2-y^2$ we have

$$\begin{aligned}\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} &= \iint_D [-0-(-3-2y)(-2y)+2z]\,dA = \iint_D [-6y-4y^2+2(5-x^2-y^2)]\,dA\\ &= \int_0^{2\pi}\int_0^2 \left[-6r\sin\theta - 4r^2\sin^2\theta + 2(5-r^2)\right] r\,dr\,d\theta = \int_0^{2\pi}\left[-2r^3\sin\theta - r^4\sin^2\theta + 5r^2 - \tfrac{1}{2}r^4\right]_{r=0}^{r=2} d\theta\\ &= \int_0^{2\pi}\left(-16\sin\theta - 16\sin^2\theta + 20 - 8\right) d\theta = 16\cos\theta - 16\left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right) + 12\theta\Big]_0^{2\pi} = 8\pi\end{aligned}$$

15. The boundary curve C is the circle $x^2+z^2=1$, $y=0$ oriented in the counterclockwise direction as viewed from the positive y-axis. Then C can be described by $\mathbf{r}(t) = \cos t\,\mathbf{i} - \sin t\,\mathbf{k}$, $0 \le t \le 2\pi$, and $\mathbf{r}'(t) = -\sin t\,\mathbf{i} - \cos t\,\mathbf{k}$. Thus $\mathbf{F}(\mathbf{r}(t)) = -\sin t\,\mathbf{j} + \cos t\,\mathbf{k}$, $\mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t) = -\cos^2 t$, and $\oint_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi}(-\cos^2 t)\,dt = -\frac{1}{2}t - \frac{1}{4}\sin 2t\Big]_0^{2\pi} = -\pi$.
Now curl $\mathbf{F} = -\mathbf{i}-\mathbf{j}-\mathbf{k}$, and S can be parametrized (see Example 16.6.10) by
$\mathbf{r}(\phi,\theta) = \sin\phi\cos\theta\,\mathbf{i} + \sin\phi\sin\theta\,\mathbf{j} + \cos\phi\,\mathbf{k}$, $0 \le \theta \le \pi$, $0 \le \phi \le \pi$. Then
$\mathbf{r}_\phi \times \mathbf{r}_\theta = \sin^2\phi\cos\theta\,\mathbf{i} + \sin^2\phi\sin\theta\,\mathbf{j} + \sin\phi\cos\phi\,\mathbf{k}$ and

$$\begin{aligned}\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} &= \iint_{x^2+z^2\le 1} \operatorname{curl}\mathbf{F}\cdot(\mathbf{r}_\phi\times\mathbf{r}_\theta)\,dA = \int_0^\pi\int_0^\pi(-\sin^2\phi\cos\theta - \sin^2\phi\sin\theta - \sin\phi\cos\phi)\,d\theta\,d\phi\\ &= \int_0^\pi(-2\sin^2\phi - \pi\sin\phi\cos\phi)\,d\phi = \left[\tfrac{1}{2}\sin 2\phi - \phi - \tfrac{\pi}{2}\sin^2\phi\right]_0^\pi = -\pi\end{aligned}$$

16. Let S be the surface in the plane $x+y+z=1$ with upward orientation enclosed by C. Then an upward unit normal vector for S is $\mathbf{n} = \frac{1}{\sqrt{3}}(\mathbf{i}+\mathbf{j}+\mathbf{k})$. Orient C in the counterclockwise direction, as viewed from above. $\int_C z\,dx - 2x\,dy + 3y\,dz$ is equivalent to $\int_C \mathbf{F}\cdot d\mathbf{r}$ for $\mathbf{F}(x,y,z) = z\,\mathbf{i} - 2x\,\mathbf{j} + 3y\,\mathbf{k}$, and the components of $\mathbf{F}$ are polynomials, which have continuous partial derivatives throughout $\mathbb{R}^3$. We have curl $\mathbf{F} = 3\,\mathbf{i}+\mathbf{j}-2\,\mathbf{k}$, so by Stokes' Theorem,

$$\begin{aligned}\int_C z\,dx - 2x\,dy + 3y\,dz &= \int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,dS = \iint_S (3\,\mathbf{i}+\mathbf{j}-2\,\mathbf{k})\cdot\tfrac{1}{\sqrt{3}}(\mathbf{i}+\mathbf{j}+\mathbf{k})\,dS\\ &= \tfrac{2}{\sqrt{3}}\iint_S dS = \tfrac{2}{\sqrt{3}}\,(\text{surface area of } S)\end{aligned}$$

Thus the value of $\int_C z\,dx - 2x\,dy + 3y\,dz$ is always $\frac{2}{\sqrt{3}}$ times the area of the region enclosed by C, regardless of its shape or location. [Notice that because $\mathbf{n}$ is normal to a plane, it is constant. But curl $\mathbf{F}$ is also constant, so the dot product curl $\mathbf{F}\cdot\mathbf{n}$ is constant and we could have simply argued that $\iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,dS$ is a constant multple of $\iint_S dS$, the surface area of S.]

17. It is easier to use Stokes' Theorem than to compute the work directly. Let S be the planar region enclosed by the path of the particle, so S is the portion of the plane $z = \frac{1}{2}y$ for $0 \le x \le 1$, $0 \le y \le 2$, with upward orientation.
curl $\mathbf{F} = 8y\,\mathbf{i} + 2z\,\mathbf{j} + 2y\,\mathbf{k}$ and

$$\begin{aligned}\oint_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D \left[-8y\,(0) - 2z\left(\tfrac{1}{2}\right) + 2y\right] dA = \int_0^1\int_0^2 \left(2y - \tfrac{1}{2}y\right) dy\,dx\\ &= \int_0^1\int_0^2 \tfrac{3}{2}y\,dy\,dx = \int_0^1 \left[\tfrac{3}{4}y^2\right]_{y=0}^{y=2} dx = \int_0^1 3\,dx = 3\end{aligned}$$

18. $\int_C (y+\sin x)\,dx + (z^2+\cos y)\,dy + x^3\,dz = \int_C \mathbf{F}\cdot d\mathbf{r}$, where $\mathbf{F}(x,y,z) = (y+\sin x)\,\mathbf{i} + (z^2+\cos y)\,\mathbf{j} + x^3\,\mathbf{k}$ $\Rightarrow$ curl $\mathbf{F} = -2z\,\mathbf{i} - 3x^2\,\mathbf{j} - \mathbf{k}$. Since $\sin 2t = 2\sin t\cos t$, C lies on the surface $z = 2xy$. Let S be the part of this surface that is bounded by C. Then the projection of S onto the xy-plane is the unit disk D $[x^2+y^2 \le 1]$. C is traversed clockwise (when viewed from above) so S is oriented downward. Using Equation 16.7.10 with $g(x,y) = 2xy$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$P = -2z = -2(2xy) = -4xy$, $Q = -3x^2$, $R = -1$ and multiplying by -1 for the downward orientation, we have

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= -\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = -\iint_D \left[-(-4xy)(2y) - (-3x^2)(2x) - 1\right] dA \\ &= -\iint_D (8xy^2 + 6x^3 - 1)\, dA = -\int_0^{2\pi}\int_0^1 (8r^3\cos\theta\sin^2\theta + 6r^3\cos^3\theta - 1)\, r\, dr\, d\theta \\ &= -\int_0^{2\pi} \left(\tfrac{8}{5}\cos\theta\sin^2\theta + \tfrac{6}{5}\cos^3\theta - \tfrac{1}{2}\right) d\theta = -\left[\tfrac{8}{15}\sin^3\theta + \tfrac{6}{5}\left(\sin\theta - \tfrac{1}{3}\sin^3\theta\right) - \tfrac{1}{2}\theta\right]_0^{2\pi} = \pi\end{aligned}$$

19. Assume S is centered at the origin with radius a and let H_1 and H_2 be the upper and lower hemispheres, respectively, of S. Then $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_{H_1} \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} + \iint_{H_2} \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \oint_{C_1}\mathbf{F}\cdot d\mathbf{r} + \oint_{C_2}\mathbf{F}\cdot d\mathbf{r}$ by Stokes' Theorem. But C_1 is the circle $x^2 + y^2 = a^2$ oriented in the counterclockwise direction while C_2 is the same circle oriented in the clockwise direction. Hence $\oint_{C_2}\mathbf{F}\cdot d\mathbf{r} = -\oint_{C_1}\mathbf{F}\cdot d\mathbf{r}$ so $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = 0$ as desired.

20. (a) By Exercise 16.5.26, $\operatorname{curl}(f\nabla g) = f\operatorname{curl}(\nabla g) + \nabla f \times \nabla g = \nabla f \times \nabla g$ since $\operatorname{curl}(\nabla g) = \mathbf{0}$. Hence by Stokes' Theorem $\int_C (f\nabla g)\cdot d\mathbf{r} = \iint_S (\nabla f \times \nabla g)\cdot d\mathbf{S}$.

(b) As in (a), $\operatorname{curl}(f\nabla f) = \nabla f \times \nabla f = \mathbf{0}$, so by Stokes' Theorem, $\int_C (f\nabla f)\cdot d\mathbf{r} = \iint_S [\operatorname{curl}(f\nabla f)]\cdot d\mathbf{S} = 0$.

(c) As in part (a),

$$\begin{aligned}\operatorname{curl}(f\nabla g + g\nabla f) &= \operatorname{curl}(f\nabla g) + \operatorname{curl}(g\nabla f) \qquad \text{[by Exercise 16.5.24]} \\ &= (\nabla f \times \nabla g) + (\nabla g \times \nabla f) = \mathbf{0} \qquad \text{[since } \mathbf{u}\times\mathbf{v} = -(\mathbf{v}\times\mathbf{u})\text{]}\end{aligned}$$

Hence by Stokes' Theorem, $\int_C (f\nabla g + g\nabla f)\cdot d\mathbf{r} = \iint_S \operatorname{curl}(f\nabla g + g\nabla f)\cdot d\mathbf{S} = 0$.

16.9 The Divergence Theorem

1. $\operatorname{div}\mathbf{F} = 3 + x + 2x = 3 + 3x$, so

$\iiint_E \operatorname{div}\mathbf{F}\, dV = \int_0^1\int_0^1\int_0^1 (3x+3)\, dx\, dy\, dz = \frac{9}{2}$ (notice the triple integral is three times the volume of the cube plus three times $\overline{x}$).



To compute $\iint_S \mathbf{F}\cdot d\mathbf{S}$, on

S_1: $\mathbf{n} = \mathbf{i}$, $\mathbf{F} = 3\mathbf{i} + y\mathbf{j} + 2z\mathbf{k}$, and $\iint_{S_1}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_1} 3\, dS = 3$;

S_2: $\mathbf{F} = 3x\mathbf{i} + x\mathbf{j} + 2xz\mathbf{k}$, $\mathbf{n} = \mathbf{j}$ and $\iint_{S_2}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_2} x\, dS = \frac{1}{2}$;

S_3: $\mathbf{F} = 3x\mathbf{i} + xy\mathbf{j} + 2x\mathbf{k}$, $\mathbf{n} = \mathbf{k}$ and $\iint_{S_3}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_3} 2x\, dS = 1$;

S_4: $\mathbf{F} = \mathbf{0}$, $\iint_{S_4}\mathbf{F}\cdot d\mathbf{S} = 0$; S_5: $\mathbf{F} = 3x\mathbf{i} + 2x\mathbf{k}$, $\mathbf{n} = -\mathbf{j}$ and $\iint_{S_5}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_5} 0\, dS = 0$;

S_6: $\mathbf{F} = 3x\mathbf{i} + xy\mathbf{j}$, $\mathbf{n} = -\mathbf{k}$ and $\iint_{S_6}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_6} 0\, dS = 0$. Thus $\iint_S \mathbf{F}\cdot d\mathbf{S} = \frac{9}{2}$.

2. $\operatorname{div}\mathbf{F} = 0 + 2z + 8z = 10z$ so, using cylindrical coordinates,



$$\begin{aligned}\iiint_E \operatorname{div}\mathbf{F}\, dV &= \iiint_E 10z\, dV = \int_0^{2\pi}\int_0^3\int_{r^2}^9 (10z)\, r\, dz\, dr\, d\theta \\ &= \int_0^{2\pi}\int_0^3 \left[5rz^2\right]_{z=r^2}^{z=9} dr\, d\theta = \int_0^{2\pi}\int_0^3 (405r - 5r^5)\, dr\, d\theta \\ &= \int_0^{2\pi} d\theta \int_0^3 (405r - 5r^5)\, dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{405}{2}r^2 - \tfrac{5}{6}r^6\right]_0^3 \\ &= 2\pi\left(\tfrac{3645}{2} - \tfrac{1215}{2}\right) = 2430\pi\end{aligned}$$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

On S_1: The surface is $z = x^2 + y^2,\ x^2 + y^2 \le 9$, with downward orientation, and $\mathbf{F}(x, y, z) = y^2z^3\,\mathbf{i} + 2yz\,\mathbf{j} + 4z^2\,\mathbf{k}$. Then

$$\begin{aligned}
\iint_{S_1} \mathbf{F}\cdot d\mathbf{S} &= -\iint_D [-(y^2z^3)(2x) - (2yz)(2y) + (4z^2)]\,dA \\
&= \iint_D \left[2xy^2(x^2+y^2)^3 + 4y^2(x^2+y^2) - 4(x^2+y^2)^2\right] dA \\
&= \int_0^{2\pi}\int_0^3 (2r^3\cos\theta\sin^2\theta\cdot r^6 + 4r^2\sin^2\theta\cdot r^2 - 4r^4)\,r\,dr\,d\theta \\
&= \int_0^{2\pi}\int_0^3 (2r^{10}\sin^2\theta\cos\theta + 4r^5\sin^2\theta - 4r^5)\,dr\,d\theta \\
&= \int_0^{2\pi} \left[\tfrac{2}{11}r^{11}\sin^2\theta\cos\theta + \tfrac{2}{3}r^6\sin^2\theta - \tfrac{2}{3}r^6\right]_{r=0}^{r=3} d\theta \\
&= \int_0^{2\pi} \left(\tfrac{354{,}294}{11}\sin^2\theta\cos\theta + 486\sin^2\theta - 486\right) d\theta \\
&= \left[\tfrac{354{,}294}{11}\cdot\tfrac{1}{3}\sin^3\theta + 486\left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right) - 486\theta\right]_0^{2\pi} \\
&= 0 + 486(\pi - 0) - 486(2\pi) = -486\pi
\end{aligned}$$

On S_2: The surface is $z = 9,\ x^2 + y^2 \le 9$, with upward orientation, so $\mathbf{F}(x, y, z) = y^2z^3\,\mathbf{i} + 2yz\,\mathbf{j} + 4z^2\,\mathbf{k}$ and

$$\begin{aligned}
\iint_{S_2} \mathbf{F}\cdot d\mathbf{S} &= \iint_D [-(y^2z^3)(0) - (2yz)(0) + (4z^2)]\,dA = \iint_D 4(9)^2\,dA \\
&= 324\,A(D) = 324\cdot\pi(3)^2 = 2916\pi
\end{aligned}$$

Thus $\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_{S_1} \mathbf{F}\cdot d\mathbf{S} + \iint_{S_2} \mathbf{F}\cdot d\mathbf{S} = -486\pi + 2916\pi = 2430\pi$.

3. $\operatorname{div}\mathbf{F} = 0 + 1 + 0 = 1$, so $\iiint_E \operatorname{div}\mathbf{F}\,dV = \iiint_E 1\,dV = V(E) = \frac{4}{3}\pi\cdot 4^3 = \frac{256}{3}\pi$.

S is a sphere of radius 4 centered at the origin which can be parametrized by $\mathbf{r}(\phi, \theta) = \langle 4\sin\phi\cos\theta, 4\sin\phi\sin\theta, 4\cos\phi\rangle$, $0 \le \phi \le \pi$, $0 \le \theta \le 2\pi$ (similar to Example 16.6.10). Then

$$\begin{aligned}
\mathbf{r}_\phi \times \mathbf{r}_\theta &= \langle 4\cos\phi\cos\theta, 4\cos\phi\sin\theta, -4\sin\phi\rangle \times \langle -4\sin\phi\sin\theta, 4\sin\phi\cos\theta, 0\rangle \\
&= \left\langle 16\sin^2\phi\cos\theta, 16\sin^2\phi\sin\theta, 16\cos\phi\sin\phi\right\rangle
\end{aligned}$$

and $\mathbf{F}(\mathbf{r}(\phi, \theta)) = \langle 4\cos\phi, 4\sin\phi\sin\theta, 4\sin\phi\cos\theta\rangle$. Thus

$\mathbf{F}\cdot(\mathbf{r}_\phi \times \mathbf{r}_\theta) = 64\cos\phi\sin^2\phi\cos\theta + 64\sin^3\phi\sin^2\theta + 64\cos\phi\sin^2\phi\cos\theta = 128\cos\phi\sin^2\phi\cos\theta + 64\sin^3\phi\sin^2\theta$

and

$$\begin{aligned}
\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iint_D \mathbf{F}\cdot(\mathbf{r}_\phi\times\mathbf{r}_\theta)\,dA = \int_0^{2\pi}\int_0^{\pi} (128\cos\phi\sin^2\phi\cos\theta + 64\sin^3\phi\sin^2\theta)\,d\phi\,d\theta \\
&= \int_0^{2\pi} \left[\tfrac{128}{3}\sin^3\phi\cos\theta + 64\left(\tfrac{1}{3}\cos^3\phi - \cos\phi\right)\sin^2\theta\right]_{\phi=0}^{\phi=\pi} d\theta \\
&= \int_0^{2\pi} \tfrac{256}{3}\sin^2\theta\,d\theta = \tfrac{256}{3}\left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi} = \tfrac{256}{3}\pi
\end{aligned}$$

4. $\operatorname{div}\mathbf{F} = 2x - 1 + 1 = 2x$, so

$$\iiint_E \operatorname{div}\mathbf{F}\,dV = \iint_{y^2+z^2\le 9} \left[\int_0^2 2x\,dx\right] dA = \iint_{y^2+z^2\le 9} 4\,dA = 4(\text{area of circle}) = 4(\pi\cdot 3^2) = 36\pi$$

Let S_1 be the front of the cylinder (in the plane $x = 2$), S_2 the back (in the yz-plane), and S_3 the lateral surface of the cylinder. S_1 is the disk $x = 2$, $y^2 + z^2 \le 9$. A unit normal vector is $\mathbf{n} = \langle 1, 0, 0\rangle$ and $\mathbf{F} = \langle 4, -y, z\rangle$ on S_1, so $\iint_{S_1} \mathbf{F}\cdot d\mathbf{S} = \iint_{S_1} \mathbf{F}\cdot\mathbf{n}\,dS = \iint_{S_1} 4\,dS = 4(\text{surface area of } S_1) = 4(\pi\cdot 3^2) = 36\pi$. S_2 is the disk $x = 0$, $y^2 + z^2 \le 9$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Here $\mathbf{n} = \langle -1, 0, 0\rangle$ and $\mathbf{F} = \langle 0, -y, z\rangle$, so $\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_2} \mathbf{F} \cdot \mathbf{n}\, dS = \iint_{S_2} 0\, dS = 0$.

S_3 can be parametrized by $\mathbf{r}(x, \theta) = \langle x, 3\cos\theta, 3\sin\theta\rangle$, $0 \le x \le 2$, $0 \le \theta \le 2\pi$. Then $\mathbf{r}_x \times \mathbf{r}_\theta = \langle 1, 0, 0\rangle \times \langle 0, -3\sin\theta, 3\cos\theta\rangle = \langle 0, -3\cos\theta, -3\sin\theta\rangle$. For the outward (positive) orientation we use $-(\mathbf{r}_x \times \mathbf{r}_\theta)$ and $\mathbf{F}(\mathbf{r}(x, \theta)) = \langle x^2, -3\cos\theta, 3\sin\theta\rangle$, so

$$\begin{aligned}\iint_{S_3} \mathbf{F} \cdot d\mathbf{S} &= \iint_D \mathbf{F} \cdot (-(\mathbf{r}_x \times \mathbf{r}_\theta))\, dA = \int_0^2 \int_0^{2\pi} (0 - 9\cos^2\theta + 9\sin^2\theta)\, d\theta\, dx \\ &= -9\int_0^2 dx \int_0^{2\pi} \cos 2\theta\, d\theta = -9\,(2)\left[\tfrac{1}{2}\sin 2\theta\right]_0^{2\pi} = 0\end{aligned}$$

Thus $\iint_S \mathbf{F} \cdot d\mathbf{S} = 36\pi + 0 + 0 = 36\pi$.

5. $\operatorname{div}\mathbf{F} = \frac{\partial}{\partial x}(xye^z) + \frac{\partial}{\partial y}(xy^2z^3) + \frac{\partial}{\partial z}(-ye^z) = ye^z + 2xyz^3 - ye^z = 2xyz^3$, so by the Divergence Theorem,

$$\begin{aligned}\iint_S \mathbf{F} \cdot d\mathbf{S} &= \iiint_E \operatorname{div}\mathbf{F}\, dV = \int_0^3 \int_0^2 \int_0^1 2xyz^3\, dz\, dy\, dx = 2\int_0^3 x\, dx \int_0^2 y\, dy \int_0^1 z^3\, dz \\ &= 2\left[\tfrac{1}{2}x^2\right]_0^3 \left[\tfrac{1}{2}y^2\right]_0^2 \left[\tfrac{1}{4}z^4\right]_0^1 = 2\left(\tfrac{9}{2}\right)(2)\left(\tfrac{1}{4}\right) = \tfrac{9}{2}\end{aligned}$$

6. $\operatorname{div}\mathbf{F} = \frac{\partial}{\partial x}(x^2yz) + \frac{\partial}{\partial y}(xy^2z) + \frac{\partial}{\partial z}(xyz^2) = 2xyz + 2xyz + 2xyz = 6xyz$, so by the Divergence Theorem,

$$\begin{aligned}\iint_S \mathbf{F} \cdot d\mathbf{S} &= \iiint_E \operatorname{div}\mathbf{F}\, dV = \int_0^a \int_0^b \int_0^c 6xyz\, dz\, dy\, dx = 6\int_0^a x\, dx \int_0^b y\, dy \int_0^c z\, dz \\ &= 6\left[\tfrac{1}{2}x^2\right]_0^a \left[\tfrac{1}{2}y^2\right]_0^b \left[\tfrac{1}{2}z^2\right]_0^c = 6\left(\tfrac{1}{2}a^2\right)\left(\tfrac{1}{2}b^2\right)\left(\tfrac{1}{2}c^2\right) = \tfrac{3}{4}a^2b^2c^2\end{aligned}$$

7. $\operatorname{div}\mathbf{F} = 3y^2 + 0 + 3z^2$, so using cylindrical coordinates with $y = r\cos\theta$, $z = r\sin\theta$, $x = x$ we have

$$\begin{aligned}\iint_S \mathbf{F} \cdot d\mathbf{S} &= \iiint_E (3y^2 + 3z^2)\, dV = \int_0^{2\pi} \int_0^1 \int_{-1}^2 (3r^2\cos^2\theta + 3r^2\sin^2\theta)\, r\, dx\, dr\, d\theta \\ &= 3\int_0^{2\pi} d\theta \int_0^1 r^3\, dr \int_{-1}^2 dx = 3\left[\theta\right]_0^{2\pi} \left[\tfrac{1}{4}r^4\right]_0^1 \left[x\right]_{-1}^2 = 3(2\pi)\left(\tfrac{1}{4}\right)(3) = \tfrac{9\pi}{2}\end{aligned}$$

8. $\operatorname{div}\mathbf{F} = 3x^2 + 3y^2 + 3z^2$, so by the Divergence Theorem,

$$\begin{aligned}\iint_S \mathbf{F} \cdot d\mathbf{S} &= \iiint_E 3(x^2 + y^2 + z^2)\, dV = \int_0^{\pi} \int_0^{2\pi} \int_0^2 3\rho^2 \cdot \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi = 3\int_0^{\pi} \sin\phi\, d\phi \int_0^{2\pi} d\theta \int_0^2 \rho^4\, d\rho \\ &= 3\left[-\cos\phi\right]_0^{\pi} \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{5}\rho^5\right]_0^2 = 3\,(2)\,(2\pi)\left(\tfrac{32}{5}\right) = \tfrac{384}{5}\pi\end{aligned}$$

9. $\operatorname{div}\mathbf{F} = e^y + (-e^y) + 0 = 0$, so by the Divergence Theorem, $\iint_S \mathbf{F} \cdot d\mathbf{S} = \iiint_E 0\, dV = 0$.

10. The tetrahedron has vertices $(0, 0, 0)$, $(a, 0, 0)$, $(0, b, 0)$, $(0, 0, c)$ and is described by $E = \left\{(x, y, z) \mid 0 \le x \le a,\ 0 \le y \le b\left(1 - \frac{x}{a}\right),\ 0 \le z \le c\left(1 - \frac{x}{a} - \frac{y}{b}\right)\right\}$. Here we have $\operatorname{div}\mathbf{F} = 0 + 1 + x = x + 1$, so

$$\begin{aligned}\iint_S \mathbf{F} \cdot d\mathbf{S} &= \iiint_E (x+1)\, dV = \int_0^a \int_0^{b\left(1-\frac{x}{a}\right)} \int_0^{c\left(1-\frac{x}{a}-\frac{y}{b}\right)} (x+1)\, dz\, dy\, dx \\ &= \int_0^a \int_0^{b\left(1-\frac{x}{a}\right)} (x+1)\left[c\left(1 - \tfrac{x}{a} - \tfrac{y}{b}\right)\right] dy\, dx = c\int_0^a (x+1)\left[\left(1-\tfrac{x}{a}\right)y - \tfrac{1}{2b}y^2\right]_{y=0}^{y=b\left(1-\frac{x}{a}\right)} dx \\ &= c\int_0^a (x+1)\left[\left(1-\tfrac{x}{a}\right)\cdot b\left(1-\tfrac{x}{a}\right) - \tfrac{1}{2b}\cdot b^2\left(1-\tfrac{x}{a}\right)^2\right] dx = \tfrac{1}{2}bc\int_0^a (x+1)\left(1-\tfrac{x}{a}\right)^2 dx \\ &= \tfrac{1}{2}bc\int_0^a \left(\tfrac{1}{a^2}x^3 + \tfrac{1}{a^2}x^2 - \tfrac{2}{a}x^2 + x - \tfrac{2}{a}x + 1\right) dx \\ &= \tfrac{1}{2}bc\left[\tfrac{1}{4a^2}x^4 + \tfrac{1}{3a^2}x^3 - \tfrac{2}{3a}x^3 + \tfrac{1}{2}x^2 - \tfrac{1}{a}x^2 + x\right]_0^a \\ &= \tfrac{1}{2}bc\left(\tfrac{1}{4}a^2 + \tfrac{1}{3}a - \tfrac{2}{3}a^2 + \tfrac{1}{2}a^2 - a + a\right) = \tfrac{1}{2}bc\left(\tfrac{1}{12}a^2 + \tfrac{1}{3}a\right) = \tfrac{1}{24}abc(a+4)\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

11. $\operatorname{div}\mathbf{F} = 6x^2 + 3y^2 + 3y^2 = 6x^2 + 6y^2$ so

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 6(x^2+y^2)\,dV = \int_0^{2\pi}\int_0^1\int_0^{1-r^2} 6r^2\cdot r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^1 6r^3(1-r^2)\,dr\,d\theta$$
$$-\int_0^{2\pi} d\theta\int_0^1(6r^3-6r^5)\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{3}{2}r^4 - r^6\right]_0^1 = 2\pi\left(\tfrac{3}{2}-1\right) = \pi$$

12. For $x^2+y^2\le 4$ the plane $z = y-2$ is below the xy-plane, so the solid E bounded by S is $E = \{(x,y,z)\mid x^2+y^2\le 4,\ y-2\le z\le 0\}$. Here $\operatorname{div}\mathbf{F} = y+2z+2y-2z = 3y$ so

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 3y\,dV = \int_0^{2\pi}\int_0^2\int_{r\sin\theta-2}^0 (3r\sin\theta)\,r\,dz\,dr\,d\theta$$
$$= \int_0^{2\pi}\int_0^2 (3r^2\sin\theta)(0-r\sin\theta+2)\,dr\,d\theta = \int_0^{2\pi}\int_0^2\left(-3r^3\sin^2\theta+6r^2\sin\theta\right)dr\,d\theta$$
$$= \int_0^{2\pi}\left[-\tfrac{3}{4}r^4\sin^2\theta+2r^3\sin\theta\right]_{r=0}^{r=2} d\theta = \int_0^{2\pi}\left(-12\sin^2\theta+16\sin\theta\right)d\theta$$
$$= \left[-12\left(\tfrac{1}{2}\theta-\tfrac{1}{4}\sin 2\theta\right)-16\cos\theta\right]_0^{2\pi} = -12\pi-16+16 = -12\pi$$

13. $\mathbf{F}(x,y,z) = x\sqrt{x^2+y^2+z^2}\,\mathbf{i} + y\sqrt{x^2+y^2+z^2}\,\mathbf{j} + z\sqrt{x^2+y^2+z^2}\,\mathbf{k}$, so

$$\operatorname{div}\mathbf{F} = x\cdot\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2x) + (x^2+y^2+z^2)^{1/2} + y\cdot\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2y) + (x^2+y^2+z^2)^{1/2}$$
$$+ z\cdot\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2z) + (x^2+y^2+z^2)^{1/2}$$
$$= (x^2+y^2+z^2)^{-1/2}\left[x^2+(x^2+y^2+z^2)+y^2+(x^2+y^2+z^2)+z^2+(x^2+y^2+z^2)\right]$$
$$= \frac{4(x^2+y^2+z^2)}{\sqrt{x^2+y^2+z^2}} = 4\sqrt{x^2+y^2+z^2}.$$

Then
$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 4\sqrt{x^2+y^2+z^2}\,dV = \int_0^{\pi/2}\int_0^{2\pi}\int_0^1 4\sqrt{\rho^2}\cdot\rho^2\sin\phi\,d\rho\,d\theta\,d\phi$$
$$= \int_0^{\pi/2}\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_0^1 4\rho^3\,d\rho = \left[-\cos\phi\right]_0^{\pi/2}\left[\theta\right]_0^{2\pi}\left[\rho^4\right]_0^1 = (1)\,(2\pi)\,(1) = 2\pi$$

14. $\mathbf{F}(x,y,z) = x(x^2+y^2+z^2)\,\mathbf{i} + y(x^2+y^2+z^2)\,\mathbf{j} + z(x^2+y^2+z^2)\,\mathbf{k}$, so

$\operatorname{div}\mathbf{F} = x\cdot 2x + (x^2+y^2+z^2) + y\cdot 2y + (x^2+y^2+z^2) + z\cdot 2z + (x^2+y^2+z^2) = 5(x^2+y^2+z^2)$. Then

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 5(x^2+y^2+z^2)\,dV = \int_0^{\pi}\int_0^{2\pi}\int_0^R 5\rho^2\cdot\rho^2\sin\phi\,d\rho\,d\theta\,d\phi$$
$$= 5\int_0^{\pi}\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_0^R \rho^4\,d\rho = 5\left[-\cos\phi\right]_0^{\pi}\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{5}\rho^5\right]_0^R = 5\,(2)\,(2\pi)\left(\tfrac{1}{5}R^5\right) = 4\pi R^5$$

15. $\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E \sqrt{3-x^2}\,dV = \int_{-1}^1\int_{-1}^1\int_0^{2-x^4-y^4}\sqrt{3-x^2}\,dz\,dy\,dx = \frac{341}{60}\sqrt{2} + \frac{81}{20}\sin^{-1}\left(\frac{\sqrt{3}}{3}\right)$

16.





By the Divergence Theorem, the flux of $\mathbf{F}$ across the surface of the cube is

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \int_0^{\pi/2}\int_0^{\pi/2}\int_0^{\pi/2}\left[\cos x\,\cos^2 y + 3\sin^2 y\,\cos y\,\cos^4 z + 5\sin^4 z\,\cos z\,\cos^6 x\right]dz\,dy\,dx = \tfrac{19}{64}\pi^2.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. For S_1 we have $\mathbf{n} = -\mathbf{k}$, so $\mathbf{F} \cdot \mathbf{n} = \mathbf{F} \cdot (-\mathbf{k}) = -x^2 z - y^2 = -y^2$ (since $z = 0$ on S_1). So if D is the unit disk, we get $\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_1} \mathbf{F} \cdot \mathbf{n}\, dS = \iint_D (-y^2)\, dA = -\int_0^{2\pi} \int_0^1 r^2 (\sin^2\theta)\, r\, dr\, d\theta = -\frac{1}{4}\pi$. Now since S_2 is closed, we can use the Divergence Theorem. Since $\operatorname{div} \mathbf{F} = \frac{\partial}{\partial x}(z^2 x) + \frac{\partial}{\partial y}\left(\frac{1}{3}y^3 + \tan z\right) + \frac{\partial}{\partial z}(x^2 z + y^2) = z^2 + y^2 + x^2$, we use spherical coordinates to get $\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \iiint_E \operatorname{div} \mathbf{F}\, dV = \int_0^{2\pi} \int_0^{\pi/2} \int_0^1 \rho^2 \cdot \rho^2 \sin\phi\, d\rho\, d\phi\, d\theta = \frac{2}{5}\pi$. Finally $\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_{S_2} \mathbf{F} \cdot d\mathbf{S} - \iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \frac{2}{5}\pi - \left(-\frac{1}{4}\pi\right) = \frac{13}{20}\pi$.

18. As in the hint to Exercise 17, we create a closed surface $S_2 = S \cup S_1$, where S is the part of the paraboloid $x^2 + y^2 + z = 2$ that lies above the plane $z = 1$, and S_1 is the disk $x^2 + y^2 = 1$ on the plane $z = 1$ oriented downward, and we then apply the Divergence Theorem. Since the disk S_1 is oriented downward, its unit normal vector is $\mathbf{n} = -\mathbf{k}$ and $\mathbf{F} \cdot (-\mathbf{k}) = -z = -1$ on S_1. So $\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_1} \mathbf{F} \cdot \mathbf{n}\, dS = \iint_{S_1} (-1)\, dS = -A(S_1) = -\pi$. Let E be the region bounded by S_2. Then $\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \iiint_E \operatorname{div} \mathbf{F}\, dV = \iiint_E 1\, dV = \int_0^1 \int_0^{2\pi} \int_1^{2-r^2} r\, dz\, d\theta\, dr = \int_0^1 \int_0^{2\pi} (r - r^3)\, d\theta\, dr = (2\pi)\frac{1}{4} = \frac{\pi}{2}$. Thus the flux of $\mathbf{F}$ across S is $\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_{S_2} \mathbf{F} \cdot d\mathbf{S} - \iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \frac{\pi}{2} - (-\pi) = \frac{3\pi}{2}$.

19. The vectors that end near P_1 are longer than the vectors that start near P_1, so the net flow is inward near P_1 and $\operatorname{div} \mathbf{F}(P_1)$ is negative. The vectors that end near P_2 are shorter than the vectors that start near P_2, so the net flow is outward near P_2 and $\operatorname{div} \mathbf{F}(P_2)$ is positive.

20. (a) The vectors that end near P_1 are shorter than the vectors that start near P_1, so the net flow is outward and P_1 is a source. The vectors that end near P_2 are longer than the vectors that start near P_2, so the net flow is inward and P_2 is a sink.

(b) $\mathbf{F}(x, y) = \langle x, y^2 \rangle \Rightarrow \operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F} = 1 + 2y$. The y-value at P_1 is positive, so $\operatorname{div} \mathbf{F} = 1 + 2y$ is positive, thus P_1 is a source. At P_2, $y < -1$, so $\operatorname{div} \mathbf{F} = 1 + 2y$ is negative, and P_2 is a sink.

21.



From the graph it appears that for points above the x-axis, vectors starting near a particular point are longer than vectors ending there, so divergence is positive. The opposite is true at points below the x-axis, where divergence is negative.

$\mathbf{F}(x, y) = \langle xy, x + y^2 \rangle \Rightarrow \operatorname{div} \mathbf{F} = \frac{\partial}{\partial x}(xy) + \frac{\partial}{\partial y}(x + y^2) = y + 2y = 3y$.

Thus $\operatorname{div} \mathbf{F} > 0$ for $y > 0$, and $\operatorname{div} \mathbf{F} < 0$ for $y < 0$.

22.



From the graph it appears that for points above the line $y = -x$, vectors starting near a particular point are longer than vectors ending there, so divergence is positive. The opposite is true at points below the line $y = -x$, where divergence is negative.

$\mathbf{F}(x, y) = \langle x^2, y^2 \rangle \Rightarrow \operatorname{div} \mathbf{F} = \frac{\partial}{\partial x}(x^2) + \frac{\partial}{\partial y}(y^2) = 2x + 2y$. Then $\operatorname{div} \mathbf{F} > 0$ for $2x + 2y > 0 \Rightarrow y > -x$, and $\operatorname{div} \mathbf{F} < 0$ for $y < -x$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

23. Since $\dfrac{\mathbf{x}}{|\mathbf{x}|^3} = \dfrac{x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}}{(x^2+y^2+z^2)^{3/2}}$ and $\dfrac{\partial}{\partial x}\left(\dfrac{x}{(x^2+y^2+z^2)^{3/2}}\right) = \dfrac{(x^2+y^2+z^2)-3x^2}{(x^2+y^2+z^2)^{5/2}}$ with similar expressions for $\dfrac{\partial}{\partial y}\left(\dfrac{y}{(x^2+y^2+z^2)^{3/2}}\right)$ and $\dfrac{\partial}{\partial z}\left(\dfrac{z}{(x^2+y^2+z^2)^{3/2}}\right)$, we have

$$\operatorname{div}\left(\frac{\mathbf{x}}{|\mathbf{x}|^3}\right) = \frac{3(x^2+y^2+z^2)-3(x^2+y^2+z^2)}{(x^2+y^2+z^2)^{5/2}} = 0, \text{ except at } (0,0,0) \text{ where it is undefined.}$$

24. We first need to find $\mathbf{F}$ so that $\iint_S \mathbf{F}\cdot\mathbf{n}\,dS = \iint_S (2x+2y+z^2)\,dS$, so $\mathbf{F}\cdot\mathbf{n} = 2x+2y+z^2$. But for S,

$\mathbf{n} = \dfrac{x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}}{\sqrt{x^2+y^2+z^2}} = x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}$. Thus $\mathbf{F} = 2\,\mathbf{i}+2\,\mathbf{j}+z\,\mathbf{k}$ and $\operatorname{div}\mathbf{F} = 1$.

If $B = \{(x,y,z) \mid x^2+y^2+z^2 \le 1\}$, then $\iint_S (2x+2y+z^2)\,dS = \iiint_B dV = V(B) = \frac{4}{3}\pi(1)^3 = \frac{4}{3}\pi$.

25. $\iint_S \mathbf{a}\cdot\mathbf{n}\,dS = \iiint_E \operatorname{div}\mathbf{a}\,dV = 0$ since $\operatorname{div}\mathbf{a} = 0$.

26. $\frac{1}{3}\iint_S \mathbf{F}\cdot d\mathbf{S} = \frac{1}{3}\iiint_E \operatorname{div}\mathbf{F}\,dV = \frac{1}{3}\iiint_E 3\,dV = V(E)$

27. $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}(\operatorname{curl}\mathbf{F})\,dV = 0$ by Theorem 16.5.11.

28. $\iint_S D_{\mathbf{n}} f\,dS = \iint_S (\nabla f\cdot\mathbf{n})\,dS = \iiint_E \operatorname{div}(\nabla f)\,dV = \iiint_E \nabla^2 f\,dV$

29. $\iint_S (f\nabla g)\cdot\mathbf{n}\,dS = \iiint_E \operatorname{div}(f\nabla g)\,dV = \iiint_E (f\nabla^2 g + \nabla g\cdot\nabla f)\,dV$ by Exercise 16.5.25.

30. $\iint_S (f\nabla g - g\nabla f)\cdot\mathbf{n}\,dS = \iiint_E \left[(f\nabla^2 g + \nabla g\cdot\nabla f) - (g\nabla^2 f + \nabla g\cdot\nabla f)\right] dV$ [by Exercise 29].

But $\nabla g\cdot\nabla f = \nabla f\cdot\nabla g$, so that $\iint_S (f\nabla g - g\nabla f)\cdot\mathbf{n}\,dS = \iiint_E (f\nabla^2 g - g\nabla^2 f)\,dV$.

31. If $\mathbf{c} = c_1\,\mathbf{i}+c_2\,\mathbf{j}+c_3\,\mathbf{k}$ is an arbitrary constant vector, we define $\mathbf{F} = f\mathbf{c} = fc_1\,\mathbf{i}+fc_2\,\mathbf{j}+fc_3\,\mathbf{k}$. Then

$\operatorname{div}\mathbf{F} = \operatorname{div} f\mathbf{c} = \dfrac{\partial f}{\partial x}c_1 + \dfrac{\partial f}{\partial y}c_2 + \dfrac{\partial f}{\partial z}c_3 = \nabla f\cdot\mathbf{c}$ and the Divergence Theorem says $\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV \;\Rightarrow$

$\iint_S \mathbf{F}\cdot\mathbf{n}\,dS = \iiint_E \nabla f\cdot\mathbf{c}\,dV$. In particular, if $\mathbf{c} = \mathbf{i}$ then $\iint_S f\mathbf{i}\cdot\mathbf{n}\,dS = \iiint_E \nabla f\cdot\mathbf{i}\,dV \;\Rightarrow$

$\iint_S fn_1\,dS = \iiint_E \dfrac{\partial f}{\partial x}\,dV$ (where $\mathbf{n} = n_1\,\mathbf{i}+n_2\,\mathbf{j}+n_3\,\mathbf{k}$). Similarly, if $\mathbf{c} = \mathbf{j}$ we have $\iint_S fn_2\,dS = \iiint_E \dfrac{\partial f}{\partial y}\,dV$,

and $\mathbf{c} = \mathbf{k}$ gives $\iint_S fn_3\,dS = \iiint_E \dfrac{\partial f}{\partial z}\,dV$. Then

$$\begin{aligned}\iint_S f\mathbf{n}\,dS &= \left(\iint_S fn_1\,dS\right)\mathbf{i} + \left(\iint_S fn_2\,dS\right)\mathbf{j} + \left(\iint_S fn_3\,dS\right)\mathbf{k}\\ &= \left(\iiint_E \frac{\partial f}{\partial x}\,dV\right)\mathbf{i} + \left(\iiint_E \frac{\partial f}{\partial y}\,dV\right)\mathbf{j} + \left(\iiint_E \frac{\partial f}{\partial z}\,dV\right)\mathbf{k} = \iiint_E \left(\frac{\partial f}{\partial x}\,\mathbf{i} + \frac{\partial f}{\partial y}\,\mathbf{j} + \frac{\partial f}{\partial z}\,\mathbf{k}\right) dV\\ &= \iiint_E \nabla f\,dV \quad \text{as desired.}\end{aligned}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. By Exercise 31, $\iint_S p\mathbf{n}\,dS = \iiint_E \nabla p\,dV$, so

$$\mathbf{F} = -\iint_S p\mathbf{n}\,dS = -\iiint_E \nabla p\,dV = -\iiint_E \nabla(\rho g z)\,dV = -\iiint_E (\rho g\,\mathbf{k})\,dV = -\rho g\Big(\iiint_E dV\Big)\,\mathbf{k} = -\rho g V(E)\,\mathbf{k}.$$

But the weight of the displaced liquid is volume × density × $g = \rho g V(E)$, thus $\mathbf{F} = -W\mathbf{k}$ as desired.

16 Review

TRUE-FALSE QUIZ

1. False; div $\mathbf{F}$ is a scalar field.

2. True. (See Definition 16.5.1.)

3. True, by Theorem 16.5.3 and the fact that div $\mathbf{0} = 0$.

4. True, by Theorem 16.3.2.

5. False. See Exercise 16.3.35. (But the assertion is true if D is simply-connected; see Theorem 16.3.6.)

6. False. See the discussion accompanying Figure 8 on page 1120 [ET 1080].

7. False. For example, $\operatorname{div}(y\,\mathbf{i}) = 0 = \operatorname{div}(x\,\mathbf{j})$ but $y\,\mathbf{i} \neq x\,\mathbf{j}$.

8. True. Line integrals of conservative vector fields are independent of path, and by Theorem 16.3.3, work $= \int_C \mathbf{F}\cdot d\mathbf{r} = 0$ for any closed path C.

9. True. See Exercise 16.5.24.

10. False. $\mathbf{F}\cdot\mathbf{G}$ is a scalar field, so $\operatorname{curl}(\mathbf{F}\cdot\mathbf{G})$ has no meaning.

11. True. Apply the Divergence Theorem and use the fact that div $\mathbf{F} = 0$.

12. False by Theorem 16.5.11, because if it were true, then div curl $\mathbf{F} = 3 \neq 0$.

13. False. By Equations 16.4.5, the area is given by $-\oint_C y\,dx$ or $\oint_C x\,dy$.

EXERCISES

1. (a) Vectors starting on C point in roughly the direction opposite to C, so the tangential component $\mathbf{F}\cdot\mathbf{T}$ is negative. Thus $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \mathbf{F}\cdot\mathbf{T}\,ds$ is negative.

(b) The vectors that end near P are shorter than the vectors that start near P, so the net flow is outward near P and div $\mathbf{F}(P)$ is positive.

2. We can parametrize C by $x = x$, $y = x^2$, $0 \le x \le 1$ so

$$\int_C x\,ds = \int_0^1 x\sqrt{1+(2x)^2}\,dx = \tfrac{1}{12}(1+4x^2)^{3/2}\Big]_0^1 = \tfrac{1}{12}\big(5\sqrt{5}-1\big).$$

3. $\int_C yz\cos x\,ds = \int_0^\pi (3\cos t)\,(3\sin t)\cos t\,\sqrt{(1)^2+(-3\sin t)^2+(3\cos t)^2}\,dt = \int_0^\pi (9\cos^2 t\,\sin t)\sqrt{10}\,dt$

$$= 9\sqrt{10}\,\big(-\tfrac{1}{3}\cos^3 t\big)\Big]_0^\pi = -3\sqrt{10}\,(-2) = 6\sqrt{10}$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

4. $x = 3\cos t \;\Rightarrow\; dx = -3\sin t\,dt$, $y = 2\sin t \;\Rightarrow\; dy = 2\cos t\,dt$, $0 \le t \le 2\pi$, so

$$\begin{aligned}\int_C y\,dx + (x+y^2)\,dy &= \int_0^{2\pi}\left[(2\sin t)(-3\sin t) + (3\cos t + 4\sin^2 t)(2\cos t)\right]dt \\ &= \int_0^{2\pi}(-6\sin^2 t + 6\cos^2 t + 8\sin^2 t\,\cos t)\,dt = \int_0^{2\pi}\left[6(\cos^2 t - \sin^2 t) + 8\sin^2 t\,\cos t\right]dt \\ &= \int_0^{2\pi}(6\cos 2t + 8\sin^2 t\,\cos t)\,dt = 3\sin 2t + \tfrac{8}{3}\sin^3 t\Big]_0^{2\pi} = 0\end{aligned}$$

Or: Notice that $\frac{\partial}{\partial y}(y) = 1 = \frac{\partial}{\partial x}(x+y^2)$, so $\mathbf{F}(x,y) = \langle y, x+y^2\rangle$ is a conservative vector field. Since C is a closed curve, $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C y\,dx + (x+y^2)\,dy = 0$.

5. $\int_C y^3\,dx + x^2\,dy = \int_{-1}^{1}\left[y^3(-2y) + (1-y^2)^2\right]dy = \int_{-1}^{1}(-y^4 - 2y^2 + 1)\,dy$

$$= \left[-\tfrac{1}{5}y^5 - \tfrac{2}{3}y^3 + y\right]_{-1}^{1} = -\tfrac{1}{5} - \tfrac{2}{3} + 1 - \tfrac{1}{5} - \tfrac{2}{3} + 1 = \tfrac{4}{15}$$

6. $\int_C \sqrt{xy}\,dx + e^y\,dy + xz\,dz = \int_0^1\left(\sqrt{t^4\cdot t^2}\cdot 4t^3 + e^{t^2}\cdot 2t + t^4\cdot t^3\cdot 3t^2\right)dt = \int_0^1(4t^6 + 2te^{t^2} + 3t^9)\,dt$

$$= \left[\tfrac{4}{7}t^7 + e^{t^2} + \tfrac{3}{10}t^{10}\right]_0^1 = e - \tfrac{9}{70}$$

7. C: $x = 1+2t \;\Rightarrow\; dx = 2\,dt$, $y = 4t \;\Rightarrow\; dy = 4\,dt$, $z = -1+3t \;\Rightarrow\; dz = 3\,dt$, $0 \le t \le 1$.

$$\begin{aligned}\int_C xy\,dx + y^2\,dy + yz\,dz &= \int_0^1[(1+2t)(4t)(2) + (4t)^2(4) + (4t)(-1+3t)(3)]\,dt \\ &= \int_0^1(116t^2 - 4t)\,dt = \left[\tfrac{116}{3}t^3 - 2t^2\right]_0^1 = \tfrac{116}{3} - 2 = \tfrac{110}{3}\end{aligned}$$

8. $\mathbf{F}(\mathbf{r}(t)) = (\sin t)(1+t)\,\mathbf{i} + (\sin^2 t)\,\mathbf{j}$, $\mathbf{r}'(t) = \cos t\,\mathbf{i} + \mathbf{j}$ and

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \int_0^{\pi}((1+t)\sin t\cos t + \sin^2 t)\,dt = \int_0^{\pi}\left(\tfrac{1}{2}(1+t)\sin 2t + \sin^2 t\right)dt \\ &= \left[\tfrac{1}{2}\left((1+t)(-\tfrac{1}{2}\cos 2t) + \tfrac{1}{4}\sin 2t\right) + \tfrac{1}{2}t - \tfrac{1}{4}\sin 2t\right]_0^{\pi} = \tfrac{\pi}{4}\end{aligned}$$

9. $\mathbf{F}(\mathbf{r}(t)) = e^{-t}\,\mathbf{i} + t^2(-t)\,\mathbf{j} + (t^2+t^3)\,\mathbf{k}$, $\mathbf{r}'(t) = 2t\,\mathbf{i} + 3t^2\,\mathbf{j} - \mathbf{k}$ and

$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1(2te^{-t} - 3t^5 - (t^2+t^3))\,dt = \left[-2te^{-t} - 2e^{-t} - \frac{1}{2}t^6 - \frac{1}{3}t^3 - \frac{1}{4}t^4\right]_0^1 = \frac{11}{12} - \frac{4}{e}$.

10. (a) C: $x = 3-3t$, $y = \frac{\pi}{2}t$, $z = 3t$, $0 \le t \le 1$. Then

$$W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1\left[3t\,\mathbf{i} + (3-3t)\,\mathbf{j} + \tfrac{\pi}{2}t\,\mathbf{k}\right]\cdot\left[-3\,\mathbf{i} + \tfrac{\pi}{2}\,\mathbf{j} + 3\,\mathbf{k}\right]dt = \int_0^1\left[-9t + \tfrac{3\pi}{2}\right]dt = \tfrac{1}{2}(3\pi - 9).$$

(b) $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{\pi/2}(3\sin t\,\mathbf{i} + 3\cos t\,\mathbf{j} + t\,\mathbf{k})\cdot(-3\sin t\,\mathbf{i} + \mathbf{j} + 3\cos t\,\mathbf{k})\,dt$

$$= \int_0^{\pi/2}(-9\sin^2 t + 3\cos t + 3t\cos t)\,dt = \left[-9\left(\tfrac{1}{2}t - \tfrac{1}{4}\sin 2t\right) + 3\sin t + 3(t\sin t + \cos t)\right]_0^{\pi/2}$$

$$= \tfrac{9\pi}{4} + 3 + \tfrac{3\pi}{2} - 3 = \tfrac{3\pi}{4}$$

11. $\frac{\partial}{\partial y}\left[(1+xy)e^{xy}\right] = 2xe^{xy} + x^2ye^{xy} = \frac{\partial}{\partial x}\left[e^y + x^2e^{xy}\right]$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$, so $\mathbf{F}$ is conservative. Thus there exists a function f such that $\mathbf{F} = \nabla f$. Then $f_y(x,y) = e^y + x^2e^{xy}$ implies $f(x,y) = e^y + xe^{xy} + g(x)$ and then

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f_x(x,y) = xye^{xy} + e^{xy} + g'(x) = (1+xy)e^{xy} + g'(x)$. But $f_x(x,y) = (1+xy)e^{xy}$, so $g'(x) = 0 \Rightarrow g(x) = K$. Thus $f(x,y) = e^y + xe^{xy} + K$ is a potential function for $\mathbf{F}$.

12. $\mathbf{F}$ is defined on all of $\mathbb{R}^3$, its components have continuous partial derivatives, and curl $\mathbf{F} = (0-0)\,\mathbf{i} - (0-0)\,\mathbf{j} + (\cos y - \cos y)\,\mathbf{k} = \mathbf{0}$, so $\mathbf{F}$ is conservative by Theorem 16.5.4. Thus there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x,y,z) = \sin y$ implies $f(x,y,z) = x\sin y + g(y,z)$ and then $f_y(x,y,z) = x\cos y + g_y(y,z)$. But $f_y(x,y,z) = x\cos y$, so $g_y(y,z) = 0 \Rightarrow g(y,z) = h(z)$. Then $f(x,y,z) = x\sin y + h(z)$ implies $f_z(x,y,z) = h'(z)$. But $f_z(x,y,z) = -\sin z$, so $h(z) = \cos z + K$. Thus a potential function for $\mathbf{F}$ is $f(x,y,z) = x\sin y + \cos z + K$.

13. Since $\frac{\partial}{\partial y}(4x^3y^2 - 2xy^3) = 8x^3y - 6xy^2 = \frac{\partial}{\partial x}(2x^4y - 3x^2y^2 + 4y^3)$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$, $\mathbf{F}$ is conservative. Furthermore $f(x,y) = x^4y^2 - x^2y^3 + y^4$ is a potential function for $\mathbf{F}$. $t = 0$ corresponds to the point $(0,1)$ and $t = 1$ corresponds to $(1,1)$, so $\int_C \mathbf{F}\cdot d\mathbf{r} = f(1,1) - f(0,1) = 1 - 1 = 0$.

14. Here curl $\mathbf{F} = \mathbf{0}$, the domain of $\mathbf{F}$ is $\mathbb{R}^3$, and the components of $\mathbf{F}$ have continuous partial derivatives, so $\mathbf{F}$ is conservative. Furthermore $f(x,y,z) = xe^y + ye^z$ is a potential function for $\mathbf{F}$. Then $\int_C \mathbf{F}\cdot d\mathbf{r} = f(4,0,3) - f(0,2,0) = 4 - 2 = 2$.

15. C_1: $\mathbf{r}(t) = t\,\mathbf{i} + t^2\,\mathbf{j}$, $-1 \le t \le 1$;
C_2: $\mathbf{r}(t) = -t\,\mathbf{i} + \mathbf{j}$, $-1 \le t \le 1$.



Then

$$\int_C xy^2\,dx - x^2y\,dy = \int_{-1}^{1}(t^5 - 2t^5)\,dt + \int_{-1}^{1} t\,dt$$
$$= \left[-\tfrac{1}{6}t^6\right]_{-1}^{1} + \left[\tfrac{1}{2}t^2\right]_{-1}^{1} = 0$$

Using Green's Theorem, we have

$$\int_C xy^2\,dx - x^2y\,dy = \iint_D \left[\frac{\partial}{\partial x}(-x^2y) - \frac{\partial}{\partial y}(xy^2)\right] dA = \iint_D (-2xy - 2xy)\,dA = \int_{-1}^{1}\int_{x^2}^{1} -4xy\,dy\,dx$$
$$= \int_{-1}^{1}\left[-2xy^2\right]_{y=x^2}^{y=1} dx = \int_{-1}^{1}(2x^5 - 2x)\,dx = \left[\tfrac{1}{3}x^6 - x^2\right]_{-1}^{1} = 0$$

16. $\int_C \sqrt{1+x^3}\,dx + 2xy\,dy = \iint_D \left[\frac{\partial}{\partial x}(2xy) - \frac{\partial}{\partial y}\left(\sqrt{1+x^3}\right)\right] dA = \int_0^1\int_0^{3x}(2y-0)\,dy\,dx = \int_0^1 9x^2\,dx = 3x^3\Big]_0^1 = 3$

17. $\int_C x^2y\,dx - xy^2\,dy = \iint_{x^2+y^2\le 4}\left[\frac{\partial}{\partial x}(-xy^2) - \frac{\partial}{\partial y}(x^2y)\right] dA = \iint_{x^2+y^2\le 4}(-y^2 - x^2)\,dA = -\int_0^{2\pi}\int_0^2 r^3\,dr\,d\theta = -8\pi$

18. curl $\mathbf{F} = (0 - e^{-y}\cos z)\,\mathbf{i} - (e^{-z}\cos x - 0)\,\mathbf{j} + (0 - e^{-x}\cos y)\,\mathbf{k} = -e^{-y}\cos z\,\mathbf{i} - e^{-z}\cos x\,\mathbf{j} - e^{-x}\cos y\,\mathbf{k}$,
div $\mathbf{F} = -e^{-x}\sin y - e^{-y}\sin z - e^{-z}\sin x$

19. If we assume there is such a vector field $\mathbf{G}$, then $\operatorname{div}(\operatorname{curl}\mathbf{G}) = 2 + 3z - 2xz$. But $\operatorname{div}(\operatorname{curl}\mathbf{F}) = 0$ for all vector fields $\mathbf{F}$. Thus such a $\mathbf{G}$ cannot exist.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

20. Let $\mathbf{F} = P_1\,\mathbf{i} + Q_1\,\mathbf{j} + R_1\,\mathbf{k}$ and $\mathbf{G} = P_2\,\mathbf{i} + Q_2\,\mathbf{j} + R_2\,\mathbf{k}$ be vector fields whose first partials exist and are continuous. Then

$$\begin{aligned}\mathbf{F}\,\mathrm{div}\,\mathbf{G} - \mathbf{G}\,\mathrm{div}\,\mathbf{F} = {} & \left[P_1\left(\frac{\partial P_2}{\partial x} + \frac{\partial Q_2}{\partial y} + \frac{\partial R_2}{\partial z}\right)\mathbf{i} + Q_1\left(\frac{\partial P_2}{\partial x} + \frac{\partial Q_2}{\partial y} + \frac{\partial R_2}{\partial z}\right)\mathbf{j} + R_1\left(\frac{\partial P_2}{\partial x} + \frac{\partial Q_2}{\partial y} + \frac{\partial R_2}{\partial z}\right)\mathbf{k}\right] \\ & - \left[P_2\left(\frac{\partial P_1}{\partial x} + \frac{\partial Q_1}{\partial y} + \frac{\partial R_1}{\partial z}\right)\mathbf{i} + Q_2\left(\frac{\partial P_1}{\partial x} + \frac{\partial Q_1}{\partial y} + \frac{\partial R_1}{\partial z}\right)\mathbf{j}\right. \\ & \left. + R_2\left(\frac{\partial P_1}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R_1}{\partial z}\right)\mathbf{k}\right]\end{aligned}$$

and

$$\begin{aligned}(\mathbf{G}\cdot\nabla)\,\mathbf{F} - (\mathbf{F}\cdot\nabla)\,\mathbf{G} = {} & \left[\left(P_2\frac{\partial P_1}{\partial x} + Q_2\frac{\partial P_1}{\partial y} + R_2\frac{\partial P_1}{\partial z}\right)\mathbf{i} + \left(P_2\frac{\partial Q_1}{\partial x} + Q_2\frac{\partial Q_1}{\partial y} + R_2\frac{\partial Q_1}{\partial z}\right)\mathbf{j}\right. \\ & \left. + \left(P_2\frac{\partial R_1}{\partial x} + Q_2\frac{\partial R_1}{\partial y} + R_2\frac{\partial R_1}{\partial z}\right)\mathbf{k}\right] \\ & - \left[\left(P_1\frac{\partial P_2}{\partial x} + Q_1\frac{\partial P_2}{\partial y} + R_1\frac{\partial P_2}{\partial z}\right)\mathbf{i} + \left(P_1\frac{\partial Q_2}{\partial x} + Q_1\frac{\partial Q_2}{\partial y} + R_1\frac{\partial Q_2}{\partial z}\right)\mathbf{j}\right. \\ & \left. + \left(P_1\frac{\partial R_2}{\partial x} + Q_1\frac{\partial R_2}{\partial y} + R_1\frac{\partial R_2}{\partial z}\right)\mathbf{k}\right]\end{aligned}$$

Hence

$$\begin{aligned}&\mathbf{F}\,\mathrm{div}\,\mathbf{G} - \mathbf{G}\,\mathrm{div}\,\mathbf{F} + (\mathbf{G}\cdot\nabla)\,\mathbf{F} - (\mathbf{F}\cdot\nabla)\,\mathbf{G} \\ &= \left[\left(P_1\frac{\partial Q_2}{\partial y} + Q_2\frac{\partial P_1}{\partial x}\right) - \left(P_2\frac{\partial Q_1}{\partial y} + Q_1\frac{\partial P_2}{\partial y}\right)\right. \\ &\qquad \left. - \left(P_2\frac{\partial R_1}{\partial z} + R_1\frac{\partial P_2}{\partial z}\right) + \left(P_1\frac{\partial R_2}{\partial z} + R_2\frac{\partial P_1}{\partial z}\right)\right]\mathbf{i} \\ &\quad + \left[\left(Q_1\frac{\partial R_2}{\partial z} + R_2\frac{\partial Q_1}{\partial z}\right) - \left(Q_2\frac{\partial R_1}{\partial z} + R_1\frac{\partial Q_2}{\partial z}\right)\right. \\ &\qquad \left. - \left(P_1\frac{\partial Q_2}{\partial x} + Q_2\frac{\partial P_1}{\partial x}\right) + \left(P_2\frac{\partial Q_1}{\partial x} + Q_1\frac{\partial P_2}{\partial x}\right)\right]\mathbf{j} \\ &\quad + \left[\left(P_2\frac{\partial R_1}{\partial x} + R_1\frac{\partial P_2}{\partial x}\right) - \left(P_1\frac{\partial R_2}{\partial x} + R_2\frac{\partial P_1}{\partial x}\right)\right. \\ &\qquad \left. - \left(Q_1\frac{\partial R_2}{\partial y} + R_2\frac{\partial Q_1}{\partial y}\right) + \left(Q_2\frac{\partial R_1}{\partial y} + R_1\frac{\partial Q_2}{\partial y}\right)\right]\mathbf{k} \\ &= \left[\frac{\partial}{\partial y}(P_1Q_2 - P_2Q_1) - \frac{\partial}{\partial z}(P_2R_1 - P_1R_2)\right]\mathbf{i} \\ &\quad + \left[\frac{\partial}{\partial z}(Q_1R_2 - Q_2R_1) - \frac{\partial}{\partial x}(P_1Q_2 - P_2Q_1)\right]\mathbf{j} \\ &\quad + \left[\frac{\partial}{\partial x}(P_2R_1 - P_1R_2) - \frac{\partial}{\partial y}(Q_1R_2 - Q_2R_1)\right]\mathbf{k} \\ &= \mathrm{curl}\,(\mathbf{F}\times\mathbf{G})\end{aligned}$$

21. For any piecewise-smooth simple closed plane curve C bounding a region D, we can apply Green's Theorem to $\mathbf{F}(x,y) = f(x)\,\mathbf{i} + g(y)\,\mathbf{j}$ to get $\int_C f(x)\,dx + g(y)\,dy = \iint_D \left[\frac{\partial}{\partial x}\,g(y) - \frac{\partial}{\partial y}\,f(x)\right]dA = \iint_D 0\,dA = 0$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

22. $\nabla^2(fg) = \dfrac{\partial^2(fg)}{\partial x^2} + \dfrac{\partial^2(fg)}{\partial y^2} + \dfrac{\partial^2(fg)}{\partial z^2}$

$$= \frac{\partial}{\partial x}\left(\frac{\partial f}{\partial x}\,g + f\,\frac{\partial g}{\partial x}\right) + \frac{\partial}{\partial y}\left(\frac{\partial f}{\partial y}\,g + f\,\frac{\partial g}{\partial y}\right) + \frac{\partial}{\partial z}\left(\frac{\partial f}{\partial z}\,g + f\,\frac{\partial g}{\partial z}\right) \quad \text{[Product Rule]}$$

$$= \frac{\partial^2 f}{\partial x^2}\,g + 2\,\frac{\partial f}{\partial x}\frac{\partial g}{\partial x} + f\,\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}\,g + 2\,\frac{\partial f}{\partial y}\frac{\partial g}{\partial y}$$

$$+ f\,\frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}\,g + 2\,\frac{\partial f}{\partial z}\frac{\partial g}{\partial z} + f\,\frac{\partial^2 g}{\partial z^2} \quad \text{[Product Rule]}$$

$$= f\left(\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 g}{\partial z^2}\right) + g\left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}\right) + 2\left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}\right\rangle \cdot \left\langle \frac{\partial g}{\partial x}, \frac{\partial g}{\partial y}, \frac{\partial g}{\partial z}\right\rangle$$

$$= f\nabla^2 g + g\nabla^2 f + 2\nabla f \cdot \nabla g$$

Another method: Using the rules in Exercises 14.6.37(b) and 16.5.25, we have

$$\nabla^2(fg) = \nabla \cdot \nabla(fg) = \nabla \cdot (g\,\nabla f + f\,\nabla g) = \nabla g \cdot \nabla f + g\nabla \cdot \nabla f + \nabla f \cdot \nabla g + f\nabla \cdot \nabla g$$
$$= g\,\nabla^2 f + f\,\nabla^2 g + 2\nabla f \cdot \nabla g$$

23. $\nabla^2 f = 0$ means that $\dfrac{\partial^2 f}{\partial x^2} + \dfrac{\partial^2 f}{\partial y^2} = 0$. Now if $\mathbf{F} = f_y\,\mathbf{i} - f_x\,\mathbf{j}$ and C is any closed path in D, then applying Green's Theorem, we get

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C f_y\,dx - f_x\,dy = \iint_D \left[\tfrac{\partial}{\partial x}\left(-f_x\right) - \tfrac{\partial}{\partial y}\left(f_y\right)\right] dA$$
$$= -\iint_D (f_{xx} + f_{yy})\,dA = -\iint_D 0\,dA = 0$$

Therefore the line integral is independent of path, by Theorem 16.3.3.

24. (a) $x^2 + y^2 = \cos^2 t + \sin^2 t = 1$, so C lies on the circular cylinder $x^2 + y^2 = 1$. But also $y = z$, so C lies on the plane $y = z$. Thus C is contained in the intersection of the plane $y = z$ and the cylinder $x^2 + y^2 = 1$; with $0 \le t \le 2\pi$ we get the entire intersection (an ellipse).



(b) Apply Stokes' Theorem, $\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S}$:

$$\operatorname{curl}\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ 2xe^{2y} & 2x^2e^{2y} + 2y\cot z & -y^2\csc^2 z \end{vmatrix} = \left\langle -2y\csc^2 z - (-2y\csc^2 z), 0, 4xe^{2y} - 4xe^{2y}\right\rangle = \mathbf{0}$$

Therefore $\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \mathbf{0}\cdot d\mathbf{S} = 0$.

25. $z = f(x, y) = x^2 + 2y$ with $0 \le x \le 1$, $0 \le y \le 2x$. Thus

$$A(S) = \iint_D \sqrt{1 + 4x^2 + 4}\,dA = \int_0^1\int_0^{2x} \sqrt{5 + 4x^2}\,dy\,dx = \int_0^1 2x\sqrt{5 + 4x^2}\,dx = \tfrac{1}{6}(5 + 4x^2)^{3/2}\Big]_0^1 = \tfrac{1}{6}(27 - 5\sqrt{5}).$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

26. (a) $\mathbf{r}_u = -v\,\mathbf{j} + 2u\,\mathbf{k}$, $\mathbf{r}_v = 2v\,\mathbf{i} - u\,\mathbf{j}$ and $\mathbf{r}_u \times \mathbf{r}_v = 2u^2\,\mathbf{i} + 4uv\,\mathbf{j} + 2v^2\,\mathbf{k}$. Since the point $(4,-2,1)$ corresponds to $u=1, v=2$ (or $u=-1, v=-2$ but $\mathbf{r}_u \times \mathbf{r}_v$ is the same for both), a normal vector to the surface at $(4,-2,1)$ is $2\,\mathbf{i} + 8\,\mathbf{j} + 8\,\mathbf{k}$ and an equation of the tangent plane is $2x + 8y + 8z = 0$ or $x + 4y + 4z = 0$.

(b)



(c) By Definition 16.6.6, the area of S is given by

$$A(S) = \iint_D |\,\mathbf{r}_u \times \mathbf{r}_v\,|\,dA = \int_0^3\int_{-3}^3 \sqrt{(2u^2)^2 + (4uv)^2 + (2v^2)^2}\,dv\,du = 2\int_0^3\int_{-3}^3 \sqrt{u^4 + 4u^2v^2 + v^4}\,dv\,du.$$

(d) By Equation 16.7.9, the surface integral is

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_D \mathbf{F}\cdot(\mathbf{r}_u \times \mathbf{r}_v)\,dA = \int_0^3\int_{-3}^3 \left\langle \frac{(u^2)^2}{1+(v^2)^2}, \frac{(v^2)^2}{1+(-uv)^2}, \frac{(-uv)^2}{1+(u^2)^2} \right\rangle \cdot \langle 2u^2, 4uv, 2v^2\rangle\,dv\,du$$

$$= \int_0^3\int_{-3}^3 \left(\frac{2u^6}{1+v^4} + \frac{4uv^5}{1+u^2v^2} + \frac{2u^2v^4}{1+u^4}\right) dv\,du \approx 1524.0190$$

27. $z = f(x,y) = x^2 + y^2$ with $0 \le x^2 + y^2 \le 4$ so $\mathbf{r}_x \times \mathbf{r}_y = -2x\,\mathbf{i} - 2y\,\mathbf{j} + \mathbf{k}$. Then

$$\iint_S z\,dS = \iint_{x^2+y^2\le 4} (x^2+y^2)\sqrt{4x^2+4y^2+1}\,dA$$

$$= \int_0^{2\pi}\int_0^2 r^3\sqrt{1+4r^2}\,dr\,d\theta = \tfrac{1}{60}\pi\left(391\sqrt{17}+1\right)$$

(Substitute $u = 1 + 4r^2$ and use tables.)

28. $z = f(x,y) = 4 + x + y$ with $0 \le x^2 + y^2 \le 4$ so $\mathbf{r}_x \times \mathbf{r}_y = -\mathbf{i} - \mathbf{j} + \mathbf{k}$. Then

$$\iint_S (x^2z + y^2z)\,dS = \iint_{x^2+y^2<4} (x^2+y^2)(4+x+y)\sqrt{3}\,dA$$

$$= \int_0^2\int_0^{2\pi} \sqrt{3}\,r^3(4 + r\cos\theta + r\sin\theta)\,d\theta\,dr = \int_0^2 8\pi\sqrt{3}\,r^3\,dr = 32\pi\sqrt{3}$$

29. Since the sphere bounds a simple solid region, the Divergence Theorem applies and

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV = \iiint_E (z-2)\,dV = \iiint_E z\,dV - 2\iiint_E dV$$

$$= 0 \left[\begin{array}{l}\text{odd function in } z\\ \text{and } E \text{ is symmetric}\end{array}\right] - 2\cdot V(E) = -2\cdot\tfrac{4}{3}\pi(2)^3 = -\tfrac{64}{3}\pi$$

Alternate solution: $\mathbf{F}(\mathbf{r}(\phi,\theta)) = 4\sin\phi\cos\theta\cos\phi\,\mathbf{i} - 4\sin\phi\sin\theta\,\mathbf{j} + 6\sin\phi\cos\theta\,\mathbf{k}$,

$\mathbf{r}_\phi \times \mathbf{r}_\theta = 4\sin^2\phi\cos\theta\,\mathbf{i} + 4\sin^2\phi\sin\theta\,\mathbf{j} + 4\sin\phi\cos\phi\,\mathbf{k}$, and

$\mathbf{F}\cdot(\mathbf{r}_\phi \times \mathbf{r}_\theta) = 16\sin^3\phi\cos^2\theta\cos\phi - 16\sin^3\phi\sin^2\theta + 24\sin^2\phi\cos\phi\cos\theta$. Then

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \int_0^{2\pi}\int_0^{\pi} (16\sin^3\phi\cos\phi\cos^2\theta - 16\sin^3\phi\sin^2\theta + 24\sin^2\phi\cos\phi\cos\theta)\,d\phi\,d\theta$$

$$= \int_0^{2\pi} \tfrac{4}{3}(-16\sin^2\theta)\,d\theta = -\tfrac{64}{3}\pi$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $z = f(x,y) = x^2 + y^2$, $\mathbf{r}_x \times \mathbf{r}_y = -2x\,\mathbf{i} - 2y\,\mathbf{j} + \mathbf{k}$ (because of upward orientation) and $\mathbf{F}(\mathbf{r}(x,y)) \cdot (\mathbf{r}_x \times \mathbf{r}_y) = -2x^3 - 2xy^2 + x^2 + y^2$. Then

$$\begin{aligned}\iint_S \mathbf{F} \cdot d\mathbf{S} &= \iint_{x^2+y^2\le 1} (-2x^3 - 2xy^2 + x^2 + y^2)\,dA \\ &= \int_0^1 \int_0^{2\pi} (-2r^3\cos^3\theta - 2r^3\cos\theta\,\sin^2\theta + r^2)\,r\,dr\,d\theta = \int_0^1 r^3(2\pi)\,dr = \tfrac{\pi}{2}\end{aligned}$$

31. Since $\operatorname{curl}\mathbf{F} = \mathbf{0}$, $\iint_S (\operatorname{curl}\mathbf{F}) \cdot d\mathbf{S} = 0$. We parametrize C: $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j}$, $0 \le t \le 2\pi$ and

$\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} (-\cos^2 t\,\sin t + \sin^2 t\,\cos t)\,dt = \frac{1}{3}\cos^3 t + \frac{1}{3}\sin^3 t\,\big]_0^{2\pi} = 0.$

32. $\iint_S \operatorname{curl}\mathbf{F} \cdot d\mathbf{S} = \oint_C \mathbf{F} \cdot d\mathbf{r}$ where C: $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} + \mathbf{k}$, $0 \le t \le 2\pi$, so $\mathbf{r}'(t) = -2\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$, $\mathbf{F}(\mathbf{r}(t)) = 8\cos^2 t\,\sin t\,\mathbf{i} + 2\sin t\,\mathbf{j} + e^{4\cos t \sin t}\,\mathbf{k}$, and $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = -16\cos^2 t \sin^2 t + 4\sin t\cos t$. Thus

$\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} (-16\cos^2 t\,\sin^2 t + 4\sin t\,\cos t)\,dt = \big[-16\big(-\frac{1}{4}\sin t\,\cos^3 t + \frac{1}{16}\sin 2t + \frac{1}{8}t\big) + 2\sin^2 t\big]_0^{2\pi} = -4\pi.$

33. The surface is given by $x + y + z = 1$ or $z = 1 - x - y$, $0 \le x \le 1$, $0 \le y \le 1 - x$ and $\mathbf{r}_x \times \mathbf{r}_y = \mathbf{i} + \mathbf{j} + \mathbf{k}$. Then

$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F} \cdot d\mathbf{S} = \iint_D (-y\,\mathbf{i} - z\,\mathbf{j} - x\,\mathbf{k}) \cdot (\mathbf{i} + \mathbf{j} + \mathbf{k})\,dA = \iint_D (-1)\,dA = -(\text{area of } D) = -\frac{1}{2}.$

34. $\iint_S \mathbf{F} \cdot d\mathbf{S} = \iiint_E 3(x^2 + y^2 + z^2)\,dV = \int_0^{2\pi}\int_0^1\int_0^2 (3r^2 + 3z^2)\,r\,dz\,dr\,d\theta = 2\pi\int_0^1 (6r^3 + 8r)\,dr = 11\pi$

35. $\iiint_E \operatorname{div}\mathbf{F}\,dV = \iiint_{x^2+y^2+z^2<1} 3\,dV = 3(\text{volume of sphere}) = 4\pi$. Then

$\mathbf{F}(\mathbf{r}(\phi,\theta)) \cdot (\mathbf{r}_\phi \times \mathbf{r}_\theta) = \sin^3\phi\,\cos^2\theta + \sin^3\phi\,\sin^2\theta + \sin\phi\,\cos^2\phi = \sin\phi$ and

$\iint_S \mathbf{F} \cdot d\mathbf{S} = \int_0^{2\pi}\int_0^{\pi} \sin\phi\,d\phi\,d\theta = (2\pi)(2) = 4\pi.$

36. Here we must use Equation 16.9.7 since $\mathbf{F}$ is not defined at the origin. Let S_1 be the sphere of radius 1 with center at the origin and outer unit normal $\mathbf{n}_1$. Let S_2 be the surface of the ellipsoid with outer unit normal $\mathbf{n}_2$ and let E be the solid region between S_1 and S_2. Then the outward flux of $\mathbf{F}$ through the ellipsoid is given by

$\iint_{S_2} \mathbf{F} \cdot \mathbf{n}_2\,dS = -\iint_{S_1} \mathbf{F} \cdot (-\mathbf{n}_1)\,dS + \iiint_E \operatorname{div}\mathbf{F}\,dV$. But $\mathbf{F} = \mathbf{r}/|\mathbf{r}|^3$, so

$\operatorname{div}\mathbf{F} = \nabla \cdot \left(|\mathbf{r}|^{-3}\,\mathbf{r}\right) = |\mathbf{r}|^{-3}(\nabla \cdot \mathbf{r}) + \mathbf{r} \cdot \left(\nabla\,|\mathbf{r}|^{-3}\right) = |\mathbf{r}|^{-3}\,(3) + \mathbf{r} \cdot \left(-3\,|\mathbf{r}|^{-4}\right)\left(\mathbf{r}\,|\mathbf{r}|^{-1}\right) = 0$. [Here we have used Exercises 16.5.30(a) and 16.5.31(a).] And $\mathbf{F} \cdot \mathbf{n}_1 = \dfrac{\mathbf{r}}{|\mathbf{r}|^3} \cdot \dfrac{\mathbf{r}}{|\mathbf{r}|} = |\mathbf{r}|^{-2} = 1$ on S_1.

Thus $\iint_{S_2} \mathbf{F} \cdot \mathbf{n}_2\,dS = \iint_{S_1} dS + \iiint_E 0\,dV = (\text{surface area of the unit sphere}) = 4\pi(1)^2 = 4\pi$.

37. Because $\operatorname{curl}\mathbf{F} = \mathbf{0}$, $\mathbf{F}$ is conservative, so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x,y,z) = 3x^2yz - 3y$ implies $f(x,y,z) = x^3yz - 3xy + g(y,z) \quad \Rightarrow \quad f_y(x,y,z) = x^3z - 3x + g_y(y,z)$. But $f_y(x,y,z) = x^3z - 3x$, so $g(y,z) = h(z)$ and $f(x,y,z) = x^3yz - 3xy + h(z)$. Then $f_z(x,y,z) = x^3y + h'(z)$ but $f_z(x,y,z) = x^3y + 2z$, so $h(z) = z^2 + K$ and a potential function for $\mathbf{F}$ is $f(x,y,z) = x^3yz - 3xy + z^2$. Hence

$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(0,3,0) - f(0,0,2) = 0 - 4 = -4.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

38. Let C' be the circle with center at the origin and radius a as in the figure. Let D be the region bounded by C and C'. Then D's positively oriented boundary is $C \cup (-C')$. Hence by Green's Theorem



$$\int_C \mathbf{F}\cdot d\mathbf{r} + \int_{-C'} \mathbf{F}\cdot d\mathbf{r} = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA = 0, \text{ so}$$

$$\int_C \mathbf{F}\cdot d\mathbf{r} = -\int_{-C'} \mathbf{F}\cdot d\mathbf{r} = \int_{C'} \mathbf{F}\cdot dr = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t))\cdot \mathbf{r}'(t)\,dt$$

$$= \int_0^{2\pi} \left[\frac{2a^3\cos^3 t + 2a^3\cos t\,\sin^2 t - 2a\sin t}{a^2}(-a\sin t) + \frac{2a^3\sin^3 t + 2a^3\cos^2 t\,\sin t + 2a\cos t}{a^2}(a\cos t)\right] dt$$

$$= \int_0^{2\pi} \frac{2a^2}{a^2}\,dt = 4\pi$$

39. By the Divergence Theorem, $\iint_S \mathbf{F}\cdot\mathbf{n}\,dS = \iiint_E \operatorname{div}\mathbf{F}\,dV = 3(\text{volume of } E) = 3(8-1) = 21$.

40. The stated conditions allow us to use the Divergence Theorem. Hence $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}(\operatorname{curl}\mathbf{F})\,dV = 0$ since $\operatorname{div}(\operatorname{curl}\mathbf{F}) = 0$.

41. Let $\mathbf{F} = \mathbf{a}\times\mathbf{r} = \langle a_1, a_2, a_3\rangle \times \langle x, y, z\rangle = \langle a_2 z - a_3 y, a_3 x - a_1 z, a_1 y - a_2 x\rangle$. Then $\operatorname{curl}\mathbf{F} = \langle 2a_1, 2a_2, 2a_3\rangle = 2\mathbf{a}$, and $\iint_S 2\mathbf{a}\cdot d\mathbf{S} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_C (\mathbf{a}\times\mathbf{r})\cdot d\mathbf{r}$ by Stokes' Theorem.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

PROBLEMS PLUS

1. Let S_1 be the portion of $\Omega(S)$ between $S(a)$ and S, and let ∂S_1 be its boundary. Also let S_L be the lateral surface of S_1 [that is, the surface of S_1 except S and $S(a)$]. Applying the Divergence Theorem we have $\iint_{\partial S_1} \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iiint_{S_1} \nabla\cdot\dfrac{\mathbf{r}}{r^3}\,dV$.

But
$$\nabla\cdot\frac{\mathbf{r}}{r^3} = \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}\right\rangle \cdot \left\langle \frac{x}{(x^2+y^2+z^2)^{3/2}}, \frac{y}{(x^2+y^2+z^2)^{3/2}}, \frac{z}{(x^2+y^2+z^2)^{3/2}}\right\rangle$$
$$= \frac{(x^2+y^2+z^2-3x^2)+(x^2+y^2+z^2-3y^2)+(x^2+y^2+z^2-3z^2)}{(x^2+y^2+z^2)^{5/2}} = 0$$

$\Rightarrow \iint_{\partial S_1} \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iiint_{S_1} 0\,dV = 0$. On the other hand, notice that for the surfaces of ∂S_1 other than $S(a)$ and S, $\mathbf{r}\cdot\mathbf{n} = 0 \Rightarrow$

$0 = \iint_{\partial S_1} \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iint_S \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS + \iint_{S(a)} \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS + \iint_{S_L} \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iint_S \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS + \iint_{S(a)} \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS \Rightarrow$

$\iint_S \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = -\iint_{S(a)} \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS$. Notice that on $S(a)$, $r = a \Rightarrow \mathbf{n} = -\dfrac{\mathbf{r}}{r} = -\dfrac{\mathbf{r}}{a}$ and $\mathbf{r}\cdot\mathbf{r} = r^2 = a^2$, so that $-\iint_{S(a)} \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iint_{S(a)} \dfrac{\mathbf{r}\cdot\mathbf{r}}{a^4}\,dS = \iint_{S(a)} \dfrac{a^2}{a^4}\,dS = \dfrac{1}{a^2}\iint_{S(a)} dS = \dfrac{\text{area of } S(a)}{a^2} = |\Omega(S)|$.

Therefore $|\Omega(S)| = \iint_S \dfrac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS$.

2. By Green's Theorem
$$\int_C (y^3 - y)\,dx - 2x^3\,dy = \iint_D \left[\frac{\partial(-2x^3)}{\partial x} - \frac{\partial(y^3-y)}{\partial y}\right] dA = \iint_D (1 - 6x^2 - 3y^2)\,dA$$

Notice that for $6x^2 + 3y^2 > 1$, the integrand is negative. The integral has maximum value if it is evaluated only in the region where the integrand is positive, which is within the ellipse $6x^2 + 3y^2 = 1$. So the simple closed curve that gives a maximum value for the line integral is the ellipse $6x^2 + 3y^2 = 1$.

3. The given line integral $\frac{1}{2}\int_C (bz - cy)\,dx + (cx - az)\,dy + (ay - bx)\,dz$ can be expressed as $\int_C \mathbf{F}\cdot d\mathbf{r}$ if we define the vector field $\mathbf{F}$ by $\mathbf{F}(x,y,z) = P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k} = \frac{1}{2}(bz - cy)\,\mathbf{i} + \frac{1}{2}(cx - az)\,\mathbf{j} + \frac{1}{2}(ay - bx)\,\mathbf{k}$. Then define S to be the planar interior of C, so S is an oriented, smooth surface. Stokes' Theorem says $\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,dS$.

Now
$$\operatorname{curl}\mathbf{F} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}\right)\mathbf{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x}\right)\mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right)\mathbf{k}$$
$$= \left(\tfrac{1}{2}a + \tfrac{1}{2}a\right)\mathbf{i} + \left(\tfrac{1}{2}b + \tfrac{1}{2}b\right)\mathbf{j} + \left(\tfrac{1}{2}c + \tfrac{1}{2}c\right)\mathbf{k} = a\,\mathbf{i} + b\,\mathbf{j} + c\,\mathbf{k} = \mathbf{n}$$

so $\operatorname{curl}\mathbf{F}\cdot\mathbf{n} = \mathbf{n}\cdot\mathbf{n} = |\mathbf{n}|^2 = 1$, hence $\iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,dS = \iint_S dS$ which is simply the surface area of S. Thus, $\int_C \mathbf{F}\cdot d\mathbf{r} = \frac{1}{2}\int_C (bz - cy)\,dx + (cx - az)\,dy + (ay - bx)\,dz$ is the plane area enclosed by C.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. The surface given by $x = \sin u$, $y = \sin v$, $z = \sin(u + v)$ is difficult to visualize, so we first graph the surface from three different points of view.



The trace in the horizontal plane $z = 0$ is given by $z = \sin(u + v) = 0 \quad \Rightarrow \quad u + v = k\pi$ [k an integer]. Then we can write $v = k\pi - u$, and the trace is given by the parametric equations $x = \sin u$, $y = \sin v = \sin(k\pi - u) = \sin k\pi \cos u - \cos k\pi \sin u = \pm \sin u$, and since $\sin u = x$, the trace consists of the two lines $y = \pm x$.

If $z = 1$, $z = \sin(u + v) = 1 \quad \Rightarrow \quad u + v = \frac{\pi}{2} + 2k\pi$. So $v = \left(\frac{\pi}{2} + 2k\pi\right) - u$ and the trace in $z = 1$ is given by the parametric equations $x = \sin u$, $y = \sin v = \sin\left(\left(\frac{\pi}{2} + 2k\pi\right) - u\right) = \sin\left(\frac{\pi}{2} + 2k\pi\right)\cos u - \cos\left(\frac{\pi}{2} + 2k\pi\right)\sin u = \cos u$. This curve is equivalent to $x^2 + y^2 = 1$, $z = 1$, a circle of radius 1. Similarly, in $z = -1$ we have $z = \sin(u + v) = -1 \quad \Rightarrow$ $u + v = \frac{3\pi}{2} + 2k\pi \quad \Rightarrow \quad v = \left(\frac{3\pi}{2} + 2k\pi\right) - u$, so the trace is given by the parametric equations $x = \sin u$, $y = \sin v = \sin\left(\left(\frac{3\pi}{2} + 2k\pi\right) - u\right) = \sin\left(\frac{3\pi}{2} + 2k\pi\right)\cos u - \cos\left(\frac{3\pi}{2} + 2k\pi\right)\sin u = -\cos u$, which again is a circle, $x^2 + y^2 = 1$, $z = -1$.

If $z = \frac{1}{2}$, $z = \sin(u + v) = \frac{1}{2} \quad \Rightarrow \quad u + v = \alpha + 2k\pi$ where $\alpha = \frac{\pi}{6}$ or $\frac{5\pi}{6}$. Then $v = (\alpha + 2k\pi) - u$ and the trace in $z = \frac{1}{2}$ is given by the parametric equations $x = \sin u$, $y = \sin v = \sin[(\alpha + 2k\pi) - u] = \sin(\alpha + 2k\pi)\cos u - \cos(\alpha + 2k\pi)\sin u = \frac{1}{2}\cos u \pm \frac{\sqrt{3}}{2}\sin u$. In rectangular coordinates, $x = \sin u$ so $y = \frac{1}{2}\cos u \pm \frac{\sqrt{3}}{2}x \quad \Rightarrow \quad y \pm \frac{\sqrt{3}}{2}x = \frac{1}{2}\cos u \quad \Rightarrow \quad 2y \pm \sqrt{3}\,x = \cos u$. But then $x^2 + \left(2y \pm \sqrt{3}\,x\right)^2 = \sin^2 u + \cos^2 u = 1 \quad \Rightarrow \quad x^2 + 4y^2 \pm 4\sqrt{3}\,xy + 3x^2 = 1 \quad \Rightarrow \quad 4x^2 \pm 4\sqrt{3}\,xy + 4y^2 = 1$, which may be recognized as a conic section. In particular, each equation is an ellipse rotated $\pm 15°$ from the standard orientation (see the following graph). The trace in $z = -\frac{1}{2}$ is similar: $z = \sin(u + v) = -\frac{1}{2} \quad \Rightarrow \quad u + v = \beta + 2k\pi$ where $\beta = \frac{7\pi}{6}$ or $\frac{11\pi}{6}$. Then $v = (\beta + 2k\pi) - u$ and the trace is given by the parametric equations $x = \sin u$, $y = \sin v = \sin[(\beta + 2k\pi) - u] = \sin(\beta + 2k\pi)\cos u - \cos(\beta + 2k\pi)\sin u = -\frac{1}{2}\cos u \pm \frac{\sqrt{3}}{2}\sin u$. If we convert to rectangular coordinates, we arrive at the same pair of equations, $4x^2 \pm 4\sqrt{3}\,xy + 4y^2 = 1$, so the trace is identical to the trace in $z = \frac{1}{2}$.

Graphing each of these, we have the following 5 traces.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.



$z = -1$ $z = -\frac{1}{2}$ $z = 0$



$z = \frac{1}{2}$ $z = 1$

Visualizing these traces on the surface reveals that horizontal cross sections are pairs of intersecting ellipses whose major axes are perpendicular to each other. At the bottom of the surface, $z = -1$, the ellipses coincide as circles of radius 1. As we move up the surface, the ellipses become narrower until at $z = 0$ they collapse into line segments, after which the process is reversed, and the ellipses widen to again coincide as circles at $z = 1$.

5. $(\mathbf{F} \cdot \nabla)\,\mathbf{G} = \left(P_1 \dfrac{\partial}{\partial x} + Q_1 \dfrac{\partial}{\partial y} + R_1 \dfrac{\partial}{\partial z}\right)(P_2\,\mathbf{i} + Q_2\,\mathbf{j} + R_2\,\mathbf{k})$

$$= \left(P_1 \frac{\partial P_2}{\partial x} + Q_1 \frac{\partial P_2}{\partial y} + R_1 \frac{\partial P_2}{\partial z}\right)\mathbf{i} + \left(P_1 \frac{\partial Q_2}{\partial x} + Q_1 \frac{\partial Q_2}{\partial y} + R_1 \frac{\partial Q_2}{\partial z}\right)\mathbf{j}$$
$$+ \left(P_1 \frac{\partial R_2}{\partial x} + Q_1 \frac{\partial R_2}{\partial y} + R_1 \frac{\partial R_2}{\partial z}\right)\mathbf{k}$$

$$= (\mathbf{F} \cdot \nabla P_2)\,\mathbf{i} + (\mathbf{F} \cdot \nabla Q_2)\,\mathbf{j} + (\mathbf{F} \cdot \nabla R_2)\,\mathbf{k}.$$

Similarly, $(\mathbf{G} \cdot \nabla)\,\mathbf{F} = (\mathbf{G} \cdot \nabla P_1)\,\mathbf{i} + (\mathbf{G} \cdot \nabla Q_1)\,\mathbf{j} + (\mathbf{G} \cdot \nabla R_1)\,\mathbf{k}$. Then

$$\mathbf{F} \times \operatorname{curl} \mathbf{G} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ P_1 & Q_1 & R_1 \\ \partial R_2/\partial y - \partial Q_2/\partial z & \partial P_2/\partial z - \partial R_2/\partial x & \partial Q_2/\partial x - \partial P_2/\partial y \end{vmatrix}$$

$$= \left(Q_1 \frac{\partial Q_2}{\partial x} - Q_1 \frac{\partial P_2}{\partial y} - R_1 \frac{\partial P_2}{\partial z} + R_1 \frac{\partial R_2}{\partial x}\right)\mathbf{i} + \left(R_1 \frac{\partial R_2}{\partial y} - R_1 \frac{\partial Q_2}{\partial z} - P_1 \frac{\partial Q_2}{\partial x} + P_1 \frac{\partial P_2}{\partial y}\right)\mathbf{j}$$
$$+ \left(P_1 \frac{\partial P_2}{\partial z} - P_1 \frac{\partial R_2}{\partial x} - Q_1 \frac{\partial R_2}{\partial y} + Q_1 \frac{\partial Q_2}{\partial z}\right)\mathbf{k}$$

[continued]

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

and

$$\mathbf{G} \times \operatorname{curl} \mathbf{F} = \left(Q_2 \frac{\partial Q_1}{\partial x} - Q_2 \frac{\partial P_1}{\partial y} - R_2 \frac{\partial P_1}{\partial z} + R_2 \frac{\partial R_1}{\partial x} \right) \mathbf{i} + \left(R_2 \frac{\partial R_1}{\partial y} - R_2 \frac{\partial Q_1}{\partial z} - P_2 \frac{\partial Q_1}{\partial x} + P_2 \frac{\partial P_1}{\partial y} \right) \mathbf{j} + \left(P_2 \frac{\partial P_1}{\partial z} - P_2 \frac{\partial R_1}{\partial x} - Q_2 \frac{\partial R_1}{\partial y} + Q_2 \frac{\partial Q_1}{\partial z} \right) \mathbf{k}.$$

Then

$$(\mathbf{F} \cdot \nabla)\, \mathbf{G} + \mathbf{F} \times \operatorname{curl} \mathbf{G} = \left(P_1 \frac{\partial P_2}{\partial x} + Q_1 \frac{\partial Q_2}{\partial x} + R_1 \frac{\partial R_2}{\partial x} \right) \mathbf{i} + \left(P_1 \frac{\partial P_2}{\partial y} + Q_1 \frac{\partial Q_2}{\partial y} + R_1 \frac{\partial R_2}{\partial y} \right) \mathbf{j} + \left(P_1 \frac{\partial P_2}{\partial z} + Q_1 \frac{\partial Q_2}{\partial z} + R_1 \frac{\partial R_2}{\partial z} \right) \mathbf{k}$$

and

$$(\mathbf{G} \cdot \nabla)\, \mathbf{F} + \mathbf{G} \times \operatorname{curl} \mathbf{F} = \left(P_2 \frac{\partial P_1}{\partial x} + Q_2 \frac{\partial Q_1}{\partial x} + R_2 \frac{\partial R_1}{\partial x} \right) \mathbf{i} + \left(P_2 \frac{\partial P_1}{\partial y} + Q_2 \frac{\partial Q_1}{\partial y} + R_2 \frac{\partial R_1}{\partial y} \right) \mathbf{j} + \left(P_2 \frac{\partial P_1}{\partial z} + Q_2 \frac{\partial Q_1}{\partial z} + R_2 \frac{\partial R_1}{\partial z} \right) \mathbf{k}.$$

Hence

$$\begin{aligned}
&(\mathbf{F} \cdot \nabla)\, \mathbf{G} + \mathbf{F} \times \operatorname{curl} \mathbf{G} + (\mathbf{G} \cdot \nabla)\, \mathbf{F} + \mathbf{G} \times \operatorname{curl} \mathbf{F} \\
&\quad = \left[\left(P_1 \frac{\partial P_2}{\partial x} + P_2 \frac{\partial P_1}{\partial x} \right) + \left(Q_1 \frac{\partial Q_2}{\partial x} + Q_2 \frac{\partial Q_1}{\partial y} \right) + \left(R_1 \frac{\partial R_2}{\partial x} + R_2 \frac{\partial R_1}{\partial x} \right) \right] \mathbf{i} \\
&\qquad + \left[\left(P_1 \frac{\partial P_2}{\partial y} + P_2 \frac{\partial P_1}{\partial y} \right) + \left(Q_1 \frac{\partial Q_2}{\partial y} + Q_2 \frac{\partial Q_1}{\partial y} \right) + \left(R_1 \frac{\partial R_2}{\partial y} + R_2 \frac{\partial R_1}{\partial y} \right) \right] \mathbf{j} \\
&\qquad + \left[\left(P_1 \frac{\partial P_2}{\partial z} + P_2 \frac{\partial P_1}{\partial z} \right) + \left(Q_1 \frac{\partial Q_2}{\partial z} + Q_2 \frac{\partial Q_1}{\partial z} \right) + \left(R_1 \frac{\partial R_2}{\partial z} + R_2 \frac{\partial R_1}{\partial z} \right) \right] \mathbf{k} \\
&\quad = \nabla(P_1 P_2 + Q_1 Q_2 + R_1 R_2) = \nabla(\mathbf{F} \cdot \mathbf{G}).
\end{aligned}$$

6. (a) First we place the piston on coordinate axes so the top of the cylinder is at the origin and $x(t) > 0$ is the distance from the top of the cylinder to the piston at time t. Let C_1 be the curve traced out by the piston during one four-stroke cycle, so C_1 is given by $\mathbf{r}(t) = x(t)\,\mathbf{i}$, $a \le t \le b$. (Thus, the curve lies on the positive x-axis and reverses direction several times.) The force on the piston is $AP(t)\,\mathbf{i}$, where A is the area of the top of the piston and $P(t)$ is the pressure in the cylinder at time t. As in Section 16.2, the work done on the piston is $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_a^b AP(t)\,\mathbf{i} \cdot x'(t)\,\mathbf{i}\, dt = \int_a^b AP(t)\, x'(t)\, dt$. Here, the volume of the cylinder at time t is $V(t) = Ax(t) \;\Rightarrow\; V'(t) = Ax'(t) \;\Rightarrow\; \int_a^b AP(t)\, x'(t)\, dt = \int_a^b P(t)\, V'(t)\, dt$. Since the curve C in the PV-plane corresponds to the values of P and V at time t, $a \le t \le b$, we have

$$W = \int_a^b AP(t)\, x'(t)\, dt = \int_a^b P(t)\, V'(t)\, dt = \int_C P\, dV$$

Another method: If we divide the time interval $[a, b]$ into n subintervals of equal length Δt, the amount of work done on the piston in the ith time interval is approximately $AP(t_i)[x(t_i) - x(t_{i-1})]$. Thus we estimate the total work done during

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

one cycle to be $\sum_{i=1}^{n} AP(t_i)[x(t_i) - x(t_{i-1})]$. If we allow $n \to \infty$, we have

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} AP(t_i)[x(t_i) - x(t_{i-1})] = \lim_{n\to\infty} \sum_{i=1}^{n} P(t_i)[Ax(t_i) - Ax(t_{i-1})] = \lim_{n\to\infty} \sum_{i=1}^{n} P(t_i)[V(t_i) - V(t_{i-1})]$$
$$= \int_C P\,dV$$

(b) Let C_L be the lower loop of the curve C and C_U the upper loop. Then $C = C_L \cup C_U$. C_L is positively oriented, so from Formula 16.4.5 we know the area of the lower loop in the PV-plane is given by $-\oint_{C_L} P\,dV$. C_U is negatively oriented, so the area of the upper loop is given by $-\left(-\oint_{C_U} P\,dV\right) = \oint_{C_U} P\,dV$. From part (a),

$$W = \int_C P\,dV = \int_{C_L \cup C_U} P\,dV = \oint_{C_L} P\,dV + \oint_{C_U} P\,dV = \oint_{C_U} P\,dV - \left(-\oint_{C_L} P\,dV\right),$$

the difference of the areas enclosed by the two loops of C.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

17 □ SECOND-ORDER DIFFERENTIAL EQUATIONS

17.1 Second-Order Linear Equations

1. The auxiliary equation is $r^2 - r - 6 = 0 \Rightarrow (r-3)(r+2) = 0 \Rightarrow r = 3, r = -2$. Then by (8) the general solution is $y = c_1 e^{3x} + c_2 e^{-2x}$.

2. The auxiliary equation is $r^2 - 6r + 9 = 0 \Rightarrow (r-3)^2 = 0 \Rightarrow r = 3$. Then by (10), the general solution is $y = c_1 e^{3x} + c_2 x e^{3x}$.

3. The auxiliary equation is $r^2 + 2 = 0 \Rightarrow r = \pm\sqrt{2}\,i$. Then by (11) the general solution is $y = e^{0x}\left(c_1 \cos\left(\sqrt{2}\,x\right) + c_2 \sin\left(\sqrt{2}\,x\right)\right) = c_1 \cos\left(\sqrt{2}\,x\right) + c_2 \sin\left(\sqrt{2}\,x\right)$.

4. The auxiliary equation is $r^2 + r - 12 = 0 \Rightarrow (r-3)(r+4) = 0 \Rightarrow r = 3, r = -4$. Then by (8) the general solution is $y = c_1 e^{3x} + c_2 e^{-4x}$.

5. The auxiliary equation is $4r^2 + 4r + 1 = 0 \Rightarrow (2r+1)^2 = 0 \Rightarrow r = -\frac{1}{2}$. Then by (10), the general solution is $y = c_1 e^{-x/2} + c_2 x e^{-x/2}$.

6. The auxiliary equation is $9r^2 + 4 = 0 \Rightarrow r^2 = -\frac{4}{9} \Rightarrow r = \pm\frac{2}{3}i$, so the general solution is $y = e^{0x}\left[c_1 \cos\left(\frac{2}{3}x\right) + c_2 \sin\left(\frac{2}{3}x\right)\right] = c_1 \cos\left(\frac{2}{3}x\right) + c_2 \sin\left(\frac{2}{3}x\right)$.

7. The auxiliary equation is $3r^2 - 4r = r(3r-4) = 0 \Rightarrow r = 0, r = \frac{4}{3}$, so $y = c_1 e^{0x} + c_2 e^{4x/3} = c_1 + c_2 e^{4x/3}$.

8. The auxiliary equation is $r^2 - 1 = (r-1)(r+1) = 0 \Rightarrow r = 1, r = -1$. Then the general solution is $y = c_1 e^{x} + c_2 e^{-x}$.

9. The auxiliary equation is $r^2 - 4r + 13 = 0 \Rightarrow r = \dfrac{4 \pm \sqrt{-36}}{2} = 2 \pm 3i$, so $y = e^{2x}(c_1 \cos 3x + c_2 \sin 3x)$.

10. The auxiliary equation is $3r^2 + 4r - 3 = 0 \Rightarrow r = \dfrac{-4 \pm \sqrt{52}}{6} = \dfrac{-2 \pm \sqrt{13}}{3}$, so $y = c_1 e^{(-2+\sqrt{13})x/3} + c_2 e^{(-2-\sqrt{13})x/3}$.

11. The auxiliary equation is $2r^2 + 2r - 1 = 0 \Rightarrow r = \dfrac{-2 \pm \sqrt{12}}{4} = \dfrac{-1 \pm \sqrt{3}}{2}$, so $y = c_1 e^{\left(-1+\sqrt{3}\right)t/2} + c_2 e^{\left(-1-\sqrt{3}\right)t/2}$.

12. The auxiliary equation is $r^2 + 6r + 34 = 0 \Rightarrow r = \dfrac{-6 \pm \sqrt{-100}}{2} = -3 \pm 5i$, so $R = e^{-3t}(c_1 \cos 5t + c_2 \sin 5t)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. The auxiliary equation is $3r^2 + 4r + 3 = 0 \Rightarrow r = \dfrac{-4 \pm \sqrt{-20}}{6} = -\frac{2}{3} \pm \frac{\sqrt{5}}{3}i$, so

$V = e^{-2t/3}\left[c_1 \cos\left(\frac{\sqrt{5}}{3}t\right) + c_2 \sin\left(\frac{\sqrt{5}}{3}t\right)\right]$.

14. The auxiliary equation is $4r^2 - 4r + 1 = (2r-1)^2 = 0 \Rightarrow r = \frac{1}{2}$, so the general solution is $y = c_1 e^{x/2} + c_2 x e^{x/2}$. We graph the basic solutions $f(x) = e^{x/2}$, $g(x) = xe^{x/2}$ as well as $y = 2e^{x/2} + 3xe^{x/2}$, $y = -e^{x/2} - 3xe^{x/2}$, and $y = 4e^{x/2} - 2xe^{x/2}$. The graphs are all asymptotic to the x-axis as $x \to -\infty$, and as $x \to \infty$ the solutions approach $\pm\infty$.



15. The auxiliary equation is $r^2 + 2r + 2 = 0 \Rightarrow$

$r = \dfrac{-2 \pm \sqrt{-4}}{2} = -1 \pm i$, so the general solution is

$y = e^{-x}(c_1 \cos x + c_2 \sin x)$. We graph the basic solutions

$f(x) = e^{-x}\cos x$, $g(x) = e^{-x}\sin x$ as well as



$y = e^{-x}(-\cos x - 2\sin x)$ and $y = e^{-x}(2\cos x + 3\sin x)$. All the solutions oscillate with amplitudes that become arbitrarily large as $x \to -\infty$ and the solutions are asymptotic to the x-axis as $x \to \infty$.

16. The auxiliary equation is $2r^2 + r - 1 = (2r-1)(r+1) = 0 \Rightarrow$

$r = \frac{1}{2}$, $r = -1$, so the general solution is $y = c_1 e^{x/2} + c_2 e^{-x}$. We graph the basic solutions $f(x) = e^{x/2}$, $g(x) = e^{-x}$ as well as $y = 2e^{x/2} + e^{-x}$, $y = -e^{x/2} - 2e^{-x}$, and $y = e^{x/2} - e^{-x}$. Each solution consists of a single continuous curve that approaches either 0 or $\pm\infty$ as $x \to \pm\infty$.



17. $r^2 + 3 = 0 \Rightarrow r = \pm\sqrt{3}\,i$ and the general solution is

$y = e^{0x}\left[c_1 \cos\left(\sqrt{3}\,x\right) + c_2 \sin\left(\sqrt{3}\,x\right)\right] = c_1 \cos\left(\sqrt{3}\,x\right) + c_2 \sin\left(\sqrt{3}\,x\right)$. Then $y(0) = 1 \Rightarrow c_1 = 1$ and, since $y' = -\sqrt{3}\,c_1 \sin\left(\sqrt{3}\,x\right) + \sqrt{3}\,c_2 \cos\left(\sqrt{3}\,x\right)$, $y'(0) = 3 \Rightarrow \sqrt{3}\,c_2 = 3 \Rightarrow c_2 = \frac{3}{\sqrt{3}} = \sqrt{3}$, so the solution to the initial-value problem is $y = \cos\left(\sqrt{3}\,x\right) + \sqrt{3}\,\sin\left(\sqrt{3}\,x\right)$.

18. $r^2 - 2r - 3 = (r-3)(r+1) = 0$, so $r = 3$, $r = -1$ and the general solution is $y = c_1 e^{3x} + c_2 e^{-x}$. Then $y' = 3c_1 e^{3x} - c_2 e^{-x}$, so $y(0) = 2 \Rightarrow c_1 + c_2 = 2$ and $y'(0) = 2 \Rightarrow 3c_1 - c_2 = 2$, giving $c_1 = 1$ and $c_2 = 1$. Thus the solution to the initial-value problem is $y = e^{3x} + e^{-x}$.

19. $9r^2 + 12r + 4 = (3r+2)^2 = 0 \Rightarrow r = -\frac{2}{3}$ and the general solution is $y = c_1 e^{-2x/3} + c_2 x e^{-2x/3}$. Then $y(0) = 1 \Rightarrow c_1 = 1$ and, since $y' = -\frac{2}{3}c_1 e^{-2x/3} + c_2\left(1 - \frac{2}{3}x\right)e^{-2x/3}$, $y'(0) = 0 \Rightarrow -\frac{2}{3}c_1 + c_2 = 0$, so $c_2 = \frac{2}{3}$ and the solution to the initial-value problem is $y = e^{-2x/3} + \frac{2}{3}xe^{-2x/3}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. $3r^2 - 2r - 1 = (3r+1)(r-1) = 0 \;\Rightarrow\; r = -\frac{1}{3}, r = 1$ and the general solution is $y = c_1 e^{-x/3} + c_2 e^x$. Then $y' = -\frac{1}{3}c_1 e^{-x/3} + c_2 e^x$, so $y(0) = 0 \;\Rightarrow\; c_1 + c_2 = 0$ and $y'(0) = -4 \;\Rightarrow\; -\frac{1}{3}c_1 + c_2 = -4$, giving $c_1 = 3$ and $c_2 = -3$. Thus the solution to the initial-value problem is $y = 3e^{-x/3} - 3e^x$.

21. $r^2 - 6r + 10 = 0 \;\Rightarrow\; r = 3 \pm i$ and the general solution is $y = e^{3x}(c_1 \cos x + c_2 \sin x)$. Then $2 = y(0) = c_1$ and $3 = y'(0) = c_2 + 3c_1 \;\Rightarrow\; c_2 = -3$ and the solution to the initial-value problem is $y = e^{3x}(2\cos x - 3\sin x)$.

22. $4r^2 - 20r + 25 = (2r-5)^2 = 0 \;\Rightarrow\; r = \frac{5}{2}$ and the general solution is $y = c_1 e^{5x/2} + c_2 x e^{5x/2}$. Then $2 = y(0) = c_1$ and $-3 = y'(0) = \frac{5}{2}c_1 + c_2 \;\Rightarrow\; c_2 = -8$. The solution to the initial-value problem is $y = 2e^{5x/2} - 8xe^{5x/2}$.

23. $r^2 - r - 12 = (r-4)(r+3) = 0 \;\Rightarrow\; r = 4, r = -3$ and the general solution is $y = c_1 e^{4x} + c_2 e^{-3x}$. Then $0 = y(1) = c_1 e^4 + c_2 e^{-3}$ and $1 = y'(1) = 4c_1 e^4 - 3c_2 e^{-3}$ so $c_1 = \frac{1}{7}e^{-4}$, $c_2 = -\frac{1}{7}e^3$ and the solution to the initial-value problem is $y = \frac{1}{7}e^{-4}e^{4x} - \frac{1}{7}e^3 e^{-3x} = \frac{1}{7}e^{4x-4} - \frac{1}{7}e^{3-3x}$.

24. $4r^2 + 4r + 3 = 0 \;\Rightarrow\; r = -\frac{1}{2} \pm \frac{\sqrt{2}}{2}i$ and the general solution is $y = e^{-x/2}\left(c_1 \cos \frac{\sqrt{2}}{2}x + c_2 \sin \frac{\sqrt{2}}{2}x\right)$. Then $0 = y(0) = c_1$ and $1 = y'(0) = \frac{\sqrt{2}}{2}c_2 - \frac{1}{2}c_1 \;\Rightarrow\; c_2 = \sqrt{2}$ and the solution to the initial-value problem is $y = e^{-x/2}\left(0 + \sqrt{2}\sin \frac{\sqrt{2}}{2}x\right) = \sqrt{2}\,e^{-x/2}\sin \frac{\sqrt{2}}{2}x$.

25. $r^2 + 16 = 0 \;\Rightarrow\; r = \pm 4i$ and the general solution is $y = c_1 \cos 4x + c_2 \sin 4x$. Then $-3 = y(0) = c_1$ and $2 = y(\pi/8) = c_2$, so the solution of the boundary-value problem is $y = -3\cos 4x + 2\sin 4x$.

26. $r^2 + 6r = r(r+6) = 0 \;\Rightarrow\; r = 0, r = -6$ and the general solution is $y = c_1 + c_2 e^{-6x}$. Then $1 = y(0) = c_1 + c_2$ and $0 = y(1) = c_1 + c_2 e^{-6}$ so $c_1 = \dfrac{1}{1-e^6}$, $c_2 = -\dfrac{e^6}{1-e^6}$. The solution of the boundary-value problem is $y = \dfrac{1}{1-e^6} - \dfrac{e^6}{1-e^6}\cdot e^{-6x} = \dfrac{1}{1-e^6} - \dfrac{e^{6-6x}}{1-e^6}$.

27. $r^2 + 4r + 4 = (r+2)^2 = 0 \;\Rightarrow\; r = -2$ and the general solution is $y = c_1 e^{-2x} + c_2 x e^{-2x}$. Then $2 = y(0) = c_1$ and $0 = y(1) = c_1 e^{-2} + c_2 e^{-2}$ so $c_2 = -2$, and the solution of the boundary-value problem is $y = 2e^{-2x} - 2xe^{-2x}$.

28. $r^2 - 8r + 17 = 0 \;\Rightarrow\; r = 4 \pm i$ and the general solution is $y = e^{4x}(c_1 \cos x + c_2 \sin x)$. But $3 = y(0) = c_1$ and $2 = y(\pi) = -c_1 e^{4\pi} \;\Rightarrow\; c_1 = -2/e^{4\pi}$, so there is no solution.

29. $r^2 - r = r(r-1) = 0 \;\Rightarrow\; r = 0, r = 1$ and the general solution is $y = c_1 + c_2 e^x$. Then $1 = y(0) = c_1 + c_2$ and $2 = y(1) = c_1 + c_2 e$ so $c_1 = \dfrac{e-2}{e-1}$, $c_2 = \dfrac{1}{e-1}$. The solution of the boundary-value problem is $y = \dfrac{e-2}{e-1} + \dfrac{e^x}{e-1}$.

30. $4r^2 - 4r + 1 = (2r-1)^2 = 0 \;\Rightarrow\; r = \frac{1}{2}$ and the general solution is $y = c_1 e^{x/2} + c_2 x e^{x/2}$. Then $4 = y(0) = c_1$ and $0 = y(2) = c_1 e + 2c_2 e \;\Rightarrow\; c_2 = -2$. The solution of the boundary-value problem is $y = 4e^{x/2} - 2xe^{x/2}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

31. $r^2+4r+20=0 \Rightarrow r=-2\pm 4i$ and the general solution is $y=e^{-2x}(c_1\cos 4x+c_2\sin 4x)$. But $1=y(0)=c_1$ and $2=y(\pi)=c_1e^{-2\pi} \Rightarrow c_1=2e^{2\pi}$, so there is no solution.

32. $r^2+4r+20=0 \Rightarrow r=-2\pm 4i$ and the general solution is $y=e^{-2x}(c_1\cos 4x+c_2\sin 4x)$. But $1=y(0)=c_1$ and $e^{-2\pi}=y(\pi)=c_1e^{-2\pi} \Rightarrow c_1=1$, so c_2 can vary and the solution of the boundary-value problem is $y=e^{-2x}(\cos 4x+c\sin 4x)$, where c is any constant.

33. (a) *Case 1* ($\lambda=0$): $y''+\lambda y=0 \Rightarrow y''=0$ which has an auxiliary equation $r^2=0 \Rightarrow r=0 \Rightarrow y=c_1+c_2x$ where $y(0)=0$ and $y(L)=0$. Thus, $0=y(0)=c_1$ and $0=y(L)=c_2L \Rightarrow c_1=c_2=0$. Thus $y=0$.

Case 2 ($\lambda<0$): $y''+\lambda y=0$ has auxiliary equation $r^2=-\lambda \Rightarrow r=\pm\sqrt{-\lambda}$ [distinct and real since $\lambda<0$] $\Rightarrow$ $y=c_1e^{\sqrt{-\lambda}x}+c_2e^{-\sqrt{-\lambda}x}$ where $y(0)=0$ and $y(L)=0$. Thus $0=y(0)=c_1+c_2$ $(*)$ and $0=y(L)=c_1e^{\sqrt{-\lambda}L}+c_2e^{-\sqrt{-\lambda}L}$ $(\dagger)$.

Multiplying $(*)$ by $e^{\sqrt{-\lambda}L}$ and subtracting $(\dagger)$ gives $c_2\left(e^{\sqrt{-\lambda}L}-e^{-\sqrt{-\lambda}L}\right)=0 \Rightarrow c_2=0$ and thus $c_1=0$ from $(*)$. Thus $y=0$ for the cases $\lambda=0$ and $\lambda<0$.

(b) $y''+\lambda y=0$ has an auxiliary equation $r^2+\lambda=0 \Rightarrow r=\pm i\sqrt{\lambda} \Rightarrow y=c_1\cos\sqrt{\lambda}\,x+c_2\sin\sqrt{\lambda}\,x$ where $y(0)=0$ and $y(L)=0$. Thus, $0=y(0)=c_1$ and $0=y(L)=c_2\sin\sqrt{\lambda}L$ since $c_1=0$. Since we cannot have a trivial solution, $c_2\neq 0$ and thus $\sin\sqrt{\lambda}\,L=0 \Rightarrow \sqrt{\lambda}\,L=n\pi$ where n is an integer $\Rightarrow \lambda=n^2\pi^2/L^2$ and $y=c_2\sin(n\pi x/L)$ where n is an integer.

34. The auxiliary equation is $ar^2+br+c=0$. If $b^2-4ac>0$, then any solution is of the form $y(x)=c_1e^{r_1x}+c_2e^{r_2x}$ where $r_1=\dfrac{-b+\sqrt{b^2-4ac}}{2a}$ and $r_2=\dfrac{-b-\sqrt{b^2-4ac}}{2a}$. But a, b, and c are all positive so both r_1 and r_2 are negative and $\lim_{x\to\infty}y(x)=0$. If $b^2-4ac=0$, then any solution is of the form $y(x)=c_1e^{rx}+c_2xe^{rx}$ where $r=-b/(2a)<0$ since a, b are positive. Hence $\lim_{x\to\infty}y(x)=0$. Finally if $b^2-4ac<0$, then any solution is of the form $y(x)=e^{\alpha x}(c_1\cos\beta x+c_2\sin\beta x)$ where $\alpha=-b/(2a)<0$ since a and b are positive. Thus $\lim_{x\to\infty}y(x)=0$.

35. (a) $r^2-2r+2=0 \Rightarrow r=1\pm i$ and the general solution is $y=e^x(c_1\cos x+c_2\sin x)$. If $y(a)=c$ and $y(b)=d$ then $e^a(c_1\cos a+c_2\sin a)=c \Rightarrow c_1\cos a+c_2\sin a=ce^{-a}$ and $e^b(c_1\cos b+c_2\sin b)=d \Rightarrow$ $c_1\cos b+c_2\sin b=de^{-b}$. This gives a linear system in c_1 and c_2 which has a unique solution if the lines are not parallel. If the lines are not vertical or horizontal, we have parallel lines if $\cos a=k\cos b$ and $\sin a=k\sin b$ for some nonzero constant k or $\dfrac{\cos a}{\cos b}=k=\dfrac{\sin a}{\sin b} \Rightarrow \dfrac{\sin a}{\cos a}=\dfrac{\sin b}{\cos b} \Rightarrow \tan a=\tan b \Rightarrow b-a=n\pi$, n any integer. (Note that none of $\cos a$, $\cos b$, $\sin a$, $\sin b$ are zero.) If the lines are both horizontal then $\cos a=\cos b=0 \Rightarrow b-a=n\pi$, and similarly vertical lines means $\sin a=\sin b=0 \Rightarrow b-a=n\pi$. Thus the system has a unique solution if $b-a\neq n\pi$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The linear system has no solution if the lines are parallel but not identical. From part (a) the lines are parallel if $b - a = n\pi$. If the lines are not horizontal, they are identical if $ce^{-a} = kde^{-b} \Rightarrow \dfrac{ce^{-a}}{de^{-b}} = k = \dfrac{\cos a}{\cos b} \Rightarrow \dfrac{c}{d} = e^{a-b}\dfrac{\cos a}{\cos b}$. (If $d = 0$ then $c = 0$ also.) If they are horizontal then $\cos b = 0$, but $k = \dfrac{\sin a}{\sin b}$ also (and $\sin b \neq 0$) so we require $\dfrac{c}{d} = e^{a-b}\dfrac{\sin a}{\sin b}$. Thus the system has no solution if $b - a = n\pi$ and $\dfrac{c}{d} \neq e^{a-b}\dfrac{\cos a}{\cos b}$ unless $\cos b = 0$, in which case $\dfrac{c}{d} \neq e^{a-b}\dfrac{\sin a}{\sin b}$.

(c) The linear system has infinitely many solution if the lines are identical (and necessarily parallel). From part (b) this occurs when $b - a = n\pi$ and $\dfrac{c}{d} = e^{a-b}\dfrac{\cos a}{\cos b}$ unless $\cos b = 0$, in which case $\dfrac{c}{d} = e^{a-b}\dfrac{\sin a}{\sin b}$.

17.2 Nonhomogeneous Linear Equations

1. The auxiliary equation is $r^2 + 2r - 8 = (r-2)(r+4) = 0 \Rightarrow r = 2, r = -4$, so the complementary solution is $y_c(x) = c_1e^{2x} + c_2e^{-4x}$. We try the particular solution $y_p(x) = Ax^2 + Bx + C$, so $y_p' = 2Ax + B$ and $y_p'' = 2A$. Substituting into the differential equation, we have $(2A) + 2(2Ax + B) - 8(Ax^2 + Bx + C) = 1 - 2x^2$ or $-8Ax^2 + (4A - 8B)x + (2A + 2B - 8C) = -2x^2 + 1$. Comparing coefficients gives $-8A = -2 \Rightarrow A = \frac{1}{4}$, $4A - 8B = 0 \Rightarrow B = \frac{1}{8}$, and $2A + 2B - 8C = 1 \Rightarrow C = -\frac{1}{32}$, so the general solution is $y(x) = y_c(x) + y_p(x) = c_1e^{2x} + c_2e^{-4x} + \frac{1}{4}x^2 + \frac{1}{8}x - \frac{1}{32}$.

2. The auxiliary equation is $r^2 - 3r = r(r-3) = 0 \Rightarrow r = 0, r = 3$, so the complementary solution is $y_c(x) = c_1 + c_2e^{3x}$. We try the particular solution $y_p(x) = A\cos 2x + B\sin 2x$, so $y_p' = -2A\sin 2x + 2B\cos 2x$ and $y_p'' = -4A\cos 2x - 4B\sin 2x$. Substitution into the differential equation gives $(-4A\cos 2x - 4B\sin 2x) - 3(-2A\sin 2x + 2B\cos 2x) = \sin 2x \Rightarrow (-4A - 6B)\cos 2x + (6A - 4B)\sin 2x = \sin 2x$. Then $-4A - 6B = 0$ and $6A - 4B = 1 \Rightarrow A = \frac{3}{26}$ and $B = -\frac{1}{13}$. Thus the general solution is $y(x) = y_c(x) + y_p(x) = c_1 + c_2e^{3x} + \frac{3}{26}\cos 2x - \frac{1}{13}\sin 2x$.

3. The auxiliary equation is $9r^2 + 1 = 0$ with roots $r = \pm\frac{1}{3}i$, so the complementary solution is $y_c(x) = c_1\cos(x/3) + c_2\sin(x/3)$. Try the particular solution $y_p(x) = Ae^{2x}$, so $y_p' = 2Ae^{2x}$ and $y_p'' = 4Ae^{2x}$. Substitution into the differential equation gives $9\left(4Ae^{2x}\right) + \left(Ae^{2x}\right) = e^{2x}$ or $37Ae^{2x} = e^{2x}$. Thus $37A = 1 \Rightarrow A = \frac{1}{37}$ and the general solution is $y(x) = y_c(x) + y_p(x) = c_1\cos(x/3) + c_2\sin(x/3) + \frac{1}{37}e^{2x}$.

4. The auxiliary equation is $r^2 - 2r + 2 = 0$ with roots $r = 1 \pm i$, so the complementary solution is $y_c(x) = e^x(c_1\cos x + c_2\sin x)$. Try the particular solution $y_p(x) = Ax + B + Ce^x$, so $y_p' = A + Ce^x$ and $y_p'' = Ce^x$. Substitution into the differential equation gives $(Ce^x) - 2(A + Ce^x) + 2(Ax + B + Ce^x) = x + e^x \Rightarrow$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$2Ax + (-2A + 2B) + Ce^x = x + e^x$. Comparing coefficients, we have $2A = 1 \Rightarrow A = \frac{1}{2}$, $-2A + 2B = 0 \Rightarrow B = \frac{1}{2}$, and $C = 1$, so the general solution is $y(x) = y_c(x) + y_p(x) = e^x(c_1 \cos x + c_2 \sin x) + \frac{1}{2}x + \frac{1}{2} + e^x$.

5. The auxiliary equation is $r^2 - 4r + 5 = 0$ with roots $r = 2 \pm i$, so the complementary solution is $y_c(x) = e^{2x}(c_1 \cos x + c_2 \sin x)$. Try $y_p(x) = Ae^{-x}$, so $y_p' = -Ae^{-x}$ and $y_p'' = Ae^{-x}$. Substitution gives $Ae^{-x} - 4(-Ae^{-x}) + 5(Ae^{-x}) = e^{-x} \Rightarrow 10Ae^{-x} = e^{-x} \Rightarrow A = \frac{1}{10}$. Thus the general solution is $y(x) = e^{2x}(c_1 \cos x + c_2 \sin x) + \frac{1}{10}e^{-x}$.

6. The auxiliary equation is $r^2 - 4r + 4 = (r-2)^2 = 0 \Rightarrow r = 2$, so the complementary solution is $y_c(x) = c_1e^{2x} + c_2xe^{2x}$. For $y'' - 4y' + 4y = x$ try $y_{p_1}(x) = Ax + B$. Then $y_{p_1}' = A$ and $y_{p_1}'' = 0$, and substitution into the differential equation gives $0 - 4A + 4(Ax + B) = x$ or $4Ax + (4B - 4A) = x$, so $4A = 1 \Rightarrow A = \frac{1}{4}$ and $4B - 4A = 0 \Rightarrow B = \frac{1}{4}$. Thus $y_{p_1}(x) = \frac{1}{4}x + \frac{1}{4}$. For $y'' - 4y' + 4y = -\sin x$ try $y_{p_2}(x) = A\cos x + B\sin x$. Then $y_{p_2}' = -A\sin x + B\cos x$ and $y_{p_2}'' = -A\cos x - B\sin x$. Substituting, we have $(-A\cos x - B\sin x) - 4(-A\sin x + B\cos x) + 4(A\cos x + B\sin x) = -\sin x \Rightarrow (3A - 4B)\cos x + (4A + 3B)\sin x = -\sin x$. Thus $3A - 4B = 0$ and $4A + 3B = -1$, giving $A = -\frac{4}{25}$ and $B = -\frac{3}{25}$, so $y_{p_2}(x) = -\frac{4}{25}\cos x - \frac{3}{25}\sin x$. The general solution is $y(x) = y_c(x) + y_{p_1}(x) + y_{p_2}(x) = c_1e^{2x} + c_2xe^{2x} + \frac{1}{4}x + \frac{1}{4} - \frac{4}{25}\cos x - \frac{3}{25}\sin x$.

7. The auxiliary equation is $r^2 - 2r + 5 = 0$ with roots $r = 1 \pm 2i$, so the complementary solution is $y_c(x) = e^x(c_1 \cos 2x + c_2 \sin 2x)$. Try the particular solution $y_p(x) = A\cos x + B\sin x$, so $y_p' = -A\sin x + B\cos x$ and $y_p'' = -A\cos x - B\sin x$. Substituting, we have $(-A\cos x - B\sin x) - 2(-A\sin x + B\cos x) + 5(A\cos x + B\sin x) = \sin x \Rightarrow (4A - 2B)\cos x + (2A + 4B)\sin x = \sin x$. Then $4A - 2B = 0$, $2A + 4B = 1 \Rightarrow A = \frac{1}{10}$, $B = \frac{1}{5}$ and the general solution is $y(x) = y_c(x) + y_p(x) = e^x(c_1 \cos 2x + c_2 \sin 2x) + \frac{1}{10}\cos x + \frac{1}{5}\sin x$. But $1 = y(0) = c_1 + \frac{1}{10} \Rightarrow c_1 = \frac{9}{10}$ and $1 = y'(0) = 2c_2 + c_1 + \frac{1}{5} \Rightarrow c_2 = -\frac{1}{20}$. Thus the solution to the initial-value problem is $y(x) = e^x\left(\frac{9}{10}\cos 2x - \frac{1}{20}\sin 2x\right) + \frac{1}{10}\cos x + \frac{1}{5}\sin x$.

8. The auxiliary equation is $r^2 - 1 = 0$ with roots $r = \pm 1$, so the complementary solution is $y_c(x) = c_1e^x + c_2e^{-x}$. Try the particular solution $y_p(x) = (Ax + B)e^{2x}$, so $y_p' = (2Ax + A + 2B)e^{2x}$ and $y_p'' = (4Ax + 4A + 4B)e^{2x}$. Substituting, we have $(4Ax + 4A + 4B)e^{2x} - (Ax + B)e^{2x} = xe^{2x} \Rightarrow (3Ax + 4A + 3B)e^{2x} = xe^{2x}$. Then $3A = 1 \Rightarrow A = \frac{1}{3}$ and $4A + 3B = 0 \Rightarrow B = -\frac{4}{9}$, and the general solution is $y(x) = y_c(x) + y_p(x) = c_1e^x + c_2e^{-x} + \left(\frac{1}{3}x - \frac{4}{9}\right)e^{2x}$. But $0 = y(0) = c_1 + c_2 - \frac{4}{9}$ and $1 = y'(0) = c_1 - c_2 - \frac{5}{9} \Rightarrow c_1 = 1, c_2 = -\frac{5}{9}$. Thus the solution to the initial-value problem is $y(x) = e^x - \frac{5}{9}e^{-x} + \left(\frac{1}{3}x - \frac{4}{9}\right)e^{2x}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. The auxiliary equation is $r^2 - r = 0$ with roots $r = 0, r = 1$ so the complementary solution is $y_c(x) = c_1 + c_2e^x$. Try $y_p(x) = x(Ax + B)e^x$ so that no term in y_p is a solution of the complementary equation. Then $y_p' = (Ax^2 + (2A + B)x + B)e^x$ and $y_p'' = (Ax^2 + (4A + B)x + (2A + 2B))e^x$. Substitution into the differential equation gives $(Ax^2 + (4A + B)x + (2A + 2B))e^x - (Ax^2 + (2A + B)x + B)e^x = xe^x \Rightarrow (2Ax + (2A + B))e^x = xe^x \Rightarrow A = \frac{1}{2}, B = -1$. Thus $y_p(x) = \left(\frac{1}{2}x^2 - x\right)e^x$ and the general solution is $y(x) = c_1 + c_2e^x + \left(\frac{1}{2}x^2 - x\right)e^x$. But $2 = y(0) = c_1 + c_2$ and $1 = y'(0) = c_2 - 1$, so $c_2 = 2$ and $c_1 = 0$. The solution to the initial-value problem is $y(x) = 2e^x + \left(\frac{1}{2}x^2 - x\right)e^x = e^x\left(\frac{1}{2}x^2 - x + 2\right)$.

10. $y_c(x) = c_1e^x + c_2e^{-2x}$. For $y'' + y' - 2y = x$ try $y_{p_1}(x) = Ax + B$. Then $y_{p_1}' = A$, $y_{p_1}'' = 0$, and substitution gives $0 + A - 2(Ax + B) = x \Rightarrow A = -\frac{1}{2}, B = -\frac{1}{4}$, so $y_{p_1}(x) = -\frac{1}{2}x - \frac{1}{4}$. For $y'' + y' - 2y = \sin 2x$ try $y_{p_2}(x) = A\cos 2x + B\sin 2x$. Then $y_{p_2}' = -2A\sin 2x + 2B\cos 2x$, $y_{p_2}'' = -4A\cos 2x - 4B\sin 2x$, and substitution gives $(-4A\cos 2x - 4B\sin 2x) + (-2A\sin 2x + 2B\cos 2x) - 2(A\cos 2x + B\sin 2x) = \sin 2x \Rightarrow A = -\frac{1}{20}$, $B = -\frac{3}{20}$. Thus $y_{p_2}(x) = -\frac{1}{20}\cos 2x + -\frac{3}{20}\sin 2x$ and the general solution is $y(x) = c_1e^x + c_2e^{-2x} - \frac{1}{2}x - \frac{1}{4} - \frac{1}{20}\cos 2x - \frac{3}{20}\sin 2x$. But $1 = y(0) = c_1 + c_2 - \frac{1}{4} - \frac{1}{20}$ and $0 = y'(0) = c_1 - 2c_2 - \frac{1}{2} - \frac{3}{10} \Rightarrow c_1 = \frac{17}{15}$ and $c_2 = \frac{1}{6}$. Thus the solution to the initial-value problem is $y(x) = \frac{17}{15}e^x + \frac{1}{6}e^{-2x} - \frac{1}{2}x - \frac{1}{4} - \frac{1}{20}\cos 2x - \frac{3}{20}\sin 2x$.

11. The auxiliary equation is $r^2 + 3r + 2 = (r + 1)(r + 2) = 0$, so $r = -1, r = -2$ and $y_c(x) = c_1e^{-x} + c_2e^{-2x}$. Try $y_p = A\cos x + B\sin x \Rightarrow y_p' = -A\sin x + B\cos x$, $y_p'' = -A\cos x - B\sin x$. Substituting into the differential equation gives $(-A\cos x - B\sin x) + 3(-A\sin x + B\cos x) + 2(A\cos x + B\sin x) = \cos x$ or $(A + 3B)\cos x + (-3A + B)\sin x = \cos x$. Then solving the equations $A + 3B = 1, -3A + B = 0$ gives $A = \frac{1}{10}, B = \frac{3}{10}$ and the general solution is $y(x) = c_1e^{-x} + c_2e^{-2x} + \frac{1}{10}\cos x + \frac{3}{10}\sin x$. The graph shows y_p and several other solutions. Notice that all solutions are asymptotic to y_p as $x \to \infty$. Except for y_p, all solutions approach either ∞ or $-\infty$ as $x \to -\infty$.



12. The auxiliary equation is $r^2 + 4 = 0 \Rightarrow r = \pm 2i$, so $y_c(x) = c_1\cos 2x + c_2\sin 2x$. Try $y_p = Ae^{-x} \Rightarrow y_p' = -Ae^{-x}$, $y_p'' = Ae^{-x}$. Substituting into the differential equation gives $Ae^{-x} + 4Ae^{-x} = e^{-x} \Rightarrow 5A = 1 \Rightarrow A = \frac{1}{5}$, so $y_p = \frac{1}{5}e^{-x}$ and the general solution is $y(x) = c_1\cos 2x + c_2\sin 2x + \frac{1}{5}e^{-x}$. We graph y_p along with several other solutions. All of the solutions except y_p oscillate around $y_p = \frac{1}{5}e^{-x}$, and all solutions approach ∞ as $x \to -\infty$.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

13. Here $y_c(x) = c_1e^{2x} + c_2e^{-x}$, and a trial solution is $y_p(x) = (Ax + B)e^x \cos x + (Cx + D)e^x \sin x$.

14. Here $y_c(x) = c_1 \cos 2x + c_2 \sin 2x$. For $y'' + 4y = \cos 4x$ try $y_{p_1}(x) = A \cos 4x + B \sin 4x$ and for $y'' + 4y = \cos 2x$ try $y_{p_2}(x) = x(C \cos 2x + D \sin 2x)$ (so that no term of y_{p_2} is a solution of the complementary equation). Thus a trial solution is $y_p(x) = y_{p_1}(x) + y_{p_2}(x) = A \cos 4x + B \sin 4x + Cx \cos 2x + Dx \sin 2x$.

15. Here $y_c(x) = c_1e^{2x} + c_2e^x$. For $y'' - 3y' + 2y = e^x$ try $y_{p_1}(x) = Axe^x$ (since $y = Ae^x$ is a solution of the complementary equation) and for $y'' - 3y' + 2y = \sin x$ try $y_{p_2}(x) = B \cos x + C \sin x$. Thus a trial solution is $y_p(x) = y_{p_1}(x) + y_{p_2}(x) = Axe^x + B \cos x + C \sin x$.

16. Since $y_c(x) = c_1e^x + c_2e^{-4x}$ try $y_p(x) = x(Ax^3 + Bx^2 + Cx + D)e^x$ so that no term of $y_p(x)$ satisfies the complementary equation.

17. Since $y_c(x) = e^{-x}(c_1 \cos 3x + c_2 \sin 3x)$ we try $y_p(x) = x(Ax^2 + Bx + C)e^{-x} \cos 3x + x(Dx^2 + Ex + F)e^{-x} \sin 3x$ (so that no term of y_p is a solution of the complementary equation).

18. Here $y_c(x) = c_1 \cos 2x + c_2 \sin 2x$. For $y'' + 4y = e^{3x}$ try $y_{p_1}(x) = Ae^{3x}$ and for $y'' + 4y = x \sin 2x$ try $y_{p_2}(x) = x(Bx + C) \cos 2x + x(Dx + E) \sin 2x$ (so that no term of y_{p_2} is a solution of the complementary equation).

Note: Solving Equations (7) and (9) in The Method of Variation of Parameters gives

$$u_1' = -\frac{Gy_2}{a\,(y_1y_2' - y_2y_1')} \qquad \text{and} \qquad u_2' = \frac{Gy_1}{a\,(y_1y_2' - y_2y_1')}$$

We will use these equations rather than resolving the system in each of the remaining exercises in this section.

19. (a) Here $4r^2 + 1 = 0 \;\Rightarrow\; r = \pm\frac{1}{2}i$ and $y_c(x) = c_1 \cos\left(\frac{1}{2}x\right) + c_2 \sin\left(\frac{1}{2}x\right)$. We try a particular solution of the form $y_p(x) = A \cos x + B \sin x \;\Rightarrow\; y_p' = -A \sin x + B \cos x$ and $y_p'' = -A \cos x - B \sin x$. Then the equation $4y'' + y = \cos x$ becomes $4(-A \cos x - B \sin x) + (A \cos x + B \sin x) = \cos x$ or $-3A \cos x - 3B \sin x = \cos x \;\Rightarrow\; A = -\frac{1}{3}, B = 0$. Thus, $y_p(x) = -\frac{1}{3} \cos x$ and the general solution is $y(x) = y_c(x) + y_p(x) = c_1 \cos\left(\frac{1}{2}x\right) + c_2 \sin\left(\frac{1}{2}x\right) - \frac{1}{3} \cos x$.

(b) From (a) we know that $y_c(x) = c_1 \cos\frac{x}{2} + c_2 \sin\frac{x}{2}$. Setting $y_1 = \cos\frac{x}{2}$, $y_2 = \sin\frac{x}{2}$, we have $y_1y_2' - y_2y_1' = \frac{1}{2}\cos^2\frac{x}{2} + \frac{1}{2}\sin^2\frac{x}{2} = \frac{1}{2}$. Thus $u_1' = -\dfrac{\cos x \sin\frac{x}{2}}{4 \cdot \frac{1}{2}} = -\frac{1}{2}\cos\left(2 \cdot \frac{x}{2}\right)\sin\frac{x}{2} = -\frac{1}{2}\left(2\cos^2\frac{x}{2} - 1\right)\sin\frac{x}{2}$ and $u_2' = \dfrac{\cos x \cos\frac{x}{2}}{4 \cdot \frac{1}{2}} = \frac{1}{2}\cos\left(2 \cdot \frac{x}{2}\right)\cos\frac{x}{2} = \frac{1}{2}\left(1 - 2\sin^2\frac{x}{2}\right)\cos\frac{x}{2}$. Then

$u_1(x) = \int \left(\frac{1}{2}\sin\frac{x}{2} - \cos^2\frac{x}{2}\,\sin\frac{x}{2}\right)dx = -\cos\frac{x}{2} + \frac{2}{3}\cos^3\frac{x}{2}$ and

$u_2(x) = \int \left(\frac{1}{2}\cos\frac{x}{2} - \sin^2\frac{x}{2}\,\cos\frac{x}{2}\right)dx = \sin\frac{x}{2} - \frac{2}{3}\sin^3\frac{x}{2}$. Thus

$$\begin{aligned} y_p(x) &= \left(-\cos\tfrac{x}{2} + \tfrac{2}{3}\cos^3\tfrac{x}{2}\right)\cos\tfrac{x}{2} + \left(\sin\tfrac{x}{2} - \tfrac{2}{3}\sin^3\tfrac{x}{2}\right)\sin\tfrac{x}{2} = -\left(\cos^2\tfrac{x}{2} - \sin^2\tfrac{x}{2}\right) + \tfrac{2}{3}\left(\cos^4\tfrac{x}{2} - \sin^4\tfrac{x}{2}\right) \\ &= -\cos\left(2 \cdot \tfrac{x}{2}\right) + \tfrac{2}{3}\left(\cos^2\tfrac{x}{2} + \sin^2\tfrac{x}{2}\right)\left(\cos^2\tfrac{x}{2} - \sin^2\tfrac{x}{2}\right) = -\cos x + \tfrac{2}{3}\cos x = -\tfrac{1}{3}\cos x \end{aligned}$$

and the general solution is $y(x) = y_c(x) + y_p(x) = c_1 \cos\frac{x}{2} + c_2 \sin\frac{x}{2} - \frac{1}{3}\cos x$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. (a) Here $r^2 - 2r - 3 = (r-3)(r+1) = 0 \Rightarrow r = 3, r = -1$ and the complementary solution is $y_c(x) = c_1 e^{3x} + c_2 e^{-x}$. A particular solution is of the form $y_p(x) = Ax + B \Rightarrow y_p' = A, y_p'' = 0$, and substituting into the differential equation gives $0 - 2A - 3(Ax + B) = x + 2$ or $-3Ax + (-2A - 3B) = x + 2$, so $A = -\frac{1}{3}$ and $-2A - 3B = 2 \Rightarrow B = -\frac{4}{9}$. Thus $y_p(x) = -\frac{1}{3}x - \frac{4}{9}$ and the general solution is $y(x) = y_c(x) + y_p(x) = c_1 e^{3x} + c_2 e^{-x} - \frac{1}{3}x - \frac{4}{9}$.

(b) In (a), $y_c(x) = c_1 e^{3x} + c_2 e^{-x}$, so set $y_1 = e^{3x}$, $y_2 = e^{-x}$. Then $y_1 y_2' - y_2 y_1' = -e^{3x}e^{-x} - 3e^{3x}e^{-x} = -4e^{2x}$ so

$u_1' = -\dfrac{(x+2)e^{-x}}{-4e^{2x}} = \frac{1}{4}(x+2)e^{-3x} \Rightarrow u_1(x) = \frac{1}{4}\int (x+2)e^{-3x}\,dx = \frac{1}{4}\left[-\frac{1}{3}(x+2)e^{-3x} - \frac{1}{9}e^{-3x}\right]$ [by parts]

and $u_2' = \dfrac{(x+2)e^{3x}}{-4e^{2x}} = -\frac{1}{4}(x+2)e^{x} \Rightarrow u_2(x) = -\frac{1}{4}\int (x+2)e^{x}\,dx = -\frac{1}{4}[(x+2)e^x - e^x]$ [by parts].

Hence $y_p(x) = \frac{1}{4}\left[\left(-\frac{1}{3}x - \frac{7}{9}\right)e^{-3x}\right]e^{3x} - \frac{1}{4}[(x+1)e^x]e^{-x} = -\frac{1}{3}x - \frac{4}{9}$ and

$y(x) = y_c(x) + y_p(x) = c_1 e^{3x} + c_2 e^{-x} - \frac{1}{3}x - \frac{4}{9}$.

21. (a) $r^2 - 2r + 1 = (r-1)^2 = 0 \Rightarrow r = 1$, so the complementary solution is $y_c(x) = c_1 e^x + c_2 x e^x$. A particular solution is of the form $y_p(x) = Ae^{2x}$. Thus $4Ae^{2x} - 4Ae^{2x} + Ae^{2x} = e^{2x} \Rightarrow Ae^{2x} = e^{2x} \Rightarrow A = 1 \Rightarrow y_p(x) = e^{2x}$. So a general solution is $y(x) = y_c(x) + y_p(x) = c_1 e^x + c_2 x e^x + e^{2x}$.

(b) From (a), $y_c(x) = c_1 e^x + c_2 x e^x$, so set $y_1 = e^x$, $y_2 = xe^x$. Then, $y_1 y_2' - y_2 y_1' = e^{2x}(1+x) - xe^{2x} = e^{2x}$ and so $u_1' = -xe^x \Rightarrow u_1(x) = -\int xe^x\,dx = -(x-1)e^x$ [by parts] and $u_2' = e^x \Rightarrow u_2(x) = \int e^x\,dx = e^x$. Hence $y_p(x) = (1-x)e^{2x} + xe^{2x} = e^{2x}$ and the general solution is $y(x) = y_c(x) + y_p(x) = c_1 e^x + c_2 x e^x + e^{2x}$.

22. (a) Here $r^2 - r = r(r-1) = 0 \Rightarrow r = 0, 1$ and $y_c(x) = c_1 + c_2 e^x$ and so we try a particular solution of the form $y_p(x) = Axe^x$. Thus, after calculating the necessary derivatives, we get $y'' - y' = e^x \Rightarrow$ $Ae^x(2+x) - Ae^x(1+x) = e^x \Rightarrow A = 1$. Thus $y_p(x) = xe^x$ and the general solution is $y(x) = c_1 + c_2 e^x + xe^x$.

(b) From (a) we know that $y_c(x) = c_1 + c_2 e^x$, so setting $y_1 = 1$, $y_2 = e^x$, then $y_1 y_2' - y_2 y_1' = e^x - 0 = e^x$. Thus $u_1' = -e^{2x}/e^x = -e^x$ and $u_2' = e^x/e^x = 1$. Then $u_1(x) = -\int e^x dx = -e^x$ and $u_2(x) = x$. Thus $y_p(x) = -e^x + xe^x$ and the general solution is $y(x) = c_1 + c_2 e^x - e^x + xe^x = c_1 + c_3 e^x + xe^x$.

23. As in Example 5, $y_c(x) = c_1 \sin x + c_2 \cos x$, so set $y_1 = \sin x$, $y_2 = \cos x$. Then $y_1 y_2' - y_2 y_1' = -\sin^2 x - \cos^2 x = -1$,

so $u_1' = -\dfrac{\sec^2 x \cos x}{-1} = \sec x \Rightarrow u_1(x) = \int \sec x\,dx = \ln(\sec x + \tan x)$ for $0 < x < \frac{\pi}{2}$,

and $u_2' = \dfrac{\sec^2 x \sin x}{-1} = -\sec x \tan x \Rightarrow u_2(x) = -\sec x$. Hence

$y_p(x) = \ln(\sec x + \tan x)\cdot \sin x - \sec x \cdot \cos x = \sin x \ln(\sec x + \tan x) - 1$ and the general solution is

$y(x) = c_1 \sin x + c_2 \cos x + \sin x \ln(\sec x + \tan x) - 1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

24. As in Exercise 23, $y_c(x) = c_1 \sin x + c_2 \cos x$, $y_1 = \sin x$, $y_2 = \cos x$, and $y_1y_2' - y_2y_1' = -1$. Then

$u_1' = -\dfrac{\sec^3 x \cos x}{-1} = \sec^2 x \;\Rightarrow\; u_1(x) = \tan x$ and $u_2' = \dfrac{\sec^3 x \sin x}{-1} = -\sec^2 x \tan x \;\Rightarrow$

$u_2(x) = -\int \tan x \sec^2 x\, dx = -\frac{1}{2}\tan^2 x$. Hence

$y_p(x) = \tan x \sin x - \frac{1}{2}\tan^2 x \cos x = \tan x \sin x - \frac{1}{2}\tan x \sin x = \frac{1}{2}\tan x \sin x$ and the general solution is $y(x) = c_1 \sin x + c_2 \cos x + \frac{1}{2}\tan x \sin x$.

25. $y_1 = e^x$, $y_2 = e^{2x}$ and $y_1y_2' - y_2y_1' = e^{3x}$. So $u_1' = \dfrac{-e^{2x}}{(1+e^{-x})e^{3x}} = -\dfrac{e^{-x}}{1+e^{-x}}$ and

$u_1(x) = \displaystyle\int -\frac{e^{-x}}{1+e^{-x}}\,dx = \ln(1+e^{-x})$. $u_2' = \dfrac{e^x}{(1+e^{-x})e^{3x}} = \dfrac{e^x}{e^{3x}+e^{2x}}$ so

$u_2(x) = \displaystyle\int \frac{e^x}{e^{3x}+e^{2x}}\,dx = \ln\left(\frac{e^x+1}{e^x}\right) - e^{-x} = \ln(1+e^{-x}) - e^{-x}$. Hence

$y_p(x) = e^x \ln(1+e^{-x}) + e^{2x}[\ln(1+e^{-x}) - e^{-x}]$ and the general solution is

$y(x) = [c_1 + \ln(1+e^{-x})]e^x + [c_2 - e^{-x} + \ln(1+e^{-x})]e^{2x}$.

26. $y_1 = e^{-x}$, $y_2 = e^{-2x}$ and $y_1y_2' - y_2y_1' = -e^{-3x}$. So $u_1' = -\dfrac{(\sin e^x)e^{-2x}}{-e^{-3x}} = e^x \sin e^x$

and $u_2' = \dfrac{(\sin e^x)e^{-x}}{-e^{-3x}} = -e^{2x}\sin e^x$. Hence $u_1(x) = \int e^x \sin e^x\,dx = -\cos e^x$ and

$u_2(x) = \int -e^{2x}\sin e^x\,dx = e^x\cos e^x - \sin e^x$. Then $y_p(x) = -e^{-x}\cos e^x - e^{-2x}[\sin e^x - e^x \cos e^x]$

and the general solution is $y(x) = (c_1 - \cos e^x)e^{-x} + [c_2 - \sin e^x + e^x \cos e^x]e^{-2x}$.

27. $r^2 - 2r + 1 = (r-1)^2 = 0 \;\Rightarrow\; r = 1$ so $y_c(x) = c_1e^x + c_2xe^x$. Thus $y_1 = e^x$, $y_2 = xe^x$ and

$y_1y_2' - y_2y_1' = e^x(x+1)e^x - xe^xe^x = e^{2x}$. So $u_1' = -\dfrac{xe^x \cdot e^x/(1+x^2)}{e^{2x}} = -\dfrac{x}{1+x^2} \;\Rightarrow$

$u_1 = -\displaystyle\int \frac{x}{1+x^2}\,dx = -\frac{1}{2}\ln(1+x^2)$, $u_2' = \dfrac{e^x \cdot e^x/(1+x^2)}{e^{2x}} = \dfrac{1}{1+x^2} \;\Rightarrow\; u_2 = \displaystyle\int \frac{1}{1+x^2}\,dx = \tan^{-1}x$ and

$y_p(x) = -\frac{1}{2}e^x \ln(1+x^2) + xe^x \tan^{-1}x$. Hence the general solution is $y(x) = e^x\left[c_1 + c_2x - \frac{1}{2}\ln(1+x^2) + x\tan^{-1}x\right]$.

28. $y_1 = e^{-2x}$, $y_2 = xe^{-2x}$ and $y_1y_2' - y_2y_1' = e^{-4x}$. Then $u_1' = \dfrac{-e^{-2x}xe^{-2x}}{x^3e^{-4x}} = -\dfrac{1}{x^2}$ so $u_1(x) = x^{-1}$ and

$u_2' = \dfrac{e^{-2x}e^{-2x}}{x^3e^{-4x}} = \dfrac{1}{x^3}$ so $u_2(x) = -\dfrac{1}{2x^2}$. Thus $y_p(x) = \dfrac{e^{-2x}}{x} - \dfrac{xe^{-2x}}{2x^2} = \dfrac{e^{-2x}}{2x}$ and the general solution is

$y(x) = e^{-2x}[c_1 + c_2x + 1/(2x)]$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17.3 Applications of Second-Order Differential Equations

1. By Hooke's Law $k(0.25) = 25$ so $k = 100$ is the spring constant and the differential equation is $5x'' + 100x = 0$.

The auxiliary equation is $5r^2 + 100 = 0$ with roots $r = \pm 2\sqrt{5}\,i$, so the general solution to the differential equation is $x(t) = c_1 \cos\left(2\sqrt{5}\,t\right) + c_2 \sin\left(2\sqrt{5}\,t\right)$. We are given that $x(0) = 0.35 \;\Rightarrow\; c_1 = 0.35$ and $x'(0) = 0 \;\Rightarrow\; 2\sqrt{5}\,c_2 = 0 \;\Rightarrow\; c_2 = 0$, so the position of the mass after t seconds is $x(t) = 0.35\cos\left(2\sqrt{5}\,t\right)$.

2. By Hooke's Law $k(0.4) = 32$ so $k = \frac{32}{0.4} = 80$ is the spring constant and the differential equation is $8x'' + 80x = 0$.

The general solution is $x(t) = c_1 \cos\left(\sqrt{10}\,t\right) + c_2 \sin\left(\sqrt{10}\,t\right)$. But $0 = x(0) = c_1$ and $1 = x'(0) = \sqrt{10}\,c_2 \;\Rightarrow\; c_2 = \frac{1}{\sqrt{10}}$, so the position of the mass after t seconds is $x(t) = \frac{1}{\sqrt{10}} \sin\left(\sqrt{10}\,t\right)$.

3. $k(0.5) = 6$ or $k = 12$ is the spring constant, so the initial-value problem is $2x'' + 14x' + 12x = 0$, $x(0) = 1$, $x'(0) = 0$.

The general solution is $x(t) = c_1 e^{-6t} + c_2 e^{-t}$. But $1 = x(0) = c_1 + c_2$ and $0 = x'(0) = -6c_1 - c_2$. Thus the position is given by $x(t) = -\frac{1}{5}e^{-6t} + \frac{6}{5}e^{-t}$.

4. (a) $k(0.25) = 13 \;\Rightarrow\; k = 52$, so the differential equation is $2x'' + 8x' + 52x = 0$ with general solution $x(t) = e^{-2t}\left[c_1 \cos\left(\sqrt{22}\,t\right) + c_2 \sin\left(\sqrt{22}\,t\right)\right]$. Then $0 = x(0) = c_1$ and $0.5 = x'(0) = \sqrt{22}\,c_2 \;\Rightarrow\; c_2 = \frac{1}{2\sqrt{22}}$, so the position is given by $x(t) = \frac{1}{2\sqrt{22}} e^{-2t} \sin\left(\sqrt{22}\,t\right)$.

(b)



5. For critical damping we need $c^2 - 4mk = 0$ or $m = c^2/(4k) = 14^2/(4 \cdot 12) = \frac{49}{12}$ kg.

6. For critical damping we need $c^2 = 4mk$ or $c = 2\sqrt{mk} = 2\sqrt{2 \cdot 52} = 4\sqrt{26}$.

7. We are given $m = 1$, $k = 100$, $x(0) = -0.1$ and $x'(0) = 0$. From (3), the differential equation is $\dfrac{d^2x}{dt^2} + c\dfrac{dx}{dt} + 100x = 0$ with auxiliary equation $r^2 + cr + 100 = 0$.

If $c = 10$, we have two complex roots $r = -5 \pm 5\sqrt{3}\,i$, so the motion is underdamped and the solution is $x = e^{-5t}\left[c_1 \cos\left(5\sqrt{3}\,t\right) + c_2 \sin\left(5\sqrt{3}\,t\right)\right]$. Then $-0.1 = x(0) = c_1$ and $0 = x'(0) = 5\sqrt{3}\,c_2 - 5c_1 \;\Rightarrow\; c_2 = -\frac{1}{10\sqrt{3}}$, so $x = e^{-5t}\left[-0.1\cos\left(5\sqrt{3}\,t\right) - \frac{1}{10\sqrt{3}} \sin\left(5\sqrt{3}\,t\right)\right]$.

If $c = 15$, we again have underdamping since the auxiliary equation has roots $r = -\frac{15}{2} \pm \frac{5\sqrt{7}}{2}\,i$. The general solution is $x = e^{-15t/2}\left[c_1 \cos\left(\frac{5\sqrt{7}}{2}t\right) + c_2 \sin\left(\frac{5\sqrt{7}}{2}t\right)\right]$, so $-0.1 = x(0) = c_1$ and $0 = x'(0) = \frac{5\sqrt{7}}{2}c_2 - \frac{15}{2}c_1 \;\Rightarrow\; c_2 = -\frac{3}{10\sqrt{7}}$. Thus $x = e^{-15t/2}\left[-0.1\cos\left(\frac{5\sqrt{7}}{2}t\right) - \frac{3}{10\sqrt{7}} \sin\left(\frac{5\sqrt{7}}{2}t\right)\right]$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

For $c = 20$, we have equal roots $r_1 = r_2 = -10$, so the oscillation is critically damped and the solution is $x = (c_1 + c_2 t)e^{-10t}$. Then $-0.1 = x(0) = c_1$ and $0 = x'(0) = -10c_1 + c_2 \Rightarrow c_2 = -1$, so $x = (-0.1 - t)e^{-10t}$.

If $c = 25$ the auxiliary equation has roots $r_1 = -5$, $r_2 = -20$, so we have overdamping and the solution is $x = c_1 e^{-5t} + c_2 e^{-20t}$. Then $-0.1 = x(0) = c_1 + c_2$ and $0 = x'(0) = -5c_1 - 20c_2 \Rightarrow c_1 = -\frac{2}{15}$ and $c_2 = \frac{1}{30}$, so $x = -\frac{2}{15}e^{-5t} + \frac{1}{30}e^{-20t}$.

If $c = 30$ we have roots $r = -15 \pm 5\sqrt{5}$, so the motion is overdamped and the solution is $x = c_1 e^{(-15+5\sqrt{5})t} + c_2 e^{(-15-5\sqrt{5})t}$. Then $-0.1 = x(0) = c_1 + c_2$ and $0 = x'(0) = \left(-15 + 5\sqrt{5}\right)c_1 + \left(-15 - 5\sqrt{5}\right)c_2 \Rightarrow$ $c_1 = \frac{-5-3\sqrt{5}}{100}$ and $c_2 = \frac{-5+3\sqrt{5}}{100}$, so $x = \left(\frac{-5-3\sqrt{5}}{100}\right)e^{(-15+5\sqrt{5})t} + \left(\frac{-5+3\sqrt{5}}{100}\right)e^{(-15-5\sqrt{5})t}$.



8. We are given $m = 1$, $c = 10$, $x(0) = 0$ and $x'(0) = 1$. The differential equation is $\dfrac{d^2x}{dt^2} + 10\dfrac{dx}{dt} + kx = 0$ with auxiliary equation $r^2 + 10r + k = 0$. $k = 10$: the auxiliary equation has roots $r = -5 \pm \sqrt{15}$ so we have overdamping and the solution is $x = c_1 e^{(-5+\sqrt{15})t} + c_2 e^{(-5-\sqrt{15})t}$. Entering the initial conditions gives $c_1 = \frac{1}{2\sqrt{15}}$ and $c_2 = -\frac{1}{2\sqrt{15}}$, so $x = \frac{1}{2\sqrt{15}}e^{(-5+\sqrt{15})t} - \frac{1}{2\sqrt{15}}e^{(-5-\sqrt{15})t}$.

$k = 20$: $r = -5 \pm \sqrt{5}$ and the solution is $x = c_1 e^{(-5+\sqrt{5})t} + c_2 e^{(-5-\sqrt{5})t}$ so again the motion is overdamped. The initial conditions give $c_1 = \frac{1}{2\sqrt{5}}$ and $c_2 = -\frac{1}{2\sqrt{5}}$, so $x = \frac{1}{2\sqrt{5}}e^{(-5+\sqrt{5})t} - \frac{1}{2\sqrt{5}}e^{(-5-\sqrt{5})t}$.

$k = 25$: we have equal roots $r_1 = r_2 = -5$, so the motion is critically damped and the solution is $x = (c_1 + c_2 t)e^{-5t}$. The initial conditions give $c_1 = 0$ and $c_2 = 1$, so $x = te^{-5t}$.

$k = 30$: $r = -5 \pm \sqrt{5}\,i$ so the motion is underdamped and the solution is $x = e^{-5t}\left[c_1 \cos\left(\sqrt{5}\,t\right) + c_2 \sin\left(\sqrt{5}\,t\right)\right]$. The initial conditions give $c_1 = 0$ and $c_2 = \frac{1}{\sqrt{5}}$, so $x = \frac{1}{\sqrt{5}}\,e^{-5t}\sin\left(\sqrt{5}\,t\right)$.

$k = 40$: $r = -5 \pm \sqrt{15}\,i$ so we again have underdamping. The solution is $x = e^{-5t}\left[c_1 \cos\left(\sqrt{15}\,t\right) + c_2 \sin\left(\sqrt{15}\,t\right)\right]$, and the initial conditions give $c_1 = 0$ and $c_2 = \frac{1}{\sqrt{15}}$. Thus $x = \frac{1}{\sqrt{15}}e^{-5t}\sin\left(\sqrt{15}\,t\right)$.



9. The differential equation is $mx'' + kx = F_0 \cos\omega_0 t$ and $\omega_0 \neq \omega = \sqrt{k/m}$. Here the auxiliary equation is $mr^2 + k = 0$ with roots $\pm\sqrt{k/m}\,i = \pm\omega i$ so $x_c(t) = c_1 \cos\omega t + c_2 \sin\omega t$. Since $\omega_0 \neq \omega$, try $x_p(t) = A\cos\omega_0 t + B\sin\omega_0 t$. Then we need $(m)(-\omega_0^2)(A\cos\omega_0 t + B\sin\omega_0 t) + k(A\cos\omega_0 t + B\sin\omega_0 t) = F_0\cos\omega_0 t$ or $A\left(k - m\omega_0^2\right) = F_0$ and

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$B(k - m\omega_0^2) = 0$. Hence $B = 0$ and $A = \dfrac{F_0}{k - m\omega_0^2} = \dfrac{F_0}{m(\omega^2 - \omega_0^2)}$ since $\omega^2 = \dfrac{k}{m}$. Thus the motion of the mass is given by $x(t) = c_1 \cos \omega t + c_2 \sin \omega t + \dfrac{F_0}{m(\omega^2 - \omega_0^2)} \cos \omega_0 t$.

10. As in Exercise 9, $x_c(t) = c_1 \cos \omega t + c_2 \sin \omega t$. But the natural frequency of the system equals the frequency of the external force, so try $x_p(t) = t(A \cos \omega t + B \sin \omega t)$. Then we need $m(2\omega B - \omega^2 At) \cos \omega t - m(2\omega A + \omega^2 Bt) \sin \omega t + kAt \cos \omega t + kBt \sin \omega t = F_0 \cos \omega t$ or $2m\omega B = F_0$ and $-2m\omega A = 0$ [noting $-m\omega^2 A + kA = 0$ and $-m\omega^2 B + kB = 0$ since $\omega^2 = k/m$]. Hence the general solution is $x(t) = c_1 \cos \omega t + c_2 \sin \omega t + [F_0 t/(2m\omega)] \sin \omega t$.

11. From Equation 6, $x(t) = f(t) + g(t)$ where $f(t) = c_1 \cos \omega t + c_2 \sin \omega t$ and $g(t) = \dfrac{F_0}{m(\omega^2 - \omega_0^2)} \cos \omega_0 t$. Then f is periodic, with period $\frac{2\pi}{\omega}$, and if $\omega \neq \omega_0$, g is periodic with period $\frac{2\pi}{\omega_0}$. If $\frac{\omega}{\omega_0}$ is a rational number, then we can say $\frac{\omega}{\omega_0} = \frac{a}{b} \quad \Rightarrow \quad a = \frac{b\omega}{\omega_0}$ where a and b are non-zero integers. Then

$$x\left(t + a \cdot \tfrac{2\pi}{\omega}\right) = f\left(t + a \cdot \tfrac{2\pi}{\omega}\right) + g\left(t + a \cdot \tfrac{2\pi}{\omega}\right) = f(t) + g\left(t + \tfrac{b\omega}{\omega_0} \cdot \tfrac{2\pi}{\omega}\right) = f(t) + g\left(t + b \cdot \tfrac{2\pi}{\omega_0}\right) = f(t) + g(t) = x(t)$$

so $x(t)$ is periodic.

12. (a) The graph of $x = c_1 e^{rt} + c_2 t e^{rt}$ has a t-intercept when $c_1 e^{rt} + c_2 t e^{rt} = 0 \quad \Leftrightarrow \quad e^{rt}(c_1 + c_2 t) = 0 \quad \Leftrightarrow \quad c_1 = -c_2 t$. Since $t > 0$, x has a t-intercept if and only if c_1 and c_2 have opposite signs.

(b) For $t > 0$, the graph of x crosses the t-axis when $c_1 e^{r_1 t} + c_2 e^{r_2 t} = 0 \quad \Leftrightarrow \quad c_2 e^{r_2 t} = -c_1 e^{r_1 t} \quad \Leftrightarrow$ $c_2 = -c_1 \dfrac{e^{r_1 t}}{e^{r_2 t}} = -c_1 e^{(r_1 - r_2)t}$. But $r_1 > r_2 \quad \Rightarrow \quad r_1 - r_2 > 0$ and since $t > 0$, $e^{(r_1 - r_2)t} > 1$. Thus $|c_2| = |c_1|\, e^{(r_1 - r_2)t} > |c_1|$, and the graph of x can cross the t-axis only if $|c_2| > |c_1|$.

13. Here the initial-value problem for the charge is $Q'' + 20Q' + 500Q = 12$, $Q(0) = Q'(0) = 0$. Then $Q_c(t) = e^{-10t}(c_1 \cos 20t + c_2 \sin 20t)$ and try $Q_p(t) = A \quad \Rightarrow \quad 500A = 12$ or $A = \frac{3}{125}$. The general solution is $Q(t) = e^{-10t}(c_1 \cos 20t + c_2 \sin 20t) + \frac{3}{125}$. But $0 = Q(0) = c_1 + \frac{3}{125}$ and $Q'(t) = I(t) = e^{-10t}[(-10c_1 + 20c_2) \cos 20t + (-10c_2 - 20c_1) \sin 20t]$ but $0 = Q'(0) = -10c_1 + 20c_2$. Thus the charge is $Q(t) = -\frac{1}{250} e^{-10t}(6 \cos 20t + 3 \sin 20t) + \frac{3}{125}$ and the current is $I(t) = e^{-10t}\left(\frac{3}{5}\right) \sin 20t$.

14. (a) Here the initial-value problem for the charge is $2Q'' + 24Q' + 200Q = 12$ with $Q(0) = 0.001$ and $Q'(0) = 0$. Then $Q_c(t) = e^{-6t}(c_1 \cos 8t + c_2 \sin 8t)$ and try $Q_p(t) = A \quad \Rightarrow \quad A = \frac{3}{50}$ and the general solution is $Q(t) = e^{-6t}(c_1 \cos 8t + c_2 \sin 8t) + \frac{3}{50}$. But $0.001 = Q(0) = c + \frac{3}{50}$ so $c_1 = -0.059$. Also $Q'(t) = I\,(t) = e^{-6t}[(-6c_1 + 8c_2) \cos 8t + (-6c_2 - 8c_1) \sin 8t]$ and $0 = Q'(0) = -6c_1 + 8c_2$ so $c_2 = -0.04425$. Hence the charge is $Q(t) = -e^{-6t}(0.059 \cos 8t + 0.04425 \sin 8t) + \frac{3}{50}$ and the current is $I(t) = e^{-6t}(0.7375) \sin 8t$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(b)



charge, $Q(t)$



current, $I(t) = Q'(t)$

15. As in Exercise 13, $Q_c(t) = e^{-10t}(c_1 \cos 20t + c_2 \sin 20t)$ but $E(t) = 12 \sin 10t$ so try $Q_p(t) = A \cos 10t + B \sin 10t$. Substituting into the differential equation gives $(-100A + 200B + 500A) \cos 10t + (-100B - 200A + 500B) \sin 10t = 12 \sin 10t \Rightarrow$ $400A + 200B = 0$ and $400B - 200A = 12$. Thus $A = -\frac{3}{250}$, $B = \frac{3}{125}$ and the general solution is $Q(t) = e^{-10t}(c_1 \cos 20t + c_2 \sin 20t) - \frac{3}{250} \cos 10t + \frac{3}{125} \sin 10t$. But $0 = Q(0) = c_1 - \frac{3}{250}$ so $c_1 = \frac{3}{250}$. Also $Q'(t) = \frac{3}{25} \sin 10t + \frac{6}{25} \cos 10t + e^{-10t}[(-10c_1 + 20c_2) \cos 20t + (-10c_2 - 20c_1) \sin 20t]$ and $0 = Q'(0) = \frac{6}{25} - 10c_1 + 20c_2$ so $c_2 = -\frac{3}{500}$. Hence the charge is given by $Q(t) = e^{-10t}\left[\frac{3}{250} \cos 20t - \frac{3}{500} \sin 20t\right] - \frac{3}{250} \cos 10t + \frac{3}{125} \sin 10t$.

16. (a) As in Exercise 14, $Q_c(t) = e^{-6t}(c_1 \cos 8t + c_2 \sin 8t)$ but try $Q_p(t) = A \cos 10t + B \sin 10t$. Substituting into the differential equation gives $(-200A + 240B + 200A) \cos 10t + (-200B - 240A + 200B) \sin 10t = 12 \sin 10t$, so $B = 0$ and $A = -\frac{1}{20}$. Hence, the general solution is $Q(t) = e^{-6t}(c_1 \cos 8t + c_2 \sin 8t) - \frac{1}{20} \cos 10t$. But $0.001 = Q(0) = c_1 - \frac{1}{20}$, $Q'(t) = e^{-6t}[(-6c_1 + 8c_2) \cos 8t + (-6c_2 - 8c_1) \sin 8t] - \frac{1}{2} \sin 10t$ and $0 = Q'(0) = -6c_1 + 8c_2$, so $c_1 = 0.051$ and $c_2 = 0.03825$. Thus the charge is given by $Q(t) = e^{-6t}(0.051 \cos 8t + 0.03825 \sin 8t) - \frac{1}{20} \cos 10t$.

(b)



17. $x(t) = A \cos(\omega t + \delta) \Leftrightarrow x(t) = A[\cos \omega t \cos \delta - \sin \omega t \sin \delta] \Leftrightarrow x(t) = A\left(\frac{c_1}{A} \cos \omega t + \frac{c_2}{A} \sin \omega t\right)$ where $\cos \delta = c_1/A$ and $\sin \delta = -c_2/A \Leftrightarrow x(t) = c_1 \cos \omega t + c_2 \sin \omega t$. [Note that $\cos^2 \delta + \sin^2 \delta = 1 \Rightarrow c_1^2 + c_2^2 = A^2$.]

18. (a) We approximate $\sin \theta$ by θ and, with $L = 1$ and $g = 9.8$, the differential equation becomes $\frac{d^2\theta}{dt^2} + 9.8\theta = 0$. The auxiliary equation is $r^2 + 9.8 = 0 \Rightarrow r = \pm\sqrt{9.8}\, i$, so the general solution is $\theta(t) = c_1 \cos\left(\sqrt{9.8}\, t\right) + c_2 \sin\left(\sqrt{9.8}\, t\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Then $0.2 = \theta(0) = c_1$ and $1 = \theta'(0) = \sqrt{9.8}\,c_2 \quad \Rightarrow \quad c_2 = \frac{1}{\sqrt{9.8}}$, so the equation is

$\theta(t) = 0.2\cos\left(\sqrt{9.8}\,t\right) + \frac{1}{\sqrt{9.8}}\sin\left(\sqrt{9.8}\,t\right)$.

(b) $\theta'(t) = -0.2\sqrt{9.8}\sin\left(\sqrt{9.8}\,t\right) + \cos\left(\sqrt{9.8}\,t\right) = 0$ or $\tan\left(\sqrt{9.8}\,t\right) = \frac{5}{\sqrt{9.8}}$, so the critical numbers are

$t = \frac{1}{\sqrt{9.8}}\tan^{-1}\left(\frac{5}{\sqrt{9.8}}\right) + \frac{n}{\sqrt{9.8}}\,\pi$ (n any integer). The maximum angle from the vertical is

$\theta\left(\frac{1}{\sqrt{9.8}}\tan^{-1}\left(\frac{5}{\sqrt{9.8}}\right)\right) \approx 0.377$ radians (or about 21.7°).

(c) From part (b), the critical numbers of $\theta(t)$ are spaced $\frac{\pi}{\sqrt{9.8}}$ apart, and the time between successive maximum values is $2\left(\frac{\pi}{\sqrt{9.8}}\right)$. Thus the period of the pendulum is $\frac{2\pi}{\sqrt{9.8}} \approx 2.007$ seconds.

(d) $\theta(t) = 0 \quad \Rightarrow \quad 0.2\cos\left(\sqrt{9.8}\,t\right) + \frac{1}{\sqrt{9.8}}\sin\left(\sqrt{9.8}\,t\right) = 0 \quad \Rightarrow \quad \tan\left(\sqrt{9.8}\,t\right) = -0.2\sqrt{9.8} \quad \Rightarrow$

$t = \frac{1}{\sqrt{9.8}}\left[\tan^{-1}\left(-0.2\sqrt{9.8}\right) + \pi\right] \approx 0.825$ seconds.

(e) $\theta'(0.825) \approx -1.180$ rad/s.

17.4 Series Solutions

1. Let $y(x) = \sum_{n=0}^{\infty} c_n x^n$. Then $y'(x) = \sum_{n=1}^{\infty} n c_n x^{n-1}$ and the given equation, $y' - y = 0$, becomes

$\sum_{n=1}^{\infty} n c_n x^{n-1} - \sum_{n=0}^{\infty} c_n x^n = 0$. Replacing n by $n+1$ in the first sum gives $\sum_{n=0}^{\infty} (n+1)c_{n+1}x^n - \sum_{n=0}^{\infty} c_n x^n = 0$, so

$\sum_{n=0}^{\infty} [(n+1)c_{n+1} - c_n]x^n = 0$. Equating coefficients gives $(n+1)c_{n+1} - c_n = 0$, so the recursion relation is

$c_{n+1} = \dfrac{c_n}{n+1}$, $n = 0, 1, 2, \ldots$. Then $c_1 = c_0$, $c_2 = \dfrac{1}{2}c_1 = \dfrac{c_0}{2}$, $c_3 = \dfrac{1}{3}c_2 = \dfrac{1}{3}\cdot\dfrac{1}{2}c_0 = \dfrac{c_0}{3!}$, $c_4 = \dfrac{1}{4}c_3 = \dfrac{c_0}{4!}$, and

in general, $c_n = \dfrac{c_0}{n!}$. Thus, the solution is $y(x) = \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} \dfrac{c_0}{n!}\,x^n = c_0 \sum_{n=0}^{\infty} \dfrac{x^n}{n!} = c_0 e^x$.

2. Let $y(x) = \sum_{n=0}^{\infty} c_n x^n$. Then $y' = xy \quad \Rightarrow \quad y' - xy = 0 \quad \Rightarrow \quad \sum_{n=1}^{\infty} n c_n x^{n-1} - x\sum_{n=0}^{\infty} c_n x^n = 0$ or

$\sum_{n=1}^{\infty} n c_n x^{n-1} - \sum_{n=0}^{\infty} c_n x^{n+1} = 0$. Replacing n with $n+1$ in the first sum and n with $n-1$ in the second

gives $\sum_{n=0}^{\infty} (n+1)c_{n+1}x^n - \sum_{n=1}^{\infty} c_{n-1}x^n = 0$ or $c_1 + \sum_{n=1}^{\infty} (n+1)\,c_{n+1}x^n - \sum_{n=1}^{\infty} c_{n-1}x^n = 0$. Thus,

$c_1 + \sum_{n=1}^{\infty} [(n+1)c_{n+1} - c_{n-1}]\,x^n = 0$. Equating coefficients gives $c_1 = 0$ and $(n+1)\,c_{n+1} - c_{n-1} = 0$. Thus, the

recursion relation is $c_{n+1} = \dfrac{c_{n-1}}{n+1}$, $n = 1, 2, \ldots$. But $c_1 = 0$, so $c_3 = 0$ and $c_5 = 0$ and in general $c_{2n+1} = 0$. Also,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$c_2 = \frac{c_0}{2}$, $c_4 = \frac{c_2}{4} = \frac{c_0}{4 \cdot 2} = \frac{c_0}{2^2 \cdot 2!}$, $c_6 = \frac{c_4}{6} = \frac{c_0}{6 \cdot 4 \cdot 2} = \frac{c_0}{2^3 \cdot 3!}$ and in general $c_{2n} = \frac{c_0}{2^n \cdot n!}$. Thus, the solution is $y(x) = \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} c_{2n} x^{2n} = \sum_{n=0}^{\infty} \frac{c_0}{2^n \cdot n!} x^{2n} = c_0 \sum_{n=0}^{\infty} \frac{\left(x^2/2\right)^n}{n!} = c_0 e^{x^2/2}$.

3. Assuming $y(x) = \sum_{n=0}^{\infty} c_n x^n$, we have $y'(x) = \sum_{n=1}^{\infty} n c_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) c_{n+1} x^n$ and $-x^2 y = -\sum_{n=0}^{\infty} c_n x^{n+2} = -\sum_{n=2}^{\infty} c_{n-2} x^n$. Hence, the equation $y' = x^2 y$ becomes $\sum_{n=0}^{\infty} (n+1) c_{n+1} x^n - \sum_{n=2}^{\infty} c_{n-2} x^n = 0$ or $c_1 + 2c_2 x + \sum_{n=2}^{\infty} \left[(n+1)c_{n+1} - c_{n-2}\right] x^n = 0$. Equating coefficients gives $c_1 = c_2 = 0$ and $c_{n+1} = \frac{c_{n-2}}{n+1}$ for $n = 2, 3, \ldots$. But $c_1 = 0$, so $c_4 = 0$ and $c_7 = 0$ and in general $c_{3n+1} = 0$. Similarly $c_2 = 0$ so $c_{3n+2} = 0$. Finally $c_3 = \frac{c_0}{3}$, $c_6 = \frac{c_3}{6} = \frac{c_0}{6 \cdot 3} = \frac{c_0}{3^2 \cdot 2!}$, $c_9 = \frac{c_6}{9} = \frac{c_0}{9 \cdot 6 \cdot 3} = \frac{c_0}{3^3 \cdot 3!}$, …, and $c_{3n} = \frac{c_0}{3^n \cdot n!}$. Thus, the solution is $y(x) = \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} c_{3n} x^{3n} = \sum_{n=0}^{\infty} \frac{c_0}{3^n \cdot n!} x^{3n} = c_0 \sum_{n=0}^{\infty} \frac{x^{3n}}{3^n n!} = c_0 \sum_{n=0}^{\infty} \frac{\left(x^3/3\right)^n}{n!} = c_0 e^{x^3/3}$.

4. Let $y(x) = \sum_{n=0}^{\infty} c_n x^n \;\Rightarrow\; y'(x) = \sum_{n=1}^{\infty} n c_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) c_{n+1} x^n$. Then the differential equation becomes

$$(x-3)\sum_{n=0}^{\infty}(n+1)c_{n+1}x^n + 2\sum_{n=0}^{\infty}c_n x^n = 0 \;\Rightarrow\; \sum_{n=0}^{\infty}(n+1)c_{n+1}x^{n+1} - 3\sum_{n=0}^{\infty}(n+1)c_{n+1}x^n + 2\sum_{n=0}^{\infty}c_n x^n = 0 \;\Rightarrow$$

$$\sum_{n=1}^{\infty} n c_n x^n - \sum_{n=0}^{\infty} 3(n+1)c_{n+1}x^n + \sum_{n=0}^{\infty} 2c_n x^n = 0 \;\Rightarrow\; \sum_{n=0}^{\infty}\left[(n+2)c_n - 3(n+1)c_{n+1}\right]x^n = 0$$

$\left[\text{since } \sum_{n=1}^{\infty} n c_n x^n = \sum_{n=0}^{\infty} n c_n x^n\right]$. Equating coefficients gives $(n+2)c_n - 3(n+1)c_{n+1} = 0$, thus the recursion relation is $c_{n+1} = \frac{(n+2)c_n}{3(n+1)}$, $n = 0, 1, 2, \ldots$. Then $c_1 = \frac{2c_0}{3}$, $c_2 = \frac{3c_1}{3(2)} = \frac{3c_0}{3^2}$, $c_3 = \frac{4c_2}{3(3)} = \frac{4c_0}{3^3}$, $c_4 = \frac{5c_3}{3(4)} = \frac{5c_0}{3^4}$, and in general, $c_n = \frac{(n+1)c_0}{3^n}$. Thus the solution is $y(x) = \sum_{n=0}^{\infty} c_n x^n = c_0 \sum_{n=0}^{\infty} \frac{n+1}{3^n} x^n$.

$\left[\text{Note that } c_0 \sum_{n=0}^{\infty} \frac{n+1}{3^n} x^n = \frac{9c_0}{(3-x)^2} \text{ for } |x| < 3.\right]$

5. Let $y(x) = \sum_{n=0}^{\infty} c_n x^n \;\Rightarrow\; y'(x) = \sum_{n=1}^{\infty} n c_n x^{n-1}$ and $y''(x) = \sum_{n=0}^{\infty} (n+2)(n+1) c_{n+2} x^n$. The differential equation becomes $\sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n + x\sum_{n=1}^{\infty} n c_n x^{n-1} + \sum_{n=0}^{\infty} c_n x^n = 0$ or $\sum_{n=0}^{\infty}\left[(n+2)(n+1)c_{n+2} + n c_n + c_n\right]x^n = 0$

$\left[\text{since } \sum_{n=1}^{\infty} n c_n x^n = \sum_{n=0}^{\infty} n c_n x^n\right]$. Equating coefficients gives $(n+2)(n+1)c_{n+2} + (n+1)c_n = 0$, thus the recursion relation is $c_{n+2} = \frac{-(n+1)c_n}{(n+2)(n+1)} = -\frac{c_n}{n+2}$, $n = 0, 1, 2, \ldots$. Then the even coefficients are given by $c_2 = -\frac{c_0}{2}$, $c_4 = -\frac{c_2}{4} = \frac{c_0}{2 \cdot 4}$, $c_6 = -\frac{c_4}{6} = -\frac{c_0}{2 \cdot 4 \cdot 6}$, and in general,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$c_{2n} = (-1)^n \dfrac{c_0}{2 \cdot 4 \cdot \cdots \cdot 2n} = \dfrac{(-1)^n c_0}{2^n\, n!}$. The odd coefficients are $c_3 = -\dfrac{c_1}{3}$, $c_5 = -\dfrac{c_3}{5} = \dfrac{c_1}{3 \cdot 5}$, $c_7 = -\dfrac{c_5}{7} = -\dfrac{c_1}{3 \cdot 5 \cdot 7}$, and in general, $c_{2n+1} = (-1)^n \dfrac{c_1}{3 \cdot 5 \cdot 7 \cdot \cdots \cdot (2n+1)} = \dfrac{(-2)^n\, n!\, c_1}{(2n+1)!}$. The solution is

$$y(x) = c_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n\, n!} x^{2n} + c_1 \sum_{n=0}^{\infty} \frac{(-2)^n\, n!}{(2n+1)!} x^{2n+1}.$$

6. Let $y(x) = \sum_{n=0}^{\infty} c_n x^n$. Then $y''(x) = \sum_{n=2}^{\infty} n(n-1)c_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n$. Hence, the equation $y'' = y$ becomes $\sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n - \sum_{n=0}^{\infty} c_n x^n = 0$ or $\sum_{n=0}^{\infty} [(n+2)(n+1)c_{n+2} - c_n]x^n = 0$. So the recursion relation is $c_{n+2} = \dfrac{c_n}{(n+2)(n+1)}$, $n = 0, 1, \ldots$. Given c_0 and c_1, $c_2 = \dfrac{c_0}{2 \cdot 1}$, $c_4 = \dfrac{c_2}{4 \cdot 3} = \dfrac{c_0}{4!}$, $c_6 = \dfrac{c_4}{6 \cdot 5} = \dfrac{c_0}{6!}$, $\ldots$, $c_{2n} = \dfrac{c_0}{(2n)!}$ and $c_3 = \dfrac{c_1}{3 \cdot 2}$, $c_5 = \dfrac{c_3}{5 \cdot 4} = \dfrac{c_1}{5 \cdot 4 \cdot 3 \cdot 2} = \dfrac{c_1}{5!}$, $c_7 = \dfrac{c_5}{7 \cdot 6} = \dfrac{c_1}{7!}$, $\ldots$, $c_{2n+1} = \dfrac{c_1}{(2n+1)!}$. Thus, the solution is $y(x) = \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} c_{2n} x^{2n} + \sum_{n=0}^{\infty} c_{2n+1} x^{2n+1} = c_0 \sum_{n=0}^{\infty} \dfrac{x^{2n}}{(2n)!} + c_1 \sum_{n=0}^{\infty} \dfrac{x^{2n+1}}{(2n+1)!}$. The solution can be written as $y(x) = c_0 \cosh x + c_1 \sinh x$ $\left[\text{or } y(x) = c_0 \dfrac{e^x + e^{-x}}{2} + c_1 \dfrac{e^x - e^{-x}}{2} = \dfrac{c_0 + c_1}{2} e^x + \dfrac{c_0 - c_1}{2} e^{-x}\right]$.

7. Let $y(x) = \sum_{n=0}^{\infty} c_n x^n \;\Rightarrow\; y'(x) = \sum_{n=1}^{\infty} n c_n x^{n-1} = \sum_{n=0}^{\infty} (n+1)c_{n+1}x^n$ and $y''(x) = \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n$. Then

$$(x-1)y''(x) = \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^{n+1} - \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n = \sum_{n=1}^{\infty} n(n+1)c_{n+1}x^n - \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n.$$

Since $\sum_{n=1}^{\infty} n(n+1)c_{n+1}x^n = \sum_{n=0}^{\infty} n(n+1)c_{n+1}x^n$, the differential equation becomes

$$\sum_{n=0}^{\infty} n(n+1)c_{n+1}x^n - \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n + \sum_{n=0}^{\infty} (n+1)c_{n+1}x^n = 0 \;\Rightarrow$$

$$\sum_{n=0}^{\infty} [n(n+1)c_{n+1} - (n+2)(n+1)c_{n+2} + (n+1)c_{n+1}]x^n = 0 \text{ or } \sum_{n=0}^{\infty} [(n+1)^2 c_{n+1} - (n+2)(n+1)c_{n+2}]x^n = 0.$$

Equating coefficients gives $(n+1)^2 c_{n+1} - (n+2)(n+1)c_{n+2} = 0$ for $n = 0, 1, 2, \ldots$. Then the recursion relation is $c_{n+2} = \dfrac{(n+1)^2}{(n+2)(n+1)} c_{n+1} = \dfrac{n+1}{n+2} c_{n+1}$, so given c_0 and c_1, we have $c_2 = \frac{1}{2}c_1$, $c_3 = \frac{2}{3}c_2 = \frac{1}{3}c_1$, $c_4 = \frac{3}{4}c_3 = \frac{1}{4}c_1$, and in general $c_n = \dfrac{c_1}{n}$, $n = 1, 2, 3, \ldots$. Thus the solution is $y(x) = c_0 + c_1 \sum_{n=1}^{\infty} \dfrac{x^n}{n}$. Note that the solution can be expressed as $c_0 - c_1 \ln(1-x)$ for $|x| < 1$.

8. Assuming $y(x) = \sum_{n=0}^{\infty} c_n x^n$, $y''(x) = \sum_{n=2}^{\infty} n(n-1)c_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n$ and $-xy(x) = -\sum_{n=0}^{\infty} c_n x^{n+1} = -\sum_{n=1}^{\infty} c_{n-1}x^n$. The equation $y'' = xy$ becomes

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$\sum_{n=0}^{\infty}(n+2)(n+1)c_{n+2}x^n - \sum_{n=1}^{\infty}c_{n-1}x^n = 0$ or $2c_2 + \sum_{n=1}^{\infty}[(n+2)(n+1)c_{n+2} - c_{n-1}]x^n = 0$. Equating coefficients gives $c_2 = 0$ and $c_{n+2} = \dfrac{c_{n-1}}{(n+2)(n+1)}$ for $n = 1, 2, \ldots$. Since $c_2 = 0$, $c_{3n+2} = 0$ for $n = 0, 1, 2, \ldots$. Given c_0, $c_3 = \dfrac{c_0}{3\cdot 2}$, $c_6 = \dfrac{c_3}{6\cdot 5} = \dfrac{c_0}{6\cdot 5\cdot 3\cdot 2}, \ldots, c_{3n} = \dfrac{c_0}{3n(3n-1)(3n-3)(3n-4)\cdot\cdots\cdot 6\cdot 5\cdot 3\cdot 2}$. Given c_1, $c_4 = \dfrac{c_1}{4\cdot 3}$, $c_7 = \dfrac{c_4}{7\cdot 6} = \dfrac{c_1}{7\cdot 6\cdot 4\cdot 3}, \ldots, c_{3n+1} = \dfrac{c_1}{(3n+1)3n(3n-2)(3n-3)\ldots 7\cdot 6\cdot 4\cdot 3}$. The solution can be written as $y(x) = c_0\sum_{n=0}^{\infty}\dfrac{(3n-2)(3n-5)\cdot\cdots\cdot 7\cdot 4\cdot 1}{(3n)!}x^{3n} + c_1\sum_{n=0}^{\infty}\dfrac{(3n-1)(3n-4)\cdot\cdots\cdot 8\cdot 5\cdot 2}{(3n+1)!}x^{3n+1}$.

9. Let $y(x) = \sum_{n=0}^{\infty}c_nx^n$. Then $-xy'(x) = -x\sum_{n=1}^{\infty}nc_nx^{n-1} = -\sum_{n=1}^{\infty}nc_nx^n = -\sum_{n=0}^{\infty}nc_nx^n$, $y''(x) = \sum_{n=0}^{\infty}(n+2)(n+1)c_{n+2}x^n$, and the equation $y'' - xy' - y = 0$ becomes $\sum_{n=0}^{\infty}[(n+2)(n+1)c_{n+2} - nc_n - c_n]x^n = 0$. Thus, the recursion relation is $c_{n+2} = \dfrac{nc_n + c_n}{(n+2)(n+1)} = \dfrac{c_n(n+1)}{(n+2)(n+1)} = \dfrac{c_n}{n+2}$ for $n = 0, 1, 2, \ldots$. One of the given conditions is $y(0) = 1$. But $y(0) = \sum_{n=0}^{\infty}c_n(0)^n = c_0 + 0 + 0 + \cdots = c_0$, so $c_0 = 1$. Hence, $c_2 = \dfrac{c_0}{2} = \dfrac{1}{2}$, $c_4 = \dfrac{c_2}{4} = \dfrac{1}{2\cdot 4}$, $c_6 = \dfrac{c_4}{6} = \dfrac{1}{2\cdot 4\cdot 6}, \ldots$, $c_{2n} = \dfrac{1}{2^n n!}$. The other given condition is $y'(0) = 0$. But $y'(0) = \sum_{n=1}^{\infty}nc_n(0)^{n-1} = c_1 + 0 + 0 + \cdots = c_1$, so $c_1 = 0$. By the recursion relation, $c_3 = \dfrac{c_1}{3} = 0$, $c_5 = 0, \ldots, c_{2n+1} = 0$ for $n = 0, 1, 2, \ldots$. Thus, the solution to the initial-value problem is $y(x) = \sum_{n=0}^{\infty}c_nx^n = \sum_{n=0}^{\infty}c_{2n}x^{2n} = \sum_{n=0}^{\infty}\dfrac{x^{2n}}{2^n n!} = \sum_{n=0}^{\infty}\dfrac{(x^2/2)^n}{n!} = e^{x^2/2}$.

10. Assuming that $y(x) = \sum_{n=0}^{\infty}c_nx^n$, we have $x^2y = \sum_{n=0}^{\infty}c_nx^{n+2}$ and $y''(x) = \sum_{n=2}^{\infty}n(n-1)c_nx^{n-2} = \sum_{n=-2}^{\infty}(n+4)(n+3)c_{n+4}x^{n+2} = 2c_2 + 6c_3x + \sum_{n=0}^{\infty}(n+4)(n+3)c_{n+4}x^{n+2}$. Thus, the equation $y'' + x^2y = 0$ becomes $2c_2 + 6c_3x + \sum_{n=0}^{\infty}[(n+4)(n+3)c_{n+4} + c_n]x^{n+2} = 0$. So $c_2 = c_3 = 0$ and the recursion relation is $c_{n+4} = -\dfrac{c_n}{(n+4)(n+3)}$, $n = 0, 1, 2, \ldots$. But $c_1 = y'(0) = 0 = c_2 = c_3$ and by the recursion relation, $c_{4n+1} = c_{4n+2} = c_{4n+3} = 0$ for $n = 0, 1, 2, \ldots$. Also, $c_0 = y(0) = 1$, so $c_4 = -\dfrac{c_0}{4\cdot 3} = -\dfrac{1}{4\cdot 3}$, $c_8 = -\dfrac{c_4}{8\cdot 7} = \dfrac{(-1)^2}{8\cdot 7\cdot 4\cdot 3}, \ldots, c_{4n} = \dfrac{(-1)^n}{4n(4n-1)(4n-4)(4n-5)\cdot\cdots\cdot 4\cdot 3}$. Thus, the solution to the initial-value problem is $y(x) = \sum_{n=0}^{\infty}c_nx^n = c_0 + \sum_{n=0}^{\infty}c_{4n}x^{4n} = 1 + \sum_{n=1}^{\infty}(-1)^n\dfrac{x^{4n}}{4n(4n-1)(4n-4)(4n-5)\cdot\cdots\cdot 4\cdot 3}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. Assuming that $y(x) = \sum_{n=0}^{\infty} c_n x^n$, we have $xy = x\sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} c_n x^{n+1}$, $x^2y' = x^2\sum_{n=1}^{\infty} nc_n x^{n-1} = \sum_{n=0}^{\infty} nc_n x^{n+1}$,

$$y''(x) = \sum_{n=2}^{\infty} n(n-1)c_n x^{n-2} = \sum_{n=-1}^{\infty} (n+3)(n+2)c_{n+3}x^{n+1} \qquad \text{[replace } n \text{ with } n+3\text{]}$$

$$= 2c_2 + \sum_{n=0}^{\infty} (n+3)(n+2)c_{n+3}x^{n+1},$$

and the equation $y'' + x^2y' + xy = 0$ becomes $2c_2 + \sum_{n=0}^{\infty} [(n+3)(n+2)c_{n+3} + nc_n + c_n]\, x^{n+1} = 0$. So $c_2 = 0$ and the recursion relation is $c_{n+3} = \dfrac{-nc_n - c_n}{(n+3)(n+2)} = -\dfrac{(n+1)c_n}{(n+3)(n+2)}$, $n = 0, 1, 2, \ldots$. But $c_0 = y(0) = 0 = c_2$ and by the recursion relation, $c_{3n} = c_{3n+2} = 0$ for $n = 0, 1, 2, \ldots$. Also, $c_1 = y'(0) = 1$, so $c_4 = -\dfrac{2c_1}{4\cdot 3} = -\dfrac{2}{4\cdot 3}$, $c_7 = -\dfrac{5c_4}{7\cdot 6} = (-1)^2\dfrac{2\cdot 5}{7\cdot 6\cdot 4\cdot 3} = (-1)^2\dfrac{2^2 5^2}{7!}, \ldots, c_{3n+1} = (-1)^n\dfrac{2^2 5^2 \cdot \cdots \cdot (3n-1)^2}{(3n+1)!}$. Thus, the solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^n = x + \sum_{n=1}^{\infty}\left[(-1)^n\frac{2^2 5^2 \cdot \cdots \cdot (3n-1)^2 x^{3n+1}}{(3n+1)!}\right].$$

12. (a) Let $y(x) = \sum_{n=0}^{\infty} c_n x^n$. Then $x^2y''(x) = \sum_{n=2}^{\infty} n(n-1)c_n x^n = \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^{n+2}$,

$xy'(x) = \sum_{n=1}^{\infty} nc_n x^n = \sum_{n=-1}^{\infty} (n+2)c_{n+2}x^{n+2} = c_1 x + \sum_{n=0}^{\infty} (n+2)c_{n+2}x^{n+2}$, and the equation

$x^2y'' + xy' + x^2y = 0$ becomes $c_1 x + \sum_{n=0}^{\infty} \{[(n+2)(n+1) + (n+2)]c_{n+2} + c_n\}x^{n+2} = 0$. So $c_1 = 0$ and the recursion relation is $c_{n+2} = -\dfrac{c_n}{(n+2)^2}$, $n = 0, 1, 2, \ldots$. But $c_1 = y'(0) = 0$ so $c_{2n+1} = 0$ for $n = 0, 1, 2, \ldots$.

Also, $c_0 = y(0) = 1$, so $c_2 = -\dfrac{1}{2^2}$, $c_4 = -\dfrac{c_2}{4^2} = (-1)^2\dfrac{1}{4^2 2^2} = (-1)^2\dfrac{1}{2^4\,(2!)^2}$, $c_6 = -\dfrac{c_4}{6^2} = (-1)^3\dfrac{1}{2^6\,(3!)^2}, \ldots$,

$c_{2n} = (-1)^n\dfrac{1}{2^{2n}\,(n!)^2}$. The solution is $y(x) = \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} (-1)^n\dfrac{x^{2n}}{2^{2n}\,(n!)^2}$.

(b) The Taylor polynomials T_0 to T_{12} are shown in the graph. Because T_{10} and T_{12} are close together throughout the interval $[-5, 5]$, it is reasonable to assume that T_{12} is a good approximation to the Bessel function on that interval.



© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

17 Review

TRUE-FALSE QUIZ

1. True. See Theorem 17.1.3.
2. False. The differential equation is not homogeneous.
3. True. $\cosh x$ and $\sinh x$ are linearly independent solutions of this linear homogeneous equation.
4. False. $y = Ae^x$ is a solution of the complementary equation, so we have to take $y_p(x) = Axe^x$.

EXERCISES

1. The auxiliary equation is $4r^2 - 1 = 0 \Rightarrow (2r+1)(2r-1) = 0 \Rightarrow r = \pm\frac{1}{2}$. Then the general solution is $y = c_1 e^{x/2} + c_2 e^{-x/2}$.

2. The auxiliary equation is $r^2 - 2r + 10 = 0 \Rightarrow r = 1 \pm 3i$, so $y = e^x(c_1 \cos 3x + c_2 \sin 3x)$.

3. The auxiliary equation is $r^2 + 3 = 0 \Rightarrow r = \pm\sqrt{3}\,i$. Then the general solution is $y = c_1 \cos(\sqrt{3}\,x) + c_2 \sin(\sqrt{3}\,x)$.

4. The auxiliary equation is $r^2 + 8r + 16 = 0 \Rightarrow (r+4)^2 = 0 \Rightarrow r = -4$, so the general solution is $y = c_1 e^{-4x} + c_2 x e^{-4x}$.

5. $r^2 - 4r + 5 = 0 \Rightarrow r = 2 \pm i$, so $y_c(x) = e^{2x}(c_1 \cos x + c_2 \sin x)$. Try $y_p(x) = Ae^{2x} \Rightarrow y_p' = 2Ae^{2x}$ and $y_p'' = 4Ae^{2x}$. Substitution into the differential equation gives $4Ae^{2x} - 8Ae^{2x} + 5Ae^{2x} = e^{2x} \Rightarrow A = 1$ and the general solution is $y(x) = e^{2x}(c_1 \cos x + c_2 \sin x) + e^{2x}$.

6. $r^2 + r - 2 = 0 \Rightarrow r = 1, r = -2$ and $y_c(x) = c_1 e^x + c_2 e^{-2x}$. Try $y_p(x) = Ax^2 + Bx + C \Rightarrow y_p' = 2Ax + B$ and $y_p'' = 2A$. Substitution gives $2A + 2Ax + B - 2Ax^2 - 2Bx - 2C = x^2 \Rightarrow A = B = -\frac{1}{2}, C = -\frac{3}{4}$ so the general solution is $y(x) = c_1 e^x + c_2 e^{-2x} - \frac{1}{2}x^2 - \frac{1}{2}x - \frac{3}{4}$.

7. $r^2 - 2r + 1 = 0 \Rightarrow r = 1$ and $y_c(x) = c_1 e^x + c_2 x e^x$. Try $y_p(x) = (Ax + B)\cos x + (Cx + D)\sin x \Rightarrow y_p' = (C - Ax - B)\sin x + (A + Cx + D)\cos x$ and $y_p'' = (2C - B - Ax)\cos x + (-2A - D - Cx)\sin x$. Substitution gives $(-2Cx + 2C - 2A - 2D)\cos x + (2Ax - 2A + 2B - 2C)\sin x = x\cos x \Rightarrow A = 0, B = C = D = -\frac{1}{2}$. The general solution is $y(x) = c_1 e^x + c_2 x e^x - \frac{1}{2}\cos x - \frac{1}{2}(x+1)\sin x$.

8. $r^2 + 4 = 0 \Rightarrow r = \pm 2i$ and $y_c(x) = c_1 \cos 2x + c_2 \sin 2x$. Try $y_p(x) = Ax\cos 2x + Bx\sin 2x$ so that no term of y_p is a solution of the complementary equation. Then $y_p' = (A + 2Bx)\cos 2x + (B - 2Ax)\sin 2x$ and $y_p'' = (4B - 4Ax)\cos 2x + (-4A - 4Bx)\sin 2x$. Substitution gives $4B\cos 2x - 4A\sin 2x = \sin 2x \Rightarrow A = -\frac{1}{4}$ and $B = 0$. The general solution is $y(x) = c_1 \cos 2x + c_2 \sin 2x - \frac{1}{4}x\cos 2x$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $r^2 - r - 6 = 0 \Rightarrow r = -2, r = 3$ and $y_c(x) = c_1e^{-2x} + c_2e^{3x}$. For $y'' - y' - 6y = 1$, try $y_{p_1}(x) = A$. Then $y'_{p_1}(x) = y''_{p_1}(x) = 0$ and substitution into the differential equation gives $A = -\frac{1}{6}$. For $y'' - y' - 6y = e^{-2x}$ try $y_{p_2}(x) = Bxe^{-2x}$ [since $y = Be^{-2x}$ satisfies the complementary equation]. Then $y'_{p_2} = (B - 2Bx)e^{-2x}$ and $y''_{p_2} = (4Bx - 4B)e^{-2x}$, and substitution gives $-5Be^{-2x} = e^{-2x} \Rightarrow B = -\frac{1}{5}$. The general solution then is $y(x) = c_1e^{-2x} + c_2e^{3x} + y_{p_1}(x) + y_{p_2}(x) = c_1e^{-2x} + c_2e^{3x} - \frac{1}{6} - \frac{1}{5}xe^{-2x}$.

10. Using variation of parameters, $y_c(x) = c_1 \cos x + c_2 \sin x$, $u'_1(x) = -\csc x \sin x = -1 \Rightarrow u_1(x) = -x$, and $u'_2(x) = \dfrac{\csc x \cos x}{x} = \cot x \Rightarrow u_2(x) = \ln|\sin x| \Rightarrow y_p = -x \cos x + \sin x \ln|\sin x|$. The solution is $y(x) = (c_1 - x)\cos x + (c_2 + \ln|\sin x|)\sin x$.

11. The auxiliary equation is $r^2 + 6r = 0$ and the general solution is $y(x) = c_1 + c_2e^{-6x} = k_1 + k_2e^{-6(x-1)}$. But $3 = y(1) = k_1 + k_2$ and $12 = y'(1) = -6k^2$. Thus $k_2 = -2$, $k_1 = 5$ and the solution is $y(x) = 5 - 2e^{-6(x-1)}$.

12. The auxiliary equation is $r^2 - 6r + 25 = 0$ and the general solution is $y(x) = e^{3x}(c_1 \cos 4x + c_2 \sin 4x)$. But $2 = y(0) = c_1$ and $1 = y'(0) = 3c_1 + 4c_2$. Thus the solution is $y(x) = e^{3x}\left(2\cos 4x - \frac{5}{4}\sin 4x\right)$.

13. The auxiliary equation is $r^2 - 5r + 4 = 0$ and the general solution is $y(x) = c_1e^x + c_2e^{4x}$. But $0 = y(0) = c_1 + c_2$ and $1 = y'(0) = c_1 + 4c_2$, so the solution is $y(x) = \frac{1}{3}(e^{4x} - e^x)$.

14. $y_c(x) = c_1 \cos(x/3) + c_2 \sin(x/3)$. For $9y'' + y = 3x$, try $y_{p_1}(x) = Ax + B$. Then $y_{p_1}(x) = 3x$. For $9y'' + y = e^{-x}$, try $y_{p_2}(x) = Ae^{-x}$. Then $9Ae^{-x} + Ae^{-x} = e^{-x}$ or $y_{p_2}(x) = \frac{1}{10}e^{-x}$. Thus the general solution is $y(x) = c_1 \cos(x/3) + c_2 \sin(x/3) + 3x + \frac{1}{10}e^{-x}$. But $1 = y(0) = c_1 + \frac{1}{10}$ and $2 = y'(0) = \frac{1}{3}c_2 + 3 - \frac{1}{10}$, so $c_1 = \frac{9}{10}$ and $c_2 = -\frac{27}{10}$. Hence the solution is $y(x) = \frac{1}{10}[9\cos(x/3) - 27\sin(x/3)] + 3x + \frac{1}{10}e^{-x}$.

15. $r^2 + 4r + 29 = 0 \Rightarrow r = -2 \pm 5i$ and the general solution is $y = e^{-2x}(c_1 \cos 5x + c_2 \sin 5x)$. But $1 = y(0) = c_1$ and $-1 = y(\pi) = -c_1e^{-2\pi} \Rightarrow c_1 = e^{2\pi}$, so there is no solution.

16. $r^2 + 4r + 29 = 0 \Rightarrow r = -2 \pm 5i$ and the general solution is $y = e^{-2x}(c_1 \cos 5x + c_2 \sin 5x)$. But $1 = y(0) = c_1$ and $-e^{-2\pi} = y(\pi) = -c_1e^{-2\pi} \Rightarrow c_1 = 1$, so c_2 can vary and the solution of the boundary-value problem is $y = e^{-2x}(\cos 5x + c \sin 5x)$, where c is any constant.

17. Let $y(x) = \sum_{n=0}^{\infty} c_nx^n$. Then $y''(x) = \sum_{n=0}^{\infty} n(n-1)c_nx^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n$ and the differential equation becomes $\sum_{n=0}^{\infty} [(n+2)(n+1)c_{n+2} + (n+1)c_n]x^n = 0$. Thus the recursion relation is $c_{n+2} = -c_n/(n+2)$ for $n = 0, 1, 2, \ldots$. But $c_0 = y(0) = 0$, so $c_{2n} = 0$ for $n = 0, 1, 2, \ldots$. Also $c_1 = y'(0) = 1$, so $c_3 = -\dfrac{1}{3}$, $c_5 = \dfrac{(-1)^2}{3 \cdot 5}$,

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

$c_7 = \dfrac{(-1)^3}{3\cdot 5\cdot 7} = \dfrac{(-1)^3 2^3 3!}{7!}, \ldots, c_{2n+1} = \dfrac{(-1)^n\, 2^n\, n!}{(2n+1)!}$ for $n = 0, 1, 2, \ldots$. Thus the solution to the initial-value problem is $y(x) = \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} \dfrac{(-1)^n\, 2^n\, n!}{(2n+1)!}\, x^{2n+1}$.

18. Let $y(x) = \sum_{n=0}^{\infty} c_n x^n$. Then $y''(x) = \sum_{n=0}^{\infty} n(n-1)\, c_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) c_{n+2} x^n$ and the differential equation becomes $\sum_{n=0}^{\infty} [(n+2)(n+1)c_{n+2} - (n+2)c_n]x^n = 0$. Thus the recursion relation is $c_{n+2} = \dfrac{c_n}{n+1}$ for $n = 0, 1, 2, \ldots$. Given c_0 and c_1, we have $c_2 = \dfrac{c_0}{1}$, $c_4 = \dfrac{c_2}{3} = \dfrac{c_0}{1\cdot 3}$, $c_6 = \dfrac{c_4}{5} = \dfrac{c_0}{1\cdot 3\cdot 5}, \ldots,$ $c_{2n} = \dfrac{c_0}{1\cdot 3\cdot 5\cdot \cdots \cdot (2n-1)} = c_0 \dfrac{2^{n-1}(n-1)!}{(2n-1)!}$. Similarly $c_3 = \dfrac{c_1}{2}$, $c_5 = \dfrac{c_3}{4} = \dfrac{c_1}{2\cdot 4}$, $c_7 = \dfrac{c_5}{6} = \dfrac{c_1}{2\cdot 4\cdot 6}, \ldots, c_{2n+1} = \dfrac{c_1}{2\cdot 4\cdot 6\cdot \cdots \cdot 2n} = \dfrac{c_1}{2^n\, n!}$. Thus the general solution is $y(x) = \sum_{n=0}^{\infty} c_n x^n = c_0 + c_0 \sum_{n=1}^{\infty} \dfrac{2^{n-1}(n-1)!\, x^{2n}}{(2n-1)!} + c \sum_{n=0}^{\infty} \dfrac{x^{2n+1}}{2^n\, n!}$. But $\sum_{n=0}^{\infty} \dfrac{x^{2n+1}}{2^n\, n!} = x \sum_{n=0}^{\infty} \dfrac{\left(\frac{1}{2}x^2\right)^n}{n!} = x e^{x^2/2}$, so $y(x) = c_1 x e^{x^2/2} + c_0 + c_0 \sum_{n=1}^{\infty} \dfrac{2^{n-1}(n-1)!\, x^{2n}}{(2n-1)!}$.

19. Here the initial-value problem is $2Q'' + 40Q' + 400Q = 12$, $Q(0) = 0.01$, $Q'(0) = 0$. Then $Q_c(t) = e^{-10t}(c_1 \cos 10t + c_2 \sin 10t)$ and we try $Q_p(t) = A$. Thus the general solution is $Q(t) = e^{-10t}(c_1 \cos 10t + c_2 \sin 10t) + \frac{3}{100}$. But $0.01 = Q'(0) = c_1 + 0.03$ and $0 = Q''(0) = -10c_1 + 10c_2$, so $c_1 = -0.02 = c_2$. Hence the charge is given by $Q(t) = -0.02e^{-10t}(\cos 10t + \sin 10t) + 0.03$.

20. By Hooke's Law the spring constant is $k = 64$ and the initial-value problem is $2x'' + 16x' + 64x = 0$, $x(0) = 0$, $x'(0) = 2.4$. Thus the general solution is $x(t) = e^{-4t}(c_1 \cos 4t + c_2 \sin 4t)$. But $0 = x(0) = c_1$ and $2.4 = x'(0) = -4c_1 + 4c_2 \quad \Rightarrow \quad c_1 = 0$, $c_2 = 0.6$. Thus the position of the mass is given by $x(t) = 0.6e^{-4t} \sin 4t$.

21. (a) Since we are assuming that the earth is a solid sphere of uniform density, we can calculate the density ρ as follows: $\rho = \dfrac{\text{mass of earth}}{\text{volume of earth}} = \dfrac{M}{\frac{4}{3}\pi R^3}$. If V_r is the volume of the portion of the earth which lies within a distance r of the center, then $V_r = \frac{4}{3}\pi r^3$ and $M_r = \rho V_r = \dfrac{Mr^3}{R^3}$. Thus $F_r = -\dfrac{GM_r m}{r^2} = -\dfrac{GMm}{R^3} r$.

(b) The particle is acted upon by a varying gravitational force during its motion. By Newton's Second Law of Motion, $m\dfrac{d^2y}{dt^2} = F_y = -\dfrac{GMm}{R^3} y$, so $y''(t) = -k^2 y(t)$ where $k^2 = \dfrac{GM}{R^3}$. At the surface, $-mg = F_R = -\dfrac{GMm}{R^2}$, so $g = \dfrac{GM}{R^2}$. Therefore $k^2 = \dfrac{g}{R}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) The differential equation $y'' + k^2 y = 0$ has auxiliary equation $r^2 + k^2 = 0$. (This is the r of Section 17.1, not the r measuring distance from the earth's center.) The roots of the auxiliary equation are $\pm ik$, so by (11) in Section 17.1, the general solution of our differential equation for t is $y(t) = c_1 \cos kt + c_2 \sin kt$. It follows that $y'(t) = -c_1 k \sin kt + c_2 k \cos kt$. Now $y(0) = R$ and $y'(0) = 0$, so $c_1 = R$ and $c_2 k = 0$. Thus $y(t) = R \cos kt$ and $y'(t) = -kR \sin kt$. This is simple harmonic motion (see Section 17.3) with amplitude R, frequency k, and phase angle 0. The period is $T = 2\pi/k$. $R \approx 3960 \text{ mi} = 3960 \cdot 5280 \text{ ft}$ and $g = 32 \text{ ft/s}^2$, so $k = \sqrt{g/R} \approx 1.24 \times 10^{-3} \text{ s}^{-1}$ and $T = 2\pi/k \approx 5079 \text{ s} \approx 85 \text{ min}$.

(d) $y(t) = 0 \quad \Leftrightarrow \quad \cos kt = 0 \quad \Leftrightarrow \quad kt = \frac{\pi}{2} + \pi n$ for some integer $n \quad \Rightarrow \quad y'(t) = -kR \sin\left(\frac{\pi}{2} + \pi n\right) = \pm kR$. Thus the particle passes through the center of the earth with speed $kR \approx 4.899 \text{ mi/s} \approx 17{,}600 \text{ mi/h}$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

APPENDIX

Complex Numbers

1. $(5-6i)+(3+2i)=(5+3)+(-6+2)i=8+(-4)i=8-4i$

2. $\left(4-\frac{1}{2}i\right)-\left(9+\frac{5}{2}i\right)=(4-9)+\left(-\frac{1}{2}-\frac{5}{2}\right)i=-5+(-3)i=-5-3i$

3. $(2+5i)(4-i)=2(4)+2(-i)+(5i)(4)+(5i)(-i)=8-2i+20i-5i^2=8+18i-5(-1)$
$=8+18i+5=13+18i$

4. $(1-2i)(8-3i)=8-3i-16i+6(-1)=2-19i$

5. $\overline{12+7i}=12-7i$

6. $2i\left(\frac{1}{2}-i\right)=i-2(-1)=2+i \quad \Rightarrow \quad \overline{2i\left(\frac{1}{2}-i\right)}=\overline{2+i}=2-i$

7. $\dfrac{1+4i}{3+2i}=\dfrac{1+4i}{3+2i}\cdot\dfrac{3-2i}{3-2i}=\dfrac{3-2i+12i-8(-1)}{3^2+2^2}=\dfrac{11+10i}{13}=\dfrac{11}{13}+\dfrac{10}{13}i$

8. $\dfrac{3+2i}{1-4i}=\dfrac{3+2i}{1-4i}\cdot\dfrac{1+4i}{1+4i}=\dfrac{3+12i+2i+8(-1)}{1^2+4^2}=\dfrac{-5+14i}{17}=-\dfrac{5}{17}+\dfrac{14}{17}i$

9. $\dfrac{1}{1+i}=\dfrac{1}{1+i}\cdot\dfrac{1-i}{1-i}=\dfrac{1-i}{1-(-1)}=\dfrac{1-i}{2}=\dfrac{1}{2}-\dfrac{1}{2}i$

10. $\dfrac{3}{4-3i}=\dfrac{3}{4-3i}\cdot\dfrac{4+3i}{4+3i}=\dfrac{12+9i}{16-9(-1)}=\dfrac{12}{25}+\dfrac{9}{25}i$

11. $i^3=i^2\cdot i=(-1)i=-i$

12. $i^{100}=(i^2)^{50}=(-1)^{50}=1$

13. $\sqrt{-25}=\sqrt{25}\,i=5i$

14. $\sqrt{-3}\sqrt{-12}=\sqrt{3}\,i\sqrt{12}\,i=\sqrt{3\cdot 12}\,i^2=\sqrt{36}\,(-1)=-6$

15. $\overline{12-5i}=12+15i$ and $|12-15i|=\sqrt{12^2+(-5)^2}=\sqrt{144+25}=\sqrt{169}=13$

16. $\overline{-1+2\sqrt{2}\,i}=-1-2\sqrt{2}\,i$ and $\left|-1+2\sqrt{2}\,i\right|=\sqrt{(-1)^2+\left(2\sqrt{2}\right)^2}=\sqrt{1+8}=\sqrt{9}=3$

17. $\overline{-4i}=\overline{0-4i}=0+4i=4i$ and $|-4i|=\sqrt{0^2+(-4)^2}=\sqrt{16}=4$

18. Let $z=a+bi$ and $w=c+di$.

(a) $\overline{z+w}=\overline{(a+bi)+(c+di)}=\overline{(a+c)+(b+d)i}=(a+c)-(b+d)i=(a-bi)+(c-di)=\overline{z}+\overline{w}$

(b) $\overline{zw}=\overline{(a+bi)(c+di)}=\overline{(ac-bd)+(ad+bc)i}=(ac-bd)-(ad+bc)i.$

On the other hand, $\overline{z}\,\overline{w}=(a-bi)(c-di)=(ac-bd)-(ad+bc)i=\overline{zw}.$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

(c) Use mathematical induction and part (b): Let S_n be the statement that $\overline{z^n} = \overline{z}^n$. S_1 is true because $\overline{z^1} = \overline{z} = \overline{z}^1$. Assume S_k is true, that is $\overline{z^k} = \overline{z}^k$. Then $\overline{z^{k+1}} = \overline{z^{1+k}} = \overline{zz^k} = \overline{z}\,\overline{z^k}$ [part (b) with $w = z^k$] $= \overline{z}^1\overline{z}^k = \overline{z}^{1+k} = \overline{z}^{k+1}$, which shows that S_{k+1} is true. Therefore, by mathematical induction, $\overline{z^n} = \overline{z}^n$ for every positive integer n.

Another proof: Use part (b) with $w = z$, and mathematical induction.

19. $4x^2 + 9 = 0 \;\Leftrightarrow\; 4x^2 = -9 \;\Leftrightarrow\; x^2 = -\frac{9}{4} \;\Leftrightarrow\; x = \pm\sqrt{-\frac{9}{4}} = \pm\sqrt{\frac{9}{4}}\,i = \pm\frac{3}{2}i.$

20. $x^4 = 1 \;\Leftrightarrow\; x^4 - 1 = 0 \;\Leftrightarrow\; (x^2-1)(x^2+1) = 0 \;\Leftrightarrow\; x^2 - 1 = 0$ or $x^2 + 1 = 0 \;\Leftrightarrow\; x = \pm 1$ or $x = \pm i$.

21. By the quadratic formula, $x^2 + 2x + 5 = 0 \;\Leftrightarrow\; x = \dfrac{-2 \pm \sqrt{2^2 - 4(1)(5)}}{2(1)} = \dfrac{-2 \pm \sqrt{-16}}{2} = \dfrac{-2 \pm 4i}{2} = -1 \pm 2i.$

22. $2x^2 - 2x + 1 = 0 \;\Leftrightarrow\; x = \dfrac{-(-2) \pm \sqrt{(-2)^2 - 4(2)(1)}}{2(2)} = \dfrac{2 \pm \sqrt{-4}}{4} = \dfrac{2 \pm 2i}{4} = \dfrac{1}{2} \pm \dfrac{1}{2}i$

23. By the quadratic formula, $z^2 + z + 2 = 0 \;\Leftrightarrow\; z = \dfrac{-1 \pm \sqrt{1^2 - 4(1)(2)}}{2(1)} = \dfrac{-1 \pm \sqrt{-7}}{2} = -\dfrac{1}{2} \pm \dfrac{\sqrt{7}}{2}i.$

24. $z^2 + \frac{1}{2}z + \frac{1}{4} = 0 \;\Leftrightarrow\; 4z^2 + 2z + 1 = 0 \;\Leftrightarrow$

$z = \dfrac{-2 \pm \sqrt{2^2 - 4(4)(1)}}{2(4)} = \dfrac{-2 \pm \sqrt{-12}}{8} = \dfrac{-2 \pm 2\sqrt{3}\,i}{8} = -\dfrac{1}{4} \pm \dfrac{\sqrt{3}}{4}i$

25. For $z = -3 + 3i$, $r = \sqrt{(-3)^2 + 3^2} = 3\sqrt{2}$ and $\tan\theta = \frac{3}{-3} = -1 \;\Rightarrow\; \theta = \frac{3\pi}{4}$ (since z lies in the second quadrant). Therefore, $-3 + 3i = 3\sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right)$.

26. For $z = 1 - \sqrt{3}\,i$, $r = \sqrt{1^2 + \left(-\sqrt{3}\right)^2} = 2$ and $\tan\theta = \frac{-\sqrt{3}}{1} = -\sqrt{3} \;\Rightarrow\; \theta = \frac{5\pi}{3}$ (since z lies in the fourth quadrant). Therefore, $1 - \sqrt{3}\,i = 2\left(\cos\frac{5\pi}{3} + i\sin\frac{5\pi}{3}\right)$.

27. For $z = 3 + 4i$, $r = \sqrt{3^2 + 4^2} = 5$ and $\tan\theta = \frac{4}{3} \;\Rightarrow\; \theta = \tan^{-1}\left(\frac{4}{3}\right)$ (since z lies in the first quadrant). Therefore, $3 + 4i = 5\left[\cos\left(\tan^{-1}\frac{4}{3}\right) + i\sin\left(\tan^{-1}\frac{4}{3}\right)\right]$.

28. For $z = 8i$, $r = \sqrt{0^2 + 8^2} = 8$ and $\tan\theta = \frac{8}{0}$ is undefined, so $\theta = \frac{\pi}{2}$ (since z lies on the positive imaginary axis). Therefore, $8i = 8\left(\cos\frac{\pi}{2} + i\sin\frac{\pi}{2}\right)$.

29. For $z = \sqrt{3} + i$, $r = \sqrt{\left(\sqrt{3}\right)^2 + 1^2} = 2$ and $\tan\theta = \frac{1}{\sqrt{3}} \;\Rightarrow\; \theta = \frac{\pi}{6} \;\Rightarrow\; z = 2\left(\cos\frac{\pi}{6} + i\sin\frac{\pi}{6}\right)$.

For $w = 1 + \sqrt{3}\,i$, $r = 2$ and $\tan\theta = \sqrt{3} \;\Rightarrow\; \theta = \frac{\pi}{3} \;\Rightarrow\; w = 2\left(\cos\frac{\pi}{3} + i\sin\frac{\pi}{3}\right)$.

Therefore, $zw = 2 \cdot 2\left[\cos\left(\frac{\pi}{6} + \frac{\pi}{3}\right) + i\sin\left(\frac{\pi}{6} + \frac{\pi}{3}\right)\right] = 4\left(\cos\frac{\pi}{2} + i\sin\frac{\pi}{2}\right)$,

$z/w = \frac{2}{2}\left[\cos\left(\frac{\pi}{6} - \frac{\pi}{3}\right) + i\sin\left(\frac{\pi}{6} - \frac{\pi}{3}\right)\right] = \cos\left(-\frac{\pi}{6}\right) + i\sin\left(-\frac{\pi}{6}\right)$, and $1 = 1 + 0i = 1(\cos 0 + i\sin 0) \;\Rightarrow$

$1/z = \frac{1}{2}\left[\cos\left(0 - \frac{\pi}{6}\right) + i\sin\left(0 - \frac{\pi}{6}\right)\right] = \frac{1}{2}\left[\cos\left(-\frac{\pi}{6}\right) + i\sin\left(-\frac{\pi}{6}\right)\right]$. For $1/z$, we could also use the formula that precedes Example 5 to obtain $1/z = \frac{1}{2}\left(\cos\frac{\pi}{6} - i\sin\frac{\pi}{6}\right)$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. For $z = 4\sqrt{3} - 4i$, $r = \sqrt{\left(4\sqrt{3}\right)^2 + (-4)^2} = \sqrt{64} = 8$ and $\tan\theta = \frac{-4}{4\sqrt{3}} = -\frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{11\pi}{6} \Rightarrow$
$z = 8\left(\cos\frac{11\pi}{6} + i\sin\frac{11\pi}{6}\right)$. For $w = 8i$, $r = \sqrt{0^2 + 8^2} = 8$ and $\tan\theta = \frac{8}{0}$ is undefined, so $\theta = \frac{\pi}{2} \Rightarrow$
$w = 8\left(\cos\frac{\pi}{2} + i\sin\frac{\pi}{2}\right)$. Therefore, $zw = 8 \cdot 8\left[\cos\left(\frac{11\pi}{6} + \frac{\pi}{2}\right) + i\sin\left(\frac{11\pi}{6} + \frac{\pi}{2}\right)\right] = 64\left(\cos\frac{\pi}{3} + i\sin\frac{\pi}{3}\right)$,
$z/w = \frac{8}{8}\left[\cos\left(\frac{11\pi}{6} - \frac{\pi}{2}\right) + i\sin\left(\frac{11\pi}{6} - \frac{\pi}{2}\right)\right] = \cos\frac{4\pi}{3} + i\sin\frac{4\pi}{3}$, and
$1 = 1 + 0i = 1(\cos 0 + i\sin 0) \Rightarrow 1/z = \frac{1}{8}\left[\cos\left(0 - \frac{11\pi}{6}\right) + i\sin\left(0 - \frac{11\pi}{6}\right)\right] = \frac{1}{8}\left[\cos\left(\frac{\pi}{6}\right) + i\sin\left(\frac{\pi}{6}\right)\right]$.
For $1/z$, we could also use the formula that precedes Example 5 to obtain $1/z = \frac{1}{8}\left(\cos\frac{11\pi}{6} - i\sin\frac{11\pi}{6}\right)$.

31. For $z = 2\sqrt{3} - 2i$, $r = \sqrt{\left(2\sqrt{3}\right)^2 + (-2)^2} = 4$ and $\tan\theta = \frac{-2}{2\sqrt{3}} = -\frac{1}{\sqrt{3}} \Rightarrow \theta = -\frac{\pi}{6} \Rightarrow$
$z = 4\left[\cos\left(-\frac{\pi}{6}\right) + i\sin\left(-\frac{\pi}{6}\right)\right]$. For $w = -1 + i$, $r = \sqrt{2}$, $\tan\theta = \frac{1}{-1} = -1 \Rightarrow \theta = \frac{3\pi}{4} \Rightarrow$
$w = \sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right)$. Therefore, $zw = 4\sqrt{2}\left[\cos\left(-\frac{\pi}{6} + \frac{3\pi}{4}\right) + i\sin\left(-\frac{\pi}{6} + \frac{3\pi}{4}\right)\right] = 4\sqrt{2}\left(\cos\frac{7\pi}{12} + i\sin\frac{7\pi}{12}\right)$,
$z/w = \frac{4}{\sqrt{2}}\left[\cos\left(-\frac{\pi}{6} - \frac{3\pi}{4}\right) + i\sin\left(-\frac{\pi}{6} - \frac{3\pi}{4}\right)\right] = \frac{4}{\sqrt{2}}\left[\cos\left(-\frac{11\pi}{12}\right) + i\sin\left(-\frac{11\pi}{12}\right)\right] = 2\sqrt{2}\left(\cos\frac{13\pi}{12} + i\sin\frac{13\pi}{12}\right)$, and
$1/z = \frac{1}{4}\left[\cos\left(-\frac{\pi}{6}\right) - i\sin\left(-\frac{\pi}{6}\right)\right] = \frac{1}{4}\left(\cos\frac{\pi}{6} + i\sin\frac{\pi}{6}\right)$.

32. For $z = 4\left(\sqrt{3} + i\right) = 4\sqrt{3} + 4i$, $r = \sqrt{\left(4\sqrt{3}\right)^2 + 4^2} = \sqrt{64} = 8$ and $\tan\theta = \frac{4}{4\sqrt{3}} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \Rightarrow$
$z = 8\left(\cos\frac{\pi}{6} + i\sin\frac{\pi}{6}\right)$. For $w = -3 - 3i$, $r = \sqrt{(-3)^2 + (-3)^2} = \sqrt{18} = 3\sqrt{2}$ and $\tan\theta = \frac{-3}{-3} = 1 \Rightarrow \theta = \frac{5\pi}{4} \Rightarrow$
$w = 3\sqrt{2}\left(\cos\frac{5\pi}{4} + i\sin\frac{5\pi}{4}\right)$. Therefore, $zw = 8 \cdot 3\sqrt{2}\left[\cos\left(\frac{\pi}{6} + \frac{5\pi}{4}\right) + i\sin\left(\frac{\pi}{6} + \frac{5\pi}{4}\right)\right] = 24\sqrt{2}\left(\cos\frac{17\pi}{12} + i\sin\frac{17\pi}{12}\right)$,
$z/w = \frac{8}{3\sqrt{2}}\left[\cos\left(\frac{\pi}{6} - \frac{5\pi}{4}\right) + i\sin\left(\frac{\pi}{6} - \frac{5\pi}{4}\right)\right] = \frac{4\sqrt{2}}{3}\left[\cos\left(-\frac{13\pi}{12}\right) + i\sin\left(-\frac{13\pi}{12}\right)\right]$, and $1/z = \frac{1}{8}\left(\cos\frac{\pi}{6} - i\sin\frac{\pi}{6}\right)$.

33. For $z = 1 + i$, $r = \sqrt{2}$ and $\tan\theta = \frac{1}{1} = 1 \Rightarrow \theta = \frac{\pi}{4} \Rightarrow z = \sqrt{2}\left(\cos\frac{\pi}{4} + i\sin\frac{\pi}{4}\right)$. So by De Moivre's Theorem,

$$\begin{aligned}(1+i)^{20} &= \left[\sqrt{2}\left(\cos\tfrac{\pi}{4} + i\sin\tfrac{\pi}{4}\right)\right]^{20} = (2^{1/2})^{20}\left(\cos\tfrac{20\cdot\pi}{4} + i\sin\tfrac{20\cdot\pi}{4}\right) = 2^{10}(\cos 5\pi + i\sin 5\pi)\\ &= 2^{10}[-1 + i(0)] = -2^{10} = -1024\end{aligned}$$

34. For $z = 1 - \sqrt{3}\,i$, $r = \sqrt{1^2 + \left(-\sqrt{3}\right)^2} = 2$ and $\tan\theta = \frac{-\sqrt{3}}{1} = -\sqrt{3} \Rightarrow \theta = \frac{5\pi}{3} \Rightarrow z = 2\left(\cos\frac{5\pi}{3} + i\sin\frac{5\pi}{3}\right)$.
So by De Moivre's Theorem,

$$\begin{aligned}\left(1 - \sqrt{3}\,i\right)^5 &= \left[2\left(\cos\tfrac{5\pi}{3} + i\sin\tfrac{5\pi}{3}\right)\right]^5 = 2^5\left(\cos\tfrac{5\cdot 5\pi}{3} + i\sin\tfrac{5\cdot 5\pi}{3}\right) = 2^5\left(\cos\tfrac{\pi}{3} + i\sin\tfrac{\pi}{3}\right)\\ &= 32\left(\tfrac{1}{2} + \tfrac{\sqrt{3}}{2}\,i\right) = 16 + 16\sqrt{3}\,i\end{aligned}$$

35. For $z = 2\sqrt{3} + 2i$, $r = \sqrt{\left(2\sqrt{3}\right)^2 + 2^2} = \sqrt{16} = 4$ and $\tan\theta = \frac{2}{2\sqrt{3}} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \Rightarrow z = 4\left(\cos\frac{\pi}{6} + i\sin\frac{\pi}{6}\right)$.
So by De Moivre's Theorem,

$$\left(2\sqrt{3} + 2i\right)^5 = \left[4\left(\cos\tfrac{\pi}{6} + i\sin\tfrac{\pi}{6}\right)\right]^5 = 4^5\left(\cos\tfrac{5\pi}{6} + i\sin\tfrac{5\pi}{6}\right) = 1024\left[-\tfrac{\sqrt{3}}{2} + \tfrac{1}{2}i\right] = -512\sqrt{3} + 512i.$$

36. For $z = 1 - i$, $r = \sqrt{2}$ and $\tan\theta = \frac{-1}{1} = -1 \Rightarrow \theta = \frac{7\pi}{4} \Rightarrow z = \sqrt{2}\left(\cos\frac{7\pi}{4} + i\sin\frac{7\pi}{4}\right) \Rightarrow$

$$(1-i)^8 = \left[\sqrt{2}\left(\cos\tfrac{7\pi}{4} + i\sin\tfrac{7\pi}{4}\right)\right]^8 = 2^4\left(\cos\tfrac{8\cdot 7\pi}{4} + i\sin\tfrac{8\cdot 7\pi}{4}\right) = 16(\cos 14\pi + i\sin 14\pi) = 16(1 + 0i) = 16.$$

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

37. $1 = 1 + 0i = 1(\cos 0 + i \sin 0)$. Using Equation 3 with $r = 1$, $n = 8$, and $\theta = 0$, we have

$$w_k = 1^{1/8}\left[\cos\left(\frac{0 + 2k\pi}{8}\right) + i \sin\left(\frac{0 + 2k\pi}{8}\right)\right] = \cos\frac{k\pi}{4} + i \sin\frac{k\pi}{4}, \text{ where } k = 0, 1, 2, \ldots, 7.$$

$w_0 = 1(\cos 0 + i \sin 0) = 1$, $w_1 = 1\left(\cos\frac{\pi}{4} + i \sin\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$,

$w_2 = 1\left(\cos\frac{\pi}{2} + i \sin\frac{\pi}{2}\right) = i$, $w_3 = 1\left(\cos\frac{3\pi}{4} + i \sin\frac{3\pi}{4}\right) = -\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$,

$w_4 = 1(\cos\pi + i \sin\pi) = -1$, $w_5 = 1\left(\cos\frac{5\pi}{4} + i \sin\frac{5\pi}{4}\right) = -\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i$,

$w_6 = 1\left(\cos\frac{3\pi}{2} + i \sin\frac{3\pi}{2}\right) = -i$, $w_7 = 1\left(\cos\frac{7\pi}{4} + i \sin\frac{7\pi}{4}\right) = \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i$



38. $32 = 32 + 0i = 32(\cos 0 + i \sin 0)$. Using Equation 3 with $r = 32$, $n = 5$, and $\theta = 0$, we have

$$w_k = 32^{1/5}\left[\cos\left(\frac{0 + 2k\pi}{5}\right) + i \sin\left(\frac{0 + 2k\pi}{5}\right)\right] = 2\left(\cos\tfrac{2}{5}\pi k + i \sin\tfrac{2}{5}\pi k\right), \text{ where } k = 0, 1, 2, 3, 4.$$

$w_0 = 2(\cos 0 + i \sin 0) = 2$

$w_1 = 2\left(\cos\frac{2\pi}{5} + i \sin\frac{2\pi}{5}\right)$

$w_2 = 2\left(\cos\frac{4\pi}{5} + i \sin\frac{4\pi}{5}\right)$

$w_3 = 2\left(\cos\frac{6\pi}{5} + i \sin\frac{6\pi}{5}\right)$

$w_4 = 2\left(\cos\frac{8\pi}{5} + i \sin\frac{8\pi}{5}\right)$



39. $i = 0 + i = 1\left(\cos\frac{\pi}{2} + i \sin\frac{\pi}{2}\right)$. Using Equation 3 with $r = 1$, $n = 3$, and $\theta = \frac{\pi}{2}$, we have

$$w_k = 1^{1/3}\left[\cos\left(\frac{\frac{\pi}{2} + 2k\pi}{3}\right) + i \sin\left(\frac{\frac{\pi}{2} + 2k\pi}{3}\right)\right], \text{ where } k = 0, 1, 2.$$

$w_0 = \left(\cos\frac{\pi}{6} + i \sin\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2} + \frac{1}{2}i$

$w_1 = \left(\cos\frac{5\pi}{6} + i \sin\frac{5\pi}{6}\right) = -\frac{\sqrt{3}}{2} + \frac{1}{2}i$

$w_2 = \left(\cos\frac{9\pi}{6} + i \sin\frac{9\pi}{6}\right) = -i$



40. $1 + i = \sqrt{2}\left(\cos\frac{\pi}{4} + i \sin\frac{\pi}{4}\right)$. Using Equation 3 with $r = \sqrt{2}$, $n = 3$, and $\theta = \frac{\pi}{4}$, we have

$$w_k = \left(\sqrt{2}\right)^{1/3}\left[\cos\left(\frac{\frac{\pi}{4} + 2k\pi}{3}\right) + i \sin\left(\frac{\frac{\pi}{4} + 2k\pi}{3}\right)\right], \text{ where } k = 0, 1, 2.$$

$w_0 = 2^{1/6}\left(\cos\frac{\pi}{12} + i \sin\frac{\pi}{12}\right)$

$w_1 = 2^{1/6}\left(\cos\frac{3\pi}{4} + i \sin\frac{3\pi}{4}\right) = 2^{1/6}\left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i\right) = -2^{1/3} + 2^{1/3}i$

$w_2 = 2^{1/6}\left(\cos\frac{17\pi}{12} + i \sin\frac{17\pi}{12}\right)$



41. Using Euler's formula (6) with $y = \frac{\pi}{2}$, we have $e^{i\pi/2} = \cos\frac{\pi}{2} + i \sin\frac{\pi}{2} = 0 + 1i = i$.

42. Using Euler's formula (6) with $y = 2\pi$, we have $e^{2\pi i} = \cos 2\pi + i \sin 2\pi = 1$.

43. Using Euler's formula (6) with $y = \dfrac{\pi}{3}$, we have $e^{i\pi/3} = \cos\dfrac{\pi}{3} + i \sin\dfrac{\pi}{3} = \dfrac{1}{2} + \dfrac{\sqrt{3}}{2}\,i$.

44. Using Euler's formula (6) with $y = -\pi$, we have $e^{-i\pi} = \cos(-\pi) + i \sin(-\pi) = -1$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. Using Equation 7 with $x = 2$ and $y = \pi$, we have $e^{2+i\pi} = e^2 e^{i\pi} = e^2(\cos\pi + i\sin\pi) = e^2(-1+0) = -e^2$.

46. Using Equation 7 with $x = \pi$ and $y = 1$, we have $e^{\pi+i} = e^\pi \cdot e^{1i} = e^\pi(\cos 1 + i\sin 1) = e^\pi\cos 1 + (e^\pi \sin 1)i$.

47. Take $r = 1$ and $n = 3$ in De Moivre's Theorem to get

$$\begin{aligned}
[1(\cos\theta + i\sin\theta)]^3 &= 1^3(\cos 3\theta + i\sin 3\theta)\\
(\cos\theta + i\sin\theta)^3 &= \cos 3\theta + i\sin 3\theta\\
\cos^3\theta + 3(\cos^2\theta)(i\sin\theta) + 3(\cos\theta)(i\sin\theta)^2 + (i\sin\theta)^3 &= \cos 3\theta + i\sin 3\theta\\
\cos^3\theta + (3\cos^2\theta\,\sin\theta)i - 3\cos\theta\,\sin^2\theta - (\sin^3\theta)i &= \cos 3\theta + i\sin 3\theta\\
(\cos^3\theta - 3\sin^2\theta\,\cos\theta) + (3\sin\theta\,\cos^2\theta - \sin^3\theta)i &= \cos 3\theta + i\sin 3\theta
\end{aligned}$$

Equating real and imaginary parts gives $\cos 3\theta = \cos^3\theta - 3\sin^2\theta\,\cos\theta$ and $\sin 3\theta = 3\sin\theta\,\cos^2\theta - \sin^3\theta$.

48. Using Formula 6,

$$e^{ix} + e^{-ix} = (\cos x + i\sin x) + [\cos(-x) + i\sin(-x)] = \cos x + i\sin x + \cos x - i\sin x = 2\cos x$$

Thus, $\cos x = \dfrac{e^{ix} + e^{-ix}}{2}$. Similarly,

$$e^{ix} - e^{-ix} = (\cos x + i\sin x) - [\cos(-x) + i\sin(-x)] = \cos x + i\sin x - \cos x - (-i\sin x) = 2i\sin x$$

Therefore, $\sin x = \dfrac{e^{ix} - e^{-ix}}{2i}$.

49. $F(x) = e^{rx} = e^{(a+bi)x} = e^{ax+bxi} = e^{ax}(\cos bx + i\sin bx) = e^{ax}\cos bx + i(e^{ax}\sin bx) \;\Rightarrow$

$$\begin{aligned}
F'(x) &= (e^{ax}\cos bx)' + i(e^{ax}\sin bx)'\\
&= (ae^{ax}\cos bx - be^{ax}\sin bx) + i(ae^{ax}\sin bx + be^{ax}\cos bx)\\
&= a[e^{ax}(\cos bx + i\sin bx)] + b[e^{ax}(-\sin bx + i\cos bx)]\\
&= ae^{rx} + b[e^{ax}(i^2\sin bx + i\cos bx)]\\
&= ae^{rx} + bi[e^{ax}(\cos bx + i\sin bx)] = ae^{rx} + bie^{rx} = (a+bi)e^{rx} = re^{rx}
\end{aligned}$$

50. (a) From Exercise 49, $F(x) = e^{(1+i)x} \;\Rightarrow\; F'(x) = (1+i)e^{(1+i)x}$. So

$$\int e^{(1+i)x}\,dx = \frac{1}{1+i}\int F'(x)\,dx = \frac{1}{1+i}F(x) + C = \frac{1-i}{2}F(x) + C = \frac{1-i}{2}e^{(1+i)x} + C$$

(b) $\int e^{(1+i)x}\,dx = \int e^x e^{ix}\,dx = \int e^x(\cos x + i\sin x)\,dx = \int e^x\cos x\,dx + i\int e^x\sin x$ **(1)**.

Also,

$$\begin{aligned}
\frac{1-i}{2}e^{(1+i)x} &= \tfrac{1}{2}e^{(1+i)x} - \tfrac{1}{2}ie^{(1+i)x} = \tfrac{1}{2}e^{x+ix} - \tfrac{1}{2}ie^{x+ix}\\
&= \tfrac{1}{2}e^x(\cos x + i\sin x) - \tfrac{1}{2}ie^x(\cos x + i\sin x)\\
&= \tfrac{1}{2}e^x\cos x + \tfrac{1}{2}e^x\sin x + \tfrac{1}{2}ie^x\sin x - \tfrac{1}{2}ie^x\cos x\\
&= \tfrac{1}{2}e^x(\cos x + \sin x) + i\left[\tfrac{1}{2}e^x(\sin x - \cos x)\right] \quad \textbf{(2)}
\end{aligned}$$

Equating the real and imaginary parts in **(1)** and **(2)**, we see that $\int e^x\cos x\,dx = \frac{1}{2}e^x(\cos x + \sin x) + C$ and $\int e^x\sin x\,dx = \frac{1}{2}e^x(\sin x - \cos x) + C$.

© 2016 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part

30692011924956

COOLEY Res 2NO

QA303.2 S75 2016 PERSONAL

Clegg, Dan. author

Complete solutions manual for Multivariable calculus, eighth edition